1 **uBin – a manual refining tool for metagenomic bins designed for educational**
2 **purposes**
3

4 Till L.V. Bornemann[1], Sarah P. Esser[1], Tom L. Stach[1], Tim Burg[2], and Alexander J. Probst[1]
5
6 1: Institute for Environmental Microbiology and Biotechnology, Department of Chemistry,
7 University Duisburg-Essen, Germany
8 2: Tim Burg, Im Acker 59, 56072 Koblenz, Germany
9
10 To whom the correspondence should be addressed:
11 alexander.probst@uni-due.de

12
13 **Abstract**
14

Resolving bacterial and archaeal genomes from metagenomes has revolutionized our
understanding of Earth's biomes, yet producing high quality genomes from assembled
fragments has been an ever-standing problem. While automated binning software and their
combination produce prokaryotic bins in high-throughput, their manual refinement has been
slow and sometimes difficult. Here, we present uBin, a GUI-based, standalone bin refiner that
runs on all major operating platforms and was specifically designed for educational purposes.
When applied to the public CAMI dataset, refinement of bins was able to improve 78.9% of
bins by decreasing their contamination. We also applied the bin refiner as a standalone binner
to public metagenomes from the International Space Station and demonstrate the recovery of
near-complete genomes, whose replication indices indicate active proliferation of microbes in
Earth's lower orbit. uBin is an easy to install software for bin refinement, binning of simple
metagenomes and communication of metagenomic results to other scientists and in
classrooms. The software is open source and available under
https://github.com/ProbstLab/uBin.

29
30 **Keywords**
31 Genome-resolved metagenomics, genomics, genome curation, education, ISS, bacteria,
32 archaea
33
34 **The authors declare no competing interest. All data is publicly available.**

Main Text

Genome-resolved metagenomics aims at recovering genomes from shotgun sequencing data of environmental DNA. The genomes allow determination of the metabolic capacities of the individual community members and provide the basis for many downstream 'omics techniques like metatranscriptomics and metaproteomics. Results from these technologies can provide important insight into the interactions of microbes within the community and with the environment [1-2]. While long-read sequencing can nowadays produce complete genomes from environmental samples [3], the percentage of closed genomes from complex ecosystems remains, however, as low as 5.3% [4]. Consequently, genomes need to be binned from metagenomes using genome-wide shared characteristics like their similar abundance pattern and k-mer frequencies [5-6]. Many automatic and semi-automatic tools have been developed to extract genomes from metagenomes [2,7–10]. The quality of the resulting bins, however, can vary greatly depending on metagenome complexity, sample type or microbial community characteristics [2]. Recent studies have shown that contamination in genomes from metagenomes in public databases is a frequent occurrence [11-12] and suggested genome curation as a mandatory analysis step prior to genome submission to public databases [13].

While established tools exist to determine the bin quality [2,14], i.e. searching candidate genomes for ubiquitous or specific marker genes to evaluate completeness and contamination, tools to improve upon the bin quality are sparse. The Anvi'o analysis and visualization platform [15] can be used to visualize genome or metagenome characteristics in a multi-level circosplot and thus allow the definition or curation of genomes. However, Anvi'o is limited in the amount of scaffolds that can be curated and needs basic knowledge in bioinformatics. Similarly, the database ggKbase [16] has a genome curation and binning functionality but is not an open platform for a wider audience. Consequently, we developed uBin as an interactive graphical-user interface that is easy to install on Mac OS, Windows, and Ubuntu for usage in, e.g., classrooms. uBin is inspired by ggKbase and enables the curation of genomes based on a combination of GC content, coverage and taxonomy and couples this to information on completeness and contamination for supervised binning. In addition, uBin can be directly used as a standalone software to bin genomes from low complexity samples.

We tested the performance of uBin (MacOS, 16 GB of RAM) on simulated datasets with varying complexity of the Critical Assessment of Metagenome Interpretation (CAMI) challenge. The pre-assembled CAMI scaffolds were binned using four automated binners (using tetranucleotide frequency and differential coverage) and the results were aggregated using DAS Tool [2] (see Supplementary Methods for details). The dereplicated bins were curated using uBin, and the quality of the bins before and after curation was compared to the correct assignment based on the CAMI dataset (see *Table S1* for Bin completeness &

contamination statistics). uBin curated bins showed a highly significant quality improvement in medium ($p < 10^{-4}$) and high complexity datasets ($p < 10^{-5}$), using both paired t-test and unpaired Kruskal-Wallis tests (*Fig 1A*). No significant difference could be detected for the low complexity dataset ($p > 0.70 / 0.65$).

Fig. 1 | Performance of uBin on simulated and real datasets with varying degrees of complexity. A: Violin plots of the F-score (mean between recall and precision) of genomes prior to uBin curation (pre uBin) and after uBin curation (post uBin) across simulated low, medium and high complexity datasets of the CAMI challenge as well as real world metagenomic datasets of medium (Tomsk) and high (SulCav AS07-7) complexity. Unpaired Kruskal-Wallis p-values are depicted. **B:** Histograms of the F-score differences for each bin

83 prior to and post uBin curation and their density distribution. Paired Welch t-test p-values are
84 shown.
85

The bin quality of the low complexity dataset was significantly higher than the bin
quality in medium (0.197 higher F-score, $p < 10^{-6}$) and high complexity (0.118 higher F-score,
$p < 10^{-4}$) datasets (ANOVA coupled to TukeyHSD, $p < 2 \times 10^{-6}$) after DAS Tool [2] bin
aggregation. Subsequent to curation with uBin the differences between these datasets were
much less pronounced (ANOVA, $p < 0.01$), with only the high to medium complexity dataset
showing a significant difference ($p < 0.01$, average 0.077 higher F-score in high complexity).
We conclude that low complexity datasets bin very well with automated binners, while medium
to high complexity datasets can greatly benefit from manual curation.

To challenge the above-mentioned conclusion, we applied uBin for the curation of bins
from environmental metagenomes of medium and high complexity. As the true genome
composition is unknown for these datasets, we used CheckM [14] to assess the completeness
and contamination of constructed genomic bins. CheckM [14] is an independent metric
compared to the marker sets used within DAS Tool [2] and uBin (see *Tab S1* for Bin statistics).
We detected a significant improvement in genome quality when using uBin curation and
directly comparing the bins in paired tests (p-values are provided in *Fig. 1*B).

Following the conclusion that binning of low complexity genomes can be achieved
easily, we tested uBin's capability as a standalone binner compared to Emergent-Self-
Organizing Maps (ESOMs) [8] on public metagenomes of the International Space Station
(ISS). uBin outperformed ESOM-based binning when used as a standalone tool and when
used as a curation tool of the ESOM bins (*Fig. 2A*, see Supplementary Material for details).
Using uBin, we successfully reconstructed 53 genomes with at least 94 percent completeness
(*Fig. 2B*) and only 6% or less contamination. When comparing their phylogenetic placement
based on 16 ribosomal proteins to the taxonomic classification of uBin, we observed
agreement between the taxonomic classification methods. The one exception was the
genome ISS_JPL_2332_S1_L003_Corynebacterium_afermentans_66_84**,** which was phylo-
genetically placed next to a *Turicella* genome [17]. This genome has since been reclassified
as *Corynebacterium otitidis* ATCC 51513 (NZ_AHAE00000000, see File S1 for the full
phylogenetic tree).

Fig. 2 | Reconstruction of genomes from the ISS, scoring of their curation and their phylogeny. A: Comparison of genome statistics after ESOM 4-mer binning, after uBin curation and after standalone binning using uBin. p-values correspond to paired Welch t-tests. **B:** Phylogenetic reconstruction based on the concatenation of 16 ribosomal proteins of 53 genomes from ISS metagenomes when using uBin as standalone binner. Branch colors indicate phyla assignments with coloring of leaves on tree displaying the sampling origin of the genomes. Genomes from PMA-treated samples (see main text) are highlighted with a red circle. The bargraphs on the right panel display completeness, contamination, genome size, GC content, coverage (relative abundance based on read-mapping) and the *in situ* replication measure (iRep [1]). **C:** Replication index dependency on flight of origin and significance testing thereof using ANOVA followed by TukeyHSD. **D:** Effect of PMA-treatment for removal of extracellular DNA on iRep of genomes from PMA-treated samples having increased iRep variance but no significant differences in iRep value based on paired Wilcoxon and paired t-tests (n=7 per group). Genomes were paired based on sample ID as well as using their shared uBin-taxonomy and GC content.

These bins represent an important step for space science since these are the first environmental genomes reconstructed from the ISS or associated transport flights. To investigate if the genomes are actively replicated under these conditions, we calculated the *in situ* replication measure iRep [1] for 43 out of 53 genomes. Across all sampling sites, the

replication rates of the recovered population genomes varied from 1.20 to 2.55, which implies an active metabolism. For instance, the lowest iRep value, which was calculated for *Methylobacterium aquaticum*, indicated that on average 20% of its sampled population was undergoing genome replication. While closely related organisms often had similar replication measures (*Fig. S3*), the main discriminatory factor for varying replication indices was the origin of the flight (*Fig. 2C*) indicating community-wide shifts in replication between the different flights. The dataset also enabled the answer to a long-standing question of indoor microbiology relating to how external DNA influences the measurements of iRep values in metagenomics. Samples of the third sampled ISS flight were analyzed using both regular metagenomics as well as metagenomics following propidium monoazide (PMA) treatment, which removes external DNA fragments and enables DNA sequencing of cells with intact membranes. When comparing the iRep values of the paired samples (n=7 per group), no significant difference could be observed (paired t- and Wilcoxon-tests, *Fig. 2D*), although the variance of the iRep values increased tremendously after PMA treatment. Equivalence testing confirmed that there are no differences between these two sample types ($p < 0.01$). We suggest that PMA-treatment can improve the accuracy of iRep measures of environmental samples and recommend its usage where appropriate.

The herein presented uBin software is designed for improvement of bins and as a standalone binner for simple metagenomes with few species. It is independent of the operating system (available for Windows, MacOS, Linux) and GUI-based so that a wide audience of non-bioinformaticians can make use of it. The initial data processing (as general metagenomic data processing) necessitates bioinformatics knowledge but respective easy-to-use wrapper scripts are provided along with the software. Thus, uBin is ideally used by bioinformaticians to communicate metagenomic data to non-bioinformatics peers and to students in classrooms. After binning or curation with uBin, the user can deploy each genome into individual fasta files. These genomes can then be further explored for metabolic analyses with, e.g., MAGE [18] or KEGG mapper [19]. Consequently, uBin represents an important software link between automated binners along with the widely-used software DAS Tool and downstream analyses including genome refinement to completion [20].

Acknowledgments

This study was funded by the Ministerium für Kultur und Wissenschaft des Landes Nordrhein-Westfalen ("Nachwuchsgruppe Dr. Alexander Probst"). We thank the students who tested and worked with uBin over the last two years in classrooms. We thank Christine Sun for her contribution to the script for calculating consensus taxonomy of scaffolds and Kasthuri Venkateswaran for input regarding sampling locations of the ISS samples.

References

1. Brown CT, Olm MR, Thomas BC, Banfield JF. Measurement of bacterial replication rates
in microbial communities. Nature Biotechnology. 2016;34:1256.

2. Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, et al. Recovery of
genomes from metagenomes via a dereplication, aggregation and scoring strategy. Nature
Microbiology. 2018;3:836–43.

3. Moss EL, Maghini DG, Bhatt AS. Complete, closed bacterial genomes from microbiomes
using nanopore sequencing. Nat Biotechnol. Nature Publishing Group; 2020;1–7.

4. Singleton CM, Petriglieri F, Kristensen JM, Kirkegaard RH, Michaelsen TY, Andersen MH,
et al. Connecting structure to function with the recovery of over 1000 high-quality activated
sludge metagenome-assembled genomes encoding full-length rRNA genes using long-read
sequencing. bioRxiv. Cold Spring Harbor Laboratory; 2020;2020.05.12.088096.

5. Teeling H, Meyerdierks A, Bauer M, Amann R, Glöckner FO. Application of
tetranucleotide frequencies for the assignment of genomic fragments. Environ Microbiol.
2004;6:938–47.

6. Albertsen M, Hugenholtz P, Skarshewski A, Nielsen KL, Tyson GW, Nielsen PH. Genome
sequences of rare, uncultured bacteria obtained by differential coverage binning of multiple
metagenomes. Nat Biotechnol. 2013;31:533–8.

7. Wu Y-W, Simmons BA, Singer SW. MaxBin 2.0: an automated binning algorithm to
recover genomes from multiple metagenomic datasets. Bioinformatics. 2016;32:605–7.

8. Dick GJ, Andersson AF, Baker BJ, Simmons SL, Thomas BC, Yelton AP, et al.
Community-wide analysis of microbial genome sequence signatures. Genome Biology.
2009;10:R85.

9. Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, Ijaz UZ, et al. Binning
metagenomic contigs by coverage and composition. Nat Methods. 2014;11:1144–6.

10. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: an adaptive
binning algorithm for robust and efficient genome reconstruction from metagenome
assemblies. PeerJ. 2019;7:e7359.

11. Shaiber A, Eren AM. Composite Metagenome-Assembled Genomes Reduce the Quality
of Public Genome Repositories. mBio [Internet]. American Society for Microbiology; 2019;
10.

12. Ballenghien M, Faivre N, Galtier N. Patterns of cross-contamination in a multispecies
population genomic project: detection, quantification, impact, and solutions. BMC Biology.
2017;15:25.

13. Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-Smith M, Doud D, Reddy TBK, et
al. Minimum information about a single amplified genome (MISAG) and a metagenome-
assembled genome (MIMAG) of bacteria and archaea. Nat Biotechnol. 2017;35:725–31.

14. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the
quality of microbial genomes recovered from isolates, single cells, and metagenomes.
Genome Res. 2015;25:1043–55.

15. Eren AM, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML, et al. Anvi'o: an
advanced analysis and visualization platform for 'omics data. PeerJ. PeerJ Inc.;
2015;3:e1319.

16. Wrighton KC, Thomas BC, Sharon I, Miller CS, Castelle CJ, VerBerkmoes NC, et al.
Fermentation, hydrogen, and sulfur metabolism in multiple uncultivated bacterial phyla.
Science. 2012;337:1661–5.

17. Brinkrolf K, Schneider J, Knecht M, Rückert C, Tauch A. Draft genome sequence of
Turicella otitidis ATCC 51513, isolated from middle ear fluid from a child with otitis media. J
Bacteriol. 2012;194:5968–9.

18. Vallenet D, Engelen S, Mornico D, Cruveiller S, Fleury L, Lajus A, et al. MicroScope: a
platform for microbial genome annotation and comparative genomics. Database (Oxford);
2009.

19. Kanehisa M, Sato Y. KEGG Mapper for inferring cellular functions from protein
sequences. Protein Sci. 2020;29:28–35.

20. Chen L-X, Anantharaman K, Shaiber A, Eren AM, Banfield JF. Accurate and Complete
Genomes from Metagenomes. bioRxiv. Cold Spring Harbor Laboratory; 2019;808410.

21. JN Fass NJ. Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ
files. https://github.com/najoshi/sickle; 2011.

22. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile
metagenomic assembler. Genome Res. 2017;27:824–34.

23. Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal:
prokaryotic gene recognition and translation initiation site identification. BMC Bioinformatics.
2010;11:119.

24. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND.
Nature Methods. 2015;12:59–60.

25. Suzek BE, Huang H, McGarvey P, Mazumder R, Wu CH. UniRef: comprehensive and
non-redundant UniProt reference clusters. Bioinformatics. 2007;23:1282–8.

26. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature Methods.
2012;9:357–9.

27. Probst AJ, Castelle CJ, Singh A, Brown CT, Anantharaman K, Sharon I, et al. Genomic
resolution of a cold subsurface aquifer community provides metabolic insights for novel
microbes adapted to high CO_2 concentrations. Environ Microbiol. 2017;19:459–74.

28. Eddy SR. Accelerated Profile HMM Searches. PLoS Comput Biol. 2011;7:e1002195.

29. Brown CT, Hug LA, Thomas BC, Sharon I, Castelle CJ, Singh A, et al. Unusual biology
across a group comprising more than 15% of domain Bacteria. Nature. 2015;523:208.

30. Bornemann TLV, Adam PS, Turzynski V, Schreiber U, Figueroa-Gonzalez PA, Rahlff J,
et al. Geological degassing enhances microbial metabolism in the continental subsurface.
bioRxiv. Cold Spring Harbor Laboratory; 2020;2020.03.07.980714.

31. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool.
J Mol Biol. 1990;215:403–10.

32. Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, et al. A new
view of the tree of life. Nature Microbiology. 2016;1:16048.

33. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic acids research. 2004;32:1792–1797.

34. Criscuolo A, Gribaldo S. BMGE (Block Mapping and Gathering with Entropy): a new
software for selection of phylogenetic informative regions from multiple sequence
alignments. BMC Evol Biol. 2010;10:210.

35. Price MN, Dehal PS, Arkin AP. FastTree: computing large minimum evolution trees with
profiles instead of a distance matrix. Molecular biology and evolution. 2009;26:1641–1650.

36. Huson DH, Richter DC, Rausch C, Dezulian T, Franz M, Rupp R. Dendroscope: An
interactive viewer for large phylogenetic trees. BMC Bioinformatics. 2007;8:460.

37. R Core Team. R: A Language and Environment for Statistical Computing. 2008;R
Foundation for Statistical Computing,   Vienna, Austria. Available from: http://www.R-
project.org

38. Welch BL. The generalisation of student's problems when several different population
variances are involved. Biometrika. 1947;34:28–35.

39. Kruskal WH, Wallis WA. Use of Ranks in One-Criterion Variance Analysis. Journal of the
American Statistical Association. Taylor & Francis; 1952;47:583–621.

40. St»hle L, Wold S. Analysis of variance (ANOVA). Chemometrics and Intelligent
Laboratory Systems. 1989;6:259–72.

41. Haynes W. Tukey's Test. In: Dubitzky W, Wolkenhauer O, Cho K-H, Yokota H, editors.
Encyclopedia of Systems Biology. New York, NY: Springer New York; 2013; p. 2303–4.

42. Wickham H. ggplot2: Elegant Graphics for Data Analysis. New York: Springer-Verlag;
2009.

43. Lakens D, Scheel AM, Isager PM. Equivalence Testing for Psychological Research: A
Tutorial: Advances in Methods and Practices in Psychological Science; 2018.

277
278
279

Supplementary Material for:

uBin – a manual refining tool for metagenomic bins designed for educational purposes

 Till L.V. Bornemann[1], Sarah P. Esser[1], Tom L. Stach[1], Tim Burg[2], and Alexander J. Probst[1]

1: Institute for Environmental Microbiology and Biotechnology, Department of Chemistry, University Duisburg-Essen, Germany

2: Tim Burg, Im Acker 59, 56072 Koblenz, Germany

To whom the correspondence should be addressed:

alexander.probst@uni-due.de

Contents

Supplementary Methods

Software implementation. uBin is written in TypeScript(3.2+)/JavaScript. It utilizes React (https://reactjs.org/) for its user interface and Redux (https://redux.js.org/) to manage the application state/data.

All imported data is stored in a local SQLite (sqlite3) database. Communication between uBin and the database is abstracted through TypeORM (https://typeorm.io/), an ORM written in TypeScript. To build the application and to provide cross-platform support, we use Electron (https://www.electronjs.org/).

The user interface uses HTML/CSS + Blueprint JS (a User-Interface (UI) toolkit, https://blueprintjs.com/) for general UI elements, react-vis (https://uber.github.io/react-vis/) for its Sunburst plot, and VX (a library for d3-based React visualization components,

https://github.com/hshoff/vx) for every other plot. Crossfilter
(https://github.com/crossfilter/crossfilter) is used to calculate the data to be plotted on-the-fly.
Metagenomic data assembly and processing. Quality control of ISS metagenome raw
reads was performed using BBduk (B Bushnell, http://jgi.doe.gov/data-and-tools/bb-tools/)
and Sickle [21]. Reads were assembled into contigs and scaffolded using metaSPAdes 3.12
[22] (see *Table S4* for read and assembly statistics). Genes were predicted for scaffolds larger
than 1 kbp using Prodigal [23] in meta mode and annotated using DIAMOND [24] against
UniRef100 (state Dec. 2017) [25], modified with NCBI taxonomic information of the respective
protein sequences (FunTaxDB, tentatively accessible through https://uni-duisburg-
essen.sciebo.de/s/pi4cuYwyZ3KJVMl). The consensus taxonomy of each scaffold was
predicted by considering the taxonomic rank of each protein on the scaffold on each taxonomic
level and choosing the lowest taxonomic rank when more than 50% of the protein taxonomies
agree. Reads were mapped to scaffolds using Bowtie2 [26] and the average scaffold coverage
was estimated along with scaffolds' length and GC content. Previously published ubiquitous
single copy genes [27] were identified using HHmer 3.2 [28] and custom tables collecting GC,
coverage, length, taxonomy and presence / absence of single copy genes of scaffolds were
generated using scripts available along with uBin under https://github.com/ProbstLab/uBin-
helperscripts.
Binning and curation. ISS assemblies were binned using Emergent Self-Organizing Maps
(ESOM) [8]. Scaffolds were fragmented using the esomWrapper.pl [8] script, using 10kbp and
5kbp as maximum and minimum fragment sizes respectively. *Streptomyces griseus*
NBRC13350 (high GC, NC_010572.1) and *Escherichia coli* K12 (low GC, NC_000913.3)
genomes were spiked in to verify successful ESOM training. For ESOM training, the starting
radius was set to 50 and the map-size was adjusted to the suggested size in the
esomWrapper.pl output. ISS data was additionally binned directly using uBin.
CAMI datasets were binned using the automatic binners abawaca [29] and MaxBin2
[7], using both 3 kbp and 5 kbp as well as 5kbp and 10kbp as minimum and maximum fragment
sizes respectively as abawaca input and using both available marker gene sets of MaxBin2
for binning. The output of the four different binners was aggregated using DAS Tool [2]. Tomsk
and SulCav binning has been described previously [30].
Tables containing Bin, GC, coverage, length, taxonomy and single copy gene
presence / absence information were loaded into uBin and used to curate draft genomes.
Coding regions and single copy genes on genomes were predicted as described, omitting the
-meta flag in prodigal.
Calculation of in situ replication indices. Bacterial *in situ* replication indices (iRep [1]) were
calculated by mapping reads on the genomes and filtered for 3 mismatches, which correspond

to 2% mismatch rate in the 150 bp reads. The rest of the settings for the iRep software were
default.

Estimation of sample complexity. The sample complexity was estimated using the diversity
of the _rpS3_ marker gene. _rpS3_ genes were annotated as described above. We are aware that
sample complexity can also stem from other factors like K-mer frequency or coverage
distribution patterns that this estimation does not take into account. However, these metrics
cannot be assessed for environmental samples easily as the real composition is unknown.
See _Table S3_ for _rpS3_ based complexity estimates across analyzed samples.

Phylogenomics. Ribosomal proteins were identified with blastp [31] (e-value 10^{-5}) against 16
ribosomal proteins set as used in [32], aligned using muscle [33] with default parameters,
trimmed with BMGE [34] and the BLOSUM30 substitution matrix and concatenated. The
phylogenetic tree was calculated using Fasttree 2.1.8 [35] with default parameters. The tree
was visualized in Dendroscope 3.7.2 [36].

Calculation of F-scores. Precision and recall of CAMI bins were determined using the known
genomic assignment of the scaffolds and where they were allocated to during binning and
curation. Genomic bins were assigned as corresponding to a CAMI genome based on the
maximum scaffolds belonging to the same CAMI genome. Precision and recall of genomes
from real-world datasets were determined using completeness as a proxy for True Positives,
1-%completeness as False Negatives, contamination as a proxy for False Positives and 1-
contamination as True Negatives. The F-score was calculated as the mean between precision
and recall.

Statistical evaluation. Statistical evaluation was performed in R [37]. Both paired and
unpaired Welch t-tests [38] as well as Kruskal-Wallis [39] tests, one- and two-way ANOVA's
[40] and TukeyHSD [41] significance tests were performed. ggplot2 [42] was used to visualize
data. The TOSTpaired.raw function within the TOSTER [43] package was used to confirm the
non-significance of PMA-related tests, using 0.1 as the equivalence bound.

Metagenome availability. Accessions to raw reads and assemblies used in this study are
listed in _Table S3_.

Software availability. The platform-independent genome curation software uBin is freely
available under the MIT license at https://github.com/ProbstLab/uBin. The installation of the
software from the OS-dedicated installers is dependency-free, while source code installation
requires a Unix-based OS and package managers like npm or yarn.

384
385
386
387

Supplementary Figures

390
Fig S1 | Comparison of bin qualities before and after uBin curation in simulated
datasets with varying complexity. Compared are the bin qualities within bins before uBin
(A) and within bins after uBin (B) in different complexities. ANOVA followed by the TukeyHSD
post-hoc test were used to identify significant differences in quality between complexity
groups.

Fig S2 | Performance of uBin on simulated and real datasets with varying degrees of complexity. A: Violin plots of the F-score (mean between recall and precision) of genomes prior to uBin curation (pre uBin) and after uBin curation (post uBin) across simulated low, medium and high complexity datasets of the CAMI challenge as well as real world metagenomic datasets of medium (Tomsk) and high (SulCav AS07-7) complexity. Unpaired Welch t-test p values are depicted. **B:** Histograms of the F-score differences for each bin prior and post uBin curation and their density distribution. Paired Welch t-test p-values are depicted.

408
409
410 **Fig S3 | iRep distribution of ISS genomes by phylum.** One- and two-way ANOVA were
411 performed for significance testing. The Betaproteobacteria were excluded for the statistical
412 analyses because of too few and highly diverging datapoints. No significant influence of the
413 species was determined while the flight of origin was a significant coefficient in the two-way
414 ANOVA analyses.
415
416
417

Supplementary Tables

Table S1 | Genome statistics of recovered ISS genomes from metagenomes based on CheckM [14].

Genome	Complete-ness	Contami-nation
ISS_JPL_2332_S1_L003_Corynebacterium_afermentans_66_84	92.99	0.44
ISS_JPL_5492_S3_L3_Azohydromonas_australica_70_64	98.22	0.62
ISS_JPL_5492_S3_L3_Caulobacter_vibrioides_69_50	99.97	1.14
ISS_JPL_5501_S5_L3_Acidovorax_69_463	99.81	2.19
ISS_JPL_5501_S5_L3_Caulobacter_vibrioides_69_106	99.97	1.14
ISS_JPL_5501_S5_L3_Sphingomonas_65_22	81.37	0.85
ISS_JPL_5629_S1_L5_Enterobacterales_58_141	100.00	0.32
ISS_JPL_5629_S1_L5_Firmicutes_33_15	98.06	0.28
ISS_JPL_5630_S2_L5_Enterobacterales_56_407	99.97	0.33
ISS_JPL_5631_S3_L5_Methylobacterium_69_176	99.06	1.25
ISS_JPL_5631_S3_L5_Methylobacterium_aquaticum_71_106	99.37	0.16
ISS_JPL_5632_S4_L5_Staphylococcus_33_30	78.70	1.14
ISS_JPL_5633_S5_L5_Pantoea_sp._3.5.1_56_354	100.00	0.12
ISS_JPL_5635_S6_L5_Methylobacterium_aquaticum_71_83	100.00	0.16
ISS_JPL_5640_S8_L5_Enterobacterales_58_121	100.00	0.34
ISS_JPL_5644_S12_L6_Pantoea_sp._3.5.1_56_351	99.94	0.12
ISS_JPL_5647_S14_L6_Methylobacterium_69_10	79.10	1.88
ISS_JPL_5647_S14_L6_Methylobacterium_72_69	95.30	0.86
ISS_JPL_5806_S15_L6_Acinetobacter_calcoaceticus_baumannii_complex_39_32	99.45	0.31
ISS_JPL_5806_S15_L6_Kocuria_71_254	99.34	0.00
ISS_JPL_5810_S18_L6_Paenibacillus_polymyxa_45_47	99.85	0.00
ISS_JPL_5810_S18_L6_Pantoea_sp._3.5.1_56_483	100.00	0.12
ISS_JPL_5813_S20_L6_Methylobacterium_extorquens_68_14	96.96	0.35
ISS_JPL_5813_S20_L6_Paenibacillus_polymyxa_45_354	99.85	0.00
ISS_JPL_5813_S20_L6_Staphylococcus_aureus_33_16	96.91	0.84
ISS_JPL_5819_S24_L7_Firmicutes_35_387	97.35	6.33
ISS_JPL_5819_S24_L7_Methylobacterium_aquaticum_71_7	70.53	0.50
ISS_JPL_5820_S25_L7_Paenibacillus_polymyxa_46_18	99.58	0.12
ISS_JPL_5820_S25_L7_Pantoea_sp._3.5.1_56_460	100.00	0.12
ISS_JPL_5822_S26_L7_Firmicutes_33_32	98.89	0.76
ISS_JPL_5822_S26_L7_Firmicutes_47_9	93.57	3.08
ISS_JPL_5822_S26_L7_Methylobacterium_aquaticum_71_87	99.37	0.16
ISS_JPL_5822_S26_L7_Sphingomonas_sanguinis	99.42	0.44
ISS_JPL_IIIF1SWP_Pantoea_57_200	99.95	0.25
ISS_JPL_IIIF1SW_Pantoea_57_295	99.95	0.33

ISS_JPL_IIIF2SWP_Actinobacteria_60_116	98.27	0.23
ISS_JPL_IIIF2SWP_Firmicutes_33_639	90.00	0.00
ISS_JPL_IIIF2SW_Firmicutes_33_339	97.96	0.00
ISS_JPL_IIIF2SW_Propionibacteriales_60_28	72.82	1.32
ISS_JPL_IIIF3SW2_Enterobacterales_58_402	98.23	2.72
ISS_JPL_IIIF3SW2_Firmicutes_34_22	87.90	0.56
ISS_JPL_IIIF3SWP_Enterobacterales_58_479	100.00	0.46
ISS_JPL_IIIF3SWP_Firmicutes_34_51	93.80	0.83
ISS_JPL_IIIF4SWP_Pantoea_dispersa_58_459	100.00	0.38
ISS_JPL_IIIF4SW_Pantoea_dispersa_58_467	100.00	0.38
ISS_JPL_IIIF5SWP_Pantoea_57_584	100.00	0.33
ISS_JPL_IIIF5SW_Pantoea_57_456	100.00	0.25
ISS_JPL_IIIF7SWP_Enterobacterales_55_52	98.18	0.21
ISS_JPL_IIIF7SWP_Pantoea_57_325	99.80	0.33
ISS_JPL_IIIF7SW_Enterobacterales_55_58	99.03	0.20
ISS_JPL_IIIF7SW_Pantoea_57_294	100.00	0.33
ISS_JPL_IIIF8SWP_Pantoea_57_601	100.00	0.33
ISS_JPL_IIIF8SW_Pantoea_57_546	100.00	0.33

425
426
427 **Table S2 | Phylogenetic characterization of recovered genomes.** 16 ribosomal proteins
428 were used to phylogenetically place the genomes. The taxonomy of the genomes represents
429 the consensus taxonomy.
430

Genome	Phylogenetic Placement
ISS_JPL_2332_S1_L003_Corynebacterium_afermentans_66_84	Turicella_otitidis_AtCC_51513
ISS_JPL_5492_S3_L3_Azohydromonas_australica_70_64	Azihydromonas_australica_DSM_1124
ISS_JPL_5492_S3_L3_Caulobacter_vibrioides_69_50	Caulobacter_segnis_ATCC_21756
ISS_JPL_5501_S5_L3_Acidovorax_69_463	Acidovorax_avenae_subsp._citrulli_AAC00_1
ISS_JPL_5501_S5_L3_Caulobacter_vibrioides_69_106	Caulobacter_segnis_ATCC_21756
ISS_JPL_5501_S5_L3_Sphingomonas_65_22	Sphingomonas japonicum_UT26S_1
ISS_JPL_5629_S1_L5_Enterobacterales_58_141	Enterobacteriaceae
ISS_JPL_5629_S1_L5_Firmicutes_33_15	Staphylococcus_aureus_502A
ISS_JPL_5630_S2_L5_Enterobacterales_56_407	Enterobacteriaceae
ISS_JPL_5631_S3_L5_Methylobacterium_69_176	Methylobacterium_extorquens_PA1

ISS_JPL_5631_S3_L5_Methylobacterium_aquaticum_71_106	Methylobacterium
ISS_JPL_5632_S4_L5_Staphylococcus_33_30	Staphylococcus_aureus_502A
ISS_JPL_5633_S5_L5_Pantoea_sp._3.5.1_56_354	Pantoea_anantis_LMG_5342
ISS_JPL_5635_S6_L5_Methylobacterium_aquaticum_71_83	Methylobacterium
ISS_JPL_5640_S8_L5_Enterobacterales_58_121	Enterobacteriaceae
ISS_JPL_5644_S12_L6_Pantoea_sp._3.5.1_56_351	Pantoea_anantis_LMG_5342
ISS_JPL_5647_S14_L6_Methylobacterium_69_10	Methylobacterium_extorquens_PA1
ISS_JPL_5647_S14_L6_Methylobacterium_72_69	Methylobacterium
ISS_JPL_5806_S15_L6_Acinetobacter_calcoaceticus_baumannii_complex_39_32	Acinetobacter_baumannii_AB30
ISS_JPL_5806_S15_L6_Kocuria_71_254	Kocuria_rhizophila_DC2201
ISS_JPL_5810_S18_L6_Paenibacillus_polymyxa_45_47	Paenibacillus_lactis_154
ISS_JPL_5810_S18_L6_Pantoea_sp._3.5.1_56_483	Pantoea_anantis_LMG_5342
ISS_JPL_5813_S20_L6_Methylobacterium_extorquens_68_14	Methylobacterium_extorquens_PA1
ISS_JPL_5813_S20_L6_Paenibacillus_polymyxa_45_354	Paenibacillus_lactis_154
ISS_JPL_5813_S20_L6_Staphylococcus_aureus_33_16	Staphylococcus_aureus_502A
ISS_JPL_5819_S24_L7_Firmicutes_35_387	Bacillus_anthracis_52_G
ISS_JPL_5819_S24_L7_Methylobacterium_aquaticum_71_7	Methylobacterium
ISS_JPL_5820_S25_L7_Paenibacillus_polymyxa_46_18	Paenibacillus_lactis_154
ISS_JPL_5820_S25_L7_Pantoea_sp._3.5.1_56_460	Pantoea_anantis_LMG_5342
ISS_JPL_5822_S26_L7_Firmicutes_33_32	Staphylococcus_aureus_502A
ISS_JPL_5822_S26_L7_Firmicutes_47_9	Bacillus_anthracis_52_G
ISS_JPL_5822_S26_L7_Methylobacterium_aquaticum_71_87	Methylobacterium
ISS_JPL_5822_S26_L7_Sphingomonas_sanguinis	Sphingomonas japonicum_UT26S_1
ISS_JPL_IIIF1SWP_Pantoea_57_200	Pantoea_anantis_LMG_5342
ISS_JPL_IIIF1SW_Pantoea_57_295	Pantoea_anantis_LMG_5342
ISS_JPL_IIIF2SWP_Actinobacteria_60_116	Propionibacterium_acnes
ISS_JPL_IIIF2SWP_Firmicutes_33_639	Staphylococcus_aureus_502A
ISS_JPL_IIIF2SW_Firmicutes_33_339	Staphylococcus_aureus_502A
ISS_JPL_IIIF2SW_Propionibacteriales_60_28	Propionibacterium_acnes
ISS_JPL_IIIF3SW2_Enterobacterales_58_402	Propionibacterium_acnes
ISS_JPL_IIIF3SW2_Firmicutes_34_22	Staphylococcus_aureus_502A
ISS_JPL_IIIF3SWP_Enterobacterales_58_479	Enterobacteriaceae
ISS_JPL_IIIF3SWP_Firmicutes_34_51	Staphylococcus_aureus_502A
ISS_JPL_IIIF4SWP_Pantoea_dispersa_58_459	Pantoea_anantis_LMG_5342
ISS_JPL_IIIF4SW_Pantoea_dispersa_58_467	Pantoea_anantis_LMG_5342
ISS_JPL_IIIF5SWP_Pantoea_57_584	Pantoea_anantis_LMG_5342

ISS_JPL_IIIF5SW_Pantoea_57_456	Pantoea_anantis_LMG_5342
ISS_JPL_IIIF7SWP_Enterobacterales_55_52	Enterobacteriaceae
ISS_JPL_IIIF7SWP_Pantoea_57_325	Pantoea_anantis_LMG_5342
ISS_JPL_IIIF7SW_Enterobacterales_55_58	Enterobacteriaceae
ISS_JPL_IIIF7SW_Pantoea_57_294	Pantoea_anantis_LMG_5342
ISS_JPL_IIIF8SWP_Pantoea_57_601	Pantoea_anantis_LMG_5342
ISS_JPL_IIIF8SW_Pantoea_57_546	Pantoea_anantis_LMG_5342

431
432
433 **Table S3 | Sample accessions and complexity based on the *rpS3* marker gene**.
434 Simulated low, medium and high complexity assembly and read datasets from the 1st CAMI
435 challenge were downloaded from https://data.cami-challenge.org/participate. Raw reads from
436 ISS flights I and II can be downloaded from the GeneLabs website (https://genelab-
437 data.ndc.nasa.gov/genelab/accession/GLDS-66/).
438
439

Sample	Read Accession	# rpS3 genes	Complexity category
CAMI_low	See table caption	30	Low
CAMI_medium_S1	See table caption	98	Medium
CAMI_high_S1	See table caption	404	High
Tomsk	SRR7102746	46	Low-Medium
SulCav_AS07-7	SRR1559028	151	Medium-High
ISS_JPL_2332_S1_L003	See table caption	9	Low
ISS_JPL_2333_S2_L003	See table caption	1	Low
ISS_JPL_5492_S3_L003	See table caption	7	Low
ISS_JPL_5501_S5_L003	See table caption	10	Low
ISS_JPL_5629_S1_L005	See table caption	4	Low
ISS_JPL_5630_S2_L005	See table caption	1	Low
ISS_JPL_5631_S3_L005	See table caption	2	Low
ISS_JPL_5632_S4_L005	See table caption	3	Low
ISS_JPL_5633_S5_L005	See table caption	1	Low
ISS_JPL_5635_S6_L005	See table caption	4	Low
ISS_JPL_5640_S8_L005	See table caption	4	Low
ISS_JPL_5644_S12_L006	See table caption	1	Low
ISS_JPL_5647_S14_L006	See table caption	2	Low
ISS_JPL_5806_S15_L006	See table caption	5	Low
ISS_JPL_5808_S16_L006	See table caption	5	Low
ISS_JPL_5810_S18_L006	See table caption	2	Low
ISS_JPL_5813_S20_L006	See table caption	3	Low
ISS_JPL_5818_S23_L007	See table caption	2	Low
ISS_JPL_5819_S24_L007	See table caption	5	Low
ISS_JPL_5820_S25_L007	See table caption	2	Low

ISS_JPL_5822_S26_L007	See table caption	4	Low
ISS_JPL_IIIF1SWP	SRX3808505	1	Low
ISS_JPL_IIIF1SW	SRX3808512	1	Low
ISS_JPL_IIIF2SWP	SRX3808504	3	Low
ISS_JPL_IIIF2SW	SRX3808511	2	Low
ISS_JPL_IIIF3SWP	SRX3808503	2	Low
ISS_JPL_IIIF3SW	SRX3808514	2	Low
ISS_JPL_IIIF4SWP	SRX3808508	1	Low
ISS_JPL_IIIF4SW	SRX3808513	1	Low
ISS_JPL_IIIF5SWP	SRX3808507	1	Low
ISS_JPL_IIIF5SW	SRX3808510	1	Low
ISS_JPL_IIIF7SWP	SRX3808529	1	Low
ISS_JPL_IIIF7SW	SRX3808509	1	Low
ISS_JPL_IIIF8SWP	SRX3808530	1	Low
ISS_JPL_IIIF8SW	SRX3808535	1	Low

440
441
442 **Table S4 | ISS metagenome assembly statistics.** Assembly statistics for SulCav AS07-7
443 and Tomsk metagenomes have been previously reported [30].
444

Sample	#Gbp reads after QC	#Mbp scaffolds	#Mbp scaffolds > 1Kbp length	N50 scaffolds > 1Kbp length
ISS_JPL_2332_S1_L003	9.5	859.2	54.7	1335
ISS_JPL_2333_S2_L003	0.3	4.3	0.4	1344
ISS_JPL_5492_S3_L003	6.7	87.6	65.9	10284
ISS_JPL_5501_S5_L003	6.5	84.5	62.7	7814
ISS_JPL_5629_S1_L005	3.6	127.2	76.4	38542
ISS_JPL_5630_S2_L005	2.1	13.7	10.2	19268
ISS_JPL_5631_S3_L005	3.6	68.5	22.5	51306
ISS_JPL_5632_S4_L005	1.9	82.4	40.4	38085
ISS_JPL_5633_S5_L005	2.5	57.6	30.1	1959
ISS_JPL_5635_S6_L005	2.6	112.0	84.5	24153
ISS_JPL_5640_S8_L005	2.7	88.5	58.4	29537
ISS_JPL_5644_S12_L006	2.6	62.6	40.6	3170
ISS_JPL_5647_S14_L006	3.5	77.5	64.9	9487
ISS_JPL_5806_S15_L006	4.1	71.0	66.7	31164
ISS_JPL_5808_S16_L006	3.4	128.2	55.8	53142
ISS_JPL_5810_S18_L006	3.2	11.8	10.8	179991
ISS_JPL_5813_S20_L006	2.6	53.6	16.1	17216
ISS_JPL_5818_S23_L007	3.4	54.2	52.9	86659

ISS_JPL_5819_S24_L007	2.6	64.8	40.0	3059
ISS_JPL_5820_S25_L007	2.8	11.4	10.8	62007
ISS_JPL_5822_S26_L007	2.1	29.0	18.9	41732
ISS_JPL_IIIF1SWP	1.0	7.3	5.4	132556
ISS_JPL_IIIF1SW	1.5	9.1	6.1	225749
ISS_JPL_IIIF2SWP	4.2	53.7	37.0	3007
ISS_JPL_IIIF2SW	2.1	36.0	10.2	1910
ISS_JPL_IIIF3SWP	2.7	14.4	7.5	168978
ISS_JPL_IIIF3SW	2.4	14.1	7.4	168987
ISS_JPL_IIIF4SWP	2.3	5.1	5.0	552627
ISS_JPL_IIIF4SW	2.4	5.8	5.1	405632
ISS_JPL_IIIF5SWP	2.9	5.1	4.9	205914
ISS_JPL_IIIF5SW	2.4	5.5	4.9	375929
ISS_JPL_IIIF7SWP	2.0	10.5	10.0	61535
ISS_JPL_IIIF7SW	1.8	11.4	10.0	59237
ISS_JPL_IIIF8SWP	3.0	6.2	5.0	248062
ISS_JPL_IIIF8SW	2.8	7.0	5.0	274906

445
446
447 **Additional Supplementary Files**
448
449
450 **File S1 | Phylogenetic tree for placement of ISS genomes based on 16 ribosomal**
451 **proteins.**
452
453 FileS1_ISS_PhyloTree.tree