Interleukin-19 alleviates experimental autoimmune encephalomyelitis by attenuating antigen-presenting cell activation

Hiroshi Horiuchi[1†], Bijay Parajuli[1†], Hiroyasu Komiya[2†], Yuki Ogawa[2], Jin Shijie[1],

Keita Takahashi[2], Yasu-Taka Azuma[3], Fumiaki Tanaka[2], Akio Suzumura[1],

and Hideyuki Takeuchi[1, 2*]

[1]Department of Neuroimmunology, Research Institute of Environmental Medicine, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, 464-8601, Japan.

[2]Department of Neurology and Stroke Medicine, Yokohama City University Graduate School of Medicine, 3-9 Fukuura, Kanazawa-ku, Yokohama 236-0004, Japan.

[3]Laboratory of Veterinary Pharmacology, Division of Veterinary Science, Osaka Prefecture University Graduate School of Life and Environmental Science, Izumisano, Osaka 598-8531, Japan.

[†]These authors contributed equally to this work.

[*]Corresponding Author: Hideyuki Takeuchi, M.D., Ph.D., Department of Neurology and Stroke Medicine, Yokohama City University Graduate School of Medicine, 3-9 Fukuura, Kanazawa-ku, Yokohama 236-0004, Japan.

Phone: +81-45-787-2725, FAX: +81-45-788-6041, E-mail: htake@yokohama-cu.ac.jp

Abstract

Interleukin-19 (IL-19) acts as an anti-inflammatory cytokine in various inflammatory diseases. Multiple sclerosis (MS) is a major neuroinflammatory disease in the central nervous system, but it remains uncertain how IL-19 contributes to MS pathogenesis. Here, we demonstrate that IL-19 deficiency aggravates experimental autoimmune encephalomyelitis (EAE), a mouse model of MS, by promoting IL-17–producing helper T cell (Th17 cell) infiltration into the central nervous system. In addition, IL-19–deficient splenic macrophages expressed elevated levels of major histocompatibility complex class II, co-stimulatory molecules, and Th17 cell differentiation–associated cytokines such as IL-1β, IL-6, IL-23, TGF-β1, and TNF-α. These observations indicated that IL-19 plays a critical role in suppression of MS pathogenesis by inhibiting macrophage antigen presentation, Th17 cell expansion, and subsequent inflammatory responses. Furthermore, treatment with IL-19 significantly abrogated EAE. Our data suggest that IL-19 could provide significant therapeutic benefits in patients with MS.

Introduction

Multiple sclerosis (MS) and experimental autoimmune encephalomyelitis (EAE), a mouse model of MS, are major autoimmune demyelinating diseases of the central nervous system (CNS)[1,2]. Various types of immune cells and soluble mediators contribute to the complex mechanisms underlying the onset and progression of both MS and EAE, and recent studies have shown that type 1 helper T (Th1) cells and interleukin-17–producing helper T (Th17) cells play pivotal roles in their pathogenesis[3, 4, 5]. In these diseases, autoreactive Th17 cells primed in the lymph nodes infiltrate the CNS and activate microglia/macrophages that induce inflammatory demyelination and subsequent neuronal damage, resulting in a wide range of clinical features, including sensory and motor paralysis, blindness, pain, incontinence, and dementia[1, 2].

Interleukin-19 (IL-19) is an IL-10 family cytokine that is homologous and highly similar to IL-20 and IL-24[6, 7]. IL-19 binds to the heterodimeric receptor consisting of IL-20Rα and IL-20Rβ, and its downstream signaling is mediated by STAT3 phosphorylation[8, 9]. IL-19 is mainly produced by activated macrophages and microglia[10, 11, 12, 13]. Recent studies showed that IL-19 exerts anti-inflammatory effects on macrophages by inhibiting inflammatory cytokine production, downregulating antigen-presenting capacity, and enhancing M2 phenotype polarization, which promotes type 2 helper T (Th2) cell differentiation and suppresses Th1 and Th17 cell differentiation[11, 12, 14, 15, 16, 17]. In fact, IL-19 plays a critical role in development of various autoimmune diseases, including asthma[18, 19], psoriasis[20, 21, 22], inflammatory bowel disease[11, 23], rheumatoid arthritis[24], and Type I diabetes[25]. However, it remains to be elucidated how IL-19 contributes to MS

pathogenesis.

Here, we examined the pathological role of IL-19 in EAE using IL-19–deficient

(IL-19$^{-/-}$) mice. IL-19 deficiency markedly exacerbated EAE, and treatment with IL-19

effectively suppressed EAE accompanied by inhibiting macrophage antigen presentation

and subsequent expansion of Th17 cells. Our findings suggest that IL-19 may provide

significant therapeutic benefits for treating MS.

Results

IL-19 deficiency exacerbates EAE

We generated myelin oligodendrocyte glycoprotein (MOG)-induced EAE in C57BL/6J

wild-type (WT) and IL-19–deficient (IL-19$^{-/-}$) mice. IL-19$^{-/-}$ mice exhibited earlier disease

onset and more severe symptoms than WT mice (Fig. 1A). Histological analysis of the

lumbar spinal cords revealed more inflammatory cell infiltration in IL-19$^{-/-}$ mice than in

WT mice (Fig. 1B). Flow cytometric analysis also disclosed that IL-19$^{-/-}$ EAE mice had

more CNS-infiltrating cells than WT EAE mice (Fig. 1C). We then chronologically

evaluated IL-19 expression levels in the spleen and lumbar spinal cord of WT mice at

pre-immunization, disease onset, and disease peak. Splenic IL-19 mRNA expression was

upregulated at disease onset, but was strongly suppressed at the disease peak (Fig. 1D).

By contrast, upregulation of IL-19 mRNA in the lumbar spinal cord was observed at

disease peak (Fig. 1E). These results suggest that endogenous IL-19 serves as a negative

regulator of EAE pathogenesis at both the induction and effector phases.

IL-19 deficiency increases Th17 cell infiltration into the CNS

Because EAE is a Th1 and Th17 cell–mediated autoimmune disease, we next assessed

whether IL-19 deficiency would increase CNS infiltration of Th1 and Th17 cells. Flow

cytometric analysis revealed that at disease peak, IL-19$^{-/-}$ mice exhibited more infiltration

of Th17 cells in the spinal cord than WT mice (Fig. 2A, B). By contrast, no difference

was observed in Th1 cell infiltration between IL-19$^{-/-}$ and WT mice (Fig. 2C). These

results indicate that IL-19 deficiency mediates elevated CNS infiltration by Th17 cells,

but not Th1 cells.

IL-19 deficiency expands Th17 cell population

To determine whether IL-19 contributes to the expansion of Th17 cells during the

induction phase of EAE, we assessed the antigen-specific expansion of Th17 cells *ex vivo*.

Splenic CD4$^+$ T cells isolated from WT and IL-19$^{-/-}$ mice at MOG-EAE onset were

stimulated with MOG peptide for 3 days *in vitro*. IL-19 deficiency significantly

upregulated *Il17a* mRNA and downregulated *Foxp3* mRNA, but it did not affect the level

of *Ifng* mRNA (Fig. 3A). Flow cytometric analysis also revealed that IL-19 deficiency

expanded the Th17 cell population (Fig. 3B, C). These results indicate that IL-19

deficiency mediates expansion of Th17 cells in the peripheral lymphoid tissues.

IL-19 deficiency skews cytokine expression profiles toward Th17 cell expansion in

macrophages

We then examined how IL-19 deficiency expands Th17 cells in the induction phase of

EAE. First, we evaluated the mRNA expression level of IL-19 receptor (heterodimer of

IL-20Rα and IL-20Rβ subunits) in the splenic immune cells such as macrophage,

dendritic cell (DC), and CD4$^+$ helper T cell. We found that both the IL-20Rα and IL-20Rβ

subunits were more highly expressed in CD11b$^+$ macrophages and CD4$^+$ helper T cells

than in CD11c$^+$ DCs (Fig. S1). These results suggest that IL-19 mainly affects

macrophages and CD4$^+$ helper T cells.

Next, we assessed whether IL-19 directly differentiates naïve CD4$^+$ T cells into

Th17 cells. Naïve T cells were polarized using immobilized CD3 and CD28 antibodies in

the presence of IL-6 and transforming growth factor β1 (TGF-β1), with or without IL-19.

Quantitative PCR (qPCR) and flow cytometry revealed that IL-19 did not alter the

differentiation of naïve T cells into Th17 cells (Fig. S2).

Because antigen-presenting cells (APCs) are crucial for differentiation of naïve

T cells into effector T cells, we evaluated the expression levels of cytokines required for

Th17 cell expansion in splenic macrophages and DCs. Interestingly, IL-19$^{-/-}$ macrophages

exhibited a significant increase in mRNA levels of the genes encoding IL-1β, IL-6, TGF-

β, IL-12 p40, IL-23 p19, and tumor necrosis factor α (TNF-α), which play pivotal roles

in Th17 cell differentiation and expansion (Fig. 4). Although a previous study showed

that IL-19 increases IL-10 expression [26], our data showed that IL-19 deficiency did not

alter the *Il10* mRNA level in macrophages (Fig. 4). By contrast, IL-19$^{-/-}$ DCs did not

exhibit a significant alteration in the expression levels of these cytokines (Fig. S3). These

findings indicated that IL-19 deficiency skews the cytokine expression profiles toward

Th17 cell differentiation and expansion in macrophages. Conversely, our data suggested

that IL-19 suppresses Th17-skewed condition by activating macrophages.

IL-19 deficiency promotes MHC class II expression in macrophages

To determine whether IL-19 signaling contributes to antigen presentation by macrophages,

we assessed the expression of major histocompatibility complex (MHC) class II (H2-Ab)

and co-stimulatory molecules (CD80 and CD86) in splenic CD11b$^+$ macrophages from

WT and IL-19$^{-/-}$ EAE mice. (Fig. 5A). IL-19 deficiency significantly enhanced expression

of the gene encoding MHC class II, whereas the genes encoding co-stimulatory molecules

CD80 and CD86 were not affected (Fig. 5A). Flow cytometric data corroborated the

enhanced presentation of MHC class II in IL-19$^{-/-}$ splenic macrophages (Fig. 5B). These

observations suggested that IL-19 also suppresses the antigen-presenting activity of

macrophages.

Treatment with IL-19 abrogates EAE

To determine whether exogenous IL-19 abolishes the effect of IL-19 deficiency in EAE,

we treated IL-19$^{-/-}$ EAE mice with recombinant mouse IL-19 protein (20 ng/g of body

weight) by intraperitoneal injection every other day, starting on day 2 post-immunization.

As shown in Figure 6A, administration of IL-19 to IL-19$^{-/-}$ mice abolished the

exacerbation of EAE (Fig. 6A, IL-19$^{-/-}$ + IL-19). We then investigated the therapeutic

effect of IL-19 on EAE. When we treated WT EAE mice with recombinant mouse IL-19

protein in the same manner, we found that IL-19 treatment almost completely inhibited

EAE (Fig. 6B). These results indicated that IL-19 represents a potential target for MS

therapy.

Discussion

In this study, we demonstrated that endogenous IL-19 negatively regulates development of EAE by inhibiting macrophage activation, and that IL-19 treatment effectively abrogates EAE. As shown in Fig. 1D and 1E, endogenous *Il19* mRNA expression was upregulated at EAE onset and downregulated at EAE peak in the spleen, whereas it was elevated at EAE peak in the CNS. We have previously identified IL-19 as a negative-feedback regulator to limit proinflammatory response of macrophages and microglia in autocrine/paracrine manners[11, 12]. From this point of view, these data imply that endogenous IL-19 increases to suppress inflammation accompanied by macrophage/microglia activation as disease progresses from the periphery to CNS, although it is insufficient to halt EAE progression.

Th17 cell infiltration in the CNS is considered critical for the development of EAE[5, 27]. Our data revealed that IL-19 deficiency increased CNS infiltration of Th17 cells, but not Th1 cells (Fig. 2). In addition, IL-19 deficiency enhanced the peripheral expansion of Th17 cells in the induction phase of EAE (Fig. 3). Conversely, our data indicated that IL-19 negatively regulates Th17 cell differentiation and expansion, which is critical for EAE development.

However, IL-19 did not directly affect differentiation of Th17 cells from naïve T cells (Fig. S2). APCs such as macrophages, microglia, and DC are also essential for effector T cell differentiation and expansion in EAE[28]. Specifically, IL-1β, IL-6, IL-23, TGF-β1, and TNF-α released by APCs play pivotal roles in Th17 cell differentiation and expansion[5, 29, 30, 31 32, 33]. In this study, we revealed that IL-19 deficiency significantly

upregulated these Th17 cell differentiation–associated cytokines in macrophages, but not

in DCs (Fig. 4 and Fig. S3). These phenomena were correlated with the expression level

of IL-19 receptor (IL-20Rα and IL-20Rβ heterodimer), which was highly expressed in

macrophages, but not in DCs or helper T cells (Fig. S1). Thus, IL-19 might negatively

regulate Th17 cell differentiation and expansion through inhibition of cytokine release

from macrophages.

Moreover, activated APCs highly express MHC class II and co-stimulatory

molecules (CD80 and CD86) on the cell surface and acquire the ability to prime T cells[34].

As shown in Fig. 5, IL-19 deficiency further enhanced MHC class II expression in

macrophages. Taken together, our findings suggest that IL-19 suppresses Th17 cell

differentiation and expansion by suppressing cytokine production and antigen

presentation in macrophages. Interestingly, a previous study reported that IL-17A induces

IL-19 production[19]. Thus, IL-19 also might serve as a negative regulator of further Th17

cell polarization.

In addition, activated macrophages and microglia directly contribute to

neuroinflammation-induced demyelination by releasing proinflammatory cytokines such

as IL-1β, IL-6, and TNF-α[28, 35, 36, 37]. In our study, IL-19 deficiency increased the levels

of these proinflammatory cytokines in macrophages and exacerbated EAE until the late

phase of the disease. We previously revealed that IL-19 secreted from activated

macrophages and microglia suppressed their proinflammatory responses in an

autocrine/paracrine manner[11, 12]. Therefore, IL-19 might suppress development of EAE

by dual inhibition of both autoreactive Th17 cell expansion and macrophage/microglia-

mediated CNS neuroinflammation.

Although the IL-19 signaling pathway has not been fully elucidated, IL-19

mediates its downstream signaling at least by STAT3 activation[8, 9, 12, 38]. However, it

remains controversial whether STAT3 activation is beneficial or harmful with regard to

autoimmune-mediated neuroinflammation. Previous studies also reported that STAT3

activation in myeloid cells (including macrophages and microglia) exacerbates EAE[39, 40].

By contrast, STAT3 ablation worsens neuroinflammation in mice with spinal cord injury[41],

and STAT3 activation alleviates cuprizone-induced CNS demyelination[42]. This

discordancy may depend on spatiotemporally specific activation of STAT3[43]. Indeed, in

contrast to macrophages, IL-19 deficiency did not affect activation of DCs. Recent

clinical trials of JAK/STAT inhibitors for autoimmune diseases have revealed a

complicated signal network of cytokine/STAT axes in multiple cell types[44]. Further

studies are needed to elucidate the precise IL-19 signaling pathway in each cell type.

In summary, we revealed that IL-19 deficiency exacerbated EAE by upregulating

Th17 cell differentiation–associated cytokines and enhancing antigen presentation in

macrophages, followed by Th17 cell expansion and infiltration in the CNS. We also

demonstrated that IL-19 administration potently prevented development of EAE.

Therefore, enhancement of IL-19 signaling represents a promising therapeutic strategy

against MS and other Th17-mediated autoimmune diseases.

Material and Methods

Reagents

MOG peptide 35–55 (MOG$_{35-55}$; MEVGWYRSPFSRVVHLYRNGK) was synthesized and purified by Operon Biotechnologies (Tokyo, Japan). Incomplete Freund's adjuvant was obtained from Sigma-Aldrich (St. Louis, MO, USA). Heat-killed *Mycobacterium tuberculosis* H37Ra was obtained from Difco (Detroit, MI, USA), and pertussis toxin was obtained from List Biological Laboratories (Campbell, CA, USA). Recombinant mouse IL-6, IL-19, and TGF-β1 were obtained from R&D Systems (Minneapolis, MN, USA).

Animals

All animal experiments were conducted under protocols approved by the Animal Experiment Committee of Nagoya University (approved numbers: 15017 and 15018) and Yokohama City University (approved number: F-A-19-036). C57BL/6J (B6) mice were purchased from Japan SLC (Hamamatsu, Japan). IL-19$^{-/-}$ mice (B6 background) [11, 12] were obtained from Regeneron Pharmaceuticals (Tarrytown, NY, USA).

EAE induction and treatment studies

MOG-EAE was induced as previously described [45, 46, 47]. In brief, 8-week-old female mice were immunized subcutaneously at the base of the tail with 0.2 ml of emulsion containing 200 μg MOG$_{35-55}$ in saline, combined with an equal volume of complete Freund's adjuvant containing 300 mg heat-killed *Mycobacterium tuberculosis* H37Ra. The mice

were intraperitoneally injected with 200 ng pertussis toxin on days 0 and 2 post-

immunization. To investigate the effect of IL-19, EAE mice were treated with mouse

recombinant IL-19 protein (20 ng/g of body weight) by intraperitoneal injection every

other day starting on day 2 post-immunization, according to a modification of a

previously reported method [48, 49]. The mice were assessed daily for clinical signs of EAE,

according to the following grading system: 0, normal; 1 - limp tail or mild hind limb

weakness; 2 - moderate hind limb weakness or mild ataxia; 3 - moderate to severe hind

limb weakness; 4 - severe hind limb weakness, mild forelimb weakness or moderate

ataxia; 5 - paraplegia with moderate forelimb weakness; and 6 - paraplegia with severe

forelimb weakness, severe ataxia, or moribundity.

Isolation of cells from spleen and lumbar spinal cord

Mononuclear cells were collected from the spleen and lumbar spinal cord as described

previously [45, 46, 47]. $CD4^+$, $CD11b^+$, and $CD11c^+$ cells were isolated using the MACS

system (Miltenyi Biotec, Bergisch Gladbach, Germany). Helper T cell differentiation was

induced as described previously [5, 32, 47]. For flow cytometric analysis, cells were stained

with PerCP/Cy5.5 or BV421-conjugated anti–mouse CD4 rat monoclonal antibody

(RM4-5; BD Biosciences, Franklin Lakes, NJ, USA). The cells were then fixed and

permeabilized with Cytofix/Cytoperm reagent (BD Biosciences) and stained with PE-

conjugated anti–mouse IFN-γ rat monoclonal antibody (XMG1.2; BD Biosciences) and

APC- or PE-conjugated anti–mouse IL-17A rat monoclonal antibody (TC11-18H10; BD

Biosciences). The samples were analyzed using a FACS Aria III system (BD Biosciences)

and the FlowJo software (FlowJo, Ashland, OR, USA).

Histological analysis

Histological analysis was performed as previously described [45]. Mice with peak EAE

were anesthetized and perfused transcardially with 4% paraformaldehyde in 0.1 M PBS.

Lumbosacral spinal cords were immediately removed, postfixed in 4% paraformaldehyde,

and embedded in paraffin. Eight-micron-thick sections were stained with hematoxylin

and eosin. Stained sections were analyzed on a NanoZoomer 2.0-RS slide scanner

(Hamamatsu Photonics, Hamamatsu, Japan).

RNA extraction and reverse-transcription polymerase chain reactions (RT-PCRs)

We evaluated the expression levels of proinflammatory factors in the lumbar spinal cords

and spleens by qPCR as described previously [12, 45]. In brief, lumbar spinal cords and

spleens were collected from EAE mice at pre-immunization, EAE onset, and EAE peak

(approximately on days 0, 10, 16 post-immunization, respectively). Total RNA was

isolated with an RNeasy Mini Kit (Qiagen, Valencia, CA, USA) and reverse transcribed

with SuperScript III (Life Technologies, Carlsbad, CA, USA). Expression levels of

mRNAs were evaluated by qPCR using SYBR Select Master Mix (Applied Biosystems,

Foster City, CA, USA) on a Rotor-Gene Q (Qiagen) or LightCycler 96 (Roche). Mouse

gene–specific primers were obtained from Life Technologies (Table 1). Gene-expression

values were determined using the $\Delta\Delta C_T$ method. Levels of mRNAs of interest were

standardized   to   the   geometric   mean   of   the   level   of   hypoxanthine

phosphoribosyltransferase 1 (*Hprt1*). Assays were carried out in three independent trials.

Statistical analysis

Statistical significance was analyzed using Student's t-test or one-way analysis of

variance (ANOVA) followed by post hoc Tukey's test in GraphPad Prism version 8

(GraphPad Software, La Jolla, CA, USA).

Data availability

The datasets generated and analyzed during the study are available from the

corresponding author upon reasonable request.

References

1.   Hemmer B, Archelos JJ, Hartung HP. New concepts in the immunopathogenesis of multiple sclerosis. *Nat Rev Neurosci* **3**, 291-301 (2002).

2.   Filippi M*, et al.* Multiple sclerosis. *Nat Rev Dis Primers* **4**, 43 (2018).

3.   Waisman A, Hauptmann J, Regen T. The role of IL-17 in CNS diseases. *Acta Neuropathol* **129**, 625-637 (2015).

4.   Takeuchi H. Midkine and multiple sclerosis. *Brit J Pharmacol* **171**, 931-935 (2014).

5.   Bettelli E*, et al.* Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells. *Nature* **441**, 235-238 (2006).

6.   Gallagher G*, et al.* Cloning, expression and initial characterisation of interleukin-19 (IL-19), a novel homologue of human interleukin-10 (IL-10). *Genes and Immunity* **1**, 442-450 (2000).

7.   Gallagher G*, et al.* Human interleukin-19 and its receptor: a potential role in the induction of Th2 responses. *Int Immunopharmacol* **4**, 615-626 (2004).

8.   Nagalakshmi ML, Murphy E, McClanahan T, de Waal Malefyt R. Expression patterns of IL-10 ligand and receptor gene families provide leads for biological characterization. *Int Immunopharmacol* **4**, 577-592 (2004).

9.   Dumoutier L, Leemans C, Lejeune D, Kotenko SV, Renauld JC. Cutting edge: STAT activation by IL-19, IL-20 and mda-7 through IL-20 receptor complexes of two types. *Journal of Immunology* **167**, 3545-3549 (2001).

10. Wolk K*, et al.* Maturing dendritic cells are an important source of IL-29 and IL-20 that may cooperatively increase the innate immunity of keratinocytes. *J Leukoc Biol* **83**, 1181-1193 (2008).

11. Azuma YT*, et al.* Interleukin-19 protects mice from innate-mediated colonic inflammation. *Inflamm Bowel Dis* **16**, 1017-1028 (2010).

12. Horiuchi H*, et al.* Interleukin-19 acts as a negative autocrine regulator of activated microglia. *PLoS One* **10**, e0118640 (2015).

13. Cooley ID, Chauhan VS, Donneyz MA, Marriott I. Astrocytes produce IL-19 in response to bacterial challenge and are sensitive to the immunosuppressive effects of this IL-10 family member. *Glia* **62**, 818-828 (2014).

337 14. Myles IA, *et al.* Signaling via the IL-20 receptor inhibits cutaneous production of
338   IL-1β and IL-17A to promote infection with methicillin-resistant Staphylococcus
339   aureus. *Nat Immunol* **14**, 804-811 (2013).
340
341 15. Ellison S, *et al.* IL-19 reduces ligation-mediated neointimal hyperplasia by
342   reducing vascular smooth muscle cell activation. *Am J Pathol* **184**, 2134-2143
343   (2014).
344
345 16. Wahl C, *et al.* IL-20 receptor 2 signaling down-regulates antigen-specific T cell
346   responses. *J Immunol* **182**, 802-810 (2009).
347
348 17. Oral HB, *et al.* Regulation of T cells and cytokines by the interleukin-10 (IL-10)-
349   family cytokines IL-19, IL-20, IL-22, IL-24 and IL-26. *Eur J Immunol* **36**, 380-
350   388 (2006).
351
352 18. Liao SC, *et al.* IL-19 induced Th2 cytokines and was up-regulated in asthma
353   patients. *J Immunol* **173**, 6712-6718 (2004).
354
355 19. Huang F, *et al.* Potentiation of IL-19 expression in airway epithelia by IL-17A and
356   IL-4/IL-13: Important implications in asthma. *Journal of Allergy and Clinical
357   Immunology* **121**, 1415-1421000 (2008).
358
359 20. Kunz S, *et al.* Interleukin (IL)-19, IL-20 and IL-24 are produced by and act on
360   keratinocytes and are distinct from classical ILs. *Exp Dermatol* **15**, 991-1004
361   (2006).
362
363 21. Ghoreschi K, *et al.* Interleukin-4 therapy of psoriasis induces Th2 responses and
364   improves human autoimmune disease. *Nat Med* **9**, 40-46 (2003).
365
366 22. Chan JR, *et al.* IL-23 stimulates epidermal hyperplasia via TNF and IL-20R2-
367   dependent mechanisms with implications for psoriasis pathogenesis. *J Exp Med*
368   **203**, 2577-2587 (2006).
369
370 23. Fonseca-Camarillo G, Furuzawa-Carballeda J, Granados J, Yamamoto-Furusho
371   JK. Expression of interleukin (IL)-19 and IL-24 in inflammatory bowel disease
372   patients: a cross-sectional study. *Clin Exp Immunol* **177**, 64-75 (2014).
373
374 24. Sakurai N, *et al.* Expression of IL-19 and its receptors in RA: potential role for
375   synovial hyperplasia formation. *Rheumatology (Oxford)* **47**, 815-820 (2008).
376
377 25. Barrett JC, *et al.* Genome-wide association study and meta-analysis find that over
378   40 loci affect risk of type 1 diabetes. *Nat Genet* **41**, 703-707 (2009).
379
380 26. Jordan WJ, *et al.* Human IL-19 regulates immunity through auto-induction of IL-
381   19 and production of IL-10. *Eur J Immunol* **35**, 1576-1582 (2005).

383 27. Xu Y*, et al.* Irgm1 is required for the inflammatory function of M1 macrophage
384   in early experimental autoimmune encephalomyelitis. *J Leukoc Biol* **101**, 507-517
385   (2017).
386

387 28. Yoshida Y*, et al.* The transcription factor IRF8 activates integrin-mediated TGF-
388   β signaling and promotes neuroinflammation. *Immunity* **40**, 187-198 (2014).
389

390 29. Ghoreschi K*, et al.* Generation of pathogenic T(H)17 cells in the absence of TGF-
391   beta signalling. *Nature* **467**, 967-971 (2010).
392

393 30. Ivanov, II*, et al.* The orphan nuclear receptor RORgammat directs the
394   differentiation program of proinflammatory IL-17+ T helper cells. *Cell* **126**, 1121-
395   1133 (2006).
396

397 31. Mangan PR*, et al.* Transforming growth factor-beta induces development of the
398   T(H)17 lineage. *Nature* **441**, 231-234 (2006).
399

400 32. Chung Y*, et al.* Critical regulation of early Th17 cell differentiation by interleukin-
401   1 signaling. *Immunity* **30**, 576-587 (2009).
402

403 33. Mufazalov IA*, et al.* IL-1 signaling is critical for expansion but not generation of
404   autoreactive GM-CSF+ Th17 cells. *The EMBO journal* **36**, 102-115 (2017).
405

406 34. Blum JS, Wearsch PA, Cresswell P. Pathways of antigen processing. *Annu Rev*
407   *Immunol* **31**, 443-473 (2013).
408

409 35. Becher B, Spath S, Goverman J. Cytokine networks in neuroinflammation. *Nat*
410   *Rev Immunol* **17**, 49-59 (2017).
411

412 36. Deng X, Sriram S. Role of microglia in multiple sclerosis. *Curr Neurol Neurosci*
413   *Rep* **5**, 239-244 (2005).
414

415 37. Luo C*, et al.* The role of microglia in multiple sclerosis. *Neuropsychiatr Dis Treat*
416   **13**, 1661-1667 (2017).
417

418 38. Uto-Konomi A*, et al.* Dysregulation of suppressor of cytokine signaling 3 in
419   keratinocytes causes skin inflammation mediated by interleukin-20 receptor-
420   related cytokines. *PLoS One* **7**, e40343 (2012).
421

422 39. Qin H*, et al.* Signal transducer and activator of transcription-3/suppressor of
423   cytokine signaling-3 (STAT3/SOCS3) axis in myeloid cells regulates
424   neuroinflammation. *Proc Natl Acad Sci U S A* **109**, 5004-5009 (2012).
425

426 40. Lu HC*, et al.* STAT3 signaling in myeloid cells promotes pathogenic myelin-

427   specific T cell differentiation and autoimmune demyelination. *Proc Natl Acad Sci*
428   *U S A* **117**, 5430-5441 (2020).
429

430 41.   Okada S*, et al.* Conditional ablation of Stat3 or Socs3 discloses a dual role for
431   reactive astrocytes after spinal cord injury. *Nat Med* **12**, 829-834 (2006).
432

433 42.   Emery B*, et al.* Suppressor of cytokine signaling 3 limits protection of leukemia
434   inhibitory factor receptor signaling against central demyelination. *Proc Natl Acad*
435   *Sci U S A* **103**, 7859-7864 (2006).
436

437 43.   Liu Y, Gibson SA, Benveniste EN, Qin H. Opportunities for Translation from the
438   Bench: Therapeutic Intervention of the JAK/STAT Pathway in
439   Neuroinflammatory Diseases. *Crit Rev Immunol* **35**, 505-527 (2015).
440

441 44.   Schwartz DM, Kanno Y, Villarino A, Ward M, Gadina M, O'Shea JJ. JAK
442   inhibition as a therapeutic strategy for immune and inflammatory diseases. *Nat*
443   *Rev Drug Discov* **16**, 843-862 (2017).
444

445 45.   Wang J*, et al.* Inhibition of midkine alleviates experimental autoimmune
446   encephalomyelitis through the expansion of regulatory T cell population. *Proc*
447   *Natl Acad Sci U S A* **105**, 3915-3920 (2008).
448

449 46.   Shimojima C*, et al.* Conditioned Medium from the Stem Cells of Human
450   Exfoliated Deciduous Teeth Ameliorates Experimental Autoimmune
451   Encephalomyelitis. *J Immunol* **196**, 4164-4171 (2016).
452

453 47.   Zhou Y*, et al.* IL-9 promotes Th17 cell migration into the central nervous system
454   via CC chemokine ligand-20 produced by astrocytes. *J Immunol* **186**, 4415-4421
455   (2011).
456

457 48.   Richards J, Gabunia K, Kelemen SE, Kako F, Choi ET, Autieri MV. Interleukin-
458   19 increases angiogenesis in ischemic hind limbs by direct effects on both
459   endothelial cells and macrophage polarization. *J Mol Cell Cardiol* **79**, 21-31
460   (2015).
461

462 49.   Guo J, Wang H, Li L, Yuan Y, Shi X, Hou S. Treatment with IL-19 improves
463   locomotor functional recovery after contusion trauma to the spinal cord. *Br J*
464   *Pharmacol* **175**, 2611-2621 (2018).
465
466
467

Acknowledgements

This work was supported by grants-in-aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan (H.T.); grants from the Ministry of Health, Labour and Welfare of Japan (A.S., H.T.); the Program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation (NIBIO) of Japan (A.S.); and a grant from the Naito Foundation (H.T.).

Author contributions

H.H., B.P., A.S., and H.T. designed the research; H.H., B.P., H.K., Y.O., J.S., K.T., Y.A., and H.T. performed the research; H.H., B.P., H.K., F.T., A.S., and H.T. analyzed the data; and H.H., B.P., and H.T. wrote the paper.

Competing interests

The authors declare no competing interests.

Figure Legends

Figure 1. IL-19 deficiency aggravates EAE.

(A) EAE clinical scores for WT (black) and IL-19$^{-/-}$ (red) mice. Data represent means ± SD (n = 10). ***, $p < 0.0001$. (B) Micrographs of hematoxylin/eosin staining of L5 lumbar spinal cords at the peak EAE of WT and IL-19$^{-/-}$ mice. The right panels show enlargements of the boxed areas in the left panels. Scale bars: 500 μm (left), 100 μm (right). (C) Flow cytometric analysis of cell infiltration in the CNS at the peak EAE (n = 5). Data represent means ± SD. **, $p < 0.01$. (D and E) chronological qPCR data for IL-19 mRNA expression level in the spleen (D) and the lumbar spinal cord (E) of WT EAE mice (n = 5). Pre, pre-immunization; onset, EAE onset; peak, EAE peak. Data represent means ± SD. **, $p < 0.01$.

Figure 2. IL-19 deficiency increases CNS infiltration of Th17 cells, but not Th1 cells.

(A) Representative flow-cytometric data for IL-17– and IFN-γ–producing CD4$^+$ T cells in the CNS at the peak EAE. (B) Percentage of IL-17–producing CD4$^+$ T cells. (C) Percentage of IFN-γ–producing CD4$^+$ T cells. Data represent means ± SD (n = 3). *, $p < 0.05$.

Figure 3. IL-19 deficiency expands the Th17 cell population.

(A) qPCR data for levels of mRNAs encoding IL-17A, IFN-γ, and FoxP3 in splenic CD4$^+$ T cells. (B) Representative flow cytometric data for IL-17–producing CD4$^+$ T cells in the

spleens of EAE mice. (C) Percentage of IL-17–producing CD4+ T cells in the spleens of

EAE mice. Data represent means ± SD (n = 3). *, $p < 0.05$.

Figure 4. IL-19 deficiency upregulates Th17 cell differentiation–associated cytokines

in macrophages.

(A) qPCR data for mRNAs encoding IL-1β, IL-6, TGF-β1, IL-12 p40, IL-23 p19, IL-10,

and TNF-α in splenic macrophages of EAE mice. Assessments were performed 7 days

after immunization. Data represent means ± SD. ***, $p < 0.0001$ (n = 6).

Figure 5. IL-19 deficiency enhances antigen-presenting activity in macrophages.

(A) qPCR data for mRNAs encoding MHC class II (H2-Ab), CD80, and CD86 in the

splenic macrophages on day 7 after immunization. Data represent means ± SD. *, $p <$

0.05 (n = 6). (B) Representative flow cytometric data for MHC class II (H2-Ab)

presentation in splenic macrophages.

Figure 6. Treatment with recombinant IL-19 alleviates EAE.

(A) EAE clinical score. WT + PBS (black): WT EAE mice treated with PBS; IL-19$^{-/-}$ +

PBS (red): IL-19$^{-/-}$ EAE mice treated with PBS; IL-19$^{-/-}$ + IL-19 (blue): IL-19$^{-/-}$ EAE mice

treated with IL-19. (B) EAE clinical score. WT + PBS (black), WT EAE mice treated

with PBS; WT + IL-19 (blue), WT EAE mice treated with IL-19. Data represent means ±

SD. *, $p < 0.05$ (n = 5).

Table 1. Primers for qPCR

Gene	Primer sequence
mouse *Il19* sense	CAACCTGCTGACATTCTACAGAG
mouse *Il19* antisense	CCTGACATCGCTCCAGAGATTT
mouse *Il17a* sense	TCATCTGTGTCTCTGATGCTGTTG
mouse *Il17a* antisense	TCGCTGCCTTCACTGT
mouse *Ifng* sense	TGGCATAGATGTGGAAGAAAAGAG
mouse *Ifng* antisense	TGCAGGATTTTCATGTCACCAT
mouse *Foxp3* sense	TTCATGCATCAGCTCTCCAC
mouse *Foxp3* antisense	CTGGACACCCATTCCAGACT
mouse *Il1b* sense	GAAATGCCACCTTTTGACAGTG
mouse *Il1b* antisense	TGGATGCTCTCATCAGGACAG
mouse *Il6* sense	TCTATACCACTTCACAAGTCGGA
mouse *Il6* antisense	GAATTGCCATTGCACAACTCTTT
mouse *Tgfb1* sense	CGAAGCGGACTACTATGCTAAAGA
mouse *Tgfb1* antisense	GTTTTCTCATAGATGGCGTTGTTG
mouse *Il10* sense	GAGAAGCATGGCCCAGAAATC
mouse *Il10* antisense	CGCATCCTGAGGGTCTTCA
mouse *Il12p40* sense	GGTGCAAAGAAACATGGACTTG
mouse *Il12p40* antisense	CACATGTCACTGCCCGAGAGT
mouse *Il23p19* sense	GCACCAGCGGGACATATGA
mouse *Il23p19* antisense	CCTTGTGGGTCACAACCATCT
mouse *Tnfa* sense	GACCCTCACACTCAGATCATCTTCT
mouse *Tnfa* antisense	CCACTTGGTGGTTTGCTACGA
mouse *Il20ra* sense	GGAAACTCAAGTCAGCCCAC
mouse *Il20ra* antisense	AGATGGACTTCTCGCCAGTT
mouse *Il20rb* sense	CCGAAATGCAACTGTCCTCAC
mouse *Il20rb* antisense	AATAACCAGATGCAGCCCATGT
mouse *Rorgt* sense	GCGACTGGAGGACCTTCTAC
mouse *Rorgt* antisense	TCCCACATTGACTTCCTCTG
mouse *H2ab1* sense	AGACGCCGAGTACTGGAACAGCCAGC
mouse *H2ab1* sense	CAGAGTGTTGTGGTGGTTGAGGGCCTC
mouse *Cd80* sense	CATCAAAGCTGACTTCTCTACCC
mouse *Cd80* antisense	GGGTTTTTCCCAGGTGAAGT
mouse *Cd86* sense	TCAGTGATCGCCAACTTCAG
mouse *Cd86* sense	GAAACTCTTGAGTGAAATTGAGAGG
mouse *Hprt1* sense	CAGTCAACGGGGACATAAA
mouse *Hprt1* antisense	GGGGCTGTACTGCTTAACCAG

Horiuchi et al, Figure. 1

Horiuchi et al, Figure. 2

Horiuchi et al, Figure. 3

Horiuchi et al, Figure. 4

A

B

Gated on CD11b

Horiuchi et al, Figure. 5

A

B

Horiuchi et al, Figure. 6