Global transcriptional regulation by cell-free supernatant of *Salmonella* Typhimurium peptide transporter mutant leads to inhibition of intra-species biofilm initiation

Kasturi Chandra[1], Prerana Muralidhara[2,‡], Sathisha Kamanna[3], Utpal S. Tatu[3], Dipshikha Chakravortty[1,4,#]

[1]Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore, India.

[2]Department of Molecular Biology, Yuvaraja's College, Mysore, Karnataka, India.

[3]Department of Biochemistry, Indian Institute of Science, Bangalore, India.

[4]Centre for BioSystems Science and Engineering, Indian Institute of Science, Bangalore, India.

[‡]Present address: Molecular Biology and Genetic Unit, Jawaharlal Nehru Centre for Advanced Scientific Research, Bangalore, India

 # Corresponding author

Dipshikha Chakravortty

Email: dipa@iisc.ac.in

Telephone: +91 80 22932842

Keywords: *Salmonella* Typhimurium, Biofilm, flagella, H-NS, EPS, yjiY, oxidative stress

Abstract

Salmonella is a genus of widely spread Gram negative, facultative anaerobic bacteria, which is known to cause ¼th of diarrheal morbidity and mortality globally. It causes typhoid fever and gastroenteritis by gaining access to the host gut through contaminated food and water. *Salmonella* utilizes its biofilm lifestyle to strongly resist antibiotics and persist in the host. Although biofilm removal or dispersal has been studied widely, the inhibition of the initiation of *Salmonella* biofilm remains elusive. This study was conducted to determine the anti-biofilm property of the cell-free supernatant obtained from a carbon-starvation inducible proline peptide transporter mutant (Δ*yjiY)* strain. Our study shows that *Salmonella* Δ*yjiY* culture supernatant primarily inhibits biofilm initiation by regulating biofilm-associated transcriptional network. This work demonstrates that highly abundant proteases such as HslV and GrpE cleave the protein aggregates, whereas global transcription regulators H-NS, FlgM regulate expression of SPIs and flagellar genes. Relatively low abundances of flavoredoxin, glutaredoxin, thiol peroxidase etc. leads to accumulation of ROS within the biofilm, and subsequent toxicity. This work further suggests that targeting these oxidative stress relieving proteins might be a good druggable choice to reduce *Salmonella* biofilm.

Importance

The enteric pathogen *Salmonella* forms biofilm in the internal organs of asymptomatic carriers, and on abiotic surfaces that leads to contamination of food and water. Biofilms are highly drug-resistant life forms that also helps in host immune evasion. Therefore, recent insurgence of drug tolerant strains necessitates development of biofilm inhibitory strategies, and finding novel

druggable targets. In this study we investigated the bioactive molecules present in the cell-free

supernatant of a biofilm deficient strain of *Salmonella* Typhimurium that inhibit biofilm

initiation by the wildtype strain. Further we showed that the supernatant treatment leads to

virulence defect *in vivo*. Collectively, our results suggest a comprehensive view of virulence

regulation in *Salmonella* by the cell-free supernatant of the biofilm deficient strain.

Introduction

Salmonella is one of the most common foodborne pathogens that causes one of the 4 major

diarrheal diseases, salmonellosis (1, 2). The biofilm life form of this bacteria makes it a hardy

human pathogen that can survive several weeks in a dry environment and many months in water

(3). A biofilm refers to a community of bacteria adherent to a biotic/abiotic substratum, held

together by its secreted extracellular polymeric substances. Biofilms are highly resistant to

antibiotics, disinfectants etc. and biofilms on medical devices, catheters, implants are a major

cause of hospital acquired infections (4, 5). Recent reports suggest that *Salmonella* can form

biofilm on the gall bladder, leading to chronic infection (6, 7). The major constituents of

Salmonella biofilm are curli, cellulose and BapA. Curli facilitates intercellular and cell to surface

interactions, cellulose enables long range contact and provides characteristic stickiness to the

biofilm and BapA strengthens curli interactions and stabilizes the biofilm (8). A major cue for

biofilm formation is nutritional stress. Previously we found that the deletion of carbon starvation

inducible gene, *yjiY*, leads to deficiency in biofilm formation (9). Antibiotic stress, DNA

damage, cold stress, acid stress etc. can also regulate the transcription of *yjiY* gene (10-12). In *S.*

Typhimurium, YjiY is known to regulate virulence by affecting flagellar class III genes, and the

deletion of *yjiY* leads to reduced colonization in mice (9, 13), and upregulation of the virulence

factor *mgtC*, leading to biofilm deficiency (14). This could indicate the possible role of YjiY in

survival and defense against host induced stress in bacteria.

Recent studies corroborate that various culture supernatants of commensal bacteria can inhibit

colonization and biofilm formation by their pathogenic counterparts. This inhibitory activity of

culture supernatants can be seen across Gram positive, and Gram negative bacteria as well as

across kingdoms. Iwase *et al.* showed that *Staphylococcus epidermidis* Esp inhibits biofilm

formation and nasal colonization by pathogenic *Staphylococcus aureus* (15). In another study,

probiotic *Escherichia coli* Nissle (EcN) inhibited biofilm formation in pathogenic EHEC, by a

secreted bifunctional (protease and chaperone) protein DegP (16). There are various other modes

of interspecies biofilm regulation such as biofilm dispersal proteins (17), indole-mediated

biofilm regulation (18), quorum sensing molecules, and antimicrobial peptides (19). The

inhibition can also be interkingdom, such as *Aspergillus* biofilm inhibition by *Pseudomonas*

culture supernatants (20). While there are a few strategies to deal with preformed biofilms (21,

22), the inhibition of biofilm initiation is comparatively less explored in human pathogens. In our

lab we found that *Salmonella* WT could not form biofilm when cocultured with Δ*yjiY* strain. This

study was designed to investigate the novel biofilm inhibitory activity of biofilm deficient strain

Salmonella Typhimurium Δ*yjiY*.

In this study, we are reporting a novel mechanism of biofilm initiation inhibition by a complex

transcriptional regulatory network. This complex network causes adhesion impairment, flagellar

motility inhibition, cleavage of protein aggregates and quorum sensing inhibition. Furthermore,

we have shown that STM Δ*yjiY* supernatant can also inhibit biofilm formation by *E. coli*. We

have also shown that the supernatant treatment can impair the invasion of the pathogen in the *C.*

elegans gut, thus reducing its virulence.

Results

Salmonella ΔyjiY culture supernatant inhibits WT biofilm formation

To understand the biofilm inhibitory property of STM Δ*yjiY* culture supernatant, we inoculated

WT bacteria in biofilm inducing low salinity media with or without the culture supernatant of

different strains. The Δ*yjiY* bacterial culture supernatant exhibited the biofilm inhibitory property

(**Fig. 1A**), which was absent in the supernatant of the strain where *yjiY* was trans-complemented

in a plasmid (STM Δ*yjiY*:pQE60-*yjiY*). Previous study showed that Δ*yjiY* is a biofilm deficient

strain (14). To check whether the biofilm inhibitory property is specific to Δ*yjiY* supernatant, we

inoculated WT strain with biofilm deficient strain Δ*csgD* (**Fig. S1A**). We observed inhibition

only in the case of Δ*yjiY* supernatant indicating that the inhibitory molecules are unique to the

Δ*yjiY* secretome, and are independent of the activity of CsgD. To further validate that the

inhibition is a cell-free phenomenon, we inoculated WT bacteria along with either live cells of

other strains along with spent culture media (coculture), supernatant-free cell pellets, cell free

supernatant, or whole cell lysate, and quantified the biofilm on solid-liquid-air interface.

Although Δ*yjiY* showed significant inhibition of biofilm formation in all the setups, the

maximum inhibition was observed with Δ*yjiY* culture supernatant (**Fig. 1B**). The supernatants

were concentrated using an Amicon ultra filter device and total protein was quantified. We found

that <u>m</u>inimum <u>b</u>iofilm <u>i</u>nhibitory <u>c</u>oncentration (**MBIC**; minimum concentration of total protein

required to inhibit biofilm formation by WT strain) falls between 15-20 ng protein/ml (**Fig. 1C**),

therefore we used supernatant containing 20 ng protein/ml of supernatant for further

experiments. We did not find any difference in the growth (**Fig. S1B**), suggesting that the

supernatant lacks any bactericidal or bacteriostatic properties. Interestingly, we observed slower,

yet longer exponential growth after supernatant treatment (**Fig. S1C**). To determine whether the

secretion of the inhibitory component(s), is dependent on the culture media, we grew the bacteria

in minimal media (M9 media supplemented with 0.5% glucose) that exerts nutrition stress and

used the supernatant to treat WT bacteria. We observed that the biofilm inhibitory molecule(s)

were active even in minimal media (**Fig. 1D**), suggesting that the production of the inhibitory

molecule(s) is not dependent on the nutritional condition and is an intrinsic property of the Δ*yjiY*

strain. Additionally we checked the temporal expression or accumulation of the inhibitory

component(s) by inoculating the WT strain with culture supernatant harvested from 2, 3, 4 and 5

day old Δ*yjiY* cultures. Our results suggest that the optimum concentration of the inhibitory

component(s) is/are reached after 3 days of growth (**Fig. S1D**), which remains unchanged on the

4^{th}-5^{th} days of growth. Therefore, we used filtered culture supernatant from 3 day old Δ*yjiY*

culture for further experiments.

Δ*yjiY* culture supernatant weakens the WT biofilm and interferes with cell structure

The characteristic EPS components are cellulose (produced by *bcsA* encoded cellulose synthase),

curli fimbriae (encoded by *csgAB*), BapA and LPS (23). We observed a significant reduction of

the EPS bound Congo red fluorescence intensity (**Fig. 2A, S2A**) and biofilm thickness (**Fig. S2B,**

S2C) upon Δ*yjiY* supernatant treatment. We also found that Δ*yjiY* supernatant treatment

significantly reduced the strength of the biofilm pellicle (**Fig. S2D, S2E, 2B**) than that of

untreated or WT supernatant treated samples. SEM and AFM analysis of the biofilm surface

showed that the Δ*yjiY* supernatant treated biofilms lack the characteristic dome shape of a proper

biofilm (**Fig. 2C, S2F**). We also noticed that the median cell length increased upon Δ*yjiY*

supernatant treatment (1.58$\pm$0.30 µm) as compared to the untreated WT cells (1.38$\pm$0.34 µm)

(**Fig. 2D, 2E, S2G**), hinting towards the presence of multiple regulatory components in the Δ*yjiY*

supernatant that can modulate multiple phenotypic effects.

The active molecule(s) is/are protein(s)

Previous studies have shown that secreted components from some bacteria can inhibit biofilm

formation by the wild type strain or closely related species (15, 16, 20). To delineate the

chemical nature of the inhibitory molecule(s) present in $\Delta yjiY$ supernatant, we treated the

supernatant with chemical agents, such as a divalent cation chelator (EDTA), protease

(Proteinase K), protease inhibitor (PMSF), RNase and DNase and quantified the biofilm

inhibition. We found that upon pre-treatment with different concentrations of EDTA, the

inhibitory property remained intact. Interestingly, 10 mM EDTA enhanced biofilm formation

with both treated and untreated WT strain (**Fig. 3A**). Since EDTA is known to chelate divalent

cations and inhibit a few proteases at higher concentrations (24-27), our data signify the

requirement of divalent cations and/or active proteases for biofilm inhibition. Upon treating the

supernatants with 20 mg/ml Proteinase K, the ability of $\Delta yjiY$ supernatant to inhibit biofilm

formation was significantly reduced, suggesting that the inhibitory molecule(s) are protein(s)

(**Fig. 3B**). Since the activity of many proteins is sensitive to even small changes in pH and

temperature, we checked the activity of $\Delta yjiY$ supernatant at different pH and temperatures.

Surprisingly, we found that the active component(s) is/are heat stable at 65 $^{\circ}$C and 95 $^{\circ}$C

temperatures (**Fig. 3C**) and stable over a wide range of acidic and alkaline pH (**Fig. 3D**),

although there was a small reduction in biofilm inhibition at pH 9.0. As recent studies show that

several small noncoding RNAs regulate biofilm formation and other virulence traits in *Vibrio*

cholerae and *Pseudomonas aeruginosa* (28, 29), we tested the stability of the component(s) after

treating the supernatant with RNase. Although the inhibition was lost upon RNase treatment and

incubation at 37°C, we found a similar loss of inhibition with only heating the supernatant at

37°C (**Fig. S3A**), suggestive of the heat, rather RNase treatment, to be the reason for the loss of

inhibition. We also fractionated the supernatant using an Amicon 3k MWCO ultra filter device,

and we found that the active component(s) of $\Delta yjiY$ supernatant is/are >3kDa molecular weight

(**Fig. S3B**), quashing the role of small molecules and ions in biofilm inhibition by $\Delta yjiY$

supernatant. Since the inhibitory activity was abolished upon proteinase treatment, we attributed

the inhibition to protein components and quantified the total protein for further experiments.

Active molecules inhibit biofilm only during the initial phases, and cannot disrupt mature

biofilm

To determine the effect of the active molecule on mature biofilm, we treated mature biofilm

pellicles with the supernatants and checked for dispersion. Our results showed that the $\Delta yjiY$

supernatant could not disrupt pre-formed biofilm, hinting towards the effect of the active

molecule(s) on biofilm initiation (**Fig. 4A**). To determine whether the supernatant treated WT

cells remained biofilm defective in the absence of the inhibitory molecules, the $\Delta yjiY$ supernatant

treated WT cells were re-inoculated in biofilm medium without the supernatant, and monitored

for pellicle formation. We observed that the inhibition was diminished in the absence of the

supernatant (**Fig. 4B**), although the defect reappeared when these cells were re-treated with the

inhibitory supernatant. Together our data suggest that supernatant mediated initial molecular

reprogramming is required for biofilm inhibition, and that inhibitory molecules do not alter the

inherent biofilm forming ability of the WT cells.

The inhibitory molecules impair flagella-mediated initial attachment to abiotic surfaces

Salmonella biofilm formation can be divided into 4 distinct steps, (i) initial attachment to the

abiotic surface, (ii) secretion of adhesive components, (iii) maturation and (iv) dispersion of

biofilm upon relief of the stress (30). Since biofilm initiation was inhibited and retained beyond

72 hours, we checked the expression of biofilm-associated genes (*csgD, bcsA, fliC*), SPI-1

effectors (*invF, sopD)* and virulence factors (*phoP, sodA, mgtC*) from untreated or treated WT

cells, 72 hours post inoculation. We found that *csgD, bcsA*, *fliC* and *invF* expression was

significantly downregulated in Δ*yjiY* supernatant treated WT cells (**Fig. 4C**). Since, flagella-

mediated initial attachment initiates biofilm formation (31, 32), we investigated the effect of

Δ*yjiY* supernatant on flagellar motility. We observed that Δ*yjiY* supernatant treatment

significantly reduced both swimming and swarming motility (**Fig. 4D, 4E** and **S4A**), indicating a

flagella-mediated motility defect. We further observed the downregulation of *fliC* occurs as early

as 4-6 hours post inoculation with the Δ*yjiY* supernatant (**Fig. S4B**). Interestingly, we observed

an increase in *fliC* expression at 12 hours post inoculation. We reasoned that adhesion deficiency

arises from the sequential effect of the initial downregulation of *fliC*, followed by an increase in

the planktonic population. SEM analysis showed that Δ*yjiY* supernatant treated bacteria had

significantly fewer flagella (**Fig. 4F, S4C**). Altogether, these results indicate a defect in flagella-

mediated initial attachment of the bacteria, sets the course for biofilm deficiency.

Δ*yjiY* supernatant reduces adhesion and virulence of *Salmonella* both *in vitro* and in *C.*

***elegans* gut**

Initial attachment of *Salmonella* on the host gut epithelial cells requires swimming through the

mucus layer (33). SPI1 encoded Invs, Sops and Sips are important in initial invasion (34). Since

Δ*yjiY* supernatant treatment downregulated *invF* and *sopD* expression, we checked the infectivity

of the supernatant treated cells in mammalian intestinal epithelial cells, Int407. Invasion assay

shows that Δ*yjiY* supernatant treated cells are defective in initial invasion, although these cells

showed significantly higher intracellular proliferation than the untreated cells (**Fig. 5A** and **5B**),

suggesting that the Δ*yjiY* supernatant only makes WT cells invasion defective by inhibiting

flagella-mediated adhesion. To check the infectivity of the Δ*yjiY* supernatant treated cells in a

systemic condition, we fed young adult *C. elegans* N2 worms RFP-STM-WT bacteria and

quantified gut colonization. The micrograph images and CFU analysis show that Δ*yjiY*

supernatant treated RFP-STM-WT cells were able to colonize the gut lumen when fed

continuously, but did not persist (**Fig. 5C, 5D, 5E**). Therefore, our data suggest that Δ*yjiY*

supernatant treatment impairs *in vitro* invasion and *in vivo* colonization.

Inhibitory molecules impede biofilm formation in closely related species

We next checked the effectiveness of STM Δ*yjiY* supernatant on the biofilms of common human

pathogens. We inoculated *E. coli* DH5α, *Pseudomonas aeruginosa* PA01, *Klebsiella*

pneumoniae, *and Staphylococcus aureus* wild-type strains with STM WT supernatant, and STM

Δ*yjiY* supernatant, and quantified the biofilm. We found that the STM Δ*yjiY* supernatant

significantly inhibited only *E. coli* DH5α biofilm which is regulated by CsgD (**Fig. 5F**).

Therefore, we concluded that the Δ*yjiY* supernatant can cross-react with closely related

Enterobacteriaceae, *E. coli*, but is not effective against distant members such as *Klebsiella*

pneumoniae or different family (*Pseudomonas aeruginosa*) or phylum (*Staphylococcus aureus*).

The active components are primarily global transcription factors that regulate multiple cellular processes

To further identify the biofilm inhibitory molecule(s), concentrated supernatants from WT and

Δ*yjiY* cultures were resolved on 10% SDS-PAGE. After colloidal CBB staining, we clearly

visualized 14 differential bands in Δ*yjiY* supernatant compared to WT supernatant (**Fig. 6A**). We

next analyzed the secretome in LC Q-TOF MS/MS. Among the 244 proteins, 188 proteins were

present in both supernatants at differential levels, while 38 and 58 proteins were enriched in the

WT supernatant and Δ*yjiY* supernatant, respectively (**Fig. 6B**). These proteins may have resulted

from cell lysis during growth or they were secreted in the supernatant via an active secretion

system. After careful data-mining, among the proteins found only in Δ*yjiY* supernatant (**Table**

S1), probable transcriptional regulatory protein YebC, anti-sigma28 factor FlgM, a serine

protease inhibitor ecotin and transcription termination/anti-termination protein NusG were

selected for further analysis. Many transcriptional regulators (H-NS, Rnk, StpA), cold shock

proteins (CspE, CspC), chaperones GrpE, ATP-dependent protease HslV, proteins related to

oxidative stress and iron homeostasis (FldA, SodB, YdhD, Tph and Ftn, Bcp, Bfr, respectively)

were differentially present in both supernatants (**Table S2**). Additionally, we treated STM WT

cells with purified ecotin, HNS, and NusG (**Fig. S5**), to delineate the effect of each of these

proteins. Interestingly, the fractions containing monomeric ecotin and NusG, as well as

homodimeric HNS (**Fig. S6A, S6B**), showed biofilm inhibition similar to that of Δ*yjiY*

supernatant, whereas the buffer (50mM Tris) treated WT cells did not have biofilm defect (**Fig.**

6C). Together, our data suggest that Δ*yjiY* supernatant perturbs various cellular processes leading

to biofilm defect by complex transcriptional regulation (**Fig. 6D**).

Discussion

Salmonella forms biofilm to evade host defense, colonize in host, persist in asymptomatic host,

and transmit to a new host (35). The transition from planktonic to biofilm mode depends upon

various stress factors. The master regulator CsgD controls the production of EPS components

such as cellulose, tafi (curli fimbriae) etc. by regulating AdrA. BapA is important both for

biofilm production and attachment to intestinal epithelium (36). In this study we have shown

how metabolic stress determines the fate of the infectious WT strain. From preliminary data, we

found that coculture of STM WT and Δ*yjiY* led to biofilm defect in the WT strain. Although

various studies have shown that interspecies and intraspecies biofilm inhibitions exist in nature

(15, 20, 37), in *Salmonella,* this phenotype is novel. Therefore, we carried out experiments to

understand the underlying mechanism.

In this study, we found that biofilm inhibition was unique to Δ*yjiY* supernatant and the inhibition

was reversed when WT cells were treated with culture supernatant from the complement strain

(STM Δ*yjiY*:pQE60-*yjiY*). Although we observed that the Δ*yjiY* supernatant had only a biofilm

inhibitory effect and lacked bactericidal properties (as observed from the growth analysis), STM

WT cells showed delayed yet prolonged exponential growth upon supernatant treatment.

Furthermore, the exopolysaccharide cellulose was sparse in the biofilm of Δ*yjiY* supernatant

treated WT cells. Since only cellulose mutant *Salmonella* Typhimurium was proficient in

adhering to tumour cells (38), our data imply the presence of multiple inhibitory factors involved

in the inhibition by the supernatant. While incubation of Δ*yjiY* supernatant at 37°C was found to

reduce the inhibition, heating at 65°C and 95°C enhanced the inhibition, indicating that the

inhibitory molecules might be heat-shock proteins or high temperature inducible stress proteins.

Fascinatingly we observed that the effect of the inhibitory molecules is temporary, and they do

not cause any genetic change to alter the inherent biofilm forming competency of the WT cells.

Δ*yjiY* supernatant was also found to exert its anti-biofilm activity only when administered at the

beginning of biofilm formation, suggesting a modification in the complex network through

which biofilm develops. In the Δ*yjiY* supernatant treated WT cells, the expression of biofilm

genes and virulence genes was reduced significantly. *sodA* downregulation suggests that these

cells are prone to ROS assault. A previous study from our group showed that YjiY depletion

upregulates *mgtC* leading to biofilm defect (14). Interestingly, we observed downregulation of

mgtC, implying that biofilm inhibition by Δ*yjiY* supernatant and the inherent biofilm defect of

the Δ*yjiY* strain follow different mechanisms. The *ΔyjiY* supernatant treated WT cells showed

significantly less invasion in Int407 cells. It was recently proposed that *Salmonella* persists in *C.*

elegans gut by forming biofilm (39). Δ*yjiY* supernatant treated WT cells showed a concomitant

reduction in colonization in the worm gut, suggesting an *in vivo* biofilm defect. We also observed

the absence of flagella and other protein aggregates upon Δ*yjiY* supernatant treatment, leading to

defects in cell aggregate formation and biofilm initiation. While there was a temporal increase in

fliC in Δ*yjiY* supernatant treated WT cells, these cells exhibited defective motility after 72 hours

of treatment, further validating the importance of flagella-mediated motility in the initial

attachment of the bacteria to the substratum. Interestingly, we found that STM Δ*yjiY* supernatant

effectively inhibited *E. coli* biofilm formation. Although *Salmonella* diverged from *E. coli* by

acquiring virulence-associated genes, they share many evolutionarily conserved cellular

pathways (40-42). Therefore this observation reiterates that the inhibitory effects are more

profound among closely related species. In the proteomics analysis of the Δ*yjiY* supernatant, we

specifically detected three potential inhibitory candidates: proteinase K sensitive ecotin (43),

anti-sigma28 factor FlgM (a negative transcriptional regulator of class III flagellar genes (44,

45)) and YebC (negatively regulates quorum sensing in *Pseudomonas aeruginosa* PA01 (46)).

The abundance of FlgM correlates with the absence of flagella in the Δ*yjiY* supernatant treated

WT cells. NusG works synergistically with the global transcriptional regulator H-NS, which

binds specifically to AT-rich SPIs in the *Salmonella* genome and represses those genes (47, 48).

Interestingly, in *E. coli,* H-NS has been linked to the cell cycle, since the cells attempt to

optimize spatial H-NS concentration by maintaining a constant ratio of H-NS to chromosomal

DNA in the cell (49), which might explain the increase in cell length after Δ*yjiY* supernatant

treatment. HslV, a heat-inducible ATP-dependent protease subunit of a proteasome-like

degradation complex (50) and GrpE, which is involved in removal of protein aggregates leading

to unsuccessful biofilm formation (51) were found in higher abundance in Δ*yjiY* supernatant.

Doyle *et al.*(52) showed that excess GrpE inhibits the interaction between DnaK and the

regulatory protein, RepA. Since RepA helps maintain the plasmid copy number in *E. coli* (53),

inactivation of RepA might explain cell elongation upon Δ*yjiY* supernatant treatment. While

protein aggregation is necessary during initial attachment of the bacteria to an abiotic surface,

upon maturation, proteolytic activity of several proteases helps in dispersal of biofilm (54).

Presence of a protease inhibitor, such as ecotin, might hinder this process, leading to

dysregulation of biofilm maturation and dispersion. Although several moonlighting functions

have been demonstrated for mycobacterial superoxide dismutase and DnaK in modulating host

response (55), any such activity for other hits remains elusive.

Oxidative stress proteins such as flavoredoxin, SodB, glutaredoxin, thiol peroxidase (Tph), and

thioredoxin-dependent thiol peroxidase (Bcp) were less abundant in Δ*yjiY* supernatant, among

which Tph and SodB help reduce oxidative stress in STEC biofilm (56). In *E. faecalis*, Tph is

required for the *in vitro* oxidative stress response and survival inside murine macrophages (57).

Iron availability has been shown to both positively and negatively regulate biofilm formation

through complex network systems (58). Although high Fe can lead to ROS production, Kang and

Kirienko suggested that iron uptake and homeostasis are essential for successful biofilm

formation in *Pseudomonas aeruginosa* (59). The Fe-storage proteins bacterioferritin (Bfr) and

ferritin A (Ftn), required to prevent ROS generation via the Fenton reaction and DNA damage,

are functionally very large proteins with a core to accommodate 3000 Fe atoms. Fe-rich

conditions induce *E. coli* FtnA and Bfr due to loss of repression by small RNA RyhB(60).

Similarly, *Salmonella* Typhimurium Bfr is involved in reducing intracellular Fe toxicity, and the

absence of Bfr causes increased intracellular free Fe^{2+} ion and oxidative stress(61).

Therefore, we conclude that Δ*yjiY* supernatant treatment inhibits biofilm formation by WT in

four major ways- (i) NusG-HNS mediated transcriptional repression of AT-rich SPI-encoded

genes, making the WT bacteria less virulent; (ii) FlgM mediated downregulation of class III

flagellar genes, impairing flagella-mediated initial attachment to abiotic surfaces, (iii) High

abundance of proteases and proteolytic molecules, hindering cell-cell adhesion, and cellular

aggregate formation in the EPS matrix, and (iv) Ecotin mediated inhibition of proteases that are

necessary during biofilm dispersal. We reasoned that the lower abundance of redox homeostasis

proteins and ferritin-like molecules in the Δ*yjiY* supernatant might facilitate the accumulation of

toxic oxidative species, causing cellular stress and toxicity. In this light, these oxidative stress

relieving proteins can be exploited as potential druggable targets to inhibit *Salmonella* biofilm

initiation.

Materials and Methods

Bacterial strains

All *Salmonella* Typhimurium strains used in this study are listed in the following section with

their genetic descriptions. *Salmonella enterica* serovar Typhimurium strain 14028S was used as

the wild type strain, and was also the parental background for all the mutant strains used in this

study, i.e. Δ*yjiY*, Δ*csgD*, Δ*fliC* and Δ*fliC* Δ*fljB*. All strains were grown and maintained in Lennox

broth (LB; 0.5% NaCl, 1% casein enzyme hydrolysate and 0.5% yeast extract) at 37˚C under

shaking conditions. STM GFP, STM mCherry (RFP-STM-WT) and STM *ΔyjiY:yjiY* were

cultured in Lennox broth with 50 µg/ml Ampicillin at 37˚C in shaking condition. *E.coli* DH5α,

Staphylococcus aureus, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae* were grown in

Lennox Broth at 37°C in shaking condition.

List of strains used in this study.

Strain name	Description	Reference
S. Typhimurium ATCC 14028S (STM_WT)	Wild type (WT)	Kind gift from Prof. M. Hensel (Division of Microbiology, University of Osnabrück, Germany)
Δ*yjiY*	Isogenic knockout strain for the gene *yjiY*; Kan[r]	Garai *et al.* (2016)
Δ*csgD*	Isogenic knockout strain for the gene *csgD*; Chl[r]	Srinandan *et al.* (2015)
Δ*fliC*	Isogenic knockout strain for the gene *fliC*; Chl[r]	Garai *et al.* (2016)
Δ*yjiY*-pQE60-*yjiY*	Isogenic complement strain for Δ*yjiY* expressing *yjiY* under the promoter of T5 present in the plasmid pQE60; Amp[r]	Garai *et al.* (2016)
Δ*fliC* Δ*fljB*	Isogenic double knockout strain for *fliC* and *fljB*; Kan[r] Chl[r]	Karmakar *et al.* (2019)

STM mCherry (RFP-STM-WT)	STM 14028s wildtype strain carrying pFPV-mCherry plasmid, Amp^r	This study
STM GFP	*S.* Typhimurium rpsM promoter driving gfpmut3 expression through pFPV25.1 plasmid, Amp^r	This study

Growth conditions for biofilm formation

LB without NaCl, i.e. 1% casein enzyme hydrolysate and 0.5% yeast extract, was used as biofilm media. Overnight cultures grown in LB were subcultured in 2 ml biofilm medium at the dilution of 1:100 in a flat-bottom 24-well polystyrene plate (Tarsons) and incubated at 28°C for 72 hours without shaking. All images of biofilm in the form of a pellicle were taken with a digital camera (Olympus Stylus VH-520).

Preparation of cell free supernatant

To 2 mL of biofilm media, 1:100 dilution of overnight grown culture was added in flat-bottom 24-well polystyrene plate (Tarsons). The plate was incubated at 28°C without shaking for 72 hours. After 72 hours, the media from each well was collected, centrifuged twice at 10000 rpm for 15 minutes, the supernatant was collected and filtered using a 0.2 µm filter and stored at -20°C. To collect supernatant from M9 (0.5% glucose) media, the cultures were grown at 37°C in shaking conditions. After 72 hours, the supernatant was collected as described above.

Bacterial biofilm inoculation with culture supernatant

To test the effect of the 3 day culture supernatant (chemically treated and untreated) on biofilm

formation, 20 µl of STM WT overnight culture and indicated volume (1% v/v and/or as indicated

in the figures) of the supernatants (WT supernatant, $\Delta yjiY$ supernatant, $\Delta yjiY:yjiY$ supernatant

and $\Delta csgD$ supernatant) was added to 2 ml of biofilm media, and incubated under biofilm

inducing condition. **Cell pellets** were collected from 3 day old biofilm culture by harvesting the

contents of the flat-bottom 24-well polystyrene plates (Tarsons), centrifuging at 10000 rpm for

15 minutes. The cells were resuspended in fresh biofilm inducing media and added at the

concentration of 0.5×10^7 CFU/ml of biofilm media and 20 µl of STM WT overnight culture, and

incubated under biofilm inducing condition. For **co-culture** experiments, the overnight cultures

of the strains were added in 1:1 ratio to 2 ml of biofilm media and incubated under biofilm

inducing condition. **Cell lysates** were prepared by adding 30 µl 1x TME buffer (25 mM Tris pH

8.0, 1 mM EDTA pH 8.0, 2 mM β-mercaptoethanol) and sonicating the bacterial samples (STM

WT, STM $\Delta yjiY$, and STM $\Delta yjiY:yjiY$). The samples were centrifuged at 12000 rpm for 10

minutes and 2 µl supernatant was used for inoculation along with 20 µl STM WT overnight

culture and 2 ml of biofilm media and incubated under biofilm inducing condition.

Crystal Violet staining

To quantify the biofilm at the solid-liquid interface, crystal violet (CV) staining was carried out

as mentioned in Chandra *et al*(14). Briefly, the protocol followed for biofilm formation was the

same as mentioned above. After 72 hours, the media in each well was discarded and the plates

were thoroughly washed with RO water to remove all planktonic cells. The plates were then

dried and 2 ml of 1% CV was added into each well. After 15 minutes, the CV was removed, and

the excessive stain was thoroughly rinsed with RO water. The stained biofilm was destained with

70% ethanol and the intensity of color of the destained solution was quantified at OD_{595} in Tecan

plate reader (Infinite Pro 200). The absorbance was plotted in GraphPad Prism 6 and significance

values determined using Student's t-test or two-way ANOVA.

Confocal microscopy

Sterile square coverslips (18 mm) were placed in flat-bottom 12-well polystyrene plate

(Tarsons). Cultures were inoculated for biofilm formation as mentioned previously. After 72

hours, biofilm appeared on the coverslip at the liquid–air interface, in the form of a thin line

spanning the width of the coverslip (18 mm). The coverslip was washed thoroughly with water to

remove planktonic cells and stained with Congo red (20 mg/ml in water) for 20 min at room

temperature. After washing with water, the coverslip was mounted on a slide and imaged for

biofilm distribution, with a laser scanning confocal microscope (Zeiss LSM 710) using a 40x

objective. Z stacks were taken to generate a three-dimensional image. The MFI of the images

were calculated using the ImageJ software. The MFI and thickness of the biofilm were plotted

using GraphPad Prism 6. Single layer of cells were imaged and cell length of ~1000 cells from

each coverslips was measured using ImageJ.

Glass bead assay

To test the strength of the pellicle at the air-liquid interface, medium sized (0.5mm to 1mm) glass

beads were added one by one onto the pellicle. The initial weight of the glass beads was noted,

and the number of glass beads added until the pellicle just collapsed was noted down and plotted

using GraphPad Prism 6.

Scanning electron microscopy

Biofilm was allowed to form on coverslips as mentioned in the previous section. After thorough

washing with water, the sample was fixed in 2.5% gluteraldehyde for 48 hours at room

temperature. Excess gluteraldehyde was removed by washing with water and the sample was

dehydrated by gradient washes in increasing concentrations of 30%, 50%, 75%, 85% and 95%

ethanol. The coverslips were then air dried under vacuum before coating with gold (JOEL-JFC-

1100E ion sputtering device) for imaging by scanning electron microscope. For checking

flagellar morphology, 20 µl of STM WT overnight culture and 20 µl of supernatant treated STM

WT were smeared on autoclaved 18mm coverslips, air dried, and processed using the

abovementioned protocol. Flagellar structure was imaged using field emission-SEM (FEI Sirion,

Eindhoven, The Netherlands) scanning electron microscope.

Atomic force microscopy

Sterile 18 mm square coverslips were placed in flat-bottom 12-well polystyrene plates (Tarsons)

and biofilm inoculation was done. After 72 hours, the coverslips were removed and washed with

sterile MilliQ water. The coverslips were dried and AFM analysis was done using XEISS AFM

systems and was analyzed using XEI software.

Supernatant conditioning

The supernatant was treated with different concentrations of EDTA (2.5 mM, 5 mM, 7.5 mM and

10 mM) and incubated at 65°C for 1 hour. To 200 µl of supernatant, 10 µl of proteinase-K (NEB,

stock 20 mg/ml) was added and incubated at 37°C for 1 hour. The proteinase was inactivated

with 0.5 mM PMSF and incubated at 28°C for 1 hour. The supernatant was heated to 37°C (60

min), 65°C (15 min) or 95°C (15 min) for indicated time and immediately frozen at -20°C. The

pH of the biofilm inducing media as well as that of the supernatant were adjusted using

concentrated HCl and 10N NaOH to obtain pH of 4, 7.4, and 9. The supernatants were also

treated with 10 µl RNase (stock 1 mg/ml) for 1 hour at 37˚C. Post treatment, the supernatants

were used to inoculate biofilm in order to check the activity of the inhibitory molecule(s).

Preformed biofilm disruption

425   STM WT was allowed to form biofilm as described earlier. 72 hours post inoculation, the formed

426   biofilms were treated with 20, 50 or 100 µl (1% (v/v), 2.5% (v/v) or 5% (v/v), respectively)

427   supernatants and was incubated at 28˚C for 72 hours and biofilm was quantified using CV

428   staining.

Recovery of biofilm formation

430   STM WT was allowed to form biofilm in the presence of supernatants as described earlier. 72

431   hours post inoculation, the cells were harvested by centrifuging the culture at 10000 rpm for

432   15min. 20 µl of these treated or untreated cells were inoculated in fresh biofilm media (with and

433   without supernatants for retreatment or recovery, respectively) and incubated at 28˚C for 72

434   hours and biofilm was quantified using CV staining.

Supernatant concentrating and separation based on molecular weight

436   The supernatants from biofilm culture were harvested and filter sterilized as previously

437   mentioned. 4 ml of the supernatants were transferred to the Amicon ultra filter device (Amicon®

438   Ultra-4 centrifugal filter device, 3k MWCO, UFC800324) and centrifuged at 4000g for 30 min in

439   a swing bucket rotor. The concentrated solute from the bottom of the filter device was collected

440   by inserting a pipette.

Quantitative RT PCR

STM WT was allowed to form biofilm in the presence or absence of supernatants as described

earlier. After 72 hours, the biofilm population as well as the planktonic cells were harvested by

thorough pipetting and centrifugation. From these cells, RNA was isolated by the TRIzol method

(Takara). cDNA was synthesized with reverse transcriptase (GCC Biotech). Quantitative PCR

was carried out using SYBR Green Q-PCR kit (Takara).

List of oligonucleotides used in this study.

Gene	RT primer sequence (5'-3')
csgD	FP: ACGATTATCCCTACCGTGAA RP: GCGGACTCGGTGCTGTTGTA
bcsA	FP: CTGCACCAGTCCGTAAAACA RP: AGTTTGTGGCGATCTTCGATT
bapA	FP: ACGGTGACTACGCCTAAAAC RP: CTTCCGCCTCAGTCACTTTT
fliC	FP: CTAAACAAACTGGGTGGCGC RP: GCACCCAGGTCAGAACGTAA
fimA	FP: TTCGGATCGCAGTCATTCAG RP: GTTGCGGCTGATCCTACTC
invF	FP: AGATCGTAAACGCTGCGAGT RP: CTGCTGCACAAACGACGAAA

phoP	FP: GATCTCTCACGCCGGGAATT RP: TGACATCGTGCGGATACTGG
sodA	FP: CCTGCCGGTTGAAGAACTGA RP: GGTTGCTGCTGCTTTTTCGA
ssaV	FP: TATTGATAGGCGCGGACGCTA RP: CGCCTTATGGGCCATGTCTTT
mgtC	FP: GGACCGAACCTAACCCTTG RP: AACTCCCAATATCCGCTGAG
16s rRNA	FP: GTGAGGTAACGGCTCACCAA RP: TAACCGCAACACCTTCCTCC

***In vitro* motility assay**

2 µl of bacterial samples (treated or untreated) were spotted onto the 0.3% agar plates

supplemented with 0.5% yeast extract, 1% casein enzyme hydrolysate, 0.5% NaCl and 0.5%

glucose (swim agar plates) or 0.5% agar plates supplemented with 0.5% yeast extract, 1% casein

enzyme hydrolysate, 0.5% NaCl and 0.5% glucose (swarm agar plates). The plates were

incubated at 37˚C and images were taken every 2 hours using a digital camera (Olympus). The

diameters of the motility halos were measured using ImageJ. At least five replicate plates were

used for each condition, and statistical significance was calculated using Student's *t*-test.

Transmission electron microscope

Flagella and fimbriae were visualised using the protocol described in Garai et al 2016(9). Briefly,

overnight STM WT were inoculated with or without supernatants in biofilm media and kept

under biofilm inducing condition for 72 hours. After 72 hours, both biofilm and planktonic cells

were collected and fixed with 2.5% glutaraldehyde overnight at 4°C. Similarly fixed overnight

cultures of STM WT and STM Δ*fliC* Δ*fljB* were used as positive and negative control,

respectively. 2 µl of the cell suspension was added on copper grid, air dried, stained with 1%

uranyl acetate for 30 sec, and visualised under transmission electron microscope.

In vitro intracellular survival assay

Int407 cells (ATCC® CCL-6™) were seeded at a density of $2x10^5$ cells per well in flat-bottom

24-well polystyrene plates (Tarsons). 72 hours old supernatant treated or untreated biofilm

cultures were sub-cultured, grown to mid-log phase and used for infection of Int407 at MOI 10.

The plates were incubated at 37°C with 5% CO_2 for 20 min to allow infection of the cells. The

cells were incubated for an hour in 100 µg/ml Gentamicin. The cells were then washed, and

grown in DMEM containing 25 µg/ml Gentamicin. After 1h, 2h and 16h, the cells were lysed

with 0.1% Triton-X100 and bacterial load was estimated by plating on SS- agar plates.

In vivo virulence in *C. elegans* N2 worms

STM-RFP bacteria were treated with the supernatants, and kept for biofilm formation for 72

hours. After that, STM-RFP cells were harvested as mentioned before, and 10^7 CFU were seeded

on NGM plates, incubated for 12 hours at 37 °C. *E. coli* OP50 fed *Caenorhabditis elegans* N2

young adults were harvested in M9 buffer, counted and approximately 150 worms were fed with

supernatant treated or untreated 10^7CFU of STM WT RFP bacteria for 12 hours. The worms

were then harvested in 1ml M9 buffer, washed 6-8 times to remove extracellular *Salmonella*. 10

worms from each sample were transferred to sterile M9 buffer containing 0.5mm sterile glass

beads. The worms were lysed, and the gut content of the worms was plated on SS-agar plates.

~20 worms were taken for laser scanning confocal microscopy analysis using Zeiss LSM 880

with Multiphoton mode. ~50 worms were transferred to *E. coli* OP50 plate and fed for another

12 hours. After 12 hours, the worms were sampled in a similar manner for CFU analysis and

imaging.

SDS-PAGE and colloidal Coomassie Blue staining

The proteins in the supernatants were quantified by Bradford's assay and equal quantity of the

STM WT and STM Δ*yjiY* supernatants were loaded onto the SDS-PAGE gel (5% stacking, 10%

resolving). The gel was then stained with colloidal Coomassie Brilliant Blue R-250 for 3 days

and then washed with water to remove excessive stain and visualize the band.

LC QTOF MS/MS analysis of the supernatants

Total protein from the 10-fold concentrated supernatants was subjected to in-solution trypsin

digestion for LC Q-TOF MS analysis. Briefly, 200 mg of total protein was resuspended in 200 µl

of 50 mM Ammonium bicarbonate. To this, 10 mM DTT was added, and incubated at 56°C for

30 min, followed by alkylation with 20 mM iodoacetamide and incubation at 37°C for 30 min in

dark. For protein digestion, trypsin solution was added to the sample to a final protease to protein

ratio of 1:50 (w/w) and incubated at 37°C for 16 hours, with frequent shaking. The reaction was

stopped by adding 10 µl of 0.1% formic acid, and samples were stored at -20°C until loaded on

the Agilent AdvanceBio Peptide Map column (2.1 ×150 mm, 2.7µ). 20 µl of the digested

samples were injected to the column, and analysed at a flow rate of 0.4 ml/min using 0.1%

formic acid in water as mobile phase A and 0.1% formic acid in acetonitrile as mobile phase B.

All the samples were analyzed using Agilent 1290 Infinity II LC System coupled with Agilent

AdvanceBio Q-TOF (6545XT). The MS and MS/MS scan were acquired in the positive mode

and stored in centroid mode. The MS data acquisition parameters were as follows- Vcap was set

at 3500V, drying gas flow rate and temperature was set at 13 ml/min and 325 ˚C, respectively. A

collision energy with a slope of 3.6 V/100 Da and an offset of 4.8 V was used for fragmentation.

The Precursor ion data were captured in a mass range of m/z 200– 2700 and fragment ions data

were acquired between m/z 50–2800. The raw data were analyzed using MaxQuant software

(v1.6.2.10) and processed through MS excel sheet. The database analysis was performed against

Salmonella Typhimurium proteome in UniProt (Proteome ID: UP000002695). Following search

parameters were used for the database analysis: Precursor mass tolerance: 10ppm, fragment mass

tolerance: 40ppm, fixed modifications: carbamidomethyl (C), variable modification: oxidation.

Cloning, expression and purification of ecotin, HNS and NusG

S. Typhimurium 14028S Ecotin (STM14_2792), HNS (STM14_2116) and NusG (STM14_4985)

were cloned with 6x His tag in pET15b vector using Gibson assembly. Positive clones were

confirmed by PCR and transformed in *E. coli* BL21 (DE3) pLysS strain. *E. coli* BL21 (DE3)

cells bearing expression plasmids were grown overnight at 37°C in LB media containing 50

µg/ml chloramphenicol and 100 µg/ml ampicillin. 1% of the primary inoculum was added to

500ml of LB media containing 100 µg/ml ampicillin and grown at 37°C till the culture attained

an OD_{600} of ~ 0.8, and induced with 1mM isopropyl β-D-1-thiogalactopyranoside (IPTG) at

30°C for 6 hours. The cells were harvested by centrifugation, and the pellet was resuspended in

20ml ice-cold resuspension buffer (0.2M Tris (pH 7.5), 0.5mM EDTA, and 1mM PMSF).

Further, the cells were lysed by sonication on ice, followed by centrifugation at 14000g. The

protein was purified from the soluble fraction of the lysate by affinity chromatography using Ni-

NTA column. Protein was eluted with linear gradient of 0.1-1M imidazole in 50mM Tris (pH

8.0). Elute fractions containing protein of interest were pooled, concentrated and subjected to

purification by size exclusion chromatography (SEC) on Superdex 75 Increase 10/300GL

column (with a flow rate of 0.5 ml/min and 500 µl sample injection volume) equilibrated with

50mM Tris (pH 8.0). The peaks obtained from the chromatogram were analyzed by subjecting

the collected fractions (which were pooled and concentrated after SEC) to SDS-PAGE, to obtain

the protein of interest in its desired monomeric form. The protein purity was also confirmed by

MALDI-TOF.

List of oligonucleotides used for cloning.

Oligo Name	Primer sequences (5'-3')
pET15b_Ecotin	**FP:**CGGCCTGGTGCCGCGCGGCAGCCATATGAAGAT GTTTGTCCCTGCCG
	RP:CTTTCGGGCTTTGTTAGCAGCCGGATCCTCATCG CGCGACGGCGTTC
pET15b_HNS	**FP:**GCGGCCTGGTGCCGCGCGGCAGCCATATGAGCG AAGCACTTAAAATTC
	RP:TTTCGGGCTTTGTTAGCAGCCGGATCCTTATTCC TTGATCAGGAAATCTTCC
pET15b_NusG	**FP:**AGCGGCCTGGTGCCGCGCGGCAGCCATATGTCTG AAGCACCTAAAAAGC
	RP:TCGGGCTTTGTTAGCAGCCGGATCCTTACGCTTT CTCAACCTGACTG

Statistical analysis

Statistical analyses were performed with GraphPad Prism software. Student's t test and one-way or two-way ANOVA tests were performed as indicated. The results are expressed as mean $\pm$ SD or mean $\pm$ SEM. Group sizes, experiment number, and p values for each experiment are described in the figure legends.

Acknowledgements

We thank the electron microscopy facility of IISc, AFMM facility of IISc, Atomic force microscopy facility of BSSE, IISc and confocal microscopy facility and real-time facility of Dept. of Microbiology and Cell Biology, IISc. We thank Mr. Priyabrata (TH lab, MRDG, IISc) for the helping with the TEM sample preparation. We thank Dr Shahbaz and Dr Jayantika (RV lab, MBU, IISc) for helping with cloning, protein expression and purification. We are very grateful to Dr Kapudeep and Ms. Meghanashree for their valuable inputs.

Funding

This work was supported by the DAE SRC fellowship (DAE00195) and DBT-IISc partnership umbrella program for advanced research in BioSystems Science and Engineering to DC. Infrastructure support from ICMR (Centre for Advanced Study in Molecular Medicine), DST (FIST), and UGC (special assistance) is acknowledged.

Author Contribution

KC and DC conceived the study and designed experiments. KC and PM performed experiments. SK performed and UST provided valuable inputs for the LC QTOF MS/MS experiment. KC analyzed the data, prepared the figures and wrote the manuscript. DC supervised the work. All the authors read and approved the manuscript.

557   **Competing Interests**

558   The authors declare no competing interests.

559   **References**

560   1.   Bakhshi B, Dehghan-Mouriaabadi A, Kiani P. 2018. Heterogeneity of Multidrug-Resistant

561   Salmonella enterica Isolates with Increasing Frequency of Resistance to Ciprofloxacin During a 4-

562   Year Period in Iran. Microb Drug Resist 24:479-488.

563   2.   Michael GB, Schwarz S. 2016. Antimicrobial resistance in zoonotic nontyphoidal Salmonella: an

564   alarming trend? Clin Microbiol Infect 22:968-974.

565   3.   WHO. 2018. Salmonella (non -typhoidal) Fact sheet. https://www.who.int/en/news-room/fact-

566   sheets/detail/salmonella-(non-typhoidal). Accessed

567   4.   Krzysciak P, Chmielarczyk A, Pobiega M, Romaniszyn D, Wojkowska-Mach J. 2017. Acinetobacter

568   baumannii isolated from hospital-acquired infection: biofilm production and drug susceptibility.

569   APMIS 125:1017-1026.

570   5.   Di Lodovico S, Cataldi V, Di Campli E, Ancarani E, Cellini L, Di Giulio M. 2017. Enterococcus hirae

571   biofilm formation on hospital material surfaces and effect of new biocides. Environ Health Prev

572   Med 22:63.

573   6.   Di Domenico EG, Cavallo I, Pontone M, Toma L, Ensoli F. 2017. Biofilm Producing Salmonella

574   Typhi: Chronic Colonization and Development of Gallbladder Cancer. Int J Mol Sci 18.

575   7.   Prouty AM, Schwesinger WH, Gunn JS. 2002. Biofilm formation and interaction with the surfaces

576   of gallstones by Salmonella spp. Infect Immun 70:2640-9.

577   8.   Grantcharova N, Peters V, Monteiro C, Zakikhany K, Romling U. 2010. Bistable expression of

578   CsgD in biofilm development of Salmonella enterica serovar typhimurium. J Bacteriol 192:456-

579   66.

9.   Garai P, Lahiri A, Ghosh D, Chatterjee J, Chakravortty D. 2016. Peptide utilizing carbon starvation

gene yjiY is required for flagella mediated infection caused by Salmonella. Microbiology

162:100-116.

10.   Khil PP, Camerini-Otero RD. 2002. Over 1000 genes are involved in the DNA damage response of

Escherichia coli. Mol Microbiol 44:89-105.

11.   Soufi B, Krug K, Harst A, Macek B. 2015. Characterization of the E. coli proteome and its

modifications during growth and ethanol stress. Front Microbiol 6:103.

12.   Goswami M, Narayana Rao A. 2018. Transcriptome Profiling Reveals Interplay of Multifaceted

Stress Response in Escherichia coli on Exposure to Glutathione and Ciprofloxacin. mSystems 3.

13.   Garai P, Chandra K, Chakravortty D. 2017. Bacterial peptide transporters: Messengers of

nutrition to virulence. Virulence 8:297-309.

14.   Chandra K, Garai P, Chatterjee J, Chakravortty D. 2017. Peptide transporter YjiY influences the

expression of the virulence gene mgtC to regulate biofilm formation in Salmonella. FEMS

Microbiol Lett 364.

15.   Iwase T, Uehara Y, Shinji H, Tajima A, Seo H, Takada K, Agata T, Mizunoe Y. 2010. Staphylococcus

epidermidis Esp inhibits Staphylococcus aureus biofilm formation and nasal colonization. Nature

465:346-9.

16.   Fang K, Jin X, Hong SH. 2018. Probiotic Escherichia coli inhibits biofilm formation of pathogenic

E. coli via extracellular activity of DegP. Sci Rep 8:4939.

17.   Hong SH, Hegde M, Kim J, Wang X, Jayaraman A, Wood TK. 2012. Synthetic quorum-sensing

circuit to control consortial biofilm formation and dispersal in a microfluidic device. Nat

Commun 3:613.

18.   Lee J, Jayaraman A, Wood TK. 2007. Indole is an inter-species biofilm signal mediated by SdiA.

BMC Microbiol 7:42.

19. Jayaraman A, Hallock PJ, Carson RM, Lee CC, Mansfeld FB, Wood TK. 1999. Inhibiting sulfate-
reducing bacteria in biofilms on steel with antimicrobial peptides generated in situ. Appl
Microbiol Biotechnol 52:267-75.

20. Mowat E, Rajendran R, Williams C, McCulloch E, Jones B, Lang S, Ramage G. 2010. Pseudomonas
aeruginosa and their small diffusible extracellular molecules inhibit Aspergillus fumigatus biofilm
formation. FEMS Microbiol Lett 313:96-102.

21. Gnanadhas DP, Elango M, Janardhanraj S, Srinandan CS, Datey A, Strugnell RA, Gopalan J,
Chakravortty D. 2015. Successful treatment of biofilm infections using shock waves combined
with antibiotic therapy. Sci Rep 5:17440.

22. Datey A, Thaha CSA, Patil SR, Gopalan J, Chakravortty D. 2019. Shockwave Therapy Efficiently
Cures Multispecies Chronic Periodontitis in a Humanized Rat Model. Front Bioeng Biotechnol
7:382.

23. Steenackers H, Hermans K, Vanderleyden J, Sigrid CJ, Keersmaecker D. 2012. Salmonella
biofilms: An overview on occurrence, structure, regulation and eradication. Food Res Int 45:502-
531.

24. Yuksel FN, Buzrul S, Akcelik M, Akcelik N. 2018. Inhibition and eradication of Salmonella
Typhimurium biofilm using P22 bacteriophage, EDTA and nisin. Biofouling 34:1046-1054.

25. Todorov SD, de Paula OAL, Camargo AC, Lopes DA, Nero LA. 2018. Combined effect of
bacteriocin produced by Lactobacillus plantarum ST8SH and vancomycin, propolis or EDTA for
controlling biofilm development by Listeria monocytogenes. Rev Argent Microbiol 50:48-55.

26. Gockerman A, Clemmons DR. 1995. Porcine aortic smooth muscle cells secrete a serine protease
for insulin-like growth factor binding protein-2. Circ Res 76:514-21.

27.   Bahgat M, Doenhoff M, Kirschfink M, Ruppel A. 2002. Serine protease and phenoloxidase
activities in hemocytes of Biomphalaria glabrata snails with varying susceptibility to infection
with the parasite Schistosoma mansoni. Parasitol Res 88:489-94.

28.   Papenfort K, Forstner KU, Cong JP, Sharma CM, Bassler BL. 2015. Differential RNA-seq of Vibrio
cholerae identifies the VqmR small RNA as a regulator of biofilm formation. Proc Natl Acad Sci U
S A 112:E766-75.

29.   Janssen KH, Diaz MR, Gode CJ, Wolfgang MC, Yahr TL. 2018. RsmV, a Small Noncoding
Regulatory RNA in Pseudomonas aeruginosa That Sequesters RsmA and RsmF from Target
mRNAs. J Bacteriol 200.

30.   Coughlan LM, Cotter PD, Hill C, Alvarez-Ordonez A. 2016. New Weapons to Fight Old Enemies:
Novel Strategies for the (Bio)control of Bacterial Biofilms in the Food Industry. Front Microbiol
7:1641.

31.   O'Toole GA, Kolter R. 1998. Flagellar and twitching motility are necessary for Pseudomonas
aeruginosa biofilm development. Mol Microbiol 30:295-304.

32.   Ammendola S, D'Amico Y, Chirullo B, Drumo R, Ciavardelli D, Pasquali P, Battistoni A. 2016. Zinc
is required to ensure the expression of flagella and the ability to form biofilms in Salmonella
enterica sv Typhimurium. Metallomics 8:1131-1140.

33.   Haraga A, Ohlson MB, Miller SI. 2008. Salmonellae interplay with host cells. Nat Rev Microbiol
6:53-66.

34.   Lou L, Zhang P, Piao R, Wang Y. 2019. Salmonella Pathogenicity Island 1 (SPI-1) and Its Complex
Regulatory Network. Front Cell Infect Microbiol 9:270.

35.   Fux CA, Costerton JW, Stewart PS, Stoodley P. 2005. Survival strategies of infectious biofilms.
Trends Microbiol 13:34-40.

36.   Fabrega A, Vila J. 2013. Salmonella enterica serovar Typhimurium skills to succeed in the host:

virulence and regulation. Clin Microbiol Rev 26:308-41.

37.   Chen H, Fang Q, Tu Q, Liu C, Yin J, Yin Y, Xia L, Bian X, Zhang Y. 2018. Identification of a contact-

dependent growth inhibition system in the probiotic Escherichia coli Nissle 1917. FEMS

Microbiol Lett 365.

38.   Simm R, Ahmad I, Rhen M, Le Guyon S, Romling U. 2014. Regulation of biofilm formation in

Salmonella enterica serovar Typhimurium. Future Microbiol 9:1261-82.

39.   Desai SK, Padmanabhan A, Harshe S, Zaidel-Bar R, Kenney LJ. 2019. Salmonella biofilms program

innate immunity for persistence in Caenorhabditis elegans. Proc Natl Acad Sci U S A 116:12462-

12467.

40.   Ochman H, Groisman EA. 1994. The origin and evolution of species differences in Escherichia coli

and Salmonella typhimurium. EXS 69:479-93.

41.   Chattopadhyay S, Paul S, Kisiela DI, Linardopoulou EV, Sokurenko EV. 2012. Convergent

molecular evolution of genomic cores in Salmonella enterica and Escherichia coli. J Bacteriol

194:5002-11.

42.   Meysman P, Sanchez-Rodriguez A, Fu Q, Marchal K, Engelen K. 2013. Expression divergence

between Escherichia coli and Salmonella enterica serovar Typhimurium reflects their lifestyles.

Mol Biol Evol 30:1302-14.

43.   Subramaniam S, Muller VS, Hering NA, Mollenkopf H, Becker D, Heroven AK, Dersch P,

Pohlmann A, Tedin K, Porwollik S, McClelland M, Meyer TF, Hunke S. 2019. Contribution of the

Cpx envelope stress system to metabolism and virulence regulation in Salmonella enterica

serovar Typhimurium. PLoS One 14:e0211584.

44.   Saito T, Ueno T, Kubori T, Yamaguchi S, Iino T, Aizawa SI. 1998. Flagellar filament elongation can

be impaired by mutations in the hook protein FlgE of Salmonella typhimurium: a possible role of

the hook as a passage for the anti-sigma factor FlgM. Mol Microbiol 27:1129-39.

45.   Chadsey MS, Karlinsey JE, Hughes KT. 1998. The flagellar anti-sigma factor FlgM actively

dissociates Salmonella typhimurium sigma28 RNA polymerase holoenzyme. Genes Dev 12:3123-

36.

46.   Liang H, Li L, Dong Z, Surette MG, Duan K. 2008. The YebC family protein PA0964 negatively

regulates the Pseudomonas aeruginosa quinolone signal system and pyocyanin production. J

Bacteriol 190:6217-27.

47.   Bossi L, Ratel M, Laurent C, Kerboriou P, Camilli A, Eveno E, Boudvillain M, Figueroa-Bossi N.

2019. NusG prevents transcriptional invasion of H-NS-silenced genes. PLoS Genet 15:e1008425.

48.   Dorman CJ. 2007. H-NS, the genome sentinel. Nat Rev Microbiol 5:157-61.

49.   Free A, Dorman CJ. 1995. Coupling of Escherichia coli hns mRNA levels to DNA synthesis by

autoregulation: implications for growth phase control. Mol Microbiol 18:101-13.

50.   Kobayashi H, Miyamoto T, Hashimoto Y, Kiriki M, Motomatsu A, Honjoh K, Iio M. 2005.

Identification of factors involved in recovery of heat-injured Salmonella Enteritidis. J Food Prot

68:932-41.

51.   Gur E, Biran D, Shechter N, Genevaux P, Georgopoulos C, Ron EZ. 2004. The Escherichia coli DjlA

and CbpA proteins can substitute for DnaJ in DnaK-mediated protein disaggregation. J Bacteriol

186:7236-42.

52.   Doyle SM, Shastry S, Kravats AN, Shih YH, Miot M, Hoskins JR, Stan G, Wickner S. 2015. Interplay

between E. coli DnaK, ClpB and GrpE during protein disaggregation. J Mol Biol 427:312-27.

53.   del Solar GH, de al Campa AG, Perez-Martin J, Choli T, Espinosa M. 1989. Purification and

694   characterization of RepA, a protein involved in the copy number control of plasmid pLS1. Nucleic

695   Acids Res 17:2405-20.

696   54.   Kaplan JB. 2010. Biofilm dispersal: mechanisms, clinical implications, and potential therapeutic

697   uses. J Dent Res 89:205-18.

698   55.   Henderson B, Martin A. 2011. Bacterial virulence in the moonlight: multitasking bacterial

699   moonlighting proteins are virulence determinants in infectious disease. Infect Immun 79:3476-

700   91.

701   56.   Kim YH, Lee Y, Kim S, Yeom J, Yeom S, Seok Kim B, Oh S, Park S, Jeon CO, Park W. 2006. The role

702   of periplasmic antioxidant enzymes (superoxide dismutase and thiol peroxidase) of the Shiga

703   toxin-producing Escherichia coli O157:H7 in the formation of biofilms. Proteomics 6:6181-93.

704   57.   La Carbona S, Sauvageot N, Giard JC, Benachour A, Posteraro B, Auffray Y, Sanguinetti M, Hartke

705   A. 2007. Comparative study of the physiological roles of three peroxidases (NADH peroxidase,

706   Alkyl hydroperoxide reductase and Thiol peroxidase) in oxidative stress response, survival inside

707   macrophages and virulence of Enterococcus faecalis. Mol Microbiol 66:1148-63.

708   58.   Wu Y, Outten FW. 2009. IscR controls iron-dependent biofilm formation in Escherichia coli by

709   regulating type I fimbria expression. J Bacteriol 191:1248-57.

710   59.   Kang D, Kirienko NV. 2018. Interdependence between iron acquisition and biofilm formation in

711   Pseudomonas aeruginosa. J Microbiol 56:449-457.

712   60.   Masse E, Gottesman S. 2002. A small RNA regulates the expression of genes involved in iron

713   metabolism in Escherichia coli. Proc Natl Acad Sci U S A 99:4620-5.

714   61.   Velayudhan J, Castor M, Richardson A, Main-Hester KL, Fang FC. 2007. The role of ferritins in the

715   physiology of Salmonella enterica sv. Typhimurium: a unique role for ferritin B in iron-sulphur

716   cluster repair and virulence. Mol Microbiol 63:1495-507.

Figure Legends

Fig 1. *Salmonella* **biofilm deficient Δ***yjiY* **inhibits biofilm formation by WT strain**

A. Biofilm formation ability of the WT strain after 72 hours of inoculation in presence or absence of WT, Δ*yjiY,* Δ*yjiY:yjiY* and Δ*csgD* culture supernatants was checked. Crystal violet staining was carried out to check the biofilm formed on the solid-liquid interphase (Data are presented as mean $\pm$ SEM of 12 independent experiments). B. Similarly biofilm formation assay was carried out in presence or absence of overnight grown different bacterial strains (co-culture), washed bacterial pellet (cell pellet), cell free supernatant (culture supernatant) and cell lysates (Data are presented as mean $\pm$ SEM of 4 independent experiments). C. Different concentration of the cell free supernatant was used to check the minimum biofilm inhibitory concentration (MBIC) of Δ*yjiY* supernatant (Data are presented as mean $\pm$ SEM of 4 independent experiments). D. Biofilm inhibitory property of the Δ*yjiY* supernatant was checked with supernatant isolated from bacteria grown in shaking condition in LB media and minimal M9 media (Data are presented as mean $\pm$ SEM of 4 independent experiments). Student's t-test was used to analyze the data; p values ****<0.0001, ***<0.001, **<0.01, *<0.05.

Fig 2. *Salmonella Δyjiy* **cell free supernatant significantly reduces the biofilm biomass by reducing cell-cell adhesion**

A. Representative images of biofilm formed on coverslips that were stained with Congo red and imaged using a confocal microscope to generate 3D images and quantify cellulose biomass. Scale is shown on the X- and Y-axes. B. The tensile strength of the biofilm was measured by glass bead assay. Weight of glass beads required to just sink the biofilm to bottom, was plotted

(Data are presented as mean ± SEM of 5 independent experiments). C. Representative scanning

electron micrograph of biofilm formed on a coverslip. Scale bar is 10 µm. D. Cell length of the

biofilm inoculated treated or untreated STM WT cells was measured using ImageJ, and plotted

(Data are presented as mean ± SEM of 1200 cells were measured from 3 independent

experiments). E. Representative confocal images of biofilm cells showing a difference in cell

length. Scale bar is 5 µm (1000-1200 cells were measured from 3 independent experiments for

each treatment). Student's t-test was used to analyze the data; p values ****<0.0001, ***<0.001,

**<0.01, *<0.05.

Fig 3. The physicochemical property of the inhibitory molecules is proteinaceous

A. The cell-free supernatants were treated with different concentrations of EDTA and checked

for their ability to inhibit biofilm formation by STM WT (Data are presented as mean ± SEM of

3 independent experiments). B. The supernatants were treated with proteinase K, and heated to

37 ˚C for 1 hour, and checked for the biofilm inhibitory activity. PMSF was used to inactivate

proteinase K after 1 hour, and was also used as a control (Data are presented as mean ± SEM of 3

independent experiments). C. The supernatants were heated to 65 ˚C and 95 ˚C to check the

thermostability of the active component(s), followed by biofilm inoculation with the treated or

untreated supernatants (Data are presented as mean ± SEM of 3 independent experiments). D.

The pH sensitivity of biofilm inhibitory action of the supernatants were checked. The pH of the

supernatants as well as the biofilm media was made acidic (pH 4.0) or alkaline (pH 9.0) with

concentrated HCl or NaOH, respectively, and checked for the activity of the active molecule(s)

(Data are presented as mean ± SEM of 3 independent experiments). Two-way ANOVA and

Student's t-test were used to analyze the data; p values ****<0.0001, ***<0.001, **<0.01,

*<0.05.

Fig 4. The supernatant interrupts biofilm formation by transcriptional switching during initiation of micro-colony formation

A. Varied concentrations of the STM Δ*yjiY* supernatant were used to pre-formed mature biofilm of STM WT and the ability of the supernatant to disintegrate the biofilm was checked (Data are presented as mean $\pm$ SEM of 5 independent experiments). B. The extent of inhibitory effect of the supernatants were checked by recovery and repeated exposure of the WT strain to the supernatant (Data are presented as mean $\pm$ SEM of 3 independent experiments). C. RNA was isolated from untreated STM WT cells or WT cells treated with the indicated supernatants for 72 hours. RTPCR was done for indicated genes, and C_T values were first normalized to 16s rRNA gene and then normalized to that of untreated WT cells. Y axis values above and below 1 denotes upregulation and downregulation, respectively (Data are presented as mean $\pm$ SEM of 3 independent experiments). (D, E). *In vitro* motility of the treated and untreated STM WT cells were checked on 0.3% swim agar plates or 0.5% swarm agar plates. The radius of the motile zone was imaged after different time, measured using ImageJ and plotted (Data are presented as mean $\pm$ SEM of 4 independent experiments). F. The flagellar status of the treated or untreated cells (approximately 50 cells in each samples, in two different experiments) was imaged using scanning electron microscope. White arrows show the flagella. Scale bar is 500 nm. Two-way ANOVA and Student's t-test were used to analyze the data; p values ****<0.0001, ***<0.001, **<0.01, *<0.05.

Fig 5. STM Δ*yjiY* supernatant treatment impairs virulence of STM WT

A. *In vitro* invasiveness of the supernatant treated WT bacteria was checked by infecting adherent human epithelial Int407 cell line and percent invasion was plotted. Log phase WT culture was used as a control (Data are presented as mean $\pm$ SEM of 3 independent experiments).

B. Survival of intracellular *Salmonella* in Int407 cells was checked by gentamicin protection assay. The intracellular bacterial fold proliferation was measured as the ratio of intracellular bacterial CFU after 16 hours to the CFU at 2 hours (Data are presented as mean $\pm$ SEM of 3 independent experiments). C. Percent gut colonization was measured by measuring the diameters of the colonized gut and total body width after feeding the worms continuously for 12 hours with mCherry-*E.coli* OP50 or supernatant treated/untreated mCherry-STM WT (approximately total 25 worms were measured in 2 independent experiments). D. Representative images of 3 days supernatant treated bacterial colonization in young adult *C. elegans* N2 worms. Bacterial colonization in the gut lumen was checked after 12 hours of continuous STM WT feeding and/or 12 hours STM WT feeding, followed by 12 hours *E. coli* OP50 feeding (approximately total 25 worms were imaged in 2 independent experiments). Scale bar is 20 µm. E. Bacterial CFU from infected *C. elegans* was determined by lysing 10 worms (in triplicates) in M9 buffer followed by dilution plating (Data are presented as mean $\pm$ SEM of 2 independent experiments). Student's t-test was used to analyze the data; p value *<0.05. F. Biofilm inhibitory effect of STM $\Delta yjiY$ supernatant on few other enterobacteriaceae pathogens was checked by inoculation the strains with 1% v/v supernatants (Ec- *Escherichia coli* DH5α, Pa01- *Pseudomonas aeruginosa* strain PA01, Kpn- *Klebsiella pneumoniae*, Sa- *Staphylococcus aureus,* Data are presented as mean $\pm$ - SEM of 3 independent experiments). Two-way ANOVA and Student's t-test were used to analyze the data; p values ****<0.0001, ***<0.001, **<0.01, *<0.05.

Fig 6. Global transcriptional regulator HNS along with NusG causes major transcriptional regulation leading to biofilm inhibition and motility inhibition

A. SDS PAGE of the STM WT and STM $\Delta yjiY$ supernatant showing presence of multiple differential bands in the supernatants. B. Venn diagram showing proteins present in the

supernatants detected in LC QTOF MS/MS analysis, blue and red circles represent number of

proteins found in STM WT and Δ*yjiY* culture supernatants, respectively. C. Biofilm formation

ability of the WT strain after 72 hours of inoculation in presence of Δ*yjiY* culture supernatant and

200ng ecotin/HNS/NusG fractions was checked. Crystal violet staining was carried out to check

the biofilm formed on the solid-liquid interphase (Data are presented as mean $\pm$ SEM of 3

independent experiments; One-way was used to analyze the data; p values ****<0.0001,

***<0.001). D. Proposed model of biofilm inhibition by Δ*yjiY* supernatant. The model depicts

that transcription factors, proteases, QS-quenching molecules, and protease inhibitors are

secreted actively or released during cell lysis in Δ*yjiY* supernatant, which upon entering into the

WT cells by either active transport or membrane diffusion, regulates biofilm formation and

motility.

A % invasion — UT WT, WT + WT sup, WT + ΔyjiY sup, Log phase WT; * , ***

B Fold Change — WT control, WT + WT sup, WT + ΔyjiY sup, Log phase WT; **, *, *

C % Gut colonization — *E. coli* OP50, WT+ WT sup, WT+ ΔyjiY sup, WT untreated; ****, ****, ****

D 12hr RFP-STM continuous feeding | 12hr STM-RFP followed by 12hr OP50

STM WT+ WT sup; STM WT+ ΔyjiY sup; STM WT untreated

E ■ 12 hr STM □ 12 hr STM+12hr OP50 — CFU/worm; WT Untreated, WT+WT sup, WT+ΔyjiY sup; *, *

F ■ Untreated ■ STM WT sup ■ STM ΔyjiY sup — OD_{595}; Ec (ns, ****), Pa01 (ns, ns), Kpn (ns, ns), Sa (ns, ns)

Swimming motility

	Wt+WT sup	WT+Δ*yjiY* sup	WT untreated	O/N WT	STM Δ*fliC*
2 hr					
8 hr					

Swarming motility

	Wt+WT sup	WT+Δ*yjiY* sup	WT untreated	O/N WT	STM Δ*fliC*
2 hr					
8 hr					

A

B

fliC expression

■ WT untreated   ▨ WT + Δyjiy sup

Fold change/16s

Time Post Biofilm Inoculation

C

Overnight WT   WT+ WT sup   WT untreated   WT+ Δ*yjiY* sup   STM Δ*fljB* Δ*fliC*

A

Protein Name	Fraction Name	Pooled Fractions	Elution Volume (ml)	Oligomeric State
Ecotin	E1	A6, A7, A8	8-9	Homodimer
	E2	B1, B2, B3, B4	10-11	
	E3	B5, B6, B7	12-13	Monomer
	E4	C2, C3	14-15	
HNS	H1	A6, A7, A8, B1	8-9	Heterotrimer (Hha)
	H3	B3, B4	11	Homodimer
	H4	B5, B6	12-13	
	H5	B7, B8	13-14	Monomer
	H6	C4, C5, C7	15-17	
NusG	N1	B2	10-10.5	Monomer
	N2	B3	10.5-11	
	N3	B4	11-11.5	
	N4	B6, B7, B8	12-13.5	
	N5	C1, C2	13.5-14.5	

B