Age structuring and spatial heterogeneity in prion protein gene (*PRNP*) polymorphism in white-tailed deer

Tyler K. Chafin[1], Marlis R. Douglas[1], Bradley T. Martin[1], Zachery D. Zbinden[1], Christopher R. Middaugh[2], Jennifer R. Ballard[2], M. Cory Gray[2], Don White Jr.[3], Michael E. Douglas[1],

[1]Department of Biological Sciences, University of Arkansas, Fayetteville, Arkansas 72701, USA; Email: (TKC) tkchafin@uark.edu; (BTM) btm002@uark.edu; (ZDZ) zdzbinde@uark.edu; (MED) med1@uark.edu; (MRD) mrd1@uark.edu

[2]Arkansas Game and Fish Commission, Research, Evaluation, and Compliance Division, Little Rock, Arkansas 72205, USA; Email: (CRM) christopher.middaugh@agfc.ar.gov; (JRB) jennifer.ballard@agfc.ar.gov; (MCG) cory.gray@agfc.ar.gov

[3]University of Arkansas Agricultural Experiment Station, Monticello, Arkansas 71656, USA; Email: whited@uamont.edu

Running head: *PRNP polymorphism in white-tailed deer in Arkansas*

ABSTRACT

2   Chronic-wasting disease (CWD) is a prion-derived fatal neurodegenerative disease that has

affected wild cervid populations on a global scale. Susceptibility has been linked unambiguously

4   to several amino acid variants within the prion protein gene (*PRNP*). Quantifying their

distribution across landscapes can provide critical information for agencies attempting to

6   adaptively manage CWD. Here we attempt to further define management implications of *PRNP*

polymorphism by quantifying the contemporary geographic distribution (i.e., phylogeography) of

8   *PRNP* variants in hunter-harvested white-tailed deer (WTD; *Odocoileus virginianus,* N=1433)

distributed across Arkansas (USA), including a focal spot for CWD since detection of the disease

10   in February 2016. Of these, *PRNP* variants associated with the well-characterized 96S non-

synonymous substitution showed a significant increase in relative frequency among older CWD-

12   positive cohorts. We interpreted this pattern as reflective of a longer life expectancy for 96S

genotypes in a CWD-endemic region, suggesting either decreased probabilities of infection or

14   reduced disease progression. Other variants showing statistical signatures of potential increased

susceptibility, however, seemingly do so as an artefact of population structure. We also showed

16   marked heterogeneity across the landscape in the prevalence of 'reduced susceptibility'

genotypes. This may indicate, in turn, that differences in disease susceptibility among WTD in

18   Arkansas are an innate, population-level characteristic that is detectable through

phylogeographic analysis.

KEYWORDS Adaptive management; chronic-wasting disease; demography; disease

22   susceptibility; haplotype; selection; white-tailed deer

Introduction

Chronic wasting disease (CWD) is a fatal neurodegenerative disorder that affects white-tailed deer (WTD; *Odocoileus virginianus*) and related cervids[1,2], with severe impacts on native wildlife, that also reverberate economically for recreational hunting and ancillary commercial enterprises[3,4]. Most CWD eradication efforts have proven unsuccessful thus far[5], leading to its continued spread and increased prevalence[6]. Given this, wildlife managers in many jurisdictions are responding by shifting long-term goals away from eradication and instead towards suppression, containment, and mitigation[7,8].

Several factors have impeded the eradication of CWD, including aspects of life history in both host and agent, as well as limited knowledge with regards to how these interact with environment to define CWD epidemiology. Pathogenicity and transmission, for example, occur via the structural transformation of a naturally occurring cellular prion protein (PrP^C) into a misfolded "pathogenic" isoform (PrP^{SC})[9]. The efficiency with which this occurs, coupled with an extensive incubation period[10], serve to confound proactive surveillance and management. Additionally, both vertical[11,12] and horizontal transmission[13,14] are seemingly involved, with prion persistence well established within "environmental reservoirs"[15–17]. As a result, surveillance and monitoring are being increasingly used by state agencies to inform harvest and selective-removal based management strategies to suppress the disease where it cannot be eradicated[18,19]. This is complicated particularly due to a potential for long-distance host dispersal[20–22], and prion 'shedding' by individuals with subclinical infections[23,24].

Environmental factors that act to reduce disease spread are of particular interest. For example, the capacity of soil as a reservoir for extra-corporeal prion persistence may hinge upon its composition[25]. Likewise, environmental factors that enhance the potential for WTD

movements also may modulate CWD transmission among herds[26,27]. Considerable effort to date

48   has focused on characterizing intrinsic susceptibility, especially with regard to the quantification

of genetic polymorphisms that encode PrP^C ($PRNP$)[28–31]. One consistent approach is to identify

50   those $PRNP$ gene variants that are associated with reduced CWD susceptibility[32,33]. The amino

acid composition of the resulting protein is thus assumed to impact disease progression[34],

52   although the exact mechanism remains unclear.

Consistent among those variants reported to be associated with reduced CWD

54   susceptibility is a non-synonymous mutation corresponding to an amino acid substitution at

position 96 (i.e., from glycine to serine; hereafter 96G versus 96S). Inoculation studies

56   employing this mutation have successfully delayed the progression of CWD in both WTD and

proxies[13,35]. Our primary interest is to characterize if (and how) these $PRNP$ alleles vary

58   spatially[30,36], as one landscape-level axis that depicts a "resistance" to CWD spread. We provide

such an analysis by employing WTD sampled state-wide as a basis for the phylogeography of the

60    $PRNP$ gene (i.e., the geographic distribution of individuals associated with a gene genealogy[37]).

We then examine spatial and age-structured patterns within this phylogeographic framework to

62   ask several questions regarding the role of $PRNP$ variants on population dynamics in CWD-

endemic areas: Do 'reduced susceptibility' variants have an effect on survivorship (e.g., as one

64   might expect if $PRNP$ polymorphisms do indeed drive susceptibility)? If so, does this leave a

detectable signature of biased fitness (e.g., as a result of increased survivability and thus

66   potential reproductive output)? What are the impacts of $PRNP$ polymorphism on population

demographics? We address these questions at two spatial scales: 1) within a dense sampling of

68   the CWD-focal area (Newton County, Arkansas) from which prevalence, and, presumably,

measurable impacts on population demography are highest; and 2) state-wide, where sampling

70   densities were lower, and in many areas within which CWD has not yet been detected (though

we note that lack of detection does not mean lack of occurrence). The county-level spatial scale

72   allowed us to examine demographic impacts within a recently detected outbreak in a novel area

(Newton County), wherein dense sampling and high prevalence allow sufficient sampling for

74   testing hypotheses of age structuring and selection on *PRNP* polymorphisms, while the state-

level spatial scale allowed broad-scale analyses of heterogeneity and phylogeographic

76   structuring. The combination of these two spatial scales allows for superimposing differential

susceptibilities and fitness across a CWD-absent landscape which could in turn facilitate the

78   creation of management scenarios to project and potentially mitigate disease spread.

80   **Results**

Data generation

82   From 2016-2019, ear and tongue tissue samples were collected from 1,720 harvested WTD

across 75 counties in Arkansas (Supplemental Table 1; Figure 1). Subsequently, tissue samples

84   from 1,460 WTD were amplified and sequenced, yielding ~800 nucleotides of the *PRNP* gene

and *PRNP^{psg}* pseudogene. From these data we obtained 1,433 sequences, of which 316 were

86   obtained from Newton County, the CWD focal area. Sequences were trimmed to 720

unambiguously scored nucleotides, with 11 sites found to be polymorphic (Table 1). Three

88   previously reported polymorphic sites (i.e., nt285, nt299 and nt372; Brandt et al., 2015, 2018)

were found to be invariable in our data, whereas one additional site had a novel synonymous

90   substitution (nt499, A/C). Three sites (i.e., nt286, nt367 and nt676) also reflected non-

synonymous substitutions, corresponding to amino acid substitutions 96S, 122T and 255K,

92   respectively.

Haplotype phasing resulted in 20 distinct alleles (Table 1; Figure 2), 4 of which are

94   novel, and 16 were previously documented in other states[29–31]. Two Arkansas haplotypes (AR 2

and AR 3) share the synonymous substitution 96S (nt286/A) associated with reduced CWD

96   susceptibility (i.e., Haplotype C[30]). Three others (I, P and V) also share the 96S amino acid

alteration (Table 1; Figure 2). However, all five were at low frequencies (<1%, except Haplotype

98   I at 1.47%; Table 2), and thus were excluded from tests of CWD association.

Targeted amplification and sequencing of the *PRNP^PSG* pseudogene was successful in

100   30% of our samples (443 of 1,459). Our comparison of the 443 *PRNP* and *PRNP^PSG* haplotypes

indicated that variability at site nt413 did not represent a biological variant of the *PRNP* gene but

102   was instead a non-targeted amplification of the *PRNP^PSG* pseudogene.

104   ***PRNP haplotype frequencies***

Haplotype frequencies (Tables 2 and S2) differed slightly from those reported in other states

106   (e.g., Wisconsin and Illinois[30,31]). The four most frequent haplotypes in Illinois and Wisconsin

(i.e., <10%), were also common among Arkansas samples (Figure 3; Haplotypes A-D). We also

108   found that Haplotype A, most common in Illinois/Wisconsin at 30%, occurred in only 15% of

our samples (Figure 3; Table 2). Haplotypes B and D were instead most common in our data

110   (each at ~23%). Haplotype C, associated with reduced CWD susceptibility, was detected at a

frequency of 15%, quite similar to that in Illinois/Wisconsin (17%). Two additional haplotypes

112   (E and G) also occurred at high frequencies in Arkansas (7% and 11%, respectively), whereas

they were found at <5% in the Illinois/Wisconsin[30]. Out of the 16 rare haplotypes with

114   previously reported frequencies ≤1% (Haplotypes K-Z[30]), seven were also observed at low

frequencies in Arkansas (i.e., Haplotypes K, L, O, P, R, T, and V). Haplotype frequencies also

116   differed across counties (Supplemental Table 1), although we note that sampling effort was

uneven across the state.

Evidence for disease-mediated selection on PRNP variants

120   Haplotypes B and C did not occur in the same frequency within CWD-positive (N = 432) and

CWD-negative deer (N = 196; Table 2; Figure 3). Haplotype B was over-represented within

122   CWD-positive deer, both state-wide (OR = 2.00, p = 0.000001), and in Newton County (OR =

1.43, p = 0.033). Haplotype C was under-represented within CWD-positive deer, both state-wide

124   (OR = 0.30, p = 0.00003) and Newton County (OR = 0.42, p = 0.015).

If Haplotypes B and C influence CWD susceptibility, then their relative frequency of

126   occurrence should vary among deer age-classes, reflecting a biased probability of reaching older

age classes. We restricted our analyses to Newton County to ensure only deer from CWD-

128   endemic locations were considered. We found the frequency of Haplotype C significantly more

common in older deer, both when measured as relative allele frequency (p=0.036; R^2=0.635;

130   Figure 4), and as an age-partitioned odds ratio (p=0.043; R^2=0.601; Figure 5). The increase in

Haplotype C remained substantial even when we considered only the relative frequency of each

132   haplotype among CWD-positive deer from Newton County (i.e., 4% of yearling/ fawn

haplotypes, and 17% of those from individuals older than 5; Supplemental Table 3). Among

134   CWD-positive deer state-wide, Haplotype C was recorded at 5.24% of all sampled haplotypes

(Table 2). By contrast, Haplotype B showed neither a discernible relationship regarding CWD

136   status, nor relative frequencies across age classes (Figure 5; Supplemental Table 3).

Given the presence of age-structuring in Haplotype C, we then hypothesized a fitness

138   differential associated with polymorphism at AA position 96. We first examined phylogenetic

conservation and stability of non-synonymous mutations in Arkansas (Table 1) and predicted

140   subsequent effects on tertiary protein structure. We found the 96S substitution to be the least

conserved of the three amino acid positions, with none of the observed substitutions significantly

142   altering tertiary shape (Supplemental Table 3). We thus hypothesized that positive selection on

96S was driving the observed patterns. As a test, we treated cohorts in Newton County as a time-

144   series and computed the magnitude of selection required to generate the observed positive

growth in frequency with increasing age (Figure 5). Our results yielded a selection coefficient (*s*)

146   of 0.1215. We note that numerous caveats are associated with population genetic assumptions

behind this calculation, as well as of the interpretation of selection coefficients[38], the result

148   seemingly supports our observations.

150   **Discussion**

Our diagnosis of variability in the *PRNP* gene across 1,433 WTD collected from 75 counties in

152   Arkansas was comparable to that found in other states (Tables 1-2). Of the 20 haplotypes we

detected, 16 were previously identified in other states, with those alleles at higher frequencies in

154   Arkansas also being most common elsewhere[30]. Four novel variants were found at low

frequencies (Table 2; Figure 2-3), with paralog artefacts as a source of this variation eliminated

156   due to our sequencing of the pseudogene. Haplotype C, characterized by the 96S substitution,

showed a significantly biased representation among CWD-positive deer, being under-represented

158   in younger deer and over-represented in older CWD-positive deer (Figure 5). This suggests that

96S-individuals tend to live longer following CWD-infection than those with alternate

160   genotypes. This could reflect either a reduced likelihood of contracting the disease or a slower

disease progression following exposure.

162   We also found Haplotype B as significantly over-represented in CWD-positive deer

(Table 2 and Figure 3) but failed to find a reciprocal impact on life expectancy (Figure 5).

164   Instead, we found Haplotype B to be most extensive within the region where CWD is currently

centered (Figure 4). This, in turn, may suggest heterogeneity in Haplotype B frequency

166   represents an artefact of population structure, a well-known confounding variable recognized in

trait-association studies[39–42]. We also note that decreased precision of aging of white-tailed deer

168   on the basis of tooth development and wear patterns may also contribute some variability,

particularly with regard to haplotype frequencies among older cohorts[43–45]. We also cannot rule

170   out the potential for linked variation as influencing the observed pattern in Haplotype B,

however our results do not find any evidence for increased susceptibility of the variant. Using

172   panels of nuclear markers, we suggest further study of patterns of spatial connectivity and

population structure in wild WTD[46] as a means of separating potential spatial and

174   phylogeographic drivers of haplotype frequencies from those driven by disease-mediated

selection.

Management implications of PRNP variation

178   Structural variants of the *PrP* protein play a role in disease progression[28,34]. However, the exact

mechanisms remain poorly understood. The primary variant we have implicated herein (i.e., 96S)

180   as influencing disease susceptibility has been supported as such in laboratory settings. For

example, Mathiason et al.[23] inoculated WTD with prion strains and examined time-to-detection

182   (via saliva) across deer in multiple cohorts. Several infected individuals having the 96S prion

gene variant remained undetected at 18 months post-inoculation, although this may represent an

184   insufficient time for the necessary *in vivo* prion protein build-up. Race et al.[35] similarly

inoculated transgenic mice expressing different white-tailed deer 96GG and 96SS *PRNP*

186   genotypes and showed that this delay in disease progression also extended to heterozygotes

(96GS genotype).

188   Despite compelling evidence for an inhibitory mechanism at the 96S allele, its use as a

management tool remains unclear. Genetically-guided selective breeding in domestic sheep

190   (*Ovis aries*) has reduced scrapie incidence [47,48]. In a captive setting such an approach may be

viable for WTD[35], although we note that the degree of protection offered by the 96S genotype to

192   CWD is likely much lower than that seen among the most 'protective' genetic variants to scrapie

in sheep[47]. However, captive deer maintained at high density and under genetic selection could

194   also drive artificial selection for emergent prion strains with heightened pathogenicity in 96S

deer, as well as potentially expanding the host range to include novel species[49].

196   The implication of 96S frequency with regard to the spread of CWD within and among

herds is likewise uncertain. An important question is whether the potential for an increased

198   incubation period [50] associated with 96S could also produce a similar extended period for

asymptomatic and subclinical transmission/prion shedding[24,51]. If so, this could increase the

200   probability of transmission by 96S deer, thus promoting its increase within populations (Figure

6). An understanding of the of prion growth kinetics across host genotypes is needed, as well as a

202   more thorough understanding of prion strain evolution[52].

204   ***Conclusion***

Our results corroborate previous research conducted with WTD in Illinois and Wisconsin:

206   reduced CWD susceptibility in *PRNP* variants associated with the non-synonymous 96S

mutation[30,31]. We demonstrated that a common haplotype (Haplotype C) harboring 96S increased

208   in relative frequency when older CWD-positive cohorts were examined.

Management implications of this research requires further epidemiological understanding

210   necessary to predict outcomes (Figure 6). The next step in understanding the current distribution

and future spread of CWD in Arkansas requires the characterization of context-specific factors,

212   including 1) population structure as a potential driver of *PRNP* trends, 2) landscape features that

modulate deer dispersal, population demography, and density, and 3) increased knowledge of

214   epidemiology, including the interactions among context-specific factors[53]. We thus advocate for

a landscape genomic framework for WTD as the next logical step to characterize CWD spread

216   such that fine-scaled deer movement patterns can be more effectively parsed and interpreted[54].

218   **Materials and Methods**

*CWD surveillance and prevalence in Arkansa*s

220   In October 2015, CWD was initially detected in Arkansas in a 2.5-year old female elk (*Cervus*

canadensis) legally harvested near Pruitt in Newton County. In February 2016, a CWD-positive

222   female WTD was found dead in Ponca in Newton County. During March 2016, biologists from

the Arkansas Game and Fish Commission (AGFC) collected 266 WTD tissue samples within a

224   50,500 ha focal region. CWD prevalence was 23% and differed by gender (female = 20%, male

= 32%). For these surveys, to include the samples collected for molecular work, aging of deer

226   was done by examination of tooth replacement and wear[55].

Subsequent state-wide monitoring, which included hunter-harvested and road-killed deer,

228   identified CWD-positive individuals outside of the initial focal region. Additional state-wide

sampling efforts, in conjunction with hunter harvested and bi-annual surveys, established a state-

230   wide baseline for occurrence of CWD. As of April 2020, 818 WTD and 23 elk have tested

positive for CWD. Given the incidence of confirmed CWD-positive deer, AGFC established a

232   CWD Management Zone (CWDMZ) that included counties within a 16km radius of identified

positives. At the completion of this study (June 2019), the CWDMZ encompassed 19 counties of

234   northwestern Arkansas (Figure 1).

There has been a concerted effort by the AGFC to proactively manage CWD prevalence

236   and potential disease spread in Arkansas, which included hunting regulations that promote

harvest of young male deer and increased harvest of female deer (e.g., removal of antler point

238   restrictions and altered bag limits). Addition restrictions included prohibiting baiting and feeding

to reduce grouping behavior in deer and to hinder human-mediated transmission via hunting and

240   subsequent removal of carcasses from with the CWDMZ. The CWDMZ has subsequently been

expanded to encompass the known distribution of CWD. During the 2018/2019 deer and elk

242   hunting seasons, 246 additional CWD-positive cervids (241 WTD and five elk) were detected.

Moreover, the AGFC has mandated a compulsory testing requirement for harvested elk, and a

244   voluntary test for WTD, facilitated by a state-wide network of deer head drop-off locations.

246   ***DNA Extraction and amplification***

Frozen ear or tongue tissue was homogenized with the TissueLyser II (QIAGEN[©] Corporation,

248   Maryland, USA), with genomic DNA subsequently extracted using the QIAamp Fast DNA

Tissue Kit protocol (QIAGEN[©] Corporation, Maryland, USA). To ascertain presence of high-

250   quality genomic DNA (i.e., molecular weight >20kb), a 5µl aliquot of the DNA extract was

separated on a 2% agarose gel and visualized using GelGreen on a blue-light transilluminator

252   (Gel Doc™ EZ Imager; Bio-Rad). DNA was then used as template material to amplify a coding

section of the *PRNP* gene, following a protocol modified from previous studies[31,32]. For the

254 functional *PRNP* gene, the forward primer (CWD-13) straddles Intron 2 and Exon 3, with the

reverse primer (CWD-LA) located 850bp downstream[32]. To ascertain if the polymorphisms were

256 indeed in the functional *PRNP* gene, we tested for the presence of the non-coding *PRNP*

pseudogene (*PRNP^{PSG}*) by using pseudogene primers 223 and 224[33].

258   Amplifications for the functional *PRNP* gene and the *PRNP^{PSG}* pseudogene were

performed in 20µl reactions consisting of 10µl Qiagen HotStart Master Mix (1unit HotStartTaq

260 DNA Polymerase, PCR Buffer with 3mM MgCl$_2$, and 400µM of each dNTP), 8µM each of the

forward and reverse primer, 7.4µl DNase-free water, and 1µl of template DNA (~50-100ng).

262 Thermocyling protocols consisted of an initial denaturation step of 15min at 95^{o}C, followed by

10 cycles of 45s denaturation at 95^{o}C, 45s annealing at 57^{o}C, and 75s extension at 72^{o}C, 25

264 cycles of 30s denaturation at 95^{o}C, 30s annealing at 55^{o}C and 60s extension at 72^{o}C, completed

with a final extension step of 5min at 72^{o}C.

266   If both *PRNP* and *PRNP^{PSG}* amplified in a sample, then each was sequenced to identify

the true polymorphism in the functional *PRNP* gene. Amplicons were enzymatically purified,

268 sequenced using BigDye v. 3.1 (Applied Biosystem Inc., Forest City, CA) dye-terminator

chemistry, and resolved on an ABI 3730XL GeneAnalyzer (University of Illinois Keck Center

270 for Functional and Comparative Genomics). Sequences were manually edited using Sequencher

(v 5.4, Gene Codes, Ann Arbor MI) and aligned against a reference database of *PRNP* gene

272 sequences obtained from the NCBI GenBank (Accession # AF156185.1; AY3600089.1;

AY3600091.1).

Haplotype phasing and network construction

276 Following alignment, sequences were phased to haplotypes (paired nuclear alleles) using the

program PHASE2[56], which reconstructs haplotypes using a probabilistic model of linkage

278   disequilibrium. Only haplotypes assigned with >90% posterior probability (N=1,433) were

retained. Scripts employed to format inputs and parse haplotype phasing are available at

280   *github.com/tkchafin/haploTools*. Haplotypes were then categorized using a published

nomenclature[31], with haplotype frequencies calculated globally, by county, and by CWD status.

282   We constructed a haplotype network to visualize similarity amongst haplotype, using the

median-joining algorithm employed by POPART[57]. Scripts to formulate these input files are found

284   at: *github.com/tkchafin/scripts*.

286   ***Analysis of PRNP variants***

We first applied spatial interpolation to examine the structure of *PRNP* haplotypes as distributed

288   across the state. Haplotype frequencies were computed by first dividing the state into non-

overlapping 'pseudo-populations' that contained between 5-10 sampling localities each. This

290   was done because our state-wide sampling process lacked *a priori* information with regard to

natural population structure. Our results yielded N=211 polygons (Supplemental Figure 1). We

292   then applied Empirical Bayesian Kriging in ARCMAP v10.7.1 (Esri, Inc.) to interpolate our

posterior probabilities.

294   To associate disease with *PRNP* variants, we followed prior studies[30,31] by computing

odds-ratios (OR). We first consider the probability of displaying an outcome (=CWD status)

296   given the presence of a focal haplotype. An OR>1 = an over-representation of the focal

haplotype among CWD-positive deer; an OR<1 indicates the focal haplotype is under-

298   represented. We identified and evaluated haplotypes separately, by examining haplotype

frequencies among age classes. We did so because a bias in relative representation can be driven

by multiple factors such as population structure, which may drive a coincident relationship. Here, we assumed that if a haplotype indeed affects the probability of survival to adulthood in CWD-

present regions, presumably by reducing disease risk and/or progression, then it should show a significant change in relative representation in older age groups.

We then computed a selection coefficient (s) as a relative fitness value. We first derived counts for each haplotype within each age class, then sampled across age classes as a time-series

sample. We then estimated selection by examining changes in allele frequencies across this series, relative to that expected via stochastic change[58]. Finally, we modelled the effects non-

synonymous mutations on protein conformation so as to test if they do indeed alter PrP^{SC} tertiary structure (as opposed to simply being linked to un-sequenced variation). To understand amino

acid sequence conservation and possible deleterious effects at each non-synonymous position, we utilized two alignment-based algorithms, PROVEAN[59] and SIFT[60]. We also predicted protein

stability using the support vector machine (SM)-based I-Mutant2.0 tool[61].

Acknowledgments

The authors thank hunters and wildlife managers statewide who collected WTD tissue samples.

316   We give particular thanks to A.J. Riggs for helping to coordinate sample collection. We also thank former students and technicians P. McDill and L. Graham for contributions to study

318   design, sample handling and organization, and DNA extraction.

320   **Disclosure statement**

No potential conflict of interest is reported by the authors.

Funding

324   This project and the preparation of this publication was funded in part by the Pittman-

Robertson Wildlife Restoration Grant Program (Grant #AR-W-F18AF00020) of the U.S.

326   Fish and Wildlife Service through an agreement with the Arkansas Game and Fish

Commission. The views and opinions expressed herein are those of the authors and do not

328   necessarily reflect the views or policies of the Arkansas Game and Fish Commission. Product

references do not constitute endorsements.

ORCID

332   Tyler K. Chafin; https://orcid.org/0000-0001-8687-5905

Michael E. Douglas; https://orcid.org/0000-0001-9670-7825

336   **References**

1.   Williams ES, Young S. Spongiform encephalopathies in Cervidae. Rev Sci Tech 1992;
338   11:551–67.

2.   Mawdsley JR. Phylogenetic patterns suggest broad susceptibility to chronic wasting
340   disease across Cervidae. Wildl Soc Bull 2020; 44:152–5.

3.   Bishop RC. The economic impacts of chronic wasting disease (CWD) in Wisconsin. Hum
342   Dimens Wildl 2004; 9:181–92.

4.   Arnot C, Laate E, Unterschultz J, Adamowicz W. Chronic wasting disease (CWD)
344   potential economic impact on cervid farming in Alberta. J Toxicol Environ Heal - Part A
        Curr Issues 2009; 72:1014–7.

346   5.   Wasserberg G, Osnas EE, Rolley RE, Samuel MD. Host culling as an adaptive
        management tool for chronic wasting disease in white-tailed deer: A modelling study. J
348   Appl Ecol 2009; 46:457–66.

6.   Rivera NA, Brandt AL, Novakofski JE, Mateus-Pinilla NE. Chronic wasting disease in

350   cervids: Prevalence, impact and management strategies. Vet Med Res Reports 2019; 10:123–39.

352   7.   Uehlinger FD, Johnston AC, Bollinger TK, Waldner CL. Systematic review of management strategies to control chronic wasting disease in wild deer populations in
354   North America. BMC Vet Res 2016; 12.

  8.   Miller MW, Fischer JR. The first five (or more) decades of chronic wasting disease:
Lessons for the five decades to come. Trans North Am Wildl Nat Resour Conf 2016; 81:1–12.

9.   Aguzzi A, Sigurdson C, Heikenwaelder M. Molecular mechanisms of prion pathogenesis. Annu Rev Pathol Mech Dis 2008; 3:11–40.

10.   Miller MW, Wild MA, Williams ES. Epidemiology of chronic wasting disease in captive Rocky Mountain elk. J Wildl Dis 1998; 34:532–8.

11.   Selariu A, Powers JG, Nalls A, Brandhuber M, Mayfield A, Fullaway S, Wyckoff CA, Goldmann W, Zabel MM, Wild MA, et al. In utero transmission and tissue distribution of
chronic wasting disease-associated prions in free-ranging Rocky Mountain elk. J Gen Virol 2015; 96:3444–55.

12.   Nalls A V., McNulty E, Powers J, Seelig DM, Hoover C, Haley NJ, Hayes-Klug J, Anderson K, Stewart P, Goldmann W, et al. Mother to offspring transmission of chronic
wasting disease in Reeves' Muntjac deer. PLoS One 2013; 8:e71844.

  13.   Mathiason CK, Powers JG, Dahmes SJ, Osborn DA, Miller K V., Warren RJ, Mason GL,
Hays SA, Hayes-Klug J, Seelig DM, et al. Infectious prions in the saliva and blood of deer with chronic wasting disease. Science 2006; 314:133–6.

14.   Potapov A, Merrill E, Pybus M, Coltman D, Lewis MA. Chronic wasting disease: Possible transmission mechanisms in deer. Ecol Modell 2013; 250:244–57.

15.   Gough KC, Maddison BC. Prion transmission: Prion excretion and occurrence in the environment. Prion 2010; 4:275–82.

16.   Johnson CJ, Phillips KE, Schramm PT, McKenzie D, Aiken JM, Pedersen JA. Prions adhere to soil minerals and remain infectious. PLoS Pathog 2006; 2:e32.

17.   Wiggins RC. Prion stability and infectivity in the environment. Neurochem Res 2009; 34:158–68.

18.   Potapov A, Merrill E, Pybus M, Lewis MA. Chronic wasting disease: Transmission mechanisms and the possibility of harvest management. PLoS One 2016; 11:1–20.

19.   Manjerovic MB, Green ML, Mateus-Pinilla N, Novakofski J. The importance of localized culling in stabilizing chronic wasting disease prevalence in white-tailed deer populations.
Prev Vet Med [Internet] 2014; 113:139–45. Available from:

http://dx.doi.org/10.1016/j.prevetmed.2013.09.011

386   20.   McCoy JE, Hewitt DG, Bryant FC. Dispersal by yearling male white-tailed deer and implications for management. J Wildl Manage 2005; 69:366–76.

388   21.   Long ES, Diefenbach DR, Rosenberry CS, Wallingford BD. Multiple proximate and ultimate causes of natal dispersal in white-tailed deer. Behav Ecol 2008; 19:1235–42.

390   22.   Kelly AC, Mateus-Pinilla NE, Douglas M, Douglas M, Brown W, Ruiz MO, Killefer J, Shelton P, Beissel T, Novakofski J. Utilizing disease surveillance to examine gene flow
392   and dispersal in white-tailed deer. J Appl Ecol 2010; 47:1189–98.

        23.   Mathiason CK, Hays SA, Powers J, Hayes-Klug J, Langenberg J, Dahmes SJ, Osborn DA,
394   Miller K V., Warren RJ, Mason GL, et al. Infectious prions in pre-clinical deer and transmission of chronic wasting disease solely by environmental exposure. PLoS One
396   2009; 4.

        24.   Tamguney G, Miller MW, Wolfe LL, Sirochman TM, Glidden D V., Palmer C, Lemus A,
398   DeArmond SJ, Prusiner SB. Asymptomatic deer extrete infectious prions in feces. Nature 2009; 461:529–32.

400   25.   Saunders SE, Bartz JC, Bartelt-Hunt SL. Soil-mediated prion transmission: Is local soil-type a key determinant of prion disease incidence? Chemosphere 2012; 87:661–7.

402   26.   Hemming-Schroeder E, Lo E, Salazar C, Puente S, Yan G. Landscape genetics: A toolbox for studying vector-borne diseases. Front Ecol Evol 2018; 6:1–11.

404   27.   Kelly AC, Mateus-Pinilla NE, Brown W, Ruiz MO, Douglas MR, Douglas ME, Shelton P, Beissel T, Novakofski J. Genetic assessment of environmental features that influence deer
406   dispersal: Implications for prion-infected populations. Popul Ecol 2014; 56:327–40.

        28.   Robinson SJ, Samuel MD, O'Rourke KI, Johnson CJ. The role of genetics in chronic
408   wasting disease of North American cervids. Prion 2012; 6:153–62.

        29.   Kelly AC, Mateus-Pinilla NE, Diffendorfer J, Jewell E, Ruiz MO, Killefer J, Shelton P,
410   Beissel T, Novakofski J. Prion sequence polymorphisms and chronic wasting disease resistance in Illinois white-tailed deer (*Odocoileus virginianus*). Prion 2008; 2:28–36.

412   30.   Brandt AL, Green ML, Ishida Y, Roca AL, Novakofski J, Mateus-Pinilla NE. Influence of the geographic distribution of prion protein gene sequence variation on patterns of chronic
414   wasting disease spread in white-tailed deer (*Odocoileus virginianus*). Prion 2018; 12:204–15.

416   31.   Brandt AL, Kelly AC, Green ML, Shelton P, Novakofski J, Mateus-Pinilla NE. Prion protein gene sequence and chronic wasting disease susceptibility in white-tailed deer
418   (*Odocoileus virginianus*). Prion 2015; 9:449–62.

        32.   Johnson C, Johnson J, Clayton M, McKenzie D, Aiken J. Prion protein gene heterogeneity

420   in free-ranging white-tailed deer within the chronic wasting disease affected region of Wisconsin. J Wildl Dis 2003; 39:576–81.

422   33. O'Rourke KI, Spraker TR, Hamburg LK, Besser TE, Brayton KA, Knowles DP. Polymorphisms in the prion precursor functional gene but not the pseudogene are
424   associated with susceptibility to chronic wasting disease in white-tailed deer. J Gen Virol 2004; 85:1339–46.

426   34. Johnson CJ, Herbst A, Duque-Velasquez C, Vanderloo JP, Bochsler P, Chappell R, McKenzie D. Prion protein polymorphisms affect chronic wasting disease progression.
428   PLoS One 2011; 6.

       35. Race B, Meade-White K, Miller MW, Fox KA, Chesebro B. *In vivo* comparison of
430   chronic wasting disease infectivity from deer with variation at prion protein residue 96. J Virol 2011; 85:9235–8.

432   36. Miller WL, Walter WD. Spatial heterogeneity of prion gene polymorphisms in an area recently infected by chronic wasting disease. Prion 2019; 13:65–76.

434   37. Avise JC. Phylogeography: Retrospect and prospect. J Biogeogr 2009; 36:3–15.

       38. Barton NH, Servedio MR. The interpretation of selection coefficients. Evolution 2015;
436   69:1101–12.

       39. Felsenstein J. Phylogenies and the comparative method. Am Nat 1985; 125:1–15.

438   40. Hoggart CJ, Parra EJ, Shriver MD, Bonilla C, Kittles RA, Clayton DG, McKeigue PM. Control of confounding of genetic associations in stratified populations. Am J Hum Genet
440   2003; 72:1492–504.

       41. Pagel M. Detecting correlated evolution on phylogenies: A general method for the
442   comparative analysis of discrete characters. Proc R Soc B Biol Sci 1994; 255:37–45.

       42. Carlson J, Kadie C, Mallal S, Heckerman D. Leveraging hierarchical population structure
444   in discrete association studies. PLoS One 2007; 2.

       43. Cook RL, Hart R V. Ages assigned known-age Texas white-tailed deer: Tooth wear
446   versus cementum analysis. In: Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies. 1979.

448   44. Mitchell CJ, Smith WP. Reliability of techniques for determining age in southern white-tailed deer. J Tennessee Adacemy Sci 1991; 66:117–20.

450   45. Gee KL, Holman JH, Causey MK, Rossi AN, Armstrong JB. Aging white-tailed deer by tooth replacement and wear: A critical evaluation of a time-honored technique. Wildl Soc
452   Bull 2002; 30:387–93.

       46. Seabury CM, Oldeschulte DL, Bhattarai EK, Legare D, Ferro PJ, Metz RP, Johnson CD,

454   Lockwood MA, Nichols TA. Accurate genomic predictions for chronic wasting disease in U.S. white-tailed deer. G3 2020; 10:1433–41.

456   47.   Hagenaars TJ, Melchior MB, Bossers A, Davidse A, Engel B, van Zijderveld FG. Scrapie prevalence in sheep of susceptible genotype is declining in a population subject to
458   breeding for resistance. BMC Vet Res 2010; 6:1–11.

        48.   Goldmann W. PrP genetics in ruminant transmissible spongiform encephalopathies. Vet
460   Res 2008; 39.

        49.   Herbst A, Velásquez CD, Triscott E, Aiken JM, McKenzie D. Chronic wasting disease
462   prion strain emergence and host range expansion. Emerg. Infect. Dis.2017; 23:1598–600.

        50.   Johnson C, Johnson J, Vanderloo JP, Keane D, Aiken JM, McKenzie D. Prion protein
464   polymorphisms in white-tailed deer influence susceptibility to chronic wasting disease. J Gen Virol 2006; 87:2109–14.

466   51.   Stein RA. Super-spreaders in infectious diseases. Int J Infect Dis 2011; 15:e510–3.

        52.   Pöschel T, Brilliantov N V., Frömmel C. Kinetics of prion growth. Biophys J 2003;
468   85:3460–74.

        53.   White LA, Forester JD, Craft ME. Disease outbreak thresholds emerge from interactions
470   between movement behavior, landscape structure, and epidemiology. Proc Natl Acad Sci U S A 2018; 115:7374–9.

54.   Blanchong JA, Robinson SJ, Samuel MD, Foster JT. Application of genetics and genomics to wildlife epidemiology. J Wildl Manage 2016; 80:593–608.

55.   Severinghaus CW. Tooth development and wear as criteria of age in white-tailed deer. J Wildl Manage 1949; 13:195–216.

56.   Stephens M, Smith NJ, Donnelly P. A new statistical method for haplotype reconstruction from population data. Am J Hum Genet 2001; 68:978–89.

57.   Leigh JW, David B, Shinichi N. popart: full□feature software for haplotype network construction. Methods Ecol Evol 2015; 6:1110–6.

58.   Mathieson I, McVean G. Estimating selection coefficients in spatially structured populations from time series data of allele frequencies. Genetics 2013; 193:973–84.

59.   Choi Y, Sims GE, Murphy S, Miller JR, Chan AP. Predicting the Functional Effect of Amino Acid Substitutions and Indels. PLoS One 2012; 7.

60.   Sim NL, Kumar P, Hu J, Henikoff S, Schneider G, Ng PC. SIFT web server: Predicting effects of amino acid substitutions on proteins. Nucleic Acids Res 2012; 40:452–7.

61.   Capriotti E, Fariselli P, Casadio R. I-Mutant2.0: Predicting stability changes upon

mutation from the protein sequence or structure. Nucleic Acids Res 2005; 33:306–10.

Table 1: *PRNP* haplotypes tabulated from 1,433 white-tailed deer tissues collected from 75 counties in Arkansas (2016-2019). Haplotypes (Hap) are identified by letter (A-V) following Brandt et al. (2015, 2018). Haplotypes denoted as AR1-4 are previously unreported. Mutations that differ from haplotype A are shaded, with green indicating synonymous substitutions (no amino acid change) and blue as non-synonymous (amino acid changed in protein; NSS). Also listed are amino acid position and nucleotide site (Brandt et al. 2015, 2018).

Hap	NSS	Amino Acid Position													
		20	51	81	95	**96**	99	108	**122**	124	126	146	166	185	**225**
		Nucleotide Site													
		60	153	243	285	**286**	299	324	**367**	372	378	438	499*	555	**676**
A		C	C	T	A	G	G	A	G	G	G	C	A	C	C
B		C	C	T	A	G	G	A	G	G	G	C	A	T	C
C	96S	C	C	T	A	A	G	A	G	G	G	C	A	T	C
D		C	T	T	A	G	G	A	G	G	G	C	A	C	C
E		C	C	T	A	G	G	A	G	G	G	T	A	C	C
G		T	C	T	A	G	G	A	G	G	G	C	A	C	C
I	96S	C	C	A	A	A	G	A	G	G	G	C	A	T	C
J		C	C	T	A	G	G	G	G	G	G	C	A	C	C
K	225K	T	C	T	A	G	G	A	G	G	G	C	A	C	A
L	122T	C	C	T	A	G	G	A	A	G	G	C	A	C	C
O		T	T	T	A	G	G	A	G	G	G	C	A	C	C
P	96S	C	C	T	A	A	G	A	G	G	G	C	A	C	C
R		C	T	T	A	G	G	A	G	G	G	C	A	T	C
T		C	T	T	A	G	G	A	G	G	A	C	A	C	C
V	96S	C	T	T	A	A	G	A	G	G	G	C	A	C	C
AR1		C	C	T	A	G	G	G	G	G	A	C	A	C	C
AR2	96S	C	C	T	A	A	G	A	G	G	G	C	C	T	C
AR3	96S	T	C	T	A	A	G	A	G	G	G	C	A	T	C
AR4		C	C	T	A	G	G	A	G	G	G	T	A	T	C

Table 2: *PRNP* haplotype frequencies and odds-ratios, as associated with CWD status of white-tailed deer in Arkansas, 2016-2019. Haplotypes were derived from unphased sequences representing 720 nucleotides of the *PRNP* gene. Haplotypes (Hap) are identified by letter (A-V) following Brandt et al. (2015, 2018). Haplotypes denoted AR1-4 are previously unreported. Listed are total numbers (N) and relative frequency (%) and associated values for deer that were CWD-negative (-), CWD-positive (+) or untested (?). Odds Ratio (OR) reflects relative representation of a haplotype in CWD-positive deer; OR>1 indicated over-representation, OR<1 under-representation. SE= standard error, CI= 95% confidence interval, Z= OR Z-score and *p*= OR *p*-value. Values in bold are significant.

Hap	Counts				Frequency (%)				Odds Ratio				
	N	*N*(-)	*N*(+)	*N*(?)	%	%(-)	%(+)	%(?)	OR	SE	CI	Z	*p*
A	435	367	34	34	15.18	15.17	13.71	17.17	0.89	0.19	[0.61-1.30]	-0.61	0.54
B	657	536	90	31	22.92	22.15	36.29	15.66	**2.00**	**0.14**	**[1.52-2.64]**	**4.93**	**0.00**
C	426	376	13	37	14.86	15.54	5.24	18.69	**0.30**	**0.29**	**[0.17-0.53]**	**-4.14**	**0.00**
D	657	547	63	47	22.92	22.60	25.40	23.74	1.17	0.15	[0.86-1.58]	1.00	0.32
E	187	160	16	11	6.52	6.61	6.45	5.56	0.97	0.27	[0.57-1.66]	-0.10	0.92
G	309	266	24	19	10.78	10.99	9.68	9.60	0.87	0.22	[0.56-1.35]	-0.63	0.53
I	42	39	1	2	1.47	1.61	0.40	1.01	0.25	1.01	[0.03-1.81]	-1.38	0.17
J	65	59	3	3	2.27	2.44	1.21	1.52	0.49	0.60	[0.15-1.57]	-1.20	0.23
K	10	9	1	0	0.35	0.37	0.40	0.00	1.08	1.06	[0.14-8.60]	0.08	0.94
L	2	2	0	0	0.07	0.08	0.00	0.00	-	-	-	-	-
O	3	2	1	0	0.10	0.08	0.40	0.00	4.89	1.23	[0.44-54.2]	1.29	0.20
P	10	8	0	2	0.35	0.33	0.00	1.01	-	-	-	-	-
R	2	2	0	0	0.07	0.08	0.00	0.00	-	-	-	-	-
T	41	30	1	10	1.43	1.24	0.40	5.05	0.32	1.02	[0.04-2.38]	-1.11	0.27
V	1	1	0	0	0.03	0.04	0.00	0.00	-	-	-	-	-
AR1	1	0	0	1	0.03	0.00	0.00	0.51	-	-	-	-	-
AR2	15	14	0	1	0.52	0.58	0.00	0.51	-	-	-	-	-
AR3	2	2	0	0	0.07	0.08	0.00	0.00	-	-	-	-	-
AR4	1	0	1	0	0.03	0.00	0.40	0.00	-	-	-	-	-

520   **Figure 1:** Sampling locations for white-tailed deer tissues from Arkansas evaluated in this study.

The red shaded area indicates the 16 counties included in the 2019 Chronic Wasting Disease

522   Management Zone (CWDMZ), with a yellow boundary surrounding a focal area encompassing

Newton County. Black dots represent collection localities for each individual tissue. Note that

524   the boundaries of the CWDMZ have since expanded.

526   **Figure 2:** Haplotype network showing relationship of prion gene variants (*PRNP*) detected

across 1,433 white-tailed deer collected from 75 counties in Arkansas (2016-2019). Data are

528   based on sequence analysis of 720 nucleotides. Circles represent 20 haplotypes (=alleles) with

size reflecting frequency of occurrence in entire data set (Table 3), and tick marks representing

530   number of mutations (=nucleotide substitutions) distinguishing one from another (Table 2).

Color codes reflect relative frequency among CWD-positive (red) and CWD-negative/

532   undetected (green). Letters correspond to haplotype names (per Brandt et al. 2015), with

haplotypes unique to Arkansas indicated with numbers (AR#). Haplotypes sharing the 96S

534   mutation are indicated with (*).

536   **Figure 3:** Frequency distribution of 2,866 *PRNP* haplotypes detected in white-tailed deer

collected in Arkansas, 2016-2019. Haplotypes were determined by phasing individual genotypes

538   derived from sequencing 1,433 deer across 720 bp of the *PRNP* gene. Letters (A through V) refer

to haplotypes identified in Brandt et al. (2015), whereas numbers (1-4) are haplotypes unique to

540   Arkansas, and thus previously unreported. Frequencies are plotted for all 1,433 samples

(=statewide) and a subset of 314 samples from Newton County (N=628 chromosomes). Color

542   codes reflect frequency among CWD-positive (CWD+) and CWD-negative (CWD-) samples;

unknown indicates samples that were not tested for CWD.

Figure 4: Topographies that represent interpolated haplotype frequencies for the *PRNP* gene in

546   Arkansas. Frequency is depicted by color, with blue reflecting low occurrence (0-5%) whereas

red indicating 46-50+% of haplotypes were of this type.

Figure 5: Relative frequency and odds ratio for two haplotypes of the prion gene *PRNP*

550   haplotypes detected in white-tailed deer age cohorts (<1 year to 5+ years) sampled in Arkansas

from 2016-2019. Prion gene variant Haplotype C (top panel) has been associated with reduced

552   susceptibility to CWD, whereas Haplotype B (lower panel) has been associated with higher

susceptibility (Brandt et al. 2018). Data are based on phased haplotypes derived from 720

554   nucleotides of the *PRNP* gene sequenced across 1,433 deer.

556   **Figure 6:** Results of a conceptual model depicting prion conversion rates for PrP^C variants 96G

and 96S. The diagram demonstrates prion conversion of 96G and 96S (reduced susceptibility)

558   prion gene (=*PRNP*) variants. Theoretical prion protein (=*PrP*) mass thresholds were imposed:

Ta = time of symptomatic expression; and Tb = theoretical time after which individuals remain

560   asymptomatic. Individual-individual transmission is possible through direct contact (saliva) or

shedding (e.g. feces). The time interval for *asymptomatic spread* (e.g., prion shedding from a

562   subclinical individual) extends prior to *PrP* conversion surpassing threshold Tb, but not

extending beyond Ta.

Supplemental Figures and Tables

566 **Supplemental Table 1**: Samples per county and frequency of 20 *PRNP* haplotypes detected in 1,433 white-tailed deer collected in 75 counties in Arkansas from 2016-2019. Phased haplotypes
568 were derived from sequence analysis of 720 nucleotides of the *PRNP* gene. Letters indicate haplotypes previously detected in other states, whereas numbers (1-4) indicate haplotypes unique
570 to Arkansas. Variable sites of haplotypes are listed in Table 2. Samples that tested positive for CWD (+) are listed separately for those counties where CWD was detected.

County	N	Haplotype																		
		A	B	C	D	E	G	I	J	K	L	O	P	R	T	V	1	2	3	4
Arkansas	11	4	5	5	4	-	3	-	-	-	-	-	-	-	1	-	-	-	-	-
Ashley	1	1	-	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-
Baxter	16	3	7	3	8	4	6	-	1	-	-	-	-	-	-	-	-	-	-	-
Benton	40	25	19	3	16	5	5	1	4	-	-	-	1	-	1	-	-	-	-	-
Boone	33	6	24	8	7	5	8	-	7	-	1	-	-	-	-	-	-	-	-	-
Boone(+)	2	-	-	-	3	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Bradley	2	1	-	2	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-
Carroll	33	18	13	4	11	4	7	-	6	-	-	-	-	-	3	-	-	-	-	-
Carroll(+)	16	5	12	2	5	3	5	-	-	-	-	-	-	-	-	-	-	-	-	-
Calhoun	14	4	9	1	6	3	2	-	1	-	-	-	1	-	1	-	-	-	-	-
Cleburne	8	2	2	2	4	2	4	-	-	-	-	-	-	-	-	-	-	-	-	-
Chicot	4	3	-	1	2	-	1	1	-	-	-	-	-	-	-	-	-	-	-	-
Clark	13	2	5	6	6	1	1	3	1	-	-	-	1	-	-	-	-	-	-	-
Conway	10	1	4	1	6	3	5	-	-	-	-	-	-	-	-	-	-	-	-	-
Columbia	4	1	1	-	4	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-
Craighead	12	5	1	5	7	1	3	-	1	-	-	-	-	-	1	-	-	-	-	-
Cross	9	4	4	3	5	-	1	1	-	-	-	-	-	-	-	-	-	-	-	-
Crittenden	8	5	1	5	4	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Crawford	9	1	3	5	6	-	1	-	-	2	-	-	-	-	-	-	-	-	-	-
Clay	14	7	4	7	3	1	2	1	-	-	-	-	1	-	2	-	-	-	-	-
Dallas	9	6	3	1	3	3	1	1	-	-	-	-	-	-	-	-	-	-	-	-
Drew	14	4	2	16	3	1	2	-	-	-	-	-	-	-	-	-	-	-	-	-
Desha	12	8	4	5	4	-	1	-	-	-	-	-	-	-	1	-	1	-	-	-
Faulkner	14	5	1	2	11	4	1	-	1	-	-	-	1	2	-	-	-	-	-	-
Franklin	33	6	23	29	1	-	5	-	-	-	-	-	-	-	-	-	-	2	-	-
Fulton	21	3	8	4	12	4	5	-	1	-	-	-	-	-	5	-	-	-	-	-
Garland	16	6	5	6	5	4	1	3	-	-	-	-	-	-	-	-	-	2	-	-
Greene	12	1	8	7	-	-	2	2	1	-	-	-	-	-	3	-	-	-	-	-
Grant	10	4	3	3	7	-	2	-	-	-	-	-	-	-	1	-	-	-	-	-
Hampstead	10	6	3	-	5	2	1	3	-	-	-	-	-	-	-	-	-	-	-	-
Howard	13	-	3	6	11	2	2	1	-	-	-	-	-	-	1	-	-	-	-	-
Hot Springs	5	2	2	2	1	1	2	-	-	-	-	-	-	-	-	-	-	-	-	-
Independence	10	-	3	5	4	1	4	-	-	-	-	-	-	-	1	-	-	2	-	-
Izard	9	-	2	-	7	6	2	-	-	-	-	-	-	-	-	-	-	1	-	-
Jackson	2	-	-	3	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Jefferson	13	6	7	2	5	1	4	-	1	-	-	-	-	-	-	-	-	-	-	-

County																				
Johnson	**45**	10	33	19	20	1	4	1	-	-	-	-	1	-	-	-	-	1	-	-
Lafayette	**15**	5	3	7	5	1	4	1	-	-	-	-	-	-	3	-	-	1	-	-
Lincoln	**9**	3	3	5	4	-	1	-	2	-	-	-	-	-	-	-	-	-	-	-
Lee	**8**	4	6	3	1	-	1	-	-	1	-	-	-	-	-	-	-	-	-	-
Lonoke	**13**	6	4	5	7	3	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Logan	**41**	12	15	3	26	12	13	-	1	-	-	-	-	-	-	-	-	-	-	-
Little River	**10**	1	4	5	8	1	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Lawrence	**16**	1	3	16	2	4	3	-	-	-	-	-	-	-	3	-	-	-	-	-
Madison	**36**	9	18	24	6	3	6	1	2	-	-	-	-	-	2	-	-	1	-	-
Madison(+)	*4*	1	1	1	-	2	-	1	2	-	-	-	-	-	-	-	-	-	-	-
Marion	**40**	10	12	11	12	10	7	-		1	-	-	2	-	1	-	-	-	-	-
Miller	**8**	3	5	2	1	1	2	-	-	-	-	-	-	-	2	-	-	-	-	-
Monroe	**9**	4	1	5	3	4	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Montgomery	**2**	1	1	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi	**2**	2	-	1	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-
Nevada	**14**	7	-	6	10	2	3	-	-	-	-	-	-	-	-	-	-	-	-	-
Newton	**216**	68	124	49	105	23	48	-	5	3	-	2	-	-	2	-	-	2	1	-
Newton(+)	*100*	28	75	10	54	11	17	-	1	1	-	1	-	-	1	-	-	-	-	1
Ouachita	**14**	4	3	3	13	3	1	1	-	-	-	-	-	-	-	-	-	-	-	-
Perry	**8**	4	3	2	2	-	2	2	1	-	-	-	-	-	-	-	-	-	-	-
Phillips	**9**	8	1	1	1	-	5	-	1	-	-	-	-	-	-	-	-	-	1	-
Pike	**8**	2	3	2	5	-	2	2	-	-	-	-	-	-	-	-	-	-	-	-
Polk	**3**	2	-	-	-	2	1	1	-	-	-	-	-	-	-	-	-	-	-	-
Poinsett	**13**	7	1	5	6	2	3	-	1	-	-	-	-	-	-	-	-	1	-	-
Pope	**61**	3	48	9	41	3	15	3	-	-	-	-	-	-	-	-	-	-	-	-
Pope(+)	*1*	-	1	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Prairie	**11**	3	5	7	3	-	1	1	-	2	-	-	-	-	-	-	-	-	-	-
Pulaski	**11**	3	3	4	5	3	3	1	-	-	-	-	-	-	-	-	-	-	-	-
Randolph	**8**	3	3	5	3	-	-	-	1	-	-	-	-	-	1	-	-	-	-	-
Saline	**12**	4	4	4	1	3	3	1	2	-	-	-	-	-	2	-	-	-	-	-
Sebastian	**16**	6	3	2	12	3	5	1	-	-	-	-	-	-	-	-	-	-	-	-
Sebastian(+)	*1*	-	1	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Scott	**2**	1	2	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Searcy	**42**	8	15	7	31	5	16	-	1	-	-	-	1	-	-	-	-	-	-	-
St Francis	**8**	4	4	3	5	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Sharp	**15**	3	6	7	8	1	1	-	1	-	-	-	-	-	3	-	-	-	-	-
Stone	**15**	6	5	3	6	1	9	-	-	-	-	-	-	-	-	-	-	-	-	-
Sevier	**10**	5	1	2	5	1	4	2	-	-	-	-	-	-	-	-	-	-	-	-
Union	**7**	3	3	1	4	1	1	-	-	-	-	-	-	-	-	-	-	1	-	-
VanBuren	**33**	7	10	2	21	8	14	2	-	-	1	-	-	-	-	1	-	-	-	-
Washington	**18**	7	5	11	9	-	2	1	1	-	-	-	-	-	-	-	-	-	-	-
Woodruff	**9**	1	2	7	2	1	4	-	-	-	-	-	-	-	-	-	-	-	1	-
White	**8**	2	4	6	-	1	2	-	1	-	-	-	-	-	-	-	-	-	-	-
Yell	**40**	9	22	5	21	10	9	3	1	-	-	-	-	-	-	-	-	-	-	-
TOTAL	**1433**	**435**	**657**	**426**	**657**	**187**	**309**	**42**	**65**	**10**	**2**	**3**	**10**	**2**	**41**	**1**	**1**	**15**	**2**	**1**

574 **Supplemental Table 2:** Association of *PRNP* haplotype frequencies and odds ratio with CWD status for Newton County, Arkansas. Haplotypes were derived from unphased sequences of 720

576 nucleotides of the *PRNP* gene. Haplotype as indicated by letters were also reported by Brandt et al. (2015, 2018), whereas AR_# indicates a haplotype unique to Arkansas. Listed are total

578 numbers (N), relative frequency f(%), and values for deer that tested either CWD-negative (-), CWD-positive (+) or were untested (?). Odds Ratio (OR) reflects relative representation of a

580 haplotype in CWD+ deer, with OR>1 = over-representation and OR<1 = under-representation; SE= standard error, CI= 95% confidence interval, Z(OR)= Z-score and p(OR)= probability.

582 **Values in bold are significant**.

Hap	Counts			Frequency (%)			Odds Ratio				
	N	N(-)	N(+)	f%	f%(-)	f%(+)	OR	SE	CI	Z(OR)	p(OR)
A	95	68	27	15.1	15.7	13.8	0.86	0.25	[053-1.38]	-0.64	0.52
B	197	124	73	31.4	28.7	37.2	**1.47**	**0.18**	**[1.03-2.11]**	**2.13**	**0.03**
C	59	49	10	9.39	11.3	5.1	**0.42**	**0.36**	**[0.21-0.85]**	**-2.42**	**0.02**
D	158	105	53	25.2	24.3	27	1.15	0.20	[0.79-1.70]	0.73	0.46
E	34	23	11	5.41	5.32	5.61	1.06	0.38	[0.50-2.21]	0.15	0.88
G	65	48	17	10.4	11.1	8.67	0.76	0.30	[0.43-1.35]	-0.93	0.35
I	-	-	-	-	-	-	-	-	-	-	-
J	6	5	1	0.96	1.16	0.51	0.44	1.10	[0.05-3.77]	-0.75	0.45
K	4	3	1	0.64	0.69	0.51	0.73	1.16	[0.08-7.09]	-0.27	0.79
L	-	-	-	-	-	-	-	-	-	-	-
O	3	2	1	0.48	0.46	0.51	1.10	1.23	[0.10-12.23]	0.08	0.94
P	-	-	-	-	-	-	-	-	-	-	-
R	-	-	-	-	-	-	-	-	-	-	-
T	3	2	1	0.48	0.46	0.51	1.10	1.23	[0.10-12.23]	0.08	0.94
V	-	-	-	-	-	-	-	-	-	-	-
AR_1	-	-	-	-	-	-	-	-	-	-	-
AR_2	2	2	-	0.32	0.46	-	-	-	-	-	-
AR_3	1	1	-	0.16	0.23	-	-	-	-	-	-
AR_4	1	-	1	0.16	-	0.51	-	-	-	-	-
Total	628	432	196								

586 **Supplemental Table 3:** *PRNP* haplotype frequencies and odds ratio for six age classes of white-tailed deer collected in Newton County, Arkansas from 2016-2019. Relative frequencies and
588 odds ratio are shown for (A) Haplotype C (associated with reduced CWD susceptibility), and (B) Haplotype B (associated with increased CWD susceptibility). Relative frequency of CWD-
590 negative [f(-)] and CWD-positive [f(+)] samples were calculated by dividing numbers CWD(+/-) with C or B, respectively, then by negatives w/ other haplotypes.
592

(A) Haplotype C

Age	N	N(-)	N(+)	f(-)	f(+)	OR
FAWN	9	8	1	0.13	0.04	0.34
Y1	5	5	0	0.08	0.00	0.40
Y2	22	18	4	0.15	0.08	0.53
Y3	16	13	3	0.21	0.09	0.40
Y4	4	3	1	0.07	0.08	1.05
Y5	8	6	2	0.19	0.17	0.89
Total	64	53	11			

596 **(B) Haplotype B**

Age	N	N(-)	N(+)	f(-)	f(+)	OR
FAWN	21	14	7	0.25	0.41	1.65
Y1	26	20	6	0.45	0.75	1.65
Y2	73	52	21	0.58	0.60	1.04
Y3	28	15	13	0.25	0.52	2.05
Y4	23	17	6	0.63	0.75	1.19
Y5	11	6	5	0.19	0.56	2.96
Total:	182	124	58			

600   **Supplemental Table 4**: Modeling results for amino acid variant effect versus three non-synonymous *PRNP* gene polymorphisms. Results represent: I-MUTANT stability index (where
602   negative indices destabilize); PROVEAN conservation score (heightened negative value = more likely to be deleterious, with -2.5 as a putative neutral cutoff); MutPred2 probability of
604   pathogenicity (with values >0.8 interpreted as pathogenic); and SIFT conservation score (where 4.32=completely conserved, and 0.0=completely polymorphic).

Mutant	Method			
	I-MUTANT	**PROVEAN**	**MutPred2**	**SIFT**
96S	-0.37	-1.250	0.366	3.88
112T	-0.27	-0.937	0.446	3.88
225K	0.13	-0.743	0.425	4.32

608 **Supplemental Figure 1:** Spatial distribution of 211 non-overlapping polygons in the state of Arkansas (U.S.A.). Each included 5-10 sampling locations. Closed red and green circles
610 represent 1,433 white-tailed deer samples employed to compute prion gene variant (*PRNP* haplotype) frequencies for interpolation. Red circles = CWD+; green circles = CWD-
612

616 **Supplemental Figure 2:** Age and sex distribution for *N*=1,433 white-tailed deer tissue samples
for which *PRNP* haplotype data was generated.

620