Mechanical overstimulation causes acute injury and synapse loss followed by fast

recovery in lateral-line neuromasts of larval zebrafish

Melanie Holmgren[1], Michael E. Ravicz[4,5], Kenneth E. Hancock[4,5], Olga Strelkova[4,5], Dorina

Kallogjeri[1], Artur A. Indzhykulian[4,5], Mark E. Warchol[1,3], Lavinia Sheets[1,2]*

1. Department of Otolaryngology, Washington University School of Medicine, St. Louis, MO, USA

2. Department of Developmental Biology, Washington University School of Medicine, St. Louis, MO, USA

3. Department of Neuroscience, Washington University School of Medicine, St. Louis, MO, USA

4. Eaton-Peabody Laboratory, Massachusetts Eye and Ear, Boston, Massachusetts, USA

5. Department of Otolaryngology–Head and Neck Surgery, Harvard Medical School, Boston,

Massachusetts, USA

* Corresponding author

Lavinia Sheets: sheetsl@wustl.edu

Abstract

Excess noise damages sensory hair cells, resulting in loss of synaptic connections with

auditory nerves and, in some cases, hair-cell death. The cellular mechanisms underlying

mechanically induced hair-cell damage and subsequent repair are not completely understood.

Hair cells in neuromasts of larval zebrafish are structurally and functionally comparable to

mammalian hair cells but undergo robust regeneration following ototoxic damage. We therefore

developed a model for mechanically induced hair-cell damage in this highly tractable system.

Free swimming larvae exposed to strong water wave stimulus for 2 hours displayed mechanical

injury to neuromasts, including afferent neurite retraction, damaged hair bundles, and reduced

mechanotransduction. Synapse loss was observed in apparently intact exposed neuromasts,

and this loss was exacerbated by inhibiting glutamate uptake. Mechanical damage also elicited

an inflammatory response and macrophage recruitment. Remarkably, neuromast hair-cell

morphology and mechanotransduction recovered within hours following exposure, suggesting

severely damaged neuromasts undergo repair. Our results indicate functional changes and

synapse loss in mechanically damaged lateral-line neuromasts that share key features of

damage observed in noise-exposed mammalian ear. Yet, unlike the mammalian ear,

mechanical damage to neuromasts is rapidly reversible.

Keywords: Hair cell, mechanical damage, ribbon synapse, inflammation, regeneration

Introduction

Hair cells are the sensory receptors of the inner ear and lateral-line organs that detect

sound, orientation, and motion. They transduce these stimuli through deflection of stereocilia,

which opens mechanically-gated cation channels (LeMasurier & Gillespie, 2005; Qiu & Muller,

2018) and drives subsequent transmission of sensory information via excitatory glutamatergic

synapses (Glowatzki & Fuchs, 2002). Excessive mechanical stimulation, such as loud noise,

can damage hair cells and their synaptic connections to afferent nerves. The degree of damage

depends on the intensity and duration of the stimulus, with higher levels of traumatic noise

producing structural damage to sensory epithelia, including hair cells (Cho et al., 2013;

Nordmann, Bohne, & Harding, 2000; Slepecky, 1986) and lower levels leading to various

pathologies, including damage to the hair-cell mechanotransduction complex/machinery and

stereocilia (Gao, Ding, Zheng, Ruan, & Liu, 1992; Husbands, Steinberg, Kurian, & Saunders,

1999), misshapen hair cells (Bullen, Anderson, Bakay, & Forge, 2019), synaptic terminal

damage, neurite retraction, and hair-cell synapse loss (Fernandez et al., 2020; Henry & Mulroy,

1995; Kujawa & Liberman, 2009; Puel, Ruel, Gervais d'Aldin, & Pujol, 1998). In addition to

directly damaging hair cells, excess noise also initiates an inflammatory response (Hirose,

Discolo, Keasler, & Ransohoff, 2005; Kaur et al., 2019). Such inflammation is mediated by

macrophages, a class of leukocyte that responds to injury by clearing cellular debris and

promoting tissue repair (Wynn & Vannella, 2016).

The variety of injury and range of severity suggests multiple mechanisms are involved in

hair-cell organ damage associated with exposure to strong stimuli. Also unknown are the

cellular processes mediating repair following such trauma. While hair cells show a partial

capacity for repair of stereocilia and synaptic connections, some sub-lethal damage to hair cells

is permanent. Numerous studies of the mammalian cochlea suggest that a subgroup of inner

hair-cell synapses are permanently lost following noise exposure (Cho et al., 2013; Hickman,

Smalt, Bobrow, Quatieri, & Liberman, 2018; Kujawa & Liberman, 2009; Shi et al., 2013).

Glutamate excitotoxicity is likely the pathological event that initiates noise-induced hair-cell

synapse loss (Hu, Rutherford, & Green, 2020; Kim et al., 2019; Puel et al., 1998), but the

downstream cellular mechanisms are still undefined. Further, the cellular mechanisms that

promote hair-cell synapse recovery following damage-induced loss are also not understood.

Zebrafish have proven to be a valuable model system for studying the molecular basis of

hair-cell injury and repair. Zebrafish sensory hair cells are structurally and functionally

homologous to mammalian hair cells (Coffin, Kelley, Manley, & Popper, 2004; Kindt & Sheets,

2018; Sebe et al., 2017). In contrast to other vertebrate model organisms, zebrafish hair cells

are optically accessible in whole larvae within the lateral-line organs. These sensory organs,

called neuromasts, contain clusters of ~ 14 hair cells each and are distributed throughout the

external surface of the fish to detect local water movements. Zebrafish can repair and

regenerate damaged tissues, including lateral-line organs (Kniss, Jiang, & Piotrowski, 2016;

Xiao et al., 2015). This capacity for organ repair combined with optical accessibility allows us to

study cellular and synaptic damage and repair *in vivo* following mechanical trauma.

In order to model physical damage to hair cell organs in zebrafish lateral line, we

developed a protocol to mechanically stimulate the lateral line of free-swimming 7-day-old

larvae. Using this protocol, we were able to induce mechanical injury to lateral-line organs that

resembled the trauma observed in the mammalian cochlea following acoustic overstimulation.

We observed synapse loss in a subset of neuromasts that appeared morphologically intact, as

well as hair-cell loss and afferent neurite retraction in a subset of neuromasts that appeared

morphologically disrupted. Hair cell mechanotransduction, as measured by uptake of the

cationic dye FM1-43, was significantly reduced after mechanical injury. We also observed an

inflammatory response similar to that observed in the mammalian cochlea after noise trauma.

Remarkably, mechanically induced lateral-line damage appeared to rapidly recover; hair-cell

number and morphology returned to normal within 4 hours following exposure, concurrent with

clearance of cellular debris by macrophages. Additionally, neuromasts showed partial recovery

of afferent innervation within 2 hours following exposure and completely recovered hair-cell

morphology and FM1-43 uptake within 4 hours. Cumulatively, these results support that

mechanically injured neuromasts show similar features of damage observed in noise exposed

ears, yet rapidly repair.

<u>Results</u>

Mechanical overstimulation of zebrafish lateral-line hair cells

To mechanically damage hair cells of lateral-line organs in free-swimming 7-day-old

zebrafish, we developed a stimulation protocol using an electrodynamic shaker to create a

strong water wave stimulus (Fig.1 A). The frequency used for mechanical stimulation was

selected and further verified (see Method Details) based on previous studies showing 60 Hz to

be within the optimal upper frequency range of mechanical sensitivity of superficial posterior

lateral-line neuromasts, which respond maximally between 10-60 Hz, but a suboptimal

frequency for hair cells of the anterior macula of the inner ear (Levi, Akanyeti, Ballo, & Liao,

2015; Trapani & Nicolson, 2010; Weeg, Fay, & Bass, 2002). Dorsal-ventral displacement of a 6-

well dish at 60 Hz and acceleration of 40.3 m/s^2 (+/-0.5m/s^2) created water flow and disturbance

of the water surface that was strong enough to trigger 'fast start' escape responses—a behavior

Figure 1: Intense water wave produced by shaker apparatus stimulates lateral-line hair cells and evokes a relevant behavior response. **(A)** The apparatus: a magnesium head expander holding a 6-well dish mounted on a vertically oriented electrodynamic shaker housed inside a sound-attenuation chamber. The stimulus consisted of a 60 Hz vertical displacement of the plate (hatched arrows) driven by an amplifier and controlled by a modified version of the Eaton-Peabody Laboratory Cochlear Function Test Suite. **(B,D)** Swimming behavior of 7-day-old larvae during exposure to the wave stimulus. Traces in (D) represent tracking of corresponding circled fish over 500 ms (1000 fps/ 500 frames). Asterisks indicate a "fast escape" response (B; inset). **(C,E)** Swimming behavior of larvae whose lateral-line neuromasts were ablated with low-dose CuSO4. Arrows in (E) indicate where a larva was swept into the waves and could no longer be tracked.

mediated in part by zebrafish lateral-line organs to escape predation (Fig. 1 B inset) (McHenry,

Feitl, Strother, & Van Trump, 2009; Nair, Azatian, & McHenry, 2015).

'Fast start' escape responses can also be activated by stimulating hair cells of the

lateral line and/or the posterior macula in the ear (Bhandiwad, Zeddies, Raible, Rubel, &

Sisneros, 2013). To verify that the observed escape responses were mediated predominantly by

flow sensed by lateral-line hair cells rather than hair cells of the macula, we exposed a group of

larvae to low dose (3µM) copper sulfate ($CuSO_4$) for 1 hour to specifically ablate lateral-line hair

cells, but leave hair cells of the ear intact (Olivari, Hernandez, & Allende, 2008). Following a 2

hour recovery after $CuSO_4$ treatment, we recorded fish behavior with a high-speed camera

during the stimulus (1000 fps for 10 s) and compared the responses of fish with lesioned lateral-

line organs with those of untreated control siblings (Fig. 1 B,C; Movie S1,2). When subjected to

intense water flow, we found that 'fast start' responses—defined as a c-bend of the body

occurring within 15 ms followed by a counter-bend (Burgess & Granato, 2007; McHenry et al.,

2009)—occurred significantly less frequently in larvae with ablated lateral line organs than in

siblings with intact lateral line (Fig. 1 C,E; avg. 'fast start' responses: : 0.4 (± 0.1)/s in lateral-line

ablated vs 1.5 (± 0.1)/s in control; 3 trials,10 s per trial; **P=0.0043). Accordingly, some $CuSO_4$-

treated fish were unable to escape the waves and were swept out of view (Fig. 1 E; arrowheads;

Movie S2). These observations indicate that the strong water wave generated by our device is

stimulating lateral-line hair cells and evoking a behaviorally relevant response.

A subset of mechanically injured neuromasts undergo physical displacement that is

position dependent but does not require mechanotransduction

To induce mechanical damage to lateral line organs, larvae (10-15 per well) were

exposed to an initial 20 minutes of strong water wave stimulus followed by 10 minutes of rest,

then 2 hours of continuous stimulus (Supplemental Fig. 1 A). The ten-minute break in exposure

was introduced early on when establishing the stimulus duration because it appeared to

enhance larval survival; it was therefore maintained throughout the study for consistency. Fish

were euthanized and fixed immediately after exposure, then processed for

immunohistofluorescent labeling of hair cells and neurons. Unexposed sibling fish served as

controls. As posterior lateral-line (pLL) neuromasts have been shown to specifically initiate

escape behavior in response to strong stimuli (Liao & Haehnel, 2012), analysis of the

morphology of pLL neuromasts L3, L4, and L5 was conducted for exposed and control larvae

(Fig. 2 A). Initially, we divided the observed neuromast morphology into two categories:

"normal", in which hair cells were radially organized with a relatively uniform shape and size,

and "disrupted", in which the hair cells were misshapen and displaced to one side, with the

apical ends of the hair cells localized anteriorly (Fig. 2 B, C; see Methods Details for measurable

criteria). Position of the neuromast along the tail was also associated with vulnerability to

disruption; we observed a gradient of damage in the pLL from rostral to caudal i.e. L5 was more

Figure 2: Morphological changes in pLL neuromast hair cells exposed to strong water wave stimulus. (A) Schematic of a larval zebrafish. Blue dots indicate neuromasts of the lateral-line organs; green lines indicate innervating afferent lateral-line nerves. pLL neuromasts L3, L4, and L5 were analyzed (dashed circles). **(B-C)** Maximum intensity dorsal top-down 2D projections of confocal images of control or stimulus-exposed neuromast hair cells (blue (B) or orange (C); Parvalbumin immunolabel). Exposed neuromast hair-cell morphology was categorized as "normal" i.e. radial hair-cell organization indistinguishable from control or "disrupted" i.e. asymmetric organization with the hair-cell apical ends oriented posteriorly. **(D)** Maximum intensity projections of supporting cells (SCs) expressing GFP (green), immunolabeled synaptic ribbons (magenta; Ribeye b) and all cell nuclei (blue; DAPI). Note that SCs underlying displaced hair cells also appear physically disrupted (indicated by white arrows). Scale bars: 5 μm **(F)** Average percentage of neuromasts with "disrupted" morphology following mechanical stimulation. Each dot represents the percentage of disrupted neuromasts (NM) in a single experimental trial. Disrupted hair-cell morphology was place dependent, with neuromasts more frequently disrupted following sustained stimulus and when localized toward the posterior end of the tail (*P=0.0386, **P=0.0049, ***P=0.0004) **(G)** Average percentage of exposed neuromasts (NM) with "disrupted" morphology in *lhfpl5b* mutants, which lack mechanotransduction specifically in lateral-line hair cells, vs. heterozygous WT. *lhfpl5b* mutants show a similar gradient of neuromast disruption following mechanical injury as WT siblings. Error Bars = SEM

susceptible to disruption than L4, which was more susceptible to disruption than L3 (Fig. 2 F;

Repeated measure One-way ANOVA *P=0.0386, **P=0.0049, ***= 0.0004).

Additionally, we compared mechanical stress from our sustained exposure to an

exposure protocol that delivered intermittent pulses of stimulus ("periodic exposure";

Supplemental Fig.1 A). We observed neuromast disruption less frequently with periodic

exposures vs. sustained exposure of the same intensity (Supplemental Figure 1 B; Unpaired t-

test **P=0.0034), supporting that displacement of neuromasts is a consequence of mechanical

injury. Additionally, we examined hair cell morphology in the ears of larvae exposed to sustained

stimulus and observed no apparent damage (Supplemental Fig. 2), indicating our

overstimulation protocol produces mechanical damage specifically to lateral-line organs.

To determine if hair-cell activity plays a key role in the displacement of neuromasts, we

exposed *lhfpl5b* mutants—fish that have intact hair cell function in the ear, but no

mechanotransduction in hair cells of the lateral line—to sustained stimulation (Erickson,

Pacentine, Venuto, Clemens, & Nicolson, 2019). We observed comparable morphological

disruption of mutant neuromasts lacking mechanotransduction (Fig. 2 G), suggesting that

displacement of lateral-line hair cells is due to physical damage from the stimulus. Further, we

observed the adjacent supporting cells in neuromasts with disrupted hair-cell morphology

appeared similarly displaced and elongated (Fig. 2 E "disrupted"; white arrows), indicating that

mechanical injury disrupts the structural integrity of the entire neuromast organ.

Hair-cell loss and reduced afferent innervation correspond to neuromast disruption

Moderate noise exposures can cause damage or loss of cochlear hair-cell synapses,

including swelling and retraction of afferent nerve fibers, while more extended and/or severe

exposures lead to hair-cell loss. To address whether mechanical damage in the lateral line

produced similar morphological changes, we surveyed the number of hair cells per neuromast

and the percentage of neuromast hair cells lacking afferent innervation in fish immediately

following exposure to sustained stimulation (Fig. 3). We observed a reduction in the number of

hair cells per neuromast immediately following exposure (Fig. 3 D, E; **Adj P=0.0019; N=13

trials) as well as a significant reduction in the percentage of hair cells per neuromast innervated

by afferent neurons (Fig. 3 F, G; ****Adj P<0.0001; N=12 trials).

As described in Fig. 2 F, we found on average ~half of L3-L5 neuromasts examined

showed "disrupted" hair-cell morphology immediately following sustained stimulus exposure. To

define the associations between overall neuromast morphology and specific structural changes

in mechanically injured neuromasts, we examined the numbers of hair cells and the percent of

hair cells contacted by afferent fibers in exposed neuromasts parsed into "normal" and

"disrupted" morphologies. With hair cell number, we observed significant loss specifically in

"disrupted" neuromasts, while "normal" neuromast hair-cell number appeared comparable to

control (Fig. 3 D, E; Adj P=0.3859 normal, ****Adj P<0.0001 disrupted). With hair-cell afferent

innervation we observed a similar trend i.e., a significant number of hair cells lacked afferent

innervation in stimulus exposed neuromasts with "disrupted" morphology, but not "normal"

neuromasts (Fig. 3 F,G; Adj P=0.7503 normal, ****Adj P<0.0001 disrupted).

Figure 3: Hair-cell loss and de-innervation is specific to "disrupted" neuromasts.
(A-C) Representative maximum intensity projection images of control (A) or exposed lateral line neuromasts with "normal" (B) or "disrupted" (C) morphology immediately following sustained strong wave exposure (0h). Synaptic ribbons (magenta; Ribeye b) and hair cells (blue; Parvalbumin) were immunolabled. Afferent neurons were expressing GFP. Scale bar: 5 μm **(D)** Hair-cell number per neuromast immediately post exposure. A significant reduction in hair cell number was observed (**Adj P=0.0019) and was specific to "disrupted" neuromasts (Adj P=0.3859 normal, ****Adj P<0.0001 disrupted). Pink box plot (Exp) represents pooled exposed neuromasts, while gray (Norm) and red (Dis) plots represent neuromasts parsed into normal and disrupted groups. Numbers beneath each plot indicate the number of neuromasts per group. Whiskers = min to max **(E)** Differences of least squares means in hair cell number per neuromast between groups. Bars represent 95% confidence interval (CI). **(F)** Percentage of neuromast hair cells innervated by afferent nerves. Numbers within each bar indicate the number of neuromasts per group. A significant portion of neuromast hair cells lacked afferent innervation following exposure (****Adj P<0.0001). Hair cells lacking afferent innervation were specifically observed in disrupted neuromasts (Adj P=0.7503 normal, ****Adj P<0.0001 disrupted). **(G)** Differences of least squares means in % hair cells innervated per neuromast between groups. Bars represent 95% CI

Mechanically overstimulated neuromasts with "normal" morphology lose a greater number of hair-cell synapses when glutamate clearance is inhibited

Hair cells contain electron-dense presynaptic specializations—called dense bodies or synaptic ribbons—apposing afferent PSDs which constitute afferent synaptic contacts (Davies, Tingley, Kachar, Wenthold, & Petralia, 2001; Sheets, Trapani, Mo, Obholzer, & Nicolson, 2011).

***Figure 4: Significant hair-cell synapse loss is observed in "normal" neuromasts following mechanical overstimulation and exacerbated by blocking glutamate uptake.* A-C)** Representative maximum intensity projection images of unexposed (A), or stimulus exposed lateral-line neuromast with "normal" (B) or "disrupted" (C) morphology. Synaptic ribbons (magenta; Ribeye b), PSDs (green; MAGUK) and hair cells (blue; Parvalbumin) were immunolabled. Scale bar: 5 µm **(D-E)** Intact synapses per neuromast hair cell. Pink box plot in D (Exp) represents pooled exposed neuromasts while, in E, gray (Norm) and red (Dis) plots represent neuromasts parsed into normal and disrupted groups. Whiskers =min to max. The average number of intact synapses per hair cell was significantly reduced in exposed neuromasts (D; **Adj P=0.0078); when parsed, this reduction was significant in the "normal" exposure group relative to control (E; **Adj P=0.0043 normal, Adj P=0.1207 disrupted). **(F)** Differences of least squares means in number of intact synapses per hair cell between groups. Bars represent 95% CI. **(G)** The number of intact synapses per hair cell in larvae co-treated with TBOA, to block glutamate clearance, or drug carrier alone during exposure. Synapse loss was significantly greater in "normal" neuromasts co-exposed to TBOA compared to fish co-exposed to the drug carrier alone (Two-way ANOVA. *P<0.0187).

In the larval zebrafish lateral line, afferent nerve fibers innervate multiple hair cells per

neuromast forming ~3-4 synaptic contacts per hair cell. To determine whether strong water

wave stimulus exposure generated lateral-line hair-cell synapse loss, we counted the number of

intact synapses (ribbons juxtaposed to PSDs; Fig. 4 A-C) in control and exposed larvae. We

observed significant reduction in the number of intact synapses per hair cell following sustained

exposure (Fig. 4 D, F; **Adj P=0.0078). When we

compared "normal" and "disrupted" neuromasts

following exposure, we observed a loss of intact

synapses per hair cell in all exposed neuromasts,

with significantly fewer synapses in "normal"

exposed neuromasts relative to control (Fig. 4 E-

F; **Adj P=0.0043 normal, Adj P=0.1207

disrupted). Additionally, significant synapse loss

was observed in the neuromasts of fish exposed

to the less mechanically damaging "periodic"

stimulus (Supplemental Fig. 1 F').

Previous studies indicate that excess

glutamate signaling may be a key factor driving

inner hair-cell synapse loss following exposure to

damaging noise (Chen, Kujawa, & Sewell, 2010;

Kim et al., 2019). We therefore inhibited glutamate

clearance from neuromast hair-cell synapses by

pharmacologically blocking uptake with the

glutamate transporter antagonist Threo-beta-

benzyloxyaspartate (TBOA) during sustained

stimulus exposure. We observed a significantly

greater degree of hair-cell synapse loss in

stimulated neuromasts with "normal" morphology

co-treated with TBOA than in stimulated

neuromasts co-treated with drug carrier alone

(Fig. 4 E; Two-way ANOVA. *P<0.0187),

suggesting glutamate excitotoxicity contributes to

hair-cell synapse loss observed in mechanically

overstimulated neuromasts with intact

morphology.

To further characterize synapse loss in

relation to afferent neurite retraction, we

examined fish in which we immunolabeled

synaptic ribbons, PSDs, afferent nerve fibers, and

Figure 5: Mechanically overstimulated neuromasts showed retracted neurites and detached synaptic debris. **(A-B)** Representative images of control (A) and exposed (B) neuromasts. Synaptic ribbons (magenta; Ribeye b) and PSDs (green; MAGUK) were immunolabeled; hair cells were also immunolabeled, but not shown for clarity. Afferent neurons (white) were labeled with GFP. Insets: Arrows indicate intact synapses adjacent to afferent neurons; arrowheads (B) indicate synaptic debris. Scale bars: 5μm (main panels), 1μm (insets). **(C)** Frequency histogram of observed synaptic debris per neuromast (NM). While control neuromasts occasionally had 1 detached synapse, exposed neuromasts were observed that had up to 5 detached synapses.

hair cells (Fig. 5 A,B). We then quantified instances where synapses, i.e. juxtaposed pre- and

postsynaptic components associated with hair cells, were no longer adjacent to an afferent

nerve terminal. As these synapses appeared detached and suspended from their associated

neurons, making them no longer functional, we refer to them as synaptic debris. While we rarely

observed synaptic debris in unexposed neuromasts, we observed a greater relative frequency

of synaptic debris in neuromasts exposed to strong water wave stimulus (Fig. 5 C; One sample

Wilcoxon test, P=0.1250 (control), *P=0.0313(normal), **P=0.0039(disrupted)). Taken together,

we observed two distinct types of morphological damage to afferent synapses in mechanically

overstimulated neuromasts: loss of synapses within neuromasts that appear intact that is

exacerbated when glutamate uptake is blocked and a higher incidence of synaptic debris that

appear detached from afferent neurites in all exposed neuromasts.

Mechanically overstimulated lateral line neuromasts show signs of hair-cell injury and

macrophage recruitment

240   The inner ears of birds and mammals possess resident populations of macrophages,

and additional macrophages are recruited after acoustic trauma or ototoxic injury (Warchol,

2019). A similar macrophage response occurs at lateral line neuromasts of larval zebrafish after

neomycin ototoxicity (Warchol, Schrader, & Sheets, 2021). Analysis of fixed specimens, as well

as time-lapse imaging of living fish (e.g., Hirose, Rutherford and Warchol, 2017), has

demonstrated that macrophages migrate into neomycin-injured neuromasts and actively

phagocytose the debris of dying hair cells. To determine whether a similar inflammatory

response also occurs after mechanical injury to the lateral line, we characterized macrophage

behavior after sustained stimulation. These studies employed Tg(*mpeg1:yfp*) transgenic fish,

which express YFP in all macrophages and microglia. Fish were fixed immediately after

exposure, or allowed to recover for 2, 4 or 8 hours. Control fish consisted of siblings that

received identical treatment but were not exposed to mechanical stimulation. Data were

obtained from the two terminal neuromasts from the pLL of each fish (Fig. 6 A). In agreement

with data shown in Fig. 3 D, we observed a modest but significant decline in hair-cell number in

specimens that were examined immediately and at 2 hours after sustained exposure (Fig. 6 B;

**P<0.003). Consistent with earlier studies (Hirose, Rutherford, & Warchol, 2017), 1-2

macrophages were typically present within a 25 μm radius of each neuromast (Fig. 6 C). In

uninjured (control) fish, those macrophages remained outside the sensory region of the

neuromast and rarely contacted hair cells. However, at 2, 4 and 8 hours after sustained

stimulus, we observed increased macrophage-hair cell contacts (Fig. 6 D; *P=0.024), as well as

the presence of immunolabeled hair-cell debris within macrophage cytoplasm (suggestive of

Figure 6. Macrophage response to mechanical overstimulation of lateral line hair cells. Experiments used fish that express YFP under regulation of the macrophage-specific *mpeg1* promoter. All images and data were collected from the two distal-most neuromasts of the posterior lateral line (Fig 2A; term). **(A)** Macrophages (green) responded to mechanical injury by entering neuromasts, contacting hair cells and internalizing otoferlin-labeled debris (arrows, magenta). Images show examples of macrophage behavior at different time points after noise trauma. **(B)** Quantification of hair cell number, which was significantly reduced at 0-2 hr after noise exposure (Mixed-effects analysis: **P<0.003). **(C)** Quantification of macrophages within a 25 μm radius of the terminal neuromasts at 0-8 hr after noise injury. Most neuromasts possessed 1-2 nearby macrophages and this number was not changed by noise exposure. **(D)** Quantification of direct contacts between macrophages and hair cells. Macrophage-hair cell contacts were counted at each survival time after noise exposure and normalized to the total number of sampled neuromasts. Increased levels of contact were observed at 2 and 4 hr after noise (*P=0.0243). **(E)** Quantification of phagocytosis as a function of post-noise survival time. Macrophages that had internalized otoferlin-labeled material were counted at each time point and normalized to the total number of sampled neuromasts. The percentage of macrophages that contained such debris was significantly increased at 0-2 hr after strong water wave stimulus (*P=0.0465; **P=0.0013). Data were obtained from 26-50 neuromasts/time point. Error Bars = SD

phagocytosis, Fig. 6 A, 4 h., arrow). Macrophage-hair cell contact and phagocytosis peaked at 2

hours after exposure (Fig. 6 E; **P= 0.0013 (2 h)). Notably, the numbers of macrophages within

a 25 μm radius of each neuromast remained unchanged at all time points after exposure,

suggesting that the inflammatory response was mediated by local macrophages and that

mechanical injury did not recruit macrophages from distant locations (Fig. 6 C). This pattern of

injury-evoked macrophage behavior is qualitatively similar to the macrophage response

observed in the mouse cochlea after acoustic trauma (Hirose et al., 2005; Kaur et al., 2019;

Kaur et al., 2015).

***Figure 7: Mechanically overstimulated neuromasts recover hair-cell morphology, hair-cell number, and innervation.* (A,B)** Average percentage of exposed neuromasts with "normal" vs. "disrupted" morphology following exposure. Each dot represents the percentage of disrupted neuromasts (L3-L5) in a single experimental trial; lines connect data points from the same cohort of exposed fish following 2 hours (A) or 48 hours (B) recovery. **(C,D)** Multilevel analysis of hair-cell number per neuromast immediately (0 h) post-exposure or after 2 or 48 hour recovery. Numbers beneath each plot indicate the number of neuromasts per group. Whiskers =min to max. Morphologically "disrupted" neuromasts has significantly fewer hair cells at 0h but not 2h following exposure (C; *Adj P=0.0321 (0h disrupted), Adj P=0.1875 (2h disrupted). Most exposed neuromasts were morphologically "normal" following 48h recovery and had a comparable number of hair cells relative to control (D; Adj P=0.4443). **(E,F)** The percentage of "disrupted" neuromast hair cells lacking afferent innervation was significant following 0h and 2h recovery (E; ****Adj P<0.0001 (0h disrupted), **Adj P=0.0016 (2h disrupted)). All hair cells were fully innervated following 48h recovery, including the few neuromasts with "disrupted" morphology (F; aligned red dots).

Mechanically injured neuromasts rapidly repair following exposure

To determine if damage to mechanically injured neuromasts was progressive, persistent,

or reversible, we compared neuromast morphology, hair-cell number, and innervation at 0

hours, 2 hours, and 48 hours following sustained exposure to strong water wave stimulus. We

observed a decrease in the percentage of neuromasts showing "disrupted" morphology 2 hours

following exposure, relative to fish fixed immediately following exposure (Fig. 7 A; 54%

disrupted (0 h) vs. 32% disrupted (2 h); N=4 trials), suggesting that physical disruption of

neuromast morphology following mechanical injury is rapidly reversible. Consistent with this

observation, the average hair-cell number per neuromast at 2 hours post-exposure appeared to

recover (Fig. 7 C; *Adj P=0.0321 (0h disrupted), Adj P=0.1875 (2h disrupted); N=6 trials).

Recovery of hair-cell number occurred within 2-4 hours (Fig. 6 A, B) and corresponded with

macrophages infiltrating neuromasts and phagocytosing hair-cell debris (Fig. 6 A, E). We also

observed, compared to immediately following exposure, a lesser degree of afferent fiber

retraction (Fig. 7 E; ****Adj P<0.0001 (0h disrupted), **Adj P=0.0016 (2h disrupted); N=6 trials)

indicating partial recovery of innervation.

A recent study characterized zebrafish lateral-line hair-cell damage induced by exposure

to ultrasonic waves and reported a delayed hair-cell death and synapse loss 48-72 hours

following exposure (Uribe et al., 2018). To determine if lateral line neuromasts exposed to the

strong wave stimulus generated by our apparatus underwent delayed hair-cell loss, we

examined hair-cell morphology, number, and innervation 48 hours following sustained stimulus

exposure. Most exposed neuromasts examined showed "normal" HC morphology (Fig. 7 B) with

no significant difference in hair-cell number (Fig. 7 D; Adj P=0.4443; N=4 trials). Hair-cell

afferent innervation after 48 hours was comparable to control fish; even the few neuromasts that

remained morphologically "disrupted" were fully innervated (Fig. 7 F).

Figure 8: Changes in synaptic ribbon and PSD sizes following sustained mechanical overstimulation. **(A-A'')**
Representative images of control (A) and exposed (A', A") neuromasts. Synaptic ribbons (magenta; Ribeye b),
PSDs (green; MAGUK), and hair cells (blue, Parvalbumin) were immunolabeled. Scale bars: 5µm (main panels),
1µm (insets). **(B-E)** Box and whisker plots of relative synapse volumes normalized to 0h control. Whiskers indicate
the min. and max. values; "+" indicates the mean value, horizontal lines indicate the relative median value of the
control. (B) Ribbon volume appeared comparable to control immediately following exposure but was reduced 2
hours after exposure (*P=0.0195). (C) Significant reduction in ribbon size relative to control was specific to disrupted
neuromasts (Kruskal-Wallis test: ***P=0.0004 (2h)). (D) Significantly larger PSDs were observed both immediately
and 2 hours following exposure (****P<0.0001). (E) Enlarged PSDs were present in both "normal" and "disrupted"
exposed neuromasts, with a greater enlargement observed 0h post-exposure (Kruskal-Wallis test: ****P<0.0001 (0h)
; ***P=0.0001, **P=0.0024 (2h)).

PSDs are enlarged in all neuromasts following mechanical overstimulation

Previous studies in mice and guinea pigs indicate moderate noise exposures modulate

the size of synaptic components (Kim et al., 2019; Song et al., 2016). To determine if pre- and

postsynaptic components were also affected in our model, we compared the relative volumes of

neuromast hair-cell presynaptic ribbons and their corresponding PSDs in control and stimulus

exposed larvae. We observed a moderate reduction in synaptic-ribbon size following exposure;

ribbon volumes were significantly reduced relative to controls following 2 hours recovery (Fig. 8

B; Kruskal-Wallis test *P=0.0195; N=3 trials), and this reduction was specific to "disrupted"

neuromasts (Fig. 8 C). While the changes in ribbon volume we observed were modest and

delayed in onset, we saw dramatic enlargement of PSDs immediately and 2 hours following

exposure (Fig. 8 D; Kruskal-Wallis test ****P<0.0001; N=3 trials). In contrast to the observed

reduction in ribbon size, relative PSD volumes were significantly enlarged in all exposed

neuromasts regardless of whether neuromast morphology was "normal" or "disrupted" (Fig. 8

E). These data reveal enlarged PSDs as the predominant structural change in mechanically

overstimulated neuromast hair-cell synapses.

Mechanically injured neuromasts have damaged kinocilia, disrupted hair-bundle

morphology, and reduced FM1-43 uptake immediately following exposure

An additional consequence of excess noise exposure in the cochlea is damage to

mechanosensitive hair bundles at the apical end of hair cells and, correspondingly, disruption of

mechanotransduction (Wagner & Shin, 2019). Larval zebrafish lateral-line hair cells each have a

hair bundle consisting of a single kinocilium flanked by multiple rows of actin-rich stereocilia

(Kindt, Finch, & Nicolson, 2012). To determine if our exposure protocol damaged apical hair-cell

structures, we used confocal imaging and scanning electron microscopy (SEM) to assess hair

bundle morphology in both unexposed control larvae and larvae fixed immediately following

sustained exposure. All neuromasts throughout the fish were evaluated, but to remain

consistent with our fluorescence imaging results, we closely assessed the appearance of the

caudal pLL neuromasts. We found the caudal neuromasts to be more damaged than the ones

positioned more rostrally: the frequency of neuromasts with apparently disrupted appearance

increased the closer its position to the tail (Supplemental Fig. 3). This is consistent with our

fluorescence observations (Fig. 2 F; Supplemental Fig. 4 B) in which L5 neuromasts were more

likely to be disrupted than more anteriorly positioned L3.

A closer examination of neuromast morphology revealed a difference of the kinocilia

length and bundling. The neuromasts of the control fish carry a bundle of long (10-15 µm),

uniformly shaped kinocilia (Fig. 9 A-E). In contrast, the neuromasts of the fish fixed immediately

Control (no exposure)

Mechanical stimulation (0hr recovery)

Figure 9: Scanning electron microscopy imaging of neuromasts following mechanical injury reveals disorganized hair cell stereocilia bundles and damaged kinocilia. **(A-E)** Representative images of tail neuromasts of control fish larvae. Each hair cell carries a kinocilium, which is visibly thicker than its neighboring actin-filled, mechanosensitive stereocilia: see panel C featuring both structures at higher magnification (the kinocilium diameter is 220 nm, while stereocilia measured 90-110 nm). The kinocilia of control neuromasts are long (10-15 µm) and bundled together, while the stereocilia bundles have an apparent staircase arrangement. **(F-K)** Representative images of damaged tail neuromasts immediately following noise exposure featuring short (F-H, yellow arrows), disorganized (G, H), and swollen (I-K, yellow arrowheads) kinocilia, and disorganized stereocilia. (K) Same stereocilia bundle as in J marked with an asterisk at higher magnification to highlight the difference in the diameter of the kinocilium (360 nm) and neighboring stereocilia (85-100 nm) for noise exposed hair cells, as compared to the control hair cells in C. Scale bars: A, B, D-J – 2 µm; C, K – 500 nm. **(L-M)** Kinocilia diameter at bundle level (L; Mann Whitney test ***P=0.0007) and 3-5 µm above bundle level (M; Welch's t test *P=0.0243). Exposed NM data in L were not normally distributed (D'Agostino-Pearson test ****P<0.0001). Error Bars = SD

after sustained exposure often appear to carry much shorter kinocilia (Fig. 9 F-H, yellow arrows), which lack bundling and, in some cases, pointing to different directions (Fig. 9 G,H). The apparent kinocilia length difference between control and overstimulated neuromasts suggests at least some kinocilia may undergo a catastrophic damage event at the time of stimulation, as their distal parts break off the hair cells. This is further supported by some

Figure 10: Hair cell mechanotransduction was significantly reduced but rapidly recovered following mechanical overstimulation. (A-C) Representative images of hair-cell stereocilia (conjugated phalloidin, gray) and FM1-43FX fluorescence intensity of the corresponding neuromast in representative control (A) or mechanically overstimulated fish immediately (B, B') or 4 hours (C) following exposure. Yellow arrows in (B') indicate phalloidin labeling that appeared tapered. **(D)** Average relative FM1-43FX fluorescence intensity measurements in control and exposed neuromasts over 4 hours of recovery. FM1-43FX uptake was significantly reduced in exposed neuromasts immediately following mechanical overstimulation but appear to completely recover by 4 hours (Tukey's multiple comparisons test ****P<0.0001 (0h), P=0.0579 (1h), P=0.8387 (2h), P=0.8387 (4h)). Dashed lines indicate FM1-43FX fluorescence intensity measurements in exposed neuromasts parsed into "normal" and "disrupted" morphologies. **(E)** Average stereocilia length of centrally localized hair bundles in control and exposed neuromasts. Dashed lines indicate measurements in exposed neuromasts parsed into "normal" and "disrupted" morphologies. Error Bars = SD **(F)** Relative FM1-43FX fluorescence in both "normal" and "disrupted" exposed neuromasts was significantly reduced immediately following exposure but recovered over time (Tukey's multiple comparisons test ****P<0.0001, *P=0.0328 (0h); ****P<0.0001, **P=0.0098 (1h); **P=0.0025 control vs. dis, **P=0.0089 normal vs. dis (2h)). Each point represents an individual neuromast. Nearly all observed exposed neuromasts appeared morphologically normal following 4 hours; note only one neuromast data point in the disrupted category of the 4h recovery graph. Data were obtained from 26-32 neuromasts per condition over three trials.

examples of kinocilia with thicker, 'swollen' proximal shafts closer to the cuticular plate of the

cell, some of which extend into a thinner distal part while others appear to lack the distal part

completely (Fig. 9 I-K, yellow arrowheads). Accordingly, the average diameter of kinocilia at the

level of the hair bundle (L2-5) was significantly larger than control, with a few kinocilia showing

dramatically thicker widths ~2x greater than the thickest control (Fig. 9 L; ***P=0.0007). When

measured ~3-5 μm above the bundle, the exposed neuromast kinocilia have a somewhat larger

average diameter relative to control, but not as dramatic as observed at the base (Fig. 9 M;

*P=0.0243). Stereocilia bundles from both groups of animals carried tip links, but we were

unable to systematically evaluate and quantify their abundance. However, we observed signs of

damaged bundle morphology following overstimulation, as they often appeared splayed, with

gaps between the rows of stereocilia (Fig. 9 I-K).

Figure 11: Neuromasts show no change in cell proliferation following mechanical overstimulation. **(A,B)**
Representative cross-section images of EdU (magenta) labeling of proliferating neuromast cells. Fish were
exposed to EdU for 4 hours following stimulus exposure. Supporting cells (SC) were expressing GFP. Scale
bars: 5μm **(C)** Average number of EdU+ cells per neuromast were comparable in control and exposed larvae.
Data were obtained from 33-34 neuromasts per condition over three trials. (Two-way ANOVA. P=0.4193) Bars
represent 95% CI.

To evaluate the effect of mechanical overstimulation on hair bundle function in relation to

hair bundle morphology, we assessed mechanotransduction by briefly exposing free-swimming

larvae to the fixable mechanotransduction-channel permeable dye FM1-43X (Holmgren &

Sheets, 2021; Toro et al., 2015). We treated control and mechanically overstimulated larvae

with FM1-43X immediately and at several time points up to 4 hours following exposure, then

fixed the larvae and co-labeled hair-cell stereocilia with fluorescently-conjugated phalloidin (Fig.

10 A-C). We observed a significant reduction in the relative intensity of FM1-43 in all exposed

neuromasts immediately following exposure (Fig. 10 D, F (0h); ****P<0.0001). While phalloidin

labeling of stereocilia revealed what appeared to be tapered hair bundles in some exposed

neuromasts (Fig. 10 B'; yellow arrows), average stereocilia length obtained from 3D interpolated

confocal image stacks was not significantly altered (Fig. 10 E). Remarkably, FM1-43FX uptake

showed recovery within 30 minutes and fully recovered over several hours. (Fig. 10 C, D). The

degree of FM1-43FX fluorescence recovery following mechanical damage appeared to

correspond with recovery of neuromast morphology; following 4 hours, nearly all neuromasts

exposed to strong wave stimulus showed "normal" morphology and relative FM1-43X

fluorescence that was comparable to control (Fig 10 D, F, 4h recovery). This observed timeline

of morphological recovery coincides with macrophage recruitment and phagocytosis hair cell

debris peaking 2 hours post exposure (Fig 6 D,E) followed by full recovery of hair cell number

between 2-4 hours post exposure (Fig 6 B). Additionally, we quantified proliferating neuromast

cells by treating control and mechanically overstimulated 5-ethynyl-2-deoxyuridine (EdU) larvae

for 4 hours following exposure and saw no difference in the number of EdU positive cells per

neuromast in each condition (Fig 11 A-C). Cumulatively, these data support that most damaged

hair cells are repaired within mechanically injured neuromasts.

Discussion

To model mechanical injury resulting from noise trauma in the zebrafish lateral line, we

describe here a method to mechanically overstimulate neuromasts of the posterior lateral line.

Using this method, we observed: i) hair-cell synapse loss in a subset of stimulus exposed

neuromasts with intact morphology, ii) morphological displacement, hair-cell loss, and afferent

deinnervation in a subset of mechanically disrupted neuromasts, iii) an inflammatory response

that peaked 2-4 hours following stimulus exposure, iv) kinocilia and hair bundle damage, and v)

reduced FM1-43 uptake in all immediately following exposure. Remarkably, mechanically

injured neuromasts rapidly recover following exposure; neuromast morphology, innervation, and

mechanotransduction showed significant recovery within an hour, and most neuromasts were

completely recovered within 4 hours post exposure.

Zebrafish lateral-line as a model for sub-lethal mechanical damage and noise-induced

synapse loss

Mechanical damage to the zebrafish lateral line induced by strong water wave stimulus

is observable immediately following exposure, is specific to the lateral-line organ, and appears

to be rapidly repaired. These observations contrast with a recently published noise damage

protocol for larval zebrafish which used ultrasonic transducers (40-kHz) to generate small,

localized shock waves (Uribe et al., 2018). They reported delayed hair-cell death and modest

synapse loss that was not apparent until 48 hours following exposure, was not accompanied by

decreased mechanotransduction, and was observed in the inner ear as well as the lateral-line

organs. Some of the features of the damage they observed—delayed onset apoptosis and hair-

cell death—may correspond to lethal damage following blast injuries. We propose features of

the damage we observe with our stimulus protocol—reduced mechanotransduction, hair-cell

synapse loss, and rapid inflammatory response—may correspond to sub-lethal noise-induced

damage of hair-cell organs. Because of differences in the nature of the stimuli in these two

studies, it is difficult to directly compare the pathological outcomes. Mechanical overstimulation

in the Uribe et al., study was induced using ultrasonic (40 kHz) actuators. Such high frequencies

are far outside those that are detected by lateral line neuromasts (Levi et al., 2015; Trapani &

Nicolson, 2010) supporting that cavitation in the water medium is likely causing the observed

damage. In contrast, our study delivered a stimulus of high intensity 60 Hz water waves directly

to the fish. This frequency is within the range of sensitivity of lateral line neuromasts of larval

zebrafish and evoked a lateral-line mediated behavior (Fig 1 B, C) suggesting that the hair cells

were being directly stimulated by the water motion. The present method more closely resembles

the techniques that are typically used to study noise damage in the mammalian cochlea, where

high intensity acoustic energy causes hair cell and synaptic injury in specific regions of the

cochlea that are best-responsive to the frequency of the stimulus. This idea is further supported

by the observation that synapse loss in hair cells exposed to strong wave stimulation is greater

when glutamate uptake is blocked (Fig. 4 E), suggesting a shared mechanism of glutamate

excitotoxicity between noise-exposed mammalian ears and strong water wave stimulus exposed

lateral-line organs (Kim et al., 2019; Sebe et al., 2017).

Disruption of neuromast morphology is a consequence of mechanical injury

Strong water wave exposure produced a percentage of pLL neuromasts that were

morphologically "disrupted". Several observations support that such "disrupted" neuromasts

represent mechanical injury to neuromast organs. One is that *lhfpl5b* mutant neuromasts, which

lack mechanotransduction specifically in lateral-line hair cells, were comparably vulnerable to

physical disruption as their wild-type siblings (Fig. 2 G; (Erickson et al., 2019)). This finding

supports that hair-cell activity during stimulation does not underly the physical displacement of

hair cells observed following strong water wave stimulus. Additionally, physical disruption of the

neuromast affects the whole organ—hair cells and their adjacent supporting cells (Fig. 2 D).

This observation contrasts with what is observed in mammalian ears exposed to high intensity

noise, where mechanical injury to outer hair cells is localized to stereocilia disruption (H. Wang,

Yin, Yu, Huang, & Wang, 2011). Speculatively, displacement of hair cells in mechanically

injured neuromasts may be due to loss of structural support from displaced supporting cells. As

the lateral-line organs are superficially localized on the surface of the skin, some of the intense

mechanical tension applied across the tail is likely coupled to neuromasts leading to the

physical displacement of a subset of exposed neuromasts. Finally, we observed reduced FM1-

43 uptake in "disrupted" neuromasts (Fig. 10 G) and measurable changes kinocilia likely

reflecting mechanical damage (Fig. 9; (Wagner & Shin, 2019)). It is notable how rapidly

mechanically disrupted lateral-line neuromasts regain normal morphology and FM1-43 uptake

(Fig. 10). The superficial lateral line's direct exposure to the environment may require more

robust mechanisms for repair, unlike hair cell organs of mammals which are encased in bone.

Hair-cell overstimulation and synapse loss

In contrast to de-innervation and modest hair-cell loss we observed in mechanically

disrupted neuromasts, we saw significant loss of hair-cell synapses in neuromasts that were

exposed to strong water wave stimulus but not mechanically disrupted i.e. "normal" (Fig. 4). Two

notable observations were made in exposed neuromasts regarding synapse loss. First, loss of

synapses in "normal" exposed neuromasts was markedly more severe when synaptic glutamate

clearance was inhibited (Fig. 4 G), suggesting that synapse loss may reflect moderate hair-cell

damage resulting from overstimulation of hair cells and excess glutamate accumulation.

Involvement of glutamate signaling as a key mediator of noise-induced synapse loss has

recently been reported in mice; loss of glutamate signaling prevents noise-induced synapse loss

and pharmacologically blocking postsynaptic Ca^{2+} permeable AMPA receptors protects against

cochlear hair-cell synapse loss from moderate noise exposure (Hu et al., 2020; Kim et al.,

2019). Second, synapse loss occurred in neuromasts that appeared to be fully innervated (Fig.

3 B, Fig. 4 B). This observation was initially surprising given that pharmacologically activating

evolutionarily conserved Ca^{2+} permeable AMPARs has been shown to drive afferent terminal

retraction in the zebrafish lateral line (Sebe et al., 2017). We propose that subtle damage to

afferents in "normal" exposed neuromasts may accompany synapse loss but not be apparent as

loss of innervation, as single afferent processes innervate multiple hair cells of the same polarity

within an individual neuromast (Dow, Jacobo, Hossain, Siletti, & Hudspeth, 2018; Faucherre,

Pujol-Marti, Kawakami, & Lopez-Schier, 2009). This idea is further supported by our observation

that the relative frequency of synaptic debris (i.e. synaptic components that appear detached

from afferent neurites) was higher in exposed neuromasts with "normal" morphology relative to

control (Fig. 5 C). We speculate that synaptic debris observed in exposed neuromasts with

"normal" morphology may be the result of excitotoxic damage at synaptic terminals (Sebe et al.,

2017) while synaptic debris observed in "disrupted" neuromasts may reflect mechanical

disruption of supporting cells and retraction of afferent innervation from a subset of hair cells

(Fig 3 C, F).

In addition, glutamate signaling may not be the only driver of synapse loss resulting from

excess stimulation. Studies in zebrafish and mice support that mitochondrial stress combined

with excessive synaptic activity may also contribute to hair-cell synapse loss (X. Wang et al.,

2018; Wong et al., 2019). Future work using this model to examine the effect of excess

mechanical stimulation on mutants with reduced glutamate release or impaired glutamate

clearance (to elevate or reduce glutamate in the synaptic cleft, respectively) combined with

modified mitochondrial function may define the relative roles of glutamate excitotoxicity and hair-

cell mitochondrial stress to synaptic loss.

Role of inflammation following mechanical injury to lateral-line organs

Our results indicate that mechanical injury to neuromasts evokes an inflammatory

response. Prior studies of larval zebrafish have shown that macrophages reside near the

borders of uninjured neuromasts and migrate into neuromasts after ototoxic injury (Hirose et al.,

2017). We found that macrophages migrate into neuromasts within ~2 hours of mechanical

injury, where they contact hair cells and, in some cases, engulf hair-cell debris. Although this

macrophage response is similar to that which occurs after ototoxic injury to neuromasts (Carrillo

et al., 2016; Hirose et al., 2017; Warchol et al., 2021), the extent of hair cell loss after

mechanical overstimulation is much less than the injury that occurs after ototoxicity. We

observed macrophage entry in 30-40% of exposed neuromasts, despite modest hair cell loss

(Fig. 6 B,D). It is possible that the morphological changes characteristic of mechanically injured

neuromasts are accompanied by the release of macrophage chemoattractants. In addition,

studies of noise exposure to the mammalian cochlea indicate that high levels of synaptic activity

(without accompanying hair cell loss) can evoke macrophage migration to the synaptic region

(Kaur et al., 2019). In either case, the signals responsible for such recruitment remain to be

identified. The observation that macrophages had internalized immunolabeled hair-cell material

further suggests that recruited macrophages engage in the phagocytosis of hair cell debris, but

it is not clear whether macrophages remove entire hair cells or target specific regions of cellular

injury (e.g., synaptic debris; Fig. 5). In any case, our data indicate that the macrophage

response to mechanical injury of zebrafish lateral-line neurons is similar to that which occurs

after noise injury to the mammalian cochlea (Warchol, 2019) and suggests that zebrafish may

be an advantageous model system in which to identify the signals that recruit macrophages to

sites of excitotoxic injury.

Hair-cell synapse morphology following mechanical overstimulation

Immediately following mechanical overstimulation, the most pronounced morphological

change we observed in hair-cell synapses was significantly enlarged PSDs (Fig. 8 D, E).

Speculatively, PSD enlargement may be a consequence of reduced glutamate release from hair

cells following sustained intense stimulation. Mice and zebrafish fish lacking hair-cell

glutamatergic transmission have enlarged postsynaptic structures (Kim et al., 2019; Sheets,

Kindt, & Nicolson, 2012), indicating that glutamate may regulate postsynaptic size. While our

data do not directly support this idea, we speculate that reduced glutamatergic transmission in

mechanically overstimulated neuromasts may be a consequence of transiently impaired

mechanotransduction (Fig. 10 B,D; (Zhang et al., 2018)). Alternatively, cholinergic efferent

feedback, which has been shown to hyperpolarize lateral-line hair cells, may reduce hair-cell

excitability during sustained strong wave exposure to protect against excess glutamate release,

excitotoxic damage, and synapse loss (Carpaneto Freixas et al., 2021). Interestingly,

presynaptic ribbons were not similarly enlarged, but instead showed a modest reduction in size

following mechanical injury. Functional imaging of zebrafish lateral line has shown that a subset

of hair cells in each neuromast are synaptically silent, and these silent hair cells can become

active following damage (Zhang et al., 2018). As ribbon size has been observed to correspond

with synaptic activity (Merchan-Perez & Liberman, 1996; Sheets et al., 2012), reduction in

ribbon size may reflect recruitment of synaptically silent hair cells following mechanically-

induced damage. Future functional studies are needed to determine if mechanical

overstimulation recruits more active hair-cell synapses, and to verify whether glutamate release

from active synapses is reduced following mechanical overstimulation.

Lateral-line neuromasts fully recover following mechanical damage

508   Previous studies indicate mammalian cochlear hair cells have some capacity for repair

following sub-lethal mechanical damage, including tip-link repair and regeneration of a subset of

ribbons synapses (Indzhykulian et al., 2013; Jia, Yang, Guo, & He, 2009; Kim et al., 2019). But

such ability is limited, and our understanding of hair-cell repair mechanisms is incomplete. By

contrast, we observe complete recovery of neuromast morphology and innvervation following

mechanical trauma to the zebrafish lateral line. The cellular mechanisms responsible for such

repair are not fully defined, but may involve regulation of neurite growth, glutamate signaling,

inflammation, and/or neurotrophic factors (Kaur et al., 2019; Kim et al., 2019; Wan, Gomez-

Casati, Gigliello, Liberman, & Corfas, 2014). Further study using this zebrafish model of

mechanical overstimulation may provide insights that will assist in the development of methods

for promoting complete hair-cell repair following damaging stimuli to the mammalian cochlea.

519   We also found that mechanical trauma resulted in a small degree of hair-cell loss in

disrupted neuromasts (Fig. 3D), but that hair-cell numbers had recovered after 2 hours (Fig.

6B). The mechanism that mediates this recovery is not clear, but it is notable that a low level of

hair cell production normally occurs in lateral line neuromasts of larval zebrafish, as part of a

process of ongoing turnover (Cruz et al., 2015; Williams & Holder, 2000). In the present study,

we observed a small amount of proliferation in neuromasts of both mechanically-damaged and

control fish. Since mechanical damage did not increase the level of cell proliferation in

neuromasts relative to control (Fig 11), we believe this observed cell division is likely associated

with the turnover process. Hair-cell regeneration in the vertebrate inner ear can also occur via

direct phenotypic conversion of supporting cells into a replacement hair cells (Warchol, 2011).

While it is conceivable that transdifferentiation of supporting cells could occur within 2 hours of

mechanical injury, such transdifferentiation has not been previously demonstrated in zebrafish

lateral line neuromasts (e.g. (Thomas et al., 2015)). Overall, these observations indicate that

mechanical trauma does not increase the rate of hair-cell production within neuromasts, further

supporting that neuromast recovery is largely due to hair-cell repair.

In summary, our data show that exposure of zebrafish lateral-line organs to strong water

wave results in mechanical injury and loss of afferent synapses, but that these injuries rapidly

recover. Our next steps will be to define the time course for synaptic recovery and to determine

how lateral-line mediated behavior is affected by mechanically induced damage. Sub-lethal

overstimulation of hair cells in the zebrafish lateral line provides a useful model for defining

mechanisms of damage and inflammation and for identifying pathways that promote hair-cell

repair following mechanically-induced injury.

Declaration of interests:

The authors declare no competing financial or non-financial interests.

Acknowledgments: This work was supported by the National Institute on Deafness and Other

Communication Disorders R01DC016066 (L.S.), R01DC017166 (A.A.I), and R01DC006283

(M.E.W.), Washington University Dept. of Otolaryngology (L.S.), and the Amelia Peabody

Charitable Fund (L.S.). We would like to thank Valentin Militchin (WashU) and Evan Foss (Mass

Eye and Ear) for engineering support and Mark Rutherford for thoughtful feedback on the

manuscript.

<u>Method Details</u>

Ethics Statement

This study was performed with the approval of the Institutional Animal Care and Use

Committee of Washington University School of Medicine in St. Louis and in accordance with

NIH guidelines for use of zebrafish.

Zebrafish

All zebrafish experiments and procedures were performed in accordance with the

Washington University Institutional Animal Use and Care Committee. Adult zebrafish were

raised under standard conditions at 27-29°C in the Washington University Zebrafish Facility.

Embryos were raised in incubators at 29°C in E3 media (5 mM NaCl, 0.17 mM KCl, 0.33 mM

$CaCl_2$, 0.33 mM $MgCl_2$; (Nüsslein-Volhard & Dahm, 2002)) with a 14 h:10 h light:dark cycle. After

4 dpf, larvae were raised in 100-200 ml E3 media in 250-ml plastic beakers and fed rotifers

daily. Sex of the animal was not considered in our studies because sex cannot be predicted or

determined in larval zebrafish.

The transgenic lines *TgBAC(neurod1:EGFP)* (Obholzer et al., 2008),

Tg(tnks1bp1:EGFP) (Behra et al., 2012), *Tg(-6myo6b:βactin-EGFP)* (Kindt et al., 2012) and

Tg(mpeg1:YFP) (Roca & Ramakrishnan, 2013) were used in this study. Fluorescent larvae were

identified at 3 dpf without anesthesia in E3 media. The mutant line *lhfpl5b$^{vo35/vo35}$* was also used

(Erickson et al., 2019).

Genotyping

To genotype *lhfpl5b$^{vo35/vo35}$* larvae and siblings after mechanical stimulation and

immunohistochemical labeling, ~1 mm tail tissue was excised, and genomic DNA was extracted

by incubation in a lysis buffer (10 mM Tris pH 8.0, 50 mM KCl, 0.3% NP-40, 0.3% Tween-20). A

genomic region of *lhfpl5b* was amplified by PCR using forward primer

GCGTCATGTGGGCAGTTTTC and reverse primer TAGACACTAGCGGCGTTGC. The

lhfpl5b^{vo35} mutation disrupts a MluCI restriction site (AATT), so PCR products were digested with

MluCI, and homo- and heterozygotes were resolved by differences in band size on a 1-1.5%

agarose gel.

Experimental Apparatus

Multi-well plates containing larvae were clamped to a custom magnesium head

expander (Vibration & Shock Technologies, Woburn, MA) on a vertically-oriented Brüel+Kjær

LDS Vibrator, V408 (Brüel and Kjær, Naerum, Denmark). An additional metal plate was fitted to

the bottom of the multi-well dish to fill a small gap between the bottoms of the wells and the

head expander to eliminate flexing of the well plate relative to the head expander. Vibrometry of

the well bottoms and the head expander with a laser-Dopper vibrometer (OFV-2600 and OFV-

501, Polytec, Irvine, CA) confirmed that the well plate and head expander motion were equal at

stimulus frequencies. This experimental apparatus was housed in a custom sound-attenuation

chamber. An Optiplex 3020 Mini Tower (Dell) with a NI PCIe-6321, X Series Multifunction DAQ

(National Instruments) running a custom stimulus generation program (modified version of

Cochlear Function Test Suite) was used to relay the stimulus signal to a Brüel+Kjær LDS

PA100E Amplifier that drove a controlled 60 Hz vibratory stimulus along the larvae's

dorsoventral axis (vertically). Two accelerometers (BU-21771, Knowles, Itasca, IL) were

mounted to the head expander to monitor the vertical displacement of the plate. The output of

the accelerometers was relayed through a custom accelerometer amplifier (EPL Engineering

Core). A block diagram for the EPL Lateral Line Stimulator can be found here:

https://www.masseyeandear.org/research/otolaryngology/eaton-peabody-

laboratories/engineering-core).

Mechanical overstimulation of lateral-line organs in free swimming larvae

At 7 dpf, free-swimming zebrafish larvae were placed in untreated 6-well plates

(Corning, Cat# 3736; well diameter: 34.8 mm; total well volume: 16.8 ml) with 6 ml E3 per well,

pre-warmed to 29°C. Up to 15 larvae were placed in each well. Individual wells were sealed with

Glad® Press 'n Seal plastic food wrap prior to placing the lid on the plate. An additional metal

plate was fitted to the bottom of the multi-well dish to fill a small gap between the bottoms of the

wells and the head expander.

Mechanical water displacement exposures (stimulus parameters: 60 Hz, 40.3 ± 0.5 m/s^2)

were conducted at room temperature (22-24°C) up to 2 hours after dawn. The frequency

selected for mechanical overexposure of lateral-line organs was based on previous studies

showing 60 Hz to be within the optimal upper frequency range of mechanical sensitivity of

superficial posterior lateral-line neuromasts (Weeg and Bass 2002, Trapani et al, 2009, Levi et

al, 2015). To confirm that 60 Hz was the optimal frequency to induce damage, we tested 45, 60,

and 75 Hz at comparable intensities. We observed at 75 Hz no apparent damage to lateral line

neuromasts while 45 Hz at a comparable intensity proved toxic i.e. it was lethal to the fish.

Exposures consisted of 20 minutes of stimulation followed by a 10-minute break and 2

hours of uninterrupted stimulation. We also tested periodic exposures that consisted of a series

of short pulses spanning 2 hours total: 2 20-minute exposures each followed by 10 minutes of

rest, followed by 30 minutes of stimulation, a 10-minute break, and a final 20 minutes of

stimulation. During the entire duration of exposure, unexposed control fish were kept in the

same conditions as exposed fish i.e. placed in a multi-well dish and maintained in the same

room as the exposure chamber. For experiments pharmacologically blocking glutamate uptake,

fish were co-exposed to 10 µM DL-TBOA (Tocris; Cat. No.1223) + 0.1% DMSO or 0.1% DMSO

alone. After exposure, larvae were either immediately fixed for histology, prepared for live

imaging, or allowed to recover for up to 2 days in an incubator at 29°C.

Ablation of lateral-line organ with CuSO$_4$

Free-swimming larvae were exposed to freshly made 3 µM CuSO$_4$ solution in E3 for 1

hour, then rinsed and allowed to recover for 2 hours to ensure complete ablation of the lateral-

line neuromasts. Neuromast ablation was confirmed by immunofluorescent labeling of hair cells.

The effects of low-dose copper exposure are likely specific to lateral-line organs; a previous

study in zebrafish determined exposure to low-dose CuSO$_4$ for 1 hour did not alter the acoustic

escape response, which is similar to the fast start response we observed but evoked by higher

frequency stimulation (100-500 Hz) of the anterior macula of inner ear (Buck, Winter, Redfern, &

Whitfield, 2012).

Fast-start escape response behavior assay

Images of larval swimming behavior (1000 frames per second) were acquired with an

Edgertronic SC1 high-speed camera (Sanstreak Corp). Image acquisition began 10 seconds

following stimulus onset. All subsequent analysis was performed using ImageJ. To track

swimming behavior, images were initially stabilized using the Image Stabilizer Plugin. In

stabilized images, the position of individual larval heads (located via the pigmented eyes) in

each frame were tracked using the Manual Tracking Plugin. Larvae were tracked over 10

seconds (10,000 frames total) per trial. 'Fast start' responses—defined as a c-bend of the body

occurring within 15 ms followed by a counter-bend— were identified manually.

Whole-mount immunohistochemistry

For visualization of zebrafish lateral-line hair cells, neurons, and synapses: 7-9 dpf

larvae were briefly sedated on ice, transferred to fixative (4% paraformaldehyde, 1% sucrose,

37.5 µM CaCl$_2$, 0.1 M phosphate buffer) in a flat-bottomed 2 ml Eppedorf tubes, and fixed for 5

hours at 4-8°C. Fixed larvae were permeabilized in ice-cold acetone for 5 minutes, then blocked

in phosphate-buffered saline (PBS) with 2% goat serum, 1% bovine serum albumin (BSA), and

1% DMSO for 2-4 hours at room temperature (RT; 22-24 °C). Larvae were incubated with

primary antibodies diluted in PBS with 1% BSA and 1% DMSO overnight at 4-8°C, followed by

several rinses in PBS/BSA/DMSO and incubation in diluted secondary antibodies conjugated to

Pacific Blue (1:400), Alexa Fluor 488 (1:1000), Alexa Fluor 555 (1:1000), Alexa Fluor 647

(1:1000; Invitrogen), or DyLight 549 (1:1000; Thermo-Fisher) for 2 hours at RT. In some

experiments, fixed larvae were stained with 2.5 ug/ml 4′,6-diamidino-2-phenylindole (DAPI;

Invitrogen) diluted in PBS to label all cell nuclei. Larvae were mounted on glass slides with

elvanol (13% w/v polyvinyl alcohol, 33% w/v glycerol, 1% w/v DABCO (1,4 diazobicylo[2,2,2]

octane) in 0.2 M Tris, pH 8.5) and #1.5 cover slips.

For visualization of inflammation and macrophage recruitment: 7 dpf larvae were

sedated on ice, transferred to 4% paraformaldehyde fixative in PBS, then fixed overnight at 4-

8°C. The next day larvae were rinsed in PBS and blocked in PBS with 5% normal horse serum

(NHS), 1% DMSO, and 1% Triton x-100 for 2 hours at RT. Larvae were incubated with primary

antibodies diluted in PBS with 5% NHS and 1% Triton-x 100 overnight at RT, rinsed several

times in PBS, then incubated in diluted secondary antibodies listed above for 2 hours at RT.

Larvae were mounted on glass slides with glycerol/PBS (9:1); coverslips were sealed with clear

nail polish.

Primary antibodies

The following commercial antibodies were used in this study: GFP (1:500; Aves Labs,

Inc; Cat# GFP-1020), Parvalbumin (1:2000; Thermo Fisher; Cat# PA1-933), Parvalbumin

(1:2000; Abcam; Cat# ab11427), Parvalbumin (1:500; Sigma-Aldrich; Cat# P3088), MAGUK

(K28/86; 1:500; NeuroMab, UC Davis; Cat# 75-029), Otoferlin (1:500; Developmental Studies

Hybridoma Bank/ HCS-1). Custom affinity-purified antibody generated against *Danio rerio*

Ribeye b (mouse IgG2a; 1:2000; (Sheets et al., 2011)) was also used.

Hair-cell labeling

To selectively label hair-cell nuclei, live zebrafish larvae were incubated with DAPI

diluted 1:2000 in E3 media for 4 minutes. Larvae were briefly rinsed 3 times in fresh E3 media,

then immediately exposed to mechanical overstimulation.

Live imaging of FM1-43 did not provide the temporal resolution needed to compare

relative uptake and fast recovery 0-4 hours following exposure. We therefore examined FM1-43

uptake using the fixable analogue. To label with FM1-43X (n-(3-triethylammoniumpropyl)-4-(4-

(dibutylamino)-styryl) pyridinium dibromide; ThermoFisher)), free-swimming larvae were

exposed to FM 1-43FX at 3 μM for 20 seconds, then rinsed 3 times in fresh E3 as previously

described (Toro et al., 2015) and immediately fixed (4% paraformaldehyde, 4% sucrose, 150

μM $CaCl_2$, 0.1 M phosphate buffer). FM 1-43FX mean signal intensity from maximum projection

images was calculated using ImageJ as the integrated pixel intensity divided by the area of the

neuromast region of interest. We verified that relative labeling of hair cells at 1-3 hours

appeared comparable between live FM1-43 and FM 1-43FX. We also verified loss of FM 1-

43FX uptake in larvae following brief treatment with 5 mM BAPTA to disrupt tip links (Kindt et

al., 2012). Following fixation, stereocilia were labeled by incubation with phalloidin conjugated to

Alexa Fluor 488 (Invitrogen) at 66 µM in PBS, washed, and mounted on slides with elvanol.

EdU labeling and quantification

To label proliferating cells, we used the Click-iT EdU Cell Proliferation Kit for Imaging,

Alexa Fluor 555 dye (Invitrogen). Following mechanical overstimulation, larvae were incubated

in 500 µM EdU with 0.5% DMSO in E3 for 4 hours at 28°C then fixed in 4% paraformaldehyde

in PBS overnight at 4°C. Larvae were washed in 3% bovine serum albumin in PBS then

permeabilized with 0.5% Triton-X in PBS. GFP signal in Tg[tnks1bp1:GFP] fish was amplified

using anti-GFP primary antibody (Aves), followed by a secondary antibody conjugated to Alexa

Fluor 488. The EdU detection reaction was performed according to manufacturer guidelines;

larvae were incubated in a reaction cocktail (4 mM CuSO4 and Alexa Fluor 555 azide in 1X

Click-iT reaction buffer with reaction buffer additive) for 1 hour at 25°C. Larvae were washed,

counterstained with Hoechst 33342, and mounted on slides with elvanol. Confocal images of

neuromasts were acquired using an LSM 700 laser scanning confocal microscope with a 63x

1.4 NA Plan-Apochromat oil-immersion objective (Carl Zeiss). The numbers of EdU+ cells per

neuromast were quantified in ImageJ.

Confocal imaging

Images of fixed samples were acquired using an LSM 700 laser scanning confocal

microscope with a 63x 1.4 NA Plan-Apochromat oil-immersion objective (Carl Zeiss). Confocal

stacks were collected with a z step of 0.3 µm over 7-10 µm with pixel size of 100 nm (x-y image

size 51 x 51 µm). Acquisition parameters were established using the brightest control specimen

such that just a few pixels reached saturation in order to achieve the greatest dynamic range in

our experiments. These parameters including gain, laser power, scan speed, dwell time,

resolution, and zoom, were kept consistent between comparisons.

Confocal image processing and analysis

All analysis was performed on blinded images. Digital images were processed using

ImageJ software (Schneider, Rasband, & Eliceiri, 2012). In order to quantitatively measure sizes

and fluorescent intensities of puncta, raw images containing single immunolabel were

subtracted for background using a 20-pixel rolling ball radius and whole neuromasts were

delineated from Parvalbumin-labeled hair cells using the freehand selection and "synchronize

windows" tools. Puncta were defined as regions of immunolabel with pixel intensity above a

determined threshold: threshold for Ribeye label was calculated using the Isodata algorithm

(Ridler & Calvard, 1978) on maximum-intensity projections, threshold for MAGUK label was

calculated as the product of 7 times the average pixel intensity of the whole neuromast region in

a maximum-intensity projection. Particle volume and intensity were measured using the 3D

object counter (Bolte & Cordelieres, 2006) using a lower threshold and a minimum size of 10

voxels. To normalize for differences in staining intensity across experimental trials, all volumes

were divided by the median control volume in each trial for each individual channel. The number

of particles above lower threshold was quantified using the ImageJ Maximum Finder plugin with

a noise tolerance of 10 on maximum-intensity projections. Intact synapses were manually

counted and defined as adjoining or overlapping maxima of Ribeye and MAGUK labels. The

number of synapses per hair cell was approximated by dividing the number of intact synapses

within an neuromast by the number of hair cells in the neuromast. Innervation of neuromast hair

cells was quantified during blinded analysis by scrolling through confocal z-stacks of each

neuromast (step size 0.3 µm) containing hair cell and afferent labeling and identifying hair cells

that were not directly contacted by an afferent neuron i.e. no discernable space between the

hair cell and the neurite. Hair cells that were identified as no longer innervated showed

measurable neurite retraction; there was generally >0.5 µm distance between a retracted

neurite and hair cell. Stereocilia length measurements were obtained in ImageJ from

interpolated 3D projections of z-stack images (step size 0.3 µm) containing phalloidin labeled

hair bundles. Three independent measurements were obtained from the base to the tips of hair

bundles at center of each neuromast, and the average length was calculated.

Quantitative data on macrophage response to mechanical injury were collected from the

two caudal-most ('terminal') neuromasts. Confocal image stacks were obtained using a Zeiss

LSM700 microscope and visualized using Volocity software. These image stacks were used to

derive three metrics from each neuromast. First, the number of macrophages within 25 µm of a

particular neuromast was determined by inscribing a circle of 25 µm radius, centered on the

neuromast, and counting the number of macrophages that were either fully or partially enclosed

by this circle. Next, the number of macrophages contacting a neuromast was determined by

scrolling through the x-y planes of each image stack (1 µm interval between x-y planes, 15 µm

total depth) and the counting macrophages that were in direct contact with Otoferlin-labeled hair

cells. Finally, the number of macrophages that had internalized Otoferlin-labeled material (hair-

cell debris) were counted and were assumed to reflect the number of phagocytic events. For

each metric, the recorded number reflected the activity of a single macrophage, i.e., a

macrophage that made contacts with multiple hair cells and/or had internalized debris from

several hair cells was still classified as a single 'event.'

Subsequent image processing for display within figures was performed using Photoshop

and Illustrator software (Adobe).

Scanning electron microscopy

To image hair-cell bundles, zebrafish larvae were exposed to strong water wave

stimulus, then anesthetized in 0.12% tricaine in E3 and immediately fixed in 2.5%

glutaraldehyde in 0.1 M sodium cacodylate buffer (Electron Microscopy Sciences)

supplemented with 2 mM $CaCl_2$. Larvae were shipped overnight in fixative, then most of the

fixative (~90-95%) was removed, replaced with distilled water, and samples were stored at 4C.

Next, larvae were washed in distilled water (Gibco), dehydrated with an ascending series of

ethanol, critical point dried from liquid CO_2 (Tousimis Aurosamdri 815), mounted on adhesive

carbon tabs (TedPella), sputter coated with 5 nm of platinum (Leica EM ACE600), and imaged

on Hitachi S-4700 scanning electron microscope. Kinocilia diameter measurements were

performed using ImageJ.

Statistical analysis

A hierarchical linear model analysis with fish used as random effects was used to

compare each of the measures between the conditions and groups. Akaike Information Criterion

(AIC) and Bayessian Information Criterion (BIC) were used to identify the best fitted covariance

structure. Tukey's adjustment was used for the alpha level to avoid type I error inflation due to

multiple comparisons. Estimated marginal mean differences and 95% Confidence Intervals

around them were explored and reported for quantification of effect size for group differences.

Graphs for data visualization and additional statistical analyses were performed Prism 8

(Graphpad Software Inc). Mixed model analysis was used to compare time-series data.

Statistical significance between synaptic ribbon and PSD volumes with was determined by

Kruskal-Wallis test (one independent variable) or Mann–Whitney U test (one independent

variable) and appropriate post-hoc tests. Based on the variance and effect sizes reported in

previous studies, the number of biological replicates were suitable to provide statistical power to

avoid Type II error (Sebe et al., 2017; Uribe et al., 2018).

References

Behra, M., Gallardo, V. E., Bradsher, J., Torrado, A., Elkahloun, A., Idol, J., . . . Burgess, S. M.
(2012). Transcriptional signature of accessory cells in the lateral line, using the
Tnk1bp1:EGFP transgenic zebrafish line. *BMC Dev Biol, 12*, 6. doi:10.1186/1471-213X-
12-6
Bhandiwad, A. A., Zeddies, D. G., Raible, D. W., Rubel, E. W., & Sisneros, J. A. (2013).
Auditory sensitivity of larval zebrafish (Danio rerio) measured using a behavioral
prepulse inhibition assay. *The Journal of experimental biology, 216*(Pt 18), 3504-3513.
doi:10.1242/jeb.087635
Bolte, S., & Cordelieres, F. P. (2006). A guided tour into subcellular colocalization analysis in
light microscopy. *J Microsc, 224*(Pt 3), 213-232. doi:10.1111/j.1365-2818.2006.01706.x
Buck, L. M., Winter, M. J., Redfern, W. S., & Whitfield, T. T. (2012). Ototoxin-induced cellular
damage in neuromasts disrupts lateral line function in larval zebrafish. *Hear Res, 284*(1-
2), 67-81. doi:10.1016/j.heares.2011.12.001
Bullen, A., Anderson, L., Bakay, W., & Forge, A. (2019). Localized disorganization of the
cochlear inner hair cell synaptic region after noise exposure. *Biol Open, 8*(1).
doi:10.1242/bio.038547
Burgess, H. A., & Granato, M. (2007). Sensorimotor gating in larval zebrafish. *The Journal of
neuroscience : the official journal of the Society for Neuroscience, 27*(18), 4984-4994.
doi:10.1523/JNEUROSCI.0615-07.2007
Carpaneto Freixas, A. E., Moglie, M. J., Castagnola, T., Salatino, L., Domene, S., Marcovich, I.,
. . . Elgoyhen, A. B. (2021). Unraveling the Molecular Players at the Cholinergic Efferent
Synapse of the Zebrafish Lateral Line. *J Neurosci, 41*(1), 47-60.
doi:10.1523/JNEUROSCI.1772-20.2020
Carrillo, S. A., Anguita-Salinas, C., Pena, O. A., Morales, R. A., Munoz-Sanchez, S., Munoz-
Montecinos, C., . . . Allende, M. L. (2016). Macrophage Recruitment Contributes to
Regeneration of Mechanosensory Hair Cells in the Zebrafish Lateral Line. *J Cell
Biochem, 117*(8), 1880-1889. doi:10.1002/jcb.25487
Chen, Z., Kujawa, S. G., & Sewell, W. F. (2010). Functional roles of high-affinity glutamate
transporters in cochlear afferent synaptic transmission in the mouse. *J Neurophysiol,
103*(5), 2581-2586. doi:10.1152/jn.00018.2010
Cho, S. I., Gao, S. S., Xia, A., Wang, R., Salles, F. T., Raphael, P. D., . . . Oghalai, J. S. (2013).
Mechanisms of hearing loss after blast injury to the ear. *PLoS One, 8*(7), e67618.
doi:10.1371/journal.pone.0067618
Coffin, A. B., Kelley, M. W., Manley, G. A., & Popper, A. N. (2004). Evolution of sensory hair
cells. In G. A. Manley, R. R. Fay, & A. N. Popper (Eds.), *Evolution of the Auditory
System*. New York: Springer-Verlag.
Cruz, I. A., Kappedal, R., Mackenzie, S. M., Hailey, D. W., Hoffman, T. L., Schilling, T. F., &
Raible, D. W. (2015). Robust regeneration of adult zebrafish lateral line hair cells reflects
continued precursor pool maintenance. *Dev Biol, 402*(2), 229-238.
doi:10.1016/j.ydbio.2015.03.019
Davies, C., Tingley, D., Kachar, B., Wenthold, R. J., & Petralia, R. S. (2001). Distribution of
members of the PSD-95 family of MAGUK proteins at the synaptic region of inner and
outer hair cells of the guinea pig cochlea. *Synapse, 40*(4), 258-268.
doi:10.1002/syn.1048
Dow, E., Jacobo, A., Hossain, S., Siletti, K., & Hudspeth, A. J. (2018). Connectomics of the
zebrafish's lateral-line neuromast reveals wiring and miswiring in a simple microcircuit.
Elife, 7. doi:10.7554/eLife.33988

Erickson, T., Pacentine, I. V., Venuto, A., Clemens, R., & Nicolson, T. (2019). The lhfpl5
Ohnologs lhfpl5a and lhfpl5b Are Required for Mechanotransduction in Distinct
Populations of Sensory Hair Cells in Zebrafish. *Front Mol Neurosci, 12*, 320.
doi:10.3389/fnmol.2019.00320
Faucherre, A., Pujol-Marti, J., Kawakami, K., & Lopez-Schier, H. (2009). Afferent neurons of the
zebrafish lateral line are strict selectors of hair-cell orientation. *PloS one, 4*(2), e4477.
doi:10.1371/journal.pone.0004477
Fernandez, K. A., Guo, D., Micucci, S., De Gruttola, V., Liberman, M. C., & Kujawa, S. G.
(2020). Noise-induced Cochlear Synaptopathy with and Without Sensory Cell Loss.
Neuroscience, 427, 43-57. doi:10.1016/j.neuroscience.2019.11.051
Gao, W. Y., Ding, D. L., Zheng, X. Y., Ruan, F. M., & Liu, Y. J. (1992). A comparison of changes
in the stereocilia between temporary and permanent hearing losses in acoustic trauma.
Hearing research, 62(1), 27-41. Retrieved from
http://www.ncbi.nlm.nih.gov/pubmed/1429249
Glowatzki, E., & Fuchs, P. A. (2002). Transmitter release at the hair cell ribbon synapse. *Nature
neuroscience, 5*(2), 147-154. doi:10.1038/nn796
Henry, W. R., & Mulroy, M. J. (1995). Afferent synaptic changes in auditory hair cells during
noise-induced temporary threshold shift. *Hearing research, 84*(1-2), 81-90. Retrieved
from http://www.ncbi.nlm.nih.gov/pubmed/7642458
Hickman, T. T., Smalt, C., Bobrow, J., Quatieri, T., & Liberman, M. C. (2018). Blast-induced
cochlear synaptopathy in chinchillas. *Sci Rep, 8*(1), 10740. doi:10.1038/s41598-018-
28924-7
Hirose, K., Discolo, C. M., Keasler, J. R., & Ransohoff, R. (2005). Mononuclear phagocytes
migrate into the murine cochlea after acoustic trauma. *The Journal of comparative
neurology, 489*(2), 180-194. doi:10.1002/cne.20619
Hirose, K., Rutherford, M. A., & Warchol, M. E. (2017). Two cell populations participate in
clearance of damaged hair cells from the sensory epithelia of the inner ear. *Hear Res,
352*, 70-81. doi:10.1016/j.heares.2017.04.006
Holmgren, M., & Sheets, L. (2021). Influence of Mpv17 on Hair-Cell Mitochondrial Homeostasis,
Synapse Integrity, and Vulnerability to Damage in the Zebrafish Lateral Line. *Front Cell
Neurosci, 15*, 693375. doi:10.3389/fncel.2021.693375
Hu, N., Rutherford, M. A., & Green, S. H. (2020). Protection of cochlear synapses from noise-
induced excitotoxic trauma by blockade of Ca(2+)-permeable AMPA receptors. *Proc Natl
Acad Sci U S A, 117*(7), 3828-3838. doi:10.1073/pnas.1914247117
Husbands, J. M., Steinberg, S. A., Kurian, R., & Saunders, J. C. (1999). Tip-link integrity on
chick tall hair cell stereocilia following intense sound exposure. *Hear Res, 135*(1-2), 135-
145. doi:10.1016/s0378-5955(99)00101-x
Indzhykulian, A. A., Stepanyan, R., Nelina, A., Spinelli, K. J., Ahmed, Z. M., Belyantseva, I. A., .
. . Frolenkov, G. I. (2013). Molecular remodeling of tip links underlies mechanosensory
regeneration in auditory hair cells. *PLoS Biol, 11*(6), e1001583.
doi:10.1371/journal.pbio.1001583
Jia, S., Yang, S., Guo, W., & He, D. Z. (2009). Fate of mammalian cochlear hair cells and
stereocilia after loss of the stereocilia. *J Neurosci, 29*(48), 15277-15285.
doi:10.1523/JNEUROSCI.3231-09.2009
Kaur, T., Clayman, A. C., Nash, A. J., Schrader, A. D., Warchol, M. E., & Ohlemiller, K. K.
(2019). Lack of Fractalkine Receptor on Macrophages Impairs Spontaneous Recovery of
Ribbon Synapses After Moderate Noise Trauma in C57BL/6 Mice. *Front Neurosci, 13*,
620. doi:10.3389/fnins.2019.00620
Kaur, T., Zamani, D., Tong, L., Rubel, E. W., Ohlemiller, K. K., Hirose, K., & Warchol, M. E.
(2015). Fractalkine Signaling Regulates Macrophage Recruitment into the Cochlea and

880   Promotes the Survival of Spiral Ganglion Neurons after Selective Hair Cell Lesion. *J*
881   *Neurosci, 35*(45), 15050-15061. doi:10.1523/JNEUROSCI.2325-15.2015

882   Kim, K. X., Payne, S., Yang-Hood, A., Li, S. Z., Davis, B., Carlquist, J., . . . Rutherford, M. A.
883   (2019). Vesicular Glutamatergic Transmission in Noise-Induced Loss and Repair of
884   Cochlear Ribbon Synapses. *The Journal of neuroscience : the official journal of the*
885   *Society for Neuroscience, 39*(23), 4434-4447. doi:10.1523/JNEUROSCI.2228-18.2019

886   Kindt, K. S., Finch, G., & Nicolson, T. (2012). Kinocilia mediate mechanosensitivity in
887   developing zebrafish hair cells. *Developmental cell, 23*(2), 329-341.
888   doi:10.1016/j.devcel.2012.05.022

889   Kindt, K. S., & Sheets, L. (2018). Transmission Disrupted: Modeling Auditory Synaptopathy in
890   Zebrafish. *Front Cell Dev Biol, 6*, 114. doi:10.3389/fcell.2018.00114

891   Kniss, J. S., Jiang, L., & Piotrowski, T. (2016). Insights into sensory hair cell regeneration from
892   the zebrafish lateral line. *Curr Opin Genet Dev, 40*, 32-40.
893   doi:10.1016/j.gde.2016.05.012

894   Kujawa, S. G., & Liberman, M. C. (2009). Adding insult to injury: cochlear nerve degeneration
895   after "temporary" noise-induced hearing loss. *The Journal of neuroscience : the official*
896   *journal of the Society for Neuroscience, 29*(45), 14077-14085.
897   doi:10.1523/JNEUROSCI.2845-09.2009

898   LeMasurier, M., & Gillespie, P. G. (2005). Hair-cell mechanotransduction and cochlear
899   amplification. *Neuron, 48*(3), 403-415. doi:10.1016/j.neuron.2005.10.017

900   Levi, R., Akanyeti, O., Ballo, A., & Liao, J. C. (2015). Frequency response properties of primary
901   afferent neurons in the posterior lateral line system of larval zebrafish. *J Neurophysiol,*
902   *113*(2), 657-668. doi:10.1152/jn.00414.2014

903   McHenry, M. J., Feitl, K. E., Strother, J. A., & Van Trump, W. J. (2009). Larval zebrafish rapidly
904   sense the water flow of a predator's strike. *Biol Lett, 5*(4), 477-479.
905   doi:10.1098/rsbl.2009.0048

906   Merchan-Perez, A., & Liberman, M. C. (1996). Ultrastructural differences among afferent
907   synapses on cochlear hair cells: correlations with spontaneous discharge rate. *J Comp*
908   *Neurol, 371*(2), 208-221. doi:10.1002/(SICI)1096-9861(19960722)371:2<208::AID-
909   CNE2>3.0.CO;2-6

910   Nair, A., Azatian, G., & McHenry, M. J. (2015). The kinematics of directional control in the fast
911   start of zebrafish larvae. *J Exp Biol, 218*(Pt 24), 3996-4004. doi:10.1242/jeb.126292

912   Nordmann, A. S., Bohne, B. A., & Harding, G. W. (2000). Histopathological differences between
913   temporary and permanent threshold shift. *Hearing research, 139*(1-2), 13-30. Retrieved
914   from http://www.ncbi.nlm.nih.gov/pubmed/10601709

915   Nüsslein-Volhard, C., & Dahm, R. (2002). *Zebrafish : a practical approach* (1st ed.). Oxford:
916   Oxford University Press.

917   Obholzer, N., Wolfson, S., Trapani, J. G., Mo, W., Nechiporuk, A., Busch-Nentwich, E., . . .
918   Nicolson, T. (2008). Vesicular glutamate transporter 3 is required for synaptic
919   transmission in zebrafish hair cells. *The Journal of neuroscience : the official journal of*
920   *the Society for Neuroscience, 28*(9), 2110-2118. doi:10.1523/JNEUROSCI.5230-
921   07.2008

922   Olivari, F. A., Hernandez, P. P., & Allende, M. L. (2008). Acute copper exposure induces
923   oxidative stress and cell death in lateral line hair cells of zebrafish larvae. *Brain Res,*
924   *1244*, 1-12. doi:10.1016/j.brainres.2008.09.050

925   Puel, J. L., Ruel, J., Gervais d'Aldin, C., & Pujol, R. (1998). Excitotoxicity and repair of cochlear
926   synapses after noise-trauma induced hearing loss. *Neuroreport, 9*(9), 2109-2114.
927   Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/9674603

928   Qiu, X., & Muller, U. (2018). Mechanically Gated Ion Channels in Mammalian Hair Cells. *Front*
929   *Cell Neurosci, 12*, 100. doi:10.3389/fncel.2018.00100

Ridler, T. W., & Calvard, S. (1978). Picture Thresholding Using an Iterative Selection Method.
3EEE TRANSACTIONS ON SYSTEMS, MAN, AND CYBERNETICS(8), 630 - 632.
doi:https://doi.org/10.1109/TSMC.1978.4310039
Roca, F. J., & Ramakrishnan, L. (2013). TNF dually mediates resistance and susceptibility to
mycobacteria via mitochondrial reactive oxygen species. *Cell, 153*(3), 521-534.
doi:10.1016/j.cell.2013.03.022
Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of
image analysis. *Nat Methods, 9*(7), 671-675. doi:10.1038/nmeth.2089
Sebe, J. Y., Cho, S., Sheets, L., Rutherford, M. A., von Gersdorff, H., & Raible, D. W. (2017).
Ca(2+)-Permeable AMPARs Mediate Glutamatergic Transmission and Excitotoxic
Damage at the Hair Cell Ribbon Synapse. *J Neurosci, 37*(25), 6162-6175.
doi:10.1523/JNEUROSCI.3644-16.2017
Sheets, L., Kindt, K. S., & Nicolson, T. (2012). Presynaptic CaV1.3 channels regulate synaptic
ribbon size and are required for synaptic maintenance in sensory hair cells. *The Journal
of neuroscience : the official journal of the Society for Neuroscience, 32*(48), 17273-
17286. doi:10.1523/JNEUROSCI.3005-12.2012
Sheets, L., Trapani, J. G., Mo, W., Obholzer, N., & Nicolson, T. (2011). Ribeye is required for
presynaptic Ca(V)1.3a channel localization and afferent innervation of sensory hair cells.
Development, 138(7), 1309-1319. doi:10.1242/dev.059451
Shi, L., Liu, L., He, T., Guo, X., Yu, Z., Yin, S., & Wang, J. (2013). Ribbon synapse plasticity in
the cochleae of Guinea pigs after noise-induced silent damage. *PloS one, 8*(12),
e81566. doi:10.1371/journal.pone.0081566
Slepecky, N. (1986). Overview of mechanical damage to the inner ear: noise as a tool to probe
cochlear function. *Hear Res, 22*, 307-321. doi:10.1016/0378-5955(86)90107-3
Song, Q., Shen, P., Li, X., Shi, L., Liu, L., Wang, J., . . . Yin, S. (2016). Coding deficits in hidden
hearing loss induced by noise: the nature and impacts. *Scientific reports, 6*, 25200.
doi:10.1038/srep25200
Thomas, A. J., Wu, P., Raible, D. W., Rubel, E. W., Simon, J. A., & Ou, H. C. (2015).
Identification of small molecule inhibitors of cisplatin-induced hair cell death: results of a
10,000 compound screen in the zebrafish lateral line. *Otology & neurotology : official
publication of the American Otological Society, American Neurotology Society [and]
European Academy of Otology and Neurotology, 36*(3), 519-525.
doi:10.1097/MAO.0000000000000487
Toro, C., Trapani, J. G., Pacentine, I., Maeda, R., Sheets, L., Mo, W., & Nicolson, T. (2015).
Dopamine Modulates the Activity of Sensory Hair Cells. *The Journal of neuroscience :
the official journal of the Society for Neuroscience, 35*(50), 16494-16503.
doi:10.1523/JNEUROSCI.1691-15.2015
Trapani, J. G., & Nicolson, T. (2010). Physiological recordings from zebrafish lateral-line hair
cells and afferent neurons. *Methods in cell biology, 100*, 219-231. doi:10.1016/B978-0-
12-384892-5.00008-6
Uribe, P. M., Villapando, B. K., Lawton, K. J., Fang, Z., Gritsenko, D., Bhandiwad, A., . . . Coffin,
A. B. (2018). Larval Zebrafish Lateral Line as a Model for Acoustic Trauma. *eNeuro,
5*(4). doi:10.1523/ENEURO.0206-18.2018
Wagner, E. L., & Shin, J. B. (2019). Mechanisms of Hair Cell Damage and Repair. *Trends
Neurosci, 42*(6), 414-424. doi:10.1016/j.tins.2019.03.006
Wan, G., Gomez-Casati, M. E., Gigliello, A. R., Liberman, M. C., & Corfas, G. (2014).
Neurotrophin-3 regulates ribbon synapse density in the cochlea and induces synapse
regeneration after acoustic trauma. *eLife, 3*. doi:10.7554/eLife.03564
Wang, H., Yin, S., Yu, Z., Huang, Y., & Wang, J. (2011). Dynamic changes in hair cell
stereocilia and cochlear transduction after noise exposure. *Biochem Biophys Res
Commun, 409*(4), 616-621. doi:10.1016/j.bbrc.2011.05.049

Wang, X., Zhu, Y., Long, H., Pan, S., Xiong, H., Fang, Q., . . . Sha, S. H. (2018). Mitochondrial
Calcium Transporters Mediate Sensitivity to Noise-Induced Losses of Hair Cells and
Cochlear Synapses. *Front Mol Neurosci, 11*, 469. doi:10.3389/fnmol.2018.00469
Warchol, M. E. (2011). Sensory regeneration in the vertebrate inner ear: differences at the
levels of cells and species. *Hear Res, 273*(1-2), 72-79. doi:10.1016/j.heares.2010.05.004
Warchol, M. E. (2019). Interactions between Macrophages and the Sensory Cells of the Inner
Ear. *Cold Spring Harb Perspect Med, 9*(6). doi:10.1101/cshperspect.a033555
Warchol, M. E., Schrader, A., & Sheets, L. (2021). Macrophages Respond Rapidly to Ototoxic
Injury of Lateral Line Hair Cells but Are Not Required for Hair Cell Regeneration. *Front*
Cell Neurosci, 14, 613246. doi:10.3389/fncel.2020.613246
Weeg, M. S., Fay, R. R., & Bass, A. H. (2002). Directionality and frequency tuning of primary
saccular afferents of a vocal fish, the plainfin midshipman (Porichthys notatus). *J Comp*
Physiol A Neuroethol Sens Neural Behav Physiol, 188(8), 631-641. doi:10.1007/s00359-
002-0338-2
Williams, J. A., & Holder, N. (2000). Cell turnover in neuromasts of zebrafish larvae. *Hear Res,*
143(1-2), 171-181. doi:10.1016/s0378-5955(00)00039-3
Wong, H. C., Zhang, Q., Beirl, A. J., Petralia, R. S., Wang, Y. X., & Kindt, K. (2019). Synaptic
mitochondria regulate hair-cell synapse size and function. *Elife, 8*.
doi:10.7554/eLife.48914
Wynn, T. A., & Vannella, K. M. (2016). Macrophages in Tissue Repair, Regeneration, and
Fibrosis. *Immunity, 44*(3), 450-462. doi:10.1016/j.immuni.2016.02.015
Xiao, Y., Faucherre, A., Pola-Morell, L., Heddleston, J. M., Liu, T. L., Chew, T. L., . . . Lopez-
Schier, H. (2015). High-resolution live imaging reveals axon-glia interactions during
peripheral nerve injury and repair in zebrafish. *Dis Model Mech, 8*(6), 553-564.
doi:10.1242/dmm.018184
Zhang, Q., Li, S., Wong, H. C., He, X. J., Beirl, A., Petralia, R. S., . . . Kindt, K. S. (2018).
Synaptically silent sensory hair cells in zebrafish are recruited after damage. *Nat*
Commun, 9(1), 1388. doi:10.1038/s41467-018-03806-8
1009

1010

Supplemental Figure 1: Fish exposed to periodic stimulus have less mechanical damage to neuromasts, but still show synapse loss. (A) Schematic of the two exposure protocols. Sustained exposure was a 20 min. pulse followed by 120 minutes uninterrupted mechanical overstimulation; periodic exposure was 90 min. exposure with intermittent 10 min. breaks totaling 120 minutes. **(B)** Periodic stimulus causes less neuromast disruption. Immediately following sustained exposure, 46% of exposed neuromasts showed a "disrupted" phenotype, whereas following a periodic exposure only 17% of the neuromasts appeared "disrupted" (Unpaired t-test **P=0.0034). **(C)** Position of the neuromast along the tail was also associated with vulnerability to disruption with both sustained and periodic stimulation. **(D-F')** Measurements of hair-cell number, innervation, and synapse number averaged from individual fish (neuromasts L3-L5). **(D-D')** Average hair cell number per neuromast following exposure. With periodic exposure, the number of hair cells was comparable to control (*P=0.108 sustained, P=0.7233 periodic). **(E-E')** % of neuromast hair cells innervated (****P<0.0001 sustained, P=0.0156 periodic). **(F-F')** Average number of intact synapses per neuromast. There is significant loss of synapses with both periodic and sustained exposures (**P=0.0045 sustained, *P=0.0236 periodic).

Supplemental Figure 2: Hair cell organs of the ear appeared undamaged in larvae exposed to sustained stimulus. Representative maximum intensity images of hair cell organs in the ears of control **(A-C)** and larvae exposed to sustained strong water wave stimulus **(A'-C')**. Hair cells in A-B were immunolabeled with an antibody against Otoferlin; posterior macula in C were immunolabled with antibodies against Parvalbumin to label hair cells and CtBP to label synaptic ribbons. Scale bars: 10 μm

all scale bars are 1μm

Supplemental Figure 3: Scanning electron microscopy imaging of tail neuromasts following mechanical injury confirms the damage is more prominent for posterior neuromasts. **(A-C)** Representative images of tail neuromasts of control fish larvae, presented as they are positioned on the larva: L1, L2 and terminal neuromasts. Each hair cell carries a tubulin-based primary cilium (kinocilium), which is thicker than the multiple mechanosensitive stereocilia arranged in a staircase. **(D-G)** Examples of tail neuromasts immediately following sustained stimulus exposure, presented as they are positioned on the larva: L2, L4, and two terminal neuromasts highlighting different levels of damage, with much more pronounced damage evident on terminal neuromasts. Scale bars: 1 μm

Supplemental Figure 4: Confocal images show damaged kinocilia following mechanical injury. Representative confocal images of hair cell kinocilia (acetylated tubulin immunolabel; magenta) and stereocilia (βactin-GFP; green) in control (A-A'') and exposed (B-B'') larval neuromast L4-6 immediately following sustained strong water wave stimulus. Kinocilia appear as intact bundles in the control neuromasts. Arrow indicated kinocilial swelling, arrowheads indicate split kinocilia in the exposed neuromasts. Scale bar: 5 μm

Supplemental Movies

Swimming behavior of 7-day-old larvae during exposure to the strong water wave stimulus shown in Figure 2 over 500ms (1000 fps/ 500 frames).

Movie S1: Swimming behavior of control fish with intact lateral line organs. Magenta circle indicates a fish prior to a"fast escape" response.

Movie S2: Swimming behavior of larvae whose lateral-line neuromasts were ablated with $CuSO_4$. Magenta circles indicate larvae that were swept into the waves and could no longer tracked.