1   Nef enhances HIV-1 replication and infectivity independently of

2   SERINC3 and SERINC5 in CEM T cells

4 Peter W. Ramirez[a,b,c*,#], Aaron A. Angerstein[a,b], Marissa Suarez[b], Thomas Vollbrecht[a,b],

5 Jared Wallace[d], Ryan M. O' Connell[d] and John Guatelli[a,b]

7 [a]Department of Medicine, University of California San Diego, La Jolla, California, USA.

8 [b]VA San Diego Healthcare System, San Diego, California, USA.

9
10 [c]Department of Biological Sciences, California State University Long Beach, Long Beach, CA.
11
12 [d]Division of Microbiology and Immunology, Department of Pathology, The University of
13 Utah, Salt Lake City, UT.
14

15 Running Head: Attenuation of HIV-1Δnef without *serinc3* and *serinc5*

17 [*]Present Address

18 [#]Address Correspondence to Peter W. Ramirez: Peter.Ramirez@csulb.edu

Abstract

The lentiviral *nef* gene encodes several discrete activities aimed at co-opting or antagonizing cellular proteins and pathways to defeat host defenses and maintain persistent infection. Primary functions of Nef include downregulation of CD4 and MHC class-I from the cell surface, disruption or mimicry of T-cell receptor signaling, and enhancement of viral infectivity by counteraction of the host antiretroviral proteins SERINC3 and SERINC5. In the absence of Nef, SERINC3 and SERINC5 incorporate into virions and inhibit viral fusion with target cells, decreasing infectivity. However, whether Nef's counteraction of SERINC3 and SERINC5 is the cause of its positive influence on viral growth-rate in CD4-positive T cells is unclear. Here, we utilized CRISPR/Cas9 to knockout SERINC3 and SERINC5 in a leukemic CD4-positive T cell line (CEM) that displays robust *nef*-related infectivity and growth-rate phenotypes. As previously reported, viral replication was severely attenuated in CEM cells infected with HIV-1 lacking Nef (HIV-1ΔNef). This attenuated growth-rate phenotype was observed regardless of whether or not the coding regions of the *serinc3* and *serinc5* genes were intact. Moreover, knockout of *serinc3* and *serinc5* failed to restore the infectivity of HIV-1ΔNef virions produced from infected CEM cells in single-cycle replication experiments using CD4-positive HeLa cells as targets. Taken together, our results corroborate a recent study using another T-lymphoid cell line (MOLT-3) and suggest that Nef modulates a still unidentified host protein(s) to enhance viral growth rate and infectivity in CD4-positive T cells.

Importance

HIV-1 Nef is a major pathogenicity factor *in vivo*. A well-described activity of Nef is the enhancement of virion-infectivity and viral propagation *in vitro.* The infectivity-effect has been attributed to Nef's ability to prevent the cellular, antiretroviral proteins SERINC3 and SERINC5 from incorporating into viral particles. While the activity of the SERINCs as inhibitors of retroviral infectivity has been well-documented, the role these proteins play in controlling HIV-1 replication is less clear. We report here that genetic disruption of SERINC3 and SERINC5 rescues neither viral replication-rate nor the infectivity of cell-free virions produced from CD4-positive T cells of the CEM lymphoblastoid line infected with viruses lacking Nef. This indicates that failure to modulate SERINC3 and SERINC5 is not the cause of the virologic attenuation of *nef*-negative HIV-1 observed using this system.

Introduction

Primate lentiviruses encode several accessory gene products that facilitate viral reproduction and persistence by providing evasion of the host immune response. The lentiviral Nef protein accelerates viral pathogenesis and progression to AIDS in SIV-infected rhesus macaques (1) and enhances disease progression in humans infected with HIV-1 (2, 3). Nef is a small, myristoylated, peripheral membrane protein with many well-conserved activities, including the downregulation of cell surface proteins such as CD4 and MHC class-I (MHC-I), the modulation of T cell activation, and the enhancement of viral infectivity and growth-rate (4).

To downregulate CD4, Nef binds the cytoplasmic domain of CD4 and links it to the clathrin Adaptor Protein 2 (AP-2) complex, internalizing CD4 from the plasma membrane and delivering it to lysosomes for degradation (5–8). The targeting of CD4 by Nef contributes to viral replication in several ways. CD4 downregulation prevents superinfection of cells and consequent premature cell-death, ensuring adequate time for viral replication (9). It also contributes to inhibiting antibody dependent cellular cytotoxicity (ADCC) that recognizes CD4-induced epitopes with Env (10). Potentially directly relevant to the work presented here, CD4 incorporates into virions and inhibits virion-infectivity if not downregulated by Nef (11, 12).

To downregulate MHC-I, Nef has been proposed to use two non-mutually exclusive mechanisms: 1) Nef and the clathrin-adaptor AP-1 intercept *de novo* synthesized MHC-I

molecules within the *trans*-Golgi network (TGN), leading to lysosomal degradation (13–

16); and/or 2) Nef retains internalized MHC-I molecules in the TGN via induction of a Src-

family kinase/phosphoinositide-3-kinase signaling cascade (17–19). In either case,

reducing the expression of MHC-I molecules at the plasma membrane protects HIV-1

infected cells from lysis by cytotoxic T-lymphocytes, contributing to immune evasion (20).

Nef interacts directly with Src-family kinases including the lymphocyte specific kinase Lck,

which it downregulates from the cell surface (21). The many studies of Nef's influence on

T cell activation are difficult to reconcile, but transcriptional profiling suggests that the

expression of Nef mimics signaling through the T cell receptor (22). With the exception of

the model of MHC-I downregulation noted above, a clear mechanistic link between the

influence of HIV-1 Nef on T cell activation and its influence on the expression of cell

surface receptors is lacking.

Nef stimulates HIV-1 replication and enhances virion-infectivity in many cell culture

systems, including various T cell lines, primary CD4+ T cells, and human lymphoid tissue

(23–26). Nef alleles from various primate lentiviruses and isolated at different stages of

HIV-1 infection in humans maintain the ability to enhance HIV-1 replication and infectivity,

suggesting that these functions are important for establishing and maintaining persistent

infection (26–28). Expression of Nef within virion-producer cells and encoded either in *cis*

or in *trans* relative to the viral genome enhances HIV-1 replication and yields virions of

greater infectivity (29, 30). The ability of Nef to enhance infectivity requires cellular

components involved in vesicular trafficking (dynamin 2, AP-2, and clathrin) and is also

determined by the Envelope (Env) glycoprotein (31, 32). A survey of cell lines identified

murine leukemia virus glycosylated Gag (MLV glycoGag), a protein structurally unrelated

to Nef, as an infectivity factor that rescues Nef-deficient HIV-1 virions (33). These and

other features suggested that Nef counteracts a cellular factor that restricts viral infectivity

and possibly replication.

Two groups identified the host transmembrane protein SERINC5, and to a lesser degree

SERINC3, as an inhibitor of HIV-1 virion-infectivity that is counteracted by Nef (34, 35).

Nef downregulates SERINC5 from the plasma membrane via a clathrin/AP-2 and Rab5/7

endo-lysosomal pathway (34, 36), reducing the incorporation of SERINC5 in HIV-1

virions. This in turn correlates with more efficient fusion of virions with target cells and

greater infectivity (34, 35). Nef's ability to counteract SERINC5 is conserved across

primate lentiviruses and correlates with the prevalence of these viruses in the wild (34,

37). Modulation of SERINC5 extends to other retroviral proteins, including S2 from equine

infectious anemia virus (EIAV) as well as MLV glycoGag (34, 35, 38). HIV-1 Env

glycoproteins are differentially sensitive to SERINC-mediated restriction when produced

from CD4-negative cells in single-round replication assays; sensitivity correlates to some

extent with the degree of Env-trimer openness and instability (34, 35, 39, 40).

SERINCs comprise a family of five genes that are evolutionarily conserved from yeast to

mammals (41). They encode multi-pass transmembrane proteins that support serine-

specific phospholipid biosynthesis (hence their name: serine incorporator) (42), yet this

function does not seem to account for their anti-retroviral activity (43). Rather, SERINC5

appears to disrupt the formation of fusion pores between HIV-1 virions and target cells

(44) in an Env-conformation and CD4-dependent manner (45).

Initial studies of the Nef-SERINC relationship focused on the Jurkat T cell line, due to the

large defect in the infectivity of Nef-negative virions produced from these cells and their

relatively high levels of SERINC5 mRNA. Studies using another CD4-positive T cell line,

MOLT-3, have recently cast doubt on whether SERINC family proteins are sufficient to

explain the virologic phenotypes of Nef (46). In support of a SERINC-dependent

mechanism, Nef does not enhance HIV-1 infectivity and replication-rate in Jurkat T cells

when SERINC3 and SERINC5 are knocked out (34, 35). Moreover, a minimal MLV

glycoGag (termed glycoMA) can functionally replace Nef with respect to viral replication-

rate and virion-infectivity when the virus is propagated using Jurkat cells (46). In contrast,

Nef, but not glycoMA, enhances HIV-1 replication in MOLT-3 cells. The Nef-effect in Molt-

3 cells persists when the cells are knocked out for SERINC3 and SERINC5, indicating

that these restriction factors are not necessary for the virologic effects of Nef in this setting

(46). Remarkably, glycoMA cannot substitute functionally for Nef with respect to

stimulating viral replication in primary CD4-positive T cells. This suggests that the growth-

rate enhancing effect of Nef in primary T cells is unrelated to SERINC-antagonism (46),

even though the virion-infectivity enhancing effect of Nef reportedly is (34).

Given these conflicting results, we aimed to further test the hypothesis that Nef enhances

HIV-1 replication in a SERINC-dependent manner. To do this, we returned to the CD4-

positive T cell line in which we originally observed a striking stimulation of growth-rate by

Nef, an effect that was associated with Nef-mediated enhancement of virion-infectivity

(CEM) (23). Here, we show that neither the attenuated replication-rate of *nef*-deficient

HIV-1 in CEM T cells nor the reduced virion-infectivity of *nef*-deficient HIV-1 produced by

these cells is rescued by CRISPR/cas9 editing of *serinc3* and *serinc5*. These results

support those recently documented using MOLT-3 and primary CD4+ T cells and suggest

that how Nef stimulates HIV-1 replication remains unclear (46).

Materials and Methods

Cell Lines and Plasmids: HEK293T (a generous gift from Dr. Ned Landau) and HeLa

TZM-bl cells (Dr. John Kappes and Xiaoyun Wu,: NIH AIDS Reagent Program, Division

of AIDS, NIAID, NIH) (47–51) were grown in DMEM media (Thermo Fisher Scientific)

supplemented with 10% FBS (Hyclone) and 1% Penicillin/Streptomycin (Thermo Fisher

Scientific). HeLa P4.R5 (obtained from Dr. Ned Landau) were cultured in 10% FBS, 1%

Penicillin/Streptomycin and 1 μg puromycin. Both HeLa cell line derivatives express CD4,

CXCR4 and CCR5 and contain either a Tat-inducible β-galactosidase (HeLa P4.R5) or

both the β-galactosidase and luciferase (HeLa TZM.bl) genes under the transcriptional

control of the HIV-1 LTR. CEM cells, a leukemic T cell clone (a generous gift from Dr.

Douglas Richman), were cultured in RPMI 1640 media plus 10% FBS and 1%

Penicillin/Streptomycin (Thermo Fisher Scientific). The proviral plasmids pNL4-3 and

pNL4-3ΔNef have been previously described (23). LentiCRISPRv2 (Addgene; Catalog #:

52961) contains a single guide RNA (sgRNA) targeting Exon 2 of SERINC3 (5'-

ATAAATGAGGCGAGTCACCG-3') and was a gift from Dr. Massimo Pizzato (34). The

lentiviral packaging (pxRSV-Rev, pMDLg/pRRE) and envelope (pMD2.G) plasmids were

kindly provided by Dr. Dan Gibbs. LentiCRISPR-GFP encodes GFP in place of

puromycin. Five sgRNAs targeting either Exon 1 or Exon 2 of SERINC5 were designed

using an online CRISPR tool (benchling.com) and cloned into LentiCRISPR-GFP using

previously described methods (52, 53). The sgRNA sequences were as follows: sgRNA-

SERINC5(1), 5'- ACA GCACTGAGCTGACATCG-3' ; sgRNA-SERINC5(2), 5'-

GCACTGAGCTGACATCGC   GG-3'   ;   sgRNA-SERINC5(3),   5'-

CTTCGTTCAAGTGTGAGCTG'3'   ;   sgRNA-SERINC5(4),   5'-

CATCATGATGTCAACAACCG-3'   ;   sgRNA-SERINC5(5),   5'-

TGAGGGACTGCCGAATCCTG-3'. Briefly, sgRNA oligos were designed to produce the

same overhangs after *BsmBI* digestion (5'- *CACCG*(sgRNA Oligo #1)-3' ; 3'-*C*(sgRNA

Oligo #2)-*CAAA*-3'). The oligos were phosphorylated (T4 Polynucleotide Kinase ; NEB)

and annealed in a thermal cycler according to the following conditions: 37°C for 30

minutes; 95°C for 5 minutes with a ramp down to 25°C at 5°C/minute. Diluted oligos

(1:200) were ligated (T4 ligase; NEB) into dephosphorylated (Fast AP; Fermentas) and

BsmBI digested (Fast *BsmBI*; Fermentas) LentiCRISPR-GFP by overnight incubation at

16°C, followed by transformation into Stbl3 bacteria (Thermo Fisher Scientific). Plasmid

DNA was isolated from overnight bacterial cultures and verified via Sanger sequencing.

Generation of stable cell lines using CRISPR-Cas9: To produce 3[rd] generation lentiviral

stocks, HEK293T cells were transfected with a total of 22.5 μg total plasmid according to

the following equimolar ratios: 10 μg LentiCRISPR transfer plasmid (empty or containing

sgRNAs against either SERINC3 or SERINC5), 5.9 μg pMDLg/pRRE, 2.8 μg pxRSV-Rev

and 3.8 μg pMD2.G. Forty-eight hours later, concentrated lentivirus-containing

supernatant was harvested following low-speed centrifugation, filtration (0.45 μm), and

mixture with Lenti-X concentrator (Takara Bio) according to the manufacturer's

instructions. Briefly, 1 volume Lenti-X concentrator was mixed with 3 volumes clarified

supernatant. The mixture was incubated for 30 minutes at 4°C, centrifuged at 1,500 x g

for 45 minutes at 4°C, resuspended in 1 ml complete DMEM media and immediately

stored at -80°C in single-use aliquots (100 μL).

CEM cells knocked out for SERINC3 (S3-KO) were generated by spinoculation of 1 x 10^6

cells with 100 μL lentivirus (LentiCRISPRv2-SERINC3 ; (34)) at 1,200 x g for 2 hours at

25°C. Puromycin (1 μg/ml) was added to cell cultures 72 hours post transduction to select

for positive clones. Two weeks post-selection, genomic DNA was isolated from mock or

lentiCRISPRv2-SERINC3 transduced CEM cells using the DNeasy Blood and Tissue Kit

(Qiagen) according to the manufacturer's instructions. Genome editing was assessed by

Tracking of Indels by Decomposition (TIDE; tide.deskgen.com). PCR amplicons

encompassing exon 2 of SERINC3 were produced using Taq 2x Master Mix (NEB) and

the following primers: 5'-CAAATTACAACCAACTTGATTAACAACGACG-3' and 5'-

CTATAAAGCCTGATTTGCCTCGCTTTCTCTTC-3'. Twenty single cell clones were

isolated from bulk edited cultures using an array dilution method (IDT), followed by

genomic DNA isolation and PCR amplification as described above. PCR products

indicative of genome editing were sub-cloned into the TOPO TA vector (Thermo Fisher

Scientific) and plasmid DNA isolated from overnight bacterial cultures. Genome editing

was verified via Sanger sequencing of at least 10 independent bacterial clones.

To generate double-knockout cells, CEM S3 KO cells were spinoculated as described

above with either empty lentivirus (LentiCRISPR-GFP) or each of the five gRNAs

targeting SERINC5. Cells were sorted (BD FACS Aria) based on GFP expression

seventy-two hours post-transduction. Isolation of genomic DNA, PCR amplification and

TIDE analysis were carried out similarly to the generation of SERINC3 KO cells. We

clonally expanded, TOPO TA cloned and sequence verified sgRNA-SERINC5(4) since

this showed abundant genome editing and produced frameshift mutations in SERINC5

Exon 2. The following primers were used to generate PCR amplicons for TOPO TA

cloning and TIDE analysis: 5'-AGTGCCTGGCCATGTTTCTT-3' and 5'-

CATAGAGCAGGCTTCAGGAA-3'.

HIV-1 production and titer: To produce replication-competent viruses, HEK293T cells (4

x 10^6 /10 cm plate) were transfected with 24 μg of an infectious molecular clone of HIV-1

(NL4-3) or a mutant lacking the *nef* gene (NL4-3ΔNef) using Lipofectamine 2000 reagent

according to the manufacturer's instructions (Thermo Fisher Scientific). Virus-containing

supernatants were collected forty-eight hours post transfection, clarified by low-speed

centrifugation, passed through 0.45 μm filters and stored at -80°C. Viral titers were

measured by infecting HeLa P4.R5 cells with diluted viral stocks in duplicate in a 48-well

format for 48 hours. The cells were then fixed (1% formaldehyde; .2% gludaraldehyde)

for 5 minutes at room-temperature, followed by overnight staining at 37°C with a solution

composed of 4 mM potassium ferrocyanide, 4mM potassium ferricyanide, 2 mM $MgCl_2$

and 0.4 mg/ml X-gal (5-bromo-4chloro-3indolyl-β-D-galactopyranoside). Infected cells

(expressed as infectious centers (IC)) were quantified using a computer image-based

method (54) The p24 antigen concentrations of each stock were measured by ELISA

(ABL Bioscience).

HIV-1 infection and replication: To conduct HIV-1 replication studies, 1 x 10^6 CEM cells

(wildtype (WT); SERINC3 knockout (S3 K.O.); SERINC3/SERINC5 knockout clone 6

(S3/S5 K.O. (6)); SERINC3/SERINC5 knockout clone 8 (S3/S5 K.O. (8))) were infected

with NL4-3 (hereafter termed Nef+) or NL4-3ΔNef (hereafter termed Nef-) at an multiplicity

of infection (MOI) = 0.001 for 4 hours at 37°C in a 24-well plate format. The cells were

then washed 3 times with 1 ml PBS (Corning), resuspended in 4 ml complete growth

media (RPMI 1640), transferred to T25 labeled flasks and incubated at 37°C in an

"upright" position for the duration of the experiment. Every three days, cultures were split

1:4 (1 ml cells ; 3 ml media) and an aliquot (1 ml viral supernatant) stored at -80°C for

quantification of HIV-1 replication (p24 antigen) by ELISA. To measure cell-associated

HIV-1 infection, mock infected or infected CEM cells were fixed and permeabilized

(Cytofix/Cytoperm; BD Biosciences) for 30 minutes at 4°C, washed (Perm Wash buffer;

BD Biosciences) and incubated with an HIV-1 core antigen(p24)-FITC antibody (clone

KC57; Beckman Coulter) for 30 minutes at 4°C. Cells were washed with Flow Buffer (1x

PBS + 3% FBS) and subjected to flow cytometry on a Novocyte 3000 (ACEA

Biosciences).

Measurement of HIV-1 infectivity: Viral infectivity was quantified from virions produced in

CEM cells infected with NL4-3 or NL4-3ΔNef at either day 9, 12 or 15 post-infection. A

20% sucrose cushion was used to concentrate virions via centrifugation according to the

following parameters: 23,500 x g ; 1 hour at 4°C. Viral pellets were resuspended in culture

medium and appropriate dilutions used to immediately infect 1.25 x 10^4 HeLa TZM.bl cells

in triplicate in a 96-well format. Forty-eight hours later, the culture medium was removed

and the cells were lysed using a luciferase reporter gene assay reagent (Britelite, Perkin

Elmer). Infectivity (luciferase activity) was measured using a luminometer with data

expressed as relative light units (RLU). These values were normalized to the p24

concentration within each sample and depicted as RLU/p24.

Data analysis and presentation: Datasets were analyzed and combined in Microsoft Excel

and GraphPad Prism 8.0 software. Where indicated, two-tailed unpaired t-tests were

performed. We utilized Adobe Photoshop and Illustrator CS6 for figure production.

Results

Generation of CEM cells lacking SERINC3 and SERINC5

We utilized CRISPR/Cas9 gene editing to determine whether the modulation of SERINC3

and SERINC5 is required for Nef to enhance HIV-1 replication in CEM cells. We chose

the CEM cell line due to its ability to support robust HIV-1 replication and virion-infectivity

in a Nef-dependent manner (23). We hypothesized that Nef would remain an important

factor in promoting viral spread whether or not CEM cells expressed SERINC3 and

SERINC5, a result in conflict with data obtained using Jurkat T cells but consistent with

data recently obtained using MOLT-3 T cells. To test this, we used lentiviral vectors

(LentiCRISPR) expressing both Cas9 and gene-specific gRNAs (52). CEM cells were

transduced with viruses generated from a LentiCRISPR vector targeting SERINC3 exon

2 (34). We enriched for transduced cells using puromycin, then isolated and expanded

clones after limiting dilution. Twenty CEM cell clones were assessed for genome editing

using Tracking of Indels by Decomposition (TIDE) (55), utilizing PCR products containing

the SERINC3 gRNA target site amplified from the genomic DNA of the individual cell

clones. We cloned some of these PCR products into a bacterial plasmid vector and

sequenced ten independent plasmid clones from each putative single-cell clone to verify

specific edits. We identified a CEM line bearing three out-of-frame indels and no wild-type

sequences; these three distinct edited sequences persisted after a second round of

single-cell cloning and sequencing (Figure 1A). We concluded that cells are likely triploid

for this chromosome, and we designated this line CEM SERINC3 knockout (CEM:S3KO).

To generate CEM cells lacking both SERINC3 and SERINC5, we designed five gRNAs

targeting either Exon 1 or Exon 2 of SERINC5. The gRNAs were subcloned into a lentiviral

vector that encodes GFP in place of puromycin (LentiCRISPR-GFP). We transduced

CEM:S3KO cells with distinct lentiviruses, each encoding a different gRNA targeting

SERINC5. Three days post-transduction, CEM cells were sorted based on GFP

expression and expanded in bulk. TIDE analysis revealed that gRNA4 targeting exon 2

yielded the most promising evidence of genome editing (data not shown). We single-cell

cloned and expanded these cells in a similar manner as described above for the editing

of SERINC3. We identified two clones containing indels within the SERINC5 target region.

PCR amplification from these putative single-cell clones and sequencing of individual

plasmid clones of the PCR products indicated that these consisted of either a mixture of

two triploid cell lines ("clone 6") and a triploid edit ("clone 8"); these cell lines were termed

SERINC3/SERINC5 knockout clone 6 (CEM:S3/S5KO (6)) and CEM SERINC3/SERINC5

knockout clone 8 (CEM:S3/S5KO (8)) (Figure 1B). In both cases the cell lines encoded

out-of-frame indels in SERINC5 exon 2 and no wild-type sequences. The nine base-pair

insertion in clone 6 generated a stop codon. These three cells lines served as the basis

for our viral replication and infectivity studies.

Optimal HIV-1 spread in CEM cells is dependent on Nef but independent of SERINC3

and SERINC5.

To test whether Nef is required to counteract SERINC3 and SERINC5 to enhance HIV-1

replication, we infected CEM wildtype (WT), S3KO, S3/S5 KO (6) and S3/S5 KO (8) cells

with either Nef expressing (hereafter termed Nef+: NL4-3) or Nef lacking (hereafter

termed Nef-: NL4-3ΔNef) HIV-1 viruses at a multiplicity of infection (MOI)=0.001 infectious

units per cell. The inocula for these growth-rate experiments were produced from

HEK293T cells, and the MOI was based on the infectivity of the virus stocks measured in

CD4-positive HeLa-P4.R5 indicator cells. Thus, to infect $1x10^6$ CEM cells, we used

amounts of Nef+ or Nef- virus stocks that yielded 1,000 infectious centers in the HeLa

indicator assay. Although the infectivity of the viruses to CEM cells might be different than

to CD4-HeLa cells, we chose this approach rather than normalizing the inocula to the

content of p24 capsid antigen to adjust for the reduced infectivity of Nef- virions produced

by the HEK293T cells (data not shown). The CEM cell cultures were split every 3-4 days

and viral replication quantified by the amount of viral capsid (p24) contained within the

supernatant (cell-free: p24 ELISA) or associated with cells (intracellular flow) (Figure 2A).

As expected, we observed low but equal p24 production among all four CEM genotypes

whether the cells were infected with either Nef + (red line) or Nef – (blue line) expressing

virus (Figure 2B) at three days post-infection. This indicated that the standardization of

the inocula was likely accurate and that the infections were initiated at the same nominal

MOI. However, after day three, the Nef+ viruses propagated much more rapidly than the

Nef-, accumulating anywhere from 10- to 50-fold more p24 antigen in the culture

supernates at 6, 9 or 13 days post-infection in CEM WT cells (Figure 2B: Top left panel).

This corroborated the importance of Nef in enhancing viral replication in this *in vitro*

system. If Nef-mediated modulation of SERINC3 and/or SERINC5 were important for this

phenotype, then the attenuated replication of Nef- virus should be "rescued" in cell lacking

SERINC3 and/or SERINC5. Instead, Nef enhanced HIV-1 replication in CEM cells

knocked out for SERINC3 (Figure 2B: top right panel) or knocked out for SERINC3 and

SERINC5 (Figure 2B: bottom two panels). We observed a similar trend when analyzing

cell-associated virus at day 9 post-infection (Figure 2C) or cell-free virus from an

independent viral spreading experiment that utilized independently produced and titered

viral stocks (Figure 2D). Taken together, these data indicated that modulation of

SERINC3 and SERINC5 is not the primary mechanism by which Nef increases viral

spread in CEM cells.

Nef enhances HIV-1 virion-infectivity in CEM cells independently of SERINC3 and

SERINC5.

Various systems have shown that Nef enhances virion-infectivity in a SERINC- and cell-

type dependent manner (34). Therefore, we next asked whether counteraction of

SERINCs is necessary for Nef to enhance the infectiousness of virions. To test this, we

collected virions from the infected CEM cells at day 9, 13 or 15 post-infection and

measured single-cycle infectivity using HeLa TZM.bl reporter cells, which express

luciferase under the transcriptional control of the HIV-1 LTR (Figure 3A). We used these

cells because the luciferase read-out is more sensitive that the infectious center read-out

of the HeLa-P4.R5 cells, and the concentration of Nef- virus produced by the CEM

cultures was relatively low. Virions produced by Nef+ virus (day 9 post-infection) were 10-

to 50-fold more infectious per amount of p24 capsid antigen than those produced by Nef-

virus (day 13 post-infection), and this trend was observed regardless of SERINC3 and/or

SERINC5 expression (Figure 3B). We confirmed this result in an independent experiment

utilizing different viral stocks and analyzing virions collected at later times (Nef+: Day 13

post-infection versus Nef-: Day 15 post-infection; Figure 2C). Altogether, the data indicate

that Nef increases virion infectivity as well as growth-rate in CEM T cells independently

of SERINC3 and SERINC5.

Discussion

In this study, we sought to determine whether Nef's ability to enhance viral replication

and/or infectivity in CEM cells is primarily linked to modulation of SERINC3 and/or

SERINC5. We present evidence that argues against this by showing that Nef increases

viral replication rate in CEM cells lacking SERINC3 and SERINC5 and that virions of Nef+

virus are much more infectious than those of Nef- virus regardless of whether SERINC3

and SERINC5 are expressed in the CEM cells that produced them.

Our results indicate that Nef enhances virion-infectivity independently of SERINCs in

CEM T cells. This contrasts with reports in Jurkat and primary CD4+ cells, where Nef-

mediated enhancement of infectivity appears to correlate with SERINC5 expression (34,

35). How might these disparate results be explained? One reason could be differences in

experimental design. Most HIV-1 infectivity assays utilize virions produced from transient

overexpression systems (34, 35). Here, we used virions produced from infected CEM

cells either expressing or lacking endogenous SERINC3 and SERINC5.

Perhaps more likely are cell-type-specific differences. Our observations herein using

CEM cells are very similar to recent results reported using MOLT-3 cells (46). In MOLT-3

cells, a chimeric HIV-1 virus bearing the SERINC5 antagonist (glycoMA) failed to

substitute for Nef in rescuing HIV-1 infectivity (a property glycoMA should have if the sole

function of Nef were to counteract SERINC5), and knockout of SERINC5 did not rescue

the reduced growth rate of Nef- virus (46). Together, these data suggest that the SERINC-

dependence of the Nef infectivity phenotype is cell-type dependent (34, 35, 46), with

Jurkat and presumable primary CD4+ T cells being SERINC-dependent and CEM and

MOLT-3 cells being SERINC-independent. Thus, comparing virions produced from

infected primary CD4+ T cells either lacking or expressing SERINC5 may be particularly

insightful to adjudicate the relevance of observations made using T cell lines.

How does Nef's ability to enhance viral replication correlate with modulation of the

SERINCs? Jurkat cells lacking SERINC3 and SERINC5 support similar viral replication

when infected with HIV-1 viruses either expressing or lacking Nef (35). Furthermore,

substituting glycoMA for Nef rescues viral replication in Jurkat cells (34, 46). This

indicates that Nef's ability to enhance viral replication in these cells rests primarily on

modulation of SERINCs. However, glycoMA failed to rescue the replication of HIV-1

lacking Nef in MOLT-3 and primary CD4+ T cells (46). Moreover, MOLT-3 cells lacking

SERINC3 and SERINC5 support Nef's ability to enhance viral replication (46). Our results

herein indicate that CEM cells lacking SERINC3 and SERINC5 similarly support Nef's

ability to enhance viral replication. Together with the MOLT-3 data, these results indicate

that Nef-mediated enhancement of viral propagation can occur in a SERINC-independent

manner. Whether primary CD4+ T cells lacking SERINC5 support the attenuated

phenotype of Nef-deficient viruses remains unknown and is an important question to

answer.

What mechanisms might explain Nef-mediated enhancement of infectivity and/or viral

replication independently of SERINCs? One potential explanation could be Nef-mediated

modulation of Src family kinases (SFKs). Nef binds several SFK members such as Lck,

Hck, Lyn and c-Src through a conserved proline-rich (PxxP) motif contained within Nef's

Src homology region 3 (SH3) binding domain (56, 57), and primary Nef isolates from

HIV-1 Group M display a conserved ability to activate SFK's (58). Nef mutants lacking the

PxxP motif were reportedly able to downregulate CD4 but unable to enhance HIV-1

replication in peripheral blood mononuclear cells (57). However, other studies reported

that mutation of Nef's PxxP motif yielded little to no difference in HIV-1 replication within

MOLT-3 or primary CD4+ T-cells, and in CEM T cells the attenuated phenotype of

mutants of the SH3 binding domain was modest and seemed attributable to reduced

expression of Nef (25, 46, 59). Analyzing HIV-1 replication in T cells lacking one or more

SFKs may be necessary to adequately assess whether Nef's ability to enhance HIV-1

replication is mediated by these interactions.

One notable similarity between MOLT-3 and CEM lymphoblastoid cells that distinguishes

them from Jurkat cells is that they do not express the T cell receptor (TCR) at their

surfaces (60, 61). Given that Nef reportedly mimics TCR-signaling (22, 62), another

possibility is that MOLT-3 and CEM cells reveal a SERINC-independent growth-rate Nef-

phenotype that is exaggerated by the absence of TCR signaling in these cells. This could

be consistent with the initial observations that the activation-state of primary CD4 T cells

affects the Nef growth-rate phenotype (24). Nonetheless, how TCR signaling would affect

virion-infectivity is obscure.

Lastly, the explanation might reside in Nef's ability to interact with components of clathrin

trafficking pathways and to modulate many cellular proteins, one of which might underlie

the infectivity phenotype in MOLT-3 and CEM cells. Nef binds AP complexes via a

conserved sorting signal near its C-terminus: 160ExxxLL164,165 (63). This "di-leucine

motif" is required for both Nef-mediated CD4 downregulation and optimal viral infectivity

in CEM cells (64). The Nef LL164/165AA mutant, which is unable to bind AP-2 (65),

replicates poorly in MOLT-3 cells (46). Residues located within Nef's core domain and

required for downregulating CD4 and interacting with Dynamin-2 are also required for

enhancement of viral replication in MOLT-3 cells (46, 66, 67). Finally, Nef enhances HIV-

1 replication in the absence of CD4 downregulation in MOLT-3 cells (46) and in CEM-

derived cells (A2.01) engineered to express a CD4 lacking a cytoplasmic domain that is

unresponsive to Nef (68). These observations regarding CD4 are critically important,

since CD4 is itself a potent inhibitor of infectivity that is counteracted by Nef (11, 12). In

the CEM experiments herein, CD4 downregulation by Nef could conceivably account for

the observed infectivity and growth-rate phenotypes. Nonetheless, even if that were the

case, the data would indicate that the contribution of CD4 to these phenotypes far

outweighs the contribution of the SERINC3 and SERINC5 in these cells.

Taken together, these results raise the intriguing possibility that Nef binds and internalizes

a novel factor within CEM as well as MOLT-3 cells that is necessary for enhancing viral

infectivity and replication. Utilizing both MOLT-3 and CEM cells might facilitate the search

for such a factor, providing a more complete understanding of the enigmatic yet important

virologic effects of Nef.

References

References

1.   Kestler HW, Ringler DJ, Mori K, Panicali DL, Sehgal PK, Daniel MD, Desrosiers RC.

1991. Importance of the nef gene for maintenance of high virus loads and for

development of AIDS. Cell 65:651–662.

2.   Kirchhoff F, Greenough TC, Brettler DB, Sullivan JL, Desrosiers RC. 1995. Absence

of Intact nef Sequences in a Long-Term Survivor with Nonprogressive HIV-1

Infection. New England Journal of Medicine 332:228–232.

3.   Deacon NJ, Tsykin A, Solomon A, Smith K, Ludford-Menting M, Hooker DJ, McPhee

DA, Greenway AL, Ellett A, Chatfield C, Lawson VA, Crowe S, Maerz A, Sonza S,

Learmont J, Sullivan JS, Cunningham A, Dwyer D, Dowton D, Mills J. 1995. Genomic

structure of an attenuated quasi species of HIV-1 from a blood transfusion donor and

recipients. Science 270:988–991.

4.   Kirchhoff F. 2010. Immune evasion and counteraction of restriction factors by HIV-1

and other primate lentiviruses. Cell Host Microbe 8:55–67.

5.   Garcia JV, Alfano J, Miller AD. 1993. The negative effect of human immunodeficiency

virus type 1 Nef on cell surface CD4 expression is not species specific and requires

the cytoplasmic domain of CD4. J Virol 67:1511–1516.

6.   Aiken C, Konner J, Landau NR, Lenburg ME, Trono D. 1994. Nef induces CD4

endocytosis: requirement for a critical dileucine motif in the membrane-proximal CD4

cytoplasmic domain. Cell 76:853–864.

7. Chaudhuri R, Lindwasser OW, Smith WJ, Hurley JH, Bonifacino JS. 2007.

Downregulation of CD4 by human immunodeficiency virus type 1 Nef is dependent

on clathrin and involves direct interaction of Nef with the AP2 clathrin adaptor. J Virol

81:3877–3890.

8. Kwon Y, Kaake RM, Echeverria I, Suarez M, Shamsabadi MK, Stoneham C, Ramirez

PW, Kress J, Singh R, Sali A, Krogan N, Guatelli J, Jia X. 2020. Structural basis of

CD4 downregulation by HIV-1 Nef. Nature Structural & Molecular Biology in

press:22.

9. Wildum S, Schindler M, Münch J, Kirchhoff F. 2006. Contribution of Vpu, Env, and

Nef to CD4 down-modulation and resistance of human immunodeficiency virus type

1-infected T cells to superinfection. J Virol 80:8047–8059.

10. Alsahafi N, Ding S, Richard J, Markle T, Brassard N, Walker B, Lewis GK, Kaufmann

DE, Brockman MA, Finzi A. 2016. Nef Proteins from HIV-1 Elite Controllers Are

Inefficient at Preventing Antibody-Dependent Cellular Cytotoxicity. J Virol 90:2993–

3002.

11. Ross TM, Oran AE, Cullen BR. 1999. Inhibition of HIV-1 progeny virion release by

cell-surface CD4 is relieved by expression of the viral Nef protein. Curr Biol 9:613–

621.

12. Lama J, Mangasarian A, Trono D. 1999. Cell-surface expression of CD4 reduces

HIV-1 infectivity by blocking Env incorporation in a Nef- and Vpu-inhibitable manner.

Curr Biol 9:622–631.

13.   Kasper MR, Roeth JF, Williams M, Filzen TM, Fleis RI, Collins KL. 2005. HIV-1 Nef

disrupts antigen presentation early in the secretory pathway. J Biol Chem

280:12840–12848.

14.   Roeth JF, Williams M, Kasper MR, Filzen TM, Collins KL. 2004. HIV-1 Nef disrupts

MHC-I trafficking by recruiting AP-1 to the MHC-I cytoplasmic tail. J Cell Biol

167:903–913.

15.   Noviello CM, Benichou S, Guatelli JC. 2008. Cooperative binding of the class I major

histocompatibility complex cytoplasmic domain and human immunodeficiency virus

type 1 Nef to the endosomal AP-1 complex via its mu subunit. J Virol 82:1249–1258.

16.   Jia X, Singh R, Homann S, Yang H, Guatelli J, Xiong Y. 2012. Structural basis of

evasion of cellular adaptive immunity by HIV-1 Nef. Nat Struct Mol Biol 19:701–706.

17.   Hung C-H, Thomas L, Ruby CE, Atkins KM, Morris NP, Knight ZA, Scholz I, Barklis

E, Weinberg AD, Shokat KM, Thomas G. 2007. HIV-1 Nef assembles a Src family

kinase-ZAP-70/Syk-PI3K cascade to downregulate cell-surface MHC-I. Cell Host

Microbe 1:121–133.

18.   Blagoveshchenskaya AD, Thomas L, Feliciangeli SF, Hung CH, Thomas G. 2002.

HIV-1 Nef downregulates MHC-I by a PACS-1- and PI3K-regulated ARF6 endocytic

pathway. Cell 111:853–866.

19.   Dikeakos JD, Atkins KM, Thomas L, Emert-Sedlak L, Byeon I-JL, Jung J, Ahn J,

Wortman MD, Kukull B, Saito M, Koizumi H, Williamson DM, Hiyoshi M, Barklis E,

Takiguchi M, Suzu S, Gronenborn AM, Smithgall TE, Thomas G. 2010. Small

molecule inhibition of HIV-1-induced MHC-I down-regulation identifies a temporally

regulated switch in Nef action. Mol Biol Cell 21:3279–3292.

20. Collins KL, Chen BK, Kalams SA, Walker BD, Baltimore D. 1998. HIV-1 Nef protein

protects infected primary cells against killing by cytotoxic T lymphocytes. Nature

391:397–401.

21. Haller C, Rauch S, Fackler OT. 2007. HIV-1 Nef employs two distinct mechanisms

to modulate Lck subcellular localization and TCR induced actin remodeling. PLoS

ONE 2:e1212.

22. Simmons A, Aluvihare V, McMichael A. 2001. Nef Triggers a Transcriptional Program

in T Cells Imitating Single-Signal T Cell Activation and Inducing HIV Virulence

Mediators. Immunity 14:763–777.

23. Chowers MY, Spina CA, Kwoh TJ, Fitch NJ, Richman DD, Guatelli JC. 1994. Optimal

infectivity in vitro of human immunodeficiency virus type 1 requires an intact nef

gene. J Virol 68:2906–2914.

24. Spina CA, Kwoh TJ, Chowers MY, Guatelli JC, Richman DD. 1994. The importance

of nef in the induction of human immunodeficiency virus type 1 replication from

primary quiescent CD4 lymphocytes. J Exp Med 179:115–123.

25. Lundquist CA, Tobiume M, Zhou J, Unutmaz D, Aiken C. 2002. Nef-mediated

downregulation of CD4 enhances human immunodeficiency virus type 1 replication

in primary T lymphocytes. J Virol 76:4625–4633.

26. Münch J, Rajan D, Schindler M, Specht A, Rücker E, Novembre FJ, Nerrienet E,

Müller-Trutwin MC, Peeters M, Hahn BH, Kirchhoff F. 2007. Nef-Mediated

Enhancement of Virion Infectivity and Stimulation of Viral Replication Are

Fundamental Properties of Primate Lentiviruses. JVI 81:13852–13864.

27. Carl S, Greenough TC, Krumbiegel M, Greenberg M, Skowronski J, Sullivan JL,

Kirchhoff F. 2001. Modulation of different human immunodeficiency virus type 1 Nef

functions during progression to AIDS. J Virol 75:3657–3665.

28. Crotti A, Neri F, Corti D, Ghezzi S, Heltai S, Baur A, Poli G, Santagostino E, Vicenzi

E. 2006. Nef Alleles from Human Immunodeficiency Virus Type 1-InfectedLong-

Term-Nonprogressor Hemophiliacs with or without Late Disease Progression Are

Defective in Enhancing Virus Replication and CD4 Down-Regulation. JVI 80:10663–

10674.

29. Aiken C, Trono D. 1995. Nef stimulates human immunodeficiency virus type 1

proviral DNA synthesis. J Virol 69:5048–5056.

30. Pandori MW, Fitch NJ, Craig HM, Richman DD, Spina CA, Guatelli JC. 1996.

Producer-cell modification of human immunodeficiency virus type 1: Nef is a virion

protein. Journal of virology 70:4283–4290.

31. Pizzato M, Helander A, Popova E, Calistri A, Zamborlini A, Palù G, Göttlinger HG.

2007. Dynamin 2 is required for the enhancement of HIV-1 infectivity by Nef. Proc

Natl Acad Sci USA 104:6812–6817.

32. Pizzato M, Popova E, Göttlinger HG. 2008. Nef can enhance the infectivity of

receptor-pseudotyped human immunodeficiency virus type 1 particles. J Virol

82:10811–10819.

33. Pizzato M. 2010. MLV glycosylated-Gag is an infectivity factor that rescues Nef-

deficient HIV-1. Proc Natl Acad Sci USA 107:9364–9369.

34. Rosa A, Chande A, Ziglio S, De Sanctis V, Bertorelli R, Goh SL, McCauley SM,

Nowosielska A, Antonarakis SE, Luban J, Santoni FA, Pizzato M. 2015. HIV-1 Nef

promotes infection by excluding SERINC5 from virion incorporation. Nature

526:212–217.

35. Usami Y, Wu Y, Göttlinger HG. 2015. SERINC3 and SERINC5 restrict HIV-1

infectivity and are counteracted by Nef. Nature 526:218–223.

36. Shi J, Xiong R, Zhou T, Su P, Zhang X, Qiu X, Li H, Li S, Yu C, Wang B, Ding C,

Smithgall TE, Zheng Y-H. 2018. HIV-1 Nef Antagonizes SERINC5 Restriction by

Downregulation of SERINC5 via the Endosome/Lysosome System. J Virol 92.

37. Heigele A, Kmiec D, Regensburger K, Langer S, Peiffer L, Stürzel CM, Sauter D,

Peeters M, Pizzato M, Learn GH, Hahn BH, Kirchhoff F. 2016. The Potency of Nef-

Mediated SERINC5 Antagonism Correlates with the Prevalence of Primate

Lentiviruses in the Wild. Cell Host Microbe 20:381–391.

38. Chande A, Cuccurullo EC, Rosa A, Ziglio S, Carpenter S, Pizzato M. 2016. S2 from

equine infectious anemia virus is an infectivity factor which counteracts the retroviral

inhibitors SERINC5 and SERINC3. Proc Natl Acad Sci USA 113:13197–13202.

39. Beitari S, Ding S, Pan Q, Finzi A, Liang C. 2017. Effect of HIV-1 Env on SERINC5

Antagonism. J Virol 91.

40. Angerstein AO, Stoneham CA, Ramirez PW, Guatelli JC, Vollbrecht T. 2020.

Sensitivity to monoclonal antibody 447-52D and an open env trimer conformation

correlate poorly with inhibition of HIV-1 infectivity by SERINC5. Virology 548:73–81.

41. Firrito C, Bertelli C, Vanzo T, Chande A, Pizzato M. 2018. SERINC5 as a New

Restriction Factor for Human Immunodeficiency Virus and Murine Leukemia Virus.

Annu Rev Virol 5:323–340.

42. Inuzuka M, Hayakawa M, Ingi T. 2005. Serinc, an activity-regulated protein family,

incorporates serine into membrane lipid synthesis. J Biol Chem 280:35776–35783.

43. Trautz B, Wiedemann H, Lüchtenborg C, Pierini V, Kranich J, Glass B, Kräusslich H-

G, Brocker T, Pizzato M, Ruggieri A, Brügger B, Fackler OT. 2017. The host-cell

restriction factor SERINC5 restricts HIV-1 infectivity without altering the lipid

composition and organization of viral particles. J Biol Chem 292:13702–13713.

44. Sood C, Marin M, Chande A, Pizzato M, Melikyan GB. 2017. SERINC5 protein

inhibits HIV-1 fusion pore formation by promoting functional inactivation of envelope

glycoproteins. J Biol Chem 292:6014–6026.

45. Zhang X, Shi J, Qiu X, Chai Q, Frabutt DA, Schwartz RC, Zheng Y-H. 2019. CD4

expression and Env conformation are critical for HIV-1 restriction by SERINC5. J

Virol JVI.00544-19, jvi;JVI.00544-19v1.

46. Wu Y, Olety B, Weiss ER, Popova E, Yamanaka H, Göttlinger H. 2019. Potent

Enhancement of HIV-1 Replication by Nef in the Absence of SERINC3 and

SERINC5. mBio 10:e01071-19, /mbio/10/3/mBio.01071-19.atom.

47. Platt EJ, Bilska M, Kozak SL, Kabat D, Montefiori DC. 2009. Evidence that ecotropic

murine leukemia virus contamination in TZM-bl cells does not affect the outcome of

neutralizing antibody assays with human immunodeficiency virus type 1. J Virol

83:8289–8292.

48. Platt EJ, Wehrly K, Kuhmann SE, Chesebro B, Kabat D. 1998. Effects of CCR5 and

CD4 cell surface concentrations on infections by macrophagetropic isolates of

human immunodeficiency virus type 1. J Virol 72:2855–2864.

49. Derdeyn CA, Decker JM, Sfakianos JN, Wu X, O'Brien WA, Ratner L, Kappes JC,

Shaw GM, Hunter E. 2000. Sensitivity of human immunodeficiency virus type 1 to

the fusion inhibitor T-20 is modulated by coreceptor specificity defined by the V3 loop

of gp120. J Virol 74:8358–8367.

50. Wei X, Decker JM, Liu H, Zhang Z, Arani RB, Kilby JM, Saag MS, Wu X, Shaw GM,

Kappes JC. 2002. Emergence of resistant human immunodeficiency virus type 1 in

patients receiving fusion inhibitor (T-20) monotherapy. Antimicrob Agents

Chemother 46:1896–1905.

51. Takeuchi Y, McClure MO, Pizzato M. 2008. Identification of gammaretroviruses

constitutively released from cell lines used for human immunodeficiency virus

research. J Virol 82:12585–12588.

52. Sanjana NE, Shalem O, Zhang F. 2014. Improved vectors and genome-wide libraries
for CRISPR screening. Nat Methods 11:783–784.

53. Shalem O, Sanjana NE, Hartenian E, Shi X, Scott DA, Mikkelson T, Heckl D, Ebert
BL, Root DE, Doench JG, Zhang F. 2014. Genome-scale CRISPR-Cas9 knockout
screening in human cells. Science 343:84–87.

54. Day JR, Martínez LE, Šášik R, Hitchin DL, Dueck ME, Richman DD, Guatelli JC.
2006. A computer-based, image-analysis method to quantify HIV-1 infection in a
single-cycle infectious center assay. Journal of Virological Methods 137:125–133.

55. Brinkman EK, Chen T, Amendola M, van Steensel B. 2014. Easy quantitative
assessment of genome editing by sequence trace decomposition. Nucleic Acids Res
42:e168.

56. Trible RP, Emert-Sedlak L, Smithgall TE. 2006. HIV-1 Nef Selectively Activates Src
Family Kinases Hck, Lyn, and c-Src through Direct SH3 Domain Interaction. J Biol
Chem 281:27029–27038.

57. Saksela K, Cheng G, Baltimore D. 1995. Proline-rich (PxxP) motifs in HIV-1 Nef bind
to SH3 domains of a subset of Src kinases and are required for the enhanced growth
of Nef+ viruses but not for down-regulation of CD4. The EMBO Journal 14:484–491.

58. Narute PS, Smithgall TE. 2012. Nef Alleles from All Major HIV-1 Clades Activate Src-
Family Kinases and Enhance HIV-1 Replication in an Inhibitor-Sensitive Manner.
PLoS ONE 7:e32561.

59. Craig HM, Pandori MW, Riggs NL, Richman DD, Guatelli JC. 1999. Analysis of the SH3-Binding Region of HIV-1 Nef: Partial Functional Defects Introduced by Mutations in the Polyproline Helix and the Hydrophobic Pocket. Virology 262:55–63.

60. Greenberg JM, Gonzalez-Sarmiento R, Arthur DC, Wilkowski CW, Streifel BJ, Kersey JH. 1988. Immunophenotypic and cytogenetic analysis of Molt-3 and Molt-4: human T-lymphoid cell lines with rearrangement of chromosome 7. Blood 72:1755–1760.

61. Royer HD, Ramarli D, Acuto O, Campen TJ, Reinherz EL. 1985. Genes encoding the T-cell receptor alpha and beta subunits are transcribed in an ordered manner during intrathymic ontogeny. Proc Natl Acad Sci USA 82:5510–5514.

62. Len ACL, Starling S, Shivkumar M, Jolly C. 2017. HIV-1 Activates T Cell Signaling Independently of Antigen to Drive Viral Spread. Cell Reports 18:1062–1074.

63. Sugden SM, Bego MG, Pham TNQ, Cohen ÉA. 2016. Remodeling of the Host Cell Plasma Membrane by HIV-1 Nef and Vpu: A Strategy to Ensure Viral Fitness and Persistence. Viruses 8:67.

64. Craig HM, Pandori MW, Guatelli JC. 1998. Interaction of HIV-1 Nef with the cellular dileucine-based sorting pathway is required for CD4 down-regulation and optimal viral infectivity. Proc Natl Acad Sci USA 95:11229–11234.

65. Bresnahan PA, Yonemoto W, Ferrell S, Williams-Herman D, Geleziunas R, Greene WC. 1998. A dileucine motif in HIV-1 Nef acts as an internalization signal for CD4 downregulation and binds the AP-1 clathrin adaptor. Curr Biol 8:1235–1238.

66. Cohen GB, Rangan VS, Chen BK, Smith S, Baltimore D. 2000. The human

thioesterase II protein binds to a site on HIV-1 Nef critical for CD4 down-regulation.

J Biol Chem 275:23097–23105.

67. Poe JA, Smithgall TE. 2009. HIV-1 Nef dimerization is required for Nef-mediated

receptor downregulation and viral replication. J Mol Biol 394:329–342.

68. Chowers MY, Pandori MW, Spina CA, Richman DD, Guatelli JC. 1995. The growth

advantage conferred by HIV-1 nef is determined at the level of viral DNA formation

and is independent of CD4 downregulation. Virology 212:451–457.

Acknowledgments

We thank Dr. Vicente Planelles for the LentiCRISPR-GFP plasmid. We thank The

Pendleton Charitable Trust for equipment. P.W.R. was partly supported by NIH grant

K12GM068524. This work was supported by NIH grant R01 AI129706 to J.G.

Figure Legends

Figure 1: CRISPR/Cas9 editing of SERINC3 and SERINC5 in CEM cells. A portion of the wildtype (WT) sequence from exon 2 of either SERINC3 or SERINC5 is shown in each panel, with the gRNA target site highlighted in bold. Inserted nucleotides are in green, stop codons in red, and substituted nucleotides in blue. Deletions are depicted as a dashed line. All mutant alleles sequenced produced frameshift mutations. A.) The genotype in SERINC3 knockout (S3KO) cells consisted of alleles bearing a 1 bp insertion, 2 bp insertion, and 19 bp deletion. B.) We identified and utilized two different clones knocked out for SERINC5: clone 8 (S3/S5KO (8)) and clone 6 (S3/S5KO (6)). S3/S5KO (8) contained the following indels: 2 bp deletion, 4 bp deletion, and a 7 bp deletion (top) while S3/S5KO (6) contained a 2 bp deletion, 3 bp deletion/1 bp insertion, 4 bp deletion, 9 bp insertion with the introduction of a stop codon, 10 bp deletion, and 31 bp deletion.

Figure 2: Nef enhances HIV-1 replication independently of SERINC3 and SERINC5 in CEM cells. A.) Schematic of viral replication studies performed in CEM wildtype (WT), SERINC3 knockout (S3KO), SERINC5 knockout clone 8 (S5KO (8)) or SERINC5 knockout clone (6 S5KO (6)) cells. Each cell line was infected with either NL4-3 (termed Nef+) or HIV-1ΔNef (termed Nef-) at a low inoculum (MOI=0.001). The cultures were split every 3-4 days and viral growth measured as indicated. B.) Viral replication quantified by p24 Capsid ELISA in the supernatants of CEM WT, S3KO, S5KO (8) and S5KO cultures (6). Results depict either three (WT, S3KO, S5KO (8)) or one (S5KO (6)) independent infections measured in duplicate at each time point. C.) Flow cytometry of p24 positive cells at day 9 post-infection. Data are compiled from three independent experiments. D.)

HIV-1 replication measured by p24 ELISA using different viral stocks in a separate

independent experiment. Two-tailed unpaired t-tests were performed where indicated.

Figure 3: HIV-1 requires Nef for optimal infectivity in CEM cells regardless of SERINC3

or SERINC5 expression. A.) Schematic depicting measurement of single-cycle infectivity

with virions produced from infected CEM WT, S3KO, S5KO (8) or S5KO cultures (6).

HeLa Tzm.bl indicator cells contain a luciferase gene under the transcriptional control of

the HIV-1 LTR. B.) Infectivity data (relative luciferase units (RLU) normalized to p24

(RLU/p24)) from virions collected from cultures of either CEM WT cells or cells knocked

out for SERINC3 or SERINC3 and SERINC5 infected with either Nef+ (day 9 post-

infection) or Nef- (day 13 post-infection) viruses. Results are representative of one

experiment performed in triplicate. C.) Infectivity data (RLU/p24) for virions produced at

day 12 (Nef+) or day 15 (Nef-) post-infection of CEM WT or S3/S5 KO(8) cells. Results

represent one experiment that was fully independent from that shown in panel B and

performed in triplicate. D.) Combined RLU/24 data from Figure 3B and 3C in CEM WT

and S3/S5 KO(8) cells. Two-tailed unpaired t-tests were performed where indicated.

Figure 1

A *SERINC3 knockout: Exon 2*

<u>Target Site</u>

Gene: AATGAAAGC***ATAAATGAGGCGAGTCACCG***TGGAATTCT

S3:KO

    AATGAAAGC***ATAAATGAGGCGAGTCA****A****CCG***TGGAATTCT

    AATGAAAGC***ATAAATGAGGCGAGTCA****TT****CCG***TGGAATTCT

    AATGAAAGC***ATAAATGAG*** – – – – – – – – – – – – – – – – – – G

B *SERINC5 knockout (clone 8): Exon 2*

<u>Target Site</u>

Gene: CCTCTGCTG***CATCATGATGTCAACAACCG***TGG

S5:KO (8)

    CCTCTGCTG***CATCATGATGTCAACA*** – – ***AG***TGG

    CCTCTGCTG***CATCATGATGTCAA*** – – – – ***CG***TGG

    CCTCTGCTG***CATCATGATGT*** – – – – – – ***CA***TGG

C *SERINC5 knockout (clone 6): Exon 2*

<u>Target Site</u>

Gene: CCTCTGCTG***CATCATGATGTCAACAACCG***TGGCTCA

S5:KO (6)

    CCTCTGCTG***CATCATGATGTCAACAA*** – – ***G***TGGCTCA

    CCTCTGCTG***CATCATGATGTCAAC*** – – – ***AA***GTGGCTCA

    CCTCTGCTG***CATCATGATGTCAAC*** – – – – ***G***TGGCTCA

    CCTCTGCTG***CATCATGATGTCA****TCTTGATTTGATCTTG***TGGCTCA

    CCTCTGCTG***CATCATGAT*** – – – – – – – – – ***G***TGGCTCA

    CC – TCA

Figure 3

37