Nef enhances HIV-1 replication and infectivity independently of

SERINC3 and SERINC5 in CEM T cells

Peter W. Ramirez[a,#,*], Thomas Vollbrecht[b,c,*], Francisco M. Acosta[a], Marissa Suarez[b,c],

Aaron A. Angerstein[b,c], Jared Wallace[d], Ryan M. O' Connell[d], and John Guatelli[b,c]

[a]Department of Biological Sciences, California State University Long Beach, Long

Beach, CA.

[b]Department of Medicine, University of California San Diego, La Jolla, California, USA.

[c]VA San Diego Healthcare System, San Diego, California, USA.

[d]Division of Microbiology and Immunology, Department of Pathology, The University of

Utah, Salt Lake City, UT.

Running Head: Attenuation of HIV-1∆Nef without *serinc3 or 5*

[#]Address Correspondence to Peter W. Ramirez: Peter.Ramirez@csulb.edu

[*]These authors contributed equally to this work.

Abstract

The lentiviral *nef* gene encodes several discrete activities aimed at co-opting or antagonizing cellular proteins and pathways to defeat host defenses and maintain persistent infection. Primary functions of Nef include downregulation of CD4 and MHC class-I from the cell surface, disruption or mimicry of T-cell receptor signaling, and enhancement of viral infectivity by counteraction of the host antiretroviral proteins SERINC3 and SERINC5. In the absence of Nef, SERINC5 incorporates into virions and inhibits viral fusion with target cells, decreasing infectivity. However, whether Nef's counteraction of SERINC5 is the cause of its positive influence on viral growth-rate in CD4-positive T cells is unclear. Here, we utilized CRISPR/Cas9 to knockout SERINC3 and SERINC5 in a leukemic CD4-positive T cell line (CEM) that displays relatively robust *nef*-related infectivity and growth-rate phenotypes. As previously reported, viral replication was attenuated in CEM cells infected with HIV-1 lacking Nef (HIV-1ΔNef). This attenuated growth-rate phenotype was observed regardless of whether the coding regions of the *serinc3* or *serinc5* genes were intact. Moreover, knockout of *serinc3* or *serinc5* failed to restore the infectivity of HIV1ΔNef virions produced from infected CEM cells. Taken together, our results corroborate a similar study using another T-lymphoid cell line (MOLT-3) and indicate that the antagonism of SERINC3 and SERINC5 cannot fully explain the virology of HIV-1 lacking Nef.

Introduction

Primate lentiviruses encode several accessory gene products that facilitate viral reproduction and persistence, in some cases by providing evasion of the host immune response. The lentiviral Nef protein accelerates viral pathogenesis and progression to AIDS in SIV infected rhesus macaques (1) and enhances disease progression in humans infected with HIV-1 (2, 3). Nef is a small, myristoylated, peripheral membrane protein with many well conserved activities, including the downregulation of cell surface proteins such as CD4 and MHC class-I (MHC-I), the modulation of T cell activation, and the enhancement of viral infectivity and growth-rate (4).

To downregulate CD4, Nef binds the cytoplasmic domain of CD4 and links it to the clathrin Adaptor Protein 2 (AP-2) complex, internalizing CD4 from the plasma membrane and delivering it to lysosomes for degradation (5–8). The targeting of CD4 by Nef contributes to viral replication in several ways. CD4 downregulation prevents superinfection of cells and consequent premature cell-death, ensuring adequate time for viral replication (9). It also contributes to inhibiting antibody dependent cellular cytotoxicity (ADCC) that recognizes CD4-induced epitopes with Env (10). Potentially directly relevant to the work presented here, CD4 incorporates into virions and inhibits virion-infectivity if not downregulated by Nef (11, 12).

To downregulate MHC-I, Nef has been proposed to use two non-mutually exclusive mechanisms: 1) Nef and the clathrin-adaptor AP-1 intercept *de novo* synthesized MHC-I molecules within the *trans*-Golgi network (TGN), leading to lysosomal degradation (13–

16) ; and/or 2) Nef retains internalized MHC-I molecules in the TGN via induction of a

Src family kinase/phosphoinositide 3-kinase signaling cascade (17–19). In either case,

reducing the expression of MHC-I molecules at the plasma membrane protects HIV-1

infected cells from lysis by cytotoxic T-lymphocytes, contributing to immune evasion

(20).

Nef interacts directly with Src-family kinases including the lymphocyte specific kinase

Lck, which it downregulates from the cell surface (21). The many studies of Nef's

influence on T cell activation are difficult to reconcile, but transcriptional profiling

suggests that the expression of Nef mimics signaling through the T cell receptor (22).

Apart from the model of MHC-I downregulation noted above, a clear mechanistic link

between the influence of HIV-1 Nef on T cell activation and its influence on the

expression of cell surface receptors is lacking.

Nef stimulates HIV-1 replication and enhances virion-infectivity in many cell culture

systems, including various T cell lines, primary CD4-positive T cells, and human

lymphoid tissue (23–26). Diverse *nef* alleles from humans and primates maintain the

ability to enhance HIV-1 replication and infectivity, suggesting these functions are

important for establishing and maintaining persistent infection (26–28). Expression of

Nef within virion-producer cells and encoded either in *cis* or in *trans* relative to the viral

genome enhances HIV-1 replication and yields virions of greater infectivity (29, 30).

Nef's ability to enhance infectivity requires cellular components involved in vesicular

trafficking (dynamin 2, AP-2, and clathrin) and is also determined by the Envelope (Env)

glycoprotein (31, 32). A survey of cell lines identified murine leukemia virus glycosylated

Gag (MLV glycoGag), a protein structurally unrelated to Nef, as an infectivity factor that

rescues Nef-deficient HIV-1 virions (33). These and other features suggested that Nef

counteracts a cellular factor that restricts viral infectivity and possibly replication.

Two groups identified the host transmembrane protein SERINC5, and to a lesser
degree

SERINC3, as an inhibitor of HIV-1 virion-infectivity that is counteracted by Nef (34, 35).

Nef downregulates SERINC5 from the plasma membrane via a clathrin/AP-2 and

Rab5/7 endo-lysosomal pathway (34, 36), reducing the incorporation of SERINC5 in

HIV-1 virions. This in turn correlates with more efficient fusion of virions with target cells

and greater infectivity (34, 35). Nef's ability to counteract SERINC5 is conserved across

primate lentiviruses and correlates with the prevalence of these viruses in the wild (34,

37). Modulation of SERINC5 extends to other retroviral proteins, including S2 from

equine infectious anemia virus (EIAV) as well as MLV glycoGag (34, 35, 38). HIV-1 Env

glycoproteins are differentially sensitive to SERINC-mediated restriction when produced

from CD4-negative cells in single-round replication assays; sensitivity correlates to

some extent with the degree of Env-trimer openness and instability (34, 35, 39, 40).

SERINCs comprise a family of five genes that are evolutionarily conserved from yeast to

mammals (41). They encode multi-pass transmembrane proteins that support serine

specific phospholipid biosynthesis (hence their name: serine incorporator) (42), yet this

function does not seem to account for their anti-retroviral activity (43). Rather, SERINC5

appears to disrupt the formation of fusion pores between HIV-1 virions and target cells

(44) in an Env-conformation and CD4-dependent manner (45).

Initial studies of the Nef-SERINC relationship focused on the Jurkat T cell line, due to

the large defect in the infectivity of Nef-negative virions produced from these cells and

their relatively high levels of SERINC5 mRNA. Studies using another CD4-positive T cell

line, MOLT-3, have recently cast doubt on whether SERINC family proteins are

sufficient to explain the virologic phenotypes of Nef (46). In support of a SERINC-

dependent mechanism, Nef does not enhance HIV-1 infectivity and replication-rate in

Jurkat T cells when SERINC3 and SERINC5 are knocked out (34, 35). Moreover, a

minimal MLV glycoGag (termed glycoMA) can functionally replace Nef with respect to

viral replication rate and virion-infectivity when the virus is propagated using Jurkat cells

(46). In contrast, Nef, but not glycoMA, enhances HIV-1 replication in MOLT-3 cells. The

Nef-effect in MOLT-3 cells persists when the cells are knocked out for SERINC3 and

SERINC5, indicating that these restriction factors are not necessary for the virologic

effects of Nef in this setting (46). Remarkably, glycoMA cannot substitute functionally for

Nef with respect to stimulating viral replication in primary CD4-positive T cells. This

suggests that the growth rate enhancing effect of Nef in primary T cells is unrelated to

SERINC-antagonism (46), even though the virion-infectivity enhancing effect of Nef

reportedly is (34).

Given these conflicting results, we aimed to further test the hypothesis that Nef

enhances HIV-1 replication in a SERINC-dependent manner. To do this, we returned to

the CD4-positive T cell line in which we originally observed a stimulation of growth-rate

by Nef, an effect that was associated with Nef-mediated enhancement of virion-

infectivity (CEM) (23). Here, we show that neither the attenuated replication-rate of *nef*-

deficient HIV-1 in CEM T cells nor the reduced virion-infectivity of *nef*-deficient HIV-1

produced by these cells is rescued by CRISPR/cas9 editing of *serinc5,* either with or

without additional editing of *serinc3.* These results support those documented using

MOLT-3 and primary CD4-positive T cells and suggest that how Nef stimulates HIV-1

replication and virion infectivity remains unclear (46).

Materials and Methods

Cell Lines and Plasmids: HEK293T (a generous gift from Dr. Ned Landau) and HeLa

TZM-bl cells (Dr. John Kappes and Xiaoyun Wu,: NIH AIDS Reagent Program, Division

of AIDS, NIAID, NIH) (47, 48)   were grown in DMEM media (Thermo Fisher Scientific)

supplemented with 10% FBS (Hyclone) and 1% Penicillin/Streptomycin (Thermo Fisher

Scientific). HeLa P4.R5 (obtained from Dr. Ned Landau) were cultured in 10% FBS, 1%

Penicillin/Streptomycin and 1 μg puromycin. Both HeLa cell line derivatives express

CD4, CXCR4 and CCR5 and contain either a Tat-inducible β-galactosidase (HeLa

P4.R5) or both the β-galactosidase and luciferase (HeLa TZM-bl) genes under the

transcriptional control of the HIV-1 LTR. CEM (a generous gift from Dr. Douglas

Richman) and JTAg cells expressing (JTAg WT) or lacking SERINC3 and SERINC5

(JTAg SERINC3/SERINC5 KO; kindly provided by Dr. Heinrich Gottlinger) are T cell

leukemic clones that were cultured in RPMI 1640 media plus 10% FBS and 1%

Penicillin/Streptomycin (Thermo Fisher Scientific). The proviral plasmids pNL4-3 and

pNL4-3ΔNef have been previously described (23). LentiCRISPRv2 (Addgene; Catalog

#: 52961) contains a single guide RNA (sgRNA) targeting Exon 2 of SERINC3

(5'ATAAATGAGGCGAGTCACCG-3') and was a gift from Dr. Massimo Pizzato (34).

The lentiviral packaging (pxRSV-Rev, pMDLg/pRRE) and envelope (pMD2.G) plasmids

were kindly provided by Dr. Dan Gibbs. LentiCRISPR-GFP encodes GFP in place of

puromycin (49). Five sgRNAs targeting either Exon 1 or Exon 2 of SERINC5 were

designed using an online CRISPR tool (benchling.com) and cloned into LentiCRISPR-

GFP using previously described methods (50, 51). The sgRNA sequences were as

follows: sgRNA-SERINC5(1), 5'- ACA GCACTGAGCTGACATCG-3' ; sgRNA-

SERINC5(2), 5'GCACTGAGCTGACATCGC GG-3' ; sgRNA-SERINC5(3), 5'-

CTTCGTTCAAGTGTGAGCTG'3'   ;   sgRNA-SERINC5(4),   5'-

CATCATGATGTCAACAACCG-3'   ;   sgRNA-SERINC5(5),   5'-

TGAGGGACTGCCGAATCCTG-3'. Briefly, sgRNA oligos were designed to produce the

same overhangs after *BsmBI* digestion (5'- *CACCG*(sgRNA Oligo #1)-3' ; 3'-*C*(sgRNA

Oligo #2)-*CAAA*-3'). The oligos were phosphorylated (T4 Polynucleotide Kinase; NEB)

and annealed in a thermal cycler according to the following conditions: 37°C for 30

minutes; 95°C for 5 minutes with a ramp down to 25°C at 5°C/minute. Diluted oligos

(1:200) were ligated (T4 ligase; NEB) into dephosphorylated (Fast AP; Fermentas) and

BsmBI digested (Fast *BsmBI*; Fermentas) LentiCRISPR-GFP by overnight incubation at

16°C, followed by transformation into Stbl3 bacteria (Thermo Fisher Scientific). Plasmid

DNA was isolated from overnight bacterial cultures and verified via Sanger sequencing.

Generation of stable cell lines using CRISPR-Cas9: To produce 3[rd] generation lentiviral

stocks, HEK293T cells were transfected with a total of 22.5 µg total plasmid according

to the following equimolar ratios: 10 μg LentiCRISPR transfer plasmid (empty or

containing sgRNAs against either SERINC3 or SERINC5), 5.9 μg pMDLg/pRRE, 2.8 μg

pxRSV-Rev and 3.8 μg pMD2.G. Forty-eight hours later, concentrated lentivirus-

containing supernatant was harvested following low-speed centrifugation, filtration (0.45

μm), and mixture with Lenti-X concentrator (Takara Bio) according to the manufacturer's

instructions. Briefly, 1 volume Lenti-X concentrator was mixed with 3 volumes clarified

supernatant. The mixture was incubated for 30 minutes at 4°C, centrifuged at 1,500 x g

for 45 minutes at 4°C, resuspended in 1 ml complete DMEM media and immediately

stored at -80°C in single-use aliquots (100 μL).

We used previously validated sgRNAs to edit SERINC3 (34), whereas each of the five

sgRNAs targeting SERINC5 were screened and the sgRNA which caused the most

efficient editing in bulk transduced cells was chosen (data not shown). To create CEM

cells knocked out for SERINC3 (S3-KO), we spinoculated 1×10^6 cells with 100 μL

lentivirus (LentiCRISPRv2-SERINC3 ; (34)) at 1,200 x g for 2 hours at 25°C. Puromycin

(1 μg/ml) was added to cell cultures 72 hours post transduction to select for positive

clones. Two weeks post-selection, genomic DNA was isolated from mock or

lentiCRISPRv2-SERINC3 transduced CEM cells using the DNeasy Blood and Tissue Kit

(Qiagen) according to the manufacturer's instructions. Genome editing was assessed by

Tracking of Indels by Decomposition (TIDE; (52)). PCR amplicons encompassing exon

2 of SERINC3 were produced using Taq 2x Master Mix (NEB) and the following primers:

5'-CAAATTACAACCAACTTGATTAACAACGACG-3'   and   5'-

CTATAAAGCCTGATTTGCCTCGCTTTCTCTTC-3'. Clonal cell lines were

isolated from bulk edited cultures using single-cell dilutions in a 96-well plate, followed

by genomic DNA isolation and PCR amplification. Genome editing was verified via TIDE

analysis.

To generate either single knockout or double-knockout cells lacking SERINC5, CEM WT

or CEM S3-KO cells were spinoculated as described above with either empty lentivirus

(LentiCRISPR-GFP) or virus containing sgRNA-SERINC5(4). Isolation of genomic DNA,

PCR amplification and TIDE analysis were carried out seventy-two hours post

transduction in a similar manner to the generation of S3-KO cells. We then expanded

cells, which yielded two single SERINC5 knockout (S5-KO) clonal lines, and one

SERINC3/SERINC5 knockout clonal cell line. We named these lines: SERINC5

knockout clone 8 (S5-KO (8)), SERINC5 knockout clone 11 (S5-KO (11)) and

SERINC3/SERINC5 knockout clone 9 (S3/S5-KO (9)). The following primers were used

to generate PCR amplicons for TIDE   analysis: 5'-

AGTGCCTGGCCATGTTTCTT-3' and   5'-CATAGAGCAGGCTTCAGGAA-3'.

HIV-1 production and titer: To produce replication-competent viruses, HEK293T cells (4

x 10^6 /10 cm plate) were transfected with 24 µg of an infectious molecular clone of HIV-

1 (NL4-3) or a mutant lacking the *nef* gene (NL4-3∆Nef) using Lipofectamine 2000

reagent according to the manufacturer's instructions (Thermo Fisher Scientific). Virus-

containing supernatants were collected forty-eight hours post transfection, clarified by

low-speed centrifugation, and stored at -80°C. Viral titers were measured by infecting

HeLa P4.R5 cells with diluted viral stocks in duplicate in a 48-well format for 48 hours.

The cells were then fixed (1% formaldehyde; 0.2% glutaraldehyde) for 5 minutes at

room-temperature, followed by overnight staining at 37°C with a solution composed of 4

mM potassium ferrocyanide, 4mM potassium ferricyanide, 2 mM $MgCl_2$ and 0.4 mg/ml

X-gal (5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside). Infected cells (expressed as

infectious centers (IC)) were quantified using a computer image-based method (53).

(ABL Bioscience).

HIV-1 infection and replication: To conduct HIV-1 replication studies, 1 x 10^6 CEM cells

(wildtype (WT); SERINC5 knockout clone 8 (S5KO (8)); SERINC5 knockout clone 11

(S5KO (11)); SERINC3/SERINC5 knockout clone 9 (S3/S5 K.O. (9))) were infected with

NL4-3 (hereafter termed Nef+) or NL4-3ΔNef (hereafter termed Nef-) at a multiplicity of

infection (MOI) of 0.01 overnight at 37°C in a 24-well plate format. The cells were then

washed 3 times with 1 ml PBS (Corning), resuspended in 4 ml complete growth media

(RPMI 1640), transferred to T25 labeled flasks and incubated at 37°C in an "upright"

position for the duration of the experiment. Every three days, cultures were split 1:4 (1

ml cells; 3 ml media) and an aliquot (1 ml viral supernatant) stored at -80°C for

quantification of HIV-1 replication (p24 antigen) by ELISA.

Measurement of HIV-1 infectivity: Viral infectivity was quantified from virions produced in

CEM cells infected with NL4-3 or NL4-3ΔNef at day 12 post-infection. A 20% sucrose

cushion was used to concentrate virions via centrifugation according to the following

parameters: 23,500 x g; 1 hour at 4°C. Viral pellets were resuspended in culture

medium and dilutions used to infect 1.25 x 10^4 HeLa TZM-bl cells in triplicate in a 96-

well format. Forty-eight hours later, the culture medium was removed, and the cells

were lysed using a luciferase reporter gene assay reagent (Britelite, Perkin Elmer).

Infectivity (luciferase activity) was measured using a luminometer with data expressed

as relative light units (RLU). These values were normalized to the p24 concentration of

each sample and shown as RLU/p24.

RT-qPCR and analysis: Total cellular RNA was extracted from 1 x 10^6 JTAg WT, JTAg

S3/S5 KO, CEM WT, CEM S5KO (8), CEM S5KO (11), and CEM S3/S5 KO cells using

a Quick-RNA miniprep kit (Zymo Research), followed by treatment with RNase-free

DNAse I (Zymo Research). Complementary DNA (cDNA) was generated from 250 ng of

all extracted RNA samples using M-MLV RT (Thermo Fisher Scientific) and treated with

RNaseOUT (Thermo Fisher Scientific). cDNA was mixed with the respective primer

pairs and SyGreen Blue Mix (PCR Biosystems) following the manufacturer's protocol in

biological triplicate and performed using a LightCycler 96 real-time PCR machine

(Roche). Quantification cycle values were normalized to a reference gene (*GAPDH*)

and relative *SERINC5* or *SERINC3* gene expression ratios were calculated using the $2^{-\Delta\Delta Ct}$

method (54). The following primers were used for analysis: *SERINC5:* 5'-

ATCGAGTTCTGACGCTCTGC-3' and 5'-GCTCTTCAGTGTCCTCTCCAC-3'; *SERINC3*

5'-AATTCAGGAACACCAGCCTC-3' and 5'- GGTTGGGATTGCAGGAACGA-3';

GAPDH 5'-TGCACCACCAACTGCTTAGC-3' and 5'-GGCATGGACTGTGGTCATGAG-

3'.

Data analysis and presentation: Datasets were analyzed and combined in Microsoft

Excel and GraphPad Prism 8.0 software. Where indicated, two-tailed unpaired or paired

t-tests were performed. We utilized Adobe Photoshop and Illustrator CS6 for figure

production.

Results

Generation of CEM cells lacking SERINC3 and SERINC5

Results

Generation of CEM cells lacking SERINC3 and SERINC5

We utilized CRISPR/Cas9 gene editing to determine whether the modulation of

SERINC3 and SERINC5 is required for Nef to enhance HIV-1 replication in CEM cells.

We chose the CEM cell line due to its ability to support robust Nef-dependent HIV-1

replication and virion-infectivity (23). We hypothesized that Nef would remain an

important factor in promoting viral spread whether or not CEM cells expressed

SERINC3 and SERINC5, a result in conflict with data obtained using Jurkat T cells but

consistent with data recently obtained using MOLT-3 T cells. To test this, we created

clonal cell lines lacking either SERINC5 or both SERINC3 and SERINC5 (described in

Methods). We identified three clones containing indels within the SERINC5 target

region: "clone 8" (S5-KO (8)), containing 4 and 13 base pair deletions, "clone 11",

containing 2, 11, and 13 base pair deletions (S5/KO (11)) and "clone 9", containing 10

and 13 base pair deletions (S3/S5KO (9)) ; Figure 1A). SERINC3 knockout cells

consisted of indels bearing a single base pair insertion and a single base pair deletion

(S3/S5 KO (9); Figure 1C).

We attempted to validate editing in these cell lines using a monoclonal antibody

targeting the extracellular domain of SERINC (55). However, this antibody did not detect

endogenous SERINC5 in either Jurkat or CEM cells (data not shown). Instead, we

reasoned that CRISPR/Cas9 editing of *serinc3* and *serinc5* would lead to a quantifiable

decrease in mRNA transcripts due to nonsense mediated decay (NMD; (56)). To test

this, we isolated RNA from CEM WT, CEM S5-KO (8), CEM S5-KO (11), CEM S3/S5-

KO (9) and, as controls, JTAg WT and JTAg S3/S5-KO cells. In both JTAg and CEM,

knockout clones expressed less *serinc5* (Figure 1B) or *serinc3* (Figure 1D) compared to

wildtype clones. Given these results, these CEM cell lines served as the basis for our

viral replication and infectivity studies.

Optimal HIV-1 spread and viral infectivity in CEM cells is dependent on Nef but

independent of SERINC5.

To test whether Nef is required to counteract SERINC5 to enhance HIV-1 replication,

we infected CEM wildtype (WT), S5-KO (8) and S5-KO (11) cells with either Nef

expressing (hereafter termed Nef+: NL4-3) or Nef lacking (hereafter termed Nef-: NL4-

3ΔNef) HIV-1 viruses at a multiplicity of infection (MOI)=0.01 infectious units per cell.

The inocula for these growth-rate experiments were produced from HEK293T cells

transfected with proviral plasmids, and the MOI was based on the infectivity of the virus

stocks measured as infectious centers in cultures of CD4-positive HeLa-P4.R5 indicator

cells (53). To infect 1×10^6 CEM cells, we used amounts of Nef+ or Nef- virus stocks that

yielded 10,000 infectious centers in the HeLa indicator assay. Although the infectivity of

the viruses to CEM cells might be different than to CD4-HeLa cells, we chose this

approach rather than normalizing the inocula to the content of p24 capsid antigen to

adjust for the reduced infectivity of Nef- virions produced by the HEK293T cells (data

not shown). The CEM cell cultures were split every 3 days and viral replication

quantified by the amount of viral capsid (p24) within the supernatant (Figure 2A)

measured by ELISA. The Nef+ viruses propagated more rapidly than the Nef- viruses,

accumulating around 8.5-fold more p24 antigen in the culture supernates at 12 days

post-infection in CEM WT cells (Figure 2B: Left panel). This corroborated the

importance of Nef in enhancing viral replication in this *in vitro* system. If Nef-mediated

320 modulation of SERINC5 were important for this phenotype, then the attenuated

321 replication of Nef- virus should be "rescued" in cells lacking SERINC5. Instead, Nef

322 enhanced HIV-1 replication in the CEM S5KO (8) and CEM S5KO (11) cell lines (Figure

323 2B: middle and right panels).

324  We next asked whether counteraction of SERINCs is necessary for Nef to

325 enhance the infectiousness of virions produced by CEM cells, since Nef enhances

326 virion-infectivity in a SERINC- and cell-type dependent manner (34). To test this, we

327 collected virions from infected CEM cells at day 12 post-infection and measured single-

328 cycle infectivity using HeLa TZM-bl reporter cells, which express luciferase under the

329 transcriptional control of the HIV-1 LTR (Figure 2C). We used these cells because the

330 luciferase read-out is more sensitive that the infectious center read-out of the HeLa-

331 P4.R5 cells, and the concentration of Nef- virus produced by the CEM cultures was

332 relatively low. Virions produced at day 12 post infection by Nef+ virus were around 8-

333 fold more infectious per amount of p24 capsid antigen than those produced by Nef-

334 virus, and this trend was observed regardless of SERINC5 expression (Figure 2D).

335 Taken together, these data indicate that modulation of SERINC5 is not the primary

336 mechanism by which Nef increases viral spread or infectivity in CEM cells.

338 *Nef enhances HIV-1 spread and virion-infectivity independently of SERINC3 in CEM*

339 *cells.*

340 Whereas CEM cells express slightly less *serinc5* RNA than Jurkats, they express

341 markedly more *serinc3* RNA (Figure 1, B and D). Thus, we sought to determine

342 whether Nef-mediated modulation of SERINC3 influences viral growth rate and

infectivity in these cells and might explain the *nef*-phenotype. We infected CEM WT and

CEM cells lacking both SERINC3 and SERINC5 (S3/SKO (9)) with Nef+ and Nef –

viruses and measured viral replication and infectivity in a similar manner as shown in

Figure 2A and 2C. The absence of SERINC3 in CEM cells did not "rescue" Nef- virus in

terms of either viral replication rate (Figure 3A, compare left and right panels) or virion

infectivity (Figure 3B). Altogether, this data indicates that Nef increases virion infectivity

as well as growth-rate in CEM T cells independently of SERINC3 and SERINC5.

Discussion

In this study, we sought to determine whether Nef's ability to enhance viral replication

and/or infectivity in CEM T cells is primarily linked to modulation of SERINC3 and/or

SERINC5. We present evidence that argues against this by showing that Nef increases

viral replication rate in CEM cells lacking SERINC3 and SERINC5 and that virions of

Nef+ virus are more infectious than those of Nef- virus regardless of whether SERINC3

and SERINC5 are expressed in the CEM cells that produced them.

Our finding that Nef enhances virion-infectivity independently of SERINCs in CEM T

cells contrasts with reports in Jurkat and primary CD4-positive cells, where Nef

mediated enhancement of infectivity appears to correlate with SERINC5 expression (34,

35). On the other hand, our observations herein using CEM cells are similar to recent

results reported using MOLT-3 cells (46). In MOLT-3 cells, a chimeric HIV-1 virus

bearing the SERINC5 antagonist (glycoMA) failed to substitute for Nef in rescuing HIV-1

infectivity (a property glycoMA should have if the sole function of Nef were to counteract

SERINC5), and knockout of SERINC5 did not rescue the reduced growth rate of Nef-

virus (46). Together, these data suggest that the SERINC-dependence of the Nef

infectivity phenotype is cell-type dependent (34, 35, 46), with Jurkat being SERINC-

dependent and CEM and MOLT-3 cells being SERINC-independent. Comparing virions

produced from infected primary CD4-positive T cells either lacking or expressing

SERINC5 seems essential to adjudicate the relevance of observations made using

these T cell lines.

What mechanisms might explain Nef-mediated enhancement of infectivity and/or viral

replication independently of SERINCs? One potential explanation could be Nef-

mediated modulation of Src family kinases (SFKs). Nef binds several SFK members

such as Lck, Hck, Lyn, and c-Src through a conserved proline-rich (PxxP) motif

contained within Nef's

Src homology region 3 (SH3) binding domain (57, 58), and primary Nef isolates from

HIV-1 Group M display a conserved ability to activate SFK's (59). Nef mutants lacking

the PxxP motif were reportedly unable to enhance HIV-1 replication in peripheral blood

mononuclear cells (58). However, other studies reported that mutation of Nef's PxxP

motif yielded little to no difference in HIV-1 replication within MOLT-3 or primary CD4-

positive T cells, and in CEM T cells the attenuated phenotype of mutants of the SH3

binding domain was modest and seemed attributable to reduced expression of Nef (25,

46, 60). Analyzing HIV-1 replication in T cells lacking one or more SFKs may be

necessary to adequately assess whether Nef's ability to enhance HIV-1 replication is

mediated by these interactions.

One notable similarity between MOLT-3 and CEM lymphoblastoid cells that

distinguishes them from Jurkat cells is that they do not express the T cell receptor

(TCR) at their surfaces (61, 62). Given that Nef reportedly mimics TCR-signaling (22,

63), another possibility is that MOLT-3 and CEM cells reveal a SERINC-independent

growth-rate Nef phenotype that is exaggerated by the absence of constitutive TCR

signaling in these cells. This could be consistent with the initial observations that the

activation-state of primary CD4-positive T cells affects the Nef growth-rate phenotype

(24). Nonetheless, how TCR signaling would affect virion-infectivity is obscure.

Lastly, the explanation might reside in Nef's ability to interact with components of

clathrin-mediated trafficking pathways and to modulate many cellular membrane

proteins, one of which might be currently unidentified but underlie the infectivity

phenotype in MOLT-3 and CEM cells. Nef binds AP complexes via a conserved sorting

signal near its C-terminus: 160ExxxLL164,165 (64). This "di-leucine motif" is required

for both Nef-mediated CD4 downregulation and optimal viral infectivity in CEM cells

(65). The Nef LL164/165AA mutant, which is unable to bind AP-2 (66), replicates poorly

in MOLT-3 cells (46). Residues located within Nef's core domain and required for

downregulating CD4 and interacting with Dynamin-2, a "pinchase" of clathrin coated

pits, are also required for enhancement of viral replication in MOLT-3 cells (46, 67, 68).

Finally, Nef enhances HIV-1 replication in the absence of CD4 downregulation in MOLT-

3 cells (46) and in CEM-derived cells (A2.01) engineered to express a CD4 lacking a

cytoplasmic domain that is unresponsive to Nef (69). These observations regarding CD4

are critically important, since CD4 is itself a potent inhibitor of infectivity that is

counteracted by Nef (11, 12). In the CEM experiments herein, CD4 downregulation by

Nef could conceivably account for the observed infectivity and growth-rate phenotypes.

Nonetheless, even if that were the case, the data would indicate that the contribution of

CD4 to these phenotypes far outweighs the contribution of SERINC3 and SERINC5 in

these cells.

Our study has several caveats. As noted above, CEM cells, like MOLT3 cells, might not

reflect the role of Nef in primary T cells or macrophages. Also, quantitative experimental

variation is evident in the CEM system, with the *nef* virion-infectivity phenotype varying

between 2- and 8-fold (compare Figures 2 and 3). Nonetheless, a *nef*-phenotype was

always apparent in these cells, and it was unaffected by knock-out of *serinc5* and

serinc3.

Overall, whether Nef provides a direct, positive effect on viral replication or instead is

counteracting a still unidentified restriction factor remains to be determined. Utilizing

both MOLT-3 and CEM cells might facilitate answering this question, providing a more

complete understanding of the enigmatic yet important virologic effects of Nef.

References

1. Kestler HW, Ringler DJ, Mori K, Panicali DL, Sehgal PK, Daniel MD, Desrosiers RC.
435
1991. Importance of the nef gene for maintenance of high virus loads and for
436
development of AIDS. Cell 65:651–662.
437

2. Kirchhoff F, Greenough TC, Brettler DB, Sullivan JL, Desrosiers RC. 1995. Absence
438
of Intact nef Sequences in a Long-Term Survivor with Nonprogressive HIV-1
439
Infection. New England Journal of Medicine 332:228–232.
440

3. Deacon NJ, Tsykin A, Solomon A, Smith K, Ludford-Menting M, Hooker DJ, McPhee
441
DA, Greenway AL, Ellett A, Chatfield C, Lawson VA, Crowe S, Maerz A, Sonza S,
442
Learmont J, Sullivan JS, Cunningham A, Dwyer D, Dowton D, Mills J. 1995.
443
Genomic structure of an attenuated quasi species of HIV-1 from a blood transfusion
444
donor and recipients. Science 270:988–991.
445

4. Kirchhoff F. 2010. Immune evasion and counteraction of restriction factors by HIV-1
446
and other primate lentiviruses. Cell Host Microbe 8:55–67.
447

5. Garcia JV, Alfano J, Miller AD. 1993. The negative effect of human
448
immunodeficiency virus type 1 Nef on cell surface CD4 expression is not species
449
specific and requires the cytoplasmic domain of CD4. J Virol 67:1511–1516.
450

6. Aiken C, Konner J, Landau NR, Lenburg ME, Trono D. 1994. Nef induces CD4
451
endocytosis: requirement for a critical dileucine motif in the membrane-proximal
452
CD4 cytoplasmic domain. Cell 76:853–864.
453

7. Chaudhuri R, Lindwasser OW, Smith WJ, Hurley JH, Bonifacino JS. 2007.

Downregulation of CD4 by human immunodeficiency virus type 1 Nef is dependent

on clathrin and involves direct interaction of Nef with the AP2 clathrin adaptor. J

Virol 81:3877–3890.

8. Kwon Y, Kaake RM, Echeverria I, Suarez M, Karimian Shamsabadi M, Stoneham C,

Ramirez PW, Kress J, Singh R, Sali A, Krogan N, Guatelli J, Jia X. 2020. Structural

basis of CD4 downregulation by HIV-1 Nef. Nat Struct Mol Biol

https://doi.org/10.1038/s41594-020-0463-z.

9. Wildum S, Schindler M, Münch J, Kirchhoff F. 2006. Contribution of Vpu, Env, and

Nef to CD4 down-modulation and resistance of human immunodeficiency virus type

1-infected T cells to superinfection. J Virol 80:8047–8059.

10. Alsahafi N, Ding S, Richard J, Markle T, Brassard N, Walker B, Lewis GK,

Kaufmann DE, Brockman MA, Finzi A. 2016. Nef Proteins from HIV-1 Elite

Controllers Are Inefficient at Preventing Antibody-Dependent Cellular Cytotoxicity. J

Virol 90:2993–3002.

11. Ross TM, Oran AE, Cullen BR. 1999. Inhibition of HIV-1 progeny virion release

by cell-surface CD4 is relieved by expression of the viral Nef protein. Curr Biol

9:613–621.

12. Lama J, Mangasarian A, Trono D. 1999. Cell-surface expression of CD4 reduces

HIV-1 infectivity by blocking Env incorporation in a Nef- and Vpu-inhibitable manner.

Curr Biol 9:622–631.

13.   Kasper MR, Roeth JF, Williams M, Filzen TM, Fleis RI, Collins KL. 2005. HIV-1

Nef disrupts antigen presentation early in the secretory pathway. J Biol Chem

280:12840–12848.

14.   Roeth JF, Williams M, Kasper MR, Filzen TM, Collins KL. 2004. HIV-1 Nef

disrupts MHC-I trafficking by recruiting AP-1 to the MHC-I cytoplasmic tail. J Cell

Biol 167:903–913.

15.   Noviello CM, Benichou S, Guatelli JC. 2008. Cooperative binding of the class I

major histocompatibility complex cytoplasmic domain and human immunodeficiency

virus type 1 Nef to the endosomal AP-1 complex via its mu subunit. J Virol 82:1249–

1258.

16.   Jia X, Singh R, Homann S, Yang H, Guatelli J, Xiong Y. 2012. Structural basis of

evasion of cellular adaptive immunity by HIV-1 Nef. Nat Struct Mol Biol 19:701–706.

17.   Hung C-H, Thomas L, Ruby CE, Atkins KM, Morris NP, Knight ZA, Scholz I,

Barklis E, Weinberg AD, Shokat KM, Thomas G. 2007. HIV-1 Nef assembles a Src

family kinase-ZAP-70/Syk-PI3K cascade to downregulate cell-surface MHC-I. Cell

Host Microbe 1:121–133.

18.   Blagoveshchenskaya AD, Thomas L, Feliciangeli SF, Hung CH, Thomas G.

2002. HIV-1 Nef downregulates MHC-I by a PACS-1- and PI3K-regulated ARF6

endocytic pathway. Cell 111:853–866.

19. Dikeakos JD, Atkins KM, Thomas L, Emert-Sedlak L, Byeon I-JL, Jung J, Ahn J,

Wortman MD, Kukull B, Saito M, Koizumi H, Williamson DM, Hiyoshi M, Barklis E,

Takiguchi M, Suzu S, Gronenborn AM, Smithgall TE, Thomas G. 2010. Small

molecule inhibition of HIV-1-induced MHC-I down-regulation identifies a temporally

regulated switch in Nef action. Mol Biol Cell 21:3279–3292.

20. Collins KL, Chen BK, Kalams SA, Walker BD, Baltimore D. 1998. HIV-1 Nef

protein protects infected primary cells against killing by cytotoxic T lymphocytes.

Nature 391:397–401.

21. Haller C, Rauch S, Fackler OT. 2007. HIV-1 Nef employs two distinct

mechanisms to modulate Lck subcellular localization and TCR induced actin

remodeling. PLoS ONE 2:e1212.

22. Simmons A, Aluvihare V, McMichael A. 2001. Nef Triggers a Transcriptional

Program in T Cells Imitating Single-Signal T Cell Activation and Inducing HIV

Virulence Mediators. Immunity 14:763–777.

23. Chowers MY, Spina CA, Kwoh TJ, Fitch NJ, Richman DD, Guatelli JC. 1994.

Optimal infectivity in vitro of human immunodeficiency virus type 1 requires an intact

nef gene. J Virol 68:2906–2914.

24. Spina CA, Kwoh TJ, Chowers MY, Guatelli JC, Richman DD. 1994. The

importance of nef in the induction of human immunodeficiency virus type 1

replication from primary quiescent CD4 lymphocytes. J Exp Med 179:115–123.

25.   Lundquist CA, Tobiume M, Zhou J, Unutmaz D, Aiken C. 2002. Nef-mediated

downregulation of CD4 enhances human immunodeficiency virus type 1 replication

in primary T lymphocytes. J Virol 76:4625–4633.

26.   Münch J, Rajan D, Schindler M, Specht A, Rücker E, Novembre FJ, Nerrienet E,

Müller-Trutwin MC, Peeters M, Hahn BH, Kirchhoff F. 2007. Nef-Mediated

Enhancement of Virion Infectivity and Stimulation of Viral Replication Are

Fundamental Properties of Primate Lentiviruses. JVI 81:13852–13864.

27.   Carl S, Greenough TC, Krumbiegel M, Greenberg M, Skowronski J, Sullivan JL,

Kirchhoff F. 2001. Modulation of different human immunodeficiency virus type 1 Nef

functions during progression to AIDS. J Virol 75:3657–3665.

28.   Crotti A, Neri F, Corti D, Ghezzi S, Heltai S, Baur A, Poli G, Santagostino E,

Vicenzi E. 2006. Nef Alleles from Human Immunodeficiency Virus Type 1-

InfectedLong-Term-Nonprogressor Hemophiliacs with or without Late Disease

Progression Are Defective in Enhancing Virus Replication and CD4 Down-

Regulation. JVI 80:10663–10674.

29.   Aiken C, Trono D. 1995. Nef stimulates human immunodeficiency virus type 1

proviral DNA synthesis. J Virol 69:5048–5056.

30.   Pandori MW, Fitch NJ, Craig HM, Richman DD, Spina CA, Guatelli JC. 1996.

Producer-cell modification of human immunodeficiency virus type 1: Nef is a virion

protein. Journal of virology 70:4283–4290.

31. Pizzato M, Helander A, Popova E, Calistri A, Zamborlini A, Palu G, Gottlinger

HG. 2007. Dynamin 2 is required for the enhancement of HIV-1 infectivity by Nef.

Proceedings of the National Academy of Sciences 104:6812–6817.

32. Pizzato M, Popova E, Göttlinger HG. 2008. Nef can enhance the infectivity of

receptor-pseudotyped human immunodeficiency virus type 1 particles. J Virol

82:10811–10819.

33. Pizzato M. 2010. MLV glycosylated-Gag is an infectivity factor that rescues Nef-

deficient HIV-1. Proc Natl Acad Sci USA 107:9364–9369.

34. Rosa A, Chande A, Ziglio S, De Sanctis V, Bertorelli R, Goh SL, McCauley SM,

Nowosielska A, Antonarakis SE, Luban J, Santoni FA, Pizzato M. 2015. HIV-1 Nef

promotes infection by excluding SERINC5 from virion incorporation. Nature

526:212–217.

35. Usami Y, Wu Y, Göttlinger HG. 2015. SERINC3 and SERINC5 restrict HIV-1

infectivity and are counteracted by Nef. Nature 526:218–223.

36. Shi J, Xiong R, Zhou T, Su P, Zhang X, Qiu X, Li H, Li S, Yu C, Wang B, Ding C,

Smithgall TE, Zheng Y-H. 2018. HIV-1 Nef Antagonizes SERINC5 Restriction by

Downregulation of SERINC5 via the Endosome/Lysosome System. J Virol 92.

37. Heigele A, Kmiec D, Regensburger K, Langer S, Peiffer L, Stürzel CM, Sauter D,

Peeters M, Pizzato M, Learn GH, Hahn BH, Kirchhoff F. 2016. The Potency of Nef-

Mediated SERINC5 Antagonism Correlates with the Prevalence of Primate

Lentiviruses in the Wild. Cell Host Microbe 20:381–391.

38.   Chande A, Cuccurullo EC, Rosa A, Ziglio S, Carpenter S, Pizzato M. 2016. S2

from equine infectious anemia virus is an infectivity factor which counteracts the

retroviral inhibitors SERINC5 and SERINC3. Proc Natl Acad Sci USA 113:13197–

13202.

39.   Beitari S, Ding S, Pan Q, Finzi A, Liang C. 2017. Effect of HIV-1 Env on

SERINC5 Antagonism. J Virol 91.

40.   Angerstein AO, Stoneham CA, Ramirez PW, Guatelli JC, Vollbrecht T. 2020.

Sensitivity to monoclonal antibody 447-52D and an open env trimer conformation

correlate poorly with inhibition of HIV-1 infectivity by SERINC5. Virology 548:73–81.

41.   Firrito C, Bertelli C, Vanzo T, Chande A, Pizzato M. 2018. SERINC5 as a New

Restriction Factor for Human Immunodeficiency Virus and Murine Leukemia Virus.

Annu Rev Virol 5:323–340.

42.   Inuzuka M, Hayakawa M, Ingi T. 2005. Serinc, an activity-regulated protein

family, incorporates serine into membrane lipid synthesis. J Biol Chem 280:35776–

35783.

43.   Trautz B, Wiedemann H, Lüchtenborg C, Pierini V, Kranich J, Glass B,

Kräusslich H-G, Brocker T, Pizzato M, Ruggieri A, Brügger B, Fackler OT. 2017. The

host-cell restriction factor SERINC5 restricts HIV-1 infectivity without altering the

lipid composition and organization of viral particles. J Biol Chem 292:13702–13713.

44.   Sood C, Marin M, Chande A, Pizzato M, Melikyan GB. 2017. SERINC5 protein

inhibits HIV-1 fusion pore formation by promoting functional inactivation of envelope

glycoproteins. J Biol Chem 292:6014–6026.

45.   Zhang X, Shi J, Qiu X, Chai Q, Frabutt DA, Schwartz RC, Zheng Y-H. 2019. CD4

expression and Env conformation are critical for HIV-1 restriction by SERINC5. J

Virol JVI.00544-19, jvi;JVI.00544-19v1.

46.   Wu Y, Olety B, Weiss ER, Popova E, Yamanaka H, Göttlinger H. 2019. Potent

Enhancement of HIV-1 Replication by Nef in the Absence of SERINC3 and

SERINC5. mBio 10:e01071-19, /mbio/10/3/mBio.01071-19.atom.

47.   Platt EJ, Wehrly K, Kuhmann SE, Chesebro B, Kabat D. 1998. Effects of CCR5

and CD4 cell surface concentrations on infections by macrophagetropic isolates of

human immunodeficiency virus type 1. J Virol 72:2855–2864.

48.   Derdeyn CA, Decker JM, Sfakianos JN, Wu X, O'Brien WA, Ratner L, Kappes

JC, Shaw GM, Hunter E. 2000. Sensitivity of human immunodeficiency virus type 1

to the fusion inhibitor T-20 is modulated by coreceptor specificity defined by the V3

loop of gp120. J Virol 74:8358–8367.

49.   Wallace JA, Kagele DA, Eiring AM, Kim CN, Hu R, Runtsch MC, Alexander M,

Huffaker TB, Lee S-H, Patel AB, Mosbruger TL, Voth WP, Rao DS, Miles RR,

Round JL, Deininger MW, O'Connell RM. 2017. miR-155 promotes FLT3-ITD–

induced myeloproliferative disease through inhibition of the interferon response.

Blood 129:3074–3086.

50.   Sanjana NE, Shalem O, Zhang F. 2014. Improved vectors and genome-wide

libraries for CRISPR screening. Nat Methods 11:783–784.

51.   Shalem O, Sanjana NE, Hartenian E, Shi X, Scott DA, Mikkelson T, Heckl D,

Ebert BL, Root DE, Doench JG, Zhang F. 2014. Genome-scale CRISPR-Cas9

knockout screening in human cells. Science 343:84–87.

52.   Brinkman EK, Chen T, Amendola M, van Steensel B. 2014. Easy quantitative

assessment of genome editing by sequence trace decomposition. Nucleic Acids Res

42:e168.

53.   Day JR, Martínez LE, Šášik R, Hitchin DL, Dueck ME, Richman DD, Guatelli JC.

2006. A computer-based, image-analysis method to quantify HIV-1 infection in a

single-cycle infectious center assay. Journal of Virological Methods 137:125–133.

54.   Schmittgen TD, Livak KJ. 2008. Analyzing real-time PCR data by the

comparative CT method. Nat Protoc 3:1101–1108.

55.   Molnar S, Wieczorek L, Zemil M, Schulte B, Martinez E, Gift S, Tang L, Streeck

H, Gramzinski RA, Michael NL, Joyce G, Polonis VR. 2020. Novel monoclonal

antibodies to the SERINC5 HIV-1 restriction factor detect endogenous and virion-

associated SERINC5. mAbs 12:1802187.

56.   Popp MW, Maquat LE. 2016. Leveraging Rules of Nonsense-Mediated mRNA

613   Decay for Genome Engineering and Personalized Medicine. Cell 165:1319–1322.

614   57.   Trible RP, Emert-Sedlak L, Smithgall TE. 2006. HIV-1 Nef Selectively Activates

615   Src Family Kinases Hck, Lyn, and c-Src through Direct SH3 Domain Interaction. J

616   Biol Chem 281:27029–27038.

617   58.   Saksela K, Cheng G, Baltimore D. 1995. Proline-rich (PxxP) motifs in HIV-1 Nef

618   bind to SH3 domains of a subset of Src kinases and are required for the enhanced

619   growth of Nef+ viruses but not for down-regulation of CD4. The EMBO Journal

620   14:484–491.

621   59.   Narute PS, Smithgall TE. 2012. Nef Alleles from All Major HIV-1 Clades Activate

622   Src-Family Kinases and Enhance HIV-1 Replication in an Inhibitor-Sensitive

623   Manner. PLoS ONE 7:e32561.

624   60.   Craig HM, Pandori MW, Riggs NL, Richman DD, Guatelli JC. 1999. Analysis of

625   the SH3-Binding Region of HIV-1 Nef: Partial Functional Defects Introduced by

626   Mutations in the Polyproline Helix and the Hydrophobic Pocket. Virology 262:55–63.

627   61.   Greenberg JM, Gonzalez-Sarmiento R, Arthur DC, Wilkowski CW, Streifel BJ,

628   Kersey JH. 1988. Immunophenotypic and cytogenetic analysis of Molt-3 and Molt-4:

629   human T-lymphoid cell lines with rearrangement of chromosome 7. Blood 72:1755–

630   1760.

62.   Royer HD, Ramarli D, Acuto O, Campen TJ, Reinherz EL. 1985. Genes encoding

the T-cell receptor alpha and beta subunits are transcribed in an ordered manner

during intrathymic ontogeny. Proc Natl Acad Sci USA 82:5510–5514.

63.   Len ACL, Starling S, Shivkumar M, Jolly C. 2017. HIV-1 Activates T Cell

Signaling Independently of Antigen to Drive Viral Spread. Cell Reports 18:1062–

1074.

64.   Sugden SM, Bego MG, Pham TNQ, Cohen ÉA. 2016. Remodeling of the Host

Cell Plasma Membrane by HIV-1 Nef and Vpu: A Strategy to Ensure Viral Fitness

and Persistence. Viruses 8:67.

65.   Craig HM, Pandori MW, Guatelli JC. 1998. Interaction of HIV-1 Nef with the

cellular dileucine-based sorting pathway is required for CD4 down-regulation and

optimal viral infectivity. Proc Natl Acad Sci USA 95:11229–11234.

66.   Bresnahan PA, Yonemoto W, Ferrell S, Williams-Herman D, Geleziunas R,

Greene WC. 1998. A dileucine motif in HIV-1 Nef acts as an internalization signal for

CD4 downregulation and binds the AP-1 clathrin adaptor. Curr Biol 8:1235–1238.

67.   Cohen GB, Rangan VS, Chen BK, Smith S, Baltimore D. 2000. The human

thioesterase II protein binds to a site on HIV-1 Nef critical for CD4 down-regulation.

J Biol Chem 275:23097–23105.

68.   Poe JA, Smithgall TE. 2009. HIV-1 Nef Dimerization Is Required for Nef-

Mediated Receptor Downregulation and Viral Replication. Journal of Molecular

Biology 394:329–342.

69.   Chowers MY, Pandori MW, Spina CA, Richman DD, Guatelli JC. 1995. The

growth advantage conferred by HIV-1 nef is determined at the level of viral DNA

formation and is independent of CD4 downregulation. Virology 212:451–457.

Author Contributions

Peter W. Ramirez: Conceptualization, Methodology, Investigation, Writing – original draft & review and editing **Francisco M. Acosta**, **Marissa Suarez**, **Aaron A. Angerstein**, **Thomas Vollbrecht**: Validation, Methodology, Investigation, Writing – review and editing **Jared Wallace, Ryan M. O' Connell**: Resources **John Guatelli:** Conceptualization, Methodology, Writing – review and editing.

Acknowledgments

We thank Dr. Vicente Planelles for the LentiCRISPR-GFP plasmid. We thank The Pendleton Charitable Trust for equipment. P.W.R. was partly supported by NIH grant K12GM068524. This work was supported by NIH grant R01 AI129706 to J.G.

Figure Legends

Figure 1: CRISPR/Cas9 editing of SERINC3 and SERINC5 in CEM cells. A.) *Left:* Chromatograms depicting a portion of exon 2 from SERINC5 wildtype (WT) or three SERINC5 knockout clones: clone 8 (S5KO (8)), clone 11 (S5KO (11)) or clone 9 (S3/S5 KO (9)). The guide RNA (gRNA) target site is shown as a blue arrow/annotation. Inserted nucleotides are highlighted red. Deletions are depicted as dashed line(s). *Right:* Tracking of Indels by Decomposition (TIDE) analysis of S5 KO clones. S5 KO (8) had 4 and 13bp deletions, S5KO (11) had 2,11 and 13 bp deletions and S3/S5 KO (9) had 10 and 13bp deletions. B.) Quantitative PCR (qPCR) showing the ratio of SERINC5 to GAPDH mRNA in JTAg and CEM cells expressing or lacking SERINC5. Results depict two independent experiments performed in triplicate. C.) *Left:* Chromatograms depicting a portion of exon 2 from SERINC3 wildtype (WT) or SERINC3/SERINC5 knockout clone 9 (S3/S5 KO (9)). The guide RNA (gRNA) target site is shown as a blue arrow/annotation. Inserted nucleotides are highlighted red. *Right:* TIDE analysis of S5 KO (9) predicted a 1bp insertion and 1bp deletion. D.) Quantitative PCR (qPCR) showing the ratio of SERINC3 to GAPDH mRNA in JTAg and CEM cells expressing or lacking SERINC3. Results depict two independent experiments performed in triplicate. Two-tailed unpaired t-tests were performed where indicated.

Figure 2: Nef enhances HIV-1 replication independently of SERINC5 in

CEM cells. A.) Schematic of viral replication studies performed in CEM wildtype (WT),

SERINC5 knockout clone 8 (S5KO (8)) or SERINC5 knockout clone 11 (S5KO (11))

cells. Each cell line was infected with either NL4-3 (termed Nef+) or HIV-1ΔNef (termed

Nef-) at an MOI of 0.01. The cultures were split every 3-4 days and viral growth

measured as indicated. Created with BioRender.com. B.) Viral replication quantified by

p24 Capsid ELISA in the supernatants of CEM WT, S5KO (8) and S5KO (11) cultures.

Results depict either two (WT, S5KO (8)) or one (S5KO (11)) independent infection

measured in duplicate at each time point. C.) Schematic depicting measurement of

single-cycle infectivity with virions produced from infected CEM WT, S5KO (8) or S5KO

(11) cultures. HeLa TZM-bl indicator cells contain a luciferase gene under the

transcriptional control of the HIV-1 LTR. Created with BioRender.com. D.) Infectivity

data (relative luciferase units (RLU) normalized to p24 (RLU/p24)) from virions collected

at day 12 post-infection from cultures of either CEM WT or S5KO (8) and S5KO (11)

cells. The virions were partially purified by centrifugation through a 20% sucrose

cushion before measuring infectivity (RLU in the HeLa TZM-bl assay) and p24

concentration (ELISA). Results are representative of one experiment performed in

quadruplicate. Two-tailed paired t-tests were performed where indicated.

Figure 3: Nef enhances HIV-1 replication independently of SERINC3 in

CEM cells. A.) Viral replication quantified by p24 Capsid ELISA in the supernatants of

CEM WT, S5KO (8) and S5KO (11) cultures. Results depict one independent infection

measured in quadruplicate at each time point. B.) Infectivity data (relative luciferase

units (RLU) normalized to p24 (RLU/p24)) from virions collected at Day 12 post-infection

from cultures of either CEM WT or S3/S5KO (9) cells. Results are representative of one

experiment performed in quadruplicate. Two-tailed paired t-tests were performed where

indicated.

Figure 1