Inhibition of p-glycoprotein does not increase the efficacy of proteasome inhibitors in multiple myeloma cells

Rachel L. Mynott[1] and Craig T. Wallington-Beddoe[1,2,3,4]

[1]College of Medicine and Public Health, Flinders University, Bedford Park SA, 5042, Australia;

[2]Flinders Medical Centre, Bedford Park SA, 5042, Australia;

[3]Centre for Cancer Biology, University of South Australia and SA Pathology, UniSA CRI Building, North Tce, Adelaide SA, 5000, Australia;

[4]Adelaide Medical School, University of Adelaide, Adelaide SA, 5000, Australia;

Corresponding author:

Dr Craig Wallington-Beddoe

Level 4, Flinders Centre for Innovation in Cancer

Flinders University

Bedford Park, SA 5042, Australia

Email: craig.wallingtonbeddoe@flinders.edu.au

Tel: +61 408 080 788;

Abstract

The aim of this study is to determine whether manipulation of the drug transporter P-glycoprotein improves the efficacy of proteasome inhibitors in multiple myeloma. P-glycoprotein is a well-known drug transporter that is associated with chemotherapy resistance in a number of cancers but its role in modulating the efficacy of proteasome inhibitors in multiple myeloma is not well understood. Research has shown that the second generation proteasome inhibitor carfilzomib is a substrate of P-glycoprotein and as such its efficacy may correlate with P-glycoprotein function. In contrast to carfilzomib, research concerning the first-in-class proteasome inhibitor bortezomib is inconsistent. Some researchers have demonstrated that inhibition of P-glycoprotein increases bortezomib cytotoxicity in multiple myeloma cells whereas others have shown no effect. Through the mining of publicly available gene expression microarrays of patient bone marrow, we show that P-glycoprotein gene expression increases with the disease stages leading to multiple myeloma. However, RNA-seq on LP-1 cells treated with bortezomib or carfilzomib demonstrated minimal basal P-glycoprotein expression which did not increase with treatment. Moreover, only one (KMS-18) of nine multiple myeloma cell lines expressed P-glycoprotein, including RPMI-8226 cells that are resistant to bortezomib or carfilzomib. We hypothesised that by inhibiting P-glycoprotein, multiple myeloma cell sensitivity to proteasome inhibitors would increase, thus providing a potential approach to improving responses and reversing resistance to these agents. However, the sensitivity of multiple myeloma cells lines to proteasome inhibition was not enhanced by inhibition of P-glycoprotein with the specific inhibitor tariquidar. In addition, targeting glucosylceramide synthase with eliglustat did not inhibit P-glycoprotein function and also did not improve proteasome inhibitor efficacy except at a high concentration. We conclude that P-glycoprotein

49 is poorly expressed in multiple myeloma cells, its inhibition does not enhance the efficacy of

50 proteasome inhibitors, and it is unlikely to be a useful avenue for further translational research.

Introduction

Multiple myeloma (MM) is an incurable malignancy of plasma cells (1). The malignant plasma

cells usually produce monoclonal immunoglobulin and the disease is characterised by one or

more of the four "CRAB" symptoms: high blood calcium, renal insufficiency, anaemia and

bone lytic lesions (1). The median age at diagnosis is 69 years, meaning comorbidities and sub-

optimal response to chemotherapies make the cancer difficult to treat in many patients (2). MM

is currently incurable and if patients do not immediately present with chemotherapy resistance,

they will eventually develop such resistance, contributing to a median overall survival of

approximately 7 years (1).

P-glycoprotein (P-gp) is a well-known drug transporter that has been shown to play a role in

drug resistance in a number of solid malignancies where it exports drugs, decreasing their

intracellular concentration and thus reducing their efficacy (3, 4). Its relevance in

chemotherapy resistance in MM was shown by Dalton *et. al.* where patients previously exposed

to chemotherapy had a higher expression of P-gp on their plasma cells compared to newly-

diagnosed patients (5). In the context of resistance to proteasome inhibitors, the most important

class of novel therapies to treat MM, there have been conflicting reports in the literature. The

second generation proteasome inhibitor carfilzomib is a known substrate of P-gp and its

efficacy was shown to be enhanced in cancer cells, including MM, when P-gp is inhibited (6-

9). In comparison, the efficacy of the first-in-class proteasome inhibitor bortezomib with P-gp

inhibition was either improved (10-12) or not affected (6, 13-15) in MM and other human

cancer cell lines. Improving the efficacy of proteasome inhibitors in MM and overcoming

resistance to these agents would be very beneficial to improve treatment options for high-P-gp

expressing patients.

The activity of P-gp can be inhibited directly, for example with the third-generation P-gp

inhibitor tariquidar that blocks the ATPase activity of P-gp (8). However, P-gp can also be

indirectly inhibited with research demonstrating that inhibition of the sphingolipid enzyme

glucosylceramide synthase (GCS) reduces both gene and protein expression of P-gp (16, 17).

GCS converts ceramide to glucosylceramide, the precursor of glycosphingolipids (GSLs) and

it has been found that these GSLs upregulate P-gp gene expression via the *cSrc* and β-catenin

signalling pathways in cancer cells (16). Eliglustat is a specific and potent inhibitor of GCS,

with an EC_{50} in the low nanomolar range, and is used to treat Gaucher's disease (18-20). As an

inhibitor of GCS, it reduces glucosylceramide levels in cells and thus GSLs, causing the same

reduction in P-gp expression as seen when GCS is either genetically or chemically inhibited

by RNA interference or D,L-Threo-1-phenyl-2-palmitoylamino-3-morpholino-1-propanol

(PPMP), respectively (21, 22). Therefore, eliglustat can be used to indirectly inhibit P-gp.

Here we report our negative findings, demonstrating P-gp does not appear to be expressed in

most MM cell lines, including those resistant to proteasome inhibitors, nor is its expression

increased with proteasome inhibitor treatment. Moreover, inhibition of P-gp in MM cell lines

does not enhance the efficacy of proteasome inhibitors and may therefore not be a useful

avenue for further translational research. Overall, these data support a lack of a role for P-pg

in modulating the effects of proteasome inhibitors in MM.

Materials and Methods

Cell lines

RPMI-8226 WT (ATC CCL-155) human MM cell line was obtained from the American Type Culture Collection (Manassas, VA). Resistant RMPI-8226 V10R (bortezomib resistant) and RPMI-8226 C10R (carfilzomib resistant) cell lines were a gift from Professor Stuart Pitson (University of South Australia). LP-1 (ACC 41) human MM cell line was purchased from the Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures GmbH (Braunschweig, Germany). KMS-11 (JCRB1179) human MM cell line was purchased from Cell Bank Australia (Sydney, Australia). MM.1S, MM.1R, NCI-H929 were kindly provided by Prof. Andrew Spencer (Monash University, Vic, Australia) and KMS-18 cells kindly provided by Prof. Junia Melo (SA Pathology, Australia). Cells were cultured in RPMI 1640 (Gibco, Waltham, MA) supplemented with 10% foetal bovine serum, 50 units/mL penicillin, 0.25 mg/mL streptomycin, 2 mM L-glutamate, 1 mM sodium pyruvate and 15 mM Hepes buffer (all Gibco). Cells were incubated at 37°C in 5% CO_2.

Drugs, reagents and antibodies

Bortezomib, carfilzomib and eliglustat were purchased from MedKoo (Morrisville, NC). Tariquidar was purchased from Selleck Chemicals (Houston, TX). Rhodamine 123 (Rho123), propidium iodide (PI) and cOmplete EDTA-free Protease Inhibitor Cocktail were purchased from Sigma Aldrich (St. Louis, MO). Tris UltraPure, hydrochloric acid, sodium chloride, glycerol, β-glycerophosphate and sodium fluoride were purchased from Chem Supply (Adelaide, Australia). FITC Annexin V apoptosis marker, Cat # 556419, AB_2665412 and 10X Annexin-V Binding Buffer were purchased from BD Biosciences (San Jose, CA). Precision Plus Protein All Blue and Kaleidoscope Pre-stained Protein Standards molecular

weight ladders were purchased from Bio-Rad (Hercules, CA). Vectashield mounting media

with 4′,6-diamidino-2-phenylindole (DAPI) was purchased from Vector Labs (Burlingame,

CA).

FITC Mouse Anti-Human P-glycoprotein (CD243) monoclonal antibody (clone 17F9, Cat #

557002, AB_396548) and FITC Mouse IgG2b κ Isotype Control monoclonal antibody (clone

27-35, Cat # 555742, AB_396085) were purchased from BD Biosciences. MDR1/ABCB1

rabbit monoclonal antibody (clone E1Y7B, Cat # 13342, AB_2631176) was purchased from

Cell Signaling (Beverly, MA). Anti-actin mouse monoclonal antibody (clone C4, Cat #

MAB1501, AB_2223041) was purchased from Millipore (Burlington, MA). Anti-alpha tubulin

mouse antibody loading control (clone DM1A, Cat # ab7291, AB_2241126) was purchased

from Abcam (Cambridge, United Kingdom). Pierce Goat Anti-Rabbit IgG (H+L) peroxidase

conjugated goat secondary antibody (Cat # 31460, AB_228341) and Pierce Goat Anti-Mouse

IgG (H+L) peroxidase conjugated goat secondary antibody (Cat # 31430, AB_228307) were

purchased from Invitrogen (Waltham, MA). Cy™5 AffiniPure Donkey Anti-Rabbit IgG

(H+L), AB_2340607 and FITC AffiniPure Donkey Anti-Mouse IgG (H+L), AB_2335588

polyclonal secondary antibodies were purchased from Jackson Laboratories (Bar Harbor, ME).

Cell viability by trypan blue

136   Trypan blue (1:2 dilution, Life Technologies, Waltham, MA) was mixed 1:1 with MM cell

137   suspension. 10 µL of this mixture was pipetted onto a haemocytometer slide and cells within

138   the central grid counted. Dead cells were observed as blue whereas live cells were unstained.

139   The concentration of cells was determined by the following equation: n x d x 10,000 = cells/mL

140   where n is the number of cells counted and d is the dilution factor (i.e. 2).

Analysis of DNA microarrays

Publicly available gene expression datasets were used to assess P-gp gene expression in CD138-positive bone marrow plasma cells obtained from healthy donors, monoclonal gammopathy of undetermined significance (MGUS), smouldering MM (SMM), newly-diagnosed MM and relapsed MM patients. The following datasets were used: GSE13591 (normal, n=5, MGUS, n=11, MM, n=142) (23), E-MTAB-363 (normal, n=5, MGUS, n=5, MM, n=155) (24) and GSE6477 (normal, n=15, MGUS, n=22, smouldering MM, n=24, new MM, n=73, relapsed MM, n=28) (25). Excel (Microsoft Corporation, Redmond, WA) was used to compare differences between two groups using the Student t test ($\log_2$ normally distributed data for GSE13591 and E-MTAB-363 or untransformed data GSE6477).

RNA sequencing and analysis

RNA extraction was undertaken using the RNeasy Mini Kit (QIAGEN, Hilden, Germany). Approximately 1 million cells (>80% viable) were treated with proteasome inhibitors for 6 and 16 hours before RNA was extracted following instructions provided by the kit. PolyA+ enriched RNAseq libraries were multiplexed and sequenced on the Illumina NextSeq 500 platform. Raw data, averaging 100 million reads per sample were analysed and quality checked using the FASTQC program (http://www.bioinformatics.babraham.ac.uk/projects/fastqc). Reads were mapped against the human (hg19) reference genomes for LP-1 cells using the STAR spliced alignment algorithm (26) (version2.7.2c with default parameters and -chimSeqmentMin 20, --quantMode GeneCounts) returning an average unique alignment rate of 80%. Differential expression analysis was evaluated from the TMM normalised gene counts using R (version 3.2.3) and edgeR (version 3.3) (27, 28) following protocols as described.

165 Alignments were visualised and interrogated using the Integrative Genomics Viewer v2.3.804.

166 Samples were processed and analysed the Australian Cancer Research Foundation Cancer

167 Genomics Facility (Adelaide, Australia).

Surface P-glycoprotein expression by flow cytometry

Approximately 100,000 – 500,000 MM cells were washed in 1X PBS to pellet. Cell pellets

were stained with 2 μL FITC P-gp antibody or matched isotype for 20 minutes in the dark at

room temperature. Cells were washed in PBS again to remove unbound antibody. Flow

cytometric analysis was undertaken, and data collected from 10,000 events on the CytoFLEX

S Flow Cytometer (Beckman Coulter, Brea, CA). CytExpert Software (Beckman Coulter) was

used for data collection and analysis.

Cell viability by flow cytometry

Approximately 100,000 – 500,000 MM cells were cultured with and without drugs of interest

for up to 72 hours. Cell suspensions were washed in 1X phosphate buffered saline (PBS) and

cell pellets stained with 2 μL FITC Annexin-V, 2 μL 1:50 PI and 96 μL 1X Annexin V Binding

Buffer for 10 minutes in the dark at room temperature. Cell viability was determined by

acquiring 10,000 events using a CytoFLEX S flow cytometer (Beckman Coulter) and analysis

of cell viability performed using CytExpert Software (Beckman Coulter). Viable cells are those

being dual Annexin-V FITC and PI negative.

P-glycoprotein function by Rhodamine 123 efflux

Approximately 100,000 – 500,000 MM cells were treated in tissue culture plates with proteasome inhibitors with and without tariquidar or eliglustat. After 22 hours, 200 ng/mL of the cell permeable P-gp substrate Rho123 was added to each well and the plates incubated at 37°C, 5% CO_2 for the remaining 2 hours. Cells were then washed twice in 1X PBS and analysed by flow cytometry. Mean fluorescence intensity (MFI) was recorded from 10,000 events on the CytoFLEX S Flow Cytometer (Beckman Coulter). The Rho123 dye is excited by the blue laser (488 nm) and its emission is detected using a 525 nm filter. CytExpert Software (Beckman Coulter) was used for data collection and analysis and data were normalised to unstained, untreated controls.

Western blotting

3 to 5 million cells were cultured with drugs of interest and then lysed in 10 mM Tris/HCl (pH 7.4), 137 mM NaCl containing 10% glycerol, 1% NP40, 10 mM β-glycerophosphate, 2 mM sodium fluoride and cOmplete EDTA-free protease inhibitor cocktail. This remained at 4°C for 10 minutes before centrifugation at 14,000 x g to clarify the lysate. Equal amounts of protein were loaded onto a Mini-PROTEAN TGX Precast Gel (Bio-Rad) and the Trans-Blot Turbo Transfer System was used to transfer protein onto a nitrocellulose membrane (Bio-Rad). Membranes were then blocked in 5% non-fat dairy milk powder in 50 mM Tris (pH 7.4), 154 mM NaCl and 1% Tween20 at room temperature for two hours. Primary and secondary antibodies were used as per the manufacturers' instructions and membranes were imaged on the ChemiDoc MP (Bio-Rad) after incubation in Clarity Western ECL Substrate (Bio-Rad). MDR1/ABCB1 and anti-actin primary antibodies were used at a dilution of 1:1000 and 1:5000,

209 respectively, and HRP-conjugated secondary antibodies were used at a dilution of 1:15,000.

210 Image Lab Software (Bio-Rad) was used for densitometric analysis of Western blot images.

Statistical Analysis

213 Comparisons between two groups were undertaken in Microsoft Excel using the Student t test

214 with a p-value of less than 0.05 considered statistically significant. Synergistic effects of drug

215 combinations were determined using the fractional product method (29) with a value of < -0.1

216 indicating synergy.

Results

Changes in P-gp gene expression with MM disease progression and in response to proteasome inhibitors in MM

Publicly available gene expression microarrays were used to analyse P-gp gene expression in

CD138-positive purified bone marrow plasma cells. In the three gene expression microarray

datasets analysed, newly-diagnosed and relapsed MM patients but not those patients with the

precursor conditions monoclonal gammopathy of undetermined significance (MGUS) and

smouldering MM (SMM) had higher expression of P-gp when compared to healthy controls

(Fig 1). Unlike previous findings (30), there was no statistically significant difference in the

expression of P-gp when comparing newly-diagnosed MM patients to patients whose MM had

relapsed (Fig 1C), however P-gp appeared to be more highly expressed in MM patients

compared to MGUS in two of the datasets (Fig 1A and C).

Fig 1. P-gp expression increases with disease progression in MM patients. Boxplots
(median and interquartile range) showing P-gp expression from the GSE13591 (A), E-MTAB-
363 (B) and GSE6477 (C) datasets. ns = not statistically significant, * p < 0.05. GSE13591
(normal, n=5, MGUS, n=11, MM, n=142), E-MTAB-363 (normal, n=5, MGUS, n=5, MM,
n=155) and GSE6477 (normal, n=15, MGUS, n=22, smouldering MM, n=24, new MM, n=73,
relapsed MM, n=28).

RNA-seq was undertaken to determine changes gene expression in LP-1 cells treated with 5
nM and 10 nM of bortezomib or carfilzomib for 6 hours. Further analysis was undertaken for
genes that had a count per million (CPM) of greater than 1 in at least two culture conditions.
Consequently, the data showed that P-gp was minimally expressed in this cell line (CPM < 1
for the control and proteasome inhibitor treatment groups) and so further normalisation and
analysis was not undertaken for this gene.

Basal P-gp protein expression in MM cell lines

Surface P-gp expression was analysed by flow cytometry. When compared to isotype, there
appeared to be no P-gp expression on the surface of any of the human MM cell lines tested (Fig
2A). However, analysis of total cell P-gp by Western blot indicated that KMS-18 cells did
express P-gp, whilst the other 8 MM cell lines examined, including the RPMI-8226
bortezomib/carfilzomib resistant cells, did not (Fig 2B). Due to the lack of P-gp in all MM cell
lines aside from KMS-18, the latter cell line was used for most subsequent experiments
exploring P-gp expression and function.

Fig 2. Basal P-gp in MM cell lines. A) Cell surface P-gp MFI of MM cell lines relative to isotype control. Data are mean ± standard deviation of three independent experiments (two independent experiments for RPMI-8226 WT, RPMI-8226 V10R, MM.1S, MM.1R and KMS-11). B) Western blot probed for total cell P-gp (170 kDa) in human MM cell lines. β-actin used as loading control. RPMI WT = RPMI-8226 WT, C10R = RPMI-8226 C10R, V10R = RPMI-8226 V10R and H929 = NCI-H929.

Changes in P-gp expression, function and cell viability in response to proteasome, P-gp and GCS inhibition

Proteasome inhibitors and changes in P-gp

LP-1 and KMS-18 MM cells cultured with bortezomib or carfilzomib for 24 hours did not appear to result in a clear increase in surface P-gp expression by flow cytometry (Fig 3A and C). Bortezomib appeared to have a greater cytotoxic effect than carfilzomib in both LP-1 and KMS-18 cells despite carfilzomib being shown to be more effective than bortezomib in MM patients (Fig 3B and D) (31). Despite not causing an increase in surface P-gp in KMS-18 cells, bortezomib did result in total P-gp expression increasing after a 16-hour treatment (Fig 3E). Carfilzomib caused a more gradual increase in total P-gp expression, requiring a higher concentration to double the level of P-gp expression compared to the control culture (Fig 3F).

Fig 3. P-gp surface and total expression, and cell viability for MM cell lines treated with proteasome inhibitors. KMS-18 cells were treated with bortezomib or carfilzomib for 24

hours and flow cytometric analysis was undertaken to determine surface P-gp expression (A)

and cell viability (B). LP-1 cells were treated with bortezomib or carfilzomib for 24 hours and

flow cytometric analysis was undertaken to determine surface P-gp expression (C) and cell

viability (D). Viability is represented as fold change relative to untreated control cultures.

KMS-18: Bortezomib data are mean ± standard deviation of duplicates from four biological

replicates for 0, 4 and 10 nM treatments and two technical replicates for the remaining

conditions. Carfilzomib data are mean ± standard deviation of duplicates of four biological

replicates for 0, 10 and 20 nM treatments and duplicates of two biological replicates for the

remaining conditions. LP-1: Bortezomib data are mean ± standard deviation of duplicates from

two biological replicates for 0, 4 and 10 nM treatments and two technical replicates for the

remaining treatments. Carfilzomib data are mean ± standard deviation of two technical

replicates. KMS-18 cells were treated with bortezomib (E) or carfilzomib (F) for 16 hours and

total P-gp expression was determined by Western blot. Associated densitometric analyses are

displayed below each blot with P-gp band intensity relative to respective β-actin loading

control.

Inhibiting P-gp directly with tariquidar

Bortezomib alone caused an increase in total P-gp expression in KMS-18 cells which was not

greatly altered by the addition of 1 μM tariquidar (Fig 4A and B), a concentration shown to

effectively inhibit P-gp in cancer cells (32). The combination of bortezomib with tariquidar did

not appear to cause a statistically significant increase in surface P-gp expression (Fig 4C).

When KMS-18 cells were cultured with bortezomib alone, Rho123 MFI decreased with

increasing concentrations of the drug, likely due to efflux of Rho123 from the cell. The addition

of tariquidar inhibited the function of P-gp as expected, causing an accumulation of Rho123 as

seen by Rho123 MFI remaining constant with increasing concentrations of bortezomib (Fig

4D). The addition of tariquidar did not appear to further sensitise cells to the cytotoxic effects

of bortezomib, as neither an additive nor synergistic increase in cell death was seen in KMS-

18 cells treated with the combination of the two drugs (Fig 4E).

Fig 4. Tariquidar functionally inhibits P-gp but does not synergise with bortezomib. A)

Western blot for KMS-18 cells treated with bortezomib and 0 or 1 µM tariquidar for 16 hours.

B) Densitometric analysis of the Western blot displayed in (A), P-gp band intensity relative to

respective β-actin loading control. KMS-18 cells were treated with bortezomib and 0, 1 or 5

µM tariquidar for 24 hours. Surface P-gp expression relative to isotype (C), Rho123 MFI

normalised to unstained and relative to untreated (first data point of blue line) controls (D) and

viability relative to untreated control (E) were analysed by flow cytometry. Flow cytometry

data are mean ± standard deviation from two independent experiments. ** $p < 0.01$.

Carfilzomib alone did not appear to greatly change total P-gp expression in KMS-18 cells at

the concentrations tested, however, the addition of 1 µM tariquidar caused a slight decrease

(Fig 5A and B). On the cell surface, 5 µM tariquidar resulted in a small increase in P-gp

expression when combined with 20 nM carfilzomib (Fig 5C). When KMS-18 cells were

cultured with carfilzomib alone, the Rho123 MFI decreased with increasing concentrations of

the drug, suggesting an increase in efflux of the dye. The treatments combining carfilzomib

with tariquidar resulted in a much smaller decrease in Rho123 MFI, suggesting the addition of

tariquidar causes an accumulation of dye in the cells when compared to carfilzomib alone,

consistent with functional inhibition of P-gp by tariquidar (Fig 5D). As with bortezomib, the

addition of tariquidar did not appear to further sensitise cells to the cytotoxic effects of

carfilzomib, as neither an additive nor synergistic reduction in cell viability was observed in

cells treated with the combination of the two drugs (Fig 5E).

Fig 5. Tariquidar functionally inhibits P-gp but does not synergise with carfilzomib. A)

Western blot for KMS-18 cells treated with carfilzomib and 0 or 1 μM tariquidar for 16 hours.

B) Densitometric analysis of Western blot displayed in (A), P-gp band intensity relative to

respective β-actin loading control. KMS-18 cells were treated with carfilzomib and 0, 1 or 5

μM tariquidar for 24 hours. Surface P-gp expression relative to isotype (C), Rho123 MFI

normalised to unstained and relative to untreated (first data point of blue line) controls (D) and

viability relative to untreated (E) were analysed by flow cytometry. Flow cytometry data are

mean ± standard deviation of two independent experiments. ** $p < 0.01$.

Inhibiting glucosylceramide synthase

The potent and specific GCS inhibitor eliglustat alone did not alter total P-gp expression in

KMS-18 cells, however, 30 μM eliglustat, the highest concentration examined, augmented the

increase in P-gp induced by bortezomib (Fig 6A and B). Intriguingly, this combination did not

result in increased surface P-gp expression (Fig 6C). P-gp function as indicated by Rho123 cell

efflux did not appear to be affected by eliglustat treatment as the MFI of the dye decreased to

the same extent when comparing bortezomib alone to bortezomib plus eliglustat treatment (Fig

6D). These data agree with previous findings demonstrating GCS downregulation by antisense

transfection did not influence Rho123 efflux in an adriamycin-resistant breast adenocarcinoma

cell line with high P-gp expression (33). The addition of eliglustat did, however, further

sensitise KMS-18 cells to bortezomib, as a synergistic reduction in cell viability was seen with

bortezomib and 30 μM eliglustat (Fig 6E) (fractional product -0.314 and -0.216 for 4 nM and

10 nM bortezomib, respectively), despite eliglustat being shown to inhibit GCS at

concentrations in the low nanomolar range (19, 20).

Fig 6. Eliglustat does not inhibit P-gp but synergises with bortezomib. A) Western blot of

KMS-18 cells treated with bortezomib and 0, 40 nM or 30 μM eliglustat for 16 hours. B)

Densitometric analysis of the Western blot displayed in (A), P-gp band intensity relative to

respective β-actin loading control. KMS-18 cells were treated with bortezomib and 0, 40 nM

or 30 μM eliglustat for 24 hours. Surface P-gp expression relative to isotype (C), Rho123 MFI

normalised to unstained and relative to untreated (first data point of blue line) controls (D) and

viability relative to untreated control (E) were analysed by flow cytometry. Flow cytometry

data are mean ± standard deviation of duplicate measurements and are representative of two

independent experiments. ** $p < 0.01$.

The addition of 30 μM eliglustat to carfilzomib treatment also produced an increase in total P-

gp (Fig 7A and B), and to a lesser extent surface P-gp (Fig 7C). Rho123 efflux and thus P-gp

function did not appear to be affected by eliglustat treatment as the MFI of the dye decreased

to the same extent when comparing carfilzomib alone to carfilzomib plus eliglustat treatment

(Fig 7D). This correlates with the bortezomib and eliglustat data (Fig 6), suggesting inhibition

of GCS by eliglustat does not inhibit P-gp function. The addition of eliglustat did, however,

appear to further sensitise the cells to carfilzomib treatment, as a synergistic reduction in cell

viability was seen with carfilzomib and 30 μM eliglustat (Fig 7E) (fractional product -0.446

and -0.390 for 10 nM and 20 nM carfilzomib, respectively).

Fig 7. Eliglustat does not inhibit P-gp but synergises with carfilzomib. A) Western blot for

KMS-18 cells treated with carfilzomib and 0, 40 nM or 30 μM eliglustat for 16 hours. B)

Densitometric analysis of the Western blot displayed in (A), P-gp band intensity relative to

respective β-actin loading control. KMS-18 cells were treated with carfilzomib and 0, 40 nM

or 30 μM eliglustat for 24 hours. Surface P-gp expression relative to isotype (C), Rho123 MFI

normalised to unstained and relative to untreated (first data point of blue line) controls (D) and

viability relative to untreated control (E) were analysed by flow cytometry. Flow cytometry

data are mean ± standard deviation of duplicate measurements and are representative of two

independent experiments. *, $p < 0.05$, ** $p < 0.01$.

As a comparison to KMS-18 cells, LP-1 cells do not express P-gp and unlike KMS-18 cells,

bortezomib treatment does not upregulate P-gp protein expression by Western blot (Fig 8A).

Rho123 MFI remains constant with bortezomib alone or in combination with eliglustat,

suggesting the dye is being retained in the cell, likely due to the absence of P-gp (Fig 8B).

However, LP-1 cells appear to be more sensitive to eliglustat treatment alone at 30 μM

compared to KMS-18, as a 20% reduction in viability in LP-1 cells was observed (Fig 8C),

compared to no significant change in viability in KMS-18 cells (Figs 6E and 7E). Moreover, a

synergistic reduction in cell viability with the combination of drugs was not evident as was

seen in KMS-18 cells which express P-gp (Fig 8C).

Fig 8. LP-1 cells do not express P-glycoprotein. A) Western blot for LP-1 cells treated with

bortezomib and 0, 40 nM or 30 μM eliglustat for 16 hours, β-actin used as a loading control.

LP-1 cells were treated with bortezomib and 0, 40 nM or 30 μM eliglustat for 24 hours and the

Rho123 MFI normalised to unstained and relative to untreated (first data point of blue line)

controls (B) and viability relative to untreated control (C) were analysed by flow cytometry.

Flow cytometry data are mean ± standard deviation of two technical replicates.

397 **Discussion**

Despite the most well-known and researched function of P-gp being that of drug efflux when

located on the plasma membrane, it has also been shown to play a role in drug resistance when

situated in intracellular organelles such as lysosomes, mitochondria and nuclei (34-36). This

may explain the strong presence of P-gp in KMS-18 whole cell lysates by Western blot whilst

flow cytometric analysis demonstrated that it was not expressed to a great extent on the cell

surface. Despite these cells not expressing high levels of P-gp on the plasma membrane before

or after proteasome inhibitor treatment, they did show a decrease in Rho123 fluorescence

which is indicative of heightened P-gp function. Whilst this may be because the low level of

P-gp already on the surface is sufficient to pump out the dye, it could also be related to the

greater whole cell P-gp expression observed after proteasome inhibitor treatment and

catabolism of the dye in intracellular compartments such as lysosomes on which P-gp is present

(37). It has also been suggested that P-gp is regulated by post-translational modifications such

phosphorylation and assessing these modifications may be useful to determine whether they

have an effect on the activity of P-gp without changing the level of protein on the surface (38).

The other eight human MM cell lines tested, including bortezomib/carfilzomib resistant RPMI-

8226 cells, did not express P-gp on the cell surface or in whole cell lysates. Furthermore, its

expression was not induced by proteasome inhibition in LP-1 or H929 cells, indicated by a lack

of protein band at 170 kDa (H929 Western blot displayed in Fig S1). The Rho123 efflux studies

confirm this as the dye was retained in LP-1 cells treated with bortezomib, a response similar

to that observed for KMS-18 cells when P-gp was inhibited (Figs 4 and 5). Our RNA-seq data

also support this, with minimal P-gp gene expression found in LP-1 cells treated with

proteasome inhibitors, therefore this cell line and the remaining 7 MM cell lines were not used

in further P-gp experiments. These data support the findings that most MM cell lines have low

P-gp expression, making it difficult to test its contribution to proteasome inhibitor cellular

efflux (12). Unfortunately, the doxorubicin resistant MM cell line (RPMI-8226/Dox40) that

has been shown to have high P-gp expression could not be obtained and would be a useful

positive control for experiments assessing P-gp expression and function (11, 39).

Tariquidar is a functional inhibitor of P-gp, blocking its ATPase activity, and as expected it did

not directly affect P-gp expression (8). Tariquidar was shown to effectively inhibit P-gp in MM

cells at a concentration of 5 µM (8) and 1 µM in other cancers (32), so these concentrations

were chosen for further investigation in KMS-18 MM cells. Both 1 µM and 5 µM tariquidar

resulted in a decrease in P-gp function when combined with bortezomib, as shown by an

accumulation of Rho123 in the KMS-18 cells. Conversely, bortezomib treatment alone resulted

in an increase in P-gp activity in these cells. These data contradict a previous report where after

24 hours in culture with bortezomib, RPMI-8226/Dox40 MM cells had a reduction in P-gp

activity shown by an accumulation of Rho123 in the cells and a reduction of P-gp protein

expression by Western blot (11). Despite tariquidar inhibiting P-gp, it did not further sensitise

the cells to bortezomib. This is consistent with previous studies showing bortezomib is not a

substrate of P-gp (6, 13-15), however, contradicts research that showed inhibition of P-gp does

improve the efficacy of bortezomib in MM cell lines (11, 12).

Carfilzomib requires higher concentrations compared to bortezomib to produce an increase in

total cell P-gp expression by Western blot. Similar to bortezomib, it does not induce P-gp

expression on the cell surface by flow cytometry, however, Rho123 fluorescence and thus P-

gp function does increase with increasing concentrations of carfilzomib. The combination of 5

µM tariquidar with 20 nM carfilzomib did cause a very small increase in surface P-gp which

may be to compensate for P-gp inhibition. Despite the literature suggesting that carfilzomib is

a substrate of P-gp (6-9), inhibition of P-gp by tariquidar did not further sensitise KMS-18 cells

to carfilzomib treatment. Taken together, these data demonstrate that P-gp inhibition does not

improve the efficacy of bortezomib or carfilzomib, suggesting drug efflux by P-gp may not be

a mechanism underpinning proteasome inhibitor resistance in MM.

Despite another research group (8) showing 5 µM tariquidar was sufficient to further sensitise

MM.1S MM cells to 5 nM bortezomib and carfilzomib, this concentration of tariquidar did not

enhance proteasome inhibitor induced cell death in KMS-18 cells. Using a higher tariquidar

concentration and longer incubation period may be valuable to confirm these results in KMS-

18 MM cells. It is important to note that whilst no changes in cell viability were observed,

changes in cell cycle and proliferation were not examined. It is entirely possible that the

combination of tariquidar with a proteasome inhibitor mediates changes in cell proliferation

rather than cell viability and this could be assessed in the future.

A link between glucosylceramides and P-gp has been established but is not well understood.

Genetic or chemical inhibition of GCS which converts ceramide to glucosylceramide has been

shown to downregulate P-gp expression (17) and so these findings led us to use the potent and

specific GCS inhibitor eliglustat as a way to indirectly inhibit P-gp. Interestingly, eliglustat did

not functionally inhibit P-gp as there were no significant changes in Rho123 fluorescence

before or after combining bortezomib/carfilzomib with eliglustat, thereby not supporting the

hypothesis that GCS inhibition downregulates P-gp expression and/or activity, at least in MM

cells. Eliglustat did, however, produce synergistic cell death when combined with bortezomib

and carfilzomib but only at a high concentration (30 µM) despite only needing a concentration

in the low nanomolar range to inhibit GCS enzymatic activity (19, 20). This high concentration

of eliglustat and the fact that it did not inhibit the function of P-gp suggests off target effects

may be causing this synergistic reduction in cell viability. One research group showed that

downregulation of GCS by siRNA-mediated knockdown slowed the clearance of ceramide

which is pro-apoptotic, and so an accumulation of ceramide may be contributing to the increase

in cell death observed in cells treated with 30 µM eliglustat and a proteasome inhibitor (17).

Such a high concentration of eliglustat is likely not clinically achievable so these data suggest

eliglustat may not be an effective drug to further sensitise MM cells to proteasome inhibition.

P-gp plays a role in chemotherapy resistance in a number of cancers, however its role in

contributing to proteasome inhibitor efficacy in MM is less well studied with conflicting reports

in the literature. Our data presented herein suggests that P-gp is poorly expressed in MM cells,

including those resistant to proteasome inhibitors, and that inhibition of P-gp does not enhance

proteasome inhibitor efficacy. Thus, a strategy to either increase the efficacy of proteasome

inhibitors or reverse resistance to these agents in MM via inhibition of P-gp may not be worthy

of further exploration.

Acknowledgements

482 CWB was supported by a Sylvia and Charles Viertel Foundation Clinical Investigator Award

483 and a Royal Australasian College of Physicians Servier 'Barry Young' Research Establishment

484 Fellowship.

References

1.   Rajkumar SV, Kumar S. Multiple Myeloma: Diagnosis and Treatment. Mayo Clinic
Proceedings. 2016;91(1):101-19.
2.   Wallington-Beddoe CT, Sobieraj-Teague M, Kuss BJ, Pitson SM. Resistance to
proteasome inhibitors and other targeted therapies in myeloma. British Journal of
Haematology. 2018;182(1):11-28.

3.   Abraham J, Salama NN, Azab AK. The role of P-glycoprotein in drug resistance in
multiple myeloma. Leukemia & Lymphoma. 2015;56(1):26-33.
4.   Nigam SK. What do drug transporters really do? Nat Rev Drug Discov. 2015;14(1):29-
44.
5.   Dalton WS. Detection of multidrug resistance gene expression in multiple myeloma.
Leukemia. 1997;11(7):1166-9.
6.   Verbrugge SE, Assaraf YG, Dijkmans BAC, Scheffer GL, Al M, den Uyl D, et al.
Inactivating PSMB5 Mutations and P-Glycoprotein (Multidrug Resistance-Associated
Protein/ATP-Binding Cassette B1) Mediate Resistance to Proteasome Inhibitors: Ex Vivo
Efficacy of (Immuno)Proteasome Inhibitors in Mononuclear Blood Cells from Patients with
Rheumatoid Arthritis. Journal of Pharmacology and Experimental Therapeutics.
2012;341(1):174-82.
7.   Besse A, Stolze SC, Rasche L, Weinhold N, Morgan GJ, Kraus M, et al. Carfilzomib
resistance due to ABCB1/MDR1 overexpression is overcome by nelfinavir and lopinavir in
multiple myeloma. Leukemia. 2018;32:391.
8.   Muz B, Kusdono HD, Azab F, de la Puente P, Federico C, Fiala M, et al. Tariquidar
sensitizes multiple myeloma cells to proteasome inhibitors via reduction of hypoxia-induced
P-gp-mediated drug resistance. Leukemia & Lymphoma. 2017;58(12):2916-25.
9.   Ao L, Wu Y, Kim D, Jang ER, Kim K, Lee D-M, et al. Development of peptide-based
reversing agents for p-glycoprotein-mediated resistance to carfilzomib. Mol Pharm.
2012;9(8):2197-205.
10.   Rumpold H, Salvador C, Wolf AM, Tilg H, Gastl G, Wolf D. Knockdown of PgP
resensitizes leukemic cells to proteasome inhibitors. Biochemical and Biophysical Research
Communications. 2007;361(2):549-54.
11.   O'Connor R, Ooi MG, Meiller J, Jakubikova J, Klippel S, Delmore J, et al. The
interaction of bortezomib with multidrug transporters: implications for therapeutic applications
in advanced multiple myeloma and other neoplasias. Cancer chemotherapy and pharmacology.
2013;71(5):1357-68.
12.   Clemens J, Seckinger A, Hose D, Theile D, Longo M, Haefeli WE, et al. Cellular uptake
kinetics of bortezomib in relation to efficacy in myeloma cells and the influence of drug
transporters. Cancer chemotherapy and pharmacology. 2015;75(2):281-91.
13.   Lu S, Wang J. The resistance mechanisms of proteasome inhibitor bortezomib.
Biomarker Research. 2013;1.
14.   Minderman H, Zhou Y, O'Loughlin KL, Baer MR. Bortezomib activity and in vitro
interactions with anthracyclines and cytarabine in acute myeloid leukemia cells are
independent of multidrug resistance mechanisms and p53 status. Cancer chemotherapy and
pharmacology. 2007;60(2):245-55.
15.   de Wilt LHAM, Jansen G, Assaraf YG, van Meerloo J, Cloos J, Schimmer AD, et al.
Proteasome-based mechanisms of intrinsic and acquired bortezomib resistance in non-small
cell lung cancer. Biochemical Pharmacology. 2012;83(2):207-17.
16.   Liu Y-Y, Gupta V, Patwardhan GA, Bhinge K, Zhao Y, Bao J, et al. Glucosylceramide
synthase upregulates MDR1 expression in the regulation of cancer drug resistance through cSrc
and β-catenin signaling. Molecular Cancer. 2010;9(1):145.
17.   Gouazé V, Liu Y-Y, Prickett CS, Yu JY, Giuliano AE, Cabot MC. Glucosylceramide
Synthase Blockade Down-Regulates P-Glycoprotein and Resensitizes Multidrug-Resistant
Breast Cancer Cells to Anticancer Drugs. Cancer Research. 2005;65(9):3861-7.
18.   Scott LJ. Eliglustat: A Review in Gaucher Disease Type 1. Drugs. 2015;75(14):1669-
78.

19.   McEachern KA, Fung J, Komarnitsky S, Siegel CS, Chuang W-L, Hutto E, et al. A
specific and potent inhibitor of glucosylceramide synthase for substrate inhibition therapy of
Gaucher disease. Molecular Genetics and Metabolism. 2007;91(3):259-67.
20.   Shayman JA. Eliglustat tartrate, a prototypic glucosylceramide synthase inhibitor.
Expert Review of Endocrinology & Metabolism. 2013;8(6):491-504.
21.   Roh J-L, Kim EH, Park JY, Kim JW. Inhibition of Glucosylceramide Synthase
Sensitizes Head and Neck Cancer to Cisplatin. Molecular Cancer Therapeutics.
2015;14(8):1907-15.
22.   Xie P, Shen Y-F, Shi Y-P, Ge S-M, Gu Z-H, Wang J, et al. Overexpression of
glucosylceramide synthase in associated with multidrug resistance of leukemia cells. Leukemia
Research. 2008;32(3):475-80.
23.   Agnelli L, Mosca L, Fabris S, Lionetti M, Andronache A, Kwee I, et al. A SNP
microarray and FISH-based procedure to detect allelic imbalances in multiple myeloma: an
integrated genomics approach reveals a wide gene dosage effect. Genes, chromosomes &
cancer. 2009;48(7):603-14.
24.   Rème T, Hose D, Theillet C, Klein B. Modeling risk stratification in human cancer.
Bioinformatics. 2013;29(9):1149-57.
25.   Chng WJ, Kumar S, Vanwier S, Ahmann G, Price-Troska T, Henderson K, et al.
Molecular dissection of hyperdiploid multiple myeloma by gene expression profiling. Cancer
Res. 2007;67(7):2982-9.
26.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 2012;29(1):15-21.
27.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics.
2010;26(1):139-40.
28.   Lun AT, Chen Y, Smyth GK. It's DE-licious: A Recipe for Differential Expression
Analyses of RNA-seq Experiments Using Quasi-Likelihood Methods in edgeR. Methods Mol
Biol. 2016;1418:391-416.
29.   Webb JL. Effect of more than one inhibitor. Enzyme and metabolic Inhibitors.
1963;1:66-79,488-512.
30.   Grogan T, Spier C, Salmon S, Matzner M, Rybski J, Weinstein R, et al. P-glycoprotein
expression in human plasma cell myeloma: correlation with prior chemotherapy. Blood.
1993;81(2):490-5.
31.   Dimopoulos M, Siegel D, White DJ, Boccia R, Iskander KS, Yang Z, et al. Carfilzomib
vs bortezomib in patients with multiple myeloma and renal failure: a subgroup analysis of
ENDEAVOR. Blood. 2019;133(2):147.
32.   Weidner LD, Fung KL, Kannan P, Moen JK, Kumar JS, Mulder J, et al. Tariquidar Is
an Inhibitor and Not a Substrate of Human and Mouse P-glycoprotein. Drug Metab Dispos.
2016;44(2):275-82.
33.   Liu Y-Y, Han T-Y, Giuliano AE, Cabot MC. Ceramide glycosylation potentiates
cellular multidrug resistance. The FASEB Journal. 2001;15(3):719-30.
34.   Yamagishi T, Sahni S, Sharp DM, Arvind A, Jansson PJ, Richardson DR. P-
glycoprotein mediates drug resistance via a novel mechanism involving lysosomal
sequestration. J Biol Chem. 2013;288(44):31761-71.
35.   Solazzo M, Fantappie O, Lasagna N, Sassoli C, Nosi D, Mazzanti R. P-gp localization
in mitochondria and its functional characterization in multiple drug-resistant cell lines.
Experimental cell research. 2006;312(20):4070-8.
36.   Baldini N, Scotlandi K, Serra M, Shikita T, Zini N, Ognibene A, et al. Nuclear
immunolocalization of P-glycoprotein in multidrug-resistant cell lines showing similar
mechanisms of doxorubicin distribution. European journal of cell biology. 1995;68(3):226-39.

37.   Forster S, Thumser AE, Hood SR, Plant N. Characterization of Rhodamine-123 as a
Tracer Dye for Use In In vitro Drug Transport Assays. PLOS ONE. 2012;7(3):e33253.
38.   Callaghan R, Luk F, Bebawy M. Inhibition of the Multidrug Resistance P-
Glycoprotein: Time for a Change of Strategy? Drug Metabolism and Disposition.
2014;42(4):623.
39.   Hawley TS, Riz I, Yang W, Wakabayashi Y, DePalma L, Chang Y-T, et al.
Identification of an ABCB1 (P-glycoprotein)-positive carfilzomib-resistant myeloma
subpopulation by the pluripotent stem cell fluorescent dye CDy1. American Journal of
Hematology. 2013;88(4):265-72.

Supporting Information

601   **Fig S1. H929 MM cells do not express P-gp.** Western blot probed for P-gp (170 kDa) in H929

602 cells treated with bortezomib ± tariquidar for 16 hours, showing H929 cells do not express P-

603 gp.

604   **Fig S2. Raw Western blot images.**

Figure 1

A

B

| Cell line: | RPMI WT | C10R | V10R | MM.1S | MM.1R | LP-1 | KMS-11 | KMS-18 | H929 |

P-gp

β-actin

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8