**Molecular and clinicopathological characterization of a prognostic immune gene**

**signature associated with MGMT methylation in glioblastoma**

Liang Zhao[1*], Shurui Xuan[2*], Jiayue Zhang[1], Zhiyuan Liu [1], Yu Wang[1], Peng Zhao[1]

[1] Department of Neurosurgery, The First Affiliated Hospital of Nanjing Medical University,

Nanjing, Jiangsu, China

[2] Department of Respiratory and Critical Care Medicine, The First Affiliated Hospital of Nanjing

Medical University, Nanjing, Jiangsu, China

**Correspondence to:** Peng Zhao, Department of Neurosurgery, The First Affiliated Hospital of

Nanjing Medical University, Nanjing, Jiangsu, 210000, China**.** Email: zhaopeng@njmu.edu.cn

**Sources of support:** This work was supported by the National Natural Science Foundation of

China under [grant number 81673210]

**Conflicts of interest:** The authors have no conflicts of interest to declare.

**Abstract**

**Background:** O6-methylguanine-DNA methyltransferase (MGMT) methylation status affects tumor chemoresistance and the prognosis of glioblastoma (GBM) patients. We aimed to investigate the role of MGMT methylation in the regulation of GBM immunophenotype and discover an effective biomarker to improve prognosis prediction of GBM patients.

**Methods:** A total of 769 GBM patients with clinical information from five independent cohorts were enrolled in the present study. Samples from the Cancer Genome Atlas (TCGA) dataset were used as the training set, whereas transcriptome data from the Chinese Glioma Genome Atlas (CGGA) RNA-seq, CGGA microarray, GSE16011, and the Repository for Molecular Brain Neoplasia (REMBRANDT) cohort were used for validation. A series of bioinformatics approaches were carried out to construct a prognostic signature based on immune-related genes. The influence of the signature on immunosuppression and remodeling of the tumor microenvironment were comprehensively investigated. Then, the utility of this immune gene signature was analyzed by the development and evaluation of a nomogram.

**Results:** We found that MGMT unmethylation was closely associated with immune-related biological processes in GBM. Sixty-nine immune genes were more highly expressed in the MGMT unmethylated than the MGMT methylated group. An immune gene-based risk model was further established to divide patients into high and low-risk groups, and the prognostic value of this signature was validated in several GBM cohorts. Functional analyses manifested a universal up-regulation of immune-related pathways in the high-risk group as compared to the low-risk group. Furthermore, the risk score was highly correlated to the immune cell infiltration, immunosuppression, inflammatory activities, as well as the expression levels of

immune checkpoints. Finally, a nomogram was developed for clinical application.

**Conclusions:** MGMT methylation is strongly related to the immune responses in GBM. The

immune gene-based signature we identified may have potential implications in predicting the

prognosis of GBM patients and mechanisms underlying the role of MGMT methylation.

**Keywords:** Glioblastoma, MGMT methylation, gene signature, prognosis, immune

microenvironment

**Introduction**

Glioma is a type of central nervous system (CNS) neoplasms and accounts for the majority of

intracranial malignant tumors in adults. Glioblastoma (GBM), which is defined as the World

Health Organization (WHO) grade IV glioma, shows a highly aggressive, heterogeneous, and

lethal phenotype. Currently, the etiology of GBM is still largely unknown. Although surgical

resection combined with chemotherapy or radiotherapy has been widely used as a routine

clinical treatment, the prognosis of patients has even not improved significantly, with a median

overall survival fewer than 15 months (1). Due to the persistent proliferation of tumor cells and

penetration into the surrounding tissues, total tumor resection is seldom possible (2).

Alkylating agents, such as temozolomide and procarbazine, are applied as first-line

chemotherapeutic drugs (3). However, many GBM patients do not respond to alkylating agents

or develop chemo-resistance after a period of chemotherapy (4). Therefore, there is an urgent

need to explore the underlying mechanisms of gliomagenesis and develop reliable therapeutic

approaches.

The intracellular DNA repair enzyme $O^6$-methylguanine-DNA methyltransferase (MGMT)

catalyzes the DNA repair process by removing the alkylation of the $O^6$ position of guanine (5).

By generating methylguanine adducts, DNA-alkylating drugs can trigger base mismatching

during DNA replication, affecting the cell cycle and promoting the death of tumor cells (6-8). As

a most striking modification form, methylation of the MGMT promoter can induce a decrease of

MGMT expression and further improve the curative effect of chemotherapy with alkylating

agents (1). In the case of GBM, clinical trials revealed that patients with MGMT methylation

benefit more than those with unmethylated tumors when received combined

radiochemotherapy (9). Furthermore, patients without MGMT methylation have a shorter

overall survival time (10). So far, these underlying molecular mechanisms have not been

exhaustively described, especially about tumor immune microenvironment.

Recently, breakthroughs in tumor immunotherapy have revolutionized the treatment of several

cancer types (11). The specific intratumoral immune microenvironment can facilitate immune

escape, drive tumorigenesis, and promote the malignant progression of the lesion (12, 13).

Immune checkpoints play an intermediary role in the tumor-immune system dynamics, and

inhibitory immune checkpoints have been studied intensively as therapeutic targets for

multiple malignant tumors. In non-small cell lung cancer (NSCLC) and melanoma, the

application of immunosuppressive agents has produced remarkable anti-tumor effects (14, 15).

The discovery of lymphatic system vessels in the CNS offered new hope for developing novel

immunotherapeutic methods for GBM (16). The first large phase III clinical trial of nivolumab

combined with ipilimumab (PD-1 inhibitors) in recurrent GBM (NCT02017717) was launched in

2014. A recent clinical trial indicated that the neoadjuvant *PD-1* blockade could upregulate the

amount of T cells, promote the expression of interferon-γ-related genes, and further offer a

promising survival benefit in recurrent GBM (17). The previous study has demonstrated that

isocitrate dehydrogenase (IDH) mutation status was closely associated with immune response

in GBM, and a prognostic model based on IDH mutation was further developed to predict the

clinical outcomes of patients (18). However, the role of MGMT methylation status in regulating

tumor immunity in GBM remains to be discussed, and an MGMT methylation-related

biomarker should be exploited.

Here, we utilized data from a large number of GBM samples from multiple publicly available

datasets to comprehensively investigate the potential relationship between MGMT status and

the immunological tumor microenvironment. We hypothesized that the shorter survival of

patients with MGMT-unmethylated GBM is partly due to the pro-tumor immune response. More

importantly, we developed a robust prognostic model based on MGMT methylation status to

predict the clinical outcomes of GBM patients.

**Methods**

**GBM patient datasets**

We retrospectively collected a large-scale profile composed of five independent datasets

involving 769 GBM patients. The RNA sequencing (RNA-seq) gene expression data for 165

GBM samples were obtained from The Cancer Genome Atlas (TCGA) database. The

microarray and RNA-seq transcriptome data for 112 and 113 GBM patients were downloaded

from the Chinese Glioma Genome Atlas (CGGA) database (www.cgga.org.cn). The microarray

gene expression profile matrix files of GSE16011 (including 159 GBM samples) and the

Repository for Molecular Brain Neoplasia (REMBRANDT) database (including 220 GBM

samples) were downloaded from the Gene Expression Omnibus (GEO) database

(www.ncbi.nlm.nih.gov/geo/). The corresponding profiles comprising detailed

clinicopathological characteristics were also generated. The TCGA RNA-seq dataset was used

as the training set and the CGGA RNA-seq dataset as the validation set. Furthermore, the

CGGA microarray profile, as well as the GSE16011 and REMBRANDT datasets, were used to

test the prognostic value of the developed risk model.

**Immune-related genes**

The 1039 known immunologically relevant genes were downloaded from ImmPort database

(www.immport.org), and 1100 immune-related genes were extracted from the

"IMMUNE_RESPONSE" gene set (Molecular Signatures Database V7.0,

http://software.broadinstitute.org/gsea/msigdb/index.jsp). Finally, after combining these two

independent gene sets, 1763 immune-related genes were selected for the next analyses

(Table S1).

**Identification of prognostic MGMT methylation related immune genes**

137 GBM patients with data on gene expression and MGMT methylation status was selected

in the TCGA cohort, and the differentially expressed genes between MGMT unmethylated and

methylated samples were identified using the "DESeq2" R package (19). The differentially

expressed immune genes were filtered using the criterion of log2 fold change > 1 and

FDR-adjusted $P < 0.05$. The prognostic value of the identified genes was further investigated

using a multivariate Cox regression model integrating clinicopathological information, including

age, Karnofsky performance score (KPS), IDH mutation, and MGMT methylation status. Then,

genes with significant prognostic value ($P < 0.05$) were screened for further analyses.

**Construction of a risk model based on immune-related genes**

The least absolute shrinkage and selection operator (LASSO) regression model with 10-fold

cross-validation was adopted to select the most useful prognostic factors for GBM patients.

The optimal lambda value was estimated based on the minimum criteria. Then, the risk score

for each sample was calculated using the expression levels of genes and corresponding

regression coefficients obtained from the multivariable Cox regression analysis. The formula of

the risk score model was as follows:

$$\text{Risk score} = \sum_{i=1}^{N} (exp_{genei} \times \beta_{genei})$$

; where N is the number of genes in the signature, $\beta_{genei}$ is the regression coefficient, and

$exp_{genei}$ represents the expression value of a specific gene in a single sample. The cutoff for

classifying GBM patients into high and low-risk groups was determined using X-title 3.6.1

software (20). Kaplan-Meier survival analysis was used to evaluate the predictive value of the

risk model, and time-dependent receiver operating characteristic (ROC) curves were drawn to

assess the efficiency of the risk model in predicting the clinical outcomes at different times.

This procedure was completed using the "timeROC" R package.

**Functional enrichment analyses**

Metascape, an easy-to-use web portal that provides a comprehensive analysis for the

functional annotation of lists of genes (21), was used to perform Gene Ontology (GO) term

enrichment and the Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis for

genes of interest. Gene set enrichment analysis (GSEA) analysis was conducted using the R

package "fgsea" (22). GSEA gene set (C5 collection) was downloaded from Molecular

Signatures Database (v7.0). To obtain robust results, we performed gene-set permutations for

10000 times during GSEA. Enriched terms with FDR-adjusted P < 0.05 were considered as

statistically significant.

**Quantification of immune-cell infiltration by single-sample gene set enrichment**

**analysis (ssGSEA)**

The estimation of overall immune cell type fractions in GBM samples was performed using the

ssGSEA algorithm in the "GSVA" R package (23), which uses 24 different types of immune cell

marker genes derived from previous research to analyze each GBM sample (24). Markers

associated with cells of the innate immune system, including natural killer (NK) cells, NK

CD56$^{dim}$ cells, NK CD56$^{bright}$ cells, dendritic cells (DCs), immature DCs (iDCs), activated DCs

(aDCs), neutrophils, mast cells, eosinophils, and macrophages, as well as those associated

with cells of the adaptive immune system, including B, T central memory (Tcm), CD8$^{+}$ T, T

effector memory (Tem), T follicular helper (Tfh), Tγδ, Th1, Th2, Th17, and Treg cells, were

included in the gene list. A numeric matrix consists of the enrichment score of each immune

cell type in a single sample was obtained via ssGSEA.

**Prediction of clinical responses to immune checkpoint blockade**

We used the computational method of Tumor Immune Dysfunction and Exclusion (TIDE) (25)

to predict the likelihood of response to immune checkpoint blockade therapy for GBM patients.

Due to a lack of sufficient data from normal brain tissues in the TCGA dataset, the TCGA

TARGET GTEx dataset from UCSC Toil RNA-seq Recompute project (26), which includes a

conjoint analysis of three independent datasets in order to make the gene expression of

samples from different datasets or batches comparable, was adopted for subsequent studies.

The gene expression values of GBM patients were normalized towards the corresponding

average values of normal brain tissues (cortex, frontal cortex (Ba9), and anterior cingulate

cortex (Ba24)) from the GTEx project before subjecting the data to the TIDE algorithm. Next,

the subclass mapping method (SubMap) (27) was used to compare the gene expression

matrices of different risk groups with the expression profile of melanoma patients who received

immunotherapy targeting *CTLA-4* and *PD-1 (28)*. This step was performed using the default

parameters in the SubMap module on the GenePattern website

(http://genepattern.broadinstitute.org/).

**Construction and validation of a nomogram**

Multivariable Cox proportional hazards regression analysis was applied with the following

clinical-relevant covariates: gender, age, MGMT methylation status, Karnofsky Performance

Status (KPS) score, IDH mutation status, and risk score. A combined nomogram was

generated as a quantitative tool for predicting the likelihood to die of each patient using the

"regplot" R package. The concordance index (C-index) was calculated to assess the

consistency between model prediction and actual clinical outcomes of patients. The calibration

plot was generated to evaluate the accuracy of the prediction for 1- and 2- year overall survival

using this nomogram by the "rms" R package.

**Statistical analysis**

The LASSO Cox regression analysis was carried out using the "glmnet" R package. Restricted

mean survival (RMS) represents the loss in average life expectancy for patients, and the RMS,

as well as time ratio for the risk model in each dataset, were calculated using the "survRM" R

package. Tumor purity, which represents the heterogeneity of each sample, was estimated by

"ESTIMATE" R package (29). Univariate and multivariate Cox regression analyses were used

to identify the risk-score-based signature as an independent prognostic factor for GBM

patients. R software (version 3.6.1, www.r-project.org) was used for all statistical analyses.

Fisher's exact test was used to compare the differences of immunotherapy response between

high- and low-risk groups and was implemented using the "fisher.test" function in R software.

Mann-Whitney U test was used to assess the statistical significance of differences between the

means of continuous data. All hypothetical tests were two-sided, and P values < 0.05 were

considered to indicate statistical significance.

**Results**

**Association between MGMT methylation and the immunological phenotype of GBM**

There is a large body of work implicating that MGMT methylation can affect the prognosis as

well as the effectiveness of chemotherapy in GBM. However, the influence of MGMT promoter

status on the immunological phenotype of GBM had, to our best knowledge, not been

investigated prior to this study. Here, we utilized the RNA-seq gene expression profiles and

clinical information of GBM patients from the TCGA cohort to explore the differences in

underlying immunological mechanisms according to the MGMT methylation status. Patients

were divided into $MGMT^U$ (n = 81) and $MGMT^M$ (n = 56) groups. GSEA analysis showed that

immune-related biological processes, such as humoral immune response (normalized

enrichment score (NES) = 1.95, FDR < 0.05), adaptive immune response (NES = 1.68, FDR <

0.05), innate immune response (NES = 1.44, FDR < 0.05), and activation of immune response

(NES = 1.30, FDR < 0.05), were significantly enriched in the $MGMT^U$ group (Figure 1A).

Notably, all immune-related GO terms with NES > 1 were only enriched in the MGMT$^U$ group

(Table S2).

To further investigate the potential immune pathways and key immune-related genes driving

this phenomenon, differentially expressed genes between the MGMT$^U$ and MGMT$^M$ groups

were identified (Figure 1B, Table S3). Sixty-nine immune-related genes were found to be more

highly expressed in the MGMT$^U$ than the MGMT$^M$ group (log2 fold change > 1, FDR-adjusted P

< 0.05), such as CXC motif chemokine ligand (CXCL1, 2, 6, 12, 13), immunoglobulin kappa

variable family (IGKV1-27, IGKV2-28, IGKV3-11, IGKV3-15), and immunoglobulin heavy

constant gamma cluster (IGHG1, 2, 3, 4).

**Construction of an immune risk model based on MGMT methylation status**

The prognostic effect of the differentially expressed immune genes was investigated by

applying multivariate Cox regression adjusted for other survival-related clinicopathological

covariates. Leukocyte immunoglobulin like receptor A5 (LILA5), immunoglobulin heavy

constant gamma 4 (IGHG4), interleukin 1 receptor type 1 (IL1R1), TNF superfamily member 9

(TNFSF9), immunoglobulin heavy constant alpha 2 (IGHA2), GATA binding protein 3 (GATA3),

vitamin D receptor (VDR), CD70, CXC motif chemokine ligand 13 (CXCL13), and TNF

receptor superfamily member 9 (TNFRSF9) were selected as independent factors associated

with the overall survival of GBM patients (Table S4). All these immune genes were regarded as

risk factors with a hazard ratio (HR) > 1. Among these candidates, LASSO Cox regression

identified six highly relevant immune-related genes: LILRA5, IGHG4, TNFSF9, GATA3, VDR,

and TNFRSF9 (Figure S1A, B). The robustness of the result was confirmed using 10-fold

cross-validation. Circos plots (30) were generated to delineate the correlations between these

genes in both the TCGA and CGGA datasets (Figure S2A, B). It is worth noting that the

relationships of these genes were quite similar in different datasets indicating that the

identified relevant immune genes had a stable positive correlation in GBM.

The risk score of each GBM patient was calculated by combining the expression level of genes

and corresponding coefficients derived from the multivariate Cox regression analysis. Patients

were divided into high- and low-risk groups according to the optimal cutoff value estimated

using X-title software. Kaplan–Meier survival analysis showed that patients from the high-risk

group had shorter overall survival than those with low risk in the TCGA cohort (HR = 2.6, 95%

confidence interval (CI): 1.75-3.87, $P < 0.001$) (Figure 2A). We also applied this computational

formula to the CGGA RNA-seq, CGGA microarray, GSE16011, and REMBRANDT datasets to

further explore the prognostic value of the identified immune-genetic signature. Consistent

with the results of the TCGA cohort, high-risk patients had significantly worse outcomes than

those with low risk (Figure 2B-2E). Significant RMS time ratios ranging from 1.465 to 1.854

were observed in the five datasets ($P < 0.05$, Table S6). In the TCGA dataset, we also checked

the efficiency of this model in patients with or without MGMT methylation (Figure 2F-2G). The

results showed that low-risk patients had a significant survival advantage over the high-risk

group in these two counterparts ($P < 0.05$). Time-dependent ROC curves were generated to

evaluate the efficiency of this risk signature in predicting 1-, 2-, and 3-year survival of GBM

patients and the risk model showed considerable predictive potential in different cohorts

(Figure 2H-2L).

To confirm whether the risk model can be used as an independent prognostic tool for GBM,

univariate and multivariate Cox regression analyses were applied to the TCGA dataset. The

risk score signature was adjusted using prognostic information of clinical factors that had been

deemed statistically significant in univariate analysis (P < 0.05). The HRs for the signature in

the univariate and multivariate analyses were 3.107 (P < 0.001, 95% CI: 2.005-4.816) and

2.476 (P < 0.001, 95% CI: 1.581-3.878), respectively. The risk model was further validated in

the CGGA RNA-seq cohort, with HRs of 1.896 (P = 0.003, 95% CI: 1.246-2.886) and 1.644 (P

= 0.03, 95% CI: 1.05-2.573), respectively (Table 1). Thus, the signature based on

immune-related genes can be used independently to predict the overall survival of GBM

patients.

**Functional analysis of the MGMT methylation-based signature**

To further identify the potential biological processes and signaling pathways associated with

the immune gene signature, GO and KEGG functional analyses were performed, and the

results were graphed using Metascape online tool. Firstly, we selected a list of genes that were

strongly correlated with the risk score (|Pearson correlation coefficient| (|R|) > 0.4, P < 0.05) in

the TCGA and CGGA RNA-seq cohorts (Table S7). Heatmaps for these genes and

corresponding detailed clinicopathological features were plotted, as shown in Figure S3A, B.

Functional analysis demonstrated that these genes were significantly enriched in immune

processes, such as lymphocyte activation, cytokine production, cytokine-mediated signaling

pathway, leukocyte migration, activation of the immune response, regulation of the

inflammatory response, and interferon-gamma production (P < 0.05) (Figure 3A, B). Similarly,

immune-related terms such as cytokine-mediated signaling pathway, myeloid leukocyte

activation, cytokine production, leukocyte migration, TNF signaling pathway, and NF-kappa B

signaling pathway were also enriched in genes derived from the CGGA RNA-seq cohort (P <

0.05) (Figure S4A, B).

**Association of the risk score with tumor purity and immune cell infiltration in the GBM**

**microenvironment**

The intratumoral niche is a complex microenvironment that consists of the tumor and

non-tumor cells, such as stromal and immune cells (31). It is now clear that a mass of

non-tumor cells plays an important role in the initiation and progression of various human

cancers (32-34). Tumor purity is defined as the proportion of cancer cells in the mixed tumor

tissues (35) and has been identified as an effective prognostic index in glioma (2). In an

attempt to clarify the relationship between the immune gene-based risk score and the tumor

purity, we used the ESTIMATE algorithm to calculate the purity of GBM samples based on the

transcriptomic gene expression data. In both the TCGA and CGGA RNA-seq cohorts, the risk

score was negatively correlated with the tumor purity (R = -0.6, P < 0.0001 in TCGA, and R =

-0.4, P <0.0001 in CGGA RNA-seq) (Figure 4A, B).

We then used the ssGSEA algorithm to evaluate the immune cell populations in the tumors

from the two independent datasets. The contents of different types of immune cells varied as

the risk score increased. The results showed that samples from the high-risk group in the

TCGA cohort exhibited a higher abundance of pro-tumor immune cells, such as plasmacytoid

dendritic cells (pDCs), immature dendritic cells (iDCs), CD56dim natural killer cells,

macrophages, neutrophils, regulatory T cells (Tregs), and mast cells (P < 0.05) (Figure 4C, E).

At the same time, B cells, which indicate a favorable prognosis of GBM patients (36), were

highly abundant in the tumor samples from the low-risk group (P < 0.05). Similarly, high-risk

patients in the CGGA RNA-seq cohort had significantly higher quantities of pro-tumor immune

cells, except for pDCs and mast cells ($P < 0.05$) (Figure 4D, F). Therefore, we concluded that

the immune-related biological processes and pathways associated with the risk score might

result from the observed significant differences in differentiation or recruitment of various

immune cell types.

**The risk score is correlated with immunosuppressive processes and the inflammatory**

**responses**

To investigate the potential mechanisms underlying the correlation between the genes related

to the immune response and the risk score, we assessed the relationship between the risk

score and glioma-associated immunosuppressors. The gene list of glioma-related

immunosuppressors, including immunosuppressive cytokines and checkpoints,

tumor-supportive macrophage chemotactic and skewing molecules, immunosuppressive

signaling pathways, and immunosuppressors were obtained from the published literature (37,

38). The expression of immunosuppressive genes in tumors from the two GBM cohorts was

plotted against the increasing risk score (Figure S5A, B). Following ssGSEA analysis, the

score of each immunosuppression-related term was calculated for each tumor sample.

Interestingly, the risk score was positively correlated with the identified

immunosuppression-related terms in both the TCGA and CGGA RNA-seq cohorts (Figure 5A,

B). This finding further corroborated that the immune gene signature is related to

immunological suppression in GBM via changes in the expression of immunosuppressors.

As demonstrated above, genes correlated with the risk score were found to be significantly

enriched in the category inflammatory response. Metagenes, i.e., expression patterns of

various inflammation-related genes, were used to depict the relationship between the risk

score and inflammatory activities in GBM. Except for interferon, most metagenes (HCK, IgG,

LCK, MHC-I, MHC-II, and STAT1) were found to be positively associated with the risk score in

both the TCGA and CGGA RNA-seq cohorts (P < 0.05) (Figure 5C, D, Figure S6A, B). These

results revealed that the risk score was intimately tied to the B- and T-cell mediated immune

response, as well as macrophage activation, but not to genes regulated by interferon during

the remodeling of the immune microenvironment.

**High-risk patients are likely to be more sensitive to immunotherapy**

Pioneering investigations revealed that immunotherapy targeting immune checkpoints offered

great hope for the clinical treatment of human cancers (39). We investigated the correlation

between the risk score and the expression levels of several well-known immune checkpoints,

both in the TCGA and CGGA RNA-seq datasets. Following Pearson correlation analysis, the

risk score was found to be tightly associated with the expression of immune checkpoint

molecules, including TIM-3, TIGIT (except in the CGGA RNA-seq cohort), LAG-3, PD-1,

CTLA-4, CD80, PD-L1, PD-L2, B7-H3, and IDO1 (P < 0.05; Figure 6A, B).

Given that the immune checkpoint blockade has not been used as a routine treatment for GBM,

we modeled the potential response of patients in the different risk groups to immunotherapy,

including antibodies against PD-1 and CTLA-4. Following the TIDE instructions, normal

specimens from the GTEx database were used to adjust the gene expression matrix of GBM

samples from the TCGA cohort. However, considering none normal sample was enrolled in the

CGGA dataset, the TIDE algorithm was only utilized in the TCGA cohort. The analysis

indicated that the high-risk group (53.164%, 42/79) was significantly more likely to benefit from

treatment with immune checkpoint inhibitors than the low-risk group (20.690%, 12/58) (P <

0.001) (Figure 6C and Table S8). SubMap is an unsupervised clustering method that can

match subclasses in two independent gene expression data sets (27). Here, SubMap analysis

further indicated that patients in the high-risk group are more likely to respond to anti-PD-1

immunotherapy (Bonferroni corrected $P = 0.024$) (Figure 6D).

**Development of a nomogram based on MGMT methylation-related signature**

To develop a quantitative tool for predicting the prognosis of GBM patients in the TCGA cohort,

we established a nomogram by integrating clinicopathological risk factors and immune gene

signature based on the multivariable Cox proportional hazards model (Figure 7A). The point

scale in the nomogram was utilized to generate point to these variables, and the risk of death

of each GBM patient was qualified by accumulating total points of all variables. The risk score

was found to have the most excellent weight among all these variables, which was consistent

with the result of the previous multivariable Cox regression analysis. The C-index of this

nomogram reached 0.704 (95%CI: 0.672-0.736). The result of the calibration plots further

confirmed the significant consistency between predicted and observed actual clinical

outcomes of GBM patients (Figure 7C). We further constructed a nomogram for the CGGA

GBM cohort based on the risk model and other available clinicopathological features, and the

feasibility of this nomogram was confirmed using the calibration plots (Figure 7B, D).

**Discussion**

In GBM, MGMT promoter methylation status is an important determinant of the

aggressiveness of the tumor. Malignant gliomas often show resistance to alkylating drugs due

to the increased expression of MGMT, which remains a significant obstacle to clinical

treatment (40). Previously, considerable efforts have been made to exploit the mechanisms of

chemo-resistance induced by MGMT promoter demethylation. Over the past few years, the

promising outcomes of immunotherapy in malignant tumors have attracted enormous attention.

However, whether tumor immunity can be affected by the methylation status of the MGMT

promoter has, to our best knowledge, not been explored before. Here, we sought to investigate

the potential immunological mechanisms associated with the methylation status of MGMT in

GBM by a comprehensive bioinformatic analysis. We addressed the critical role of MGMT

methylation in regulating the remodeling of the tumor microenvironment of GBM.

The solid tumor mass of GBM is composed of both cancer cells and non-cancerous cells, such

as immune cells, endothelial cells, fibroblasts, as well as some CNS-specific cell types such as

astrocytes and microglia (41, 42). The discovery that multiple immune cell types can infiltrate

the blood-brain barrier and the presence of reactivity against antigens derived from the CNS

has made the view of the brain as an immunologically privileged site obsolete (43, 44). The

most abundant immune cells, macrophages, comprise up to 30% of the total cells inside a

glioma tumor (45). Tumor-associated macrophages suppress T cell responses due to

insufficient expression of molecules associated with T cell co-stimulation (46). Dendritic cells

are responsible for presenting antigens to T cells to induce immune responses in the CNS.

The application of dendritic cell-based vaccination in GBM patients has produced survival

benefits due to enhanced stimulated T-cell responses (47). Additionally, several studies

demonstrated that reprogramming immunosuppressive T-cell subsets can stimulate anti-tumor

immunity in glioma (46, 48). In this study, we discovered that genetic signatures of

immune-related biological processes and signaling pathways were highly enriched in MGMT

unmethylated specimens. Then, we filtered differentially expressed immune-related genes

according to the MGMT promoter status and generated a signature based on six genes that

can predict the clinical outcomes of GBM patients. At the same time, we investigated the

contents of immune cells in the bulk tumor to predict immune infiltration using a marker

gene-based approach based on transcriptomic data. Notably, we found that the infiltrating

immune cell types differed significantly between the high and low-risk groups. In agreement

with our hypothesis, the tumors in the high-risk group were severely infiltrated by immune cells

characterized as belonging to pro-tumor subsets. Thus, this risk model provides an

immunological interpretation of the effect of MGMT methylation status on the prognosis of

GBM patients.

There is a large body of work that confirms the immunity-related roles of the genes identified in

this study. For example, the transcription factor GATA3 has been characterized as a critical

regulator in the development of T, B, NK, and innate-like lymphoid cells (49). GATA3 is

responsible for maintaining the function of mature T and Treg cells via the IL-7 and IL-2

signaling pathways, respectively (50). Additionally, ectopic expression of GATA3 was found to

promote the function of human type-2 innate lymphoid cells (51). TNSRSF9, also known as

CD137, is a member of the TNF-receptor family that can activate cytokine production in

cytotoxic T lymphocytes (CTLs) and Th1 cells by binding to its native ligand TNFSF9 (52).

Recently, clinical trials with antibodies targeting CD137 have been launched with the

expectation to improve cancer immunotherapy (53). VDR negatively regulates Forkhead box

M1 (FOXM1) signaling and further suppresses the development of pancreatic ductal

adenocarcinoma (54). In the immune system, VDR expression regulates the proliferation,

differentiation, and function of CTLs (55). Stromal VDR activation reduces tumor-associated

fibrosis and inhibits tumor-supportive signaling events in pancreatic cancer (56).

Immune suppression has been recognized as a hindrance to successful antitumor

immunotherapy. Tumors create a microenvironment that sustains tumor growth as well as

reduces adaptive immunity to tumor-derived antigens (57). We found a strong correlation

between the risk score and immunosuppressive cytokines, tumor-supportive macrophages, as

well as immunosuppressive signaling pathways. This finding implied that the risk score reflects

the immunosuppressive microenvironment of GBM. Monoclonal antibodies that block immune

checkpoint signaling have shown clinical benefits in several malignant tumors, including

melanoma, NSCLC, Hodgkin lymphoma, and urothelial carcinoma. Such antibodies include

pembrolizumab, nivolumab, and pidilizumab, which target PD-1, ipilimumab and

tremelimumab for CTLA-4, or atezolizumab and durvalumab for PD-L1. Additionally, many

clinical trials have been launched to evaluate the application of checkpoint inhibitors in GBM. A

randomized phase III study (NCT02017717) was carried out to test the efficacy and safety of

nivolumab alone versus bevacizumab (58). Clinical treatment of newly diagnosed GBM

patients with ipilimumab and nivolumab is still under investigation (NCT02311920) (59).

Therefore, we explored the relationship between the risk score and the expression levels of

several well-known immune checkpoint molecules. The strong positive correlation implied that

the identified signature mediates immune escape mechanisms in GBM, which provides a

foundation for identifying novel therapeutic targets.

It is noteworthy that not all probands will respond to immune checkpoint blockade effectively in

most cancer types (60). Various factors were reported as predictors for evaluating

immunotherapy effectiveness, such as the mutation burden (61), tumor infiltration by cytotoxic

T cells (62), PD-L1 expression (63), and tumor aneuploidy (64). The known mechanisms of

tumor immune evasion include infiltration by CTLs and the prevention of T cell infiltration (65)

(66). The TIDE method underscores these two distinct aspects and can predict the outcome of

melanoma patients receiving immunotherapy more accurately than other factors (25). We

demonstrated that more patients could benefit from immune checkpoint blockade in the high

than in the low-risk group. Under a strict criterion of adjusted P-values, SubMap analysis

further confirmed that anti-PD-1 therapy might produce favorable outcomes in high-risk

patients.

**Conclusions**

In summary, our study revealed that MGMT promoter methylation status is correlated with

immunological processes and the remodeling of the tumor microenvironment in GBM. The

developed immune-related gene signature could serve as a useful prognostic tool for GBM

patients as well as for predicting which patients would benefit from immunotherapy. These

findings may provide new insights into fundamental aspects of the critical role of MGMT

methylation status in GBM.

**Acknowledgments**

This work was supported by the National Natural Science Foundation of China under [grant

number 81673210]. All authors would like to thank all contributors to the TCGA and CGGA

project, especially.

**References**

1. Stupp R, Mason WP, van den Bent MJ, et al. Radiotherapy plus concomitant and adjuvant

temozolomide for glioblastoma. N Engl J Med 2005;352:987-96.

2. Zhang C, Cheng W, Ren X, et al. Tumor Purity as an Underlying Key Factor in Glioma. Clin

Cancer Res 2017;23:6279-91.

3. Hegi ME, Liu L, Herman JG, et al. Correlation of O6-methylguanine methyltransferase

(MGMT) promoter methylation with clinical outcomes in glioblastoma and clinical strategies to

modulate MGMT activity. Journal of clinical oncology 2008;26:4189-99.

4. Wick W, Weller M, van den Bent M, et al. MGMT testing—the challenges for

biomarker-based glioma treatment. Nature Reviews Neurology 2014;10:372.

5. Pegg AE. Mammalian O6-alkylguanine-DNA alkyltransferase: regulation and importance in

response to alkylating carcinogenic and therapeutic agents. Cancer research

1990;50:6119-29.

6. Karran P, Bignami M. Self-destruction and tolerance in resistance of mammalian cells to

alkylation damage. Nucleic acids research 1992;20:2933.

7. Liu L, Gerson SL. Targeted modulation of MGMT: clinical implications. Clinical cancer

research 2006;12:328-31.

8. Dolan ME, Pegg AE. O6-benzylguanine and its role in chemotherapy. Clinical Cancer

Research 1997;3:837-47.

9. Hegi ME, Diserens A-C, Gorlia T, et al. MGMT gene silencing and benefit from

temozolomide in glioblastoma. New England Journal of Medicine 2005;352:997-1003.

10. Molenaar RJ, Verbaan D, Lamba S, et al. The combination of IDH1 mutations and MGMT

methylation status predicts survival in glioblastoma better than either IDH1 or MGMT alone.

Neuro-oncology 2014;16:1263-73.

11. Fukumura D, Kloepper J, Amoozgar Z, et al. Enhancing cancer immunotherapy using

antiangiogenics: opportunities and challenges. Nature reviews Clinical oncology 2018;15:325.

12. Heimberger AB, Abou-Ghazal M, Reina-Ortiz C, et al. Incidence and prognostic impact of

FoxP3+ regulatory T cells in human gliomas. Clin Cancer Res 2008;14:5166-72.

13. Nduom EK, Weller M, Heimberger AB. Immunosuppressive mechanisms in glioblastoma.

Neuro Oncol 2015;17 Suppl 7:vii9-vii14.

14. Rizvi NA, Mazières J, Planchard D, et al. Activity and safety of nivolumab, an anti-PD-1

immune checkpoint inhibitor, for patients with advanced, refractory squamous non-small-cell

lung cancer (CheckMate 063): a phase 2, single-arm trial. The Lancet Oncology

2015;16:257-65.

15. Hodi FS, O'Day SJ, McDermott DF, et al. Improved survival with ipilimumab in patients with

metastatic melanoma. New England Journal of Medicine 2010;363:711-23.

16. Louveau A, Smirnov I, Keyes TJ, et al. Structural and functional features of central nervous

system lymphatic vessels. Nature 2015;523:337-41.

17. Cloughesy TF, Mochizuki AY, Orpilla JR, et al. Neoadjuvant anti-PD-1 immunotherapy

promotes a survival benefit with intratumoral and systemic immune responses in recurrent

glioblastoma. Nat Med 2019;25:477-86.

18. Qian Z, Li Y, Fan X, et al. Molecular and clinical characterization of IDH associated immune

signature in lower-grade gliomas. Oncoimmunology 2018;7:e1434466.

19. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome biology 2014;15:550.

20. Camp RL, Dolled-Filhart M, Rimm DL. X-tile: a new bio-informatics tool for biomarker

assessment and outcome-based cut-point optimization. Clin Cancer Res 2004;10:7252-9.

21. Zhou Y, Zhou B, Pache L, et al. Metascape provides a biologist-oriented resource for the

analysis of systems-level datasets. Nature Communications 2019;10:1523.

22. Sergushichev A. An algorithm for fast preranked gene set enrichment analysis using

cumulative statistic calculation. BioRxiv 2016:060012.

23. Subramanian A, Tamayo P, Mootha VK, et al. Gene set enrichment analysis: a

knowledge-based approach for interpreting genome-wide expression profiles. Proceedings of

the National Academy of Sciences 2005;102:15545-50.

24. Bindea G, Mlecnik B, Tosolini M, et al. Spatiotemporal dynamics of intratumoral immune

cells reveal the immune landscape in human cancer. Immunity 2013;39:782-95.

25. Jiang P, Gu S, Pan D, et al. Signatures of T cell dysfunction and exclusion predict cancer

immunotherapy response. Nature medicine 2018:1.

26. Vivian J, Rao AA, Nothaft FA, et al. Toil enables reproducible, open source, big biomedical

data analyses. Nature Biotechnology 2017;35:314-6.

27. Hoshida Y, Brunet J-P, Tamayo P, et al. Subclass mapping: identifying common subtypes in

independent disease data sets. PloS one 2007;2:e1195.

28. Roh W, Chen P-L, Reuben A, et al. Integrated molecular analysis of tumor biopsies on

sequential CTLA-4 and PD-1 blockade reveals markers of response and resistance. Science

translational medicine 2017;9:eaah3560.

29. Yoshihara K, Shahmoradgoli M, Martínez E, et al. Inferring tumour purity and stromal and

immune cell admixture from expression data. Nature communications 2013;4:2612.

30. Gu Z, Gu L, Eils R, et al. circlize Implements and enhances circular visualization in R.

Bioinformatics 2014;30:2811-2.

31. Golebiewska A, Bougnaud S, Stieber D, et al. Side population in human glioblastoma is

non-tumorigenic and characterizes brain endothelial cells. Brain 2013;136:1462-75.

32. Mao Y, Keller ET, Garfield DH, et al. Stromal cells in tumor microenvironment and breast

cancer. Cancer and Metastasis Reviews 2013;32:303-15.

33. Bussard KM, Mutkus L, Stumpf K, et al. Tumor-associated stromal cells as key contributors

to the tumor microenvironment. Breast Cancer Research 2016;18:84.

34. Wang M, Zhao J, Zhang L, et al. Role of tumor microenvironment in tumorigenesis. Journal

of Cancer 2017;8:761.

35. Zhang N, Wu H-J, Zhang W, et al. Predicting tumor purity from methylation microarray data.

Bioinformatics 2015;31:3401-5.

36. Li B, Severson E, Pignon JC, et al. Comprehensive analyses of tumor immunity:

implications for cancer immunotherapy. Genome Biol 2016;17:174.

37. Doucette T, Rao G, Rao A, et al. Immune heterogeneity of glioblastoma subtypes:

extrapolation from the cancer genome atlas. Cancer Immunol Res 2013;1:112-22.

38. Cai J, Chen Q, Cui Y, et al. Immune heterogeneity and clinicopathologic characterization of

IGFBP2 in 2447 glioma samples. Oncoimmunology 2018;7:e1426516.

39. Sharma P, Allison JP. The future of immune checkpoint therapy. Science 2015;348:56-61.

40. Sarkaria JN, Kitange GJ, James CD, et al. Mechanisms of chemoresistance to alkylating

agents in malignant glioma. Clinical cancer research 2008;14:2900-8.

41. Quail DF, Joyce JA. The Microenvironmental Landscape of Brain Tumors. Cancer Cell

2017;31:326-41.

42. Wang Q, Hu B, Hu X, et al. Tumor Evolution of Glioma-Intrinsic Gene Expression Subtypes

Associates with Immunological Changes in the Microenvironment. Cancer Cell 2017;32:42-56

e6.

43. Louveau A, Harris TH, Kipnis J. Revisiting the mechanisms of CNS immune privilege.

Trends in immunology 2015;36:569-77.

44. Weiss N, Miller F, Cazaubon S, et al. The blood-brain barrier in brain homeostasis and

neurological diseases. Biochimica et Biophysica Acta (BBA)-Biomembranes

2009;1788:842-57.

45. Graeber MB, Scheithauer BW, Kreutzberg GW. Microglia in brain tumors. Glia

2002;40:252-9.

46. Hussain SF, Yang D, Suki D, et al. The role of human glioma-infiltrating

microglia/macrophages in mediating antitumor immune responses. Neuro-oncology

2006;8:261-79.

47. Prins RM, Soto H, Konkankit V, et al. Gene expression profile correlates with T-cell

infiltration and relative survival in glioblastoma patients vaccinated with dendritic cell

immunotherapy. Clinical cancer research 2011;17:1603-15.

48. Fecci PE, Sweeney AE, Grossi PM, et al. Systemic anti-CD25 monoclonal antibody

administration safely enhances immunity in murine glioma without eliminating regulatory T

cells. Clinical Cancer Research 2006;12:4294-305.

49. Wan YY. GATA3: a master of many trades in immune regulation. Trends in immunology

2014;35:233-42.

50. Ma A, Koka R, Burkett P. Diverse functions of IL-2, IL-15, and IL-7 in lymphoid homeostasis.

Annu Rev Immunol 2006;24:657-79.

51. Mjösberg J, Bernink J, Golebski K, et al. The transcription factor GATA3 is essential for the

function of human type 2 innate lymphoid cells. immunity 2012;37:649-59.

52. Ascierto PA, Simeone E, Sznol M, et al., editors. Clinical experiences with anti-CD137 and

anti-PD1 therapeutic antibodies. Seminars in oncology; 2010: Elsevier.

53. Yonezawa A, Dutt S, Chester C, et al. Boosting cancer immunotherapy with anti-CD137

antibody therapy. Clinical Cancer Research 2015;21:3113-20.

54. Li Z, Jia Z, Gao Y, et al. Activation of vitamin D receptor signaling downregulates the

expression of nuclear FOXM1 protein and suppresses pancreatic cancer cell stemness.

Clinical Cancer Research 2015;21:844-53.

55. Sarkar S, Hewison M, Studzinski GP, et al. Role of vitamin D in cytotoxic T lymphocyte

immunity to pathogens and cancer. Critical reviews in clinical laboratory sciences

2016;53:132-45.

56. Sherman MH, Ruth TY, Engle DD, et al. Vitamin D receptor-mediated stromal

reprogramming suppresses pancreatitis and enhances pancreatic cancer therapy. Cell

2014;159:80-93.

57. Hurwitz AA, Watkins SK. Immune suppression in the tumor microenvironment: a role for

dendritic cell-mediated tolerization of T cells. Cancer Immunology, Immunotherapy

2012;61:289-93.

58. Sampson JH, Vlahovic G, Desjardins A, et al. Randomized phase IIb study of nivolumab

(anti-PD-1; BMS-936558, ONO-4538) alone or in combination with ipilimumab versus

bevacizumab in patients (pts) with recurrent glioblastoma (GBM). American Society of Clinical

Oncology; 2014.

59. Binder DC, Davis AA, Wainwright DA. Immunotherapy for cancer in the central nervous

system: current and future directions. Oncoimmunology 2016;5:e1082027.

60. Sharma P, Hu-Lieskovan S, Wargo JA, et al. Primary, Adaptive, and Acquired Resistance

to Cancer Immunotherapy. Cell 2017;168:707-23.

61. Snyder A, Makarov V, Merghoub T, et al. Genetic basis for clinical response to CTLA-4

blockade in melanoma. New England Journal of Medicine 2014;371:2189-99.

62. Van Allen EM, Miao D, Schilling B, et al. Genomic correlates of response to CTLA-4

blockade in metastatic melanoma. Science 2015;350:207-11.

63. Nishino M, Ramaiya NH, Hatabu H, et al. Monitoring immune-checkpoint blockade:

response evaluation and biomarker development. Nature reviews Clinical oncology

2017;14:655.

64. Davoli T, Uno H, Wooten EC, et al. Tumor aneuploidy correlates with markers of immune

evasion and with reduced response to immunotherapy. Science 2017;355:eaaf8399.

65. Gajewski TF, Schreiber H, Fu Y-X. Innate and adaptive immune cells in the tumor

microenvironment. Nature immunology 2013;14:1014.

66. Joyce JA, Fearon DT. T cell exclusion, immune privilege, and the tumor microenvironment.

Science 2015;348:74-80.

**Supplementary material**

**Table S1.** List of immune genes obtained from the ImmPort database and

"GO_IMMUNE_RESPONSE" term.

**Table S2.** Gene enrichment in GBM patients with MGMT unmethylation in the TCGA cohort.

**Table S3.** Differentially expressed genes between MGMT unmethylated and methylated

samples in the TCGA GBM cohort.

**Table S4.** Multivariate Cox regression analysis of differentially expressed immune genes.

**Table S5.** Supplementary figures and corresponding figure legends.

**Table S6.** Restricted mean survival (RMS) time ratio between low- and high-risk groups in

different datasets.

**Table S7.** Genes strongly correlated with risk score in TCGA and CGGA RNA-seq datasets.

**Table S8.** Predicted outcomes of immunotherapy for GBM patients (TIDE algorithm).

**Figure Legends**

**Figure 1.** Functional annotation of MGMT methylation status in the TCGA dataset. **A**. Gene

Set Enrichment Analysis showing genes upregulated in MGMT unmethylated samples

evaluated in the gene sets representative for immune-related biological processes and

pathways. Normalized enrichment score (NES) and FDR are generated for each term. **B**.

Volcano plot of fold changes of genes in groups with different MGMT methylation status.

Overexpressed genes in MGMT unmethylated group were plotted as red dots while blue dots

represent downregulated genes. The x and y-axis represent fold-change and the P-values

estimated by DESeq2.

**Figure 2.** Prognostic analysis of the immune gene-based signature. Kaplan-Meier analysis of

overall survival based on high *vs.* low risk in different cohorts (**A-E**). The prognostic value of

this model was further validated in subgroups with MGMT methylation (**F**) and unmethylation

(**G**) from the TCGA dataset. P values were obtained from the log-rank test. Time-dependent

ROC curves analysis of this signature in the TCGA (**H**), CGGA RNA-seq (**I**), CGGA microarray

(**J**), GSE16011 (**K**), and REMBRANDT (**L**) datasets.

**Figure 3.** Biological functions of the immune gene signature in GBM in the TCGA dataset. **A.**

Bar plot showing the top 20 terms derived from the gene set enrichment of risk score. The

x-axis represents statistical significance. **B.** The enrichment network plot visualizing the

relationship between a set of representative terms. Each term is assigned with a unique color.

**Figure 4.** The risk signature is associated with tumor purity and immune infiltration in the

microenvironment of GBM. Correlation of risk score and tumor purity in GBM samples in the

TCGA (**A**) and CGGA RNA-seq (**B**) cohorts. The coefficient and P-value were obtained from

Pearson correlation analysis. Heatmaps delineate the relationship between risk score and the

contents of 24 types of immune cells infiltrated in the tumor tissues in the TCGA (**C**) and CGGA

RNA-seq (**D**) cohorts. Box plots showing significantly different immune cells among high- and

low-risk patients in the TCGA (**E**) and CGGA RNA-seq (**F**) datasets. * p < 0.05, ** p < 0.01, ***

p < 0.001, **** p < 0.0001, with Mann-Whitney U test.

**Figure 5.** The risk signature is associated with immunosuppression and inflammatory activities

in GBM. Scatterplots showing the correlation between risk score and each

immunosuppressive activity in the TCGA (**A**) and CGGA RNA-seq (**B**) cohorts. The coefficients

and P-values were calculated from Pearson correlation analyses. Corrplots showing the

correlation between risk score and seven types of inflammatory activities in the TCGA (**C**) and

CGGA RNA-seq (**D**) cohorts. The black cross indicates no statistical significance.

**Figure 6.** The risk signature may be tightly associated with the responses of immunotherapy.

Corrplots showing the correlation between risk score and the expression levels of several

immune checkpoint molecules in the TCGA (**A**) and CGGA RNA-seq (**B**) cohorts. The black

cross indicates no statistical significance. **C.** Comparison of the predicted immunotherapeutic

responses in different risk groups using the TIDE algorithm. P-value was obtained from Fisher

exact test. **D.** The SubMap analysis concluded that high-risk patients might be more sensitive

to anti-PD-1 immunotherapy.

**Figure 7.** Developed nomogram to predict the probability of survival in GBM patients.

Nomogram built with clinicopathological factors incorporated estimating 1-, 2- and 3-year

overall survival for GBM patients in the TCGA (**A**) and CGGA RNA-seq (**B**) cohorts. The

calibration curves describing the consistency between predicted and observed overall survival

at different time points in the TCGA (**C**) and CGGA RNA-seq (**D**) cohorts. The estimated

survival was plotted on the x-axes, and the actual outcomes were plotted on the y-axes. The

gray 45-degree dotted line represents an ideal calibration mode.

703 **Table 1. Univariate and multivariate analyses of clinicopathological characteristics and**
704 **immune gene-based signature with overall survival in GBM cohorts**
705

| Variable | Univariate analysis | | Multivariate analysis | |
|---|---|---|---|---|
| | HR (95 % CI) | *P* | HR (95 % CI) | *P* |
| TCGA (n = 137) | | | | |
| Age >=60 *vs.* <60 | 1.62 (1.09-2.408) | *0.017* | 1.398 (0.921-2.121) | 0.116 |
| Gender Male *vs.* female | 1.165 (0.783-1.735) | 0.451 | | |
| KPS >=80 *vs.* <80 | 0.846 (0.519-1.379) | 0.503 | | |
| IDH status Wild-type *vs.* mutation | 4.544 (1.656-12.467) | *0.003* | 2.444 (0.848-7.047) | 0.098 |

| | | | | | |
|---|---|---|---|---|---|
| MGMT methylation | | | | | |
| Unmethylated *vs.* methylated | | 1.861 (1.23-2.815) | ***0.003*** | 1.715 (1.125-2.613) | ***0.012*** |
| Risk score | | | | | |
| High *vs.* low | | 3.107 (2.005-4.816) | ***<0.001*** | 2.476 (1.581-3.878) | ***<0.001*** |
| CGGA RNA-seq (n = 105) | | | | | |
| Age | | | | | |
| >=60 *vs.* <60 | | 1.99 (1.21-3.273) | ***0.007*** | 1.657 (0.983-2.793) | 0.058 |
| Gender | | | | | |
| Male *vs.* female | | 1.142 (0.744-1.752) | 0.545 | | |
| IDH status | | | | | |
| Wild-type *vs.* mutation | | 1.082 (0.636-1.839) | 0.772 | | |
| Chemotherapy | | | | | |
| Yes *vs.* no | | 0.481 (0.303-0.765) | ***0.002*** | 0.528 (0.33-0.845) | ***0.008*** |
| Radiotherapy | | | | | |
| Yes *vs.* no | | 0.923 (0.511-1.669) | 0.792 | | |
| Risk score | | | | | |
| High *vs.* low | | 1.896 (1.246-2.886) | ***0.003*** | 1.644 (1.05-2.573) | ***0.03*** |

706    Note: Bold italics indicate statistically significant values (P < 0.05).
707

**A**

GO:0046649: lymphocyte activation
GO:0002366: leukocyte activation involved in immune response
GO:0001816: cytokine production
GO:0019221: cytokine-mediated signaling pathway
GO:0002250: adaptive immune response
GO:0050900: leukocyte migration
GO:0002253: activation of immune response
GO:0002683: negative regulation of immune system process
GO:0046631: alpha-beta T cell activation
GO:0009617: response to bacterium
GO:0050727: regulation of inflammatory response
R-HSA-1280218: Adaptive Immune System
GO:0031349: positive regulation of defense response
GO:0002573: myeloid leukocyte differentiation
GO:0071706: tumor necrosis factor superfamily cytokine production
M54: PID IL12 2PATHWAY
R-HSA-1474244: Extracellular matrix organization
GO:0006909: phagocytosis
hsa05150: Staphylococcus aureus infection
GO:0032609: interferon-gamma production

-log10(P)

**B**

**A** TCGA

**B** CGGA RNA-seq

**C**

Response to immunotherapy

*P.fisher = 1.7e−04*

|  | High risk | Low risk |
|---|---|---|
| False | 37 | 46 |
| True | 42 | 12 |

**D**

Nominal *P* value
Bonferroni corrected

| *P* value | CTAL4-noR | CTLA4-R | PD1-noR | PD1-R | |
|---|---|---|---|---|---|
| | | | | | high |
| | | | | | low |
| | | | | *P* = 0.024 | high |
| | | | | | low |