Title

Time-dependent inhibition of covert shift of attention

Authors

Antimo Buonocore[1,2*], Niklas Dietze[3,4,5*], Robert D. McIntosh[3]

*equal contribution

1. Werner Reichardt Centre for Integrative Neuroscience, Tübingen University, Tübingen,

BW, 72076, Germany

2. Hertie Institute for Clinical Brain Research, Tübingen University, Tübingen, BW, 72076,

Germany

3. Human Cognitive Neuroscience, University of Edinburgh, Edinburgh, UK

4. Neuro-cognitive Psychology, Department of Psychology, Bielefeld University, Bielefeld,

NRW, 33501, Germany

5. Cognitive Interaction Technology Cluster of Excellence (CITEC), Bielefeld University,

Bielefeld, NRW, 33501, Germany

Corresponding author:

Antimo Buonocore

Werner Reichardt Centre for Integrative Neuroscience

Otfried-Müller Str. 25

Tübingen, 72076, Germany

Tel: +49 7071 29 88821

antimo.buonocore@cin.uni-tuebingen.de

Abstract

Visual transients can interrupt overt orienting by abolishing the execution of a planned eye

movement due about 90 ms later, a phenomenon known as saccadic inhibition (SI). It is

not known if the same inhibitory process might influence covert orienting in the absence of

saccades, and consequently alter visual perception. In Experiment 1 (n=14), we measured

orientation discrimination during a covert orienting task in which an uninformative

exogenous visual cue preceded the onset of an oriented probe by 140 to 290 ms. In half of

the trials the onset of the probe was accompanied by a brief irrelevant flash, a visual

transient that would normally induce SI. We report a time-dependent inhibition of covert

orienting in which the irrelevant flash impaired orientation discrimination accuracy when

the probe followed the cue by 190 and 240 ms. The interference was more pronounced

when the cue was incongruent with the probe location, suggesting an impact on the

reorienting component of the attentional shift. In Experiment 2 (n=12), we tested whether

the inhibitory effect of the flash could occur within an earlier time range, or only within the

later, reorienting range. We presented probes at congruent cue locations in a time window

between 50 to 200 ms. Similarly to Experiment 1, discrimination performance was altered

at 200 ms after the cue. We suggest that covert attention may be susceptible to similar

inhibitory mechanisms that generate SI, especially in later stages of attentional shifting (>

200ms after a cue), typically associated with reorienting.

Keywords: covert orienting, visual interference, saccadic inhibition

Introduction

The primary function of visual processing is to guide efficient interactions with the surrounding world. This requires us to rapidly integrate new sensory information with ongoing motor behaviour to generate appropriate responses such as inhibiting a planned action in order to orient to a novel stimulus. One of the most striking examples of the fast integration of visual information into a motor command is saccadic inhibition (SI) (Bompas and Sumner 2011; Buonocore and McIntosh 2008; Buonocore and McIntosh 2012; Edelman and Xu 2009; Reingold and Stampe 1999; Reingold and Stampe 2002). It has been consistently observed that visual transient events interrupt ongoing eye movement behaviour, such that saccades that would otherwise be launched about 90 ms later are inhibited, or delayed. This is visualized as a distinct dip in saccadic frequency around 90 ms after the onset of the visual event, and followed by a rebound period in which the probability of making an eye movement is increased (Reingold and Stampe 2002). From a neuronal perspective, SI might be the result of an interruption in the build-up activity of superior colliculus (SC) neurons (Buonocore et al. 2021; Dorris et al. 2007; Munoz and Istvan 1998) coding for the location in space to which the saccade is planned (Munoz and Guitton 1991; Munoz and Wurtz 1993; Munoz and Wurtz 1995).

The exact mechanisms behind SI are still unknown, but there are strong hypotheses suggesting the involvement of omnipause neurons (OPN) in the nucleus raphe interpositus (Büttner-Ennever et al. 1988). OPNs are a small group of neurons that fire tonically during fixation and pause their activity abruptly just before and during saccades (Cohen and Henn 1972; Keller 1974; Luschei and Fuchs 1972). Sudden reactivation of OPNs following the onset of a visual transient might send an inhibitory signal interfering with the build-up activity in upstream oculomotor areas (Buonocore et al. 2020). Despite being one of the most reliable phenomena in oculomotor behaviour, the ecological function of SI is not fully understood.

We recently asked whether this interruption of ongoing oculomotor behaviour might

have functional benefits in terms of facilitating overt reorienting to a new location in space

(Buonocore et al. 2017c). For this purpose, we adapted the well-known double-step

paradigm (Becker and Jürgens 1979; Lisberger et al. 1975), asking participants to make

an eye movement in response to a sudden-onset visual target, which sometimes jumped

to a new location before the first saccade could be launched. In half of the trials the jump

was accompanied by the presentation of a brief (30 ms) visual flash at the top and bottom

of the screen. These "flash-jump" trials induced strong saccadic inhibition, and led to a

higher rate of successful reorienting to the new target location. We suggested that SI

allowed the oculomotor system the time for a decisional process to change the planned

response. Following this empirical demonstration, new models of saccadic inhibition have

been proposed, aiming to unify inhibition with the response inhibition observed in

countermanding tasks (Bompas et al. 2020; Salinas and Stanford 2018). These findings

and models have advanced our understanding of the oculomotor system by suggesting

the existence of a common inhibitory signal, driven by new visual onsets, capable of

interrupting ongoing orienting. Nonetheless, they have been concerned exclusively with

overt orienting (saccades), leaving open the question of whether covert orienting could be

similarly affected.

Moving the eyes is not the only way to modulate sampling of visual information from

the surroundings. Numerous studies have shown perceptual benefits at locations that

have been previously cued, even when no eye movement is made, confirming that it is

possible to orient attention covertly (Carrasco 2011; Posner 1980; see: Posner 2015 for a

review). These studies have mostly shown speeding of simple detection responses, but

covert orienting can also improve the discimination of spatial frequency and lower the

contrast threshold for orientation discimination (Barbot et al. 2012; Cameron et al. 2002;

Carrasco 2011; Fernández et al. 2019; Lee et al. 1999; Pestilli and Carrasco 2005;

Solomon 2004). These changes in visual sensitivity are tightly coupled with the planning of

eye movements (Buonocore et al. 2017b; Deubel and Schneider 1996; Hoffman and

Subramaniam 1995; Remington 1980; Shepherd et al. 1986; Zhao et al. 2012) as implied

by the observation of pre-saccadic perceptual enhancements at the target location of an

upcoming saccade (Hanning et al. 2019; Kowler et al. 1995; Li et al. 2016; Li et al. 2019;

Montagnini and Castet 2007; Ohl et al. 2017; Rolfs and Carrasco 2012). Indeed, the

influential premotor theory of attention posits that covert orienting is nothing other than the

preparation of an eye movement without its execution (premotor theory of attention,

Rizzolatti et al. 1987; Sheliga et al. 1994; Sheliga et al. 1995).

In support of this tight coupling, closer inspection reveals that covert attention shifts

may not be entirely covert. Although large scale eye movements may be absent during

covert attention tasks, much smaller microsaccades can often be detected (for a review on

the topic see: Hafed et al. 2021; Martinez-Conde et al. 2004), and have been suggested

as a biomarker for covert attentional shifts (Engbert and Kliegl 2003; Hafed and Clark

2002; Laubrock et al. 2005; Pastukhov and Braun 2010; Tian et al. 2016; Tian et al. 2018).

The precise nature of the coupling of covert and overt orienting processes is still disputed

(for a review on the topic see: Belopolsky and Theeuwes 2012; Hunt et al. 2019; Klein and

Pontefract 1994; Smith and Schenk 2012), but there are compelling similarities between

the two ways of directing attention in space, which suggest that the underlying

mechanisms might substantially overlap.

Given this theoretical context, we ask whether the phenomenon of SI, which arises

with striking regularity in overt responses, extends to covert orienting. We operationalise

covert orienting in terms of its perceptual consequences, specifically modulations in the

ability to discriminate visual orientation at a cued or an uncued location (Fernández et al.

2019; Pestilli and Carrasco 2005). The key question was whether an irrelevant flash, which

would induce SI in overt tasks, interferes with perceptual discrimination in a covert task.

We adapted our previously-used double-step saccadic task (Buonocore et al. 2017c), to create a covert orienting task suitable for testing the inhibitory influence of an irrelevant visual transient. This effectively combined a classic task for the exogenous cueing of covert attention (Posner 1980) with a SI paradigm. We assume that the onset of the cue starts building-up activity at its location in space (i.e. covert orienting), which may subsequently be followed by reorienting to the opposite (uncued) location. Our initial hypotheses were based on the results of our earlier saccadic study (Buonocore et al. 2017c). We predict that the transient flash will interfere with covert orienting and that this effect will depend on the state of the attentional system, affecting performance only at specific CTOAs. If the pattern of covert orienting follows that of overt orienting, then we may expect this interruption to improve the ability to re-orient to the opposite (uncued) location, giving a relative enhancement of perceptual discrimination for invalidly cued targets at later CTOAs.

Experiment 1 – Materials and Methods

Participants

Fourteen participants (7 female), aged between 21 and 35 years (mean = 25.6, *SD* = 3.86), were included in the data analysis of Experiment 1. Four participants were excluded from the analysis based on poor perceptual performance (see Procedure). All participants were free from neurological and visual impairments. Experiment 1 was conducted with the approval of the University of Edinburgh Psychology Research Ethics Committee. Participants provided written, informed consent, and were financially compensated (at £10 per hour).

Sample size considerations

SI in overt behavioural tasks is an extremely robust phenomenon, usually observed in every participant. Our previous studies of SI, like those of other researchers, have used relatively few participants (6-10), but many trials per condition (≥ 240) (e.g. Buonocore and McIntosh 2008; Buonocore and McIntosh 2012; Buonocore and McIntosh 2013; Buonocore et al. 2016; Buonocore et al. 2017c). These large trial numbers per condition are required for the estimation of SI parameters from a full distributional analysis of saccadic latencies; far fewer trials are necessary to estimate global performance measures (e.g. average latency).

The present covert experiment was designed as an exploratory follow up to our overt saccadic study, in which the addition of a flash was found to make saccadic reorienting more successful (Buonocore et al. 2017c). This effect was observed for all six participants in that study, with the flash increasing the probability of successful reorienting by an average of 10 percentage points (SD 3.4). This is a standardised effect size (dz) of 2.9, a huge effect statistically. Given uncertainty over how this effect would translate into orientation discrimination accuracy in a covert task in the present design, we did not perform a formal *a priori* analysis, but we aimed to at least double the sample size of our prior study. Eighteen participants were initially tested in Experiment 1, providing a final group of 14; no data analysis was conducted prior to termination of testing, except to determine participant exclusions.

Experiment 1 – Apparatus and stimuli

Stimuli were generated using MATLAB R2017b (MathWorks, Inc.) and Psychophysics Toolbox 3.0.14 (Brainard 1997). All the stimuli were in grey scale on a grey background (20.6 cd/m^2), presented on a 19-inch CRT monitor with a refresh rate of 75 Hz (13.3 ms temporal resolution). Participants were seated in front of the monitor at a viewing distance of 79.5 cm with their head on a chinrest and their eyes aligned with the centre of the

screen. The fixation point was a small white dot of 0.05 degrees radius (84.9 cd/m^2). The

cue was a filled white circle of 0.2 degrees radius at full contrast (84.9 cd/m^2), at an

eccentricity of 10 degrees from the fixation point and 1.5 degrees above the centre of the

possible probe location on that side (e.g. Pestilli and Carrasco 2005). The probe stimulus

was a tilted Gabor patch with a radius of 0.55 degrees, a contrast of 0.2 and a spatial

frequency of 1.78 c/deg, 10 degrees to the left or right of the fixation point. Eye

movements were monitored with a tower-mount EyeLink 1000 system tracking the right

eye at a sampling rate of 1000 Hz. Manual responses were recorded on a two-button

response pad. The room was dark, except for the display monitor, and the operator

monitor located behind the participant and facing away from them.

Each participant completed two testing sessions on separate days lasting about one

hour each including breaks. The first testing session involved a practice block, a QUEST

procedure for orientation threshold (repeated up to three times), and 38 blocks of

experimental trials. The second testing session comprised 38 blocks of experimental trials.

Experiment 1 – Procedure

At the beginning of each block, a 9-point calibration was conducted. Calibrations were

repeated if the average error across all points was greater than 0.5 degrees, and after

every 200 trials. In both the QUEST and the main experimental trials, participants were

instructed to fixate the fixation point throughout the trial and to report the orientation of the

Gabor patch (left or right tilted) by pressing the button under their left or right index finger

respectively. We used discrimination of orientation, rather than spatial detection of the

probe location, as a measure of change in sensitivity across the different conditions. In this

way, the spatial propriety of the exogenous cue (location) was orthogonal to the features

of the stimulus (orientation). Speed of responding was not emphasised. If participants

moved their eyes outside of a fixation window of 3 degrees radius, the trial was aborted

and randomly reshuffled into the remaining trial sequence.

In the first session, to familiarise with the basic task, participants first performed a

practice block of 16 trials in which the onset of a cue was followed after a random delay by

the onset of a probe at the same spatial location (congruent cue condition). The practice

block was followed by a QUEST staircase block (Watson and Pelli 1983), a commonly

used adaptive method for psychometric threshold. The QUEST block was used to identify

a suitable orientation per participant that produced around 75% correct responses. Based

on prior estimates of the psychometric function and the responses to different tilt

intensities a final threshold was estimated.

The trial sequence for the QUEST followed the same structure as the subsequent

experimental trials (description below), for a maximum of 80 trials, but only congruent cue

conditions were used. The QUEST parameters were set to a 75% performance criterion, a

beta of 1.5, and a grain of 1. If the estimated threshold orientation was greater than 15

degrees, the QUEST block was run again up to a maximum of three times. Participants

were excluded from the main experiment if they still had an outcome greater than 15

degrees after the third QUEST block, or if their average performance in experimental trials

was below 60% at the end of the whole first session. Four participants were excluded on

this basis.

The experimental task was an adaptation of the classic cue Posner paradigm for

covert perceptual judgements, modified to mimic the structure of Experiment 3 in

Buonocore et al. (2017c) for overt eye movement responses (Fig. 1). Each trial began with

the onset of a white fixation dot. After a random interval between 800 ms and 1200 ms, the

cue was presented for 53.3 ms to the left or right of fixation. The probe was presented for

106.6 ms after a cue-target onset asynchrony (CTOA) selected on every trial from a

uniform random distribution between 120 and 306.6 ms, with equal numbers of trials at the

cued location (*congruent cue condition*) or at the uncued location (*incongruent cue*

condition). On half of the trials, a black flash (0.34 cd/m^2) covering the bottom and top

thirds of the screen was presented for 13.3 ms simultaneously with the onset of the probe.

This technique was introduced by Reingold and Stampe (2002) as a way to induce

saccadic inhibition without interfering with the localization of a saccade target. It was

recently adapted to saccade tasks requiring a concurrent perceptual response (Buonocore

et al. 2016; Buonocore and Melcher 2015), establishing a lack of masking effects of this

remote flash on probe perception.

Participants completed 76 experimental blocks, for a total of 1216 valid trials,

across the two test sessions. Within each block, there were 16 valid trials, created by the

factorial combination of cue type (congruent, incongruent) by flash (no flash, flash) by

probe side (left, right) by probe tilt (left, right). The first two factors (cue, flash) were of

theoretical interest, whilst the latter two factors (probe side and tilt) were not, and were

simply counterbalanced. The CTOA on each trial was drawn randomly from a uniform

distribution between 120 and 306.6 ms. For data analysis, four CTOA categories were

formed by grouping the CTOA in bins of 50 ms (centred at 140, 190, 240 and 290 ms). On

average, each participant completed 76 trials per CTOA category for each combination of

cue type (congruent, incongruent) by flash (no flash, flash).

Figure 1. Trial scheme. Each trial started with the onset a fixation point. After a random interval
between 800 to 1200 ms, a white dot cue was presented for 53.3 ms at 10 degrees to one side of
fixation. After a random interval between 120 to 306.6 ms after cue onset (CTOA), the probe was
presented for 106.6 ms either at the same (congruent) or the opposite (incongruent) location as the
cue. On half of the trials, a flash appeared with the onset of the probe covering the bottom and top
thirds of the screen for 13.3 ms. Here we show the example of an incongruent plus flash trial.
Stimuli are not drawn to scale. Experiment 2 was identical to Experiment 1 but now in half of the
trials the cue could be presented at the centre of the screen. We also used fixed CTOAs of 47,
82.3, 105.8 and 200 ms. The duration of the cue was slightly shorter than in Experiment 1, 35.2
ms, to allow an early CTOA of 47 ms. Flash duration was 11.3 ms and probe duration was 90 ms.
258

Experiment 1 – Data processing and analysis

Eye movements, saccades and microsaccades, were detected automatically based on

velocity and acceleration thresholds of 15 deg/s and 450 deg/s^2 respectively, then all trials

were manually inspected and adjusted by an experienced researcher (AB). Samples in

which the eye signal was unstable, lost or affected by blinks were flagged as "bad data"

2.5%. From the trials with good eye movement signals, we further excluded trials with

manual reaction times less than 200 ms (4.1%) or more that 3.5 standard deviation above

each participants average response latency (1.7%), leaving a total of 15682 good trials. To

remove possible confounds in our results due to microsaccades (e.g. Hafed 2013), we

also removed all trials with microsaccades detected between 200 ms before cue onset and

200 ms after probe onset (1604 trials removed, 10.22% of the good trials). This defined an

interval of stable fixation of minimum duration 515 ms and maximum duration 715 ms,

depending upon the CTOA. We checked if microsaccades were equally balanced in the congruent and incongruent cue condition and we found that on 47.5% microsaccades were executed in the congruent and 52.5% in the incongruent cue condition. Similarly, 48.1% microsaccades were detected in no flash trials and 51.9% in flash trials. For the remaining trials (N = 14078), incorrect responses were coded as 0 and correct responses as 1. We then used a mixed-effects logistic regression to test the influence of Cue type (congruent, incongruent), Flash (no flash, flash), and CTOA (four bins centred on 140, 190, 240, 290 ms) on perceptual performance.

To asses whether our stimuli were inducing inhibition, we also analysed microsaccade rates relative to both cue and probe onset. For each participant, we calculated for every trial the absolute frequency of microsaccades in bins of 20 ms within a time window spanning 200 ms before cue/probe onset and 500 ms after cue/probe onset. The absolute frequency was averaged across trials and converted to microsaccade rate by dividing it for the bin size (and transformed in seconds). Each microsaccade rate curve was then averaged across participants.

All data pre-processing and statistical analyses were conducted with custom scripts in MATLAB R2019a (MathWorks, Inc.) and R (R Core Team 2020) (lme4, lmerTest). The entire dataset is uploaded in the Open Science Framework archive at the following link:

https://osf.io/9fnh4/?view_only=14b0ae67d49c4c1aa4898f66f20b473b

Experiment 1 – Results

The present experiment follows up the recent findings in which we reported that inducing saccadic inhibition during oculomotor programming in a variant of a double-step paradigm (Becker and Jürgens 1979) could facilitate saccade reorienting behaviour (Buonocore et al. 2017c). Based on the findings for overt saccade responses, we asked if covert

orienting, that we conceptualize in the framework of premotor preparation (Rizzolatti et al. 1987; Sheliga et al. 1994; Sheliga et al. 1995), could be subjected to similar inhibitory processes triggered by flash onset and which repercussions such inhibition of covert orienting would have on perceptual judgements. In our attentional version of the paradigm, cue onset, rather than accumulating build-up activity to trigger an eye movement, would initiate a similar build-up process that might increase visual sensitivity at its location. Usually this "shift in attention" is maximal after 100 ms relative to cue onset (e.g. Cheal and Lyon 1991) and it is followed by a reorienting of attention that decrease sensitivity at that location around 200 ms after cue (Cheal and Chastain 1999; Cheal et al. 1998; Posner and Cohen 1984; Pratt and Abrams 1999). Our expectation is that flash onset might hinder covert orienting capabilities in a specific time window that depends on the state of the attentional process, similar to how the sudden presentation of a visual stimulus can reset saccade programming only if it is presented at a critical time of the motor program, about 90 ms before its execution (Buonocore and McIntosh 2008; Reingold and Stampe 1999; Reingold and Stampe 2002). To test this idea, we ran a mixed-effects logistic regression to investigate how perceptual response accuracy (0,1) was modulated by *Cue* type (congruent, incongruent), *Flash* (no flash, flash), *CTOA* (140 ms, 190 ms, 240 ms, 290 ms, representing bin center) and their interactions. The model is summarized in Equation 1 in Wilkinson notation (Wilkinson and Rogers 1973):

$$\text{Eq. 1}: Response = Cue * Flash * CTOA + (1 \mid Subjects)$$

We found a strong main effect of *Cue* (β = -0.166, 95% CI = [-0.245, -0.086], t = -4.083, *p* = 0.001) confirming better orientation discrimination for congruently cued than for incongruently cued probes (Cameron et al. 2002). Overall, the cueing effect corresponded to an increase of about 3.5% in discrimination performance when cue location matched

the probe location. For clarity, in Figure 2A we show the mean accuracy for each condition

across all CTOAs. From the figure it is clear that on average the congruent cue conditions

(green and blue lines) produced better orientation discrimination that the incongruent cue

condition (yellow and purple lines). The analysis also revealed an effect of *Flash* (β = -0.1,

95% CI = [-0.177, -0.018], t = -2.406, *p* = 0.016) and an interaction between *Flash* and

CTOA such that perceptual performance was disrupted in flash trials (Fig. 2A, blue and

purple lines) compared to no flash ones (Fig. 2A, green and yellow lines), when the cue

preceded the probe with a delay of 190 ms (β = -0.254, 95% CI = [-0.469, -0.039], t = -

2.314, *p* = 0.021) and 240 ms (β = -0.244, 95% CI = [-0.476, -0.012], t = -2.0636, *p* =

0.039). Figure 2B illustrates this interaction, plotting the difference between flash and no

flash trials (irrespective of cue). Perceptual discrimination in the middle bins of the CTOA

range was reduced by about 4%.

The data for each condition in Figure 2A suggests that the effect of the flash may be

mostly driven by the incongruent cue condition (purple line), rather than the congruent cue

condition (blue line). We explored this pattern by running two separate mixed-effects

logistic regressions to test the influence of Flash and CTOA, and their interaction, on the

congruent and incongruent trials. There were no significant effects for the congruent cue

condition at any of the CTOAs, while there was an interaction in the incongruent cue

condition with a strong effect of the flash at 190 ms (β = -0.331, 95% CI = [-0.632, -0.03], t

= -2.155, *p* = 0.031) and 240 ms (β = -0.401, 95% CI = [-0.728, -0.074], t = -2.404, *p* =

0.016). While this analysis does not formally establish a three-way interaction, it is

consistent with a stronger inhibitory influence of the flash in incongruent trials, relatively

specific to the middle two time bins (190, 240 ms).

Figure 2. Perceptual performance in Experiment 1. **(A)** Congruent cues (green and blue lines) lead
to better performance than incongruent cues (yellow and purple lines). Flash onset (blue and
purple lines) interfered with orienting behaviour reducing accuracy relative to trials in which the
flash was absent (green and yellow lines respectively). The effect was more pronounced when the
cue was presented 190 to 240 ms before probe onset and for the incongruent condition (purple
line). **(B)** Relative effect of the flash calculated as accuracy for flash trials minus no flash trials. A
clear decrease in perceptual discrimination is visible for the middle range CTOAs. Error bars
represent one standard error of the mean.

This pattern is evident in Figure 3, where we show the discrimination accuracies of our

individual participants, contrasting the no flash against flash condition at each CTOA for

the congruent (Fig.3A, top row panels) and the incongruent (Fig.3B, bottom row panels)

cue conditions. The figure clarifies that perceptual performance in the congruent cue

condition was not strongly affected at any CTOA. On the other hand, in the incongruent

cue condition at the mid CTOAs there was a clear shift in perceptual performance

(decrease in performance observed in 11 and 12 out of 14 participants respectively).

Figure 3. Raw data points of individual participants contrasting the no flash condition against flash condition at each CTOA for the congruent **(A)** and incongruent **(B)** cue condition. Dotted lines represents the unity slope line. At CTOAs of 190 ms and 240 ms (red arrow) there was a strong decrease in discrimination performance in the incongruent plus flash cue condition. Early and late CTOAs did not show any difference for trials with or without flash.

Experiment 1 – Microsaccade analysis

In the previous section we showed that covert orienting was perturbed only when a flash was presented 190 to 240 ms after cue onset, in the total absence of eye movements. In the following analysis, we asked if the stimuli used in our paradigm were actually capable to inhibit overt responses and induce SI in a similar time period. To answer this question, we looked at microsaccades, the tiny eye movements (smaller than 1 degree) that continuously happen during covert attention tasks (Engbert and Kliegl 2003; Laubrock et al. 2005; Pastukhov and Braun 2010).

We detected a total of 3363 microsaccades in 2785 trials in a time period of 200 ms before cue onset to 500 ms after probe onset (1.2 microsaccades per trial within this time period). In Figure 4A, we aligned microsaccades to cue onset and, as expected, a strong microsaccadic inhibition took place between 200 to 400 ms after the cue (Buonocore et al. 2017a; Malevich et al. 2021; Rolfs et al. 2008; Tian et al. 2016; Tian et al. 2018),

suggesting that the cue was strongly inhibiting overt orienting responses. We note that

microsaccade rate was decreasing with a negative trend even before cue onset. This

"microsaccade suppression" during the pre-cue interval indicates the effort of participants

make to maintain fixation while waiting for the trial stimuli (e.g. Denison et al. 2019). In

Figure 4B we aligned the same data to probe onset and calculated microsaccade rate

separately for both no flash (green line) and flash (blue line) conditions. Overall, it is clear

that microsaccadic inhibition induced by the cue was already so strong that microsaccades

did not have time to recover by the time the probe was presented. Nonetheless, the

inhibition was further enhanced 100 ms after probe onset and it was extended up to 300

ms thereafter (100 ms after the minimum reaction time allowed). Comparing microsaccadic

rate in no flash (green) and flash (blue) conditions suggests that the flash did not have any

additional measurable effect on the inhibition since it was already at ceiling. It is important

to note that the lack of observable influence of the flash on overt behaviour is due to

microsaccadic inhibition being total, but this does not mean that the flash should not have

an influence on covert activation. Rather the fact that even the cue and probe were able to

cause microsaccadic inbhibition implies that the much more salient flash should provoke

even greater effects. Overall, the data confirm that our stimuli were indeed very powerful in

inducing microsaccadic inhibition in a time period largely overlapping with the inhibition of

covert orienting reported above.

Figure 4. Microsaccadic inhibition in Experiment 1. **(A)** Microsaccade times aligned to cue onset.
Each dot represents a microsaccade in an interval between 200 ms before to 500 ms after cue
onset. Each colour band represents microsaccades for one participant. About 200 ms after cue
onset, microsaccades are almost completely abolished. The black curve represents the averaged
microsaccade rate across participants. The grey shaded area represents the time at which we
recorded inhibition of covert orienting in absence of eye movements **(B)** Same as **(A)** but for
microsaccades aligned to probe onset. Microsaccade rate is calculated separately for no flash
(green line) and flash (blue line) conditions. Microsaccadic inhibition reached ceiling after the
presentation of the probe alone, rendering difficult to see any clear additional effect of the flash.

Experiment 1 – Interim discussion

The results of Experiment 1 suggest that covert orienting is inhibited by the flash, but only

at certain time points, when the flash is presented between 190 to 240 ms after cue onset.

This suggest that the only the reorienting component of the attentional shift, i.e. when

attention is directed away from the previously cued location, was sensitive to the flash

interference. In particular, we report that the effect was present only in the incongruent cue

condition, when such reorienting process was exaggerated by the need to reorient

attention to the opposite side of the visual field relative to the cue. The fact that we do not

find any effect in the congruent cue condition at any of the CTOA suggests that the covert

shift of attention was already completed by the time we presented the flash (e.g. Cheal

and Lyon 1991) and attention was sufficiently allocated at the cue location rendering the

discrimination process impervious to flash presentation. These results parallel the case in

overt orienting in which a flash is presented at a time close to saccade onset, when the

motor program for a saccade has passed a 'point-of-no-return', and is too advanced to be

cancelled or delayed (Buonocore et al. 2016). One question that arises for the present

covert case is whether the flash presented at the earliest CTOA (140 ms) completely

missed the initial component of the orienting shift, not allowing any interference to take

place, or if the early stage of covert orienting is somehow more resistant to inhibition.

To clarify at which stage of the attentional shift it is possible to inhibit orienting, we

planned Experiment 2 where we tested the effect of the flash at the very early stages of

the covert shift as well as at a later stage when reorienting is expected. Experiment 2

focused on the congruent cue condition only, by using a 100% informative cue (as

opposed to 50% informative cues of Experiment 1). This design aim to mimic a condition in

which the onset of a cue is immediately followed by a reflexive saccade. In the covert

case, the cue acts as the stimulus to start attentional allocation to a particular point in

space, similarly to a visual target starting the build-up activity for an eye movement. The

subsequent flash aims to interfere with the ongoing premotor program. By using an early

range of CTOAs compared to Experiment 1, we targeted an earlier window during the

covert attentional shift.

Experiment 2 – Material and Methods

Twelve participants (8 female), aged between 20 and 35 years old (mean = 25, *SD* = 4.2),

were included in the data analysis of Experiment 2. Two participants were excluded from

the analysis based on floor and ceiling effects (see *Experiment 2 – Procedure* below). All

participants were free from neurological and visual impairments. Experiment 2 was

conducted with the approval of relevant ethics committees at the Medical Faculty of

Tuebingen University and of the University of Bielefeld. Subjects provided written,

informed consent, and participated either as unpaid volunteers or for financial
compensation (€10 per testing session).

Sample size considerations

Experiment 2 was conducted during COVID restrictions in 2020-21, which made
recruitment and testing opportunities extremely limited. We aimed to test equivalent
participant numbers to Experiment 1, and managed to test 14 participants between two
laboratories, though only 12 participants remained after exclusions. This was the
maximum achievable sample size under the conditions that existed at the time.

Experiment 2 – Apparatus and stimuli

Experiment 2 was similar to that for Experiment 1 but run in a different laboratory
(Physiology of Active Vision Laboratory, Werner Reichardt Centre for Integrative
Neuroscience, University of Tuebingen), with minor differences in setup[1]. The monitor was
a 19-inch CRT monitor with a refresh rate of 85Hz (11.8 ms temporal resolution)
positioned at a viewing distance of 57 cm. Stimuli were presented on a grey background
(42.2 cd/m^2). The fixation point was 0.01 degrees radius (97.3 cd/m^2), and the cue (74.9
cd/m^2) was presented either 1 degrees above the fixation point or at an eccentricity of 10
degrees from the fixation point and 1 degrees above the centre of the possible probe
location on that side. All other stimulus parameters were identical to those in Experiment 1.
Eye movements were monitored with a desktop mount EyeLink 1000 system, tracking the
right eye at a sampling rate of 1000 Hz. Manual responses were recorded with two (left or
right) of the five-button on a standard response pad (RESPONSEPixx, VPixx Technologies

---

[1] Five out of the fourteen participants were collected at the University of Bielefeld with the setup matched as closely as possible, but with a different monitor refresh rate (100 Hz).

Inc.*).* Participants completed one testing session with one practice block and 16 blocks of

experimental trials.

Experiment 2 – Procedure

Experiment 2 procedure was similar to Experiment 1 except for a few critical modifications.

After calibration, the experiment started with a practice block of 64 trials and no QUEST

procedure was required. Instead, we used a probe stimulus with an orientation of either

84, 85, 86 degrees (right tilt) or 94, 95, 96 degrees (left tilt) chosen at random at the start

of each trial. After a random interval between 800 ms and 1200 ms, the cue was

presented for 35.2 ms either 10 degrees to the left or right of fixation or at the centre of the

screen one degree above of the fixation point. The probe was presented for 90 ms after

one of four randomly selected CTOAs: 47, 82, 106 and 200 ms, with equal numbers of

trials at the cued location (*congruent cue condition*) or at the neutral location (*neutral cue*

condition). The neutral cue condition was introduced as a control for any generalised

temporal "warning effect" of the congruent cue (Coull and Nobre 1998; Hackley and Valle-

Inclán 2003). On half of the trials, a black flash covering the bottom and top thirds of the

screen for 11.3 ms was onset simultaneously with the probe.

Participants were asked to complete 16 blocks of experimental trials within the 75

minutes testing session. Within each block, there were 64 valid trials, one for each

combination of cue type (congruent, incongruent) by flash (no flash, flash) by probe side

(left, right) by CTOA (47, 82, 106 and 200 ms) by probe tilt (left, right). The first three

factors (cue, flash, CTOA) were of theoretical interest, whilst the latter two factors (probe

side and tilt) were not. Each participant completed 64 trials per CTOA for each

combination of cue type (congruent, neutral) by flash (no flash, flash), except that some of

the participants did not finish the last block due to time constrains (across subjects a total

of 182 trials were not completed). Two of an original 14 participants were excluded from

the analysis because their overall mean performance was either at floor (48.6%) or at

ceiling (93.9%).

Experiment 2 – Data processing and analysis

We collected a total of 12112 trials across all included participants in Experiment 2. We

used identical criteria to Experiment 1 for trial exclusion. We rejected 1% of trials because

of "bad data", 0.17% trials with manual reaction times less than 200 ms and 1% of trials

with manual reaction times more than 3.5 standard deviation above the participant's

average. We also remove 2.3% of trials in which critical stimulus flips were not properly

timed with the monitor refresh rate. Our sample was of 11578 trials, from which we

removed 1699 trials because of microsaccades (14.7%), leaving 9879 of microsaccade

free trials in an interval between -200 ms before cue onset and 200 ms after probe onset

(minimum saccade free interval equal to 250 ms and maximum saccade free interval equal

to 400 ms). From the total of the excluded microsaccades, 50% belonged to the congruent

and 50% in the incongruent cue condition. Similarly, 52.6% microsaccade were detected

in no flash trials and 47.3% in flash trials. We ran a mixed-effects logistic regression to test

the influence of Cue type (congruent, neutral), Flash (no flash, flash), and CTOA (47, 82,

105 and 200 ms) on perceptual performance. Paired-sample t-test were used to follow-up

statistical analysis with an alpha level of 0.05. Microsaccade analysis was identical to

Experiment 1.

All data pre-processing and statistical analyses follow the procedure described in

Experiment 1 and the entire dataset is uploaded in the Open Science Framework (see

Experiment 1 - Data processing and analysis).

Experiment 2 – **Results**

In Experiment 2 we tested whether the inhibitory effect of the flash could occur within the

range relevant to early orienting or if it was only effective in the later (reorienting) range. To

do so, we tested orientation discrimination for probes presented at congruent cue locations

in a time window between 50 to 200 ms. To control for possible effects driven only by the

temporal aspects of the cue, rather than its location in space, we also tested a neutral cue

condition in which a spatially non-informative cue was presented centrally. Based on the

results of Experiment 1, we expected to see modulations of discrimination performance at

the time of reorienting, when attention moves away from the cued location, and/or at early

CTOAs, when facilitation is expected. To explore these patterns, we ran a mixed-effects

logistic regression with *Cue* type (congruent, neutral), *Flash* (no flash, flash), *CTOA* (47

ms, 82 ms, 106 ms, 200 ms), and their interactions (same model as Eq. 1).

The first significant effect was an interaction between *Flash* and *CTOA* (Fig. 5A). In

flash trials there was an enhancement of about 4% in discrimination performance when the

probe was presented 47 ms after the cue compared to later CTOAs (CTOA 82 ms: β = -

0.036, 95% CI = [-0.32, 0.247], t = -0.252, *p* = 0.801 ; CTOA 106 ms: β = -0.287, 95% CI =

[-0.566,-0.008], t = -2.017, *p* = 0.044; COTA 200 ms: β = -0.242, 95% CI = [-0.528, 0.042],

t = -1.67, *p* = 0.095). Since the interaction was irrespective of cue type, i.e. congruent or

neutral, it suggests that the flash was having a general "warning effect", increasing visual

sensitivity when presented immediately after the cue (Coull and Nobre 1998; Hackley and

Valle-Inclán 2003).

We also found an interaction between *Cue* and *CTOA* (Fig. 5B). While performance

in the neutral cue condition was stable at about 80% (grey line), the congruent cue

condition (red line) was modulated across CTOAs with a decrease in discrimination

performance of about 4% taking place 200 ms after cue onset ms (β = 0.298, 95% CI =

[0.013, 0.583], t = 2.049, *p* = 0.041). Numerically, perceptual performance was slightly

better at CTOA 82 ms, but overall we did not see a significant facilitation effect at any of

the early CTOAs. This result suggested that in our paradigm the attentional shift triggered

by the cue at early CTOAs was not strong enough to boost performance significantly for

this task. On the other hand, attentional allocation started reducing at cue location at

around 200 ms after cue onset, compatible with previous reports (Cheal and Chastain

1999; Cheal et al. 1998; Posner and Cohen 1984; Pratt and Abrams 1999) and our results

from Experiment 1.

Figure 5. Perceptual performance. **(A)** At the early CTOA of 47 ms, performance is boosted by the
presence of the flash (blue lines) compared to no flash trials (green lines) and irrespective of the
cue type **(B)** Neutral cues (grey lines) do not modulate perceptual performance across CTOAs. On
the other hand, congruent cues (red line) have an impact on discrimination performance for the
late CTOA of 200 ms, where visual sensitivity for the incoming probe is impaired. **(C)** At the time of
reorienting, i.e. 200 ms, the presentation of the flash (blue line) affects perceptual discrimination
capabilities by impeding reorienting and maintaining visual sensitivity higher compared to no flash
trials (green line).

The significant decrease in discrimination performance observed in the congruent

cue condition at 200 ms after cue onset (Fig. 5B) was suggestive of attention moving away

from the cue location (reorienting). This result motivated us to explore in more detail if the

flash could interfere with this part of the orienting process, as observed for the incongruent

cue condition in Experiment 1. To do so, we computed a measure of cue-benefit by

subtracting accuracy in the neutral cue condition from that in the congruent cue condition

in both no flash and flash trials at each CTOA. Paired-sample t-test showed that only at

200 ms was there a difference between flash and no flash conditions (t(11) = -2.94,

p=0.013), with discrimination performance being about 6% higher in flash trials (Fig. 5C).

This result supports the hypothesis that the flash has a temporal window of action in which

it can inhibit attention moving away from cue location, similarly to the result reported in

Experiment 1.

Experiment 2 – Microsaccade analysis

We detected a total of 3454 microsaccades over 3133 trials in a time period of 200 ms

before cue onset to 500 ms after probe onset (1.1 microsaccades per trial within this time

period). Similarly to Experiment 1, microsaccade rate (black line) was completely disrupted

about 100 ms after cue onset (Fig. 6A). When data were aligned to probe onset (Fig. 6B)

and microsaccade rate was split by flash condition (green: no flash, blue: flash), despite

microsaccadic rate being close to zero for both conditions, there is indication of a slightly

larger inhibition for flash trials. As in Experiment 1, microsaccadic inhibition after the probe

was long lasting, with the rebound period starting only at around 300 ms after probe onset.

This results confirms that our stimuli could abolish any subsequent eye movement in a

time period compatible with the covert effect recorded in absence of eye movement.

Figure 6. Microsaccadic inhibition. Same colour conventions as Figure 5. **(A)** After cue onset,
microsaccades are almost completely abolished (black line). The vertical grey line indicates the

CTOA were we recorded the inhibition of covert reorienting. **(B)** Same as **(A)** but for microsaccades aligned to probe onset. There is a slightly stronger inhibition in flash (blue line) compared to no flash (green line) trials.

Experiment 2 – Interim discussion

The results from Experiment 2 confirm that perceptual discrimination can be impaired by the flash during the reorienting stage of the attentional shift. At the early CTOAs we did not observe any specific effect of the flash in the congruent cue condition. This result is surprising because within this time range covert orienting should mostly overlap with the preparation of an eye movement to a lateralised cue, similarly to what happens in pre-saccadic shifts of attention (Buonocore et al. 2017b; Deubel and Schneider 1996; Hoffman and Subramaniam 1995; Kowler et al. 1995; Li et al. 2016; Li et al. 2019; Ohl et al. 2017; Remington 1980; Rolfs and Carrasco 2012; Shepherd et al. 1986). By presenting a flash at this stage of motor programming should be possible to strongly interfere with the build-up activity at the cued location, as we clearly see in SI paradigms. One possible explanation is that the early cue was not powerful enough to induce strong orienting toward the cue location, as indicated by the absence of a facilitation effect for early cues. On the contrary, we see a general facilitation effect in flash trials for CTOA equal to 47 ms. The fact that we see such facilitation speaks instead of a temporal "warning effect" of the flash, independent of cue type, that increased visual sensitivity for the incoming probe (Coull and Nobre 1998; Hackley and Valle-Inclán 2003). This facilitation might have overshadowed any inhibitory signal associated with flash presentation.

The effect recorded at CTOA equal 200 ms in the congruent cue condition might be the result of the flash introducing a small delay in the process of attentional reorienting. Consequently, a higher level of visual sensitivity was maintained at the cued location for a longer time, increasing perceptual performance. This result is consistent with the flash interfering with the incongruent cue conditions of Experiment 1. In this case, visual sensitivity was maintained higher at the incongruent cue location, leading to a decrease in

performance for a probe presented at the opposite cue side. Taken together, Experiment 1

and 2 suggest that the later reorienting component can be effectively perturbed by a

generalised flash, while the early orienting component of the attentional shift may be less

susceptible to inhibition.

General conclusions

In the present manuscript, we uncovered a new phenomenon within a classic cueing

paradigm (Cameron et al. 2002; Posner 1980) in which the visual discrimination of a probe

stimulus was deteriorated by the simultaneous presentation of a brief flash event, but only

when the flash was preceded by the exogenous cue between 190 to 240 ms. In

Experiment 1, the effect was particularly emphasised for the condition in which cue

location was incongruent with the location of the incoming probe impeding reorienting

toward the probe and decreasing perceptual discrimination. In Experiment 2, a similar

inhibitory effect was found in a congruent cue condition, by inhibiting reorienting away from

the cue and maintaining performance slightly higher. Taken together, we suggest that the

build-up of attentional resources at a cue location has a specific temporal window in which

an inhibitory signal can interfere, leading to an inhibition of covert reorienting. However,

the exact time course and aspects of the covert orienting process subject to interference

require further investigation.

We assume that the mechanisms behind the inhibitory process might share similar

characteristics with the well-known phenomenon of SI for overt responses in which a flash

(or other transient event) interferes with the preparation of any eye movement that is due

about 90 ms later (Bompas and Sumner 2011; Buonocore and McIntosh 2008; Buonocore

and McIntosh 2012; Edelman and Xu 2009; Reingold and Stampe 1999; Reingold and

Stampe 2002). Despite the differences in the time course of SI and the inhibition of covert

orienting reported here, we suggest that the two phenomena might rely on the same

neural circuitries, in accordance with the mechanistic overlap between covert and overt

orienting proposed by the premotor theory of attention (Rizzolatti et al. 1987; Sheliga et al.

1994; Sheliga et al. 1995). From a neuronal perspective, the onset of the flash might

decrease the firing rate of the premotor build-up SC neurons coding for saccades to a

particular location in space (Buonocore et al. 2021; Dorris et al. 2007; Munoz and Istvan

1998) and consequently introduce a small delay in the premotor plan. The source of the

inhibitory signal might be an increased in activation of OPNs following flash onset

(Buonocore et al. 2020; Buonocore et al. 2017a), which is then broadcast to the SC and

other upstream visual and visuo-motor areas. Higher order visual areas tuned for stimulus

features such as orientation (e.g. V1, Hubel and Wiesel 1959) might then be affected by

the temporal delays introduced in the early premotor process for target selection, affecting

perceptual discrimination.

One might ask why if the neural mechanisms are shared between covert and overt

processing there are temporal differences between the two inhibitory effects. We suggest

that perhaps overt SI has an intrinsic "all-or-nothing" nature, because a saccade either is

or is not launched within a specific time window. Any depression of the premotor activation

can prevent the saccade from launching at the expected time, leading to a large SI effect

90 ms after flash onset. On the other hand, covert orienting might depend on a more

gradual readout of premotor activity, which transiently changes following flash onset but

continues to accumulate until a discrimination is made. For this covert process to provide

sufficient cumulative activation for discrimination could require a longer integration period

compared to the process to trigger saccade execution. Moreover, while SI can be almost

entirely modelled within a small network within subcortical oculomotor areas such as SC

and the lower brainstem, visual discrimination is a process that involves higher order

cortical visual area that contribute to reshape the temporal profile of the effect. This

account is speculative, but the more general argument is that superficial differences in

timing do not necessarily preclude an underlying identity of the covert and overt inhibition

phenomena, because the measured consequences may be strongly shaped by the very

different readout processes associated with saccade execution and perceptual

discrimination.

Based on the general SI framework (Reingold and Stampe 2002) and our previous

findings on saccades (Buonocore et al. 2017c), we expected flash onset to interfere with

covert orienting behaviour in a specific time window, inducing an SI-like inhibition effect.

Beyond this, if the pattern of covert orienting was strictly following that of overt orienting as

observed in our previous study, we would have expected this interruption to improve the

ability to re-orient to the opposite (uncued) location, giving a relative enhancement of

perceptual discrimination for incongruently cued targets. In this respect, the effect would

have been the perceptual counterpart to the higher rate of successful reorienting saccades

observed when SI was boosted in "flash-jump" trials in the overt task. The other prediction

was that perceptual performance at congruent cue locations might have been slightly

impaired. Although our results confirm that the effect of the flash depended upon the stage

of evolution of the covert shift, the direction of the effect was contrary to our initial

expectation. Rather than impairing early orienting behaviour, the flash impaired later

reorienting. This finding can perhaps be informed by the idea that reorienting following

interruption after a transient event carries a small temporal cost (see reorienting latency for

Experiment 3, Table 3 in: Buonocore et al. 2017c). In our covert paradigm the flash might

have introduced a similar delay during covert orienting, requiring more time to redirect to

the probe location. For example, in Experiment 1, given the brevity of the probe stimulus to

be discriminated (106 ms), this small delay may have been enough to reduce resources at

the probe location and consequently deteriorate sensitivity for the probe stimulus (Salinas

and Stanford 2018). That is, while orienting (to the cued location) in our design might have

been mostly or wholly completed within 100 ms (e.g. Cheal and Lyon 1991), and so

invulnerable to interruption by a flash, reorienting (in incongruent trials) occurred later, and

was vulnerable to interruption, reducing the opportunity to process the brief probe

stimulus. This mechanism can also accommodate the outcome of Experiment 2, where the

flash inhibited reorienting in the congruent cue condition. In this case, the orienting

component of the shift was also completed by 100 ms and attentional allocation might

have started moving away from the cued location. Introducing a small interruption in this

reorienting process by presenting a brief flash might have maintained more resources at

the cued location, increasing perceptual performance.

It is important to emphasise that the interruption effect was restricted to later CTOA

times (190-240 ms in Experiment 1 and 200 ms in Experiment 2), despite the fact that the

flash was always simultaneous with the probe. This rules out the possibility that passive

masking mechanisms (Alpern 1952; Breitmeyer and Ogmen 2000) could account for the

deterioration of probe discrimination; if masking were responsible, then the appearance of

the flash simultaneous with the probe would always lead to the same impairment, across

all experimental conditions. Moreover, in Experiment 2 we found that the flash presented

at early CTOAs enhanced performance, acting as a general "warning" signal for the

incoming probe and increasing the overall sensitivity of the visual system. This result is

also incompatible with possible passive masking effect and speaks in favour of the flash

interference being specific for the attentional state.

Taking together these observations, we can draw some predictions from our data to

apply to future research work on inhibition of covert orienting. One main hypothesis is that

the flash would always alter perceptual performance when it occurs during a sensitive

period of the covert process triggered by the cue. The sensitive period could be predicted

by estimating the time at which the covert shift would engage or disengage from the cue.

This feature is in fact similar to the time-locked interference of the flash relative to saccadic

reaction times. Finally, inhibition of covert orienting is then expected to also extend to

paradigms in which pre-saccadic shift of attention are involved, potentially altering the

strong benefits of coupling covert orienting with eye movements. We suggest that this

paradigm may lead to new insights into how the attentional shift develops over time, and

help to investigate the extent to which covert and overt processes share underlying neural

processes.

Conflict of interest

On behalf of all authors, the corresponding author states that there is no conflict of
interest.
740

References

Alpern M (1952) Metacontrast; historical introduction Am J Optom Arch Am Acad Optom
29:631-646
Barbot A, Landy MS, Carrasco M (2012) Differential effects of exogenous and
endogenous attention on second-order texture contrast sensitivity J Vis 12
doi:10.1167/12/8/6
Becker W, Jürgens R (1979) An analysis of the saccadic system by means of double step
stimuli Vision Research 19:967-983 doi:10.1016/0042-6989(79)90222-0
Belopolsky AV, Theeuwes J (2012) Updating the premotor theory: the allocation of
attention is not always accompanied by saccade preparation J Exp Psychol Hum
Percept Perform 38:902-914 doi:10.1037/a0028662
Bompas A, Campbell AE, Sumner P (2020) Cognitive control and automatic interference in
mind and brain: A unified model of saccadic inhibition and countermanding Psychol
Rev doi:10.1037/rev0000181
Bompas A, Sumner P (2011) Saccadic inhibition reveals the timing of automatic and
voluntary signals in the human brain J Neurosci 31:12501-12512
doi:10.1523/JNEUROSCI.2234-11.2011
Brainard DH (1997) The Psychophysics Toolbox Spatial Vision 10:433-436
doi:10.1163/156856897X00357
Breitmeyer BG, Ogmen H (2000) Recent models and findings in visual backward masking:
a comparison, review, and update Percept Psychophys 62:1572-1595
doi:10.3758/bf03212157
Buonocore A, Baumann MP, Hafed ZM (2020) Visual pattern analysis by motor neurons
(Abstract) Computational and Systems Neuroscience (Cosyne) 147
Buonocore A, Chen C-Y, Tian X, Idrees S, Münch TA, Hafed ZM (2017a) Alteration of the
microsaccadic velocity-amplitude main sequence relationship after visual transients:
implications for models of saccade control. Journal of Neurophysiology 117:1894-
1910 doi:10.1152/jn.00811.2016
Buonocore A, Fracasso A, Melcher D (2017b) Pre-saccadic perception: Separate time
courses for enhancement and spatial pooling at the saccade target PLoS ONE
12:e0178902-0178923 doi:10.1371/journal.pone.0178902
Buonocore A, McIntosh RD (2008) Saccadic inhibition underlies the remote distractor
effect Experimental brain research Experimentelle Hirnforschung Expérimentation
cérébrale 191:117-122 doi:10.1007/s00221-008-1558-7
Buonocore A, McIntosh RD (2012) Modulation of saccadic inhibition by distractor size and
location Vision Research 69:32-41 doi:10.1016/j.visres.2012.07.010
Buonocore A, McIntosh RD (2013) Attention modulates saccadic inhibition magnitude
Quarterly journal of experimental psychology (2006) 66:1051-1059
doi:10.1080/17470218.2013.797001
Buonocore A, McIntosh RD, Melcher D (2016) Beyond the point of no return: effects of
visual distractors on saccade amplitude and velocity Journal of Neurophysiology
115:752-762 doi:10.1152/jn.00939.2015

Buonocore A, Melcher D (2015) Interference during eye movement preparation shifts the timing of perisaccadic compression Journal of Vision 15:3 doi:10.1167/15.15.3

Buonocore A, Purokayastha S, McIntosh RD (2017c) Saccade Reorienting Is Facilitated by Pausing the Oculomotor Program Journal of cognitive neuroscience 29:2068-2080 doi:10.1162/jocn_a_01179

Buonocore A, Tian X, Khademi F, Hafed ZM (2021) Instantaneous movement-unrelated midbrain activity modifies ongoing eye movements Elife 10 doi:10.7554/eLife.64150

Büttner-Ennever JA, Cohen B, Pause M, Fries W (1988) Raphe nucleus of the pons containing omnipause neurons of the oculomotor system in the monkey, and its homologue in man. Journal of Comparative Neurology 267:307-321 doi:10.1002/cne.902670302

Cameron EL, Tai JC, Carrasco M (2002) Covert attention affects the psychometric function of contrast sensitivity Vision Res 42:949-967 doi:10.1016/s0042-6989(02)00039-1

Carrasco M (2011) Visual attention: the past 25 years Vision Res 51:1484-1525 doi:10.1016/j.visres.2011.04.012

Cheal M, Chastain G (1999) Inhibition of return: support for generality of the phenomenon J Gen Psychol 126:375-390 doi:10.1080/00221309909595372

Cheal M, Chastain G, Lyon DR (1998) Inhibition of Return in Visual Identification Tasks Visual Cognition 5:365-388 doi:10.1080/713756786

Cheal M, Lyon DR (1991) Central and peripheral precuing of forced-choice discrimination Q J Exp Psychol A 43:859-880 doi:10.1080/14640749108400960

Cohen B, Henn V (1972) Unit activity in the pontine reticular formation associated with eye movements vol 46. Brain research. doi:10.1016/0006-8993(72)90030-3

Coull JT, Nobre AC (1998) Where and when to pay attention: the neural systems for directing attention to spatial locations and to time intervals as revealed by both PET and fMRI J Neurosci 18:7426-7435 doi:10.1523/jneurosci.18-18-07426.1998

Denison RN, Yuval-Greenberg S, Carrasco M (2019) Directing Voluntary Temporal Attention Increases Fixational Stability J Neurosci 39:353-363 doi:10.1523/jneurosci.1926-18.2018

Deubel H, Schneider WX (1996) Saccade target selection and object recognition: Evidence for a common attentional mechanism Vision Research 36:1827-1837 doi:10.1016/0042-6989(95)00294-4

Dorris MC, Olivier E, Munoz DP (2007) Competitive integration of visual and preparatory signals in the superior colliculus during saccadic programming J Neurosci 27:5053-5062 doi:10.1523/JNEUROSCI.4212-06.2007

Edelman JA, Xu KZ (2009) Inhibition of Voluntary Saccadic Eye Movement Commands by Abrupt Visual Onsets Journal of Neurophysiology 101:1222-1234 doi:10.1152/jn.90708.2008

Engbert R, Kliegl R (2003) Microsaccades uncover the orientation of covert attention Vision Research 43:1035-1045 doi:10.1016/S0042-6989(03)00084-1

Fernández A, Li H-H, Carrasco M (2019) How exogenous spatial attention affects visual representation Journal of Vision 19:4-4 doi:10.1167/19.11.4

Hackley SA, Valle-Inclán F (2003) Which stages of processing are speeded by a warning signal? Biol Psychol 64:27-45 doi:10.1016/s0301-0511(03)00101-7

Hafed ZM (2013) Alteration of Visual Perception prior to Microsaccades Neuron 77:775-786 doi:10.1016/j.neuron.2012.12.014

Hafed ZM, Clark JJ (2002) Microsaccades as an overt measure of covert attention shifts Vision Research 42:2533-2545 doi:10.1016/S0042-6989(02)00263-8

Hafed ZM, Yoshida M, Tian X, Buonocore A, Malevich T (2021) Dissociable Cortical and Subcortical Mechanisms for Mediating the Influences of Visual Cues on

834    Microsaccadic Eye Movements Frontiers in Neural Circuits 15
835    doi:10.3389/fncir.2021.638429

836 Hanning NM, Deubel H, Szinte M (2019) Sensitivity measures of visuospatial attention J
837    Vis 19:17 doi:10.1167/19.12.17

838 Hoffman JE, Subramaniam B (1995) The role of visual attention in saccadic eye
839    movements Perception & Psychophysics 57:787-795 doi:10.3758/bf03206794

840 Hubel DH, Wiesel TN (1959) Receptive fields of single neurones in the cat's striate cortex
841    J Physiol 148:574-591 doi:10.1113/jphysiol.1959.sp006308

842 Hunt AR, Reuther J, Hilchey MD, Klein RM (2019) The Relationship Between Spatial
843    Attention and Eye Movements Curr Top Behav Neurosci 41:255-278
844    doi:10.1007/7854_2019_95

845 Keller EL (1974) Participation of Medial Pontine Reticular-Formation in Eye-Movement
846    Generation in Monkey Journal of Neurophysiology 37:316-332

847 Klein RM, Pontefract A (1994) Does oculomotor readiness mediate cognitive control of
848    visual attention? Revisited! In: Attention and performance 15: Conscious and
849    nonconscious information processing. Attention and performance series. The MIT
850    Press, Cambridge, MA, US, pp 333-350

851 Kowler E, Anderson E, Dosher B, Blaser E (1995) The role of attention in the programming
852    of saccades Vision Res 35:1897-1916 doi:10.1016/0042-6989(94)00279-u

853 Laubrock J, Engbert R, Kliegl R (2005) Microsaccade dynamics during covert attention
854    Vision Res 45:721-730 doi:10.1016/j.visres.2004.09.029

855 Lee DK, Itti L, Koch C, Braun J (1999) Attention activates winner-take-all competition
856    among visual filters Nat Neurosci 2:375-381 doi:10.1038/7286

857 Li H-H, Barbot A, Carrasco M (2016) Saccade Preparation Reshapes Sensory Tuning
858    Current Biology 26:1564-1570 doi:10.1016/j.cub.2016.04.028

859 Li H-H, Pan J, Carrasco M (2019) Presaccadic attention improves or impairs performance
860    by enhancing sensitivity to higher spatial frequencies Scientific Reports 9:2659
861    doi:10.1038/s41598-018-38262-3

862 Lisberger SG, Fuchs AF, King WM, Evinger LC (1975) Effect of mean reaction time on
863    saccadic responses to two-step stimuli with horizontal and vertical components
864    Vision Research 15:1021-1025 doi:10.1016/0042-6989(75)90245-X

865 Luschei ES, Fuchs AF (1972) Activity of brain stem neurons during eye movements of
866    alert monkeys J Neurophysiol 35:445-461 doi:10.1152/jn.1972.35.4.445

867 Malevich T, Buonocore A, Hafed ZM (2021) Dependence of the stimulus-driven
868    microsaccade rate signature in rhesus macaque monkeys on visual stimulus size
869    and polarity Journal of Neurophysiology 125:282-295 doi:10.1152/jn.00304.2020

870 Martinez-Conde S, Macknik SL, Hubel DH (2004) The role of fixational eye movements in
871    visual perception Nature Reviews Neuroscience 5:229-240 doi:10.1038/nrn1348

872 Montagnini A, Castet E (2007) Spatiotemporal dynamics of visual attention during saccade
873    preparation: Independence and coupling between attention and movement planning
874    J Vis 7:8.1-16 doi:10.1167/7.14.8

875 Munoz DP, Guitton D (1991) Control of orienting gaze shifts by the tectoreticulospinal
876    system in the head-free cat. II. Sustained discharges during motor preparation and
877    fixation J Neurophysiol 66:1624-1641 doi:10.1152/jn.1991.66.5.1624

878 Munoz DP, Istvan PJ (1998) Lateral inhibitory interactions in the intermediate layers of the
879    monkey superior colliculus. Journal of Neurophysiology 79:1193-1209

880 Munoz DP, Wurtz RH (1993) Fixation cells in monkey superior colliculus. I. Characteristics
881    of cell discharge J Neurophysiol 70:559-575 doi:10.1152/jn.1993.70.2.559

882 Munoz DP, Wurtz RH (1995) Saccade-related activity in monkey superior colliculus. I.
883    Characteristics of burst and buildup cells. Journal of Neurophysiology 73:2313-2333

Ohl S, Kuper C, Rolfs M (2017) Selective enhancement of orientation tuning before
saccades Journal of Vision 17:2-11 doi:10.1167/17.13.2
Pastukhov A, Braun J (2010) Rare but precious: microsaccades are highly informative
about attentional allocation Vision Res 50:1173-1184
doi:10.1016/j.visres.2010.04.007
Pestilli F, Carrasco M (2005) Attention enhances contrast sensitivity at cued and impairs it
at uncued locations Vision Res 45:1867-1875 doi:10.1016/j.visres.2005.01.019
Posner MI (1980) Orienting of Attention Quarterly Journal of Experimental Psychology
32:3-25 doi:10.1080/00335558008248231
Posner MI (2015) Orienting of attention: Then and now Quarterly journal of experimental
psychology (2006):1-12 doi:10.1080/17470218.2014.937446
Posner MI, Cohen Y (1984) Components of visual orienting Attention and Performance X
32:531-556
Pratt J, Abrams RA (1999) Inhibition of return in discrimination tasks J Exp Psychol Hum
Percept Perform 25:229-242 doi:10.1037//0096-1523.25.1.229
R Core Team (2020) R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria, http://www.R-project.org/
Reingold EM, Stampe DM (1999) Saccadic Inhibition in Complex Visual Tasks. In., vol
Chapter 35. Springer US, Boston, MA, pp 249-255. doi:10.1007/978-1-4757-3054-
8_35
Reingold EM, Stampe DM (2002) Saccadic Inhibition in Voluntary and Reflexive Saccades
Journal of cognitive neuroscience 14:371-388 doi:10.1162/089892902317361903
Remington RW (1980) Attention and saccadic eye movements J Exp Psychol Hum
Percept Perform 6:726-744 doi:10.1037//0096-1523.6.4.726
Rizzolatti G, Riggio L, Dascola I, Umiltà C (1987) Reorienting attention across the
horizontal and vertical meridians: Evidence in favor of a premotor theory of attention
Neuropsychologia 25:31-40 doi:10.1016/0028-3932(87)90041-8
Rolfs M, Carrasco M (2012) Rapid simultaneous enhancement of visual sensitivity and
perceived contrast during saccade preparation J Neurosci 32:13744-13752a
doi:10.1523/jneurosci.2676-12.2012
Rolfs M, Kliegl R, Engbert R (2008) Toward a model of microsaccade generation: The
case of microsaccadic inhibition Journal of Vision 8:5-5 doi:10.1167/8.11.5
Salinas E, Stanford TR (2018) Saccadic inhibition interrupts ongoing oculomotor activity to
enable the rapid deployment of alternate movement plans Scientific Reports
8:14163 doi:10.1038/s41598-018-32224-5
Sheliga BM, Riggio L, Rizzolatti G (1994) Orienting of attention and eye movements
Experimental brain research Experimentelle Hirnforschung Expérimentation
cérébrale 98 doi:10.1007/BF00233988
Sheliga BM, Riggio L, Rizzolatti G (1995) Spatial attention and eye movements
Experimental brain research Experimentelle Hirnforschung Expérimentation
cérébrale 105 doi:10.1007/BF00240962
Shepherd M, Findlay JM, Hockey RJ (1986) The relationship between eye movements and
spatial attention The Quarterly Journal of Experimental Psychology Section A
38:475-491 doi:10.1080/14640748608401609
Smith DT, Schenk T (2012) The Premotor theory of attention: time to move on?
Neuropsychologia 50:1104-1114 doi:10.1016/j.neuropsychologia.2012.01.025
Solomon JA (2004) The effect of spatial cues on visual sensitivity Vision Res 44:1209-
1216 doi:10.1016/j.visres.2003.12.003
Tian X, Yoshida M, Hafed ZM (2016) A Microsaccadic Account of Attentional Capture and
Inhibition of Return in Posner Cueing Frontiers in Systems Neuroscience 10
doi:10.3389/fnsys.2016.00023

Tian X, Yoshida M, Hafed ZM (2018) Dynamics of fixational eye position and
microsaccades during spatial cueing: the case of express microsaccades Journal of
Neurophysiology 119:1962-1980 doi:10.1152/jn.00752.2017
Watson AB, Pelli DG (1983) Quest: A Bayesian adaptive psychometric method Perception
& Psychophysics 33:113-120 doi:10.3758/BF03202828
Wilkinson GN, Rogers CE (1973) Symbolic Description of Factorial Models for Analysis of
Variance Journal of the Royal Statistical Society Series C (Applied Statistics)
22:392-399 doi:10.2307/2346786
Zhao M, Gersch TM, Schnitzer BS, Dosher BA, Kowler E (2012) Eye movements and
attention: The role of pre-saccadic shifts of attention in perception, memory and the
control of saccades Vision Research 74:40-60
doi:https://doi.org/10.1016/j.visres.2012.06.017
947