Large-scale study reveals regional fungicide applications as a major determinant of

resistance evolution in the wheat pathogen *Zymoseptoria tritici* in France.

Running head:

Resistance selection driven by regional fungicide use in French *Zymoseptoria tritici*

populations.

Article type:

Original research article – Full paper

Authors:

Maxime GARNAULT[1,2], Clémentine DUPLAIX[1], Pierre LEROUX[1], Gilles COULEAUD[3], Olivier

DAVID[2]*, Anne-Sophie WALKER[1]* and Florence CARPENTIER[1,2]*

[1] Université Paris-Saclay, INRAE, AgroParisTech, UMR BIOGER, 78850, Thiverval-Grignon,

France

[2] Université Paris-Saclay, INRAE, MaIAGE, 78350, Jouy-en-Josas, France

[3] Arvalis-Institut du Végétal, 91720 Boigneville, France

* These authors contributed equally to this study

Author for correspondence:

Florence Carpentier

Tel: +331 44 08 86 35

E-mail: florence.carpentier@inrae.fr

Summary:

• Research rationale: In modern cropping systems, the quasi-systematic use of plant

protection products selects for resistance in pest populations. The emergence and

evolution of this adaptive trait threaten treatment efficacy. We identified determinants

of fungicide resistance evolution and quantified their effects at a large spatiotemporal

scale.

• Methods: We focused on *Zymoseptoria tritici*, which causes leaf blotch in wheat.

Phenotypes of qualitative or quantitative resistance to various fungicides were

monitored annually, from 2004 to 2017, at about 70 sites throughout France. We

modelled changes in resistance frequency with regional anti-*Septoria* fungicide use,

yield losses due to the disease and the regional area under organic wheat.

• Key results: We found that the major driver of resistance dynamics was fungicide use

at the regional scale. We estimated its effect on the increase in resistance and

apparent relative fitness of each resistance phenotype. The predictions of the model

replicated the spatiotemporal patterns of resistance observed in field populations

fairly accurately.

• Main conclusion: The evolution of fungicide resistance is determined at very large

scales. There is therefore a need for the collective management of resistance, which

could be guided by the results of studies like this.

Keywords:

Mycosphaerella graminicola, *Zymoseptoria tritici*, Septoria leaf blotch,

Resistance determinants, Large-scale evolution, Regional fungicide use, Resistance

dynamics, Mathematical modelling

1 INTRODUCTION

The efficacy of pesticides and drugs has been compromised by the rapid and widespread evolution of resistance, increasing the use of pesticides and drugs to maintain control levels (Georghiou & Mellon, 1983; Russell, 2005; Gould *et al.*, 2018). The management of resistance evolution is essential for human health, biodiversity and food security, given the rapid emergence and spread of resistance and the lack of new modes of action (MoA) (Palumbi, 2001; Grimmer *et al.*, 2014). Many studies have investigated the effects of various factors on the evolution of resistance: fitness cost (Andersson, 2003), mutation rate (Martinez & Baquero, 2000; Gressel, 2011), population size (Sisterson *et al.*, 2004), strength of selection pressure and its mitigation in anti-resistance strategies (Oz *et al.*, 2014; van den Bosch *et al.*, 2014). A number of studies have advocated further studies on the relative impact of these factors on a given pest and of the interactions between these factors (Berendonk *et al.*, 2015; Hughes & Andersson, 2015), with a view to promoting large-scale strategies (Okeke *et al.*, 2005; Menalled *et al.*, 2016).

Several studies have shown how agricultural selection pressures affect the large-scale structure of pest populations at national scale. For instance, the national distribution of resistance varieties shapes the adaptation of pathogen populations to cultivars (Tyutyunov *et al.*, 2008; Papaïx *et al.*, 2011). Historical herbicide applications have been shown to drive the evolution of herbicide resistance at a national scale (Hicks *et al.*, 2018). For fungicide resistance, theoretical studies have revealed that combining effective MoAs over time and space can delay resistance evolution (REX Consortium, 2013; van den Bosch *et al.*, 2014) and that large-scale management strategies may differ and interact with in-field strategies (Parnell *et al.*, 2006). However, so far, in the absence of large-scale studies, recommendations about fungicide use mostly stem from empirical studies conducted in local field trials assessing the impact of different spraying strategies (Rosenzweig *et al.*, 2008; Dooley *et al.*, 2016a,b; Heick *et al.*, 2017).

The aim of this study was to highlight the determinants of fungicide resistance evolution at

the national scale in France. We investigated the main potential drivers of evolution: (*i*) the

selection pressure effect, as assessed by regional fungicide use, (*ii*) the genetic drift effect,

which is modulated by population size (Maxwell *et al.*, 1990; Sisterson *et al.*, 2004), using

yield losses as a proxy and (*iii*) the refuge effect, including the fraction of wheat fields

unsprayed with fungicides over the territory (Parnell *et al.*, 2006; Tabashnik *et al.*, 2008),

assessed by determining the area under organic wheat.

We focused our analysis on *Zymoseptoria tritici* (formerly *Septoria tritici* and *Mycosphaerella*

graminicola as teleomorph), an ascomycete responsible for septoria leaf blotch (STB) on

winter wheat. *Z. tritici* has many features facilitating the emergence of resistance: high

genome plasticity, a large population size, high genetic diversity, asexual and sexual

reproduction, an ability to disperse over large distances (Zhan & McDonald, 2004; Croll &

McDonald, 2012). STB is one a major wheat disease that can cause yield losses of up to

50% worldwide (Ponomarenko *et al.*, 2011; Torriani *et al.*, 2015). In western Europe, up to

70% of all fungicide use is linked to STB control (Fones & Gurr, 2015). As a result, various

degrees of resistance to all authorised unisite inhibitors (*i.e.* targeting a single cellular site)

have been observed in France.

We previously proposed an initial analysis of the Performance trial network dataset, in which

phenotypes of resistance to four fungicide MoAs were monitored annually, from 2004 to

2017, at about 70 sites throughout France (Garnault *et al.* 2019). We found significant

differences between resistance phenotypes in terms of changes in spatial distribution and/or

growth rates. Major differences in population structure and dynamics were highlighted

between the north and south of France.

We develop here an explanatory model for identifying the determinants of these regional

spatiotemporal heterogeneities in resistance evolution according to resistance phenotype.

We investigated the effect of annual fungicide use, pathogen population size and the fraction

of refuges, all at the regional scale. Our analysis shows that the evolution of resistance can

be assessed at regional scale, and that the major determinant of resistance is the selection

pressure exerted by fungicide applications in the preceding year. This study provides

empirical results for regional resistance management, at a level intermediate between field

and national recommendations. A sound understanding of resistance evolution is required for

smart resistance management and should ultimately help to reduce pesticide use in

agrosystems.

2 MATERIALS AND METHODS

2.1 Data description

2.1.1 Sampling of *Z. tritici* populations and estimation of resistance frequency

The "Performance network" is supervised by ARVALIS-Institut du Végétal and the INRAE

research institute at Thiverval-Grignon. It carried out field trials on wheat throughout France

between 2004 and 2017. The frequency of resistant phenotypes in *Z. tritici* populations

sampled annually in these trials is recorded in the associated dataset (see Garnault *et al*.,

2019 for further information).

The frequency of resistant phenotypes was estimated by collecting bulk pycnidiospores from

30-40 infected leaves to ensure that the sample was representative of local populations.

Phenotypes were distinguished on the basis of their germination or growth on Petri dishes

containing discriminatory doses of fungicides, optimised on individual genotyped isolates

(see Leroux & Walker, 2011 and Garnault *et al*., 2019 for more details). We then considered:

(*i*) the phenotype displaying specific qualitative resistance to strobilurins (or QoIs; inhibitors

of respiration complex III), hereafter referred to as the StrR phenotype, (*ii*) the group of

phenotypes with moderate quantitative resistance to DMIs (sterol 14α-demethylation

inhibitors), hereafter referred to as TriMR phenotypes, (*iii*) the group of phenotypes with a

high quantitative resistance to DMIs, hereafter referred to as TriHR phenotypes. The TriMR

group encompasses the TriR6 and TriR7-TriR8 phenotypes, which were also included in the

analysis (TriR6 strains were recognised on the basis of their growth on low doses of prochloraz, contrasting with the lack of growth of TriR7–TriR8 strains in these conditions; Leroux & Walker, 2011).

Region, year, sampling date and cultivar grown were recorded for each sample. We considered only populations from unsprayed plots for this study. The plots were sampled at two time points: at "S1" in April-May, at about the Z32 wheat stage (n=1320, from 2006 to 2011), and at "S2" in May-June, at about the Z39-Z55 wheat stage (n=2407, from 2004 to 2017).

In this study, we focused on the phase of resistance selection. We therefore extracted from the Performance dataset the time periods during which resistance frequencies were increasing. These periods were 2004 to 2012 (n=852, 16 regions) for the StrR phenotype, 2005 to 2011 (n=754, 16 regions) for the TriMR phenotype group, and 2010 to 2017 (n=360, 14 regions) for the TriHR phenotype group. We also included data from 2006 to 2017 for the TriR6 and TriR7-TriR8 phenotypes (n=910 and n=851, respectively), for analysis of the spatial heterogeneity of their frequencies. The data are summarised in Table 1.

2.1.2 Regional fungicide use

Every year, Bayer Crop Science uses field surveys to estimate the area of wheat sprayed with fungicides containing anti-STB active ingredients (AIs) in each region of France. These data do not include information about the dose used in the application. They provide information only about the areas sprayed with the AIs concerned, and the number of sprayings. These areas are expressed in deployed hectares.

We retained the most widely used AIs for each MoA (AIs accounting cumulatively for more than 95% of the use of the MoA), to prevent background noise from AIs with a limited impact on STB control. The model therefore included pyraclostrobin (26%), azoxystrobin (19%),

trifloxystrobin (15%), kresoxim-methyl (15%), fluoxastrobin (12%) and picoxystrobin (12%)

for QoIs; and epoxiconazole (30%), prochloraz (17%), tebuconazole (13%), cyproconazole

(11%), prothioconazole (10%), propiconazole (7%), metconazole (7%), fluquinconazole (2%)

and hexaconazole (1%) for DMIs.

We took the regional heterogeneity in wheat production between regions (and, hence, in the

area sprayed with fungicides) into account, by dividing the number of deployed hectares by

the regional area under conventionally farmed wheat. The latter was calculated by

subtracting the area under organic wheat (see section 2.1.4) from the total area under wheat

(from the AGRESTE online data: agreste.agriculture.gouv.fr) for each year and region. This

new variable unit was named $ha_{\frac{D}{C}}$ (D for deployed and C for cropped hectares), and was

proportional to the mean number of times each AI was used over a cropping season in a

given region. The national trend and the regional heterogeneity of fungicide use expressed in

$ha_{\frac{D}{C}}$ are shown for DMIs and QoIs in Fig. 1. Henceforth, this variable is denoted F_{itf}, with f

corresponding to the AI, t to the year and i to the region.

2.1.3 Yield losses induced by STB

ARVALIS-Institut du Végétal assessed annual yield loss by conducting paired plot

experiments throughout France (a mean 80 trials annually, from 2004 to 2017, in 20 French

regions) (Arvalis, 2019). In each trial, we considered modalities cropped with STB-

susceptible wheat cultivars, in both unsprayed plots and sprayed plots (providing maximum

protection against diseases). Yield losses due to STB were calculated by subtracting the

yield in the unsprayed plot from that in the sprayed plot. We predicted regional yield losses

for each year with a linear model (fixed effects: year, region; random effects: wheat cultivar,

trial). The national trend and the regional heterogeneity of yield losses, expressed in decitons

per hectare, are shown in Fig. 1. This variable is denoted P_{it} hereafter, with t corresponding

to the year and i to the region.

2.1.4 Proportion of the total area under wheat farmed organically

The area under organically farmed wheat crops was recorded by AgenceBIO (the French national platform for the promotion and development of organic farming) and ARVALIS-Institut du Végétal. We collected regional data from 2007 onwards, and national data from 2004 onwards. The regional areas under organic wheat between 2004 and 2006 were assessed from the observed mean proportions of the regional area under organic wheat in subsequent years and from national data for 2004 to 2006. We used the regional proportion of wheat under organic farming in our models. This proportion was calculated by dividing the regional area under organic wheat by the total area under wheat in the same region, based on AGRESTE online data. The national trend and the regional heterogeneity of the area under organic wheat, expressed in hectares, are shown in Fig. 1. This variable is denoted R_{it} hereafter, with t corresponding to the year and i to the region.

2.2 Statistical modelling

We modelled the change in frequency for each resistance phenotype in French populations. The model took into account (*i*) the different phases of resistance dynamics (see below), (*ii*) the effects of previously described potential regional determinants and finally (*iii*) variability due to the sampling design (sampling date and wheat cultivar).

Phases in resistance dynamics. We distinguished three phases in resistance dynamics: "no resistance" (frequency equal to 0), "resistance selection" and "generalized resistance" (frequency equal to 100). During the "resistance selection" phase, observations were modelled with binomial random variables with a sample size of 100 (mean number of observed spores used to determine frequencies). The probabilities that a population was in the "no resistance", "generalized resistance" or "resistance selection" phase depended on the year t. These probabilities were referred as π_{0t}, π_{100t} and $(1 - \pi_{0t} - \pi_{100t})$, respectively. Thus, Y_{itjkn}, the n^{th} frequency observed in region i, in year t, on cultivar j and at sampling

217   date *k* followed a zero-and-one inflated binomial distribution (Eqn 1).

219   **Eqn 1**

$$Y_{itjkn} \begin{cases} = 0 & with\ probability\ \pi_{0t} \\ \sim \mathcal{B}\big(100, p_{itjkn}\big) & with\ probability\ 1 - \pi_{0t} - \pi_{100t} \\ = 100 & with\ probability\ \pi_{100t} \end{cases}$$

221   ***Resistance evolution.*** Using a logit transformation (Eqn 2), the proportion p_{itjkn} of resistant

222   phenotypes during resistance evolution was modelled by the regional dynamics D_{it} and the

223   variability due to sampling design ζ_{itjkn}.

225   **Eqn 2**

$$logit\big(p_{itjkn}\big) = ln\left(\frac{p_{itjkn}}{1 - p_{itjkn}}\right) = D_{it} + \zeta_{itjkn}$$

Regional dynamics. The change in resistance frequencies depended on regional-scale

variables: fungicide use, yield losses and areas under organic farming. The regional

dynamics $D_{i(t+1)}$ in region *i* at year *t+1* was obtained by adding the regional dynamics of the

previous year D_{it} to the additive effects of fungicide use ϕ_{it}, yield loss ρP_{it} and wheat area

under organic farming κR_{it} in year *t* (Eqn 3). D_{i1} is related to the logit of the initial resistance

frequency in region *i*. The parameter β is a constant corresponding to a continuous shift in

resistance frequency.

Eqn 3

$D_{i(t+1)} = D_{it} + \beta + \phi_{it} + \rho P_{it} + \kappa R_{it} \quad with\ t \geq 1$

We derived two models from Eqn 3: one in which the use of fungicides is specified for each

AI within MoAs, and another in which only the global use of each MoA (*i.e.* sum of AIs uses)

is considered.

In the first model, **M$_{AI}$,** the term ϕ_{it} from Eqn 3 was defined as in Eqn 4:

Eqn 4

$$\phi_{it} = \sum_{f \in \mathcal{F}} v_f F_{itf}$$

F_{itf} corresponds to the fungicide use for a specific AI f, in region i and year t (see section

2.1.2). The set $\mathcal{F}$ included all AIs associated with considered resistance: QoIs for the StrR

phenotype, DMIs for the TriR phenotypes. We assumed that fungicide use positively selected

resistant phenotypes over susceptible or less sensitive phenotypes. Thus, the parameters

associated with fungicide use were, by definition, positive (*i.e.* $v_f \geq 0$), except for the TriR6

and TriR7-TriR8 phenotypes, which were not the most DMI-resistant phenotypes over their

study period.

In the second model, **M$_{MoA}$,** we considered: $F_{it.} = \sum_{f \in \mathcal{F}} F_{itf}$, the regional use of a given MoA.

The term ϕ_{it} from Eqn 3 was simplified and written: $\phi_{it} = v * F_{it}$. This model was run only for

the StrR, TriMR and TriHR phenotypes, with, as above, the constraint $v \geq 0$.

Observation variability. The variability ζ_{itkln} of observations in Eqn 2 was modelled with a

mixed model (Equation 5).

Equation 5

$\zeta_{itjkn} = \gamma_j + \delta_k + \varepsilon_{itjkn}$

The parameter γ_j corresponds to the random effect of wheat cultivar (γ_j drawn from a

centered Gaussian distribution with standard deviation σ_γ). The parameter δ_k is the fixed

effect of sampling date k (*i.e.* "S1" or "S2", see section 2.1.1, with contrast $\delta_{S2} = 0$). Finally,

ε_{itjkn} is the overdispersion, modelled as a random individual effect with a mean of 0 and a

standard deviation of σ.

2.3 Parameter expression

The explanatory variables had different units (*e.g.* proportion of wheat under organic farming

vs. fungicide use). Moreover, the interpretation of the parameters of the zero-one-inflated

logistic regression was not straightforward. We simplified the interpretation, by defining the

expected frequency change (EFC) for each variable and phenotype. The EFC described the

frequency shift due to the mean value of this variable over a population with the mean

resistance frequency (*i.e.* the difference between p^e, the *expected frequency*, and $\bar{p}$, the

mean frequency of the resistance phenotype in the data). The expected frequency was

computed with:

• for v_f, the effect of fungicide use: $logit\left(p_{v_f}^e\right) = logit(\bar{p}) + \hat{v}_f \bar{F}_f$, where $\bar{F}_f$ is the mean

annual use of fungicide f over all regions, and $\hat{v}_f$ is the estimate of v_f;

• for β, the effect of constant growth: $logit\left(p_\beta^e\right) = logit(\bar{p}) + \hat{\beta}$;

• for ρ, the effect of yield losses due to STB: $logit\left(p_\rho^e\right) = logit(\bar{p}) + \hat{\rho}\bar{P}$;

• for κ, the effect of the area of wheat under organic farming: $logit(p_\kappa^e) = logit(\bar{p}) +$

$\hat{\kappa}\bar{R}$;

• for δ, the effect of sampling date: $logit(p_\delta^e) = logit(\bar{p}) + \hat{\delta}_{T0}$;

• for σ_γ, the standard deviation from the cultivar effect : $logit\left(p_{\sigma_\gamma}^e\right) = logit(\bar{p}) \pm \hat{\sigma}_\gamma$.

The *expected frequency change* was then calculated as $EFC = (p^e - \bar{p}) * 100$.

2.4 Bayesian analysis

Statistical analyses were performed with *R* software (R Development Core Team, 2008), in a

Bayesian framework, with the *rjags* package (Plummer, 2013).

Prior and posterior densities. Non-informative prior distributions were used (Supporting

Information, Eqn S1). Posterior distributions were estimated by Monte Carlo-Markov chain

(MCMC) methods. Five MCMC chains were run, over 1 000 000 iterations, with a burn-in of

100 000 and a thinning every 1 000 for the variable selection phase (see the following

section), followed by 500 000 iterations with a burn-in of 50 000 and a thinning every 500 for

the final parameter estimation. Convergence was assessed with the Gelman and Rubin $\hat{R}$

statistic (Gelman *et al.*, 2004). Credible intervals of the highest posterior density were

calculated from posterior densities with the *HDI* package (Dezeure *et al.*, 2015). Parameter

estimates were considered significant at the 5% level (or the 2.5% or 0.1% level), if their 95%

credible interval (97.5% and 99.9%, respectively) did not contain the 0 value.

Variable selection. We used a selection procedure to identify the relevant variables in each

model for each resistance phenotype. We used a method based on indicator variables (Kuo

& Mallick, 1998), in which each predictor was multiplied by a dummy variable with a prior

distribution corresponding to a Bernoulli distribution with parameter p = 0.5. A predictor was

retained in the model if the posterior expectation of its indicator variable was greater than

0.75 (thus, greater than its prior expectation of 0.5).

Predictive check. We assessed the fit of the model to the data by posterior predictive

checks (Gelman *et al.*, 2004). Replicated data (y_{itjkn}^{rep}) generated during the MCMC algorithm

from the model posterior densities were compared to observed data (y_{itjkn}). The mean

posterior value of $PP_{itjkn}^{check} = P(y_{itjkn} - y_{itjkn}^{rep} < 0 \mid Y)$, where Y is the vector of observations,

was calculated and denoted PP_{check}. This value indicated the goodness of fit of the model,

with a good fit corresponding at $PP_{check} = 0.5$.

Variable weight. We assessed the influence of each explanatory variable θ by calculating its

weight (W_θ). The weight W_θ was defined as the ratio of $RSS_{full-\theta}$ to RSS_{full}, where RSS_{full}

and $RSS_{full-\theta}$ are the residual sums of squares of the full model (*i.e.* including all the

explanatory variables selected by the variable selection procedure) and of this same model

but without the explanatory variable θ, respectively. RSS should be minimal for the full model,

so removing an explanatory variable should increase RSS: the more information θ

contributes, the greater the increase in RSS and the higher the value of W_θ. Conversely, if the

information provided by θ is negligible, RSS is unaffected and W_θ is minimal (*i.e.* close to 1).

In the result tables, we have calculated the relative weights by dividing individual variable

weights by the sum of the weights of all variables.

Model comparison. For comparison of the M*AI* and *M*MoA models, we calculated the deviance

information criterion, DIC (Plummer, 2013), and the coefficient of determination, R^2.

Predicted data. We computed predictions of the resistance frequencies for each phenotype

for a given region *l*, and a given year *T* (Equation 6).

Equation 6

$\hat{Y} = [1 - (\hat{\pi}_{0T} + \hat{\pi}_{100T})] * logit^{-1}[\hat{D}_{l1} + \hat{\beta}(T-1) + \sum_{t=1}^{T-1}(\hat{\rho}P_{lt} + \hat{\kappa}R_{lt} + \sum_{f \in \mathcal{F}} \hat{v}_f F_{ltf})] + \hat{\pi}_{100T}$

where parameter-hat are parameter estimates (*i.e.* their posterior mean). We therefore built

maps of resistance status, for known initial frequencies, use of fungicides, yield losses and

areas under organic wheat, from year 1 to year *T-1* in region *l*.

3 RESULTS

3.1 Overview of model fits

The convergence of the MCMC chain was satisfactory for all models (*i.e.* the Gelman and

Rubin indicator $\hat{R}$ was below 1.1 for all parameters, in all models) and model fit was good

(PP_{check} always between 0.498 and 0.511).

After the selection procedure, no effect was retained for the following models: M_{AI} for the

TriHR resistance phenotype, and M_{MoA} for the TriMR and the TriHR groups of resistance

phenotypes. Thus, the effects of fungicide use, yield losses, and areas of wheat under

organic farming were not significant in these models. The only remaining parameter was the

growth constant (already studied in Garnault *et al.*, 2019). As all estimates of the parameters

of interest were equal to 0, we do not discuss the results of these models, and they do not

appear in the result tables.

Finally, with Spearman's method, a few significant correlations were found between some AI

uses (F_{itf}) in model inputs, but no significant correlation between estimates ($\hat{v}_f$) was found in

model outputs (Supporting Information, Fig. S1).

3.2 Ranking of variable weight

Regional fungicide use appeared to be the major factor driving resistance evolution. In the

M_{AI} models, which explicitly considered each AI, regional fungicide uses systematically had

the highest relative weight. For the StrR and TriMR phenotypes, it accounted for 87.4% and

72.6%, respectively. It accounted for 53.1% and 43.3% for the TriR6 and TriR7-TriR8

phenotypes, respectively (Table 2). For the M_{MoA} models, fungicide use, considered as the

sum of AI uses within the same MoA, was also the major determinant of the StrR phenotype

(associated with qualitative resistance to QoIs), accounting for 79.4% of the sum of variable

weights (Table 3). For the TriMR and TriHR groups of phenotypes, no explanatory variables

were selected for the M_{MoA} models.

The growth constant was also a major parameter, albeit to a lesser extent. In M_{AI} models, the

weight of the growth constant was lower than that of regional fungicide use by a factor of 5.7

times for TriMR phenotypes, 1.32 for the TriR6 phenotype, and 1.08 for the TriR7-TriR8

phenotypes (Table 2). For the StrR phenotype, the growth constant ranked third, with a

weight lower than that of fungicide use by a factor of almost 20, for both models (Tables 2

and 3).

The proportion of the area under wheat farmed organically had a high relative weight for the

StrR and TriMR phenotypes, but was not selected for the TriR6 and TriR7-TriR8 phenotypes.

For the StrR phenotype, its effect was ranked second on the basis of relative weight, at

14.6% and 6.8% in the M_{MoA} and the M_{AI} models, respectively (Tables 2 and 3). For TriMR

phenotypes, the relative weight of the wheat area under organic farming was about 13.3%, a

value very similar to that for the growth constant (Table 2). Yield loss was systematically

excluded during the selection procedure, for all models and all phenotypes.

Sampling data and wheat cultivar, variables reflecting local variability in trials, had only a low

relative weight in models, with values always below 5%, except for the wheat cultivar variable

for the TriR7-TriR8 phenotype, for which the value was 13.4% (Table 2).

3.3 Effect of variables at the regional scale

3.3.1 Regional fungicide use

For the StrR phenotype, in the M_{MoA} model, the effect of the overall use of QoI fungicides was

highly significant ($P < 0.001$) and the expected frequency change (EFC) was estimated at

6.64%. Thus, on average, the use of QoIs led to a 6.64% annual increase in the frequency of

StrR (Table 3).

In the M_{AI} model, two fungicides from the six QoI AIs were selected: kresoxim-methyl and

pyraclostrobin. Their EFCs were similar: 4.26% ($P < 0.001$) and 3.29% ($P < 0.025$),

respectively (Table 2). The M_{MoA} and M_{AI} models also had similar adequacies to data,

according to their DIC (3480.2 and 3468.8 respectively) and R^2 values (0.82 and 0.81,

respectively).

For TriMR phenotypes, the effect of DMI use was estimated only in the M_{AI} model, as no

explanatory variable was selected in the M_{MoA} model. One AI of the nine DMI fungicides was

selected: epoxiconazole ($P < 0.025$) with an estimated positive EFC of 4.82% (Table 2).

As the TriR6 and TriR7-TriR8 phenotypes were not the most resistant to DMI over the study

period, the effect of fungicide use was not constrained to be null or positive, and was

estimated only in the M_{AI} model. For the TriR6 phenotype, three AIs from the nine DMI

fungicides were selected: prochloraz, with a positive EFC of 4.73% ($P < 0.001$),

propiconazole with a negative EFC of -2.48% ($P < 0.025$) and tebuconazole with a negative

EFC of -8.38% ($P < 0.001$). Thus, mean annual prochloraz use increased the frequency of

TriR6 by 4.73%, whereas mean annual use of tebuconazole counterselected TriR6 strains,

leading to an 8.38% decrease in their frequency. For the TriR7-TriR8 phenotype, three AIs

from the nine DMI fungicides were selected: cyproconazole and tebuconazole, with positive

EFCs of 2.6% ($P < 0.025$) at 2.38% ($P < 0.05$), respectively, and prochloraz, with a negative

EFC of -3.99% ($P < 0.001$). Prochloraz and tebuconazole clearly had opposite selection

effects on TriR6 and TriR7-TriR8 phenotypes.

3.3.2 Growth constant

The growth constant EFC quantified the change in resistance in the absence of fungicide use

and unsprayed refuges, for a mean potential yield loss. It therefore represented the relative

apparent fitness (referred to hereafter simply as fitness) of the resistant phenotype

considered (*i.e.* how much faster the resistant phenotype would grew compared to the rest of

the population (Hartl & Clark, 1997) in a year without fungicide treatment). A negative growth

constant indicates a fitness cost, whereas a positive growth constant indicates a fitness gain.

For the StrR phenotype, the growth constant provided an indication of the fitness of the

resistant strains relative to the sensitive strains. The estimated fitness costs for this

phenotype were similar in the M_{AI} (-3.74%; $P < 0.05$; Table 2) and M_{MoA} (-3.23%; not

significant; Table 3) models. DMIs selected a large diversity of phenotypes (TriLR, TriMR and

TriHR groups), and no sensitive strains were detected during the study period. For TriMR

phenotypes, the model was estimated with data from 2007 to 2011, when the frequency of

the TriHR phenotype was still negligible (Fig. 1). Thus, the growth constant for TriMR

phenotypes mostly compared their fitness with that of TriLR phenotypes. It was estimated at

-3.94%, of borderline significance ($P < 0.1$), and was associated with a relative weight of

12.7% (Table 2). The TriMR group included the TriR6 and TriR7-TriR8 phenotypes. For the

TriR6 and TRiR7-TriR8 strains, the model was estimated with data from 2006 to 2017.

However, the TriHR phenotype has been non-negligible since 2014 (Figure 1). Thus, the

growth constant for TriR6 strains compared their fitness with that of all the other phenotypes

in the population: TriLR, TriR7-TriR8 and TriHR. The TriR6 growth constant was estimated at

3.58% ($P < 0.025$; Table 2). This result may reflect a balance between a fitness cost of TriR6

relative to TriLR and TriR7-TriR8 strains, and a fitness gain relative to the TriHR phenotype.

This rationale also applies to the apparent fitness gain of TriR7-TriR8, estimated at +1.63%

($P < 0.1$; Table 2).

3.3.3 Wheat area under organic farming

The proportion of the area under wheat management by organic farming methods increased

resistance frequency, with an EFC estimated at 3.95% and 3.26% in the M_{MoA} and M_{AI}

models, respectively, for the StrR phenotype ($P < 0.001$; Tables 2 and 3), and at 2.52% for

the TriMR phenotype in the M_{AI} model ($P < 0.001$; Table 2). This variable was not selected

for the TriR6 and TriR7-TriR8 phenotypes.

3.3.4 Yield losses

The selection procedure did not retain the yield loss variable in any of the models.

3.4 Prediction maps

We mapped the predicted resistance frequencies for each year and for all phenotypes. Our

maps predicting StrR phenotype dynamics were mostly based on regional QoI fungicide use

and, to a lesser extent, the area under organic wheat, as explanatory variables (Fig. 2).

Predictions mimicked the spatial propagation from the north to the south of France observed

between 2004 and 2011 (as described by Garnault *et al.*, 2019). Based on regional DMI use,

our model accurately predicted (Fig. 3) the observed spatial partitioning of TriR6 strains,

which were found mostly in the north east of France, and TriR7-TriR8 phenotypes, which

were mostly localised in the south west (as described by Garnault *et al.*, 2019).

4 DISCUSSION

In this study, we developed a model for identifying the determinants of fungicide resistance

evolution at the regional scale. The candidate explanatory variables were regional fungicide

use, pathogen population size (approximated by potential yield losses) and the fraction of

refuges (approximated by the fraction of fields under organic wheat in a region). We

analysed resistance frequencies in *Z. tritici* populations on winter wheat. Frequencies were

monitored by the Performance trial network over the national territory of French (2004–2017;

~70 locations each year). We studied resistance against fungicides with two modes of action:

QoIs, associated with qualitative resistance (StrR phenotype), and DMIs, associated with

quantitative resistance (a continuum of multiple resistance phenotypes forming three main

groups: TriLR, TriMR and TriHR, with low, medium and high levels of resistance to DMIs,

respectively). The TriMR phenotypes encompassed two phenotypes: TriR6 and TriR7-TriR8.

The regional use of fungicides is the main driver of resistance evolution at the plot

scale

We demonstrated that regional fungicide use was a major determinant of the evolution of

resistance. Fungicide use data at regional level, without taking into account the use of

particular fungicides in particular fields, was sufficiently informative to explain resistance

dynamics. Fungicide selection is, therefore, a large-scale process, and an understanding of

the evolution of fungicide resistance requires consideration of the regional use of these

compounds.

For qualitative resistance (e.g. StrR phenotype), the global use of a MoA (*i.e.* summed uses

of the AIs of this MoA), can be considered as a good predictor of resistance evolution and

distinguishing the effects of individual AIs within the MoA did not improve the fit of the model.

By contrast, for quantitative resistance, it appeared to be necessary to consider each AI

within the MoA separately, as they may select for different phenotypes (*e.g.* antagonist

effects of prochloraz, tebuconazole on TriR phenotypes).

Interestingly, the AIs selected by the model were consistent with the history of fungicide use

and/or with the patterns of cross-resistance described for some phenotypes. For instance,

kresoxim-methyl was one of the first QoIs authorised in France (commercialised in 1997,

www.ephy.anses.fr) and was selected to explain the evolution of StrR strains, whereas more

recent QoIs were not. The fungicides most used over the study period (*i.e.* epoxiconazole for

DMIs, pyraclostrobin and kresoxim-methyl for QoIs; Supporting Information, Fig. S2-S3),

were also selected in models. The large-scale effect of AIs was also linked to the cross-

resistance pattern of the phenotypes. The estimated selection effect of epoxiconazole was

consistent with resistance factors (RFs) of TriMR phenotypes to epoxiconazole, being higher

than those of TriLR phenotypes which were the second most frequent resistance phenotype

during this period (RFs described in Supporting Information, Table S1 from Leroux & Walker,

2011). The selection for TriR6 and counterselection for TriR7-TriR8 strains induced by the

prochloraz were consistent with RFs (6.7 and <1.5, respectively), as well as the effect of

counterselection for TriR6 and selection for TriR7-TriR8 strains by tebuconazole (RF=74 and

91, respectively for TriR6 and TriR8 strains, TriR7 strains being unfrequent (Huf *et al.*,

2018)). Propiconazole and cyproconazole were often used together as a mixture, so were

strongly correlated (Supporting Information, Fig. S1). The effects of counterselection for

TriR6 strains by propiconazole and of selection for TriR7-TriR8 by cyproconazole may

indicate that their mixture globally promoted the selection of TriR7-TriR8 strains over TriR6

strains. Again, this is consistent with RFs to propiconazole (35 and 54, respectively for TriR6

and TriR8 strains) and to cyproconazole (11 and 13, respectively). These findings also

confirm that the laboratory characterisation of strains can be good predictor of resistance

evolution in the field, if properly used, as reported by Blake *et al.* (2018).

Our findings highlight the importance of defining precise resistance profile phenotypes.

Indeed, the model fitted slightly better the frequencies on more homogeneous phenotypes

(R^2 = 0.48 for TriMR vs. 0.56 for TriR6 and TriR7-TriR8 phenotypes). In addition, no

significant effect of the different DMIs was found for the TriHR phenotype group. Indeed,

TriHR strains encompass multiple heterogeneous phenotypes, resulting from combinations

of target alteration, target overexpression and enhanced efflux, as resistance mechanisms

(Leroux & Walker, 2011; Huf *et al.*, 2018). The tremendous diversity and redundancy of

phenotypes observed in the field, especially for TriHR strains, making it possible to classify

strains only approximately. There is therefore a need to develop molecular tools for

quantifying genotypes rather than phenotypes. Multi-trait high-throughput genotyping

provides a more accurate resistance frequency, and should ultimately lead to improvements

in our ability to predict resistance evolution.

The proportion of the wheat area under organic farming may be still too limited to

mitigate the evolution of resistance via a refuge effect

Wheat areas managed under organic farming systems are not treated with synthetic

fungicides. They could, therefore, act as refuges, for "wild" susceptible or less resistant

individuals, which may reproduce, delaying the evolution of resistance by a dilution effect in

mobile species (Gould, 2000). But refuges may also provide a heterogeneous environment,

promoting sink-source dynamics: selection-free wheat areas acting as a source of

susceptible strains that migrate towards areas farmed conventionally. These opposite effects

have been described in studies of the resistance to transgenic crops expressing *Bacillus*

thuringiensis (*Bt*) toxins, where non-*Bt* crops may act as refuges diluting the selection

(Huang *et al.*, 2011), or promoting the evolution of resistance (Caprio, 2001).

The refuge effect for *Z. tritici* is, theoretically, weak due to the haploid nature of this organism

(Shaw, 2009), but this has never been studied experimentally. We did not validate the

beneficial effect of refuges, approximated by the area under wheat farmed organically. On

the contrary, we estimated that the selection of StrR and TriMR phenotypes would increase

with wheat areas under organic farming. However, the weight of this explanatory variable

remained much lower than that for regional fungicide use. This effect was not found

significant for the other phenotypes.

According to Huang *et al.* (2011), three conditions must be satisfied for refuge strategies to

be successful: selection at "high dose", a very low initial frequency of resistance, and

sufficient refuge areas located nearby. In our study, initial frequencies of resistance (*i.e.* at

the very beginning of our study) were already quite high (up to 85% in some regions, Fig. 1).

In addition, the area under organic farming may still be too small within the landscape

(generally less than 1% until 2010). Nevertheless, since the 2010s, the proportion of wheat

under organic farming has steadily increased (Fig. 1). The effect of the area under organic

wheat may become detectable in those areas, and should be investigated further, particularly

for emerging resistance phenotypes.

The growth constant reveals a fitness penalty of resistant phenotypes

The growth constant represents the evolution of resistant phenotypes in the absence of

fungicide treatment. It represents the apparent fitness of the phenotype relative to that of the

susceptible phenotype (or other resistant phenotypes, in the case of quantitative resistance).

The term "apparent" is used because this quantification takes place in current crop

conditions. The fitness cost of resistance to drugs is known to be a key parameter driving

effective anti-resistance strategies (Andersson & Hughes, 2010; Melnyk *et al.*, 2015;

Mikaberidze & McDonald, 2015). If there is no fitness cost, regardless of the strategy used, it

will always result in the irreversible selection of resistance. However, it remains difficult to

infer the global fitness cost of a mutation throughout the entire life cycle of a pathogen

(Hollomon, 2015).

We inferred an apparent relative fitness penalty for StrR phenotypes, resulting in an annual

decrease of 3.74%. This was consistent with the fitness cost described by Hagerty & Mundt

(2016) using virulence comparison tests. The impact of cyp51 alterations, leading to TriR

phenotypes, on fitness is often evoked to explain the evolution of azole resistance in *Z. tritici*

populations (Cools *et al.*, 2013; Blake *et al.*, 2018) but it has not been quantified as yet. Our

model is consistent with these assumptions as it also inferred an apparent fitness penalty for

the TriMR group (-3.94% per year). Decreases in the use of QoI and DMI fungicides, and/or

the implementation of strategies favouring the expression of a resistance cost, may help to

slow the evolution of resistance. From our estimates, we can extrapolate a theoretical

equilibrium between resistance cost and selection, which could have led to a null growth of

the StrR and TriMR phenotypes by reducing by 50% the use of QoIs and 18% for DMIs.

For the TriR6 and TriR7-TriR8 phenotypes, the growth constant is an integrative value, as

these phenotypes were studied alongside more susceptible phenotypes (TriLR before 2010)

and more resistant phenotypes (TriHR after 2010). The positive growth constants we

estimated for these two phenotypes may indicate a fitness benefit of TriMR relative to TriLR

phenotypes and/or fitness cost of TriHR (*i.e.* also fitness benefit of TriMR relative to TriHR

phenotypes). Our model could be extended to determine the relative fitnesses of each

phenotype in cases of quantitative resistance. The global informative indicator provided by

our model could be used to guide the design of optimal large-scale fungicide deployment

strategies.

The yield losses caused by STB do not affect resistance evolution

Population size, a major parameter in population adaptation (Good *et al.*, 2012), is generally

positively correlated with resistance evolution (Weber & Diggins, 1990; Anderson, 2005; zur

Wiesch *et al.*, 2011). Indeed, a large population size increases the number of mutants

generated and decreases genetic drift (Linde *et al.*, 2002). We inferred this effect using

potential yield losses caused by STB as a proxy, but no significant effect was detected for

any resistance phenotype. Population size may not be limiting for resistance evolution in *Z.*

tritici (Zhan *et al.*, 2001; Mikaberidze *et al.*, 2017), particularly at larger scales. Population

size may not be described accurately enough as our proxy variable also depends on the

timing of infection (Shaw & Royle, 1993) and on stubble management (McDonald & Mundt,

2016).

Predicting resistance evolution over years

Prediction maps can be computed from our model, using only the initial regional frequencies

of the resistant phenotypes, the history of fungicide use (between the initial year and the year

to be predicted) and the history of area under organic farming. Predictions for the StrR

phenotype from 2004 to 2011 highlighted the same colonisation front structure from the

North to the South of France (Fig. 2) as reported by Garnault *et al.* (2019) albeit the regions

are assumed to be fully independent. Predictions for the TriR6 and TriR7-TriR8 phenotypes

also yielded stable spatial distributions between North-East and South-West France (Fig. 3),

as previously observed in Garnault *et al.* (2019).

Further analysis will be required to assess the prediction quality of the model. Nevertheless,

this finding supports the global validity of our model and paves the way for an original

approach to predicting resistance evolution in a heterogeneous landscape.

Conclusion

We developed a new model of resistance dynamics, which identified regional fungicide use

as the major determinant of fungicide resistance evolution, and the area under organic

farming as a much weaker explanatory variable. We estimated the apparent relative fitness

of resistant phenotypes, a key parameter for the development of sustainable resistance

management strategies. We also identified active ingredients as most tightly linked to

resistance evolution.

In conclusion, we showed here that the determination of resistance evolution occurs at a truly

large scale and demonstrating that concerted collective action is required, to reinforce

individual initiatives to tackle resistances effectively.

5 ACKNOWLEDGEMENTS

This work was funded by the INRAE SMaCH metaprogramme (FONDU project) and by

Bayer Crop Science (PhD scholarship of M. Garnault; CIFRE no. 2016-0695). We warmly

thank all partners of the Performance network for providing samples since 2004, and Arvalis-

Institut du Végétal for supervising the network. We are grateful to the INRAE MIGALE

bioinformatics facility (MIGALE, INRAE, 2020. Migale bioinformatics Facility, doi:

10.15454/1.5572390655343293E12) for providing computing resources. We thank Eva

Lacarce (AgenceBio) and Phillipe Viaux (Arvalis-Institut du Végétal) for providing information

about the historical changes in the area under organic wheat, and Clarisse Payet (Bayer

Crop Science) for providing data on fungicide use in France and for helpful discussions.

6 AUTHOR CONTRIBUTIONS

GC was responsible for Performance network administration and supervision and for

establishing the dataset. PL, ASW and CD carried out laboratory manipulations on the

network samples to determine resistance frequencies in *Z. tritici* populations. FC and OD

were involved in the development of the methodology and provided theoretical support for

statistical aspects for data analyses. MG handled data, coded the model, performed

statistical analyses and generated most of the results (figures, maps and tables) during his

PhD work. The article was written by MG, FC, ASW and OD.

7 REFERENCES

Anderson JB. **2005**. Evolution of antifungal-drug resistance: mechanisms and pathogen

fitness. *Nature Reviews Microbiology* **3**: 547–556.

Andersson DI. **2003**. Persistence of antibiotic resistant bacteria. *Current opinion in microbiology* **6**: 452–456.

Andersson DI, Hughes D. **2010**. Antibiotic resistance and its cost: is it possible to reverse resistance? *Nature Reviews Microbiology* **8**: 260–271.

Arvalis. **2019**. *Choisir & Décider, Synthèse Nationale⎕: Céréales à paille, Interventions de printemps.*

Berendonk TU, Manaia CM, Merlin C, Fatta-Kassinos D, Cytryn E, Walsh F, Bürgmann H, Sørum H, Norström M, Pons M-N, *et al.* 2015. Tackling antibiotic resistance: the environmental framework. *Nature Reviews Microbiology* **13**: 310.

Blake JJ, Gosling P, Fraaije BA, Burnett FJ, Knight SM, Kildea S, Paveley ND. **2018**. Changes in field dose--response curves for demethylation inhibitor (DMI) and quinone outside inhibitor (QoI) fungicides against *Zymoseptoria tritici*, related to laboratory sensitivity phenotyping and genotyping assays. *Pest management science* **74**: 302–313.

van den Bosch F, Oliver R, van den Berg F, Paveley N. **2014**. Governing principles can guide fungicide-resistance management tactics. *Annual Review of Phytopathology* **52**: 175–195.

Caprio MA. **2001**. Source-sink dynamics between transgenic and non-transgenic habitats and their role in the evolution of resistance. *Journal of economic entomology* **94**: 698–705.

Cools HJ, Hawkins NJ, Fraaije BA. **2013**. Constraints on the evolution of azole resistance in plant pathogenic fungi. *Plant Pathology* **62**: 36–42.

Croll D, McDonald BA. **2012**. The accessory genome as a cradle for adaptive evolution in pathogens. *PLoS pathogens* **8**.

Dezeure R, Bühlmann P, Meier L, Meinshausen N. **2015**. High-Dimensional Inference: Confidence Intervals, p-values and {R}-Software {hdi}. *Statistical Science* **30**: 533–558.

Dooley H, Shaw MW, Spink J, Kildea S. **2016a**. Effect of azole fungicide mixtures, alternations and dose on azole sensitivity in the wheat pathogen *Zymoseptoria tritici*. *Plant Pathology* **65**: 124–136.

Dooley H, Shaw MW, Spink J, Kildea S. **2016b**. The effect of succinate dehydrogenase

inhibitor/azole mixtures on selection of *Zymoseptoria tritici* isolates with reduced sensitivity.

Pest management science **72**: 1150–1159.

Fones H, Gurr S. **2015**. The impact of *Septoria tritici* Blotch disease on wheat: an EU

perspective. *Fungal Genetics and Biology* **79**: 3–7.

Garnault M, Duplaix C, Leroux P, Couleaud G, Carpentier F, David O, Walker A-S. **2019**.

Spatiotemporal dynamics of fungicide resistance in the wheat pathogen *Zymoseptoria tritici*

in France. *Pest management science* **75**: 1794–1807.

Gelman A, Carlin JB, Stern HS, Rubin DB. **2004**. Bayesian data analysis 2nd edn

Chapman & Hall. *CRC, Boca Raton FL*.

Georghiou GP, Mellon RB. **1983**. Pesticide resistance in time and space. In: Pest

resistance to pesticides. Springer, 1–46.

Good BH, Rouzine IM, Balick DJ, Hallatschek O, Desai MM. **2012**. Distribution of fixed

beneficial mutations and the rate of adaptation in asexual populations. *Proceedings of the*

National Academy of Sciences **109**: 4950–4955.

Gould F. **2000**. Testing *Bt* refuge strategies in the field. *Nature Biotechnology* **18**: 266.

Gould F, Brown ZS, Kuzma J. **2018**. Wicked evolution: Can we address the sociobiological

dilemma of pesticide resistance? *Science* **360**: 728–732.

Gressel J. **2011**. Low pesticide rates may hasten the evolution of resistance by increasing

mutation frequencies. *Pest management science* **67**: 253–257.

Grimmer MK, van den Bosch F, Powers SJ, Paveley ND. **2014**. Evaluation of a matrix to

calculate fungicide resistance risk. *Pest management science* **70**: 1008–1016.

Hagerty CH, Mundt CC. **2016**. Reduced virulence of azoxystrobin-resistant *Zymoseptoria*

tritici populations in greenhouse assays. *Phytopathology* **106**: 884–889.

Heick TM, Justesen AF, Jørgensen LN. **2017**. Anti-resistance strategies for fungicides

against wheat pathogen *Zymoseptoria tritici* with focus on DMI fungicides. *Crop protection*

99: 108–117.

Hicks HL, Comont D, Coutts SR, Crook L, Hull R, Norris K, Neve P, Childs DZ,

Freckleton RP. **2018**. The factors driving evolved herbicide resistance at a national scale.

Nature ecology & evolution **2**: 529.

Hollomon DW. **2015**. Fungicide resistance: facing the challenge-a review. *Plant Protection*

Science **51**: 170–176.

Huang F, Andow DA, Buschman LL. **2011**. Success of the high-dose/refuge resistance

management strategy after 15 years of *Bt* crop use in North America. *Entomologia*

Experimentalis et Applicata **140**: 1–16.

Huf A, Rehfus A, Lorenz KH, Bryson R, Voegele RT, Stammler G. **2018**. Proposal for a

new nomenclature for CYP 51 haplotypes in *Zymoseptoria tritici* and analysis of their

distribution in Europe. *Plant Pathology* **67**: 1706–1712.

Hughes D, Andersson DI. **2015**. Evolutionary consequences of drug resistance: shared

principles across diverse targets and organisms. *Nature Reviews Genetics* **16**: 459.

Kuo L, Mallick B. **1998**. Variable selection for regression models. *Sankhya: The Indian*

Journal of Statistics, Series B: 65–81.

Leroux P, Walker A-S. **2011**. Multiple mechanisms account for resistance to sterol 14α-

demethylation inhibitors in field isolates of *Mycosphaerella graminicola*. *Pest management*

science **67**: 44–59.

Linde CC, Zhan J, McDonald BA. **2002**. Population structure of *Mycosphaerella*

graminicola: from lesions to continents. *Phytopathology* **92**: 946–955.

Martinez JL, Baquero F. **2000**. Mutation frequencies and antibiotic resistance. *Antimicrobial*

agents and chemotherapy **44**: 1771–1777.

Maxwell BD, Roush ML, Radosevich SR. **1990**. Predicting the evolution and dynamics of

herbicide resistance in weed populations. *Weed technology* **4**: 2–13.

McDonald BA, Mundt CC. **2016**. How knowledge of pathogen population biology informs

management of *Septoria tritici* blotch. *Phytopathology* **106**: 948–955.

Melnyk AH, Wong A, Kassen R. **2015**. The fitness costs of antibiotic resistance mutations.

Evolutionary applications **8**: 273–283.

Menalled FD, Peterson RKD, Smith RG, Curran WS, Páez DJ, Maxwell BD. **2016**. The

eco-evolutionary imperative: revisiting weed management in the midst of an herbicide

resistance crisis. *Sustainability* **8**: 1297.

Mikaberidze A, McDonald BA. **2015**. Fitness cost of resistance: impact on management. In:

Fungicide Resistance in Plant Pathogens. Springer, 77–89.

Mikaberidze A, Paveley ND, Bonhoeffer S, van den Bosch F. **2017**. Emergence of

resistance to fungicides: the role of fungicide dose. *Phytopathology* **107**: 545–560.

Okeke IN, Klugman KP, Bhutta ZA, Duse AG, Jenkins P, O'Brien TF, Pablos-Mendez A,

Laxminarayan R. **2005**. Antimicrobial resistance in developing countries. Part II: strategies

for containment. *The Lancet infectious diseases* **5**: 568–580.

Oz T, Guvenek A, Yildiz S, Karaboga E, Tamer YT, Mumcuyan N, Ozan VB, Senturk GH,

Cokol M, Yeh P, *et al.* **2014**. Strength of selection pressure is an important parameter

contributing to the complexity of antibiotic resistance evolution. *Molecular biology and*

evolution **31**: 2387–2401.

Palumbi SR. **2001**. Humans as the world's greatest evolutionary force. *Science* **293**: 1786–

1790.

Papaïx J, Goyeau H, Du Cheyron P, Monod H, Lannou C. **2011**. Influence of cultivated

landscape composition on variety resistance: an assessment based on wheat leaf rust

epidemics. *New Phytologist* **191**: 1095–1107.

Parnell S, van den Bosch F, Gilligan CA. **2006**. Large-scale fungicide spray heterogeneity

and the regional spread of resistant pathogen strains. *Phytopathology* **96**: 549–555.

Plummer M. **2013**. Rjags: bayesian graphical models using MCMC. *R package version* **3**.

Ponomarenko A, Goodwin SB, Kema GHJ. **2011**. *Septoria tritici* blotch (STB) of wheat.

Plant Health Instructor **10**.

R Development Core Team. **2008**. R: A Language and Environment for Statistical

Computing.

REX Consortium. **2013**. Heterogeneity of selection and the evolution of resistance. *Trends*

in ecology & evolution **28**: 110–118.

Rosenzweig N, Atallah ZK, Olaya G, Stevenson WR. **2008**. Evaluation of QoI fungicide

application strategies for managing fungicide resistance and potato early blight epidemics in Wisconsin. *Plant Disease* **92**: 561–568.

Russell PE. **2005**. A century of fungicide evolution. *The Journal of Agricultural Science* **143**: 11–25.

Shaw MW. **2009**. Fungicide resistance: the dose rate debate. *Outlooks on Pest Management* **20**: 100–103.

Shaw MW, Royle DJ. **1993**. Factors determining the severity of epidemics of *Mycosphaerella graminicola* (*Septoria tritici*) on winter wheat in the UK. *Plant Pathology* **42**: 882–899.

Sisterson MS, Antilla L, Carrière Y, Ellers-Kirk C, Tabashnik BE. **2004**. Effects of insect population size on evolution of resistance to transgenic crops. *Journal of Economic Entomology* **97**: 1413–1424.

Tabashnik BE, Gassmann AJ, Crowder DW, Carrière Y. **2008**. Insect resistance to *Bt* crops: evidence versus theory. *Nature biotechnology* **26**: 199.

Torriani SFF, Melichar JPE, Mills C, Pain N, Sierotzki H, Courbot M. **2015**. *Zymoseptoria tritici*: a major threat to wheat production, integrated approaches to control. *Fungal Genetics and Biology* **79**: 8–12.

Tyutyunov Y, Zhadanovskaya E, Bourguet D, Arditi R. **2008**. Landscape refuges delay resistance of the European corn borer to *Bt*-maize: a demo-genetic dynamic model. *Theoretical Population Biology* **74**: 138–146.

Weber KE, Diggins LT. **1990**. Increased selection response in larger populations. II. Selection for ethanol vapor resistance in *Drosophila melanogaster* at two population sizes. *Genetics* **125**: 585–597.

zur Wiesch PA, Kouyos R, Engelstädter J, Regoes RR, Bonhoeffer S. **2011**. Population biological principles of drug-resistance evolution in infectious diseases. *The Lancet infectious diseases* **11**: 236–247.

Zhan J, McDonald BA. **2004**. The interaction among evolutionary forces in the pathogenic fungus *Mycosphaerella graminicola*. *Fungal Genetics and Biology* **41**: 590–599.

793   **Zhan J, Mundt CC, McDonald BA**. **2001**. Using restriction fragment length polymorphisms

794   to assess temporal variation and estimate the number of ascospores that initiate epidemics

795   in field populations of *Mycosphaerella graminicola*. *Phytopathology* **91**: 1011–1017.

Phenotype		First year observed	Number of years observed	Number of regions monitored	Number of regions monitored yearly	Number of observations
StrR		2004	9	16	9	852
TriMR	Total	2005	7	16	13	754
	TriR6	2006	12	16	6	910
	TriR7-TriR8	2006	12	16	6	851
TriHR		2010	8	14	6	360
Total						**3727**

797   **Table 1** Data used for the statistical analysis.

798   All regions were monitored for at least 80% of the years studied.

| | Phenotype resistant to QoIs | | | | Phenotype resistant to DMIs | | | | | | | | |
| | StrR (2004-2012) | | | | TriMR (≤2011) | | | TriR6 (2006-2017) | | | TriR7-TriR8 (2006-2017) | | |
Variable	Estimate ± 1 SD	EFC	Relative weight		Estimate ± 1 SD	EFC	Relative weight	Estimate ± 1 SD	EFC	Relative weight	Estimate ± 1 SD	EFC	Relative weight
Growth constant (β)	-0.2 * ± 0.09	-3.74	0.039		-0.2 · ± 0.11	-3.94	0.127	0.15 ** ± 0.05	3.58	0.402	0.09 · ± 0.05	1.63	0.402
Fungicide use (ν_f)			0.874				0.726			0.531			0.433
QoI Kresoxim-methyl	0.7 *** ± 0.14	4.26		DMI Cyproconazole	0			0			0.58 ** ± 0.2	2.6	
Pyraclostrobin	0.5 ** ± 0.17	3.29		Epoxiconazole	0.56 ** ± 0.18	4.82		0			0		
Azoxystrobin	0			Prochloraz	0			0.85 *** ± 0.16	4.73		-1.17 *** ± 0.17	-3.99	
Fluoxastrobin	0			Propiconazole	0			-0.43 ** ± 0.14	-2.48		0		
Picoxystrobin	0			Tebuconazole	0			-1.04 *** ± 0.19	-8.38		0.44 * ± 0.21	2.38	
Trifloxystrobin	0			Fluquinconazole	0			0			0		

	StrR				TriMR			TriR6			TriR7-TriR8		
				Hexaconazole	0			X			X		
				Metconazole	0			0			0		
				Prothioconazole	0			0			0		
Yield losses (ρ)	0		**0**		0		**0**	0		**0**	0		**0**
Area under organic farming (κ)	0.64 *** ± 0.14	*3.26*	**0.068**		0.37 *** ± 0.01	*2.52*	**0.133**	0		**0**	0		**0**
Wheat cultivar (σ_γ)	0.26 ** ± 0.08	*±4.68*	**0.005**		0.14 * ± 0.06	*±2.64*	**0.006**	0.21 ** ± 0.07	*±5.19*	**0.024**	0.37 *** ± 0.09	*±6.18*	**0.134**
Sampling date (δ_{S1})	-0.41 *** ± 0.1	*-8.16*	**0.014**		-0.17 ** ± 0.06	*-3.34*	**0.008**	-0.33 *** ± 0.08	*-8.23*	**0.043**	0.15 · ± 0.08	*2.62*	**0.031**

Table 2 Estimates from the M_{AI} model for the StrR, TriMR, TriR6 and TriR7-TriR8 resistance phenotypes. For each phenotype, the three subcolumns show parameter estimates (*i.e.* posterior mean) and their variability (*i.e.* posterior standard deviation), expected frequency change (EFC, *i.e.* the change in frequency due to the mean value of the variable in a mean frequency population), and relative weights (*i.e.* the contribution of the variable to data variability). The significance thresholds are 0.1, 0.05, 0.025 and 0.001 denoted by ".", "*", "**" and "***", respectively. Rows represent the different parameters of the models, ordered as follows: growth constant, regional scale explanatory variables (fungicide use, yield losses and area of wheat under organic farming) and local variation factors (wheat cultivar and sampling date). "0" means that the parameter was not selected during variable selection. "X" means that the variable was not considered in the model for the given phenotype. The results for the TriHR phenotypes are not shown as no explanatory variables were

808   retained by the selection procedure.

Variable	StrR (2004-2012)		
	Estimate ± 1 SD	EFC	Relative weight
Growth constant (β)	-0.17 ± 0.13	-3.23	0.041
Fungicide use (ν)	1.07 *** ± 0.26	6.64	0.794
Yield losses (ρ)	0		0
Area under organic farming (κ)	0.78 *** ± 0.16	3.95	0.146
Wheat cultivar (σ_γ)	0.31 ** ± 0.09	±5.57	0.009
Sampling date (δ_{S1})	-0.35 *** ± 0.1	-6.87	0.01

Table 3 Estimates from the M_{MoA} model for the StrR phenotype.

The three subcolumns show the parameter estimates (*i.e.* posterior mean) and their variability (*i.e.* posterior standard deviation), expected frequency change (EFC, *i.e.* the change in frequency due to the mean value of the variable in a mean frequency population), and relative weights (*i.e.* the contribution of the variable to data variability). The significance thresholds are 0.1, 0.05, 0.025 and 0.001 denoted by ".", "*", "**" and "***", respectively. Rows represent the different parameters of models, ordered as follows: growth constant, regional scale explanatory variables (fungicide use, yield losses and area of wheat under organic farming) and local variation factors (wheat cultivar and sampling date). "0" means that the parameter was not selected during variable selection. The results for TriMR and TriHR resistance phenotypes are not shown as no explanatory variable was retained by the

selection procedure.

Fig. 1 Changes in resistance frequencies (left) and explanatory variables (right, top to

bottom, fungicide use, yield losses and area of wheat under organic farming).

Bold lines: mean regional values. Shaded areas: quantiles of regional values (*i.e.* regional

variability), 25% and 75% (dark grey), 2.5% and 97.5% (light grey). Dotted lines: regional

minimum and maximum values. Fungicide use is expressed in $ha\frac{a_D}{c}$ corresponding to the

mean number of times each mode of action was used for spraying over a cropping season,

regardless of the dose used. Yield losses are expressed in decitons (quintals) per hectare.

Fig. 2 Maps of observed and predicted frequencies of the StrR resistance phenotype from

2004 to 2011.

Real observations are represented by dots. The colour within the dot indicates the observed

frequency in trials. The background map color shows the regional prediction from the M_{AI}

model.

Fig. 3 Maps of observed and predicted frequencies of TriR6 and TriR7-TriR8 resistance

phenotypes in 2008.

Real observations are represented by dots. The colour within the dot indicates the observed

frequency in trials. The background map colour shows the regional prediction from the M_{AI}

model. Top: TriR6. Bottom: TriR7-TriR8.

TriR6

TriR7-TriR8

100km

Frequency (%)
0   25   50   75   100