**Large-scale study validates that regional fungicide applications are major**

**determinants of resistance evolution in the wheat pathogen *Zymoseptoria tritici* in**

**France.**

**Running head:**

Resistance selection driven by regional fungicide use in French *Zymoseptoria tritici*

populations.

**Article type:**

Original research article – Full paper

**Authors:**

Maxime GARNAULT[1,2], Clémentine DUPLAIX[1], Pierre LEROUX[1], Gilles COULEAUD[3], Olivier

DAVID[2]*, Anne-Sophie WALKER[1]* and Florence CARPENTIER[1,2]*

[1] Université Paris-Saclay, INRAE, AgroParisTech, UMR BIOGER, 78850, Thiverval-Grignon,

France

[2] Université Paris-Saclay, INRAE, MaIAGE, 78350, Jouy-en-Josas, France

[3] Arvalis-Institut du Végétal, 91720 Boigneville, France

* These authors contributed equally to this study

**Author for correspondence:**

*Florence Carpentier*

*Tel: +331 44 08 86 35*

*E-mail: florence.carpentier@inrae.fr*

**Summary:**

• Research rationale: In modern cropping systems, the quasi-systematic use of plant

protection products selects for resistance in pest populations. The emergence and

evolution of this adaptive trait threaten treatment efficacy. We identified determinants

of fungicide resistance evolution and quantified their effects at a large spatiotemporal

scale.

• Methods: We focused on *Zymoseptoria tritici*, which causes leaf blotch in wheat.

Phenotypes of qualitative or quantitative resistance to various fungicides were

monitored annually, from 2004 to 2017, at about 70 sites throughout 22 regions of

France (territorial units of 25 000km² on average). We modelled changes in

resistance frequency with regional anti-*Septoria* fungicide use, yield losses due to the

disease and the regional area under organic wheat.

• Key results: The major driver of resistance dynamics was fungicide use at the

regional scale. We estimated its effect on the increase in resistance and relative

apparent fitness of each resistance phenotype. The predictions of the model

replicated the spatiotemporal patterns of resistance observed in field populations ($R^2$

from 0.56 to 0.82).

• Main conclusion: The evolution of fungicide resistance is mainly determined at the

regional scale. This study therefore showed that collective management at the

regional scale could effectively complete local actions.

**Keywords:**

*Mycosphaerella graminicola*, *Zymoseptoria tritici*, Septoria leaf blotch,

Resistance determinants, Large-scale evolution, Regional fungicide use, Resistance

dynamics, Mathematical modelling

**1 INTRODUCTION**

The efficacy of pesticides and drugs has been compromised by the rapid and widespread

evolution of resistance, increasing their use to maintain control (Georghiou & Mellon, 1983;

Russell, 2005; Gould *et al.*, 2018). Resistance management is essential for human health,

biodiversity and food security, given the rapid emergence and spread of resistance and the

lack of new modes of action (MoA) (Palumbi, 2001; Grimmer *et al.*, 2014). Many studies

have investigated the effects of various factors on the evolution of resistance: fitness cost

(Andersson, 2003), mutation rate (Martinez & Baquero, 2000; Gressel, 2011), population size

(Sisterson *et al.*, 2004), strength of selection pressure and its mitigation in strategies (Oz *et*

*al.*, 2014; van den Bosch *et al.*, 2014). A number of studies have advocated further work on

the relative impact of these factors on a given pest and of the interactions between these

factors (Berendonk *et al.*, 2015; Hughes & Andersson, 2015), with a view to promoting large-

scale strategies (Okeke *et al.*, 2005; Menalled *et al.*, 2016).

Several studies have shown how agricultural selection pressures affect the large-scale

structure of pest populations at national scale. For instance, the national distribution of

resistance varieties shapes the adaptation of pathogen populations to cultivars (Tyutyunov *et*

*al.*, 2008; Papaïx *et al.*, 2011). Historical herbicide applications drive the evolution of

herbicide resistance at a national scale (Hicks *et al.*, 2018). For fungicide resistance,

theoretical studies have revealed that combining effective MoAs over time and space can

delay resistance evolution (REX Consortium, 2013; van den Bosch *et al.*, 2014) and that

large-scale management strategies may differ from and interact with in-field strategies

(Parnell *et al.*, 2006). However, so far, in the absence of large-scale studies,

recommendations about fungicide use mostly stem from empirical studies conducted in local

field trials assessing the impact of different spraying strategies (Rosenzweig *et al.*, 2008;

Dooley *et al.*, 2016a,b; Heick *et al.*, 2017). In France, the metropolitan area was until 2016

subdivided into 22 administrative regions. Their average area of 25,000 square kilometres is

consistent with the long-distance dispersal capabilities of pathogens, e.g. between ten to

hundred kilometres for airborne fungi (Brown & Hovmøller, 2002; Linde *et al.*, 2002).

Homogeneous agronomic properties and pedoclimatic conditions generally lead to similar

crop sequences in regions (Xiao *et al.*, 2014). Therefore, this scale seems relevant for both

studying the evolution of resistance and implementing large-scaled resistance management.

The aim of this study was to highlight the determinants at the regional scale of fungicide

resistance evolution in France. First, we considered (i) the regional use of fungicides, since

the quantity of fungicide sprayed increases the selection of resistances (van den Bosch *et*

*al.*, 2014), (ii) the refuge effect (Parnell *et al.*, 2006; Tabashnik *et al.*, 2008) by including the

area under organic farming, as a proxy of unsprayed fields, since  the heterogeneity of the

selection may reduce the resistance growth (REX Consortium, 2010, 2013)   (iii) the

population size, using yield losses as a proxy, because it may affect the evolution of

resistance alleles through genetic drift (Maxwell *et al.*, 1990; Sisterson *et al.*, 2004). We did

not consider local field-scaled strategies (e.g. adjustment of spray timing, fungicides mixture

or alternation).

We focused our analysis on *Zymoseptoria tritici* (formerly *Septoria tritici* and *Mycosphaerella*

*graminicola* as teleomorph), an ascomycete responsible for septoria leaf blotch (STB) on

winter wheat. *Z. tritici* has many features facilitating the emergence of resistance: high

genome plasticity, a large population size, high genetic diversity, asexual and sexual

reproduction, an ability to disperse over large distances (Zhan & McDonald, 2004; Croll &

McDonald, 2012). STB is a major wheat disease that can cause yield losses of up to 50%

(Ponomarenko *et al.*, 2011; Torriani *et al.*, 2015). In western Europe, up to 70% of all

fungicide use is linked to STB control (Fones & Gurr, 2015). As a result, various degrees of

resistance to all authorised unisite inhibitors (*i.e.* exhibiting a single molecular mode of

action) have been observed in France (Garnault *et al*., 2019).

We previously published an initial analysis of the Performance trial network dataset, in which

phenotypes of resistance to four fungicide MoAs were monitored annually, from 2004 to

2017, at about 70 sites throughout France (Garnault *et al.* 2019). We found significant

differences between phenotypes in terms of changes in spatial distribution and/or growth

rates. Major differences in population structure and dynamics were highlighted between the

North and South of France.

We develop here an explanatory model for identifying the determinants of these regional

spatiotemporal heterogeneities in resistance evolution according to resistance phenotype.

We investigated the effect of three potential drivers of selection pressure through the (*i*)

regional fungicide uses and (*ii*) fraction of sprayed fields, and we also considered (*iii*) the

effect of the pathogen population size which may affect evolution through genetic drift. Our

analysis shows that the inter-annual change in resistance frequency can be assessed at

regional scale, and that the major determinant of resistance is the selection pressure exerted

by fungicide applications in the preceding year. This study provides empirical results for

regional resistance management, at a level intermediate between field and national

recommendations. A sound understanding of resistance evolution and of its determinants

would help optimizing resistance management and applying them at sound spatiotemporal

scales. It should ultimately help to reduce pesticide use in agrosystems.

**2 MATERIALS AND METHODS**

**2.1 Data description**

**2.1.1 Sampling of *Z. tritici* populations and estimation of resistance frequency**

The "Performance network" is supervised by ARVALIS-Institut du Végétal and the INRAE

research institute at Thiverval-Grignon. It carried out field trials on wheat throughout France

between 2004 and 2017 with a mean of 70 trials annually (4 to 5 trials per region and year on

average; 90% credible interval is 1 to 10). The frequency of resistant phenotypes in *Z. tritici*

populations sampled annually in these trials is recorded in the associated dataset (see

Garnault *et al.*, 2019 for further information).

Wheat trials were carried out in a randomized block design with 3 to 4 replicates. The

frequency of resistant phenotypes in populations was estimated by collecting bulk

pycnidiospores from 30 to 40 upper leaves that were randomly sampled within each plot and

showed STB symptoms. Cropped cultivars were predominantly STB-sensitive to promote the

presence of the disease. A total of 124 different wheat cultivars were cropped over the

dataset. Phenotypes were distinguished on the basis of their germination or growth on Petri

dishes containing discriminatory doses of fungicides, optimised on individual genotyped

isolates (see Leroux & Walker, 2011 and Garnault *et al.*, 2019 for more details). We then

considered: (*i*) the phenotype displaying specific qualitative resistance to strobilurins (or

QoIs; inhibitors of respiration complex III), hereafter referred to as the StrR phenotype, (*ii*) the

group of phenotypes with moderate quantitative resistance to DMIs (sterol 14α-demethylation

inhibitors), hereafter referred to as TriMR phenotypes, (*iii*) the group of phenotypes with a

high quantitative resistance to DMIs, hereafter referred to as TriHR phenotypes. The TriMR

group encompasses the TriR6 and TriR7-TriR8 phenotypes, which were also included in the

analysis (TriR6 strains grew on low doses of prochloraz, contrasting with the lack of growth of

TriR7–TriR8 strains in these conditions; Leroux & Walker, 2011).

Region, year, sampling date and cultivar grown were recorded for each sample. We

considered only populations from unsprayed plots for this study. The plots were sampled two

times: at "S1" in April-May, at about the Z32 wheat stage ($n$=1320, from 2006 to 2011), and

at "S2" in May-June, at about the Z39-Z55 wheat stage ($n$=2407, from 2004 to 2017).

In this study, we focused on  resistance selection (*i.e.* after emergence and before complete

fixation). For each phenotype, we defined a time interval grouping the years for which at least

a quarter of the regions exhibited frequencies differing from 0% and 100%. We excluded the

years following the "decline" of resistance,  identified as the year when the frequency of

resistance was no longer significantly higher than the average frequency. These frequency

differences were estimated in Garnault *et al.* (2019). The year 2012 was the first year of

decline for TriMR (other phenotypes did not show a decline). Time periods were respectively

2004 to 2012 ($n$=852, 16 regions) for the StrR phenotype, 2005 to 2011 ($n$=754, 16 regions)

for the TriMR phenotype group, and 2010 to 2017 ($n$=360, 14 regions) for the TriHR

phenotype group. We also included data from 2006 to 2017 for the TriR6 and TriR7-TriR8

phenotypes ($n$=910 and $n$=851, respectively), for analysis of the spatial heterogeneity of their

frequencies. The data are summarised in Table 1.

**2.1.2 Regional fungicide use**

Every year, Bayer Crop Science uses field surveys to estimate the area of wheat sprayed

with fungicides containing anti-STB active ingredients (AIs) in each region. These data do not include dose information but provide information about the areas sprayed with each AI and the number of sprayings (areas result from the multiplication of these two values). These areas are expressed in deployed hectares.

We retained the most widely used AIs for each MoA (AIs accounting cumulatively for more than 95% of the use of the MoA), to prevent background noise from AIs with a limited impact on STB control. The model therefore included pyraclostrobin (26%), azoxystrobin (19%), trifloxystrobin (15%), kresoxim-methyl (15%), fluoxastrobin (12%) and picoxystrobin (12%) for QoIs; and epoxiconazole (30%), prochloraz (17%), tebuconazole (13%), cyproconazole (11%), prothioconazole (10%), propiconazole (7%), metconazole (7%), fluquinconazole (2%) and hexaconazole (1%) for DMIs.

We took the regional heterogeneity in wheat production between regions (and, hence, in the area sprayed with fungicides) into account, by dividing the number of deployed hectares by the regional area under conventionally farmed wheat. The latter was calculated by subtracting the area under organic wheat (see section 2.1.4) from the total area under wheat (from the AGRESTE online data: agreste.agriculture.gouv.fr) for each year and region. This new variable unit was named $ha\frac{D}{C}$ ($D$ for deployed and $C$ for cropped hectares), and was proportional to the mean number of times each AI was used over a cropping season in a given region. The national trend and the regional heterogeneity of fungicide use expressed in $ha\frac{D}{C}$ are shown for DMIs and QoIs in Fig. 1. Henceforth, this variable is denoted $P_{itf}$, with $f$ corresponding to the AI, $t$ to the year and $i$ to the region.

### 2.1.3 Yield losses induced by STB

ARVALIS-Institut du Végétal assessed annual yield loss by conducting paired plot experiments throughout France with a mean of 80 trials (21 observations per region and year

on average, distributed among 3 to 4 trials; 90% credible interval is 9 to 48), from 2004 to

2017, in 20 French regions (Arvalis, 2019). In each trial, we considered modalities cropped

with STB-susceptible wheat cultivars, in both unsprayed plots and sprayed plots (providing

maximum protection against diseases). These cultivars were moderately to highly resistant to

rusts and yield losses were mostly attibuted to STB. Yield losses due to STB were calculated

by subtracting the yield in the unsprayed plot from that in the sprayed plot. Based on these

data, we predicted regional yield losses for each year with a linear model (fixed effects: year,

region; random effects: wheat cultivar, trial). The national trend and the regional

heterogeneity of yield losses, expressed in decitons per hectare, are shown in Fig. 1. This

variable is denoted $P_{it}$ hereafter, with $t$ corresponding to the year and $i$ to the region.

### 2.1.4 Proportion of the total area under wheat farmed organically

The area under organically farmed wheat crops was recorded by AgenceBIO (the French

national platform for the promotion and development of organic farming) and ARVALIS-

Institut du Végétal. We collected regional data from 2007 onwards, and national data from

2004 onwards. The regional areas under organic wheat between 2004 and 2006 were

assessed from the observed mean proportions of the regional area under organic wheat in

subsequent years and from national data for 2004 to 2006. We used the regional proportion

of wheat under organic farming in our models and calculated it by dividing the regional area

under organic wheat by the total area under wheat in the same region, based on AGRESTE

online data. The national trend and the regional heterogeneity of the area under organic

wheat, expressed in hectares, are shown in Fig. 1. This variable is denoted $R_{it}$ hereafter,

with $t$ corresponding to the year and $i$ to the region.

### 2.2 Statistical modelling

We modelled the change in frequency for each resistance phenotype in French populations.

The model took into account (*i*) the different phases of resistance dynamics (see below), (*ii*)

the effects of previously described potential regional determinants and finally (*iii*) variability

due to the sampling design (sampling date and wheat cultivar).

***Phases in resistance dynamics.*** We distinguished three phases in resistance dynamics:

"no resistance" (frequency equal to 0), "resistance selection" and "generalized resistance"

(frequency equal to 100). During the "resistance selection" phase, observations were

modelled with binomial random variables with a sample size of 100 (mean number of

observed spores used to determine frequencies). The probabilities that a population was in

the previous phases depended on the year $t$. These probabilities were referred as $\pi_{0t}$, $\pi_{100t}$

and $(1 - \pi_{0t} - \pi_{100t})$, respectively. Thus, $Y_{itjkn}$, the $n^{th}$ frequency observed in region $i,$ in

year $t,$ on cultivar $j$ and at sampling date $k$ followed a zero-and-one inflated binomial

distribution (Eqn 1).

**Eqn 1**

237
$$Y_{itjkn} \begin{cases} = 0 & with\,probability\,\pi_{0t} \\ \sim \mathcal{B}(100, p_{itjkn}) & with\,probability\,1 - \pi_{0t} - \pi_{100t} \\ = 100 & with\,probability\,\pi_{100t} \end{cases}$$

239    ***Resistance evolution.*** Using a logit transformation (Eqn 2), the proportion $p_{itjkn}$ of resistant

240    phenotypes during resistance evolution was modelled by the regional dynamics $D_{it}$ and the

241    variability due to sampling design $\zeta_{itjkn}$.

243    **Eqn 2**

244
$$logit(p_{itjkn}) = ln\left(\frac{p_{itjkn}}{1 - p_{itjkn}}\right) = D_{it} + \zeta_{itjkn}$$

***Regional dynamics.*** The change in resistance frequencies depended on regional-scale

variables: fungicide use, yield losses and areas under organic farming. The regional

dynamics $D_{i(t+1)}$ in region *i* at year *t+1* was obtained by adding the regional dynamics of the

previous year $D_{it}$ to the additive effects of fungicide use $\phi_{it}$, yield loss $\rho P_{it}$ and wheat area

under organic farming $\kappa R_{it}$ in year *t* (Eqn 3). $D_{i1}$ is related to the logit of the initial resistance

frequency in region *i*. We kept raw values for the fungicide use and area under organic wheat

(*i.e.* neither centered nor reduced), while we centered values for the yield losses. Thus, the

parameter β, the growth constant, corresponded to a continuous shift in resistance frequency

in the absence of fungicides and refuges, with an average population size. This parameter

will be interpreted as the relative apparent fitness.

**Eqn 3**

$$D_{i(t+1)} = D_{it} + \beta + \phi_{it} + \rho P_{it} + \kappa R_{it} \quad with\ t \geq 1$$

We derived two models from Eqn 3: one in which the use of fungicides is specified for each

AI within MoAs, and another in which only the global use of each MoA (*i.e.* sum of AIs uses)

is considered.

In the first model, **M**$_{AI}$, the term $\phi_{it}$ from Eqn 3 was defined as in Eqn 4:

**Eqn 4**

$$\phi_{it} = \sum_{f \in F} v_f F_{itf}$$

$F_{itf}$ corresponds to the fungicide use for a specific AI *f,* in region *i* and year *t* (see section

2.1.2). The set $F$ included all AIs in a specific MoA associated with considered resistance:

QoIs for the StrR phenotype, DMIs for the TriR phenotypes. We assumed that fungicide use

positively selected resistant phenotypes over susceptible or less sensitive phenotypes. Thus,

the parameters associated with fungicide use were, by definition, positive (*i.e.* $v_f \geq 0$), except

for the TriR6 and TriR7-TriR8 phenotypes, which were not the most DMI-resistant

phenotypes over their study period.

In the second model, $\mathbf{M}_{MoA}$, we considered: $F_{it.} = \sum_{f \in \mathcal{F}} F_{itf}$, the regional use of a given MoA.

The term $\phi_{it}$ from Eqn 3 was simplified and written: $\phi_{it} = \nu * F_{it.}$ This model was run only for

the StrR, TriMR and TriHR phenotypes, with, as above, the constraint $\nu \geq 0$.

**Observation variability.** The variability $\zeta_{itfkm}$ of observations in Eqn 2 was modelled with a

mixed model (Equation 5).

**Equation 5**

$$\zeta_{itfkm} = \gamma_j + \delta_k + \varepsilon_{itfkm}$$

The parameter $\gamma_j$ corresponds to the random effect of the 124 wheat cultivars ($\gamma_j$ drawn from

a centered Gaussian distribution with standard deviation $\sigma_\gamma$). The parameter $\delta_k$ is the fixed

effect of sampling date $k$ (*i.e.* "S1" or "S2", see section 2.1.1, with contrast $\delta_{S2} = 0$). Finally,

$\varepsilon_{itfkm}$ is the overdispersion, modelled as a random individual effect with a mean of 0 and a

standard deviation of $\sigma$.

**2.3 Parameter expression**

The explanatory variables had different units (*e.g.* proportion of wheat under organic farming

*vs.* fungicide use). Moreover, the interpretation of the parameters of the zero-one-inflated

logistic regression was not straightforward. We simplified the interpretation, by defining the

*expected frequency difference* (EFD) for each variable and phenotype. The EFD described

the frequency shift due to the mean value of this variable over a population with the mean

resistance frequency. It is calculated as $EFD = (p^e - \bar{p}) * 100$ (*i.e.* the difference between

$p^e$, the *expected frequency*, and $\bar{p}$, the *mean frequency* of the resistance phenotype in the

data). The computation of the expected frequencies are detailed in Table 2.

**2.4 Bayesian analysis**

Statistical analyses were performed with *R* software (R Development Core Team, 2008), in a

Bayesian framework, with the *rjags* package (Plummer, 2013).

***Prior and posterior densities.*** Non-informative prior distributions were used (Supporting

Information, Eqn S1). Posterior distributions were estimated by Monte Carlo-Markov chain

(MCMC) methods. Five MCMC chains were run, over 1 000 000 iterations, with a burn-in of

100 000 and a thinning every 1 000 for the variable selection phase (see the following

section), followed by 500 000 iterations with a burn-in of 50 000 and a thinning every 500 for

the final parameter estimation. Convergence was assessed with the Gelman and Rubin $\hat{R}$

statistic (Gelman *et al.*, 2004). Credible intervals of the highest posterior density were

calculated from posterior densities with the *HDI* package (Dezeure *et al.*, 2015). Parameter

estimates were considered significant at the 5% level (or the 2.5% or 0.1% level), if their 95%

credible interval (97.5% and 99.9%, respectively) did not contain the 0 value.

***Variable selection***. A selection procedure identified the relevant variables in each model for

each resistance phenotype. We used a method based on indicator variables (Kuo & Mallick,

1998), in which each predictor was multiplied by a dummy variable with a prior distribution

corresponding to a Bernoulli distribution with parameter $p$ = 0.5. A predictor was retained in

the model if the posterior expectation of its indicator variable was greater than 0.75 (thus,

greater than its prior expectation of 0.5).

***Predictive check.*** We assessed the fit of the model to the data by posterior predictive

checks (Gelman *et al.*, 2004). Replicated data ($y_{icjkw}^{rep}$) generated during the MCMC algorithm

from the model posterior densities were compared to observed data ($y_{icjkn}$). The mean value

of $PP_{icjkn}^{check} = P(y_{icjkn} - y_{icjkn}^{rep} < 0 \mid Y)$, where $Y$ is the vector of observations, was calculated

and denoted $PP_{check}$. This value indicated the goodness of fit of the model, with a good fit

corresponding to $PP_{check} = 0.5$, *i.e.* with equal probabilities of over- and under-estimation

***Variable weight.*** We assessed the influence of each explanatory variable $\theta$ by calculating its

weight ($W_\theta$). The weight $W_\theta$ was defined as the ratio of $RSS_{full-\theta}$ to $RSS_{full}$, where $RSS_{full}$

and $RSS_{full-\theta}$ are the residual sums of squares of the full model (*i.e.* including all the

explanatory variables selected by the variable selection procedure) and of this same model

but without the explanatory variable $\theta$, respectively. $RSS$ should be minimal for the full model,

so removing an explanatory variable should increase $RSS$: the more information $\theta$

contributes, the greater the increase in $RSS$ and the higher the value of $W_\theta$. Conversely, if the

information provided by $\theta$ is negligible, $RSS$ is unaffected and $W_\theta$ is minimal (*i.e.* close to 1).

In the result tables, we have calculated the relative weights by dividing individual variable

weights by the sum of the weights of all variables.

***Model comparison.*** For comparison of the $M_{AI}$ and $M_{MoA}$ models, we calculated the deviance

information criterion, DIC (Plummer, 2013), and the coefficient of determination, $R^2$.

***Predicted data.*** We computed predictions of the resistance frequencies for each phenotype

for a given region $I$, and a given year $T$ (Equation 6).

**Equation 6**

$$\Pr(\widetilde{Y}_{IT} = 1) = [1 - (\hat{\pi}_{0T} + \hat{\pi}_{100T})] \times logit^{-1}[\hat{D}_{I1} + \hat{\beta}(T-1) + \sum_{t=1}^{T-1}(\hat{\beta}P_{It} + \hat{R}R_{It} + \sum_{f \in F}\hat{\nu}_f F_{Itf})] + \hat{\pi}_{100T}$$

where parameter-hat are parameter estimates (*i.e.* their posterior mean). We therefore built

maps of resistance status, for known initial frequencies, use of fungicides, yield losses and

areas under organic wheat, from year 1 to year *T-1* in region *I*.

**3 RESULTS**

**3.1 Overview of model fits**

The convergence of the MCMC chain was satisfactory for all models (*i.e.* the Gelman and

Rubin indicator $\hat{R}$ was below 1.1 for all parameters, in all models) and model fit was good

($PP_{check}$ always between 0.498 and 0.511).

After the selection procedure, no effect was retained for the following models: $M_{AI}$ for the

TriHR  phenotyeand $M_{MoA}$ for the TriMR and the TriHR groups of phenotypes. The effects of

fungicide use, yield losses, and areas of wheat under organic farming were not significant.

As all estimates of the parameters of interest were equal to 0, we do not show and discuss

the results of these models.

Finally, with Spearman's method, a few significant correlations were found between some AI

uses ($F_{itf}$) in model inputs, but no significant correlation between estimates ($\hat{v}_F$) was found in

model outputs (Supporting Information, Fig. S1). No correlation exceeded 0.4 in absolute

value between AI uses and organic surfaces, nor between AI uses and yield losses.

**3.2 Ranking of variable weight**

Regional fungicide use appeared to be the major factor driving resistance evolution. In the

$M_{AI}$ models, which explicitly considered each AI, regional fungicide use systematically had

the highest relative weight. For the StrR and TriMR phenotypes, it accounted for 87.4% and

72.6%, respectively. It accounted for 53.1% and 43.3% for the TriR6 and TriR7-TriR8

phenotypes, respectively (Table 3).  For the $M_{MoA}$ models, fungicide use, considered as the

sum of AI uses within the same MoA, was also the major determinant of the StrR phenotype

(associated with qualitative resistance to QoIs), accounting for 79.4% of the sum of variable

weights (Table 4). For the TriMR and TriHR groups of phenotypes, no explanatory variables

were selected for the $M_{MoA}$ models.

The growth constant was also an important parameter, albeit to a lesser extent. In $M_{AI}$

models, the weight of the growth constant was lower than that of regional fungicide use by a

factor of 5.7 times for TriMR phenotypes, 1.32 for the TriR6 phenotype, and 1.08 for the

TriR7-TriR8 phenotypes (Table 3). For the StrR phenotype, the growth constant ranked third,

with a weight lower than that of fungicide use by a factor of almost 20, for both models

(Tables 3 and 4).

The proportion of the area under wheat farmed organically had a high relative weight for the

StrR and TriMR phenotypes, but was not selected for TriR6 and TriR7-TriR8. For the StrR

phenotype, its relative weight was ranked second, at 14.6% and 6.8% in the $M_{MoA}$ and the $M_{AI}$

models, respectively (Tables 3 and 4). For TriMR phenotypes, the relative weight of the

wheat area under organic farming was about 13.3%, a value very similar to that for the

growth constant (Table 3). Yield loss was systematically excluded during the selection

procedure, for all models and phenotypes.

Sampling data and wheat cultivar, variables reflecting local variability, had only a low relative

weight, with values always below 5%, except for the wheat cultivar variable for the TriR7-

TriR8 phenotype, for which the value was 13.4% (Table 3).

**3.3 Effect of variables at the regional scale**

**3.3.1 Regional fungicide use**

For the StrR phenotype, in the $M_{MoA}$ model, the effect of the overall use of QoI fungicides was

highly significant ($\nu$ = 1.07, $P$ < 0.001) and the expected frequency difference (EFD) was

estimated at 6.64%. Thus, a year of an average use of QoIs would have led to an increase of

6.64 frequency point on an average population composed of $\bar{p}$=76% of StrR (Table 4).

In the $M_{AI}$ model, two fungicides from the six QoI AIs were selected: kresoxim-methyl and

pyraclostrobin. Their EFDs were similar: 4.26% ($\nu$ = 0.7, $P$ < 0.001) and 3.29% ($\nu$ = 0.5, $P$ <

0.025), respectively (Table 3). The $M_{MoA}$ and $M_{AI}$ models also had similar adequacies to data,

according to their DIC (3480.2 and 3468.8 respectively) and $R^2$ values (0.82 and 0.81,

respectively).

For TriMR phenotypes, the effect of DMI use was estimated only in the $M_{AI}$ model, as no

explanatory variable was selected in the $M_{MoA}$ model. One AI of the nine DMI fungicides was

selected: epoxiconazole ($\nu$ = 0.56, $P$ < 0.025) with an estimated positive EFD of 4.82% (with

$\bar{p}$=0.75 ,Table 3).

As the TriR6 and TriR7-TriR8 phenotypes were not the most resistant to DMI over the study

period, the effect of fungicide use was not constrained to be null or positive, and was

estimated only in the $M_{AI}$ model. For the TriR6 phenotype, three AIs from the nine DMI

fungicides were selected: prochloraz, with a positive EFD of 4.73% ($\nu$ = 0.85, $P$ < 0.001,

$\bar{p}$=0.54), propiconazole with a negative EFD of -2.48% ($\nu$ = -0.43, $P$ < 0.025) and

tebuconazole with a negative EFD of -8.38% ($\nu$ = -1.04, $P$ < 0.001). Thus, prochloraz use

increased the frequency of TriR6, whereas tebuconazole counterselected TriR6 strains. For

the TriR7-TriR8 phenotype, three AIs from the nine DMI fungicides were selected:

cyproconazole and tebuconazole, with positive EFDs of 2.6% ($\nu$ = 0.58, $P$ < 0.025) at 2.38%

($\nu$ = 0.44, $P$ < 0.05), respectively, and prochloraz, with a negative EFD of -3.99% ($\nu$ = -1.17,

$P$ < 0.001). Prochloraz and tebuconazole clearly had opposite selection effects on TriR6 and

TriR7-TriR8 phenotypes.

**3.3.2 Growth constant**

The growth constant EFD quantified the change in resistance in the absence of regional

fungicide use and unsprayed refuges, for a mean potential yield loss. It was therefore an

extrapolation of the relative apparent fitness (referred to hereafter simply as fitness) of the

phenotype considered (*i.e.* how much faster the phenotype would grew compared to the rest

of the population (Hartl & Clark, 1997) in a year without fungicide treatment). A negative

growth constant indicates a fitness cost, whereas a positive growth constant indicates a

fitness gain. For the StrR phenotype, the growth constant provided an indication of the

fitness of the resistant strains relative to the sensitive strains. The estimated fitness costs for

this phenotype were similar in the $M_{AI}$ (-3.74%; $\beta$ = -0.2, *P* < 0.05; Table 3) and $M_{MoA}$ (-

3.23%; $\beta$ = -0.17, not significant; Table 4) models. DMIs selected a large diversity of

phenotypes (TriLR, TriMR and TriHR groups), and no sensitive strains were detected during

the study period. For TriMR phenotypes, the model was estimated with data from 2007 to

2011, when the frequency of the TriHR phenotype was still negligible (Fig. 1). Thus, the

growth constant for TriMR phenotypes mostly compared their fitness with that of TriLR

phenotypes. It was estimated at -3.94%, of borderline significance ($\beta$ = -0.2, *P* < 0.1), and

was associated with a relative weight of 12.7% (Table 3). The TriMR group included the

TriR6 and TriR7-TriR8 phenotypes. For the TriR6 and TriR7-TriR8 strains, the model was

estimated with data from 2006 to 2017. However, the TriHR phenotype has been non-

negligible since 2014 (Figure 1). Thus, the growth constant for TriR6 strains compared their

fitness with that of all the other phenotypes in the population: TriLR, TriR7-TriR8 and TriHR.

The TriR6 growth constant was estimated at 3.58% ($\beta$ = 0.15, *P* < 0.025; Table 3). This

result may reflect a balance between a fitness cost of TriR6 relative to TriLR and TriR7-TriR8

strains, and a fitness gain relative to the TriHR phenotype. This rationale also applies to the

apparent fitness gain of TriR7-TriR8, estimated at +1.63% ($\beta$ = 0.09, *P* < 0.1; Table 3).

**3.3.3 Wheat area under organic farming**

The proportion of the area under wheat management by organic farming methods increased

resistance frequency, with an EFD estimated at 3.95% and 3.26% in the $M_{MoA}$ and $M_{AI}$

models, respectively, for the StrR phenotype ($\kappa = 0.78$ and 0.64, $P < 0.001$; Tables 3 and 4),

and at 2.52% for the TriMR phenotype in the $M_{AI}$ model ($\kappa = 0.37$, $P < 0.001$; Table 3). This

variable was not selected for the TriR6 and TriR7-TriR8 phenotypes.

**3.3.4 Yield losses**

The selection procedure did not retain the yield loss variable in any of the models.

**3.4 Prediction maps**

We mapped the predicted resistance frequencies for each year and for all phenotypes. Our

maps predicting StrR phenotype dynamics were mostly based on regional QoI fungicide use

and, to a lesser extent, the area under organic wheat, as explanatory variables (Fig. 2).

Predictions mimicked the spatial propagation from the north to the south of France observed

between 2004 and 2011 (as described by Garnault *et al.*, 2019). Based on regional DMI use,

our model accurately predicted (Fig. 3) the observed spatial partitioning of TriR6 strains,

which were found mostly in the north east of France, and TriR7-TriR8 phenotypes, which

were mostly localised in the south west (as described by Garnault *et al.*, 2019).

**4 DISCUSSION**

Here, we developed a model for identifying the determinants of fungicide resistance evolution

at the regional scale. The candidate explanatory variables were regional fungicide use,

pathogen population size (approximated by potential yield losses) and the fraction of refuges

(approximated by the fraction of fields under organic wheat in a region). We analysed

resistance frequencies in *Z. tritici* populations on winter wheat. Frequencies were monitored

by the Performance trial network over the national territory of French (2004–2017; $\sim$70

locations each year). We studied resistance against fungicides with two modes of action:

QoIs, associated with qualitative resistance (StrR phenotype), and DMIs, associated with

quantitative resistance (a continuum of multiple resistance phenotypes forming three main

groups: TriLR, TriMR and TriHR, with low, medium and high levels of resistance to DMIs,

respectively). The TriMR phenotypes encompassed two phenotypes: TriR6 and TriR7-TriR8.

***The regional use of fungicides is the main driver of resistance evolution at the plot***

***scale***

We demonstrated that regional fungicide use was a major determinant of the evolution of

resistance. Fungicide use data at regional level, without taking into account the use of

particular fungicides in particular fields, was sufficiently informative to explain resistance

dynamics. Fungicide selection is, therefore, a large-scale process, and an understanding of

the evolution of fungicide resistance requires consideration of the regional use of these

compounds.

For qualitative resistance (e.g. StrR phenotype), the global use of a MoA (*i.e.* summed uses

of the AIs of this MoA), can be considered as a good predictor. Distinguishing the effects of

individual AIs within the MoA did not improve the fit of the model. By contrast, for quantitative

resistance, it appeared to be necessary to consider each AI within the MoA separately, as

they may select for different phenotypes (*e.g.* antagonist effects of prochloraz, tebuconazole

on TriR phenotypes).

The AIs selected by the model were consistent with the history of fungicide use and/or with

the patterns of cross-resistance described for some phenotypes. For instance, kresoxim-

methyl was one of the first QoIs authorised in France (commercialised in 1997,

www.ephy.anses.fr) and was selected to explain the evolution of StrR phenotypes, whereas

more recent QoIs were not. The fungicides most used over the study period (*i.e.*

epoxiconazole for DMIs, pyraclostrobin and kresoxim-methyl for QoIs; Supporting

Information, Fig. S2-S3), were also selected in models. The large-scale effect of AIs was also

linked to the cross-resistance pattern of the phenotypes. The estimated selection effect of

epoxiconazole was consistent with resistance factors (RFs) of TriMR phenotypes to

epoxiconazole, being higher than those of TriLR phenotypes which were the second most

frequent resistance phenotype during this period (RFs described in Supporting Information,

Table S1 from Leroux & Walker, 2011). The selection for TriR6 and counterselection for

TriR7-TriR8 phenotypes induced by the prochloraz were consistent with RFs (6.7 and <1.5,

respectively), as well as the effect of counterselection for TriR6 and selection for TriR7-TriR8

phenotypes by tebuconazole (RF=74 and 91, respectively for TriR6 and TriR8, TriR7 being

unfrequent (Huf *et al.*, 2018)). Propiconazole and cyproconazole were often used together as

a mixture, so were strongly correlated (Supporting Information, Fig. S1). The effects of

counterselection for TriR6 phenotypes by propiconazole and of selection for TriR7-TriR8 by

cyproconazole may indicate that their mixture globally promoted the selection of TriR7-TriR8

strains over TriR6 strains. Again, this is consistent with RFs to propiconazole (35 and 54,

respectively for TriR6 and TriR8 strains) and to cyproconazole (11 and 13, respectively).

These findings also confirm that the laboratory characterisation of strains can be good

predictor of resistance evolution in the field, if properly used, as reported by Blake *et al.*

(2018).

Our findings highlight the importance of defining precise resistance profile phenotypes.

Indeed, the model fitted slightly better the frequencies on more homogeneous phenotypes

($R^2$ = 0.48 for TriMR vs. 0.56 for TriR6 and TriR7-TriR8 phenotypes). In addition, no

significant effect of the different DMIs was found for the TriHR phenotype group. Indeed,

TriHR strains encompass multiple heterogeneous phenotypes, resulting from combinations

of target alteration, target overexpression and enhanced efflux, as resistance mechanisms

(Leroux & Walker, 2011; Huf *et al.*, 2018). The tremendous diversity and redundancy of

phenotypes observed in the field, especially for TriHR phenotypes, made it possible to

classify strains only approximately. There is therefore a need to develop molecular tools for

quantifying genotypes rather than phenotypes. Multi-trait high-throughput genotyping should

quantify resistance frequency more accurately and ultimately lead to improvements in our

ability to predict resistance evolution.

***The proportion of the wheat area under organic farming may be still too limited to***

***mitigate the evolution of resistance*** *via a refuge effect*

Wheat areas managed under organic farming systems are not treated with synthetic

fungicides. As some conventional fields may be unsprayed with a given MoA, organic areas

represent a lower bound of the surfaces that were unsprayed with this MoA (only true for

QoIs and SDHIs, since DMIs are quasi-systematic in conventional wheat farming).They could

act as refuges for "wild" susceptible or less resistant individuals, which may reproduce,

delaying the evolution of resistance by a dilution effect in mobile species (Gould, 2000). But

refuges may also provide a heterogeneous environment, promoting sink-source dynamics:

selection-free wheat areas acting as a source of susceptible strains that migrate towards

areas farmed conventionally. These opposite effects have been described in studies of the

resistance to transgenic crops expressing *Bacillus thuringiensis* (*Bt*) toxins, where non-*Bt*

crops may act as refuges diluting the selection (Huang *et al.*, 2011), or promoting the

evolution of resistance (Caprio, 2001).

The refuge effect for *Z. tritici* is, theoretically, weak due to the haploid nature of this organism

(Shaw, 2009), but this has never been studied experimentally. We did not validate the

beneficial effect of refuges, approximated by the area under wheat farmed organically. On

the contrary, we estimated that the selection of StrR and TriMR phenotypes would increase

with wheat areas under organic farming.  However, the weight of this explanatory variable

remained much lower than that for regional fungicide use. In addition, the evolution of

surfaces under organic wheat was very different between South and North of France, with

both higher proportions and faster increase in Southern regions (Supporting Information Fig.

S4). Therefore, the effect of refuges remains difficult to distinguish from other potential

factors reflecting differences between Northern and Southern agriculture (e.g. drier climate,

higher proportion of durum wheat,  in Southern regions). This effect was not found significant

for the other phenotypes.

According to Huang *et al.* (2011), three conditions must be satisfied for refuge strategies to

be successful: selection at "high dose", a very low initial frequency of resistance, and

sufficient refuge areas located nearby. In our study, initial frequencies of resistance (*i.e.* at

the very beginning of our study) were already quite high (up to 85% in some regions, Fig. 1).

In addition, the area under organic farming may still be too small within the landscape

(generally less than 1% until 2010). Nevertheless, its proportion has steadily increased since

the 2010s (Fig. 1). The effect of refuges may become detectable in those areas, and should

be investigated further, particularly for emerging resistance phenotypes.

***The growth constant reveals a relative fitness penalty of phenotypes***

The growth constant represents the evolution of  phenotypes in the absence of fungicide

treatment. It represents the apparent fitness of the phenotype relative to that of the

susceptible phenotype (or other resistant phenotypes, in the case of quantitative resistance).

The term "apparent" is used because this quantification takes place in current crop

conditions. The fitness cost of resistance to drugs is known to be a key parameter driving

effective anti-resistance strategies (Andersson & Hughes, 2010; Melnyk *et al.*, 2015;

Mikaberidze & McDonald, 2015). If there is no fitness cost, regardless of the strategy used,

resistance will always be selected  irreversibly. However, it remains difficult to infer the global

fitness cost of a mutation throughout the entire life cycle of a pathogen (Hollomon, 2015).

We inferred an apparent relative fitness penalty for StrR phenotypes, resulting in an annual

decrease of 3.74%. This was consistent with the fitness cost described by Hagerty & Mundt

(2016) using virulence comparison tests. The impact of cyp51 alterations, leading to TriR

phenotypes, on fitness is often evoked to explain the evolution of azole resistance in *Z. tritici*

populations (Cools *et al.*, 2013; Blake *et al.*, 2018) but it has not been quantified as yet. Our

model is consistent with these assumptions as it also inferred an apparent fitness penalty for

the TriMR group (-3.94% per year). Decreases in the use of QoI and DMI fungicides, and/or

the implementation of strategies favouring the expression of a resistance cost, may help to

slow resistance evolution. From our estimates, we can extrapolate a theoretical equilibrium

between resistance cost and selection, which could have led to a null growth of the StrR and

TriMR phenotypes by reducing by 50% the use of QoIs and 18% for DMIs.

For the TriR6 and TriR7-TriR8 phenotypes, the growth constant is an integrative value, as

these phenotypes were studied alongside more susceptible phenotypes (TriLR before 2010)

and more resistant phenotypes (TriHR after 2010). The positive growth constants we

estimated for these two phenotypes may indicate a fitness benefit of TriMR relative to TriLR

phenotypes and/or fitness cost of TriHR (*i.e.* also fitness benefit of TriMR relative to TriHR

phenotypes). Our model could be extended to determine the relative fitnesses of each

phenotype in cases of quantitative resistance. The global informative indicator provided by

our model could be used to guide the design of optimal large-scale fungicide deployment

strategies.

***The yield losses caused by STB do not affect resistance evolution***

Population size, a major parameter in population adaptation (Good *et al.*, 2012), is generally

positively correlated with resistance evolution (Weber & Diggins, 1990; Anderson, 2005; zur

Wiesch *et al.*, 2011). Indeed, a large population size increases the number of mutants

generated and defavours genetic drift (Linde *et al.*, 2002). We inferred this effect using

potential yield losses caused by STB as a proxy, but no significant effect was detected for

any resistance phenotype. Population size may not be limiting for resistance evolution in *Z.*

*tritici*, since the number of individuals remains important even in years with little disease

(Zhan *et al.*, 2001; Mikaberidze *et al.*, 2017), particularly when considering large scales.

Population size may not be described accurately enough as our proxy variable also depends

on the timing of infection (Shaw & Royle, 1993) and on stubble management (McDonald &

Mundt, 2016).

***Predicting resistance evolution over years***

Prediction maps can be computed from our model, using only the initial regional frequencies of the phenotypes, the history of fungicide use (between the initial year and the year to be predicted) and the history of area under organic farming. Predictions for the StrR phenotype from 2004 to 2011 highlighted the same colonisation front structure from the North to the South of France (Fig. 2) as reported by Garnault *et al.* (2019) albeit the regions are assumed to be fully independent. Besides, the way StrR propagated from North to South may indicate that integrating regional interdependencies in the model would improve its performance. However, this would increase its complexity and require more data to estimate inter-regional fluxes. Predictions for the TriR6 and TriR7-TriR8 phenotypes also yielded stable spatial distributions between North-East and South-West France (Fig. 3), as previously observed in Garnault *et al.* (2019).

Further analysis will be required to assess the prediction quality of the model. Nevertheless, this finding supports the global validity of our model and paves the way for an original approach to predicting resistance evolution in a heterogeneous landscape.

***Conclusion***

We developed a model of resistance dynamics, which identified regional fungicide use as the major determinant of fungicide resistance evolution, and the area under organic farming as a much weaker explanatory variable. We estimated the apparent relative fitness of resistant phenotypes, a key parameter for the development of sustainable resistance management strategies. We also identified AIs whose use drove resistance evolution.

As variability of resistance evolution is mainly explained by the regional intensity of selection induced by fungicides (without information on local application strategies), concerted collective action on the regional or national fungicide use could improve control of the evolution of resistance, especially in the most at-risk regions.

**5 ACKNOWLEDGEMENTS**

This work was funded by the INRAE SMaCH metaprogramme (FONDU project) and by Bayer Crop Science (PhD scholarship of M. Garnault; CIFRE no. 2016-0695). We warmly thank all partners of the Performance network for providing samples since 2004, and Arvalis-Institut du Végétal for supervising the network. We are grateful to the INRAE MIGALE bioinformatics facility (MIGALE, INRAE, 2020. Migale bioinformatics Facility, doi: 10.15454/1.5572390655343293E12) for providing computing resources. We thank Eva Lacarce (AgenceBio) and Phillipe Viaux (Arvalis-Institut du Végétal) for providing information about the historical changes in the area under organic wheat, and Clarisse Payet (Bayer Crop Science) for providing data on fungicide use in France and for helpful discussions.

**6 AUTHOR CONTRIBUTIONS**

GC was responsible for Performance network administration and supervision and for establishing the dataset. PL, ASW and CD carried out laboratory manipulations on the network samples to determine resistance frequencies in *Z. tritici* populations. FC and OD were involved in the development of the methodology and provided theoretical support for statistical aspects for data analyses. MG handled data, coded the model, performed statistical analyses and generated most of the results (figures, maps and tables) during his PhD work. The article was written by MG, FC, ASW and OD.

**7 REFERENCES**

**Anderson JB**. **2005**. Evolution of antifungal-drug resistance: mechanisms and pathogen fitness. *Nature Reviews Microbiology* **3**: 547–556.

**Andersson DI**. **2003**. Persistence of antibiotic resistant bacteria. *Current opinion in microbiology* **6**: 452–456.

**Andersson DI, Hughes D**. **2010**. Antibiotic resistance and its cost: is it possible to reverse resistance? *Nature Reviews Microbiology* **8**: 260–271.

**Arvalis**. **2019**. *Choisir & Décider, Synthèse Nationale : Céréales à paille, Interventions de*

*printemps.*

**Berendonk TU, Manaia CM, Merlin C, Fatta-Kassinos D, Cytryn E, Walsh F, Bürgmann**

**H, Sørum H, Norström M, Pons M-N,** *et al.* **2015**. Tackling antibiotic resistance: the

environmental framework. *Nature Reviews Microbiology* **13**: 310.

**Blake JJ, Gosling P, Fraaije BA, Burnett FJ, Knight SM, Kildea S, Paveley ND**. **2018**.

Changes in field dose--response curves for demethylation inhibitor (DMI) and quinone

outside inhibitor (QoI) fungicides against *Zymoseptoria tritici*, related to laboratory sensitivity

phenotyping and genotyping assays. *Pest management science* **74**: 302–313.

**van den Bosch F, Oliver R, van den Berg F, Paveley N**. **2014**. Governing principles can

guide fungicide-resistance management tactics. *Annual Review of Phytopathology* **52**: 175–

195.

**Brown JKM, Hovmøller MS**. **2002**. Aerial dispersal of pathogens on the global and

continental scales and its impact on plant disease. *Science* **297**: 537–541.

**Caprio MA**. **2001**. Source-sink dynamics between transgenic and non-transgenic habitats

and their role in the evolution of resistance. *Journal of economic entomology* **94**: 698–705.

**Cools HJ, Hawkins NJ, Fraaije BA**. **2013**. Constraints on the evolution of azole resistance

in plant pathogenic fungi. *Plant Pathology* **62**: 36–42.

**Croll D, McDonald BA**. **2012**. The accessory genome as a cradle for adaptive evolution in

pathogens. *PLoS pathogens* **8**.

**Dezeure R, Bühlmann P, Meier L, Meinshausen N**. **2015**. High-Dimensional Inference:

Confidence Intervals, p-values and R-Software hdi. *Statistical Science* **30**: 533–558.

**Dooley H, Shaw MW, Spink J, Kildea S**. **2016a**. Effect of azole fungicide mixtures,

alternations and dose on azole sensitivity in the wheat pathogen *Zymoseptoria tritici*. *Plant*

*Pathology* **65**: 124–136.

**Dooley H, Shaw MW, Spink J, Kildea S**. **2016b**. The effect of succinate dehydrogenase

inhibitor/azole mixtures on selection of *Zymoseptoria tritici* isolates with reduced sensitivity.

*Pest management science* **72**: 1150–1159.

**Fones H, Gurr S**. **2015**. The impact of *Septoria tritici* Blotch disease on wheat: an EU

perspective. *Fungal Genetics and Biology* **79**: 3–7.

**Garnault M, Duplaix C, Leroux P, Couleaud G, Carpentier F, David O, Walker A-S**. **2019**. Spatiotemporal dynamics of fungicide resistance in the wheat pathogen *Zymoseptoria tritici* in France. *Pest management science* **75**: 1794–1807.

**Gelman A, Carlin JB, Stern HS, Rubin DB**. **2004**. Bayesian data analysis 2nd edn Chapman & Hall. *CRC, Boca Raton FL*.

**Georghiou GP, Mellon RB**. **1983**. Pesticide resistance in time and space. In: Pest resistance to pesticides. Springer, 1–46.

**Good BH, Rouzine IM, Balick DJ, Hallatschek O, Desai MM**. **2012**. Distribution of fixed beneficial mutations and the rate of adaptation in asexual populations. *Proceedings of the National Academy of Sciences* **109**: 4950–4955.

**Gould F**. **2000**. Testing *Bt* refuge strategies in the field. *Nature Biotechnology* **18**: 266.

**Gould F, Brown ZS, Kuzma J**. **2018**. Wicked evolution: Can we address the sociobiological dilemma of pesticide resistance? *Science* **360**: 728–732.

**Gressel J**. **2011**. Low pesticide rates may hasten the evolution of resistance by increasing mutation frequencies. *Pest management science* **67**: 253–257.

**Grimmer MK, van den Bosch F, Powers SJ, Paveley ND**. **2014**. Evaluation of a matrix to calculate fungicide resistance risk. *Pest management science* **70**: 1008–1016.

**Hagerty CH, Mundt CC**. **2016**. Reduced virulence of azoxystrobin-resistant *Zymoseptoria tritici* populations in greenhouse assays. *Phytopathology* **106**: 884–889.

**Heick TM, Justesen AF, Jørgensen LN**. **2017**. Anti-resistance strategies for fungicides against wheat pathogen *Zymoseptoria tritici* with focus on DMI fungicides. *Crop protection* **99**: 108–117.

**Hicks HL, Comont D, Coutts SR, Crook L, Hull R, Norris K, Neve P, Childs DZ, Freckleton RP**. **2018**. The factors driving evolved herbicide resistance at a national scale. *Nature ecology & evolution* **2**: 529.

**Hollomon DW**. **2015**. Fungicide resistance: facing the challenge-a review. *Plant Protection Science* **51**: 170–176.

**Huang F, Andow DA, Buschman LL**. **2011**. Success of the high-dose/refuge resistance
management strategy after 15 years of Bt crop use in North America. *Entomologia*
*Experimentalis et Applicata* **140**: 1–16.

**Huf A, Rehfus A, Lorenz KH, Bryson R, Voegele RT, Stammler G**. **2018**. Proposal for a
new nomenclature for CYP 51 haplotypes in *Zymoseptoria tritici* and analysis of their
distribution in Europe. *Plant Pathology* **67**: 1706–1712.

**Hughes D, Andersson DI**. **2015**. Evolutionary consequences of drug resistance: shared
principles across diverse targets and organisms. *Nature Reviews Genetics* **16**: 459.

**Kuo L, Mallick B**. **1998**. Variable selection for regression models. *Sankhya: The Indian*
*Journal of Statistics, Series B*: 65–81.

**Leroux P, Walker A-S**. **2011**. Multiple mechanisms account for resistance to sterol 14α-
demethylation inhibitors in field isolates of *Mycosphaerella graminicola*. *Pest management*
*science* **67**: 44–59.

**Linde CC, Zhan J, McDonald BA**. **2002**. Population structure of *Mycosphaerella*
*graminicola*: from lesions to continents. *Phytopathology* **92**: 946–955.

**Martinez JL, Baquero F**. **2000**. Mutation frequencies and antibiotic resistance. *Antimicrobial*
*agents and chemotherapy* **44**: 1771–1777.

**Maxwell BD, Roush ML, Radosevich SR**. **1990**. Predicting the evolution and dynamics of
herbicide resistance in weed populations. *Weed technology* **4**: 2–13.

**McDonald BA, Mundt CC**. **2016**. How knowledge of pathogen population biology informs
management of *Septoria tritici* blotch. *Phytopathology* **106**: 948–955.

**Melnyk AH, Wong A, Kassen R**. **2015**. The fitness costs of antibiotic resistance mutations.
*Evolutionary applications* **8**: 273–283.

**Menalled FD, Peterson RKD, Smith RG, Curran WS, Páez DJ, Maxwell BD**. **2016**. The
eco-evolutionary imperative: revisiting weed management in the midst of an herbicide
resistance crisis. *Sustainability* **8**: 1297.

**Mikaberidze A, McDonald BA**. **2015**. Fitness cost of resistance: impact on management. In:
Fungicide Resistance in Plant Pathogens. Springer, 77–89.

**Mikaberidze A, Paveley ND, Bonhoeffer S, van den Bosch F**. **2017**. Emergence of

resistance to fungicides: the role of fungicide dose. *Phytopathology* **107**: 545–560.

**Okeke IN, Klugman KP, Bhutta ZA, Duse AG, Jenkins P, O'Brien TF, Pablos-Mendez A,**

**Laxminarayan R**. **2005**. Antimicrobial resistance in developing countries. Part II: strategies

for containment. *The Lancet infectious diseases* **5**: 568–580.

**Oz T, Guvenek A, Yildiz S, Karaboga E, Tamer YT, Mumcuyan N, Ozan VB, Senturk GH,**

**Cokol M, Yeh P,** *et al.* **2014**. Strength of selection pressure is an important parameter

contributing to the complexity of antibiotic resistance evolution. *Molecular biology and*

*evolution* **31**: 2387–2401.

**Palumbi SR**. **2001**. Humans as the world's greatest evolutionary force. *Science* **293**: 1786–

1790.

**Papaïx J, Goyeau H, Du Cheyron P, Monod H, Lannou C**. **2011**. Influence of cultivated

landscape composition on variety resistance: an assessment based on wheat leaf rust

epidemics. *New Phytologist* **191**: 1095–1107.

**Parnell S, van den Bosch F, Gilligan CA**. **2006**. Large-scale fungicide spray heterogeneity

and the regional spread of resistant pathogen strains. *Phytopathology* **96**: 549–555.

**Plummer M**. **2013**. Rjags: bayesian graphical models using MCMC. *R package version* **3**.

**Ponomarenko A, Goodwin SB, Kema GHJ**. **2011**. *Septoria tritici* blotch (STB) of wheat.

*Plant Health Instructor* **10**.

**R Development Core Team**. **2008**. R: A Language and Environment for Statistical

Computing.

**REX Consortium**. **2010**. The skill and style to model the evolution of resistance to pesticides

and drugs. *Evolutionary Applications* **3**: 375–390.

**REX Consortium**. **2013**. Heterogeneity of selection and the evolution of resistance. *Trends*

*in ecology & evolution* **28**: 110–118.

**Rosenzweig N, Atallah ZK, Olaya G, Stevenson WR**. **2008**. Evaluation of QoI fungicide

application strategies for managing fungicide resistance and potato early blight epidemics in

Wisconsin. *Plant Disease* **92**: 561–568.

**Russell PE**. **2005**. A century of fungicide evolution. *The Journal of Agricultural Science* **143**:

11–25.

**Shaw MW**. **2009**. Fungicide resistance: the dose rate debate. *Outlooks on Pest Management*

**20**: 100–103.

**Shaw MW, Royle DJ**. **1993**. Factors determining the severity of epidemics of

*Mycosphaerella graminicola* (*Septoria tritici*) on winter wheat in the UK. *Plant Pathology* **42**:

882–899.

**Sisterson MS, Antilla L, Carrière Y, Ellers-Kirk C, Tabashnik BE**. **2004**. Effects of insect

population size on evolution of resistance to transgenic crops. *Journal of Economic*

*Entomology* **97**: 1413–1424.

**Tabashnik BE, Gassmann AJ, Crowder DW, Carrière Y**. **2008**. Insect resistance to *Bt*

crops: evidence versus theory. *Nature biotechnology* **26**: 199.

**Torriani SFF, Melichar JPE, Mills C, Pain N, Sierotzki H, Courbot M**. **2015**. *Zymoseptoria*

*tritici*: a major threat to wheat production, integrated approaches to control. *Fungal Genetics*

*and Biology* **79**: 8–12.

**Tyutyunov Y, Zhadanovskaya E, Bourguet D, Arditi R**. **2008**. Landscape refuges delay

resistance of the European corn borer to *Bt*-maize: a demo-genetic dynamic model.

*Theoretical Population Biology* **74**: 138–146.

**Weber KE, Diggins LT**. **1990**. Increased selection response in larger populations. II.

Selection for ethanol vapor resistance in *Drosophila melanogaster* at two population sizes.

*Genetics* **125**: 585–597.

**zur Wiesch PA, Kouyos R, Engelstädter J, Regoes RR, Bonhoeffer S**. **2011**. Population

biological principles of drug-resistance evolution in infectious diseases. *The Lancet infectious*

*diseases* **11**: 236–247.

**Xiao Y, Mignolet C, Mari J-F, Benoit M**. **2014**. Modeling the spatial distribution of crop

sequences at a large regional scale using land-cover survey data: A case from France.

*Computers and Electronics in Agriculture* **102**: 51–63.

**Zhan J, McDonald BA**. **2004**. The interaction among evolutionary forces in the pathogenic

fungus *Mycosphaerella graminicola. Fungal Genetics and Biology* **41**: 590–599.

**Zhan J, Mundt CC, McDonald BA**. **2001**. Using restriction fragment length polymorphisms

to assess temporal variation and estimate the number of ascospores that initiate epidemics

in field populations of *Mycosphaerella graminicola. Phytopathology* **91**: 1011–1017.

| Phenotype | | First year observed | Number of years observed | Number of regions monitored | Number of regions monitored yearly | Number of observations |
|---|---|---|---|---|---|---|
| StrR | | 2004 | 9 | 16 | 9 | 852 |
| TriMR | Total | 2005 | 7 | 16 | 13 | 754 |
| | TriR6 | 2006 | 12 | 16 | 6 | 910 |
| | TriR7-TriR8 | 2006 | 12 | 16 | 6 | 851 |
| TriHR | | 2010 | 8 | 14 | 6 | 360 |
| Total | | | | | | 3727 |

**Table 1** Data from the Performance network database used for the statistical analysis.

Each observation corresponds to a resistance frequency measurement. "First year

observed": first year of observation in the Performance network. "Number of years

observed": total number of successive years of monitoring, including the first year. "Number

of regions monitored": total number of sampled French regions. "Number of regions

monitored yearly": total number of French regions with at least one observation each year. All

regions were monitored for at least 80% of the studied time range. The last column sums up

the total number of observed frequencies for each phenotype.

| Variable | Parameter | Expected frequency, $\gamma^\varepsilon$ | |
|---|---|---|---|
| Fungicide use | $v_f$ | $logit\left(p^\varepsilon_{v_f}\right) = logit(\bar{p}) + \hat{v}_f\bar{F}_f$ | $\bar{F}_f$, the mean annual use of fungicide $f$ over all regions $\hat{v}_f$, the estimate of $v_f$ $\bar{p}$, the *mean frequency* of the resistance phenotype |
| Growth constant | $\beta$ | $logit(p^\varepsilon_\beta) = logit(\bar{p}) + \hat{\beta}$ | |
| Yield losses due to STB | $\rho$ | $logit(p^\varepsilon_\rho) = logit(\bar{p}) + \hat{\rho}\bar{P}$ | $\bar{P}$, the mean yield losses due to STB |
| Area of wheat under organic farming | $\kappa$ | $logit(p^\varepsilon_\kappa) = logit(\bar{p}) + \hat{\kappa}\bar{R}$ | $\bar{P}$, the mean yield losses due to STB |
| Sampling date | $\delta$ | $logit(p^\varepsilon_\delta) = logit(\bar{p}) + \hat{\delta}_{T0}$ | |
| Cultivar effect | $\sigma_\gamma$ | $logit\left(p^\varepsilon_{\sigma_\gamma}\right) = logit(\bar{p}) \pm \hat{\sigma}_\gamma$ | |

**Table 2** Computation of expected frequencies of resistant phenotype of an average population submitted to an average effect of an explanatory variable. The differences between the expected and the mean frequencies described the Expected Frequency Difference (EFD ) of a given variable.

| Variable | Phenotype resistant to QoIs | | | | Phenotype resistant to DMIs | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | StrR (2004-2012) | | | | TriMR (≤2011) | | | TriR6 (2006-2017) | | | TriR7-TriR8 (2006-2017) | | |
| | Estimate ± 1 SD | EFD ($\bar{p}$=76%) | Relative weight | | Estimate ± 1 SD | EFD ($\bar{p}$=75%) | Relative weight | Estimate ± 1 SD | EFD ($\bar{p}$=54%) | Relative weight | Estimate ± 1 SD | EFD ($\bar{p}$=21%) | Relative weight |
| Growth constant ($\beta$) | -0.2 * ± 0.09 | -3.74 | 0.039 | | -0.2 · ± 0.11 | -3.94 | 0.127 | 0.15 ** ± 0.05 | 3.58 | 0.402 | 0.09 · ± 0.05 | 1.63 | 0.402 |
| Fungicide use ($v_f$) | | | 0.874 | | | | 0.726 | | | 0.531 | | | 0.433 |
| QoI  Kresoxim-methyl | 0.7 *** ± 0.14 | 4.26 | | DMI  Cyproconazole | 0 | | | 0 | | | 0.58 ** ± 0.2 | 2.6 | |
| Pyraclostrobin | 0.5 ** ± 0.17 | 3.29 | | Epoxiconazole | 0.56 ** ± 0.18 | 4.82 | | 0 | | | 0 | | |
| Azoxystrobin | 0 | | | Prochloraz | 0 | | | 0.85 *** ± 0.16 | 4.73 | | -1.17 *** ± 0.17 | -3.99 | |
| Fluoxastrobin | 0 | | | Propiconazole | 0 | | | -0.43 ** ± 0.14 | -2.48 | | 0 | | |
| Picoxystrobin | 0 | | | Tebuconazole | 0 | | | -1.04 *** ± 0.19 | -8.38 | | 0.44 * ± 0.21 | 2.38 | |

| | StrR | | | | TriMR | | | TriR6 | | | TriR7-TriR8 | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Trifloxystrobin | 0 | | | Fluquinconazole | 0 | | | 0 | | | 0 | | |
| | | | | Hexaconazole | 0 | | | X | | | X | | |
| | | | | Metconazole | 0 | | | 0 | | | 0 | | |
| | | | | Prothioconazole | 0 | | | 0 | | | 0 | | |
| Yield losses ($\mu$) | 0 | | **0** | | 0 | | **0** | 0 | | **0** | 0 | | **0** |
| Area under organic farming ($\kappa$) | 0.64 *** ± 0.14 | 3.26 | **0.068** | | 0.37 *** ± 0.01 | 2.52 | **0.133** | 0 | | **0** | 0 | | **0** |
| Wheat cultivar ($\sigma_T$) | 0.26 ** ± 0.08 | ±4.68 | **0.005** | | 0.14 * ± 0.06 | ±2.64 | **0.006** | 0.21 ** ± 0.07 | ±5.19 | **0.024** | 0.37 *** ± 0.09 | ±6.18 | **0.134** |
| Sampling date ($\delta_{SL}$) | -0.41 *** ± 0.1 | -8.16 | **0.014** | | -0.17 ** ± 0.06 | -3.34 | **0.008** | -0.33 *** ± 0.08 | -8.23 | **0.043** | 0.15 · ± 0.08 | 2.62 | **0.031** |

**Table 3** Estimates from the $M_{AI}$ model for the StrR, TriMR, TriR6 and TriR7-TriR8 resistance phenotypes.

For each phenotype, the three subcolumns show parameter estimates (*i.e.* posterior mean) and their variability (*i.e.* posterior standard

deviation), expected frequency difference (EFD, *i.e.* the change in frequency due to the mean value of the variable on the reference population)

, and relative weights (*i.e.* the contribution of the variable to data variability). Note that the EFD therefore depends on the resistance frequency

in the reference population $\bar{p}$, that we set at the average frequency observed in data. The significance thresholds are 0.1, 0.05, 0.025 and 0.001

denoted by ".", "*", "**" and "***", respectively. Rows represent the different parameters of the models, ordered as follows: growth constant,

regional scale explanatory variables (fungicide use, yield losses and area of wheat under organic farming) and local variation factors (wheat

cultivar and sampling date). "0" means that the parameter was not selected during variable selection. "X" means that the variable was not

considered in the model for the given phenotype. The results for the TriHR phenotypes are not shown as no explanatory variables were retained

by the selection procedure.

| Variable | StrR (2004-2012) | | |
|---|---|---|---|
| | Estimate ± 1 SD | EFD ($\bar{p}$=76%) | Relative weight |
| Growth constant ($\beta$) | -0.17 ± 0.13 | -3.23 | 0.041 |
| Fungicide use ($\nu$) | 1.07 *** ± 0.26 | 6.64 | 0.794 |
| Yield losses ($\rho$) | 0 | | 0 |
| Area under organic farming ($\kappa$) | 0.78 *** ± 0.16 | 3.95 | 0.146 |
| Wheat cultivar ($\sigma_Y$) | 0.31 ** ± 0.09 | ±5.57 | 0.009 |
| Sampling date ($\delta_{S1}$) | -0.35 *** ± 0.1 | -6.87 | 0.01 |

**Table 4** Estimates from the M$_{MoA}$ model for the StrR phenotype.

The three subcolumns show the parameter estimates (*i.e.* posterior mean) and their variability (*i.e.* posterior standard deviation), expected frequency difference (EFD, *i.e.* the change in frequency due to the mean value of the variable on the reference population), and relative weights (*i.e.* the contribution of the variable to data variability). Note that the EFD therefore depends on the resistance frequency in the reference population $\bar{p}$, that we set at the average frequency observed in data. The significance thresholds are 0.1, 0.05, 0.025 and 0.001 denoted by ".", "*", "**" and "***", respectively. Rows represent the different parameters of models, ordered as follows: growth constant, regional scale explanatory variables (fungicide use, yield losses and area of wheat under organic farming) and local variation factors (wheat cultivar and sampling date). "0" means that the parameter was not selected

during variable selection. The results for TriMR and TriHR resistance phenotypes are not

shown as no explanatory variable was retained by the selection procedure.

**Fig. 1** Changes in resistance frequencies (left) and explanatory variables (right, top to

bottom, fungicide use, yield losses and area of wheat under organic farming).

Bold lines: mean regional values. Shaded areas: quantiles of regional values (*i.e.* regional

variability), 25% and 75% (dark grey), 2.5% and 97.5% (light grey). Dotted lines: regional

minimum and maximum values. Fungicide use is expressed in $\frac{ha}{c}$ corresponding to the

mean number of times each mode of action was used for spraying over a cropping season,

regardless of the dose used. Yield losses are expressed in decitons (quintals) per hectare.

**Fig. 2** Maps of observed and predicted frequencies of the StrR resistance phenotype from

2004 to 2011.

Real observations are represented by dots. The colour within the dot indicates the observed

frequency in trials. The background map color shows the regional prediction from the $M_{Al}$

model.

**Fig. 3** Maps of observed and predicted frequencies of TriR6 and TriR7-TriR8 resistance

phenotypes in 2008.

Real observations are represented by dots. The colour within the dot indicates the observed

frequency in trials. The background map colour shows the regional prediction from the $M_{Al}$

model. Top: TriR6. Bottom: TriR7-TriR8.

200 km

Frequency (%)