ORIGINAL MANUSCRIPT

The impact of commercially available ale and lager yeast strains on the

fermentative diversity of beers

Diego Bonatto[a],*

[a]Brewing Yeast Research Group, Centro de Biotecnologia da Universidade Federal do Rio Grande
do Sul, Departamento de Biologia Molecular e Biotecnologia, Universidade Federal do Rio Grande
do Sul, Porto Alegre, RS, Brazil

Short title: Yeast diversity in brewing

***Corresponding author:**

Diego Bonatto

Centro de Biotecnologia da UFRGS - Sala 107

Departamento de Biologia Molecular e Biotecnologia

Universidade Federal do Rio Grande do Sul – UFRGS

Avenida Bento Gonçalves 9500 - Prédio 43421

Caixa Postal 15005

Porto Alegre – Rio Grande do Sul

BRAZIL

91509-900

Phone: (+55 51) 3308-7765

E-mail: diego.bonatto@ufrgs.br

Contract/grant sponsor: CNPq

Abstract

Yeasts from the species *Saccharomyces cerevisiae* (ale yeast) and *Saccharomyces*

pastorianus (lager yeast) are the main component of beer fermentation. It is known that different

beer categories depend on the use of specific ale or lager strains, where the yeast imprint its

distinctive fermentative profile to the beer. Despite this, there are no studies reporting how diverse,

rich, and homogeneous the beer categories are in terms of commercially available brewing yeast

strains. In this work, the diversity, richness, and evenness of different beer categories and

commercial yeast strains available for brewing were evaluated by applying quantitative concepts of

ecology analysis in a sample of 121,528 beer recipes. For this purpose, the frequency of ale or lager

and dry or liquid yeast formulations usage was accessed and its influence in the fermentation

temperature, attenuation profile, and number of recipes for a beer category were analyzed. The

results indicated that many beer categories are preferentially fermented with dry yeast strains

formulations instead of liquid yeasts, despite considering the high number of available liquid yeast

formulations. Moreover, ale dry strains are preferentially used for lager brewing. The preferential

use of specific yeast formulations drives the diversity, richness, and evenness of a beer category,

showing that many yeast strains are potentially and industrially underexplored.

Keywords: Brewing yeasts; Beer categories; Quantitative ecology analysis; Preferential yeast

usage; Fermentation profile; Data mining.

1. Introduction

Beer, a major alcoholic beverage obtained from malt-derived worts, is the product of fermentative metabolism of yeast strains that convert the sugars present in the wort into ethanol and CO_2 (Rai and Jeyaram, 2017). The flavor impact of a specific yeast strain during beer fermentation is also important; in fact, many of the flavors found in a glass of beer are derived from metabolic by-products released by yeast cells during fermentation, like esters, lactones, thiol compounds, higher alcohols, and phenolics (Carrau et al., 2015; Praet et al., 2012; Tran et al., 2015). Additionally, yeasts convert hop and malt-derived glycosylated metabolites to aglycones by the action of β-D-glucosidases during beer fermentation (Gamero et al., 2011); also, yeasts biotransform small molecules found in wort (e.g., amino acids and fatty acids) into flavor components (Carrau et al., 2015). Besides flavor, the visual aspects of a beer category are directly influenced by the yeast strains used for fermentation. For example, the clarity of the beer is a consequence of the flocculation ability of a yeast strain (Vidgren & Londesborough, 2011), while beer foam stability is also dependent on a series of glycoproteins present in the surface of yeast cell wall (Blasco & Viñas, 2011). Thus, the quality of beer is directly dependent on the yeast strain used.

Since the isolation and development of brewing yeast pure cultures from the works of Emil Christian Hansen in the end of 19[th] century (Lodolo et al., 2008; Rank et al., 1988), and the identification of the yeast species that are responsible for bottom (lager) beer fermentation and top (ale) beer fermentation, the brewing industry has benefited from the use of yeast monocultures to give reproducible and consistent products over time. Two major yeast monocultures are employed in breweries nowadays, which are the *Saccharomyces cerevisiae,* mainly responsible for ale fermentation, and *Saccharomyces pastorianus,* a hybrid species responsible for lager fermentation (Lodolo et al., 2008). In this sense, cellular and molecular techniques are allowing researchers to design lager yeast strains for breweries (Mertens et al., 2015) and there is potential for the use of conventional (*S. cerevisiae*) and non-conventional yeast strains (e.g., *Saccharomyces eubayanus*)

isolated from different environments niches for the design of new beers (Cubillos et al., 2019;

Marongiu et al., 2015). Thus, the development of new hybrid strains or the use of environmental

isolated yeast strains allow the brewer to explore different metabolic pathways and aggregate flavor

diversity to beer (Cubillos et al., 2019). However, the applicability of new yeast strains in brewing

industry could be impaired due to the genome and phenotype instabilities induced by the high

selective and specific conditions of beer fermentations (Gorter de Vries et al., 2019), and brewers

preferentially employ commercial yeast strains for beer fermentation due to the high fermentation

efficiency and control (Bellissimi & Ingledew, 2005).

Therefore, considering the commercial available yeast strains for brewing it can be asked

how diverse, rich, and homogeneous beer categories are in terms of different ale and lager yeast

usage found in both dry and liquid formulations. For this purpose, a quantitative ecology analysis of

diversity, richness, and evenness of commercial brewing yeast usage in different beer categories

was performed by considering a sample of 121,528 beer recipes obtained from Brewer's Friend web

site (https://www.brewersfriend.com). In addition, the influence of fermentative parameters (e.g.,

lower and higher recommended fermentation temperature, and attenuation), yeast type (ale or

lager), and formulation (dry or liquid) of commercial yeast strains used in beer categories

fermentation were evaluated. The data gathered showed that beer categories can be classified as

"cold fermented" and "hot fermented" considering the fermentation temperature profile of

commercial yeast strains. Additionally, it was observed that there is a preferential use of dry yeast

strains formulations for beer fermentation instead of liquid strains, even considering the high

number of commercial yeast strains available in liquid formulations. Finally, it was observed that

the preferential use of specific yeast type and/or formulation impacts the diversity, richness, and

evenness of a beer category fermentative profile.

2. Material and methods

2.1. Commercial yeast strain data prospection and analysis

Data regarding yeast strains commercially available for breweries were obtained from

Brewer's Friend (https://www.brewersfriend.com; last access on May, 2020) with the direct consent

of the web page administrator. Initially, the Lynx web browser (https://lynx.browser.org) was used

to map all links associated with commercial yeast data strains, recipes, and different beer categories

from Brewer's Friend. Once obtained, the library rvest (https://github.com/tidyverse/rvest) from R

software (https://www.r-project.org) was used to scrap recipe and commercial yeast data

information for different beer categories from Brewer's Friend links. The raw yeast and recipe data

obtained were filtered and commercial yeast formulations containing the keywords "Wilds &

Sours", "Wine", "S. boulardii", "Mead", "Cider", "Champagne", "Bretts and Blends", "Bacterial

Cultures", "B. bruxellensis", "Sake", "Sour", "Brett", "Bug", "Lactobacillus", "Blend", and

"Saccharomycodes ludwigii" were removed from data. The resulting filtered yeast data containing

information about manufacture company/laboratory, yeast strain brand name, type (ale or lager),

formulation (dry or liquid), alcohol tolerance, flocculation, attenuation percentage, and lower and

higher fermentation temperatures (in °C) were merged with beer category information. Finally, the

definitions of beer categories as well as the country or geographical region from which they

originated were obtained from the 2015 Beer Judge Certification Program (BJCP) Style Guidelines

(https://dev.bjcp.org).

2.2. Statistical and quantitative ecology data analysis and preferential use of yeast strains

The R software (https://www.r-project.org) was used for all statistical and quantitative

ecology data analysis. Data normality for quantitative variables for each beer category was

evaluated by univariate Shapiro-Wilk normality test implemented in rstatix library (https://cran.r-

project.org/web/packages/rstatix/index.html). Correlations between the number of yeast strains,

recipes, lower and higher values of original and final gravity (OG and FG, respectively),

international bitter units (IBUs), and alcohol by volume (ABV) were analyzed with corrplot library

(https://github.com/taiyun/corrplot) by applying Spearman's ρ statistic. All correlations with a *p-*

value < 0.05 were considered statistically significant and were classified as follow: $|r| = 0$, null; $0 < |r| \leq 0.3$, weak; $0.3 < |r| \leq 0.6$, regular; $0.6 < |r| \leq 0.9$, strong; $0.9 < |r| < 1.0$, very strong; $|r| = 1.0$, perfect. The library ggstatsplot (https://cran.r-project.org/web/packages/ggstatsplot/index.html) was used for comparing and plotting the lower and higher fermentation temperature as well as the attenuation percentage for brewing ale and lager yeasts strains in both dry and liquid formulations with the following parameters: display significant pairwise comparisons, Yuen's method for robust estimation and hypothesis testing (Yuen, 1974), display confidence interval ($CI_{95\%}$) and estimated average value (μ), pairwise display all, evaluation of pairwise significance comparison by exact p-value, and false discovery rate adjustment method for p-values.

In order to evaluate the impact of the number of recipes in the lower and higher fermentation temperatures (LF_T and HF_T, respectively) of a beer category, a weighted arithmetic mean value was determined considering the number of recipes for each beer category and the total number of recipes gathered from Brewer's Friend web page using the following equations (1 and 2):

$$\overline{LF_T} = (\Sigma LF_T \times \Sigma R_C)/R$$

$$(1)$$

$$\overline{HF_T} = (\Sigma HF_T \times \Sigma R_C)/R$$

$$(2)$$

where $\overline{LF_T}$ and $\overline{HF_T}$ are the weighted arithmetic mean values for the lower and higher fermentation temperatures for each beer category, ΣLF_T and ΣHF_T represent the sum of lower and higher fermentation temperatures, respectively, for a given beer category, ΣR_C is the sum of the number of beer recipes for a given beer category, and R represents the total number of recipes available in Brewer's Friend web page as obtained in May, 2020. Beer categories that display $\overline{HF_T}$ values above the average were classified as "hot fermented" beers, while beer categories with $\overline{HF_T}$ values below the average were classified as "cold fermented" beers. A linear regression analysis was performed

147 in order to determine the correlation of $\overline{\mathrm{LF}}_T$ and $\overline{\mathrm{HF}}_T$ in different beer categories with the library

148 ggpmisc (https://cran.r-project.org/web/packages/ggpmisc/index.html).

149   The preferential use of a specific brewing yeast strain (ale dry or liquid, and lager dry or

150 liquid) in comparison to all different yeast strains reported for a beer category (P_Y) was calculated as

151 follow (equation 3):

152
$$P_Y = \frac{\sum R_{YTF}}{\sum R_C} \times \frac{\sum Y_C}{\sum Y_{TF}}$$

153   (3)

where ΣR_{YTF} is the total number of beer category-associated recipes that use a specific brewing yeast

strain (ale dry or liquid, and lager dry or liquid), ΣR_c and ΣY_c are the total number of recipes and

yeast strains for a given beer category, respectively, and ΣY_{TF} is the total number of beer category-

associated specific brewing yeast strain (ale dry or liquid, and lager dry or liquid).

Quantitative ecology data analysis was performed in R environment with the vegan library

(Dixon, 2003). In this sense, the frequency of a unique yeast strain in a beer category was used to

estimate the parameters of richness, diversity, and evenness. For richness estimation, the Menhinick

index (Mi) (Cazzolla Gatti et al., 2020) was applied with the equation (4):

162
$$Mi = \frac{n}{\sqrt{N}}$$

163   (4)

164 where n is the frequency of unique yeast strains for a given beer category and N is the number of

165 recipes for a beer category. By its turn, the Simpson's diversity (D^S) (Thukral, 2017) of brewing

166 yeast strains in different beer categories was determined by using the Simpson's index (λ) described

167 in equations 5 and 6:

168
$$\lambda = \sum \frac{n_i(n_i - 1)}{N(N-1)}$$

169   (5)

$$D^S = 1 - \lambda$$

(6)

where n_i is the frequency of each i brewing yeast strain in a given beer category and N is the number

of recipes for a beer category. Finally, the evenness of a specific brewing yeast strain among

different beer categories was determined by the Pielou index (J) (Thukral, 2017) as follow

(equation 7):

$$J = \frac{\lambda}{\ln(S)}$$

(7)

where λ is the diversity Simpson's index as described in equation 5 and S indicates the total number

of brewing strains for each beer category.

3. Results and Discussion

3.1. Beer categories and commercial brewing yeast strains data analysis

The craft beer revolution is a well characterized movement inside beer industry that can be

roughly defined as the origin, development, and spread of local microbreweries as the consequence

of the large-scale, homogeneous mildly beer brands that dominated the beer market in the late 20[th]

century followed by the increasing demand of new beer styles (Garavaglia & Swinnen, 2017). In

addition, the craft beer industry can also be defined by consumers that drink less beer but are

willing to pay more for special and pricier beers with different textures and flavors (Donadini &

Porretta, 2017). Thus, it becomes clear that the major force that drives the craft beer revolution is

the development of beer categories with a high diversity in the use of ingredients, where beers

produced with local raw materials and yeast characterize the so called "beer du terroir" (Budroni et

al., 2017). As pointed by Budroni et al. (2017), the use of local yeast strains or even the

development of tailor-made yeast strains by different cellular and molecular techniques (Cubillos et

al., 2019; Gibson et al., 2017) is a relatively unexplored tool for the diversification of local beers.

However, and despite the academic or industrial initiatives to promote the use of local ingredients,

yeast manufacturers still have a major role in providing the main yeast strains used in breweries and

then directly impacting the beer quality that is consumed. In order to understand the roles and the

influence of commercial yeast strains in the fermentative aspects of different beer categories, the

Brewer's Friend, a large repository of beer recipes and yeast strain data was chosen to evaluate the

specific parameters related to yeast strain richness and diversity as well as the preference of

producers in the use of specific yeast strains for fermentation. It should be noted that beer categories

that use lactic acid bacteria and non-conventional yeast genera and species (e.g., *Brettanomyces*

bruxellensis) were excluded from this work. Thus, a total of 121,528 beer recipes divided into 34

major beer categories were downloaded from Brewer's Friend web page. In addition, 476

commercially available yeast strains were analyzed in terms of type (ale or lager), formulation (dry

or liquid), minimum and maximum fermentation temperature, and attenuation.

The data collected from Brewer's Friend website showed that 14 beer categories have a high

number of yeast strains in comparison to the average number of yeast strains employed for brewing

(μ = 144.44 unique yeast strains $\times$ beer category^{-1}, Figure 1A) and include relevant specialty craft

beers, like India Pale Ale (IPA), Standard American Beer, Pale American Beer, and Belgian and

Strong Belgian Ales (Figure 1A) (Haugland, 2014; Poelmans & Swinnen, 2018).

Additionally, eight beer categories with a high number of yeast strains (IPA, Standard American

Beer, Pale American Beer, Belgian and Strong Belgian Ales, American Porter and Stout, Dark

British Beer, Amber and Brown American Beer) also display a high number of recipes compared to

the average number of recipes by beer category (μ = 3574.35 beer recipes $\times$ beer category^{-1}, Figure

1B). By its turn, the number of yeast strains and recipes used for beer categories related to lager

family or specialty beers is low (Figures 1A and B). This initial data analysis prompted to question

if the different beer categories parameters (e.g., IBU, OG, FG, and ABV) correlate with the number

of yeast strains and recipes (Figure 1C). In fact, the number of yeast strains and recipes observed for

a specific beer category did not show any correlation with OG, FG, and ABV level; however, it was

observed a significant correlation of IBU level with the number of yeast strains and recipes (Figure

1C). This correlation could be partially explained by the increasing consumer preference for hoppier

beer as well as the development of new hop cultivars that aggregate different flavors to the beer

(Gabrielyan et al., 2014; Madsen et al., 2020), and thus directing the brewer's preference for the

design of beer recipes that made use of high amount of hops for bitterness or flavor. Moreover, a

significant correlation of IBU, OG, FG, and ABV was also observed (Figure 1C).

Considering the total number of yeast strains evaluated in each beer category (Figure 1A), it

was asked how many distinct yeast ale and lager strains in dry or liquid formulations were

employed by the brewers in different beer categories (Figures 2A and B). From the total number of

yeast strains annotated, it was observed that the number of liquid yeast strains counted for each beer

category was higher than the number of dry yeast formulations (Figure 2A). Additionally, the

number of distinct yeast ale strains determined for each beer category was higher than the number

of lager strains (Figure 2B). These data could be supported by the fact that the number of

commercially available yeast liquid formulations is higher than dry yeasts as observed from

Brewer's Friend website data (397 liquid versus 79 dry yeast strains) and the number of ale strains

commercially available is also higher than lager strains (390 ale versus 86 lager yeast strains). An

explanation about why there are many more liquid strains in comparison to dry strains (and the

same for ale versus lager strains) was not completely addressed until now, but it can be

hypothesized that many brewing yeast strains have a low tolerance to the industrial drying process,

despite the fact that different methods to dehydrate yeast cells have been developed since the 18[th]

century (Gélinas, 2019). Supporting this hypothesis, it has been reported that lager yeasts strains

have different desiccation tolerances (Layfield et al., 2011).

Thus, how similar are the commercial available brewers' yeast strains in terms of

fermentation temperature and attenuation? Considering the lower fermentation temperature reported

by the yeast manufacturers for the 390 unique ale strains (68 dry and 322 liquid yeast formulations)

used in brewing, it was observed that dry yeast strains have a significant lower mean fermentation temperature (μ = 16.42 °C; Figure 3A) in comparison to liquid yeast formulations (μ = 18.52 °C; Figure 3A). On the other hand, no significant difference was observed in the higher fermentation temperature reported by yeast manufacturers for ale dry (μ = 24.39 °C) and liquid (μ = 24.33 °C) formulations (Figure 3B). By its turn, from the 86 commercially available lager yeast strains (11 dry and 75 liquid formulations), it was not observed any significant difference in the mean lower fermentation temperature for lager dry (μ = 10.51 °C) and liquid (μ = 10.10 °C) strains (Figure 3C), while a significant difference was observed in the mean higher fermentation temperature for lager dry (μ = 17.85 °C) and liquid (μ = 14.52 °C) formulations (Figure 3D). Data regarding attenuation showed that ale yeast strains are similar in both dry (μ = 76.95%) and liquid (μ = 75.85%) forms (Figure 4A), while dry lager strains are significantly more attenuative (μ = 79.18%) than liquid strains (μ = 74.00%) (Figure 4B). To date, this is the first study that compared the attenuation and the higher and lower fermentation temperature of commercially available yeast ale and lager strains in dry and liquid formulations.

Attenuation and fermentation temperature are the two main variables that significantly impact the beer, where the efficient use of malt-derived sugars by yeast strains (resulting in high ethanol yields) and the absence of off-flavors is desirable for any beer category (Powell et al., 2003). In this sense, it becomes clear from the data collected for this study that ale and lager yeasts strains in dry and liquid formulations are phenotypically similar considering the 95% confidence intervals ($CI_{95\%}$) for temperature (Figures 3A to D) and attenuation (Figures 4A and B). However, some outliers could be observed in ale strains with high fermentation temperature profiles (Figures 3A and B) which correspond to norwegian kveik and belgian hybrid saison strains (González et al., 2008; Preiss et al., 2018) as well as Kölsch/Altbier-associated yeast strains. By its turn, high fermentation temperature profile in lager yeast was observed for strains employed in the California Common beer style (Figures 3C to D). Regarding attenuation, the outliers found in ale strain data

correspond to different belgian yeast strains that express the *STA1* gene (Krogerus & Gibson, 2020),

leading to beer overattenuation (Figure 4A).

Fermentation temperature control is critical for many beer categories, as the yeast

performance and the development of specific flavors are directly linked to fermentation

temperature, especially modulating the production of esters and higher alcohol (Olaniran et al.,

2011; Pires et al., 2014). Considering the data heterogeneity of yeast strains and recipes by each

beer category analyzed (Figures 1A and B), what is the impact of brewer's preference on lower and

higher fermentation temperature for a given beer category? To answer this question, a weighted

arithmetic mean value for the lower and higher fermentation temperatures ($\overline{\overline{LF}}_T$ and $\overline{\overline{HF}}_T$,

respectively) for each beer category was determined (Figure 5). Interestingly, two major groups of

beer could be discriminated by considering the mean value of $\overline{\overline{HF}}_T$ ($\mu = 22.56\ °C$), which were

defined as "cold fermented beer" ($\overline{\overline{HF}}_T < 22.56\ °C$) and "hot fermented beer" ($\overline{\overline{HF}}_T > 22.56\ °C$)

(Figure 5). The cold fermented beers correspond to all lager family-associated beer categories,

while the hot fermented beer group contains all ale family-associated categories (Figure 5). A strong

and positive correlation could be observed between $\overline{\overline{LF}}_T$ and $\overline{\overline{HF}}_T$, where the Czech Lager and

Strong Belgian Ale categories correspond to the extremes of $\overline{\overline{LF}}_T$ and $\overline{\overline{HF}}_T$ values (Figure 5).

3.2. The brewers' preferences for yeast strain usage

The use of dry yeast is gathering popularity over yeast liquid formulations for brewing due

to the fact that dry formulations occupy smaller volume and do not need refrigeration in comparison

to liquid yeasts, resulting in lower costs associated with logistic and yeast storage. Moreover, dry

yeast formulations can be kept for many years without loss of vitality (Rapoport, 2017). Thus, there

is a natural tendency of brewers to employ dry yeasts in beer fermentation, despite the low number

of ale and lager dry strains commercially available (68 and 11 strains, respectively). Interestingly,

the preferential use of dry yeast rather than of liquid yeast for beer fermentation could be clearly

observed from Brewer's Friend data (Figure 6). The preferential yeast usage or P_Y was higher in all

beer categories where dry ale and lager yeast strains were employed, while liquid ale and lager

formulations were less preferred (Figure 6). Some hot fermented beer categories, like Standard

American Beer, American Porter and Stout, Pale American Ale, Pale Commonwealth Beer, and IPA

have high P_Y values for dry ale yeast strains (Figure 6). High P_Y values for dry lager strains were

also observed for all cold fermented beer categories, with the exception of Pale Bitter European

Beer category, which has a preferential use for liquid ale strains (Figure 6). Considering the

brewer's preferential yeast usage (Table 1), how this variable impacts the evenness, richness, and

diversity of beer categories?

3.3. Measuring the evenness, richness, and diversity distribution of commercial yeast strains in beer

categories

To evaluate the impact of brewer's preferential yeast usage in beer categories, a quantitative

ecology analysis was performed. This analysis consider the concepts of evenness, richness, and

diversity that are employed in different fields (Xu et al., 2020). For this work, the diversity concept

is a variable that depends on the richness of different yeast strains found within a beer category,

how evenness (homogeneous) are those strains distributed among beer recipes found in a category

as well as the number of beer recipes found in a given category. Thus, a beer category with a high

diversity has an elevated number of different yeast strains with an evenness distribution of those

strains among a high number of beer recipes found within the beer category.

Initially, beer category diversity and richness were evaluated by using the Simpson's (λ) and

Menhinick (Mi) indexes, respectively, for cold and hot fermented beer (Figure 7A). By using the

mean values of λ ($\mu_{\lambda Cold} = 0.93$ and $\mu_{\lambda Hot} = 0.91$) and *Mi* ($\mu_{MiCold} = 3.35$ and $\mu_{MiHot} = 3.26$) (Figure 7A)

it was possible to classify beer categories into four major groups: (i) beer categories that have a high

richness and diversity, (ii) beer categories with low richness and high diversity, (iii) beer categories

with high richness and low diversity, and (iv) beer categories with low richness and low diversity

(Figure 7A). A similar analysis was made considering λ diversity and Pielou evenness index (J),

where the mean values for λ ($\mu_{\lambda Cold}$ = 0.93 and $\mu_{\lambda Hot}$ = 0.91; Figure 7A) and J ($\mu_{J Cold}$ = 0.19 and $\mu_{J Hot}$ =

0.18; Figure 7B) allow to group beer categories into four types: (i) high richness and evenness, (ii)

low richness and high evenness, (iii) high richness and low evenness, and (iv) low richness and low

evenness (Figure 7B).

Considering cold fermented beer group, it was observed that Pale Bitter European Beer,

Czech Lager, Pale Malty European Lager, International Lager, and Dark European Lager have low

yeast strain diversity and richness (Figure 7A and Table 1), meaning that brewers preferentially use

a small number of yeast strains, especially dry lager yeasts, to ferment beers that fall within these

categories (Figure 6). Moreover, the evenness of yeast strains usage for Pale Bitter European Beer

and International Lager is low (Figure 7B and Table 1), also pointing to a preferential use of yeasts

type and formulation as seen in the previous analysis (Figure 6). On the other hand, Amber Malty

European Lager and Amber Bitter European Lager have a high diversity and evenness, but a low

richness (Figures 7A and B; Table 1), which can be explained by the extensive use of dry lager

strains (Figure 6).

Noteworthy, from 25 hot fermented beer categories analyzed, ten categories display low

values of richness, evenness, and diversity, like Pale American Ale, IPA, Strong American Ale,

Amber and Brown American Ale, among others (Figures 7A and B; Table 1). This result indicates

that brewers preferentially use a very low number of yeast strains to ferment beers that fall within

these categories, corroborating the P_Y data that favor the use of dry ale formulations for these

categories (Figure 6). Interestingly, Belgian Ale, Strong Belgian Ale, Trappist Ale, and Brown

British Beer have a low richness and high diversity, pointing to the fact that the number of specific

strains used in these categories is not high despite being evenly distributed (Figures 7A and B; Table

1). Additionally, the number of recipes described for these categories is also high (Figure 1A),

which contributes to the diversity values observed for Belgian Ale, Strong Belgian Ale, Trappist

Ale, American Porter and Stout, and Brown British Beer categories.

Specialty beers like Fruit Beer, European Sour Ale, Spiced or Wood Beers, and Strong

British Ale have a high diversity and richness (Figure 7A and Table 1), indicating that brewers are

prone to use a high diverse set of yeast strains to ferment beer that fall within these categories.

However, the evenness of yeast strain usage among these beer categories can be variable, where

Fruit Beer, European Sour Ale, Strong British Ale, and Spiced beers display low evenness, while

Wood Beer has a high evenness value (Figure 7B; Table 1).

4. Conclusion

The data gathered in this work showed that brewers have a preference for a small set of

yeast strains, indicating that there is an underexplored potential for developing new beers by using

the commercial yeast strains that are already available and have low usage. Despite the efforts of

researchers to develop new yeast strains (Cubillos et al., 2019; Hittinger et al., 2018; Mertens et al.,

2015; Saerens et al., 2010; Steensels et al., 2014), there is an ingrained brewing culture for using

conventional yeast strains to ferment beer, especially dry ale formulations. As pointed before, dry

yeast formulations have a series of advantages when compared to liquid yeast strains (Bellissimi &

Ingledew, 2005), but the low number of dry yeast strains is a major disadvantage that brewers

should consider on the development of new products. On the other hand, the low number of

available dry yeast strains also indicates a potential and unexplored industrial field for the

development of new dry yeast strains. For example, the design of strains with high biotransforming

activity of hop-derived compounds is a major trend in brewing (Praet et al., 2012; Steyer et al.,

2017; Tran et al., 2015) and can aggregate value to beer (Gabrielyan et al., 2014). Additionally,

yeasts with increased resistance to osmotic pressure and high attenuation are gathering attention

from brewers to develop new beers (Krogerus & Gibson, 2020).

In conclusion, yeasts are an underexplored resource in the brewing industry, with a large

space for designing and repurposing commercially available yeast strains for the creation of new

beers.

Funding: This work was supported by the Conselho Nacional de Desenvolvimento

Científico e Tecnológico – CNPq [grant number 302969/2016–0]. The sponsor had no role in the

study design; collection, analysis, and interpretation of data; writing of the report; or decision to

submit the article for publication.

Declarations of interest: None

Human and animal rights: The experiments included in this manuscript did not involve

any animal or human participants.

References

Bellissimi, E., & Ingledew, W. M. (2005). Analysis of Commercially Available Active Dry Yeast
Used for Industrial Fuel Ethanol Production. *Journal of the American Society of Brewing
Chemists*, *63*(3), 107–112. https://doi.org/10.1094/ASBCJ-63-0107

Blasco, lucía, & Viñas, M. (2011). Proteins influencing foam formation in wine and beer: the role
of yeast. *International Microbiology*, *14*, 61–76. https://doi.org/10.2436/20.1501.01.136

Budroni, M., Zara, G., Ciani, M., & Comitini, F. (2017). Saccharomyces and Non-Saccharomyces
Starter Yeasts. *Brewing Technology*. https://doi.org/10.5772/intechopen.68792

Carrau, F., Gaggero, C., & Aguilar, P. S. (2015). Yeast diversity and native vigor for flavor
phenotypes. *Trends in Biotechnology*, *33*(3), 148–154.
https://doi.org/10.1016/j.tibtech.2014.12.009

Cazzolla Gatti, R., Amoroso, N., & Monaco, A. (2020). Estimating and comparing biodiversity with
a single universal metric. *Ecological Modelling*, *424*, 109020.
https://doi.org/10.1016/j.ecolmodel.2020.109020

Cubillos, F. A., Gibson, B., Grijalva-Vallejos, N., Krogerus, K., & Nikulin, J. (2019).
Bioprospecting for brewers: Exploiting natural diversity for naturally diverse beers. *Yeast*,
36(6), 383–398. https://doi.org/10.1002/yea.3380

Dixon, P. (2003). VEGAN, a package of R functions for community ecology. *Journal of Vegetation
Science*, *14*(6), 927–930. https://doi.org/10.1111/j.1654-1103.2003.tb02228.x

Donadini, G., & Porretta, S. (2017). Uncovering patterns of consumers' interest for beer: A case
study with craft beers. *Food Research International*, *91*, 183–198.
https://doi.org/10.1016/j.foodres.2016.11.043

Gabrielyan, G., McCluskey, J. J., Marsh, T. L., & Ross, C. F. (2014). Willingness to Pay for Sensory
Attributes in Beer. *Agricultural and Resource Economics Review*, *43*(1), 125–139.
https://doi.org/10.1017/S1068280500006948

Gamero, A., Manzanares, P., Querol, A., & Belloch, C. (2011). Monoterpene alcohols release and
bioconversion by Saccharomyces species and hybrids. *International Journal of Food
Microbiology*, *145*(1), 92–97. https://doi.org/10.1016/j.ijfoodmicro.2010.11.034

Garavaglia, C., & Swinnen, J. (2017). The craft beer revolution: An international perspective. *Choices, 32*(3), 1–8.

Gélinas, P. (2019). Active Dry Yeast: Lessons from Patents and Science. *Comprehensive Reviews in Food Science and Food Safety, 18*(4), 1227–1255. https://doi.org/10.1111/1541-4337.12445

Gibson, B., Geertman, J.-M. A., Hittinger, C. T., Krogerus, K., Libkind, D., Louis, E. J., Magalhães, F., & Sampaio, J. P. (2017). New yeasts—new brews: modern approaches to brewing yeast design and development. *FEMS Yeast Research, 17*(4). https://doi.org/10.1093/femsyr/fox038

González, S. S., Barrio, E., & Querol, A. (2008). Molecular Characterization of New Natural Hybrids of Saccharomyces cerevisiae and S. kudriavzevii in Brewing. *Applied and Environmental Microbiology, 74*(8), 2314–2320. https://doi.org/10.1128/AEM.01867-07

Gorter de Vries, A. R., Voskamp, M. A., van Aalst, A. C. A., Kristensen, L. H., Jansen, L., van den Broek, M., Salazar, A. N., Brouwers, N., Abeel, T., Pronk, J. T., & Daran, J.-M. G. (2019). Laboratory Evolution of a Saccharomyces cerevisiae × S. eubayanus Hybrid Under Simulated Lager-Brewing Conditions. *Frontiers in Genetics, 10,* 1:18. https://doi.org/10.3389/fgene.2019.00242

Haugland, J. E. (2014). The origins and diaspora of the India Pale Ale. In *The Geography of Beer* (pp. 119–129). Springer.

Hittinger, C. T., Steele, J. L., & Ryder, D. S. (2018). Diverse yeasts for diverse fermented beverages and foods. *Current Opinion in Biotechnology, 49,* 199–206. https://doi.org/10.1016/j.copbio.2017.10.004

Krogerus, K., & Gibson, B. (2020). A re-evaluation of diastatic Saccharomyces cerevisiae strains and their role in brewing. *Applied Microbiology and Biotechnology, 104*(9), 3745–3756. https://doi.org/10.1007/s00253-020-10531-0

Layfield, J. B., Phister, T. G., & Sheppard, J. D. (2011). Characterization of Hybrid Strains of *Saccharomyces Pastorianus* for Desiccation Tolerance. *Journal of the American Society of Brewing Chemists, 69*(2), 108–115. https://doi.org/10.1094/ASBCJ-2011-0301-01

Lodolo, E. J., Kock, J. L. F., Axcell, B. C., & Brooks, M. (2008). The yeast Saccharomyces cerevisiae– the main character in beer brewing. *FEMS Yeast Research*, *8*(7), 1018–1036. https://doi.org/10.1111/j.1567-1364.2008.00433.x

Madsen, E. S., Gammelgaard, J., & Hobdari, B. (2020). *New Developments in the Brewing Industry: The Role of Institutions and Ownership*. Oxford University Press.

Marongiu, A., Zara, G., Legras, J.-L., Del Caro, A., Mascia, I., Fadda, C., & Budroni, M. (2015). Novel starters for old processes: use of Saccharomyces cerevisiae strains isolated from artisanal sourdough for craft beer production at a brewery scale. *Journal of Industrial Microbiology & Biotechnology*, *42*(1), 85–92. https://doi.org/10.1007/s10295-014-1525-1

Mertens, S., Steensels, J., Saels, V., Rouck, G. D., Aerts, G., & Verstrepen, K. J. (2015). A Large Set of Newly Created Interspecific Saccharomyces Hybrids Increases Aromatic Diversity in Lager Beers. *Applied and Environmental Microbiology*, *81*(23), 8202–8214. https://doi.org/10.1128/AEM.02464-15

Olaniran, A. O., Maharaj, Y. R., & Pillay, B. (2011). Effects of fermentation temperature on the composition of beer volatile compounds, organoleptic quality and spent yeast density. *Electronic Journal of Biotechnology*, *14*(2), 5–5.

Pires, E. J., Teixeira, J. A., Brányik, T., & Vicente, A. A. (2014). Yeast: the soul of beer's aroma—a review of flavour-active esters and higher alcohols produced by the brewing yeast. *Applied Microbiology and Biotechnology*, *98*(5), 1937–1949. https://doi.org/10.1007/s00253-013-5470-0

Poelmans, E., & Swinnen, J. (2018). Belgium: Craft Beer Nation? In C. Garavaglia & J. Swinnen (Eds.), *Economic Perspectives on Craft Beer* (pp. 137–160). Springer International Publishing. https://doi.org/10.1007/978-3-319-58235-1_5

Powell, C., Quain, D., & Smart, K. (2003). The impact of brewing yeast cell age on fermentation performance, attenuation and flocculation. *FEMS Yeast Research*, *3*(2), 149–157. https://doi.org/10.1016/S1567-1356(03)00002-3

Praet, T., Van Opstaele, F., Jaskula-Goiris, B., Aerts, G., & De Cooman, L. (2012). Biotransformations of hop-derived aroma compounds by Saccharomyces cerevisiae upon fermentation. *Cerevisia*, *36*(4), 125–132. https://doi.org/10.1016/j.cervis.2011.12.005

Preiss, R., Tyrawa, C., Krogerus, K., Garshol, L. M., & van der Merwe, G. (2018). Traditional Norwegian Kveik Are a Genetically Distinct Group of Domesticated *Saccharomyces cerevisiae* Brewing Yeasts. *Frontiers in Microbiology, 9.* https://doi.org/10.3389/fmicb.2018.02137

Rai, A. K., & Jeyaram, K. (2017). Role of Yeasts in Food Fermentation. In T. Satyanarayana & G. Kunze (Eds.), *Yeast Diversity in Human Welfare* (pp. 83–113). Springer. https://doi.org/10.1007/978-981-10-2621-8_4

Rank, G. H., Casey, G., & Xiao, W. (1988). Gene transfer in industrial *Saccharomyces* yeasts. *Food Biotechnology, 2*(1), 1–41. https://doi.org/10.1080/08905438809549674

Rapoport, A. (2017). Anhydrobiosis and Dehydration of Yeasts. In A. A. Sibirny (Ed.), *Biotechnology of Yeasts and Filamentous Fungi* (pp. 87–116). Springer International Publishing. https://doi.org/10.1007/978-3-319-58829-2_4

Saerens, S. M. G., Duong, C. T., & Nevoigt, E. (2010). Genetic improvement of brewer's yeast: current state, perspectives and limits. *Applied Microbiology and Biotechnology, 86*(5), 1195–1212. https://doi.org/10.1007/s00253-010-2486-6

Steensels, J., Snoek, T., Meersman, E., Nicolino, M. P., Voordeckers, K., & Verstrepen, K. J. (2014). Improving industrial yeast strains: exploiting natural and artificial diversity. *FEMS Microbiology Reviews, 38*(5), 947–995. https://doi.org/10.1111/1574-6976.12073

Steyer, D., Tristram, P., Clayeux, C., Heitz, F., & Laugel, B. (2017). Yeast Strains and Hop Varieties Synergy on Beer Volatile Compounds. *BrewingScience, Volume 70,* 131–141. https://doi.org/10.23763/BrSc17-13Steyer

Thukral, A. K. (2017). A review on measurement of Alpha diversity in biology. *Agricultural Research Journal, 54*(1), 1. https://doi.org/10.5958/2395-146X.2017.00001.1

Tran, T. T. H., Cibaka, M.-L. K., & Collin, S. (2015). Polyfunctional Thiols in Fresh and Aged Belgian Special Beers: Fate of Hop S-Cysteine Conjugates. *Journal of the American Society of Brewing Chemists, 73*(1), 61–70. https://doi.org/10.1094/ASBCJ-2015-0130-01

Vidgren, V., & Londesborough, J. (2011). 125th Anniversary Review: Yeast Flocculation and Sedimentation in Brewing. *Journal of the Institute of Brewing, 117*(4), 475–487. https://doi.org/10.1002/j.2050-0416.2011.tb00495.x

Xu, S., Böttcher, L., & Chou, T. (2020). Diversity in biology: definitions, quantification and models. *Physical Biology, 17*(3), 031001. https://doi.org/10.1088/1478-3975/ab6754

Yuen, K. K. (1974). The two-sample trimmed t for unequal population variances. *Biometrika, 61*(1), 165–170. https://doi.org/10.1093/biomet/61.1.165

 Tables

 Table 1. Classification of major beer categories into fermentation types and analysis of evenness,

 richness, diversity, and preferential yeast usage.

Category	Fermentation type	Evenness (Pielou index)	Richness (Menhinick index)	Diversity (Simpson index)	Preferential yeast usage (*Py*)
Smoked Beer	Cold fermented	High	High	High	Ale Dry, Lager Dry
Amber Bitter European Lager	Cold fermented	High	Low	High	Lager Dry
Amber Malty European Lager	Cold fermented	High	Low	High	Lager Dry
Strong European Beer	Cold fermented	Low	High	High	Ale Dry, Lager Dry
Dark European Lager	Cold fermented	High	Low	Low	Lager Dry, Lager Liquid
Czech Lager	Cold fermented	High	Low	Low	Lager Dry
Pale Malty European Lager	Cold fermented	High	Low	Low	Lager Dry
International Lager	Cold fermented	Low	Low	Low	Lager Dry
Pale Bitter European Beer	Cold fermented	Low	Low	Low	Ale Dry, Ale Liquid, Lager Dry
American Wild Ale	Hot fermented	High	High	High	Ale Dry
Historical Beer	Hot fermented	High	High	High	Ale Dry
Strong British Ale	Hot fermented	High	High	High	Ale Dry
Wood Beer	Hot fermented	High	High	High	Ale Dry
Brown British Beer	Hot fermented	High	Low	High	Ale Dry
Trappist Ale	Hot fermented	High	Low	High	Ale Liquid
European Sour Ale	Hot fermented	Low	High	High	Ale Dry
Fruit Beer	Hot fermented	Low	High	High	Ale Dry
Specialty Beer	Hot fermented	Low	High	High	Ale Dry
Spiced Beer	Hot fermented	Low	High	High	Ale Dry
Belgian Ale	Hot fermented	Low	Low	High	Ale Dry
Strong Belgian Ale	Hot fermented	Low	Low	High	Ale Dry
American Porter and Stout	Hot fermented	Low	Low	High	Ale Dry
Alternative Fermentables Beer	Hot fermented	High	High	Low	Ale Dry
Scottish Ale	Hot fermented	High	Low	Low	Ale Dry, Ale Liquid
British Bitter	Hot fermented	High	Low	Low	Ale Dry
Standard American Beer	Hot fermented	Low	High	Low	Ale Dry, Lager Dry
Amber and Brown American Beer	Hot fermented	Low	Low	Low	Ale Dry
Dark British Beer	Hot fermented	Low	Low	Low	Ale Dry
German Wheat Beer	Hot fermented	Low	Low	Low	Ale Dry, Ale Liquid
IPA	Hot fermented	Low	Low	Low	Ale Dry
Irish Beer	Hot fermented	Low	Low	Low	Ale Dry
Pale American Ale	Hot fermented	Low	Low	Low	Ale Dry
Pale Commonwealth Beer	Hot fermented	Low	Low	Low	Ale Dry
Strong American Ale	Hot fermented	Low	Low	Low	Ale Dry

Figure legends

Figure 1. Determination of the number of unique yeast strains (A) and recipes (B) by each beer category. The dashed line in the graphics (A) and (B) indicates the mean value of the number of yeast strains and recipes, respectively. An amplified view of specific beer categories is indicated by the inset and dotted lines in graphic (B). In (C), correlation analysis of the number of yeast strains and recipes with major parameters associated to beer categories, like the lower and higher values of international bitter units (IBU_lower and IBU_higher, respectively), final gravity (FG_lower and FG_higher), original gravity (OG_lower and OG_higher), and alcohol by volume (ABV_lower and ABV_higher). The color scale in (C) indicates the pattern of correlation (negative or positive), from -1 (red) to 1 (blue).

Figure 2. Number of dry or liquid (A) and ale or lager (B) yeast strains observed in each beer category. The total number of yeast strains in each beer category is indicated by the dark square in each column.

Figure 3. Evaluation of lower and higher fermentation temperatures (°C) for different ale (A, B) and lager (C, D) yeast strains (type) commercially available for brewers. The number of yeast strains observed for each formulation (n) as well as the statistical data analysis are indicated in the graphics.

Figure 4. Evaluation of attenuation percentage for different ale (A) and lager (B) yeast strains (type) commercially available for brewers. The number of yeast strains observed for each formulation (n) as well as the statistical data analysis are indicated in the graphics.

Figure 5. Linear regression of weighted lower and higher fermentation temperature (ºC) observed for each beer category. The dashed line indicates the average value for weighted higher fermentation temperature. The coefficient of determination (R^2) and the equation of linear regression are indicated in the figure. The dotted line and the respective the gray area indicates the regression line and confidence intervals, respectively.

Figure 6. Preferential yeast usage (Py) analysis of ale and lager yeast strains in dry or liquid

formulations for each beer category. Red bars indicate beer categories that are cold fermented, while

blue bars indicate beer categories that are hot fermented.

Figure 7. Analysis of brewing yeast strain richness and diversity (A), and evenness and

diversity (B) for each beer category. Dashed lines indicate the average values of richness, evenness,

and diversity. Abbreviations: High Richness-High Diversity$_{Simpson}$ (HRHDSMP), High Richness-

Low Diversity$_{Simpson}$ (HRLDSMP), Low Richness-High Diversity$_{Simpson}$ (LRHDSMP), Low

Richness-Low Diversity$_{Simpson}$ (LRLDSMP), High Evenness-High Diversity$_{Simpson}$ (HEHDSMP),

High Evenness-Low Diversity$_{Simpson}$ (HELDSMP), Low Evenness-High Diversity$_{Simpson}$

(LEHDSMP), Low Evenness-Low Diversity$_{Simpson}$ (LELDSMP).

Figure 1.

Figure 2.

Figure 3.

424   **Figure 4.**

Type: Ale

$t_{\text{Yuen}}(65.79) = 1.13$, $p = 0.262$, $\hat{\xi} = 0.13$, $\text{CI}_{95\%}$ [0.00, 0.36], $n_{\text{obs}} = 390$

Type: Lager

$t_{\text{Yuen}}(10.49) = 5.91$, $p = <0.001$, $\hat{\xi} = 0.90$, $\text{CI}_{95\%}$ [0.61, 0.99], $n_{\text{obs}} = 86$

427   **Figure 5.**

430   **Figure 6.**

Figure 7.