# Neuroanatomical dissociations of syntax and semantics revealed by lesion-symptom mapping

Short title: neuroanatomical dissociations of syntax and semantics

William Matchin[1]*, Alexandra Basilakos[1], Dirk-Bart den Ouden[1], Brielle C. Stark[2], Gregory Hickok[3], & Julius Fridriksson[1]

[1]Dept. of Communication sciences and Disorders, University of South Carolina
[2]Dept. of Speech and Hearing Sciences, Program in Neuroscience, Indiana University Bloomington
[3]Dept. of Cognitive Sciences, Dept. of Language Science, University of California, Irvine

*Corresponding author

915 Greene St., Discovery 1, Room 202D, Columbia, SC 29208, USA
Email: matchin@mailbox.sc.edu
Tel: +1 803-777-9240

## Abstract

In the early and mid 1800s, scientists debated whether the human brain was functionally differentiated with respect to cognition. The issue was largely resolved when specific language impairments were identified following focal patterns of brain damage. However, neuroimaging has revived this discussion, as many studies find similar syntactic and semantic effects across the set of brain regions implicated in language. Here we address this modern debate using lesion-symptom mapping in two large, partially-overlapping groups of people with left hemisphere brain damage due to stroke (N=121, N=92). We identified multiple measure by region interaction effects, associating damage to the posterior middle temporal gyrus with syntactic comprehension deficits, damage to posterior inferior frontal gyrus with expressive agrammatism, and damage to inferior angular gyrus with semantic category word fluency deficits. Our results are inconsistent with recent hypotheses that regions of the language network play similar roles in high-level linguistic processing.

## Introduction

At its core, language consists of pairings of form and meaning. Linguistic form at the sentence level consists of syntactic structure: sequences of words and their hierarchical arrangement. Linguistic meaning consists of concepts denoted by words and the interpretation of their combination. These domains are further subdivided into distinct types, such as the distinction between hierarchical and linear syntax (Berwick & Chomsky, 2016; Matchin & Hickok, 2020)

and between taxonomic and thematic features (Binder & Desai, 2011; Estes et al., 2011; Lin & Murphy, 2001; Schwartz et al., 2011). Despite this differentiation, neuroimaging experiments typically find that contrasts of syntax and semantics highly overlap in a set of perisylvian brain regions in association cortex (Diachek et al., 2020; Fedorenko, Nieto-Castañon, et al., 2012; Fedorenko et al., 2020; Humphries et al., 2006, 2007; Matchin et al., 2017, 2019). In response to this, some researchers have suggested that higher-level linguistic processing is distributed throughout language-related cortex, and that individual brain regions are not differentiated with respect to syntactic and semantic function (Blank et al., 2016, p. 201; Fedorenko et al., 2020; Fedorenko, Nieto-Castañón, et al., 2012; Fedorenko & Blank, 2020; Mollica et al., 2020). However, while syntax and semantics are two separate aspects of language, they are systematically connected: inverting the order of words in a sentence like *dog bites man* results in a far more surprising meaning than the original. Thus, any experimental manipulation of syntax is likely to affect semantic interpretation, and vice versa. This tight connection poses a serious obstacle to identifying the potentially distinct brain bases of higher-level linguistic functions through the use of neuroimaging experiments.

Lesion-symptom mapping can help resolve this conundrum by identifying brain regions causally involved in linguistic functions (Bates et al., 2003; Wilson, 2017). Previous lesion-symptom mapping studies have associated different language-related brain regions with different syntactic and semantic processes (Ding et al., 2020; Dronkers et al., 2004; Gleichgerrcht et al., 2016; Kristinsson et al., 2020; Magnusdottir et al., 2013; Matchin et al., 2020; Mesulam et al., 2015; Pillay et al., 2017; Riccardi et al., 2020; Rogalsky et al., 2018; Schwartz et al., 2011; Thothathiri et al., 2012; Wilson, Dronkers, et al., 2010; Wilson, Henry, et al., 2010). However, few studies report task by region interaction effects which would provide strong evidence for functional dissociations across regions. Guided by a recent theoretical model of syntax and semantics in the brain (Matchin & Hickok, 2020), we assessed three measures tapping into distinct linguistic processes: syntactic comprehension, expressive agrammatism, and semantic category word fluency, in a lesion-symptom mapping study with two partially overlapping cohorts of people with chronic left hemisphere damage following stroke (N = 121 and N=92). We hypothesized that we would find significant measure by region interaction effects, supporting selectivity for distinct higher-level linguistic processes in different brain regions.

## Materials & Methods

### *Subjects & Measures*

In two, partially overlapping groups of subjects, we assessed three different measures tapping into different aspects of syntax and semantics: (i) syntactic comprehension (Group 1, N=121), (ii) expressive agrammatism (Group 2, N=92), and (iii) semantic category word fluency (Group 2, N=92). Group 1 subjects were the same as reported in den Ouden et al. (2019) and Kristinsson et al. (2020). 47 subjects were included solely in den Ouden et al. (2019), 48 subjects were included solely in Kristinsson et al. (2020), and 26 subjects were included in both studies. Group 2 subjects were the same as reported in den Ouden et al., (2019, N = 39) and Matchin et al., (2020, N = 53).

The *syntactic comprehension* measure was derived from two different sentence-picture matching tasks reported in more detail elsewhere (den Ouden et al., 2019; Kristinsson et al., 2020). The tasks involved a range of constructions, but our focus here is on complex, semantically reversible sentences with non-canonical word order. These included object-extracted clefts (Kristinsson et al., 2020; e.g., *it is the boy that chases the cat*), object-extracted relative clauses (den Ouden et al., 2019; e.g., *the boy that the girl chased is happy*), and object-extracted Wh-questions (both studies; e.g. *which boy did the girl chase?*). Sentences of this sort have a long history in research on syntactic ability in comprehension because (i) lacking canonical English subject-verb-object word order and (ii) lacking semantic plausibility constraints (c.f., *which apple did the boy eat?*), they require syntactic analysis for determining who is doing what to whom (Caramazza & Zurif, 1976). Performance on such sentences is often compared to comprehension of simple active-voice sentences (i.e., the noun verbed the noun) as a control for speech perception, lexical processing, basic ability to infer an event structured based on a sequence of words, and working memory or decision-making resources involved in performing a sentence-picture matching task. Thus, our syntactic comprehension measure was computed as the average performance on complex sentences with performance on simple, semantically reversible active sentences covaried out using linear regression. For subjects who performed both studies (N = 26), in order to provide the most reliable estimate, scores were averaged across the two studies, which contained the same number of trials (5 trials per sentence type).

The *expressive agrammatism* measure was derived from samples of connected speech production elicited either by (i) describing the Cookie Theft picture (Goodglass & Kaplan, 1983, as reported in den Ouden et al., 2019) or (ii) retelling the story of Cinderella in their own words (MacWhinney et al., 2011, as reported in Matchin et al., 2020). The presence of expressive agrammatism was determined as described in Matchin et al., (2020). Briefly, production samples were rated independently by speech and language experts for the systematic simplification of sentence structure and omission of function words and morphemes. This resulted in a categorical assessment for each subject, either agrammatic or not. We included speech rate as a covariate using logistic regression (words per minute during the task) in order to focus on residual morphosyntactic sequencing abilities rather than general speech production abilities.

The *semantic category word fluency* measure came from the Word Fluency subtest of the Western Aphasia Battery - Revised (Kertesz, 2007), as administered by a licensed speech language pathologist. Subjects were assessed at a variable number of timepoints (mean = 2.02 assessments, SD = 1.07); we averaged their performance across these assessments in order to provide the most reliable estimate. The Word Fluency subtest involves asking the subject to name as many animals as possible within one minute (maximum score is 20). Word fluency roughly assesses two broad categories of abilities: access to conceptual-semantic representations and more general search, selection, and articulatory abilities (Chertkow & Bub, 1990; Troyer et al., 1997; Unsworth et al., 2011). Notably, a previous lesion-symptom mapping study by Baldo et al. (2006) found that deficits in the same Word Fluency task were associated with temporal and inferior parietal damage, whereas letter fluency was associated with frontal and anterior parietal damage (see Libon et al., 2009 for similar findings). This suggests that Word Fluency may be less impacted by non-linguistic mechanisms than other fluency measures, consistent with the fact that patients with semantic dementia perform worse on semantic category fluency measures than letter fluency relative to patients with Alzheimer's disease (Marczinski & Kertesz,

2006). Regardless, to focus our measure more neatly on conceptual-semantic access, we incorporated the same speech rate covariate we used for the expressive agrammatism analysis to control for non-semantic factors.

All subjects were recruited through local advertisement. They provided informed consent to participate in this study, which was approved by the Institutional Review Boards at the University of South Carolina and the Medical University of South Carolina. All subjects had at least one ischemic stroke to the left hemisphere at least six months prior to study inclusion and were also pre-morbidly right handed (self- disclosed). Demographic information for the two groups of subjects is shown in Table 1.

| | **Group 1** | **Group 2** |
|---|---|---|
| *Measures* | Syntactic Comprehension | Semantic Category Word Fluency<br><br>Expressive Agrammatism |
| *Total number of subjects* | N = 121 | N = 92 |
| *Sex* | 79 male, 42 female | 60 male, 32 female |
| *Age (years)* | 59.6 (SD = 10.6) | 58.7 (SD = 11.1) |
| *Months post-stroke at initial testing* | 47.5 (SD = 52.4) | 46.4 (SD = 49.6) |
| *Education (years)* | 15.4 (SD = 2.3) | 15.3 (SD = 2.3) |
| *Lesion Volume ($mm^3$)* | 112,652 (SD = 94,945) | 96,175 (SD = 84,514) |
| *WAB AQ* | 66 (SD = 26.9) | 75.8 (SD = 19.9) |

Table 1. Subject information for the two partially overlapping groups of subjects.

## *Brain Imaging & Lesion mapping*

We acquired anatomical MRIs and performed lesion mapping using the same parameters and procedures as described in Fridriksson et al. (2018). Neuroimaging data were collected at the University of South Carolina and the Medical University of South Carolina. Lesions were demarcated onto each subject's T2 image by an expert technician or an expert neurologist blind to the behavioral data.

Lesion overlap maps for both groups are shown in Figure 1 (a & b). Overall, there was good coverage in perisylvian cortex, including all of the selected regions of interest (described below).

## *Region of Interest (ROI) definition*

To define ROIs for language-related brain regions involved in higher-level linguistic processing, we used the statistical maps associated with a previous fMRI study on sentence processing (Matchin et al., 2017). This study compared multiple conditions, including full natural sentences (e.g., *the poet might recite a verse*) and jabberwocky sentences, which involve the substitution of

pseudowords for content words (e.g., *the tevill will sawl a pand*). The contrast of natural sentences > jabberwocky sentences highlighted a number of language-related brain regions in association cortex that are frequently identified in brain imaging studies of syntax and semantics. We therefore used this contrast with a reduced voxel-wise threshold of p < 0.01 (Figure 1c). We selected clusters corresponding to six left hemisphere regions that have been previously implicated in higher-level linguistic processing (Figure 1d), including most prominently syntax and semantics: the inferior angular gyrus (iAG), the posterior middle temporal gyrus (pMTG), the anterior superior temporal sulcus (aSTS), the temporal pole (TP), the posterior inferior frontal gyrus (pIFG), and the anterior inferior frontal gyrus (aIFG). These clusters were contiguous at local minima, and so were manually separated at these junctures to form six separate ROIs.

Figure 1. (a) Lesion overlap map for the Group 1 (N = 121), assessed for *syntactic comprehension*. The lower boundary of 12 corresponds to regions where at least 10% of subjects had damage. (b) (a) Lesion overlap map for Group 2 group (N = 92), assessed for *semantic category word fluency* and *expressive agrammatism*. The lower boundary of 9 corresponds to regions where at least 10% of subjects had damage. (c) Statistical contrast of natural sentences > jabberwocky sentences from Matchin et al., (2017) at the uncorrected voxel-wise threshold of p < 0.01. (d) Selected ROIs using the clusters derived from the contrast of natural sentences > jabberwocky sentences from Matchin et al., (2017). Red: inferior angular gyrus (iAG); blue: posterior middle temporal gyrus (pMTG); cyan: anterior superior temporal sulcus (aSTS); orange: temporal pole (TP); magenta: anterior inferior frontal gyrus (aIFG); green: posterior inferior frontal gyrus (pIFG).

## *Lesion Analyses*

We performed ROI-based univariate regression analyses using the set of six ROIs and correcting for multiple comparisons using permutation tests (10,000 permutations) in NiiStat

(https://www.nitrc.org/projects/niistat/). We performed additional uncorrected whole-hemisphere voxel-based regression analyses in NiiStat in order to ensure that our ROI analyses did not obscure the true lesion distribution associated with the effects. All univariate analyses were performed within regions that contained at least 10% of sample with lesions in that location and incorporated lesion volume as a covariate in both behavioral and brain measures (DeMarco & Turkeltaub, 2018).

We also performed three hypothesis-driven interaction analyses, one for each measure. The neuroanatomical model of Matchin & Hickok (2020) predicts that pMTG is critically involved in hierarchical syntactic processing for both comprehension and production, whereas the pIFG is involved in morpho-syntactic sequencing primarily for production. With respect to *syntactic comprehension*, we tested the interaction between the pMTG and pIFG. We expected that the pMTG would be implicated, but not pIFG. By contrast, with respect to *expressive agrammatism*, we tested the interaction between pMTG and pIFG, with the expectation that this would primarily reveal deficits in morpho-syntactic sequencing abilities in production, hypothesized to be processed by the pIFG (see Matchin & Hickok, 2020 for a review). With respect to *semantic category word fluency*, we tested the interaction between iAG and aSTS, as both have been strongly associated with conceptual-semantic interpretation, but have been hypothesized to house distinct conceptual representations (Binder & Desai, 2011; Matchin & Hickok, 2020; Schwartz et al., 2011). To test these interactions, we first calculated proportion damage to each ROI and adjusted the data using a rationalized arcsine transform (Studebaker, 1985), and then computed residual damage values by covarying out the effect of lesion volume. We then assessed the region by measure interaction effect in linear regression for each of the three measures of interest in SPSS. We corrected for multiple comparisons using a Bonferroni correction with an adjusted alpha threshold of $p < 0.0167$ for each of the three independent comparisons, controlling the total error at $p < 0.05$.

## Results & Discussion

ROI results are shown in Figure 2a. For *syntactic comprehension*, two ROIs showed a significant negative association between percent damage and behavioral scores: pMTG ($Z = -3.23$, $p = 0.00087$) and aSTS ($Z = -2.56$, $p = 0.00695$). For *expressive agrammatism*, two ROIs showed a significant positive association between percent damage and behavioral scores: pIFG ($Z = 3.08$, $p = 0.00135$) and aIFG ($Z = 2.54$, $p = 0.00554$). For the regression analysis of *semantic category word fluency*, one ROI showed a significant negative association between percent damage and behavioral scores: iAG ($Z = -3.73$, $p = 0.000096$). Uncorrected analyses using an individual voxel-level threshold of $p < 0.05$ generally support our ROI analyses (Figure 2b), although the strongest localization of *expressive agrammatism* appears to have been in the inferior frontal sulcus (IFS) rather than the IFG itself.

Figure 2. Corrected ROI-based and uncorrected whole-brain analyses for each of the three measures of syntax/semantics. Each measure is identified at top in bold, corresponding to all of the figures underneath. (a) Regression analyses within the selected ROIs, corrected for multiple comparisons using permutation tests (p < 0.05). (b) Uncorrected voxel-wise analyses (p < 0.05). (c) Scatter plots illustrating three hypothesis-driven *region by measure* interaction analyses, corrected for multiple comparisons using a Bonferroni correction (individual p < 0.0167). pMTG = posterior middle temporal gyrus; pIFG = posterior inferior frontal gyrus; iAG = inferior angular gyrus; aSTS = anterior superior temporal sulcus.

Scatterplots illustrating the *region by measure* interactions are shown in Figure 2c. For *syntactic comprehension*, there was a significant interaction between the pMTG and pIFG, such that damage to pMTG was significantly more associated with deficits than the pIFG, F(1,119) = 8.928, p = 0.003. For *expressive agrammatism*, there was nonsignificant interaction between the effects in pMTG and pIFG, such that damage to pIFG was more associated with agrammatic

speech than the pMTG, F(1,90) = 3.25, p = 0.075. For *semantic category word fluency*, there was a significant interaction between the effects in iAG and aSTS, such that damage to iAG was significantly more associated with deficits than the aSTS (F(1,119) = 6.799, p = 0.011).

These results narrow down possible functions of these brain regions with respect to syntax and semantics, and suggest that neuroimaging research critically needs to incorporate insights from lesion symptom mapping in order to understand the architecture of language in the brain. Namely, the *region by measure* interactions we identified present strong challenges to the hypothesis that language-related brain regions underlie a similar function with respect to syntactic or and/or higher-level linguistic processing (Blank et al., 2016; Caplan et al., 1996; Dick et al., 2001; Fedorenko et al., 2020; Fedorenko & Blank, 2020; Mollica et al., 2020).

Our results are much more consistent with neuroanatomical models that posit distinct syntactic and semantic functions to different regions within the language network (Bornkessel-Schlesewsky & Schlesewsky, 2013; Friederici, 2017; Hagoort, 2014; Matchin & Hickok, 2020; Tyler & Marslen-Wilson, 2008) than with the hypothesis of undifferentiated linguistic function of language-related brain regions. In particular, the model presented by Matchin & Hickok (2020) provides clear explanations for the identified effects and interactions. The model posits that the posterior middle temporal gyrus (pMTG) is critical for processing hierarchical structure for both production and comprehension. Damage to the pMTG was significantly associated with semantic comprehension deficits even when compared to the pIFG. This reinforces previous lesion-symptom mapping studies that have found an association between residual syntactic comprehension scores (regressing out single word comprehension or production) with damage to the posterior temporal lobe (Kristinsson et al., 2020; Pillay et al., 2017; Rogalsky et al., 2018).

Matchin & Hickok suggest that the pIFG is hypothesized to underlie a linear morpho-syntactic system that sequences words and morphemes during speech production, but is not critical for lexical retrieval or syntactic structure building in comprehension. As we have shown previously in subsets of these data (den Ouden et al., 2019; Matchin et al., 2020), and consistent with previous research (Sapolsky et al., 2010; Wilson, Henry, et al., 2010), damage to the pIFG was significantly associated with agrammatic speech deficits, but was not associated with any comprehension deficits. The fact that the voxel-wise analysis identified the main locus of damage with the IFG, and that the interaction between pIFG and pMTG only approached significance, suggests that the role for middle frontal regions in syntax should be further investigated.

The only region significantly associated with deficits in semantic category word fluency was the iAG, similar to a previous study which also identified temporal lobe regions (Baldo et al., 2006). Following previous authors (Binder & Desai, 2011; Mesulam et al., 2015; Schwartz et al., 2011), the Matchin & Hickok model posits that the aSTS the iAG are involved in distinct aspects of conceptual-semantic processing. Namely, the iAG is hypothesized to support relations among concepts, critically encoded by verbs, whereas the aSTS is hypothesized to support identifying features of individual entities. In addition, the iAG has been associated with episodic memory (Binder et al., 2009; Maguire, 2012; Svoboda et al., 2006). With respect to semantic category word fluency (regressing out general speech rate), we observed a robust interaction effect between *semantic category word fluency* and region such that the iAG was much more strongly

associated with reduced scores than the aSTS. This suggests that subjects may navigate the task by retrieving animal names based on episodic memories involving animals and/or course thematic relations among entities, without needing to retrieve finer taxonomic features. This would predict that a similar task, for instance requiring subjects to identify all of the features of a particular animal or specific subtypes of an animal, would result in greater association between lower scores and damage to the aSTS.

While we find that different language-related brain regions are associated with different syntactic and semantic deficits, note that we are not claiming that we will find specific brain regions that correspond to some syntactic tasks but not to *any* semantic tasks, or vice versa. Rather, many supposedly "syntactic" or "semantic" tasks likely engage both mechanisms. Thus, we suggest that a key goal for future research in identifying functional dissociations among regions is to more deeply interrogate the possible syntactic and semantic functions elicited by particular experimental tasks.

## Funding & Acknowledgments

This research was supported by National Institute on Deafness and Other Communication Disorders grants P50 U01 DC011739 and R01 DC014664 awarded to Julius Fridriksson, and a National Institute on Deafness and Other Communication Disorders grant awarded to Alexandra Basilakos, T32 DC 014435. We would like to thank Grigori Yourganov and Jonathan Venezia for advice on statistical analysis. We would also like to thank Leigh Ann Spell, Allison Croxton, Anna Doyle, Michele Martin, Katie Murphy, and Sara Sayers for their assistance with data collection, and graduate student clinicians in the Aphasia Lab for transcribing and coding speech samples.

## Competing Interests

The authors declare no competing interests.

## References

Baldo, J. V., Schwartz, S., Wilkins, D., & Dronkers, N. F. (2006). Role of frontal versus temporal cortex in verbal fluency as revealed by voxel-based lesion symptom mapping. *Journal of the International Neuropsychological Society*, *12*(06). https://doi.org/10.1017/S1355617706061078

Bates, E., Wilson, S. M., Saygin, A. P., Dick, F., Sereno, M. I., Knight, R. T., & Dronkers, N. F. (2003). Voxel-based lesion–symptom mapping. *Nature Neuroscience*, *6*(5), 448–450. https://doi.org/10.1038/nn1050

Berwick, R. C., & Chomsky, N. (2016). *Why only us: Language and evolution*. MIT press.

Binder, J. R., & Desai, R. H. (2011). The neurobiology of semantic memory. *Trends in Cognitive Sciences*, *15*(11), 527–536. https://doi.org/10.1016/j.tics.2011.10.001

Binder, J. R., Desai, R. H., Graves, W. W., & Conant, L. L. (2009). Where Is the Semantic System? A Critical Review and Meta-Analysis of 120 Functional Neuroimaging Studies. *Cerebral Cortex*, *19*(12), 2767–2796. https://doi.org/10.1093/cercor/bhp055

Blank, I., Balewski, Z., Mahowald, K., & Fedorenko, E. (2016). Syntactic processing is distributed across the language system. *NeuroImage*, *127*, 307–323. https://doi.org/10.1016/j.neuroimage.2015.11.069

Bornkessel-Schlesewsky, I., & Schlesewsky, M. (2013). Reconciling time, space and function: A new dorsal–ventral stream model of sentence comprehension. *Brain and Language*, *125*(1), 60–76. https://doi.org/10.1016/j.bandl.2013.01.010

Caplan, D., Hildebrandt, N., & Makris, N. (1996). Location of lesions in stroke patients with deficits in syntactic processing in sentence comprehension. *Brain*, *119*(3), 933–949. https://doi.org/10.1093/brain/119.3.933

Caramazza, A., & Zurif, E. B. (1976). Dissociation of algorithmic and heuristic processes in language comprehension: Evidence from aphasia. *Brain and Language*, *3*(4), 572–582. https://doi.org/10.1016/0093-934X(76)90048-1

Chertkow, H., & Bub, D. (1990). Semantic Memory Loss in Dementia of Alzheimer's Type: What do Various Measures Measure? *Brain*, *113*(2), 397–417. https://doi.org/10.1093/brain/113.2.397

DeMarco, A. T., & Turkeltaub, P. E. (2018). A multivariate lesion symptom mapping toolbox and examination of lesion☐volume biases and correction methods in lesion☐symptom mapping. *Human Brain Mapping*, *39*(11), 4169–4182. https://doi.org/10.1002/hbm.24289

den Ouden, D., Malyutina, S., Basilakos, A., Bonilha, L., Gleichgerrcht, E., Yourganov, G., Hillis, A. E., Hickok, G., Rorden, C., & Fridriksson, J. (2019). Cortical and structural☐connectivity damage correlated with impaired syntactic processing in aphasia. *Human Brain Mapping*, *40*(7), 2153–2173. https://doi.org/10.1002/hbm.24514

Diachek, E., Blank, I., Siegelman, M., Affourtit, J., & Fedorenko, E. (2020). The domain-general multiple demand (MD) network does not support core aspects of language comprehension: A large-scale fMRI investigation. *The Journal of Neuroscience*, JN-RM-2036-19. https://doi.org/10.1523/JNEUROSCI.2036-19.2020

Dick, F., Bates, E., Wulfeck, B., Utman, J. A., Dronkers, N., & Gernsbacher, M. A. (2001). Language deficits, localization, and grammar: Evidence for a distributive model of language breakdown in aphasic patients and neurologically intact individuals. *Psychological Review*, *108*(4), 759–788. https://doi.org/10.1037/0033-295X.108.4.759

Ding, J., Chen, K., Liu, H., Huang, L., Chen, Y., Lv, Y., Yang, Q., Guo, Q., Han, Z., & Lambon Ralph, Matthew. A. (2020). A unified neurocognitive model of semantics language social behaviour and face recognition in semantic dementia. *Nature Communications*, *11*(1), 2595. https://doi.org/10.1038/s41467-020-16089-9

Dronkers, N. F., Wilkins, D. P., Van Valin, R. D., Redfern, B. B., & Jaeger, J. J. (2004). Lesion analysis of the brain areas involved in language comprehension. *Cognition*, *92*(1–2), 145–177. https://doi.org/10.1016/j.cognition.2003.11.002

Estes, Z., Golonka, S., & Jones, L. L. (2011). Thematic Thinking. In *Psychology of Learning and Motivation* (Vol. 54, pp. 249–294). Elsevier. https://doi.org/10.1016/B978-0-12-385527-5.00008-5

Faroqi☐Shah, Y., & Milman, L. (2018). Comparison of animal, action and phonemic fluency in aphasia. *International Journal of Language & Communication Disorders*, *53*(2), 370–384. https://doi.org/10.1111/1460-6984.12354

Fedorenko, E., & Blank, I. A. (2020). Broca's Area Is Not a Natural Kind. *Trends in Cognitive Sciences*, *24*(4), 270–284. https://doi.org/10.1016/j.tics.2020.01.001

Fedorenko, E., Blank, I. A., Siegelman, M., & Mineroff, Z. (2020). Lack of selectivity for syntax relative to word meanings throughout the language network. *Cognition*, *203*, 104348. https://doi.org/10.1016/j.cognition.2020.104348

Fedorenko, E., Nieto-Castañón, A., & Kanwisher, N. (2012). Syntactic processing in the human brain: What we know, what we don't know, and a suggestion for how to proceed. *Brain and Language*, *120*(2), 187–207. https://doi.org/10.1016/j.bandl.2011.01.001

Fedorenko, E., Nieto-Castañon, A., & Kanwisher, N. (2012). Lexical and syntactic representations in the brain: An fMRI investigation with multi-voxel pattern analyses. *Neuropsychologia*, *50*(4), 499–513. https://doi.org/10.1016/j.neuropsychologia.2011.09.014

Fridriksson, J., den Ouden, D.-B., Hillis, A. E., Hickok, G., Rorden, C., Basilakos, A., Yourganov, G., & Bonilha, L. (2018). Anatomy of aphasia revisited. *Brain*, *141*(3), 848–862. https://doi.org/10.1093/brain/awx363

Friederici, A. D. (2017). *Language in our brain: The origins of a uniquely human capacity*. MIT Press.

Gleichgerrcht, E., Fridriksson, J., Rorden, C., Nesland, T., Desai, R., & Bonilha, L. (2016). Separate neural systems support representations for actions and objects during narrative speech in post-stroke aphasia. *NeuroImage: Clinical*, *10*, 140–145. https://doi.org/10.1016/j.nicl.2015.11.013

Goodglass, H., & Kaplan, E. (1983). *Boston Diagnostic Aphasia Examination (BDAE)* (2nd ed.). Lippincott Williams & Wilkins.

Hagoort, P. (2014). Nodes and networks in the neural architecture for language: Broca's region and beyond. *Current Opinion in Neurobiology*, *28*, 136–141. https://doi.org/10.1016/j.conb.2014.07.013

Humphries, C., Binder, J. R., Medler, D. A., & Liebenthal, E. (2006). Syntactic and Semantic Modulation of Neural Activity during Auditory Sentence Comprehension. *Journal of Cognitive Neuroscience*, *18*(4), 665–679. https://doi.org/10.1162/jocn.2006.18.4.665

Humphries, C., Binder, J. R., Medler, D. A., & Liebenthal, E. (2007). Time course of semantic processes during sentence comprehension: An fMRI study. *NeuroImage*, *36*(3), 924–932. https://doi.org/10.1016/j.neuroimage.2007.03.059

Kertesz, A. (2007). *Western Aphasia Battery- Revised*. Grune and Stratton.

Kristinsson, S., Thors, H., Yourganov, G., Magnusdottir, S., Hjaltason, H., Stark, B. C., Basilakos, A., den Ouden, D.-B., Bonilha, L., Rorden, C., Hickok, G., Hillis, A., & Fridriksson, J. (2020). Brain Damage Associated with Impaired Sentence Processing in Acute Aphasia. *Journal of Cognitive Neuroscience*, *32*(2), 256–271. https://doi.org/10.1162/jocn_a_01478

Libon, D. J., McMillan, C., Gunawardena, D., Powers, C., Massimo, L., Khan, A., Morgan, B., Farag, C., Richmond, L., Weinstein, J., Moore, P., Coslett, H. B., Chatterjee, A., Aguirre, G., & Grossman, M. (2009). Neurocognitive contributions to verbal fluency deficits in frontotemporal lobar degeneration. *Neurology*, *73*(7), 535–542. https://doi.org/10.1212/WNL.0b013e3181b2a4f5

Lin, E. L., & Murphy, G. L. (2001). Thematic Relations in Adults' Concepts. *Journal of Experimental Psychology: General*, *130*(1), 3–38.

MacWhinney, B., Fromm, D., Forbes, M., & Holland, A. (2011). AphasiaBank: Methods for studying discourse. *Aphasiology*, *25*(11), 1286–1307. https://doi.org/10.1080/02687038.2011.589893

Magnusdottir, S., Fillmore, P., den Ouden, D. B., Hjaltason, H., Rorden, C., Kjartansson, O., Bonilha, L., & Fridriksson, J. (2013). Damage to left anterior temporal cortex predicts impairment of complex syntactic processing: A lesion-symptom mapping study: ATL Damage Impairs Complex Syntax Processing. *Human Brain Mapping*, *34*(10), 2715–2723. https://doi.org/10.1002/hbm.22096

Maguire, E. A. (2012). Neuroimaging studies of autobiographical event memory. *Current Biology*, *22*(24), 1441–1451. https://doi.org/10.1016/j.cub.2012.10.007

Marczinski, C. A., & Kertesz, A. (2006). Category and letter fluency in semantic dementia, primary progressive aphasia, and Alzheimer's disease. *Brain and Language*, *97*(3), 258–265. https://doi.org/10.1016/j.bandl.2005.11.001

Matchin, W., Basilakos, A., Stark, B. C., den Ouden, D.-B., Fridriksson, J., & Hickok, G. (2020). Agrammatism and paragrammatism: A cortical double dissociation revealed by lesion-symptom mapping. *Neurobiology of Language*, 1–47. https://doi.org/10.1162/nol_a_00010

Matchin, W., Brodbeck, C., Hammerly, C., & Lau, E. (2019). The temporal dynamics of structure and content in sentence comprehension: Evidence from fMRI-constrained MEG. *Human Brain Mapping*, *40*(2), 663–678. https://doi.org/10.1002/hbm.24403

Matchin, W., Hammerly, C., & Lau, E. (2017). The role of the IFG and pSTS in syntactic prediction: Evidence from a parametric study of hierarchical structure in fMRI. *Cortex*, *88*, 106–123. https://doi.org/10.1016/j.cortex.2016.12.010

Matchin, W., & Hickok, G. (2020). The Cortical Organization of Syntax. *Cerebral Cortex*, *30*(3), 1481–1498.

Mesulam, M.-M., Thompson, C. K., Weintraub, S., & Rogalski, E. J. (2015). The Wernicke conundrum and the anatomy of language comprehension in primary progressive aphasia. *Brain*, *138*(8), 2423–2437. https://doi.org/10.1093/brain/awv154

Mollica, F., Siegelman, M., Diachek, E., Piantadosi, S. T., Mineroff, Z., Futrell, R., Kean, H., Qian, P., & Fedorenko, E. (2020). Composition is the Core Driver of the Language-selective Network. *Neurobiology of Language*, *1*(1), 104–134. https://doi.org/10.1162/nol_a_00005

Pillay, S. B., Binder, J. R., Humphries, C., Gross, W. L., & Book, D. S. (2017). Lesion localization of speech comprehension deficits in chronic aphasia. *Neurology*, *88*(10), 970–975. https://doi.org/10.1212/WNL.0000000000003683

Riccardi, N., Yourganov, G., Rorden, C., Fridriksson, J., & Desai, R. (2020). Degradation of Praxis Brain Networks and Impaired Comprehension of Manipulable Nouns in Stroke. *Journal of Cognitive Neuroscience*, *32*(3), 467–483. https://doi.org/10.1162/jocn_a_01495

Rogalsky, C., LaCroix, A. N., Chen, K.-H., Anderson, S. W., Damasio, H., Love, T., & Hickok, G. (2018). The Neurobiology of Agrammatic Sentence Comprehension: A Lesion Study. *Journal of Cognitive Neuroscience*, *30*(2), 234–255. https://doi.org/10.1162/jocn_a_01200

Sapolsky, D., Bakkour, A., Negreira, A., Nalipinski, P., Weintraub, S., Mesulam, M.-M., Caplan, D., & Dickerson, B. C. (2010). Cortical neuroanatomic correlates of symptom severity in primary progressive aphasia. *Neurology*, *75*(4), 358–366. https://doi.org/10.1212/WNL.0b013e3181ea15e8

Schwartz, M. F., Kimberg, D. Y., Walker, G. M., Brecher, A., Faseyitan, O. K., Dell, G. S., Mirman, D., & Coslett, H. B. (2011). Neuroanatomical dissociation for taxonomic and

thematic knowledge in the human brain. *Proceedings of the National Academy of Sciences*, *108*(20), 8520–8524. https://doi.org/10.1073/pnas.1014935108

Studebaker, G. A. (1985). A "Rationalized" Arcsine Transform. *Journal of Speech and Hearing Research*, *28*(03), 455–462.

Svoboda, E., McKinnon, M. C., & Levine, B. (2006). The functional neuroanatomy of autobiographical memory: A meta-analysis. *Neuropsychologia*, *44*(12), 2189–2208. https://doi.org/10.1016/j.neuropsychologia.2006.05.023

Thothathiri, M., Kimberg, D. Y., & Schwartz, M. F. (2012). The Neural Basis of Reversible Sentence Comprehension: Evidence from Voxel-based Lesion Symptom Mapping in Aphasia. *Journal of Cognitive Neuroscience*, *24*(1), 212–222. https://doi.org/10.1162/jocn_a_00118

Troyer, A. K., Moscovitch, M., & Winocur, G. (1997). Clustering and Switching as Two Components of Verbal Fluency: Evidence from Younger and Older Healthy Adults. *Neuropsychology*, *11*(1), 138–146.

Tyler, L. K., & Marslen-Wilson, W. (2008). Fronto-temporal brain systems supporting spoken language comprehension. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *363*(1493), 1037–1054. https://doi.org/10.1098/rstb.2007.2158

Unsworth, N., Spillers, G. J., & Brewer, G. A. (2011). Variation in verbal fluency: A latent variable analysis of clustering, switching, and overall performance. *Quarterly Journal of Experimental Psychology*, *64*(3), 447–466. https://doi.org/10.1080/17470218.2010.505292

Wilson, S. M. (2017). Lesion-symptom mapping in the study of spoken language understanding. *Language, Cognition and Neuroscience*, *32*(7), 891–899. https://doi.org/10.1080/23273798.2016.1248984

Wilson, S. M., Dronkers, N. F., Ogar, J. M., Jang, J., Growdon, M. E., Agosta, F., Henry, M. L., Miller, B. L., & Gorno-Tempini, M. L. (2010). Neural Correlates of Syntactic Processing in the Nonfluent Variant of Primary Progressive Aphasia. *Journal of Neuroscience*, *30*(50), 16845–16854. https://doi.org/10.1523/JNEUROSCI.2547-10.2010

Wilson, S. M., Henry, M. L., Besbris, M., Ogar, J. M., Dronkers, N. F., Jarrold, W., Miller, B. L., & Gorno-Tempini, M. L. (2010). Connected speech production in three variants of primary progressive aphasia. *Brain*, *133*(7), 2069–2088. https://doi.org/10.1093/brain/awq129