Visual Effort Moderates Postural Cascade Dynamics

Madhur Mangalam,[1] I-Chieh Lee,[2] Karl M. Newell,[3] and Damian G. Kelty-Stephen[4]

[1]Department of Physical Therapy, Movement and Rehabilitation Sciences, Northeastern University, Boston, MA 02115, USA; [2]UNC-NC State Joint Department of Biomedical Engineering, UNC-Chapel Hill, Chapel Hill, NC 27514, USA; [3]Department of Kinesiology, University of Georgia, Athens, GA 30602, USA; [4]Department of Psychology, Grinnell College, Grinnell, IA 50112, USA

Address correspondence to Madhur Mangalam, Department of Physical Therapy, Movement and Rehabilitation Sciences, Northeastern University, Boston, MA 02115. Electronic mail: m.mangalam@northeastern.edu

Abstract—Standing still and focusing on a visible target in front of us is a preamble to many coordinated behaviors (e.g., reaching an object). Hiding behind its apparent simplicity is a deep layering of texture at many scales. The task of standing still laces together activities at multiple scales: from ensuring that a few photoreceptors on the retina cover the target in the visual field on an extremely fine-scale to synergies spanning the limbs and joints at smaller scales to the mechanical layout of the ground underfoot and optic flow in the visual field on the coarser scales. Here, we used multiscale probability distribution function (PDF) analysis to show that postural fluctuations exhibit similar statistical signatures of cascade dynamics as found in fluid flow. Critically, the oculomotor strain of visually fixating different distances moderates postural cascade dynamics. Visually fixating at a comfortable viewing distance elicited posture with a similar cascade dynamics as posture with eyes closed. Greater viewing distances known to stabilize posture showed more diminished cascade dynamics. In contrast, nearest and farthest viewing distances requiring greater oculomotor strain to focus on targets elicited a dramatic strengthening postural cascade dynamics, reflecting active postural adjustments. Critically, these findings suggest that vision stabilizes posture by reconfiguring the prestressed poise that prepares the body to interact with different spatial layouts.

Keywords—Center of pressure, Ergonomics, Inverse power law, Postural control, Quiet stance, Lévy walk

INTRODUCTION

Standing still and visually fixating on a target in front of us is a seemingly simple task.
However, hiding within this apparent simplicity is a deep layering of texture at many scales.
On an extremely fine-scale, task completion depends only on ensuring that a few
photoreceptors on the retina cover the entire target in the visual field. Standing still and
visually fixating at much coarser scales rests on the mechanical layout of the ground
underfoot and optic flow in the visual field before you. Between optic flow and mechanical
layout, standing still relies on synergies spanning the limbs and joints at smaller scales and
subtle postural perturbations, triggering reflex arcs and muscle twitches, at yet smaller scales.
The task of standing still and visually fixating laces together activities at these multiple scales,
entailing a close connection between subtle postural fluctuations and photoreceptor support
of the target's retinal image. Consequently, the postural center of pressure (CoP) fluctuations
—which directly relate to sway—exhibit similar statistical signatures of cascade dynamics as
found in fluid flow.[37] The metaphor of 'cascades,' as in tumbling water that accelerates and
splashed down a rockface, captures some critical aspects of the dynamics of emergent
behavior.[21,33,35] Destabilizing posture emphasizes the postural cascade dynamics.[36,38] Here,
we propose that perturbing the visual system has a similar effect of strengthening postural
cascade dynamics, in which the optical flow varies with fixation distance, as a meandering
stream joins other streams and gathers momentum into rushing rapids, becoming more
turbulent along the way. Whereas multiscale entropy addresses the difference of variation
across different scales,[9,10] cascade dynamics explicitly implicates nonlinear interactions lacing
these scales together.

Visually guided posture presents a curious case in which light entails mechanical
change in bodily center of mass.[12,15,42,48,50,51] Beyond minuscule photoreceptor responses,
large-scale layout of visible surfaces supports optic flow with bodily sway. Optic flow is the
expansion, contraction, or rotation of visible surfaces as the body moves forward or
backward, or turns to the side, respectively. Relative accelerations of visible surfaces provide
the standing participant with rich information about the layout of objects. Placing a fixation
target at different viewing distances perturbs the optic flow: changing the relative
accelerations as well as imposing retinal-specific constraints of oculomotor convergence (i.e.,
oculomotor strain in specific or visual effort in general),[14,28] prompting new head- and torso-
sway (Fig. 1). Comfortable viewing distances (50-100cm) requiring least oculomotor strain
should most resemble posture with eyes closed.[40,49] Medium viewing distances beyond most-
comfortable viewing in the range of 200 cm stabilize posture.[20] Extremely distant or extremely
near targets will be harder to focus on and have a much slower or faster optic flow,
respectively, requiring more sway to resolve the image.

Testing these hypotheses requires examining the divergence of postural fluctuations
from Gaussian form. Gaussian distributions are classically the result of adding very many
independent (i.e., non-interacting) variables. Interactivity amongst the participating variables
promotes persistent if also intermittent, uneven growth of variance with timescale. Under the
central limit theorem, homogeneity of variance keeps measurements within short, thin tails of
a Gaussian distribution. But intermittent growth in variance manifests in distributions with
longer, heavier tails specifically due to nonlinear temporal correlations.[8,31] Testing these tails is
even more difficult because they are by definition only populated by extreme and relatively
rare values. However, fortunately, hydrodynamic-cascade research has yielded an analytical
method for estimating non-Gaussianity based on querying better-populated regions of the
distribution.[5] Applying this method to physiological measures produces estimates of non-

Gaussianity that predict clinical outcomes[16,22–24,53] and that change with variation in both
endogenous postural control and exogenous postural demands.[36,38] Here, we used this
method to investigate how constraints imposed by visual effort affect postural cascade
dynamics. Specifically, we predicted that postural cascade dynamics while fixating at a 50-cm
distance would resemble eyes-closed posture (Hypothesis 1), and that postural cascade
dynamics would be weakest while fixating at medium (i.e., 130- and 220-cm) distances
(Hypothesis 2) but strongest at extremely near (i.e., 25-cm) and extremely far (i.e., 305-cm)
distances (Hypothesis 3).

89   **MATERIALS AND METHODS**
90   The present study is a reanalysis of data from a previous study examining postural
91 control in different viewing conditions.[29]

92   *Participants*
93   Seven men and eight women (18–40 years old) participated after providing
94 institutionally-approved informed consent.

95   *Experimental Setup and Procedure*
Each participant stood barefoot on a pair of force plates (AMTI Inc., Watertown, MA)—
one foot on each force plate, 25 cm apart, in front of a white background spanning his/her
field of view (Fig. 1 A). A square white screen (5×5 cm) mounted on a tripod—to be used as a
reference for postural adjustment—was placed at specific distances in front of the participant.
This setting ensured that the participant could not use a secondary object/surface in the
environment as a reference for postural adjustment. From the rear side of the participant, a
laser pen projected a static light point on the center of the screen. The 3D moments and
ground reaction forces were measured using the two force plates at 100 Hz.

Each participant was tested in six different viewing conditions: the eyes closed and
eyes fixated on the light point on the screen placed at a distance of 25, 50, 135, 220, and 305
cm in the front. Each of the six viewing conditions was repeated twice. In each trial, each
participant was instructed to minimize the postural sway by maintaining visual fixation at the
light point projected on the screen for 120 s.

Each participant completed 18 trials (6 conditions × 3 trials) in a single 90-min session.
The order of the trials was randomized for each participant. Each participant was allowed to
take a break if requested, and rest for at least 5 min was provided between every six trials to
avoid fatigue.

113   *Data Processing*
All data processing was performed in MATLAB 2019b (Matlab Inc., Natick, MA). Trial-
by-trial ground reaction forces yielded a 2D center of pressure (CoP) series, each dimension
describing CoP position along anterior-posterior (AP) and medial-lateral (ML) axes. Each 60-s
trial yielded 12000-sample 2D CoP series and 11999-sample 2D CoP displacement series.
Finally, a 1D CoP sample-to-sample planar Euclidean displacement (PED) series described
postural fluctuations along the transverse plane of the body. Iterated Amplitude Adjusted
Fourier Transformation (IAAFT) provided phase-randomized surrogates using original series'
spectral amplitudes to preserve only linear temporal correlations.[47]

122   *Canonical Indices of Endogenous Postural Fluctuations*

Three linear indices were computed for each CoP PED series: (i) Mean of all
fluctuations (CoP_PED_*Mean*), (ii) Standard deviation (CoP_PED_*SD*), and (iii) Root mean
square (CoP_PED_*RMSE*).

Four nonlinear indices were computed for each CoP PED series: (i) Sample entropy
(CoP_PED_*SampEn*), indexing the extent of complexity in the signal.[45] We used m = 2 and r
= 0.2 for computing the *SampEn*.[26] (ii) Multiscale sample entropy (CoP_PED_*MSE*), indexing
signal complexity over multiple timescales.[9] We used m = 2, r = 0.2, and $τ$ = 20 for computing
the *MSE*.[26] (iii) We used detrended fluctuation analysis[44] to compute Hurst's exponent, H_{fGn},
indexing temporal correlations in original CoP PED series (CoP_PED_$H_{fGn\_Original}$) and shuffled
versions of each series (CoP_PED_$H_{fGn\_Shuffled}$) using the scaling region: 4, 8, 12,… 1024.[34]
Shuffling destroys the temporal structure of a signal. (iv) Multifractal spectrum width, $Δα$,
indexing the extent of multifractal temporal correlations in the signal. Chhabra and Jensen's
direct method[6] was used to compute $Δα$ for the original CoP series (CoP_PED_$Δα_{Original}$) and
the IAAFT surrogate (CoP_PED_$Δα_{Surrogate}$).[35]

Multiscale Probability Density Function (PDF) Analysis

Multiscale PDF analysis characterizes the distribution of abrupt changes in CoP PED
series $\{b(t)\}$ using the PDF tail. The first step is to generate $\{B(t)\}$ by integrating $\{b(t)\}$ after
centering by the mean b_{ave} (Fig. 2A):

141
$$B(t)=\sum_{i=1}^{t/\Delta t} b(i\,\Delta t)-b_{ave}.$$

A a 3rd order polynomial detrends $\{B(t)\}$ within k overlapping windows of length $2s$, s being the
timescale (Fig. 2B). Intermittent deviation $\Delta_s B(t)$ in k^{th} window from $1+s(k-1)$ to sk in the
detrended time series $\{B^d(t)=B(t)-f_{fit}(t)\}$ is computed as $\Delta_s B^d(t)=B^d(t+s)-B^d(t)$, where
$1+s(k-1)\leq t\leq sk$ and $f_{fit}(t)$ is the polynomial representing the local trend of $\{B(t)\}$, of which the
elimination assures the zero-mean probability density function in the next step (Fig. 2C).
Finally, $\Delta_s B$ is normalized by the SD (i.e., variance is set to one) to quantify the PDF.

To quantify the non-Gaussianity of $\Delta_s B$ at timescale s, the standardized PDF
constructed from all the $\Delta_s B(t)$ values is approximated by the Castaing model,[5] with λ_s as a
single parameter characterizing the non-Gaussianity of the PDF. λ_s is estimated as

151
$$\lambda_s^2=\frac{2}{q(q-2)}\left[\ln\left(\sqrt{\pi}\left\langle|\Delta_s B|^q\right\rangle\right)+\ln\Gamma\left(\frac{q+1}{2}\right)-\frac{q}{2}\ln 2\right],$$

where $\left\langle|\Delta_s B|^q\right\rangle$ denotes an estimated value of q^{th} order absolute moment of $\{\Delta_s B\}$. As λ_s^2
increases, the PDF becomes increasingly peaked and fat-tailed (Fig. 3A). λ_s^2 can be estimated
by Eq. (7) based on q^{th} order absolute moment of a time series independent of q. Estimating
λ_s^2 based on 0.2th moment (q= 0.2) emphasizes the center part of the PDF, reducing the
effects of extreme deviations due to heavy-tails and kurtosis. We used 0.2th moment because
estimates of λ_s^2 for a time series of ~ 12000 samples are more accurate at lower values of q.[25]

Cascade-type multiplicative processes yield the inverse relationship $\lambda_s^2 \sim -\log s$ (Fig.
3B).[23] For the present purposes, we quantified λ_s^2 for each original CoP PED series and
corresponding IAAFT surrogate at timescales 5 to 1000 samples (i.e., 50 ms to 10 s) at steps
of 5 samples (50 ms; Fig. 4).

To test explicitly for Lévy-likeness, we used Akaike Information Criterion (AIC) weights
obtained via the more common maximum likelihood estimation (MLE) to determine whether
the PDF at each timescale was a power law, lognormal, exponential, or gamma distribution.
We include this analysis also to complement multiscale PDF analysis to portray the analytical
effect of sensitivity to different portions of the distribution.

167   *Statistical Analysis*
To test for crossovers in non-Gaussianity, a linear mixed-effect (LME) model using
lmer[1] in R package *lme4* tested λ_s^2 vs. log-timescale curves for orthogonal linear and quadratic
polynomials, for interactions with grouping variables (Viewing condition×Original, where
Original encoded differences in original series from surrogates) and with indices of
endogenous postural fluctuations (Section 2.4.). Statistical significance was assumed at the
alpha level of 0.05 using R package *lmerTest*.[27] To test how lognormality changed with log-
timescale, a generalized linear mixed-effect (GLME) fit changes in Lognormality as a
dichotomous variable using orthogonal linear, quadratic, and cubic polynomials and tested
interaction effects of grouping variables (Viewing condition×Original) with those polynomials
using *glmer* in lme4.[30]

179   **RESULTS**
The eyes-closed condition elicited nonlinearity in non-Gaussianity with log-timescale,
showing a stronger linear increase in λ_s^2 up to a maximum at roughly 75 samples or 750 ms (*b*
= 0.95, *p* = 0.031) and a stronger negative quadratic, downward-facing parabolic form (*b* = –
3.78, *p* < 0.001) for the original CoP PED series than the surrogates (Table S1; Fig. 5A).
Fixating the eyes at 25 and 305 cm elicited a stronger linear increase (*b* = 2.43, *p* < 0.001;
and *b* = 3.09, *p* < 0.001, receptively) in the 25- and 305- cm viewing conditions while
reversing the downward-facing parabolic form of the eyes-closed condition more weakly (*b* =
2.43, *p* < 0.001; and *b* = 1.85, *p* = 0.003, respectively) of λ_s^2 vs. log-timescale curves (Figs. 5B
and 5F). Fixating the eyes at 135 and 220 cm elicited a significant reversal of the linear form
(*b* = –2.98, *p* < 0.001; and *b* = –2.32, *p* < 0.001, respectively) and a significant reversal of the
downward-facing parabolic form (*b* = 1.81, *p* = 0.004; and *b* = 2.00, *p* = 0.001, respectively) of
λ_s^2 vs. log-timescale curves in Figs. 5D and 5E. In contrast, fixating the eyes at 50 cm neither
affected the linear form of λ_s^2 vs. log-timescale curves (*b* = –0.09, *p* = 0.879) nor the λ_s^2 vs. log-
timescale curves the downward-facing parabolic form (*b* = 0.73, *p* = 0.244; Fig. 5C). Thus, the
λ_s^2 vs. log-timescale curves differ between the eye-closed condition and all the eyes-open
conditions except the eyes-fixated-at-50-cm condition. Furthermore, whereas the 50- and
305-cm viewing conditions show significant growth of the λ_s^2 vs. log-timescale curves, the 135-
and 220-cm viewing conditions exhibit significant decay of this curve with scale.

Linear and nonlinear indices of endogenous postural fluctuations known to capture
effects of aging and disease governed the timescale-dependence of λ_s^2.[13,26,29,34,35] All indices,
except sample entropy of postural fluctuations, showed predictive effects on λ_s^2. λ_s^2 decreased
with increases in root mean square, and increased with increases in mean, standard
deviation, and multiscale entropy of postural fluctuations (Table S2). Nonetheless, greater
'original-minus-shuffled/surrogate' differences in fractality and multifractality diminished λ_s^2.
Hence, evidence of multifractality with weak difference from the surrogates suggests heavy-
tailed distributions and thus non-Gaussianity.[16,22]

Interactions of these indices with linear and quadratic terms in the first model
significantly improved the model fit (χ_{315} = 153961, $p < 0.001$; Table S2). Most variability in the
quadratic form of the λ_s^2 vs. log-timescale curves depended on these indices, as all but two of
41 significant ($p < 0.05$) quadratic effects relate to interactions of these indices with log-
timescale. That is, the parabolic tempering of non-Gaussianity across the viewing conditions
was a function of endogenous postural control.

The λ_s^2 vs. log-timescale curves reflect differences in postural stability across the
different viewing conditions. Postural instability typically weakens the growth of lognormality in
postural fluctuations with log-timescale.[36,38] The original CoP PED series exhibited less
lognormality than the surrogates (b = –4.28, $p < 0.001$; Table S3; Fig. 6). Lognormality in
postural fluctuations grew with log-timescale; the growth of lognormality showed stronger
significant linear increase (b = 2219.40, p = 0.016), quadratic decrease (b = –1678.38, p =
0.052), and cubic increase (b = 559.57, p = 0.051) in the original CoP PED series than the
surrogates. However, polynomial profiles of the growth of lognormality with log-timescale did
not differ between the eyes-closed condition and any of the eyes-open conditions, suggesting
that the observed differences in the λ_s^2 vs. log-timescale curves reflect differences in the bulk
of the postural-fluctuations distributions rather than in the sparser tail.

224   **DISCUSSION**

Visual effort moderates postural cascade dynamics, given that standing still and
visually fixating at shorter and longer distances unleashed stronger cascade dynamics than
standing still (Hypothesis 3) and visually fixating at an optimal distance or standing still
without visually fixating (i.e., with eyes closed; Hypotheses 1 & 2). Importantly, these effects
are not reducible to the mere availability of visual information, as standing still and visually
fixating at an optimal distance and standing still with eyes closed showed similar, if not
identical cascade dynamics. Nor is this effect reducible to postural stability, as the growth of
lognormality in postural fluctuations with timescale did not depend on the viewing condition.
Instead, these effects reflect endogenous postural control, as known indices of endogenous
postural fluctuations moderated the effect of viewing condition on postural cascade dynamics.
In short, the moderating of postural cascade dynamics by visual effort is an endogenous
phenomenon that is difficult to counteract.

Past work has shown that variability in postural fluctuations is reduced when visually
fixating at near as opposed to distant targets,[20,43,50,51] an effect that has been directly attributed
to the precision demands of viewing distance.[29] The present finding significantly adds to this
work by showing that this effect is not linear. Visually fixating at too-close and too-far targets
may introduce ambiguity between sway and optic flow: too-close targets making head-sway
more likely to destabilize the posture, and too-far targets might recruit more torso-sway.
However, visually fixating at an optimally placed target may resolve this ambiguity as the body
courses with fluctuations ferrying information across the body. Furthermore, the stark
similarities in the cascade dynamics between standing still and visually fixating at an optimal
distance and standing still with eyes closed adds a novel dimension to the family of
perception-action couplings in postural control. Postural control does not depend on vision per
se. Instead, it depends on how visual information is laced together with activities at multiple
scales: subtle fluctuations of the head, torso, and the whole body and photoreceptor support
of the target's retinal image.

Postural cascade dynamics is rooted in the known physiology. Indeed, the biophysical
substrate supporting postural control is imagined as a neurally-tunable bodywide multifractal
tensegrity (MFT) network consisting of the connective-tissue net and extra-cellular matrix
(ECM) in which components hang together by tensional and compression forces at multiple
scales, from micro (e.g., cells and membranes) to macro (e.g., muscles and bones).[3,17–19,41,46,52]
This network cooperates directly and uninterruptedly with the nervous system and exhibits
similar cascade-like dynamics as in "neuronal avalanches."[7,39] It is no small coincidence that
tensegrity and avalanche models all converge on multiplicativity[11,54] that inspired multiscale
quantification of non-Gaussianity.[23,24,53] Hence, MFT approaches suggest multiple points of
entry through which stimulation elicits cascades to spread across scales—from the surface
underfoot[36] to the loads at hand[38] and now from the incoming visual information. As the
cascade metaphor entails, multiple scales are not merely coexisting but explicitly interacting,
with influences of one spreading to others. If mechanically and visually perturbing posture
engender cascade dynamics similarly,[36,38] then postural control may emerge from the
perceptuomotor sensitivity of specifically multiplicative fluctuations to the environment. The
methods introduced here would enable future investigations into how different suprapostural
tasks lace together deep texture at very many scales.

In short, the present findings suggest that visual perception by no means is a passive
function proceeding in parallel to still posture—the poise carried by the body as it stands and
looks suggests that moving visual targets nearer or farther brings a reorganization of postural
interactions across scales. This standing-and-looking preamble to action brims with dynamic
redistributions of tensional and compression forces. Indeed, the participants completed no
task beyond standing still and maintaining visual fixation. But these cascades affecting a
dynamic balance of forces across the body seem to be essential foundations for how the body
reaches out beyond itself to engage with the environment.[2,4,32,33] So, the present findings show
how vision plays not only upon the photoreceptors in the eyes but also across the full-body
postural system.

AUTHOR CONTRIBUTIONS

M.M. I.-C.L., K.M.N., and D.G.K-S. conceived and designed research; I.-C.L. performed
experiments; M.M. and D.G.K-S. analyzed data; M.M. and D.G.K-S. interpreted results of
experiments; M.M. prepared figures; M.M. and D.G.K-S. drafted manuscript; M.M., I.-C.L.,
K.M.N., and D.G.K-S. edited and revised manuscript; M.M., I.-C.L., K.M.N., and D.G.K-S.
approved final version of manuscript.

CONFLICT OF INTEREST

The authors declare that they have no competing interest.

REFERENCES

1.   Bates, D., M. Mächler, B. Bolker, and S. Walker. Fitting linear mixed-effects models
using lme4. *arXiv* 5823, 2014.doi:arXiv:1406.5823

2.   Bell, C., N. Carver, J. Zbaracki, and D. Kelty-Stephen. Nonlinear amplification of
variability through interaction across scales supports greater accuracy in manual
aiming: Evidence from a multifractal analysis with comparisons to linear surrogates in
the Fitts task. *Front. Physiol.* 10:998, 2019.

3.   Cabe, P. A. All perception engages the tensegrity-based haptic medium. *Ecol. Psychol.*
31:1–13, 2018.

4.   Carver, N. S., D. Bojovic, and D. G. Kelty-Stephen. Multifractal foundations of visually-
guided aiming and adaptation to prismatic perturbation. *Hum. Mov. Sci.* 55:61–72,
2017.

5.   Castaing, B., Y. Gagne, and E. J. Hopfinger. Velocity probability density functions of
high Reynolds number turbulence. *Phys. D Nonlinear Phenom.* 46:177–200, 1990.

6.   Chhabra, A., and R. V Jensen. Direct determination of the *f*(\alpha) singularity
spectrum. *Phys. Rev. Lett.* 62:1327–1330, 1989.

7.   Chialvo, D. R. Emergent complex neural dynamics. *Nat. Phys.* 6:744–750, 2010.

8. Codling, E. A., M. J. Plank, and S. Benhamou. Random walk models in biology. *J. R. Soc. Interface* 5:813–834, 2008.

9. Costa, M., A. L. Goldberger, and C.-K. Peng. Multiscale entropy analysis of complex physiologic time series. *Phys. Rev. Lett.* 89:68102, 2002.

10. Costa, M., A. L. Goldberger, and C.-K. Peng. Multiscale entropy analysis of biological signals. *Phys. Rev. E* 71:21906, 2005.

11. Das, A., and A. Levina. Critical neuronal models with relaxed timescale separation. *Phys. Rev. X* 9:21062, 2019.

12. Dault, M. C., A. C. H. Geurts, T. W. Mulder, and J. Duysens. Postural control and cognitive task performance in healthy participants while balancing on different support-surface configurations. *Gait Posture* 14:248–255, 2001.

13. Duarte, M., and V. M. Zatsiorsky. On the fractal properties of natural human standing. *Neurosci. Lett.* 283:173–176, 2000.

14. Fiorelli, C. M., P. F. Polastri, S. T. Rodrigues, A. M. Baptista, T. Penedo, V. A. I. Pereira, L. Simieli, and F. A. Barbieri. Gaze position interferes in body sway in young adults. *Neurosci. Lett.* 660:130–134, 2017.

15. Giveans, M. R., K. Yoshida, B. Bardy, M. Riley, and T. A. Stoffregen. Postural sway and the amplitude of horizontal eye movements. *Ecol. Psychol.* 23:247–266, 2011.

16. Hayano, J., K. Kiyono, Z. Struzik, Y. Yamamoto, E. Watanabe, P. Stein, L. Watkins, J. Blumenthal, and R. Carney. Increased non-Gaussianity of heart rate variability predicts cardiac mortality after an acute myocardial infarction. , 2011.

17. Ingber, D. E. Cellular mechanotransduction: Putting all the pieces together again. *FASEB J.* 20:811–827, 2006.

18. Ingber, D. E. Tensegrity-based mechanosensing from macro to micro. *Prog. Biophys. Mol. Biol.* 97:163–179, 2008.

19. Ingber, D. E. From cellular mechanotransduction to biologically inspired engineering. *Ann. Biomed. Eng.* 38:1148–1161, 2010.

20. Kapoula, Z., and T.-T. Lê. Effects of distance and gaze position on postural stability in young and old subjects. *Exp. Brain Res.* 173:438–445, 2006.

21. Kelty-Stephen, D. G., K. Palatinus, E. Saltzman, and J. A. Dixon. A tutorial on multifractality, cascades, and interactivity for empirical time series in ecological science. *Ecol. Psychol.* 25:1–62, 2013.

22. Kiyono, K., J. Hayano, S. Kwak, E. Watanabe, and Y. Yamamoto. Non-gaussianity of low frequency heart rate variability and sympathetic activation: Lack of increases in multiple system atrophy and parkinson disease. *Front. Physiol.* 3:34, 2012.

23.   Kiyono, K., J. Hayano, E. Watanabe, Z. R. Struzik, and Y. Yamamoto. Non-Gaussian
heart rate as an independent predictor of mortality in patients with chronic heart failure.
Hear. Rhythm 5:261–268, 2008.

24.   Kiyono, K., Z. R. Struzik, N. Aoyagi, S. Sakata, J. Hayano, and Y. Yamamoto. Critical
scale invariance in a healthy human heart rate. *Phys. Rev. Lett.* 93:178103, 2004.

25.   Kiyono, K., Z. R. Struzik, and Y. Yamamoto. Estimator of a non-Gaussian parameter in
multiplicative log-normal models. *Phys. Rev. E* 76:41113, 2007.

26.   Ko, J.-H., and K. M. Newell. Aging and the complexity of center of pressure in static and
dynamic postural tasks. *Neurosci. Lett.* 610:104–109, 2016.

27.   Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen. lmerTest package: Tests in
linear mixed effects models. *J. Stat. Softw.* 82:1548–7660, 2017.

28.   Lê, T.-T., and Z. Kapoula. Role of ocular convergence in the Romberg quotient. *Gait*
Posture 27:493–500, 2008.

29.   Lee, I.-C., M. M. Pacheco, and K. M. Newell. The precision demands of viewing
distance modulate postural coordination and control. *Hum. Mov. Sci.* 66:425–439, 2019.

30.   Lee, W., and K. J. Grimm. Generalized linear mixed-effects modeling programs in R for
binary outcomes. *Struct. Equ. Model. A Multidiscip. J.* 25:824–828, 2018.

31.   Lomholt, M. A., K. Tal, R. Metzler, and K. Joseph. Lévy strategies in intermittent search
processes are advantageous. *Proc. Natl. Acad. Sci.* 105:11055–11059, 2008.

32.   Mangalam, M., N. S. Carver, and D. G. Kelty-Stephen. Global broadcasting of local
fractal fluctuations in a bodywide distributed system supports perception via effortful
touch. *Chaos, Solitons & Fractals* 135:109740, 2020.

33.   Mangalam, M., N. S. Carver, and D. G. Kelty-Stephen. Multifractal signatures of
perceptual processing on anatomical sleeves of the human body. *J. R. Soc. Interface*
17:20200328, 2020.

34.   Mangalam, M., R. Chen, T. R. McHugh, T. Singh, and D. G. Kelty-Stephen. Bodywide
fluctuations support manual exploration: Fractal fluctuations in posture predict
perception of heaviness and length via effortful touch by the hand. *Hum. Mov. Sci.*
69:102543, 2020.

35.   Mangalam, M., and D. G. Kelty-Stephen. Multiplicative-cascade dynamics supports
whole-body coordination for perception via effortful touch. *Hum. Mov. Sci.* 70:102595,
2020.

36.   Mangalam, M., and D. G. Kelty-Stephen. Hypothetical control of postural sway. *bioRxiv*
104760, 2020.doi:10.1101/2020.05.19.104760

37. Manor, B., M. D. Costa, K. Hu, E. Newton, O. Starobinets, H. G. Kang, C. K. Peng, V.
Novak, and L. A. Lipsitz. Physiological complexity and system adaptability: Evidence
from postural control dynamics of older adults. *J. Appl. Physiol.* 109:1786–1791, 2010.

38. Mariusz, P. F., M. Mangalam, D. G. Kelty-Stephen, and G. Juras. Postural instability
recruits shorter-timescale processes into the non-Gaussian cascade processes.
bioRxiv 136895, 2020.doi:10.1101/2020.06.05.136895

39. Miller, S. R., S. Yu, and D. Plenz. The scale-invariant, temporal profile of neuronal
avalanches in relation to cortical γ–oscillations. *Sci. Rep.* 9:16403, 2019.

40. Musolino, M. C., P. J. Loughlin, P. J. Sparto, and M. S. Redfern. Spectrally similar
periodic and non-periodic optic flows evoke different postural sway responses. *Gait
Posture* 23:180–188, 2006.

41. Nelson, C. M., R. P. Jean, J. L. Tan, W. F. Liu, N. J. Sniadecki, A. A. Spector, and C. S.
Chen. Emergent patterns of growth controlled by multicellular form and mechanics.
Proc. Natl. Acad. Sci. U. S. A. 102:11594–11599, 2005.

42. Oullier, O., B. G. Bardy, T. A. Stoffregen, and R. J. Bootsma. Postural coordination in
looking and tracking tasks. *Hum. Mov. Sci.* 21:147–167, 2002.

43. Paulus, W., A. Straube, S. Krafczyk, and T. Brandt. Differential effects of retinal target
displacement, changing size and changing disparity in the control of anterior/posterior
and lateral body sway. *Exp. Brain Res.* 78:243–252, 1989.

44. Peng, C.-K., S. V Buldyrev, S. Havlin, M. Simons, H. E. Stanley, and A. L. Goldberger.
Mosaic organization of DNA nucleotides. *Phys. Rev. E* 49:1685–1689, 1994.

45. Richman, J. S., and J. R. Moorman. Physiological time-series analysis using
approximate entropy and sample entropy. *Am. J. Physiol. Circ. Physiol.* 278:H2039–
H2049, 2000.

46. Schleip, R., F. Mechsner, A. Zorn, and W. Klingler. The bodywide fascial network as a
sensory organ for haptic perception. *J. Mot. Behav.* 46:191–193, 2014.

47. Schreiber, T., and A. Schmitz. Improved surrogate data for nonlinearity tests. *Phys. Rev.
Lett.* 77:635–638, 1996.

48. Stoffregen, T. A., B. G. Bardy, C. T. Bonnet, and R. J. Pagulayan. Postural stabilization
of visually guided eye movements. *Ecol. Psychol.* 18:191–222, 2006.

49. Stoffregen, T. A., P. Hove, J. Schmit, and B. G. Bardy. Voluntary and involuntary
postural responses to imposed optic flow. *Motor Control* 10:24–33, 2006.

50. Stoffregen, T. A., R. J. Pagulayan, B. G. Bardy, and L. J. Hettinger. Modulating postural
control to facilitate visual performance. *Hum. Mov. Sci.* 19:203–220, 2000.

51.   Stoffregen, T. A., L. J. Smart, B. G. Bardy, and R. J. Pagulayan. Postural stabilization of
418   looking. *J. Exp. Psychol. Hum. Percept. Perform.* 25:1641–1658, 1999.

419   52.   Turvey, M. T., and S. T. Fonseca. The medium of haptic perception: A tensegrity
420   hypothesis. *J. Mot. Behav.* 46:143–187, 2014.

421   53.   West, B. J., and M. Turalska. Hypothetical control of heart rate variability. *Front.*
422   *Physiol.* 10:1078, 2019.

423   54.   Zorick, T., J. Landers, A. Leuchter, and M. A. Mandelkern. EEG multifractal analysis
424   correlates with cognitive testing scores and clinical staging in mild cognitive impairment. *J.*
425   *Clin. Neurosci.* 76:195–200, 2020.

FIGURE 1. Schematic of the task and effects of eye-to-target distance on postural sway. (**A**) The suprapostural viewing task of standing quietly with the eyes fixated at a distant visual element. (**B** and **C**) Visual angle gain for short vs. long eye-to-target distances along the anterior-posterior (AP) and medial-lateral (ML) axes. (**D** and **E**) Visual angle gain as a function of eye-to-target distance for different sway magnitudes. Closer targets increase *AP* sway, whereas farther targets increase *ML* sway.

Fig. 2. Schematic illustration of multiscale probability density function (PDF) analysis. (**A**) The first step is to generate $\{B(t)\}$ by integrating $\{b(t)\}$ after centering by the mean b_{ave}: $B(t)=\sum_{i=1}^{t/\Delta t} b(i\,\Delta t)-b_{ave}$. (**B**) A a 3rd order polynomial detrends $\{B(t)\}$ within k overlapping windows of length $2s$, s being the timescale. (**C**) Intermittent deviation $\Delta_s B(t)$ in k^{th} window from $1+s(k-1)$ to sk in the detrended time series $\{B^d(t)=B(t)-f_{fit}(t)\}$ is computed as $\Delta_s B^d(t)=B^d(t+s)-B^d(t)$, where $1+s(k-1)\le t\le sk$ and $f_{fit}(t)$ is the polynomial representing the local trend of $\{B(t)\}$, of which the elimination assures the zero-mean probability density function in the next step.

Fig. 3. Schematic illustration of non-Gaussianity index λ_s. (**A**) The relationship between λ_s^2 and shapes of PDF plotted in linear-log coordinates for λ_s^2 = 0.9, 0.33, and 0.13 (from top to bottom). For clarity, the PDFs have been shifted vertically. As λ_s^2 increases, the PDF becomes increasingly peaked and fat-tailed. As λ_s decreases, the PDF increasingly resembles the Gaussian (dashed line), assuming a perfect Gaussian as λ_s^2 approaches 0. (**B**) Cascade-type multiplicative processes yield the inverse relationship $\lambda_s^2 \sim -\log s$.

IGURE 4. Multiscale PDF characterization of postural fluctuations in a representative
participant maintaining quiet stance for 120 s with eyes closed, and eyes fixated at a point at
a distance of 25, 50, 135, 220, and 305 cm from the eyes. From to bottom: CoP trajectories
along the anterior-posterior (AP) and medial-lateral (ML) axes. CoP PED series. $\{\Delta_s B(i)\}$ for s
= 0.5, 2, and 8 s. Standardized PDFs (in logarithmic scale) of $\{\Delta_s B(i)\}$ for s = 0.5, 2, and 8 s
(from top to bottom), where σ_s denotes the SD of $\{\Delta_s B(i)\}$ for s = 0.5, 2, and 8 s (from top to
bottom), where σ_s denotes the SD of $\{\Delta_s B(i)\}$.

FIGURE 5. Log-timescale dependence of the non-Gaussianity index λ_s^2. Mean values of λ_s^2 for the participants standing quietly for 120 s in different viewing conditions. (**A**) Eyes closed. (**B**) Eyes fixated at 25 cm. (**C**) Eyes fixated at 50 cm. (**D**) Eyes fixated at 135 cm. (**E**) Eyes fixated at 220 cm. (**F**) Eyes fixated at 305 cm. Vertical bars indicate ±1 *s.e.m.* (*n* = 15). Notice that the λ_s^2 vs. log-timescale curves differ between the eye-closed condition all the eyes-open conditions except the eyes-fixated-at-50-cm condition.

FIGURE 6. Log-timescale dependence of the mean proportion of trials with a lognormal distribution for the participants standing quietly for 120 s in different viewing conditions. (**A**) Eyes closed. (**B**) Eyes fixated at 25 cm. (**C**) Eyes fixated at 50 cm. (**D**) Eyes fixated at 135 cm. (**E**) Eyes fixated at 220 cm. (**F**) Eyes fixated at 305 cm. Vertical bars indicate ±1*s.e.m.* (*n* = 15). Notice that the polynomial profiles of the growth of lognormality with log-timescale do not differ between the eyes-closed condition and any of the eyes-open conditions.