Multifractal roots of suprapostural dexterity

Damian G. Kelty-Stephen[a,*], I-Chieh Lee[b], Nicole S. Carver[c], Karl M. Newell[d], and Madhur Mangalam[e,*]

[a]*Department of Psychology, Grinnell College, Grinnell, IA 50112, USA*

[b]*UNC-NC State Joint Department of Biomedical Engineering, UNC-Chapel Hill, Chapel Hill, NC 27514, USA*

[c]Department of Psychology, University of Cincinnati, Cincinnati, Ohio, United States of America

[d]Department of Kinesiology, University of Georgia, Athens, GA 30602, USA

[e]Department of Physical Therapy, Movement and Rehabilitation Sciences, Northeastern University, Boston, MA 02115, USA

ORCIDs:

Damian G. Kelty-Stephen (0000-0001-7332-8486)

I-Chieh Lee (0000-0001-7094-6270)

Nicole S. Carver (0000-0002-1107-0595)

Karl Newell (0000-0002-2931-4244)

Madhur Mangalam (0000-0001-6369-0414)

*Corresponding authors: keltysda@grinnell.edu (D. G. Kelty-Stephen); m.mangalam@northeastern.edu (M. Mangalam).

Data accessibility. All data analyzed in the present study are available upon request.

Author contributions: D.G.K-S., I.-C.L., K.M.N., and M.M. conceived and designed research; I.-C.L. performed experiments; D.G.K-S., N.S.C., and M.M. analyzed data; D.G.K-S. and M.M. interpreted results of experiments; D.G.K-S. and M.M. prepared figures; D.G.K-S. and M.M. drafted manuscript; D.G.K-S. and M.M. edited and revised manuscript; D.G.K-S., I.-C.L., N.S.C., K.M.N., and M.M. approved final version of manuscript.

Competing interests. The authors have no competing interests to declare.

Funding. No funding was received for this study.

Electronic supplementary material

Dataset S1. Data used for mixed-effect regression modeling of impulse-response functions relating CoM-H_{fGn},
CoP-SD, and CoP-H_{fGn}

Dataset S2. Data used for mixed-effect regression modeling of impulse-response functions incorporating CoP-
t_{MF} along with the prior set of CoM-H_{fGn}, CoP-SD, and CoP-H_{fGn}

Table S1. Full output for mixed-effect regression modeling of impulse-response functions relating CoM-H_{fGn},
CoP-SD, and CoP-H_{fGn}

Table S2. Full output of mixed-effect regression modeling of impulse-response functions incorporating CoP-t_{MF}
along with the prior set of CoM-H_{fGn}, CoP-SD, and CoP-H_{fGn}

Abstract

Visually guided postural control emerges in response to task constraints. Task constraints generate physiological fluctuations that foster the exploration of available sensory information at many scales. Temporally correlated fluctuations quantified using fractal and multifractal metrics have been shown to carry perceptual information across the body. The risk of temporally correlated fluctuations is that stable sway appears to depend on a healthy balance of standard deviation (*SD*): too much or too little *SD* entails destabilization of posture. This study presses on the visual guidance of posture by prompting participants to quietly stand and fixate at distances within, less than, and beyond comfortable viewing distance. Manipulations of the visual precision demands associated with fixating nearer and farther than comfortable viewing distance reveals an adaptive relationship between *SD* and temporal correlations in postural fluctuations. Changing the viewing distance of the fixation target shows that increases in temporal correlations and *SD* predict subsequent reductions in each other. These findings indicate that the balance of *SD* within stable bounds may depend on a tendency for temporal correlations to self-correct across time. Notably, these relationships became stronger with greater distance from the most comfortable viewing and reaching distance, suggesting that this self-correcting relationship allows the visual layout to press the postural system into a poise for engaging with objects and events. Incorporating multifractal analysis showed that all effects attributable to monofractal evidence were better attributed to multifractal evidence of nonlinear interactions across scales. These results offer a glimpse of how current nonlinear dynamical models of self-correction may play out in biological goal-oriented behavior. We interpret these findings as part of the growing evidence that multifractal nonlinearity is a modeling strategy that resonates strongly with ecological-psychological approaches to perception and action.

Keywords: biotensegrity, center of mass, center of pressure, fractal, Hurst exponent, postural sway

1. Introduction

1.1. Stability of suprapostural visual activities at longer scales rests on fluctuating behavior at shorter scales

Standing quietly and focusing on a target in front of us is the preamble to various coordinated behaviors —we lean forward and reach or track the target's progress and bat it away. Less recognized is the fact that this starting position is not merely the preamble to action but is already a rich wellspring of the action itself, exhibiting a continuous stream of intermittent fluctuations. These fluctuations in our bodily center of mass (CoM) and center of pressure (CoP, where ground reaction forces meet our lower extremities) are crucial to maintaining a quiet stance unless they pitch the CoP beyond the base of support (Winter, 1995; Zatsiorsky & Duarte, 1999). These fluctuations offer the body a subtle and flexible command of the mechanical surface underfoot (Chen et al., 2008; Rajachandrakumar et al., 2018), exemplifying long-respected proposals that what looks 'noisy' can actually stabilize nonlinear-dynamical systems (Shinbrot and Muzzio, 2001; Turing, 1952). Critically, what looks noisy may instead be richly structured movement variability (Newell and Corcos, 1993; Riccio, 1993; Slifkin and Newell, 1999, 1998; Stergiou et al., 2006)—with richness entailing high-dimensional nonlinear dynamics (Lovejoy, 2019; Schertzer and Lovejoy, 2004), enabling movement systems to sustain their dexterous contact with task context through multifractal foundations (Cavanaugh et al., 2017). This work examines how the visual precision demands reshape the pattern of interactions amongst fractal and multifractal fluctuations in postural control.

We begin with Section 1.1. to define some of the technical terms ahead. We take up these terms beginning in Section 1.2. to develop our rationale motivating this empirical work.

1.1. Preliminary definitions of terms for describing movement variability: monofractality, multifractality, and flow of mono- or multifractal fluctuations

At the outset, we wish to define these terms so that any new readers can understand what is at stake in this portrait of movement variability. Briefly, 'fractal fluctuations' refer to time series with temporally-correlated sequencing, meaning that each current value is relatively more predictable from previous values than one might find in a completely random sequence (e.g., in additive white Gaussian noise). Additive white Gaussian noise is classically 'uncorrelated,' implying no statistical relationship across time between its consecutive values. Specifically, fractal fluctuations entail a temporal correlation in which the variance of current values is strongly correlated with variance of recent values and in which the correlation between the present and past drops off very slowly. Fractal fluctuations present an intriguing case of a non-zero, non-vanishing correlation between current variance and variance in the most distance past. Hence, fractal fluctuations reflect a case of 'long memory' in a time series. The clearest signature of this long memory is when the *SD* of a measured time series grows relatively fast. For white noise series without long memory, *SD* grows according to the square root of time (i.e., time with exponent 0.5). For fractally-fluctuating series, *SD* grows linearly with time (time raised to exponent 1). The exponent on time will appear in this work as a 'Hurst exponent' H abbreviated as 'H_{fGn},' because the first step of fractal analysis is to assume that fractal (*f*) fluctuations appear as Gaussian noises (Gn). Hence, monofractal evidence diagnoses 'fractal Gaussian noise' (fGn) in terms of values of H_{fGn} deviating away from 0.5 and closer to 1.

Movement variability often exhibits fractal fluctuations in response to monofractal analyses that test for only one fractal exponent H_{fGn} at a time. However, with multiple monofractal analyses, movement variability exhibits multiple H_{fGn}. So obviously, movement variability is not 'monofractal.' Using multiple monofractal analyses reveals the texture hidden in the very many degrees of fractal patterning across space and time in the movement system. Hence, movement variability is 'multifractal,' which can be analyzed as the temporal or spatial variation of monofractal evidence.

A growing body of evidence suggests that movement variability exhibits a flow of fractal and multifractal fluctuations from one anatomical location or form into another (e.g., Mangalam, Carver, et al., 2020a, 2020b; Stephen et al., 2012). That is, a single case of monofractal and multifractal fluctuations in one part of the body is just the tip of the metaphorical iceberg. This 'flow' indicates that we can statistically test how increases in H_{fGn} at one location or one point in time predicts subsequent changes in H_{fGn} in the same or a different location. We can equally well statistically test how increases in multifractal variety predict subsequent changes in multifractal variety. This work will open up a field of inquiry where we can test for relationships

109 between monofractal estimates (H_{fGn}), multifractal estimates, and *SD* of postural fluctuations—each of these
110 measures of movement variability can change, and those changes can predict subsequent changes in each other
111 measure of movement variability. The present work will examine how task constraints of quietly standing and
112 visually fixating at a distance prompt different relationships amongst these aspects of postural control.

1.2. Stability of suprapostural visual activities at longer scales rests on fluctuating behavior at shorter scales

Eyes lace the postural system into the visual field, locking focus on a distant point in the world; this lock
is fluid, nonetheless. Fixations are regularly recognized as a class of movement. Just as upright posture is never
entirely still, the 'fixations' of the eyes are not stasis (Montesano et al., 2018). Fixations consist of a deep,
fluctuating foundation of smaller movements called 'microsaccades' (Meyberg et al., 2017). These
microsaccades stabilize images that would otherwise fade on a static retina (Martinez-Conde et al., 2013), thus
playing a similar role to the visual system that fluctuations play for the postural system. Indeed, the exploratory
role of fluctuating movements extends to the bodily extremities besides and intermediating between upright
posture and vision (Murnaghan et al., 2016), suggesting a vital role for postural sway in supporting visual
perception (Stoffregen, 2010; Stoffregen et al., 2000, 1999). Hence, even when quietly standing and fixating at a
distant target, the body is coursing with fluctuations ferrying information across the body.

Visually engaging with the world contributes to the fray of ongoing bodywide work to stay on task and
dexterously responsive to context. Fixations require precision like all other movements, stabilizing our position
in the world. By 'visual precision demands' we do not mean the laboring of visual-cognitive mechanisms (Wang
et al., 2013), but instead the body-wide muscular work maintaining fixations straight ahead at a static stimulus.
This term can be undressed of possible confusion with the visual-cognitive workload by substitution with 'ocular
precision demands,' which does not necessarily denote maintaining fixation to single body part. Indeed, no less
than the postural system seeking stability amidst wide-ranging constraints, the precision needed to maintain
fixation is the work of the entire body positioned amongst task constraints like ambient light and surface
(Cachinero-Torre et al., 2017; Kaldy and Blaser, 2020).

Postural fluctuations produce translations and rotations of visible surfaces specific to the objects' spatial
relationships in the visual field. These movement-induced translations and rotations compose an optic flow that
provides visual support for the subsequent movement. Thus, subtle sway offers the sighted organism a rich
source of information about objects' layout in the visual field. The visual layout itself then presses the postural
system into a poise for engaging with objects and events. Visual targets appearing closer to or farther from the
looking postural system impose retina-specific oculomotor convergence constraints (Jaschinski-Kruza, 1991;
Jaschinski, 2002) and chromatic aberration (Párraga et al., 2002). Consequently, changes in a target's size
relative to other aspects of the visual field entail optic flow (Fig. 1). These multiscale factors affect postural
stability, reflecting changes in postural configurations and the flow of information needed to organize these
postures. Although how these multiscale factors enact their effects remain controversial, observations of across-
scale interactions in sway align with the proposal that the coordination of posture is a fluid, cascading process in
which relatively coarse, longer-scale constraints (e.g., the task) redirect progressively finer-grained motoric
degrees of freedom without stipulating them (Latash, 2020, 2012). This cascade operates through the loose
nesting of finer motoric details within larger motoric degrees of freedom (e.g., motor neurons within bundles of
muscle fiber, muscles within joint-centered synergies, and so on) that allow dexterous behavior to adapt flexibly
to constraints changing on the way to task completion.

1.3. Visual precision demands on postural stability could reshape intrapostural interactivity

Movement depends on the bodywide network of connective and nervous tissues, forming flexible
relationships that balance tensions with compressions at multiple scales (Cabe, 2018; Ingber, 2006, 2008, 2010;
Nelson et al., 2005; Schleip et al., 2014; Turvey & Fonseca, 2014). This organization entails a multifractal
geometry that embodies multiple scale-invariant patterns of behavior (e.g., microsaccades within the saccades
that intersperse larger saccades by the eye and turns by the head) across time and space. CoP multifractality has
repeatedly borne a consistent relationship to perceptual judgments of visual and haptic stimuli (Doyon et al.,
2019; Hajnal et al., 2018; Mangalam et al., 2020c; Mangalam and Kelty-Stephen, 2020; Palatinus et al., 2014,
2013). Research on perceptual tasks (e.g., manually wielding an object to judge its heaviness or length has
shown that a bodywide flow of multifractal fluctuations precedes and shapes the verbal articulation of perceptual

judgments (Mangalam et al., 2020a, 2020b). Hence, the flow of multifractality within posture supports
information flow and might provide a glimpse of the control policy emerging from the bodily situation in task
constraints.

Indeed, the intermodal perception supporting standing and fixating makes an inquiry into multifractal
roots particularly valuable. Multifractal geometry appears not simply important for wholesale pickup of
information but is specifically predictive of how different tissues of the body coordinate different aspects of
information about the task context (Carver et al., 2017; Doyon et al., 2019; Hajnal et al., 2018; Kelty-Stephen
and Dixon, 2014). Multifractal aspects of sway also affect the coordination of mechanical and visual information
as people stand still while visually fixating. Changing the movement system's relationship to visual information
(e.g., increasing precision demands or closing eyes) while standing on the same mechanical surface could
prompt the body to arrive at new body-wide coordination. We expect that changing this coordination of visual
and mechanical information would depend on multifractal foundations. Hence, this work investigates fractal
evidence that would speak to ongoing deliberations about clinical applications to supporting upright stance (e.g.,
Hove et al., 2012; Rodger & Craig, 2016; Uchitomi et al., 2013). It also connects fractal results and discourse to
multifractal processes underlying fractal evidence and supporting the more generic intermodal coordination.

1.4. Visual precision demands for fixating across different viewing distances could reveal patterns of intrapostural interactivity

1.4.1. The task

This work is a reanalysis of a previous study that manipulated viewing distance and measured CoM and
CoP (Lee et al., 2019). Healthy adults stood and maintained a quiet stance for 120 s under six different
conditions: a control condition with eyes closed and five conditions with eyes open and fixating on a red laser
point projected on surfaces at 20, 50, 135, 220, and 305 cm distance. These five non-equally progressively
spaced distances captured the power-law like relationship between viewing distance and resulting angular
deviation under the assumption of 2-cm postural sway in either *AP* or *ML* direction (Figs. 1D and 1E). The
authors referred to this nonlinear relationship between viewing distance as 'visual angle gain' and used it to
express the range of optic flow expectable from standing as still as possible. The task follows in a long,
productive tradition of examining how visual perception of near or far targets influences posture (Bles et al.,
1980; Lee & Lishman, 1975; Mayo et al., 2010; Munafo, Curry, et al., 2016; Munafo, Wade, et al., 2016;
Stoffregen et al., 1999, 2000).

The original data collection allowed Lee et al. (2019) to examine the frequency of intermittent postural
corrections observed through increased bursts of posture–head movement coordination. Critically for our
reanalysis, this task and dataset offer the opportunity to examine the novel issue of intrapostural interactivity.
Indeed, postural sway can produce unintended movements that nonetheless serve the perceptual goals, and as
Lee et al. (2019) themselves demonstrated, the task-oriented aspect of these movements may manifest in any of
many different statistical measures of postural sway (Munafo et al., 2016b, 2016a; Palatinus et al., 2014, 2013).
We can undoubtedly examine concurrent correlations amongst these different statistical measures, but in this
work, our curiosity is specifically about the exchange of fluctuations as they change with the visual precision
demands. To our knowledge, the long tradition of studying postural sway in visual tasks does not yet include
causal modeling of movement variability to determine which fluctuations precede and potentially influence
subsequent changes in other fluctuations.

For our purpose, the two explicit precision demands from Lee et al.'s (2019) allowed an elegant
experimental clarity. Asking participants to fixate their gaze and minimize postural sway nearly froze not one but
two linkages to the task context, leaving the participant's movement system taut between static visual and static
mechanical supports. This manipulation reflects the ecological fact that sway is never due to movement in the
visual or mechanical stimulation, and changes in postural sway are most clearly due to the ongoing cooperations
and competitions amongst postural degrees of freedom. Hence, the double-precision command offers a unique
opportunity to delve into intrapostural interactivity. The manipulation of viewing distance is effectively a
manipulation of the visual precision demands. The 50-cm distance is within an ideal range to focus on red light
(Párraga et al., 2002) and prompts the least straining oculomotor convergence (Jaschinski-Kruza, 1991;
Jaschinski, 2002). We predicted that increasing precision demand by requiring viewing distances farther from

50-cm viewing distance would accentuate relationships between sway *SD* and fractality (i.e., *SD* and fractality of
postural fluctuations).

1.4.2. Modeling strategy

To accomplish our goal, we reanalyzed data from Lee et al. (2019b) in three steps: (i) carving individual
trials under each viewing condition into non-overlapping sub-trial segments, (ii) estimating *SD* and fractal
temporal structure within each segment, and (iii) using vector autoregression to test the interactions among these
descriptors across segments within each trial.

SD has been an intuitive way to model stability, but as noted above, it has only limited heuristic value as
an inverse measure of stability. Fractal and multifractal indices have drawn attention in postural-control research
only as a point of contact with nonlinear dynamics, cascades, and self-organized criticality. The fractal temporal
structure of bodywide fluctuations has been implicated in predicting variability in perceptual judgments or
responses capable of deviating from instructed or intended accuracy/veridicality. Whatever value *SD* serves for
postural control goals, we can take *SD* as an analogous measure of perceptual outcome: sway *SD* serves as a
measure of how well participants were able to follow the instructions to minimize their sway. Minimizing sway
is by no means the same as an explicit perceptual judgment, but like a perceptual response to instructions, it can
be done with varying degrees of accuracy.

Our choice of descriptors is not entirely divorced from that of Lee et al.'s (2019) choice of modeling
both *SD* and correlation dimension. The correlation dimension is a fractal dimension, but instead of emphasizing
temporal sequence, it emphasizes spatial recurrence of a phase-space trajectory. For instance, the power-law
scaling of %recurrence with a radius in recurrence quantification analysis estimates the correlation dimension.
Thus, our choice of dependent variables is not entirely novel, but it differs to align this work with other postural
and non-postural cases in which fractal fluctuations support perceptual responses.

1.4.2.1. Why we are modeling multifractality as temporal variation in fractal analysis, instead of estimating only a multifractal spectrum

Our choice to model the relationship between *SD* and fractality sits within our foregoing interest in
applying the multifractal formalism to perception and action. In the context of discussing multifractality, the
term 'fractal' does entail 'monofractality,' and the prefix 'mono-' does mean 'not multi-.' The use of
monofractality after just having trumpeted multifractality may seem jarring to seasoned readers. However, our
choice here has three points of motivations.

First, to our knowledge, all current empirically supported claims implicating multifractality as an
important factor in perception and action began with or were inspired by monofractal analysis as a preliminary
step (Mangalam et al., 2020c, 2020a; Palatinus et al., 2013). So, because we ask a new question about postural
control while fixating at a distant target and not merely aiming to examine multiple dependent measures about
postural sway separately, we hew to the tradition of modeling the predictive effects of variation in
monofractality. It is worth noting that, given the long shadow cast by information-processing accounts offering
more familiar, more intuitive models of goal-directed behavior, many readers and stakeholders in this literature
are still coming to grips with monofractality. Leaping straight to multifractal analysis is a step of complexity that
might well bury any usefulness of this work.

'Stochastic resonance' is an essential example of monofractal roots of postural control. The clinical
application of 'stochastic resonance' technologies aim to enhance movement dexterity using mechanical or
acoustic stimulations patterned according to white-noise or 1/f noise. Both are defined as the lack or presence of
monofractality. We are excited to see that research into this application of temporally-structured stimulation has
recently begun to look beyond lack-vs.-presence of monofractality and, in the vein of recent theorizing (e.g.,
Cavanaugh et al., 2017; Stergiou et al., 2006), to recognize that the fluctuations supporting the movement are not
'noise' at all (Hove et al., 2012; Rodger and Craig, 2016; Uchitomi et al., 2013). Indeed, our modeling strategy
resonates with these calls, suggesting that we may need to give voice to postural variations in monofractal
estimates rather than a simple lack-vs.-presence dichotomy. Nonetheless, the fact remains that monofractality
remains foundational to research interest in postural fluctuations. Interest in multifractality strikes us as the need
for the judicious situation of monofractal analysis sooner than a reason to eschew monofractal analysis.

Second, variation in monofractality is itself multifractality. Temporal or trial-by-trial variation of monofractal evidence is one of the most accessible ways in which to observe and communicate multifractality (Ihlen, 2012). Multifractality can be estimated as a spectrum of fractal-scaling dimensions, and preceding research on postural control has examined multifractality through this very multifractal spectrum (Koslucher et al., 2016; Munafo et al., 2016a). However, despite this constructive precedent, we still find value in examining multifractality as the variations of monofractal evidence for a third reason.

Finally, although the multifractal spectrum results from variations in monofractal evidence across time, it is not informative about the exact sequence of those variations. Variations of monofractality deliver two important insights that a single multifractal spectrum aggregating those variations could not. The first is a direct view of the multifractal spectrum width, albeit spread out over time. The second is a direct view of how the increases and reductions of monofractality could predict subsequent changes in *SD*. Similar-sized changes in monofractality in opposite directions would give us the same multifractal spectrum. The direction of change in monofractality could be predictive of subsequent *SD* that multifractal spectrum width itself could not predictive. If that were the case, then leaping to multifractal spectrum at the cost of examining variation in monofractal evidence would risk a false-negative result.

1.4.2.2. Why we are modeling multifractal-spectrum width as well to test whether nonlinear interactions across scales support the subsequent changes in SD and monofractal variation

Far from suggesting that estimating the multifractal spectrum is by itself a poor choice, this work uses it to quantify the strength of nonlinear interactions across scales. The strength of nonlinear interactions across scales can be quantified by comparing a measured series' multifractal spectrum width to the multifractal spectrum widths of surrogate series, preserving the linear structure of the original measurement but destroying the originally measured sequence. Indeed, the *t*-statistic comparing original multifractality to surrogate multifractality (henceforth, 'multifractal nonlinearity' t_{MF}) appears to be negatively correlated with concurrent *SD* in precision tasks and in postural sway (Bell et al., 2019; Jacobson et al., 2020; Kelty-Stephen, 2018).

Although we can estimate monofractality and multifractality separately, the control of goal-directed behavior reflects a balance between the two. The multifractal spectrum width reflects the flexible capacity for the body to embody multiple scale-invariant patterning, but this width itself only speaks to variety and not to exact direction of relationships across time. We expect bidirectional relationships between monofractal evidence and multifractal nonlinearity. (Note: we refer here to 'monofractal evidence' to reflect the fact that only conducting monofractal analyses can generate monofractal evidence without any indication that the analyzed system is itself monofractal and not multifractal). Changes in monofractal evidence do entail changes in multifractal structure (Ihlen, 2012), and thus changes in that monofractal evidence should predict subsequent changes in ongoing multifractal nonlinearity. Multifractality gains its causal force in shaping goal-directed behaviors not simply through the diversity of monofractality but also from the fact that the individual monofractality specify a degree of correlation. Monofractal temporal correlations have a valence to them, with greater or weaker correlations indicating the longer or shorter propagation of events across time scales (Kuznetsov and Wallot, 2011). Indeed, we might conceive multifractality as a reservoir of flexible possibility for nonlinear interactions across timescales, but temporal correlations are an important, supportive constraint in guiding multifractal behavior (Cavanaugh et al., 2017; Wallot et al., 2015).

At the outset (Section 1.2.), we noted that multifractal fluctuations bore a specific relevance to predicting how movement systems coordinate intermodal perception as in standing and visually fixating. We expect that multifractal nonlinearity is the foundation supporting monofractal variation and supporting any relationship between monofractal variation and *SD* of sway. Hence, we will aim in this study to, first, establish that monofractality might hover around a set point as posture ranges between the center and the margin of the support surface and, second, that increases in monofractality might precede subsequent reductions in *SD*. However, the concluding goal of this work is to demonstrate that multifractal nonlinearity is a crucial factor in both these features of monofractality in postural control.

To clarify, and to foreshadow our hypotheses, we expect that multifractal nonlinearity is the crucial guarantee of the capacity for monofractal variation. And we further expect that any effects of monofractality on *SD* of postural sway depend on nonlinear interactions across scales. That is, despite their being conceptually

308 separate, multifractal nonlinearity and monofractality hang together statistically. So, we expect that multifractal
309 nonlinearity will predict subsequent reductions in *SD*. However, we also predict that any evidence of
310 monofractality predicting subsequent reductions in *SD* may be better explained by multifractal nonlinearity.

1.4.2.3. *What multifractal nonlinearity could mean for intrapostural interactivity supporting the suprapostural task of quietly standing under various visual precision demands*

Although variation in monofractal evidence indicates interactions across scales that support dexterity,
multifractal nonlinearity is a stronger operationalization of these interactions (Kelty-Stephen and Wallot, 2017).
Hence, we propose that multifractal nonlinearity is more central than monofractal evidence to the task- or
context-dependence of goal-directed actions. This task-dependence appears here as visual precision demands.
We introduced above the idea that visual precision demands would moderate the association between prior
monofractal evidence and subsequent reduction in *SD* of postural fluctuations. Certainly, in the absence of
estimates of multifractal nonlinearity, monofractal evidence will be a strong first pass at modeling organismal
sensitivity to task constraints. However, when we introduce multifractal nonlinearity into the model, we expect
this sensitivity to task constraints much more in multifractal nonlinearity than in monofractal evidence.

We expected that incorporating multifractal nonlinearity into the vector-autoregressive relationships
would radically reshape our view of the intrapostural interactivity supporting postural control. First, because
multifractality is itself the generalization of monofractal structure, we expected it to show more refined
sensitivity to task constraints. For instance, whereas we predicted that the effects of monofractal evidence on
subsequent *SD* would be sensitive to visual precision demands, we expected that multifractal nonlinearity would
better predict these effects of visual precision demand and, further, predict effects of straightforward availability
of visual information (i.e., eyes-open vs. eyes-closed) on subsequent *SD*.

Multifractality increases as the variety of monofractal patterns in movement exceeds the monofractal
variety in linear models of that movement. Hence, multifractal nonlinearity is an asset in tasks that require
ongoing exploration (Booth et al., 2018; Carver et al., 2017) but a liability in tasks that require freezing out
degrees of freedom (Bell et al., 2019; Jacobson et al., 2020; Kelty-Stephen, 2018). We expected multifractal
nonlinearity to show both these faces in this work: as a stabilizing factor with increasing visual precision
demands and destabilizing factor under abrupt changes in the informational support for quiet stance. Specifically,
visual precision demands should accentuate an *SD*-reducing effect of multifractal nonlinearity—increasing the
visual precision demands might prompt a reduction of variability in average positions or trajectories, but in order
to rein in the SD of postural positions (i.e., of the center of pressure, CoP), the movement system still needs
exploratory behaviors to use the precision demand. Indeed, precision-related reductions in *SD* of position can
coincide with stronger signatures of complexity, suggesting that exploratory behavior does not live solely in the
bulk of movement variability but also in its patterning (Haddad et al., 2010, 2008). After all, extrinsic constraints
from the task context are rarely stipulated impositions and much more often available supports for organisms to
engage with through exploration or not engage at all (Stoffregen et al., 2017). Precision demands can be no
different but are more likely available features of the context with effects only contingent on the organism's
exploratory behaviors. We expect that organisms can meet progressively stronger precision demands only
through progressive—and prior—increases in their exploratory behaviors' multifractal nonlinearity.

Closing the eyes should introduce an *SD*-increasing effect of multifractal nonlinearity. Closing eyes
abruptly changes the visual support for posture (Stoffregen, 1986), destabilizing the movement system, and the
priority for the postural system to maintain quiet stance is sooner to damp out fluctuations and freeze degrees of
freedom. Hence, if we destabilize posture, postural sway (e.g., as measured by *SD* of CoP fluctuations) generally
increases with greater multifractality (Kelty-Stephen, 2018). These prior observations include a failure to find a
concurrent correlation of multifractal-spectrum width with *SD* of CoP position. Multifractal variability has
elsewhere varied inversely with stable postural outcomes, with narrower multifractal spectra appearing in
younger adults and in participants less likely to experience motion sickness (Koslucher et al., 2016; Munafo et
al., 2016a). However, previous work has never before integrated the t_{MF} statistic, vector-autoregressive effects of
one variable on subsequent values of another, and the potential moderation by visual precision. Hence, the
present test of the effects of multifractal nonlinearity on subsequent *SD* of CoP fluctuations with greater
precision demands remains so far untested in the literature.

As the second entailment of including multifractal nonlinearity is the possibility of predicting the apparently 'self-correcting' aspect of monofractal evidence. For example, as noted above, the need for postural sway to reverse course from the base of support's margins has been a long-standing explanation of why postural control shows a transition from stronger and weaker monofractal evidence by turn (Collins and De Luca, 1995, 1993). That is, we expect that increases in multifractal nonlinearity would prompt oscillations in subsequent monofractal evidence. Changes in monofractal evidence may well contribute to multifractal nonlinearity, but because of the centrality of multifractal nonlinearity to task-sensitivity, we suspect that the effects of monofractal evidence on subsequent multifractal nonlinearity would be strongest in less demanding tasks. The corollary is that greater visual precision demands would minimize the contribution of monofractal evidence to multifractal nonlinearity, emphasizing multifractal nonlinearity to the exclusion of any effects of monofractal evidence.

The final entailment of including multifractal nonlinearity is a possible displacement of the effects linking monofractal evidence to subsequent *SD* and linking *SD* to subsequent monofractal evidence. Previous work does not predict whether the effects of monofractal evidence on subsequent *SD* might go to zero or even reverse. However, given the more central and substantive meaning of multifractal nonlinearity supporting dexterous goal-directed behavior, we expected multifractal nonlinearity to influence subsequent *SD* and, furthermore, show subsequent changes due to prior changes in *SD*.

In summary, elaborating from mono- to multifractal formalisms will allow us to delve deeper into the multifractal structure supporting postural control. This move aims to situate monofractal evidence—which can be modeled entirely linearly—within a broader framework of nonlinear relationships. Instead of simply reporting a correlation table of concurrent relationships, we aim to model how SD, monofractal evidence, and multifractal nonlinearity predict subsequent changes in one another.

1.4.3. Hypotheses

Hypothesis-1: Weaker intrapostural interactivity while quietly standing with eyes closed

First, respecting the evidence that maintaining fixation recruits visual precision demands that can perturb posture (Fiorelli et al., 2017; Lê and Kapoula, 2008), we expected that the addition of a fixation task to quiet standing would generally accentuate intrapostural interactivity, suggesting that quietly standing with eyes closed would show weaker evidence of intrapostural interactivity.

Hypothesis-2: Self-correction of sway monofractality across 10-s segments

Postural sway is more correlated at shorter timescales on the order of 10 s and more anticorrelated over longer timescales (Collins and De Luca, 1993). For instance, posture roams freely about a fixed average CoP position but self-corrects at the margins of the base of support to maintain a quiet stance (Collins and De Luca, 1995, 1993; Gilfriche et al., 2018). In this sense, we expected that monofractal evidence itself would show similar self-corrections over time, with prior increases in fractal evidence followed by subsequent alternating reductions and increases in monofractal evidence. This self-correcting feature would be a crucial part of any control system in which temporal correlations stabilize sway.

Hypothesis-3: Inverse relationship between sway monofractality and SD across 10-s segments

Because monofractality entails long-range temporally organized responsivity to mechanical perturbations (Duarte and Sternad, 2008; Duarte and Zatsiorsky, 2001), we expected that either or both of CoP and CoM monofractality would show an inverse relationship with CoP *SD*. That is, increases in CoP and CoM monofractality would predict subsequent reductions in CoP *SD* (Hypothesis-3a), as well as increases in CoP *SD* would predict subsequent reductions in CoP and CoM monofractality (Hypothesis-3b). This feature could reflect a second key component of task-dependent emergent postural control.

Hypothesis-4: The inverse relationship between sway monofractality and SD is stronger with eyes open than eyes closed and becomes progressively stronger at viewing distances closer to 50 cm

The 50-cm viewing distance is within the comfortable viewing distance, ideal for the human eyes' focus of red light (Párraga et al., 2002), as well as ideal for requiring the least straining oculomotor convergence (Jaschinski-Kruza, 1991; Jaschinski, 2002). Thus, we expected that the 50-cm viewing condition would yield

intrapostural interactivity most closely resembling that in the eyes-closed condition. Because 50-cm imposes the weakest precision demand, the effects among CoM monofractality, CoP fractality, and CoP SD would show the least differences from the eye-closed condition. We also expected that the increasing visual precision demand as viewing distance deviates from 50 cm would accentuate the inverse relationship between prior monofractality and subsequent CoP SD. That is, increases in CoP monofractality would lead to less subsequent CoP SD at viewing distances 20 and 135 cm than at viewing distances of 220 and 305 cm.

Hypothesis-5: The influence of sway multifractal nonlinearity t_{MF} on subsequent increases in sway SD through the bidirectional relationship of t_{MF} with monofractal evidence H_{fGn}

We expected that sway multifractal nonlinearity t_{MF} would displace monofractal evidence H_{fGn} as the stronger predictor of subsequent reductions in SD with greater precision demands. Specifically, we expected that increases in t_{MF} would predict later reductions in SD and progressively larger reductions with progressively stronger visual-precision demands (Hypothesis-5a).

We expected that multifractal nonlinearity would also predict a difference between eyes-closed condition and 50-cm viewing condition that multifractality would not. Specifically, we expected that, for the eyes-closed condition, increases in multifractal nonlinearity would predict subsequent increases in SD of postural fluctuations (Hypothesis-5b).

We also expected that (i) multifractal nonlinearity would infiltrate the earlier-predicted bidirectional effects only between monofractal evidence and SD (e.g., in Hypotheses-3 and -4); (ii) multifractality nonlinearity would predict subsequent oscillatory reversals in monofractal evidence (Hypothesis-5c); (iii) monofractal evidence would predict subsequent increases in multifractal nonlinearity with progressively lower visual precision demands (Hypothesis-5d); (iv) including multifractal nonlinearity in our modeling would reverse the monofractal-only Hypothesis-4 as follows: with the inclusion of multifractal nonlinearity, monofractal evidence would fail to predict subsequent precision-dependent reductions in SD, yielding instead potentially only partially negative or potentially positive subsequent changes in SD (Hypothesis-5e). Finally, we expected that including multifractal nonlinearity in our modeling would reverse the monofractal-only Hypothesis-3 as follows: with the inclusion of multifractal nonlinearity, prior increases in SD would predict subsequent reductions in multifractal nonlinearity—and no longer any subsequent changes in monofractal evidence (Hypothesis-5f).

2. Materials and methods

2.1. Participants

Seven adult men and nine adult women ($M\pm1SD$ age = 23.8±3.9 years) without any skeletal or neuromuscular disorder voluntarily participated after providing verbal and written consent approved by the Institutional Review Board (IRB) at the University of Georgia (Athens, GA).

2.2. Experimental task and procedure

Each participant stood barefoot with one foot on each of two force plates (AMTI Inc., Watertown, MA), 25 cm apart (figure 1*a*). From behind the participant, a laser pen projected a static point-light on the center of a 5×5" white tripod-mounted screen in front of a white visual-field-filling background. The two force plates measured 3D moments and ground reaction forces. The full-body motion of each participant was measured using VICON Plug-in Gait full-body 39 marker set and an 8-camera VICON motion tracking system (VICON Inc., Los Angeles, CA). The kinetic and kinematic data were synchronously sampled at 100 Hz.

Each participant was instructed to maintain a standing position and minimize postural sway while looking at a point projected in front of them at their eye level with their head in a neutral position. Each participant was tested under six different viewing conditions: eyes closed and the point-light point positioned at 25, 50, 135, 220, and 305 cm distance in front. Each participant completed 18 trials (6 conditions × 3 120-s trials) in a single 90-min session with randomized trial order and with breaks on request and between every six trials.

2.3. Data processing

All data processing was performed in MATLAB 2019b (Matlab Inc., Natick, MA). The position of the
bodily center of mass (CoM) was estimated by submitting segment lengths of the head, trunk, pelvis, and left
and right hand, forearm, upper arm, thigh, shank and foot to the equations provided by Zatsiorsky and Seluyanov
(Zatsiorsky and Seluyanov, 1985), which yielded a 3D center of pressure (CoP) series describing CoM position
along the participant's anterior-posterior (*AP*), medial-lateral (*ML*), and superior-inferior axes. 3D moments and
ground reaction forces measured on each trial yielded a 2D center of pressure (CoP) series describing CoP
position along the participant's AP and ML axes. Over 120 s duration, each trial yielded a 3D CoM series and a
2D CoP series each of 120 s or 12000 samples, divided into 12 segments of 10 s or 1000 samples each. Each
segment yielded two 999-sample one-dimensional series: a CoM spatial Euclidean displacement series
describing the amplitude of CoM displacement and a CoP planar Euclidean displacement series describing the
amplitude of CoP displacement (Figs. 2A to 2F). We estimated the SD of CoP planar Euclidean displacement
series (henceforth, CoP-*SD*).

2.4. Detrended fluctuation analysis (DFA)

DFA estimates Hurst exponent, H, describing the growth of root mean square (RMS) fluctuations with
time for first-order displacements known as fractional ('fractal' for short) Gaussian noises (fGn) (Peng et al.,
1995, 1994). First, it integrates time series $x(t)$ with N samples to produce $y(t)$:

$$y(t) = \sum_{i=1}^{N} x(t) - \overline{x(t)}. \tag{1}$$

Next, DFA computes RMS residuals from the linear trend $y_n(t)$ over nonoverlapping n-length bins of $y(t)$ to
build a fluctuation function $f(N)$:

$$f(N) = \sqrt{(1/N) \sum_{i=1}^{N} \left(x(t) - \overline{x(t)} \right)^2}, \tag{2}$$

for $n < N/4$. On standard scales, $f(N)$ is a power law, that is, $f(N) \sim^{\square} n^H$, where H is the scaling exponent. H
is estimated as the slope of $f(N)$ in log-log plots:

$$\log f(N) = H \log(n). \tag{3}$$

DFA estimated H_{fGn} for the original version (i.e., unshuffled) and a shuffled version (i.e., a version with
the temporal information destroyed) of each segment of CoM spatial-displacement series (henceforth, CoM-H_{fGn})
and CoP planar-displacement series (henceforth, CoP-H_{fGn}) over the following bin sizes: 4, 8, 12,… 128).

H_{fGn} for all original series fell in the fractal range ($0.5 < H_{fGn} < 1$; Fig. 2G and 2H) and significantly
exceeded H_{fGn} for all corresponding shuffled series ($Ps < 0.0001$), indicating evidence of monofractality in both
CoM and CoP (e.g., Figs. 2G and 2H). Crossovers between shorter- (bin sizes: 4, 8, 12,… 64) and longer-scale
(bin sizes: 4, 8, 12,… 128) behavior exhibited no reliable differences, as H_{fGn} estimates from only shorter-scale
behavior correlated strongly with H_{fGn} estimates from the entire DFA fluctuation function (CoM: Spearman's ρs
= 0.96, 0.91, 0.94, 0.94, 0.93, and 0.94 for the eyes-closed and 25-, 50-, 135-, 220-, and 305-cm viewing
conditions, respectively, $Ps < 0.0001$; CoP: ρs = 0.96, 0.89, 0.93, 0.92, 0.91, and 0.94, $Ps < 0.0001$).

2.5. Direct estimation of multifractal spectra

The multifractal spectrum is a relationship between two fractal dimensions α and $f(\alpha)$. One multifractal
elaboration of DFA uses DFA-estimated H_{fGn} to estimate both α and $f(\alpha)$ (Kantelhardt et al., 2002). However,
we use Chhabra and Jensen's (1989) direct method because it does not rest on fragile assumptions. Specifically,
multifractal DFA can detrend series only with polynomial trends and is not as effective for sinusoidal trends at a
variety of frequencies (Bashan et al., 2008; Ignaccolo et al., 2010). Furthermore, the validity of estimating both
α and $f(\alpha)$ from the same H is untested in many biological examples (Zamir, 2003). We prefer Chhabra and
Jensen's (1989) method exactly because it estimates α and $f(\alpha)$ independently, providing an estimate of
multifractal-spectrum width without the risk of amplifying any errors in multifractal DFA's double use of the
same H_{fGn} estimates.

This Chhabra and Jensen's method (1989) samples non-negative series $u(t)$ at progressively larger scales
such that proportion of signal $P_i(L)$ falling within the i^{th} bin of scale L is

$$P_i(L) = \frac{\sum_{k=(i-1)L+1}^{iL} u(k)}{\sum u(t)}$$ (4)

As L increases, $P_i(L)$ represents progressively larger proportion of $u(t)$,

$$P(L) \propto L^{\alpha}$$ (5)

suggesting growth of proportion according to one 'singularity' strength α. $P(L)$ exhibits multifractal dynamics
when it grows heterogeneously across timescales L according to multiple potential fractional singularity
strengths, such that

$$P_i(L) \propto L^{\alpha_i}$$ (6)

whereby each i^{th} bin may show a distinct relationship of $P(L)$ with L. The spectrum of singularities is itself the
multifractal spectrum, and its width $\Delta\alpha(\alpha_{max} - \alpha_{min})$ indicates the heterogeneity of these relationships between
proportion and timescale (Halsey et al., 1986; Mandelbrot, 1999).

This method estimates $P(L)$ for N_L nonoverlapping bins of L-sizes and accentuates higher or lower
$P(L)$ by using parameter $q > 1$ and $q < 1$, respectively, as follows

$$\mu_i(q, L) = \frac{[P_i(L)]^q}{\sum_{i=1}^{N_L} [P_i(L)]^q}.$$ (7)

See *figure* 2.

$\alpha(q)$ is the singularity for $\mu(q)$-weighted $P(L)$ estimated by

$$\alpha(q) = -\lim_{L \to \infty} \frac{1}{\ln L} \sum_{i=1}^{N} \mu_i(q, L) \ln P_i(L)$$

$$¿\lim_{L \to 0} \frac{1}{\ln L} \sum_{i=1}^{N} \mu_i(q, L) \ln P_i(L).$$ (8)

Estimates $\alpha(q)$ belong to the multifractal spectrum if Shannon entropy of $\mu(q, l)$ scales with L according to a
dimension $f(q)$, where

$$f(q) = -\lim_{L \to \infty} \frac{1}{\ln L} \sum_{i=1}^{N} \mu_i(q, L) \ln \mu_i(q, L)$$

$$¿\lim_{L \to 0} \frac{1}{\ln L} \sum_{i=1}^{N} \mu_i(q, L) \ln \mu_i(q, L).$$ (9)

For q generating scaling relationships (Eqs. 8 and 9) with correlation coefficient, $r > 0.9$, the parametric curve
$(\alpha(q), f(q))$ or $(\alpha, f(\alpha))$ constitutes the multifractal spectrum with width $\Delta\alpha = \alpha_{max} - \alpha_{min}$. $\Delta\alpha$ was estimated
for each segment of CoM spatial-displacement series and CoP planar-displacement series (Figs. 2I and 2J).

To identify if nonzero $\Delta\alpha$ reflected nonlinear temporal correlations, $\Delta\alpha$ of each original series, was
compared to $\Delta\alpha$ of 32 surrogate series obtained using Iterated Amplitude Adjusted Fourier Transformation
(IAAFT) (Ihlen, 2012; Mandic et al., 2008). IAAFT generates surrogates with randomized phase ordering of the
series' spectral amplitudes but preserved linear temporal correlations. When $\Delta\alpha$ of the original series exceeds 1

95% confidence interval from $\Delta\alpha$ of IAAFT series (i.e., $p < 0.05$), the original series is considered to have
nonzero nonlinear correlations, easily quantifiable using the one-sample t-statistic (henceforth, CoP-t_{MF})
comparing $\Delta\alpha$ of the original series to that of the surrogates.

Finally, across all data (15 participants × 6 conditions/participant × 3 trials/condition × 12 segments/trial
= 3240 segments), 3232/3240 or 99.75% CoM segments and 2935/3240 or 90.59% CoP segments exhibited
multifractal spectrum that were significantly wider for the corresponding IAAFT surrogates (e.g., Fig. 3).

531 *2.6. Vector autoregression analysis*

Vector autoregression captures linear interdependencies amongst concurrent series and here modeled
intrapostural effects of CoP-SD, CoP-H_{fGn}, and CoM-H_{fGn} in one segment on CoP-SD, CoP-H_{fGn}, and CoM-H_{fGn}
in subsequent segments (Hypotheses 2-4), and effects of CoP-SD, CoP-H_{fGn}, and CoM-t_{MF} in one segment on
CoP-SD, CoP-H_{fGn}, and CoM-t_{MF} in subsequent segments (Hypothesis-5; Fig. 4). Vector autoregression describes
each variable based on its own lagged value and that of each other variable. Lag is ideally increased until the
residuals appear independently and are distributed identically (Sims, 1980).

Vector autoregression allows forecasting unique effects of endogenous variables on later values of each
other through impulse-response functions (IRFs). Impulse-response functions evaluate the relationship between
$f(t)$ and $g(t+\tau)$, or between $g(t)$ and $f(t+\tau)$, where τ is a whole number corresponding to a segment within a
trial. Provided vector-autoregression residuals are independent and identically distributed (i.i.d.),
orthongonalizing these residuals allows simulating an 'impulse' to the system by adding $1\,SEM$ to any single
variable, and using vector-autoregression coefficients to describe propagation of later 'responses' across all
endogenous variables. The impulse-response function describes how an impulse in one series changes later
predicted values in a different time series (Hatemi-J, 2004; Lutkepohl, 2007). All vector autoregression models
converged with lag 1, with residuals passing all tests for i.i.d. status. We performed vector autoregression
analysis using the function VAR() from the *vars* package (Pfaff et al., 2018) in *RStudio* (Team, 2013).

548 *2.7. Statistical analysis of pairwise impulse-response functions*

549 *2.7.1. Predictors*

Mixed-effect regression models (Singer and Willett, 2003) treated impulse-response functions between
each pair of postural descriptors as the dependent measure, testing the main effects of 'Eyes Closed' (EC;
equaling 1 for the eyes-closed condition and 0 for all other conditions), and 'Comfortable Viewing Distance'
(CVD; equaling 1 for 50-cm viewing condition and 0 for all other conditions), and 'Distance From Comfortable
Viewing Distance' (DF_CVD: the absolute distance of viewing conditions from 50 for viewing conditions, i.e.,
25, 0, 85, 170, and 255, as well as 0 for the eyes-closed condition), as well as each of the interaction of each of
these main effects with the full-factorial family of interactions Trial × Segment × Impulse × Response. Impulse
and Response served as class variables encoding the different descriptors (CoM-H_{fGn}, CoP-SD, and CoP-H_{fGn} for
Hypotheses-1–4; and including CoP-t_{MF} for Hypothesis 5; in both cases, CoM-H_{fGn} served as the baseline level
for both Impulse and Response) serving as impulse and as response variables, respectively.

We thus used two mixed-effect regression models of the impulse-response functions: the first with the
first set of impulse-response functions and the second with both sets of impulse-response functions (i.e., with
and without multifractal contributions, respectively). In both cases, CoM-H_{fGn} served as the baseline level for
both Impulse and Response. In the second regression model, we distinguished CoP-H_{fGn} as values of Impulse and
CoP-SD as values of Response specifically to CoP-t_{MF} between the two vector autoregressions. ('*' after H_{fGn} and
SD indicate the impulse-response functions from the second vector autoregression). The Impulse × Response
terms highlighted significant differences of specific impulse-response functions from the global patterns. The
description of results was made simpler by by the fact that the effects of multifractal nonlinearity outweighed all
effects of monofractal evidence, as the second model showed no effects among the impulse-response functions
from the first vector autoregression.

Each model was fit using the function lmer() from the *lme4* package (Bates et al., 2015) in *RStudio*
(Team, 2013), with only a random-effect intercept for each participant. In the second model, a unique intercept
was used for each participant in each vector autoregression to control for individual differences in impulse-

response functions within each vector autoregression. Pearson's correlation of residuals with model predictions
was near zero for each of the two models ($r = 0.0027$, $p = 0.679$; and $r = 0.0083$, $p = 0.067$).

2.7.2 Mixed-effect regression modeling for Hypotheses 1-4 and then Hypothesis 5 addressed impulse-response functions for only one vector-autoregressive model and both vector-autoregressive models, respectively

Hypotheses 1–4 involved modeling the impulse-response functions from a vector-autoregressive model,
including only CoM-H_{fGn}, CoP-SD, and CoP-H_{fGn} (Dataset S1). Including CoP-t_{MF} as a fourth variable led to a
vector-autoregressive model that failed to converge with i.i.d. residuals. Therefore, because CoM-H_{fGn} ultimately
exhibited little influence on the other variables, we omitted it from the second vector-autoregressive model,
which included only CoP-SD, CoP-H_{fGn}, and CoP-t_{MF}. Hypothesis-5 involved modeling the cumulative set of
impulse-response functions from both vector-autoregressive models (Dataset S2). This modeling choice reflects
our interest in developing Hypothesis-5 as an elaboration of Hypothesis-1–4 without amplifying Type-I error.
Hypothesis-5 requires additional impulse-response functions that Hypotheses-1–4 does not. Pooling all impulse-
response functions allowed testing Hypothesis-5 above and beyond the earlier-hypothesized structure instead of
testing it separately in parallel.

2.7.3. Orthogonal polynomials for segments within trial

Following a convention that we had developed in earlier work (Mangalam et al., 2020a, 2020b),
orthogonal linear, quadratic, and cubic polynomials of Segment modeled how impulse-response functions
changed over 999-sample segments within a trial. The cubic polynomial was used to capture the nonlinear decay
of impulse-response functions across later segments. The interactions with Segment indicated changes in these
effects with different third-order polynomial responses over subsequent trials. However, the cubic polynomial
failed to improve model fit, so was omitted. The second regression model was much larger to model impulse-
response functions from two separate vector autoregressions. The inclusion of multifractal nonlinearity within
vector-autoregressive modeling had an enormous effect on reshaping the impulse-response functions. One of the
consequences was the elaboration of the polynomial structure. The maximum-order polynomial for modeling
across impulse-response values for each of 10 subsequent segments was 9, and the 9th-order polynomial
significantly improved model fit ($\chi^2(384) = 869.64$, $p < 0.0001$). Because this 9th-order polynomial interacted
only with t_{MF} as Impulse and as Response, we used 9th-order polynomials for the second model but not the first
model. However, the second model omitted interactions of Trial with EC and DF_CVD because these
interactions no longer improved model fit when testing Hypothesis-5—this failure of Trial interactions to
improve model fit was equally true whether the second model included only 3rd- or all 9th-order polynomials,
($\chi^2(256) = 164.04$ and $\chi^2(640) = 294.56$, $Ps = 1.000$).

2.7.4 Addressing multiple comparisons by nesting hypotheses and testing new hypotheses while preserving data and model structure for earlier hypothesis

Despite the abundance of coefficients reported below, we stress that we report below the outcomes of
only two regression models. Hypothesis-5 was nested both within the data and the model for Hypotheses-1–4—
and not in a separate, new model. We inferred significance using P-values, with alpha set to the conventional
0.05 level in each model. Vector autoregression and impulse-response modeling appear here only to provide
average measures, and while it is possible to estimate p-values for these VAR- and impulse-response-produced
coefficients, they were not used here. Instead, differences among impulse-response model-generated coefficients
were tested through a cumulative multiple-regression modeling procedure that retained, for Hypothesis-5, all
previous structures for testing Hypotheses-1–4.

Such multiple regression allows simultaneously testing multiple predictors, precluding the need for
addition correction (Williams, 1971). Furthermore, traditional corrections (e.g., Bonferroni) are apt when the
dependent variable is correlated. However, because the test of Hypothesis-5 operates on both new impulse-
response functions and old impulse-response functions appearing in the test of Hypotheses-1–4, the dependent
variables for these distinct hypotheses cannot be correlated in any standard way. Retaining the impulse-response
functions for Hypotheses-1–4 in the dataset while testing additional impulse-response functions ensured that the
second model incorporated all of the structure of the first model.

3. Results

3.1. Regression modeling of impulse-response functions relating CoM-H_{fGn}, CoP-SD, and CoP-H_{fGn}

3.1.1. Maintaining a quiet stance with eyes closed weakened intrapostural interactivity (Hypothesis-1)

The only significant effect in the eyes-closed condition on the intercept of the impulse-response
functions was the one representing responses of CoP-H_{fGn} impulses (EC × Impulse(CoP-H_{fGn}): $B = 7.06×10^{-3}$, P
$= 2.88×10^{-2}$). This coefficient canceled out the negative responses of CoP-H_{fGn} impulses (Impulse(CoP-H_{fGn}): $B =$
$-6.84×10^{-3}$, $P = 1.73×10^{-3}$). All other significant effects in the regression refereed to polynomial changes in
impulse-response functions (Tables 1 & 2), suggesting that the eyes-closed condition yielded impulse-response
functions varying according to a polynomial centered on a zero mean.

3.1.2. CoM-H_{fGn} and CoP-H_{fGn} self-corrected from segment to segment within a trial but showed sparse effects
on each other (Hypothesis-2)

In viewing conditions, increases in CoM-H_{fGn} and CoP-H_{fGn} predicted subsequent increases and
reductions in alternation over segments (Figs. 5A and 5B). Visually fixating prompted monofractality to fall in
and out of zero change or to cycle around zero change with negative and positive changes following each other,
suggesting that viewing conditions prompted a sort of self-correcting maintenance of monofractality within CoM
and CoP. The model did not yield significant impulse-response functions between CoM-H_{fGn} and CoP-H_{fGn} either
(Figs. 5C and 5D): only the 220-cm viewing condition was accompanied by a CoM-H_{fGn} impulses that predicted
CoP-H_{fGn} responses (Fig. 5C).

3.1.3. Increases in CoP-H_{fGn} and CoP-SD predicted subsequent reductions in each other across trials
(Hypothesis-3)

In viewing conditions, CoP-H_{fGn} and CoP-*SD* impulses both predicted subsequent reductions in CoP-*SD*
and CoP-H_{fGn}, respectively (Fig. 6). These impulse-response functions remained robust with Trial, more so for
the effects of CoP-*SD* impulses on CoP-H_{fGn} responses (Fig. 5B) than for the effects of CoP-H_{fGn} impulses on
CoP-*SD* responses (Fig. 5A). 305-cm viewing condition failed to show a significant effect of CoP-*SD* impulses
on CoP-H_{fGn} responses on only one trial. The 25-, 135-, and 220-cm viewing conditions each exhibited one, one,
and two trials, respectively, that failed to show the effects of CoP-H_{fGn} impulses on CoP-*SD* responses.

3.1.4. Impulse-response relationship between increases in CoP-H_{fGn} on subsequent reductions in CoP-SD was
weakest for 50-cm viewing condition and became progressively stronger as viewing distance deviated from 50
cm (Hypothesis-4)

Of all viewing conditions, the 50-cm viewing condition prompted the least evidence that increases in
CoP-H_{fGn} predicted subsequent reductions in CoP-*SD* (Table 3, Fig. 6A). For this comfortable viewing distance
of 50 cm (i.e., Distance from Comfortable Viewing Distance, DF_CVD = 0), the intercept for impulse-response
relating CoP-H_{fGn} impulses to CoP-*SD* responses was effectively zero. The effect of CoP-H_{fGn} impulses on CoP-
SD responses (Impulse(CoP-H_{fGn}) × Response(CoP-*SD*): $B = 7.24×10^{-3}$, $P = 1.91×10^{-2}$) canceled out the general
effect of CoP-H_{fGn} impulses (Impulse(CoP-H_{fGn}): $B = -6.84×10^{-3}$, $P = 1.73×10^{-3}$). In fact, the only intrapostural
interactivity that appeared for the comfortable viewing distance was that an increase in CoP-*SD* predicted a
subsequent increase in CoM-H_{fGn} (CVD × Impulse(CoP-*SD*): $B = 9.30×10^{-3}$, $P = 3.98×10^{-3}$). The effects of CoP-
SD impulses on subsequent CoP-H_{fGn} and CoP-*SD* responses canceled out this positive effect (CVD ×
Impulse(CoP-*SD*) × Response(CoP-H_{fGn}): $B = -1.03×10^{-2}$, $P = 2.36×10^{-2}$; CVD × Impulse(CoP-*SD*) ×
Response(CoP-*SD*): $B = -1.00×10^{-2}$, $P = 2.85×10^{-2}$, Fig. 6B). The 50-cm viewing distance resembled the eyes-
closed case in that it carried many of the polynomial terms, suggesting impulse-response functions that varied
nonlinearly around a zero mean.

For all other viewing distances, an increase in CoP-H_{fGn} predicted a subsequent reduction in CoP-*SD*,
and it was a larger reduction as viewing distance deviated in either direction (DF_CVD × Impulse(CoP-H_{fGn}): B
$= 5.03×10^{-5}$, $P = 2.36×10^{-4}$; DF_CVD × Impulse(CoP-H_{fGn}) × Response(CoP-*SD*): $B = -5.33×10^{-5}$, $P = 5.88×10^{-}$
^{3}, Table 4, Fig. 6A). This reduction in CoP-*SD* following an increase in CoP-H_{fGn} became progressively larger
across trials (DF_CVD × Trial × Impulse(CoP-H_{fGn}): $B = -1.43×10^{-5}$, $P = 2.36×10^{-2}$). For instance, the reduction
in CoP-*SD* following an increase in CoP-H_{fGn} ranged from $-3.83×10^{-5}$ to $-1.13×10^{-2}$, for 25-cm viewing
condition on Trial-1 to 255 cm on Trial-3, respectively (Table 1).

3.2. Regression modeling of impulse-response functions incorporating CoP-t_{MF} along with the prior set of
CoM-H_{fGn}, CoP-H_{fGn}, and CoP-SD

This second regression model pooled impulse-response functions from both vector autoregressions.
Hence, before we detail specific effects in this model, it is noteworthy that this regression model found no effects
within the impulse-response functions from the first model that excluded multifractal nonlinearity but only
effects among the impulse-response functions from the vector autoregression that included multifractal
nonlinearity. Hence, this initial summary point reflects that multifractal nonlinearity exerted much stronger
effects on monofractal evidence or *SD* than either had exerted on the other.

We also note that the use of random-effect intercept for participant-by-participant vector autoregression
will necessarily make the regression model structure diverge from the simpler plot of Segment-by-Segment
means of the impulse-response functions. We regret this divergence for how it can make the results more
challenging to digest. However, this random-effect intercept term's usefulness addressed the fact that impulse-
response functions varied substantially across participants. We have elsewhere documented that the early values
of impulse-response functions predict individual differences in perceptual responses and, consequently, in the
use of available information (Mangalam et al., 2020a, 2020b). The regression model takes a deeper grasp of the
impulse-response functions below the more idiosyncratic variation in raw magnitude with its 9th-order
polynomial structure.

With the preceding caveat, we share Segment-by-Segment participant-wise averages ($N = 15$) of
impulse-response functions in figures with individual panels for each condition (Figs. 7 and 8). The following
results will speak to the model coefficients and illustrated through the model-predicted form of the impulse-
response functions in each case (Figs. 9–11).

3.2.1. Increasing visual-precision demand led multifractal nonlinearity to predict larger subsequent decreases
in SD of postural fluctuations (Hypothesis-5a)

In 50-cm viewing condition, increases in multifractal nonlinearity predicted oscillations in subsequent
SD (Fig. 9A). After predicting an initial reduction in subsequent *SD* (2 segments ahead), increases in multifractal
nonlinearity predicted a zig-zag pattern of subsequent changes reaching briefly back into positive values. That is,
increasing multifractal nonlinearity predicted subsequent increases in *SD* (only 3 and 5 segments ahead). But this
bobbling of subsequent changes in *SD* became progressively more negative, consistently predicting reductions in
subsequent *SD*. This negative-then-briefly-positive-and-negative-again structure was specified by the 9th-order
polynomials reflecting the impulse from CoP-t_{MF} (Impulse(CoP-t_{MF}), $B = 9.75 \times 10^{-1}$, $P = 1.12 \times 10^{-2}$; 9th-order
Segment × Impulse(CoP-t_{MF}), polynomial coefficients from linear to nonic, $Ps < 0.05$, Table 5; CVD × 9th-order
Segment × Impulse(CoP-t_{MF}), polynomial coefficients from cubic to nonic, $Ps < 0.05$, Table 6) and the response
from CoP-*SD* in 50-cm viewing condition (Response(CoP-*SD**), $B = -9.78 \times 10^{-1}$, $P = 7.0010^{-3}$; 9th-order
Segment all $Ps < 0.05$, Table 5; Response(CoP-*SD**), polynomial coefficients from linear to nonic, $Ps < 0.05$,
Table 5; and CVD × 9th-order Segment × Response(CoP-*SD*) polynomial coefficients from linear to nonic, $Ps <$
0.05, Table 6).

As visual precision demands increased, increases in multifractal nonlinearity predicted progressively
more stable reductions in subsequent *SD* (Fig. 9A). That is, compared to 50-cm viewing condition, subsequent
changes in *SD* showed progressively less oscillation and more consistent reductions in subsequent *SD* with
progressively more deviation from 50-cm viewing distance. This effect was indicated by the 9th-order
polynomials reflecting the impulse from CoP-t_{MF} with increasing distance from 50 cm (DF_CVD × 9th-order
Segment × Impulse(CoP-t_{MF}), polynomial coefficients from linear to nonic, $Ps < 0.05$, Table 7) and the response
from CoP-*SD* with increasing distance from 50 cm (DF_CVD × 9th-order Segment × Response(CoP-*SD**),
polynomial coefficients from linear to nonic, $Ps < 0.05$, Table 7).

3.2.2. Closing eyes led multifractal nonlinearity to predict subsequent increases in SD of postural fluctuations
(Hypothesis-5b)

Compared to the eyes-open conditions, the closed-eyes condition led multifractal nonlinear to predict
more increases in *SD* of postural fluctuations in 3 to 7 segments ahead and reductions in *SD* in 8 to 10 segments
ahead (Fig. 9A). However, this late reduction in subsequent *SD* was consistently weaker than the reduction of
SD in the eyes-open conditions. This effect was indicated by the 9th-order polynomials reflecting the impulse

from CoP-t_{MF} with eyes closed (EC × Impulse(CoP-t_{MF}), $B = 2.26 \times 10^0$, $P = 1.31 \times 10^{-4}$; EC × 9th-order Segment ×
Impulse(CoP-t_{MF}), polynomial coefficients from linear to nonic, Ps < 0.05, Table 8) and the response from CoP-
SD with eyes closed (EC × Response(CoP-SD*), $B = -2.26 \times 10^0$, $P = 1.86 \times 10^{-4}$; EC × 9th-order Segment ×
Response(CoP-SD*), polynomial coefficients from linear to nonic except quadratic and octic, Ps < 0.05).

3.2.3. Increases in multifractal nonlinearity predicted subsequent reductions in monofractal evidence in CoP fluctuations (Hypothesis-5c)

Increases in multifractal nonlinearity prompted an initial increase in monofractal evidence 1 Segment
ahead for all the eyes-open conditions, after which the subsequent change went to practically zero at 2 Segments
ahead (Fig. 9B). All the eyes-open conditions showed subsequent changes in monofractal evidence oscillating
from more negative to more positive change by turn beginning with 3 Segments ahead. At the extremes of
precision demand—both lowest (50-cm) and highest (220- and 305-cm)—the oscillations were actually weakest,
with subsequent changes in monofractal evidence almost always positive. At the intermediary precision demands
(25- and 135-cm), subsequent changes in monofractal evidence oscillated much more and with progressively
larger amplitude with progressive segments, indicating negative changes 5, 7, and 9 segments ahead and
indicating positive changes 4, 6, 8, and 10 segments ahead.

This effect for 50-cm viewing condition was indicated by 9th-order polynomials reflecting the impulses
of CoP-t_{MF} and of CVD × Impulse(CoP-t_{MF}) noted in Section 3.2.1 and the response from CoP-H_{fGn} to impulse
from t_{MF} (Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn}), $B = -9.74 \times 10^{-1}$, $P = 3.11 \times 10^{-2}$; 9th-order Segment ×
Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn}), polynomial coefficients from quadratic to nonic, Ps < 0.05, Table 5;
CVD × 9th-order Segment × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn}), polynomial coefficients from quartic to
nonic, Ps < 0.05, Table 6). The effect for varying precision demands was indicated by the 9th-order polynomials
reflecting the impulses of CoP-t_{MF} and of DF_CVD × Impulse(CoP-t_{MF}) noted in Section 3.2.1. and the response
from CoP-H_{fGn} with Impulse(CoP-t_{MF}) for values of DF_CVD (DF_CVD × 9th-order Segment × Impulse(CoP-
t_{MF}) × Response(CoP-H_{fGn}), polynomial coefficients from cubic to nonic, Ps < 0.05, Table 7).

The eyes-closed condition diverged from model-predictions for the eyes-open conditions. It exhibited a
large positive change in subsequent monofractal evidence 2 segments ahead, followed by smaller but relatively
sustained and non-oscillatory changes in subsequent segments. This effect was indicated by the 9th-order
polynomials reflecting the impulses from CoP-t_{MF} and EC × Impulse(CoP-t_{MF}) noted in Section 3.2.2. and the
response from CoP-H_{fGn} with impulse from CoP-t_{MF} (EC × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn}), B = –
2.26×10^0, $P = 1.80 \times 10^{-3}$; EC × 9th-order Segment × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn}), polynomial
coefficients from linear to nonic except quadratic, cubic, and octic, Ps < 0.05, Table 8).

3.2.4. Increases in monofractal evidence predicted subsequent increases in multifractal nonlinearity but less so for higher visual precision demands (Hypothesis-5d)

An increase in monofractal evidence predicted an increase in multifractal nonlinearity on the first
segment ahead, resembling the pattern of effects in Section 3.2.3. on the first segment ahead. Hence, it appeared
that increases in monofractal evidence and multifractal nonlinearity predicted subsequent increases in each other,
but only in the short term. However, this resemblance to the effects of multifractal nonlinearity on monofractal
evidence in Section 3.2.3 ended here.

Later than 1 segment ahead, monofractal evidence predicted subsequent increases in multifractal
nonlinearity for all the eyes-open conditions until 5 segments ahead (Fig. 10A). From 5 segments ahead and
onward, subsequent changes in monofractal nonlinearity began oscillating with progressively larger amplitude,
reaching negative values 6 segments ahead in all conditions except 50-cm and 220-cm viewing conditions. The
greatest subsequent changes appeared in 25- and 130-cm viewing conditions. Oscillations dwindled in 50-cm
and 220-cm viewing conditions and then reversed in 305-cm viewing condition. In summary, monofractal
evidence looked much like the inverse of results in Section 3.2.3. That is, when multifractal nonlinearity had
predicted subsequent increases or reductions in monofractal evidence (i.e., in Section 3.2.3.), monofractal
evidence predicted reductions or increases, respectively, in multifractal nonlinearity.

The foregoing effects were indicated by 9th-order polynomials reflecting the impulse from CoP-H_{fGn} for
all values of DF_CVD (9th-order Segment × Impulse(CoP-H_{fGn}*), quintic, septic, and nonic polynomial
coefficients, Table 7) and the response from CoP-t_{MF} for all values of DF_CVD (Response(CoP-t_{MF}), $B = -$

$5.57×10^{-1}$, $P = 3.64×10^{-2}$; 9th-order Segment × Response(CoP-t_{MF}), nonic polynomial coefficient, $P < 0.05$, Table
5; DF_CVD × Response(CoP-t_{MF}), $B = B = -7.30×10^{-3}$, $P = 6.82×10^{-6}$; DF_CVD × 9th-order Segment ×
Response(CoP-t_{MF}), polynomial coefficients from linear to nonic, Table S2).

As in Section 3.2.3, 50-cm viewing condition yielded greater subsequent changes in multifractal
nonlinearity comparable to that in 220-cm viewing condition (Fig. 10A). These effects were indicated by the 9th-
order polynomials reflecting the impulse of CoP-H_{fGn} for values of CVD (CVD × 9th-order Segment ×
Impulse(CoP-H_{fGn*}), nonic polynomial coefficient, Table 6) and reflecting Response(CoP-t_{MF}) for values of CVD
(CVD × 9th-order Segment × Response(CoP-t_{MF}), polynomial coefficients from linear to nonic, Table S2).

The eyes-closed conditions showed the weakest relationship between monofractal evidence and
subsequent changes in multifractal nonlinearity. Besides positive subsequent changes at 1 and 4 segments ahead
and a negative subsequent change 3 segments ahead, subsequent changes in monofractal evidence was near zero.
These effects were indicated by the 9th-order polynomials reflecting the impulse from CoP-H_{fGn} for values of EC
(EC × Impulse(H_{fGn}*), $B = 1.75×10^{0}$, $P = 3.15×10^{-3}$; EC × 9th-order Segment × Impulse(CoP-H_{fGn}*), nonic
polynomial coefficient, $P < 0.05$, Table 8) and reflecting the response from CoP-t_{MF} for values of EC (EC ×
Response(CoP-t_{MF}), $B = -8.56×10^{-1}$, $P = 3.13×10^{-7}$; EC × 9th-order Segment × Response(CoP-t_{MF}), polynomial
coefficients from linear to nonic, Table S2).

3.2.5. Increases in monofractal evidence predicted subsequent increases in SD for lower visual-precision demands but subsequent decreases for only the highest visual-precision demands (Hypothesis-5e)

788
789

Following an increase in monofractal evidence, the subsequent changes in *SD* were negative for all cases
during the first 3 segments ahead—except the 25-cm viewing condition, which exhibited a subsequent increase
in *SD* on the third segment ahead (Fig. 10B). Subsequently, the subsequent changes in SD oscillated for all the
eyes-open conditions, remaining mostly negative for the greater precision demand (220- and 305-cm viewing
conditions), but oscillating between positive and negative for intermediate precision demands (25- and 135-cm
viewing conditions), with larger oscillations for the weaker precision demands (i.e., 25-cm). These effects were
indicated by the 9th-order polynomials reflecting the response from CoP-*SD* for all values of DF_CVD
(DF_CVD × Impulse(CoP-H_{fGn}) × Response(CoP-*SD*), $B = -7.09×10^{-3}$, $P = 1.16×10^{-2}$; DF_CVD × 9th-order
Segment × Impulse(CoP-H_{fGn}) × Response(CoP-*SD*), nonic polynomial coefficient, $P < 0.05$, Table 7).

Meanwhile 50-cm viewing condition and eyes-closed conditions predicted subsequent increases in *SD*
throughout beginning at 5 and 4 segments ahead, respectively (Fig. 10B). These effects were indicated by the 9th-
order polynomials reflecting the response from CoP-*SD* for all values of CVD (CVD × 9th-order Segment ×
Impulse(CoP-H_{fGn}) × Response(CoP-*SD*) polynomial coefficients from linear to nonic, Table S2) and for all
values of EC (EC × Impulse(CoP-H_{fGn}) × Response(CoP-*SD*), $B = -1.75×10^{0}$, $P < 1.59×10^{-2}$; EC × ×
Impulse(CoP-H_{fGn}) × Response(CoP-*SD*), polynomial coefficients from linear to nonic, Table S2).

3.2.6. Increases in SD predicted subsequent reductions in multifractal nonlinearity, with progressively larger reduction for progressively higher visual precision demands and with greatest reduction for the eyes-closed condition (Hypothesis-5f)

805
806
807

Greater *SD* in postural fluctuation predicted subsequent reductions in multifractal nonlinearity. This
reduction was smallest for 50-cm viewing condition and progressively larger for conditions with higher visual
precision demands (Fig. 11). These effects were carried entirely by linear components of the polynomial for the
responses from CoP-t_{MF} for DF_CVD (Response(CoP-t_{MF}), $B = -5.57×10^{-1}$, $P = 3.64×10^{-2}$; 9th-order Segment ×
Response(CoP-t_{MF}), linear coefficient, $P < 0.05$, Table 5; DF_CVD × Response(CoP-t_{MF}), $B = -7.30×10^{-3}$, $P =$
$6.82×10^{-6}$; DF_CVD × 9th-order Segment × Response(CoP-t_{MF}), linear to nonic coefficients, Table S2) and CVD
(CVD × Response(CoP-t_{MF}), $B = 1.7185×10^{-2}$, $P = 2.14×10^{1}$; CVD × 9th-order Segment × Response(CoP-t_{MF}),
linear to nonic coefficients, Table S2).

The eyes-closed condition prompted the largest subsequent reductions in multifractal nonlinearity
following increases in *SD* (EC × Response(CoP-t_{MF}), $B = -2.14×10^{1}$, $P = 2.14×10^{1}$; EC × 9th-order Segment ×
Response(CoP-t_{MF}), linear to nonic coefficients, Table S2, Fig. 11).

4. Discussion

We tested five hypothesis concerning how visual precision demands might moderate intrapostural
interactivity. We predicted that: Standing quietly with the eyes closed would exhibit weaker intrapostural
interactivity (Hypothesis-1). Monofractal evidence in CoM and CoP, CoM-H_{fGn} and CoP-H_{fGn}, respectively,
would self-correct over time (Hypothesis-2). An inverse relationship between monofractality and *SD* over time,
that is, that increases in CoM-H_{fGn} and CoP-H_{fGn} would prompt subsequent reductions in CoP-*SD* (Hypothesis-
3a) and that increases in CoP-*SD* would prompt subsequent reductions in CoM-H_{fGn} and CoP-H_{fGn} (Hypothesis-
3b). These intrapostural interactions in the 50-cm viewing condition would most closely resemble intrapostural
interactions in the eyes-closed condition (Hypothesis-4). Increases in multifractal nonlinearity CoP-t_{MF} would
displace the role and response of monofractal evidence CoP-H_{fGn} appearing in Hypotheses-3a and -4, leaving
monofractal evidence to show less precision-dependent reduction of CoP-*SD* and to contribute to multifractality
for lower visual precision demands (Hypothesis-5). Results supported all hypotheses with the only exception
being the failure of CoM-H_{fGn} to participate in the relationships predicted in Hypothesis-3. We discuss the
implications of these findings below.

4.1. Ratcheting up our lens from monofractality to multifractality allows movement science to make contact with the nonlinear interactions across scales entailed by goal-directed and task-dependent behavior

This work vindicates the suggestion that multifractal dynamics might underlie the coordination
supporting postural control. Multifractal dynamics and the mathematical, geometrical framework for
investigating this proposal remains a novel and underdeveloped theme in movement science. Elements of this
framework have been successfully introduced in terms of the popularization of monofractal geometry (e.g., 1/*f*
noise as a key feature of movement variability; Newell & Jordan, 2007; Vaillancourt & Newell, 2002). However,
the more recent calls have been to ratchet the modeling up to a new precision level. We briefly review the bigger
picture below, on the way to explaining how this work supports a longer-range trajectory of developing
multifractal perspectives on movement science. The outcome of these deliberations will be twofold. Monofractal
modeling is an effective first step into multifractal modeling so far as multifractality follows from trial-by-trial
variation in monofractal structure. However, monofractal modeling has limits, and the further modeling of
multifractal spectra with the multifractal elaborations of monofractal modeling offers a deeper way to make
inferential contact with nonlinear interactions across scales. The value of this work lies in showing that
elaborating multifractal modeling beyond simply monofractal modeling—whether in isolation or sequence—
reveals new insights into how postural control resolves the intermodality of organism-wide perception-action.

First, monofractal modeling has always been an excellent first step, but notably not the last step.
Critically for motivating multifractal perspectives, the relationship between monofractal evidence and
'nonlinearity' is not as deep as initially hoped (Ihlen & Vereijken, 2010; Kelty-Stephen & Wallot, 2017). Yes, the
power-law shape implicated in 1/f noise or any single fractal dimension is not linear, and yes, the model system
once heralded as an explanation for 1/f noise (i.e., self-organized criticality) did enlist an interaction across
behaviors situated on two different scales (Bak et al., 1987; Jensen, 1998). However, monofractal evidence has
been insufficient on two fronts: it neither depends necessarily on self-organized criticality nor does it reach the
full scope of academic aspirations motivating much of its interest. To the first point, the appearance of a single
power law in our measurements is entirely consistent with and predictable from linear models that do not
necessarily have any interactions across scales (Granger and Joyeux, 1980; Wagenmakers et al., 2004). To the
second point, the theoretical discourse, particularly in ecological-psychological corners, has long sought to
implicate multiple (i.e., more than two and even potentially 'all') scales of biological and task variability
(Johnston & Turvey, 1980; Turvey & Fitzpatrick, 1993; Turvey, 2009; Wagman et al., 2016a, 2016b, 2017;
Wagman & Miller, 2003; Wagman & Stoffregen, 2020). To sum up, self-organized criticality is just one case of a
family of cascade models explicitly situating causal status in nonlinear interactions across scales, and choosing
monofractal modeling to the exclusion of multifractal considerations risks missing that goal-directed behavior
rests on cascade-driven causality beyond self-organized criticality (Stephen et al., 2012).

The issue of multifractal vs. monofractal modeling is about the lens we use to investigate goal-directed
behavior. Monofractal modeling can provide evidence consistent with cascade-like nonlinear interactions across
scales—cascades are one important source of monofractal structure (Mandelbrot, 1982). However, an isolated
case of monofractal evidence is nothing that the linear model cannot predict (Granger and Joyeux, 1980), and
any choice to interpret monofractal evidence as arising from cascades requires ruling out the null hypothesis that

it arises from the linear model (Wagenmakers et al., 2004). The linear model's capacity to predict monofractal
evidence is difficult to imagine as a plausible alignment of cause. Specifically, it appeals to a model called
'fractional integration' contrived to permit all past timescales of behavior to contribute to the current behavior—
with the puzzling entailment that none of these unbounded past behaviors interact with past behaviors at other
scales. Hence, for some researchers, the implausibility of causal interpretations from this linear model
originating from $1/f$ noise may be enough to warrant interpreting monofractal analysis as satisfactory evidence of
cascades. This compromise exemplifies what Gilden (2009) warned of as 'gross opportunism' for the analytical
sake of fitting the time-series data.

Monofractal analysis can presume interactions across scales, but multifractal analysis allows us to
estimate the strength of these interactions and specific roles of nonlinearity in supporting them. The mere
absurdity of linear explanation of monofractality does not somehow make monofractal evidence a better lens on
nonlinearity. Estimating the time-variation of monofractal evidence is the first step towards multifractality and
beyond the linear model of goal-directed behavior. If goal-directed behavior were strictly linear, then time-
variation of monofractal evidence should be without structure and incapable of predicting perception-action
outcomes (e.g., Section 1.3.2.1). However, as we highlighted in Section 1.3.2.2., analyses that are explicitly
multifractal can estimate a 'multifractal spectrum' directly without the intermediary step of many sequential
monofractal analyses. This spectrum width covaries with what you would find with sequential monofractal
analysis, but whereas the spectrum width glosses over the actual sequences in this temporal variation, the
multifractal spectrum offers the most compact method to estimate the nonlinear across-scales interactivity of a
single measured time series. Sequential multifractal estimation of this spectrum, that is, for multiple time series
in an experiment-wide sequence, thus reveals how nonlinear interactions across scales change alongside ongoing
measures of goal-directed behavior.

Monofractal evidence and multifractal nonlinearity inform about 'temporal structure' and 'nonlinear
support for that structure,' respectively. We can examine how the monofractal evidence and multifractal
nonlinearity both change alongside more universally interesting outcome measures. Hence, the multifractal
approach allows human movement science a wider set of features for articulating how nonlinear interactions
across scales could matter for dexterous goal-directed behavior. Certainly, monofractal evidence involves a
power-law relationship that unfolds across multiple scales, and the 'self-similarity' of this form is evocative of an
interaction among the implicated scales. But multifractal-spectrum width compared to surrogate spectra provides
the non-overlapping insight about whether the similar behavior at all scales reflect actual links between those
scales.

The advantage of multifractal modeling is that it resolves heterogeneity and continuity in single
framework. A major shortcoming of the monofractal modeling has been that a single estimate cannot necessarily
specify all wider patterning of a distributed system (Likens et al., 2019, 2015). 'Self-similarity' is certainly a
feature of power-law relationships supporting fractal and multifractal modeling, but we do not think it warrants a
denial of heterogeneity. Nonlinear dynamics are fully compatible with a difference and even competition (e.g.,
Kelso, 1995). As Bernstein (1967) suggested, "movements are not chains of details but structures which are
differentiated into details; they are structurally whole" (p. 69). The challenge is to maintain the possibility that
heterogeneity does not violate the organism's structural integrity (Stoffregen et al., 2017). Multifractal modeling
affects this balance and may provide further details about the nesting of behaviors within the organism. For
instance, multifractal nonlinear is a feature of physiological functioning (Ivanov et al., 2001), but manipulations
of the nested structure in the task context (e.g., Wagman et al., 2016a, 2016b, 2017; Wagman & Stoffregen,
2020) may prompt organisms to embody the nesting relationships quite apart from what physiological needs
demand. Multifractal analysis can reveal how nested structure in movement variability is sensitive to the nested
structure of the task context (Eddy & Kelty-Stephen, 2015; Stephen & Dixon, 2011).

This work shows the progression from monofractal to multifractal modeling in practice with postural
control as a model system. The findings can be summarized in the following way. Through the monofractal lens,
monofractal evidence suggests that $1/f$-like 'temporal structure' appears to reduce the *SD* of postural fluctuations,
suggesting as well that this 'temporal structure' self-corrects, keeping itself around some happy medium of the
temporal structure without excess in either direction. However, the multifractal lens shows us that $1/f$-like
temporal correlations are equally likely to increase or reduce the *SD* of postural fluctuations. The growth or

decay of *SD* can be better explained in terms of prior changes in how organisms blend behaviors across different
timescales. Any capacity of the organism to stabilize its monofractality inheres in its reservoir of nonlinear
interactions across scales.

4.2. Glimpses of a possible postural control policy for visually guided quiet stance

The present results offer insights into a control policy for postural stability that balances CoP
monofractality with excess CoP-*SD*. If left to *SD* alone, posture would lean towards higher variability without
clear bound: any increase in *SD* would predict subsequent increases, and those subsequent increases would, in
turn, predict further subsequent increases, and so on. The predicted subsequent reductions in monofractality
would then only serve to promote greater *SD*. It is only the corrective aspect of monofractality that might allow
posture to rein in the self-promoting and unbounded *SD*. For instance, any reductions in monofractality
following increases in *SD* might trigger subsequent increases in monofractality that would induce a negative
check on *SD*. Note that this causal interpretation aims only to offer a possible control policy that these results
could reflect. A rigorous test of this causal interpretation warrants manipulations of *SD* and CoP monofractality
through a balance board or vibrotactile stimulation.

The present results echoes the past findings that explicit feedback to participants completing a
perceptuomotor task can weaken temporal correlations in movement variability (Kelty-Stephen & Dixon, 2014;
Kuznetsov & Wallot, 2011; Stephen & Hajnal, 2011). In the task of counting seconds by tapping a finger
(Kuznetsov and Wallot, 2011), the feedback provided with each tap allowed offsetting deviations, preventing
errors from propagating from one tap to the next. At first glance, this finding seems at odds with the present
finding that better performance—standing more quietly with less sway *SD*—would follow from and contribute
to stronger and not weaker temporal correlations. However, it is possible that, in postural tasks, greater sway is
endogenous, implicit feedback signaling the postural system that corrections have been appropriately
implemented. In this way, if some proportion of *SD* reflects sway that triggers postural corrections (e.g., Hagio et
al., 2018), then the present findings would align with the past findings of feedback decorrelating movement
variability.

This control policy may resolve long-standing questions about how fluctuations support movement
stability, explicitly speaking to the 'loss of complexity' hypothesis: monofractality in sway might reflect stability,
suggestive of young, healthy, and typically developing physiology. This hypothesis has proven provocative but
controversial. Indeed, clinical research has found that temporal correlations might wander within but also
beyond the monofractal range. Additionally, the clinical implications of sway monofractality, with some results
indicating stronger temporal correlations in sway for younger and healthier participants (Duarte and Sternad,
2008; Thurner et al., 2002) and other results indicating the opposite (Ko and Newell, 2016; Lipsitz, 2002;
Thurner et al., 2002). Meanwhile, work applying 'white-noise'—that is, temporally uncorrelated—mechanical
vibration to the feet found that this, by definition, non-fractal and so non-complex signal stabilized sway
(Priplata et al., 2002; Priplata et al., 2003). In finding that greater temporal correlations were associated with
subsequent reductions in sway *SD*, the present results align poorly with the finding that uncorrelated stimulation
reduced sway.

Results have curiously diverged within the same research paradigms in this vein, making matters seem
even more paradoxical. A reanalysis of Priplata et al.'s (2003) data yielded two details (Kelty-Stephen and
Dixon, 2013). Firstly, white-noise stimulation reduced temporal correlations in sway. So, in an unhealthy level of
complexity, decorating overly correlated fluctuations may be a useful clinical strategy. Secondly and less
straightforwardly, this reanalysis showed that white-noise stimulation elicited a stronger sway reduction in
participants exhibiting stronger temporal correlations. In a sense, white-noise stimulation seems to wipe out its
efficacy by counteracting the very conditions of endogenous postural fluctuations that give it a stabilizing effect,
which seemed quite puzzling. This work solves some of this puzzle by finding predictive effects of fluctuation
patterns as evident in the association among concurrent variables for the same postural measurement series. The
significant contributions of examining prior effects and subsequent responses are twofold: first, monofractality
self-corrects, and second, stronger monofractality reduced CoP-*SD*. Together, these two points provide a
framework in which the present findings align neatly with research on white-noise stimulation stabilizing
posture (Priplata et al., 2002; Priplata et al., 2003) and explain how fractal temporal correlations could be
sometimes stabilizing and sometimes destabilizing.

Self-correction of fractal temporal correlations has been the key feature missing from the portrayal of
postural stability until now. Lack of self-correction would mean that temporal correlations and *SD* might push
each other to opposite extremes, subverting postural stability (Chen et al., 2008; Rajachandrakumar et al., 2018).
Posture lacking sufficient variability would be unstable and overly temporally correlated; posture with too much
variability would be unstable and have too little temporal correlations. Hence, understanding self-correction in
temporal correlations might then be one of the key directions for future work. Intraindividual differences in
temporal correlations are well-known (Collins and De Luca, 1995, 1993) and often replicated (Gilfriche et al.,
2018). The novelty of this study lies in recognizing that weaker temporal correlations at longer timescales might
hold only on average, consisting of the ebbing and flowing of stronger temporal correlations at shorter
timescales. Across 10-s segments within a single 120-s trial, the vector autoregression found a sequence of
fleeting (e.g., 10-s) bouts of postural sway predicting subsequent alternations between more or less temporal
correlations.

The exact basis of this alternation of temporal correlations warrants further investigation. The question
of whether these spatial constraints govern self-correction of temporal correlations would benefit from rigorous
test in the 'rambling-trembling' framework that recognizes that the fixed reference point anchoring CoP within
the base of support drifts slowly (Zatsiorsky & Duarte, 1999, 2000). Thus, it is important to model the rise and
fall of temporal correlations as a function of the distance between the fixed reference point and the edges of the
base of support (Kelty-Stephen et al., 2020). The space that 'rambling' leaves open for stable, 'trembling' may
govern how quickly temporal correlations alternate. Such modeling could add to the ongoing elaborations of the
rambling-trembling framework to include visual constraints (Ferronato and Barela, 2011; Yamagata et al., 2019).

In summary, the present results offer a glimpse of how current nonlinear dynamical models of self-
correction (Boyd et al., 2017) may play out in biological goal-oriented behavior. Naturally, the correlational
analysis results are prone to mischaracterizing the measured variables as actual causal variables. Fractal
exponents and *SD* are measurements commonly thought to be essential state variables in postural control.
Certainly, fractal exponents are not inherently physiological features but emergent properties of the organism's
interaction with the task. However, the actual postural control parameters may be no less emergent from task
constraints—one of the rare agreements between current cognitivist theorizing about visual attention (Vecera et
al., 2014) and long-standing views in ecological psychology (Bardy et al., 1999).

4.3. Postural control policy through the multifractal lens

4.3.1. Multifractal nonlinearity drives the monofractal results and reveals novel distinction between eyes-closed and 50-cm viewing distance

Our inclusion of multifractal nonlinearity for modeling postural control was devastating to our findings
related to monofractal evidence. No matter the validity of calling sequential monofractal evidence a case of
'multifractality' (Ihlen, 2012), the nonlinearity estimable by comparing multifractal spectra between original and
surrogate series knocked through our monofractal modeling like a meteorite through the statistical roof. Indeed,
from Hypothesis-4 to Hypothesis-5, the differences apparent in impulse-response functions were stark: not one
of the impulse-response functions found for testing up to Hypothesis-4 survived with the inclusion of impulse-
response functions with multifractality nonlinearity. This disparity indicates how much the landscape of modeled
relationships could vary. So, although the emergent control may not have the same labels for its control
parameters, similar observed relationships could explain a host of previous results, as discussed above. Indeed,
multifractal nonlinearity appeared to step into relationships with *SD* that were most similar in form. It predicted
subsequent reductions in *SD* with greater visual-precision demands, and it also reduced following increases in
SD. So, through Hypothesis-4, the effects of monofractal evidence likely carried the deeper effects of yet-
unestimated multifractal nonlinearity.

'Self-correcting' of monofractal evidence was not a feature of monofractality itself but instead reflected
a close relationship between monofractality and multifractal nonlinearity. As we noted above in Section 1.3.2.2.,
monofractal structure and multifractal nonlinearity are mathematically distinguishable but closely related. The
former indicates a correlation across time, and the latter indicates how diversity in this correlation. Now we see
that monofractal evidence's self-correction depends on an ongoing mutual push and pull between monofractality
and multifractality. Because of the nonlinearity inherent in the movement system, changes in monofractal

temporal correlations depend on and reshape the profile of interactions across scales. Nonlinearity supports changes in monofractal scaling, and changes in monofractal scaling could stir up new nonlinearity. This push and pull have two exciting features. First, it appears to be a case of mutual tempering, as the regression model's cubic polynomial gives voice to the oscillatory pattern of impulse-response functions. Second, it diminishes with greater visual precision demands in a suprapostural fixation task. Whether this relationship holds in other task settings is unknown, but it highlights how monofractal evidence from multifractal nonlinearity could be separate variables reflecting distinct dimensions in a parameter space governing goal-directed behavior (Saltzman and Munhall, 1992).

The relationship between monofractality and multifractality is unequal, as multifractality appears to predict monofractality to a greater extent than the extent to which monofractality predicts multifractality. Furthermore, the multifractal model revealed more than the monofractal model about the research design supporting the present dataset. For instance, the monofractal model predicted average changes in subsequent *SD* only in viewing conditions other than the 50-cm viewing condition. Including multifractality reshaped the impulse-response functions to reveal positive effects of monofractal evidence on subsequent *SD* for both the 50-cm viewing and eyes-closed conditions. Besides reshaping the apparent predictive role of monofractal evidence, multifractal nonlinearity also predicted a subsequent increase in *SD* and even estimated unique effects, situating the 50-cm viewing condition as an intermediary between the SD-increasing effect in the eyes-closed condition and *SD*-reductive effect in higher-precision conditions.

These results offer the latest caveat against simple answers to a long-running curiosity for simple resolution of "whether more (multi)fractality is better" in all cases. The capacity of multifractal nonlinearity to predict task-sensitive changes in opposite directions within the same research design is not new (Bell et al., 2019; Mangalam and Kelty-Stephen, 2020). Hence, we see that abruptly changing the support for posture, particularly in removing the task prompt for participants to reach out and invest exploratory behavior beyond the bounds of support, may make multifractal nonlinearity a liability. When the priority is to draw degrees of freedom inward (e.g., to stand as still as possible), then the variability entailed by multifractal nonlinearity becomes a source of greater spatial instability as indicated by *SD*. Certainly, the profiles of a subsequent change in *SD* show the 50-cm viewing condition in between the eyes-closed and higher-precision conditions. This point is reassuring in so far as the 50-cm viewing distance does represent a case where the visible layout beyond the eyelids is available but not demanding exploratory engagement in the way that other viewing distances do. Mere current availability of the visible layout may be enough to coax degrees of freedom 'outward.'

Prior remarks are not a conclusion but a proposal to inquire into the behavioral value of multifractality. What counts as 'inward' and how to chart 'outwardness' becomes an immediate challenge to such reasoning, but it may be a way to begin organizing a set of findings of a deep set of observables. If multifractal nonlinearity is pervasive support for context- and task-sensitive goal-directed behavior, then we need an open mind to the sensitivity of multifractal nonlinearity to various task parameters over different time scales. For instance, consider a seemingly more formal organizing theme of 'perturbation.' Changes in the availability of visual information certainly could be construed as a perturbation, but then quick conclusions that, for example, "perturbations make multifractality worse for motor outcomes" will be too fragile to hold—on a very short, fleeting timescale, multifractal nonlinearity can support a constriction of SD in motor performance—both in past research and in the '2 segments later' predicted value for the present case of our eyes-closed condition.

We expect that the unique rooting of multifractal nonlinearity in interactions across scales makes it critical for task sensitivity. The apparent links between multifractal nonlinearity and performance outcomes are thus likely to change with the task. Indeed, a challenge is that tasks are always deeper than catch-all factors like perturbation. These seemingly task-general terms like perturbation do not speak to the relevance of multiple scales in a seemingly straightforward and simple task. For instance, we recognize that the short timescale of segment and trials in the present reanalysis sets a yet mostly undetermined boundary on how far these exact findings generalize. Not only does multifractality opens up the possibility of variability with time, we know for sure that longer timescales can show different relations between multifractal profiles and behavioral or clinical outcomes (Bell et al., 2019; Ivanov et al., 2001).

Similarly, related to timescales, one can argue that these results do not generalize to situations in which participants are not instructed to minimize sway or fixate gaze. However, we already know from past studies that

multifractal nonlinearity carries significant predictive weight as well when participants can comfortably move as
they prefer in tasks that do not prompt such tight restraint of the degrees of freedom (Carver et al., 2017; Carver
& Kelty-Stephen, 2017; Jacobson et al., 2020; Kelty-Stephen & Dixon, 2014; Stephen et al., 2012; Teng et al.,
2016). The imprint of multifractality on goal-directed behavior may change its form depending on task
constraints (e.g., our reported difference between eyes-closed and 50-cm viewing conditions), but the role of
multifractal nonlinearity will not simply vanish when the participants break out of instructions to restrict
movement.

Contrived as task settings may be, they still constitute a thicket of nesting relationships and relationships
across scales (e.g., Wimsatt, 1994). Our concern is not naturalism (multifractality remains a useful predictor of
goal-directed behavior in outdoor, nonlaboratory settings; Teng et al., 2016). Instead, we are concerned that
scientists are still delving into this issue of many nested constraints in a task—well beyond the explicit
instructions and simple presentations of stimuli. For all the power of controlling task constraints, this control
comes with critical blind spots. Indeed, focusing on a specific task manipulation makes ignoring other factors
almost inevitable. This concern is not new (e.g., Wachtel, 1973), but we suspect that, given its sensitivity to
interactions across scales, multifractal nonlinearity would serve as an essential flashlight into the nestings that
support observed behavior—potentially even into nesting relationships that were not intentionally built into task
settings.

The organism participating in a task setting is no less a causal thicket, and a relatively new direction in
multifractal modeling is to examine how multifractal fluctuations might flow through the body as different parts
differently make contact with different aspects of globally available information. This approach suggests a more
constructive question of "what sort of flow of multifractal fluctuations supports perception?" Through this lens,
there is no real sense in which multifractality is abstractly good or abstractly bad. We already have evidence that
the flow of multifractal fluctuations across the body predicts the accuracy of perceptual judgments (Mangalam et
al., 2020a, 2020b). Of course, we look forward to the eventual question of which kinds of flow are better than
others and for which tasks. Our point, for now, is that multifractality might sooner be a language in which a
bodywide architecture like tensegrity may speak in ways we can listen to and learn from. There might yet be a
reason to consider whether multifractal fluctuations provide a useful domain for synergies whose manifestation
in abundant variability is a good and expectable 'bliss' (Latash, 2020, 2012).

For instance, multifractal nonlinearity vindicates the long-cherished idea of vision as a form of action,
even in this sparse setting. It is worth noting that pre-multifractal modeling of this dataset construed 'visual
precision demand' as being highest for 25-cm and 50-cm viewing conditions (Lee et al., 2019, p. 434).
Multifractal modeling of this data from the perspective of cascading interactions across scales vindicates a
wholly distinct form of visual precision demands, which is the departure of viewing distances in either direction
from a resting point for lens accommodation roughly 50-70 cm before the participants' eyes (Iwasaki and
Tawara, 2002; Korge and Krueger, 1984; Leibowitz and Owens, 1978; Rempel et al., 2007; Sakamoto et al.,
2012). Beyond the idea that vision and action are simply coexisting functionally related processes (Hayhoe,
2017), the multifractal nonlinearity of posture is thus laying bare the bodywide entailments of precision
prompted from the intentional engagement of the lens to fixate at a distance (e.g., Gibson, 1988; Kim & Turvey,
1999). Certainly, ecological approaches respect that whole organisms are worth more than just the lens to muster
in the service of visual perception (Gibson, 1966, 1979). Multifractal nonlinearity offers a quantitative grasp of
how perception-action thrives on interactions between organism-scaled events with lens-scaled events. Indeed,
multifractal nonlinearity may provide a substantial advance beyond the presumed rigor of explanatory appeals to
strictly reductive treatments of physiology, giving force to progress in the directions suggested by ecological
psychology.

The capacity of multifractal nonlinearity as a core feature of the movement system's task-sensitivity
raises an important possibility about which question is more apt. In the first place, authors must present novel
descriptors to layout their plain meaning. So, we have real sympathy with the ongoing challenges in
understanding 'what multifractality means.' But we also see a real benefit in not overpromising in the directions
of how successful or accurate 'more fractal' or 'more multifractal' goal-directed behavior will be. It might be
more constructive to discuss multifractal nonlinearity in conjunction with any other fractal-related descriptors as
quantitative flashlights can bring relationships nested across scales. So long as these relationships might go

unnoticed in an experimental design intended for other purposes, they reflect not just poetic acknowledgments of
other factors not yet explored—but also theoretical questions currently dormant. So, multifractal nonlinearity
may eventually provide a road map for deconstructing our ongoing experimental procedures and learning how to
critically evaluate and then to design nesting relationships for newer research and applications.

4.3.2. Implications for stimulation-based rehabilitation

One of the most accessible potential avenues for application of multifractality is simulation-based
rehabilitation in which the so-called 'noise' is used to nourish the movement system with more of its healthy or
optimal variability (e.g., Cavanaugh et al., 2017). The preceding in Section 4.2. speaks to a discourse on
simulation-based rehabilitation that, to date, remains poised mainly at the monofractal lens (Ducharme & van
Emmerik, 2018; Kelty-Stephen & Dixon, 2013; Priplata et al., 2002; Priplata et al., 2003; Rhea et al., 2014;
Sotirakis et al., 2020; Vaz et al., 2020). It perches at the frontiers of movement science, and its promise lies in
envisioning that multiple monofractal profiles at one time—even if it is just the contrast between two
monofractal profiles: one of the organisms and the other of the stimulation. However, the caveat on the
interpretation of correlational analyses remains in force, as a strictly monofractal discourse risks failing to
acknowledge any of possible multifractality-based variables. This work asserts clearly that, if the movement
system thrives on stochastic fluctuations, then we might need to prescribe variable stimulation with no less
multifractal structure.

The restructuring of our impulse-response functions and the expansion of our nonlinear-dynamical space
for modeling goal-directed behavior resonate with the current calls for new directions in simulation-based
rehabilitation. Recent work has examined the role of the auditory stimulus with elaborations beyond fractal
structure in supporting gait (Hove et al., 2012; Hunt et al., 2014; Rodger and Craig, 2016; Uchitomi et al., 2013).
Despite never explicitly naming 'multifractal' structure as focus, these studies propose to build gait-scaffolding
stimulation from the foundation of biological patterns (e.g., gait and music) that marry multi-scale aspect of
fractal fluctuations with complex phase relationships amongst participating timescales. This blend of features
guarantees multifractality in these structures (Muñoz-Diosdado, 2005; Telesca and Lovallo, 2011). Likewise, if
we study the postural system to identify factors that can minimize sway, it becomes evident that the postural
system itself exhibits a multifractal structure. Indeed, the stochastic resonance that the scientific community has
been trying to build might be more accessible by using multifractal stimulation.

The multifractal reanalysis could inform considerations about how endogenous fluctuations inform
tailoring stimulation for rehabilitation, potentially moving it past the frontiers of monofractal formalism. The
limitation of fractal temporal correlations is that fractal models are all too symmetrical, and postural sway is
known to be intermittent (Lee et al., 2018, 2019). Monofractal stimulation may be ill-suited to the multifractal
landscape of postural control. Future work could benefit from considering what multifractal configurations of
posture will be most receptive to stimulation, and multifractal geometry may offer the plainest framework for
tuning stimuli to multifractal control of posture. In pointing to the importance of variation in fractal temporal
correlations, this work is a step towards understanding the multisensory aspect of posture and providing
multisensory support for its clinical support and rehabilitation.

To sum up, the present findings on quietly standing and fixating situates intermodal use of information
on multifractal foundations. Visually fixating brings a prestressed quiet stance into one with informational
coupling with the visual stimulus. Past work has repeatedly implicated multifractal fluctuations in the head and
upper torso for using visual information to organize action (Bell et al., 2019; Carver et al., 2017; Eddy and
Kelty-Stephen, 2015; Kelty-Stephen and Dixon, 2014; Teng et al., 2016). Visual inspection of impulse-response
plots indicated rare instances of effects from or responses from CoM-H_{fGn} on or to CoP-SD (1 trial in the 50-,
135-, and 305-cm conditions), but the regression modeling indicated no stable relationship. This absence of
significant impulse-response functions for CoM-H_{fGn} on CoP-SD is puzzling. However, past evidence suggests
that fluctuations in the upper body moderate visual information beyond and possibly in collaboration with
microsaccades. CoP and movements of the upper extremities exhibit a close mutual predictive relationship in
fractal and multifractal fluctuations even without involving a significant role of torso fluctuations (Mangalam et
al., 2020a, 2020b). Indeed, should tensegrity-themed metaphors for the movement system be apt (Turvey &
Fonseca, 2014), then we can expect relatively less local relationships, and CoM may be one of the multiple
intermediary links in the anatomical change that need not always participate in controlling posture. Fuller-body

set of measurements may allow clearer portrayal of causal relationships knitting retinal fluctuations with CoP
fluctuations.

1179

5. Conclusions and future directions

*T*his work exemplifies that visual-postural dynamics raises new questions only definable at the
boundaries among physics of complex systems, psychology, and neuroscience (Bronstein, 2019). This work may
inform rehabilitation strategies with clinical populations that benefit from visual control of posture (e.g., patients
with Parkinson's disease). For instance, understanding what form of postural fluctuations support stable visually-
guided posture could clarify what form of variability should be built into visual information presented by head-
mounted displays in therapeutic sessions (Lei et al., 2019) that could fit neatly within decades-long discussions
on when or how stimulation with monofractal-or-not variability might offer useful clinical applications
(Ducharme & van Emmerik, 2018; Kelty-Stephen & Dixon, 2013; Priplata et al., 2002; Priplata et al., 2003;
Rhea et al., 2014; Sotirakis et al., 2020; Vaz et al., 2020).

Recent work has examined the role of the auditory stimulus with elaborations beyond fractal structure in
supporting gait (Hove et al., 2012; Hunt et al., 2014; Rodger and Craig, 2016; Uchitomi et al., 2013). Despite
never explicitly naming 'multifractal' structure as its focus, these studies investigating the auditory stimuli
propose to build gait-scaffolding stimulation from the foundation of biological patterns (e.g., gait and music) that
marry multi-scale aspect of fractal fluctuations with complex phase relationships amongst participating
timescales. This blend of features guarantees that these structures show multifractality (Muñoz-Diosdado, 2005;
Telesca and Lovallo, 2011). Likewise, if we study the postural system to identify factors that can minimize sway,
it becomes evident that the postural system itself exhibits a multifractal structure. Indeed, the stochastic
resonance that the scientific community has been trying to build might be more accessible by using multifractal
stimulation. The limitation of fractal temporal correlations is that fractal models are all too symmetrical, and
postural sway is known to be intermittent (Lee et al., 2018, 2019). Monofractal stimulation may be ill-suited to
the multifractal landscape of postural control. Future work could benefit from considering what multifractal
configurations of posture will be most receptive to stimulation, and multifractal geometry may offer the plainest
framework for tuning stimuli to multifractal control of posture. In pointing to the importance of variation in
fractal temporal correlations, this work is a step towards understanding the multisensory aspect of posture and
providing multisensory support for its clinical support and rehabilitation.

In the longer theoretical run, we aim to align this multifractal perspective with the ecological notion that
the perceiving-acting organism is an integral whole (Gibson, 1966; Stoffregen et al., 2017). We feel that the
tensegrity approaches noted above align with this perspective. Hence, our references to intermodality do not rely
on a deeper premise of partitioning a global array of stimulation into piecemeal Aristotelian senses. Instead, we
envision that multifractal geometry is an apt modeling framework that respects the fundamental continuity of a
measured system. In short, multifractality allows us to model the diversity of the movement system as it
encounters a multifaceted layout of surfaces, and while our experimental design adds or subtracts a manipulated
factor (e.g., visual precision demands or closing eyes), multifractal modeling allows investigating the measured
behavior without assuming that the measured behavior is itself the addition or subtraction of causal factors. On
the contrary, any seeming 'subsystems' within the perception-action system (e.g., eyes and hands for visual and
haptic subsystems, respectively) might be sooner better understood through the same sort of nesting that links
the whole organism to context. We imagine that different tissues within the body have different forms allowing
different sorts of behaviors, but those different tissues contribute meaning to perception-action through their
nesting in the organism. Hence, the way that a unitary perception-action system finds through the task
environment thrives from nesting relationships within the organism and at larger scales extending beyond the
organism (Kelty-Stephen, 2017). Modeling nesting relationships across many different scales at once is exactly
what multifractal geometry accomplishes, and so it may be uniquely relevant to the goal of the ecological
approaches to perception-action.

References

Bak, P., Tang, C., Wiesenfeld, K., 1987. Self-organized criticality: An explanation of the $1/f$ noise. Phys. Rev. Lett. 59, 381–384. https://doi.org/10.1103/PhysRevLett.59.381

Bardy, B.G., Marin, L., Stoffregen, T.A., Bootsma, R.J., 1999. Postural coordination modes considered as emergent phenomena. J. Exp. Psychol. Hum. Percept. Perform. 25, 1284–1301. https://doi.org/10.1037/0096-1523.25.5.1284

Bashan, A., Bartsch, R., Kantelhardt, J.W., Havlin, S., 2008. Comparison of detrending methods for fluctuation analysis. Phys. A Stat. Mech. its Appl. 387, 5080–5090. https://doi.org/10.1016/j.physa.2008.04.023

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting linear mixed-effects models using lme4. J. Stat. Softw. 1, 1–48. https://doi.org/10.18637/jss.v067.i01

Bell, C., Carver, N., Zbaracki, J., Kelty-Stephen, D., 2019. Nonlinear amplification of variability through interaction across scales supports greater accuracy in manual aiming: Evidence from a multifractal analysis with comparisons to linear surrogates in the Fitts task. Front. Physiol. 10, 998. https://doi.org/10.3389/fphys.2019.00998

Bernstein, N.A., 1967. The Co-ordination and Regulation of Movements. Pergamon Press, New York, NY.

Bles, W., Kapteyn, T.S., Brandt, T., Arnold, F., 1980. The mechanism of physiological height vertigo: II. Posturography. Acta Otolaryngol. 89, 534–540. https://doi.org/10.3109/00016488009127171

Booth, C.R., Brown, H.L., Eason, E.G., Wallot, S., Kelty-Stephen, D.G., 2018. Expectations on hierarchical scales of discourse: Multifractality predicts both short- and long-range effects of violating gender expectations in text reading. Discourse Process. 55, 12–30. https://doi.org/10.1080/0163853X.2016.1197811

Boyd, A.B., Mandal, D., Crutchfield, J.P., 2017. Correlation-powered information engines and the thermodynamics of self-correction. Phys. Rev. E 95, 12152. https://doi.org/10.1103/PhysRevE.95.012152

Bronstein, A.M., 2019. A conceptual model of the visual control of posture. Prog. Brain Res. 248, 285–302. https://doi.org/10.1016/bs.pbr.2019.04.023

Cabe, P.A., 2018. All perception engages the tensegrity-based haptic medium. Ecol. Psychol. 31, 1–13. https://doi.org/10.1080/10407413.2018.1526037

Cachinero-Torre, A., Díaz-Pulido, B., Asúnsolo-del-Barco, Á., 2017. Relationship of the lateral rectus muscle, the supraorbital nerve, and binocular coordination with episodic tension-type headaches frequently associated with visual effort. Pain Med. 18, 969–979. https://doi.org/10.1093/pm/pnw292

Carver, N.S., Bojovic, D., Kelty-Stephen, D.G., 2017. Multifractal foundations of visually-guided aiming and adaptation to prismatic perturbation. Hum. Mov. Sci. 55, 61–72. https://doi.org/10.1016/j.humov.2017.07.005

Carver, N.S., Kelty-Stephen, D.G., 2017. Multifractality in individual honeybee behavior hints at colony-specific social cascades: Reanalysis of radio-frequency identification data from five different colonies. Phys. Rev. E 95, 22402. https://doi.org/10.1103/PhysRevE.95.022402

Cavanaugh, J.T., Kelty-Stephen, D.G., Stergiou, N., 2017. Multifractality, interactivity, and the adaptive capacity of the human movement system: A perspective for advancing the conceptual basis of neurologic physical therapy. J. Neurol. Phys. Ther. 41, 245–251. https://doi.org/10.1097/NPT.0000000000000199

Chen, L.-C., Metcalfe, J.S., Chang, T.-Y., Jeka, J.J., Clark, J.E., 2008. The development of infant upright posture: Sway less or sway differently? Exp. Brain Res. 186, 293–303. https://doi.org/10.1007/s00221-007-1236-1

Chhabra, A., Jensen, R. V, 1989. Direct determination of the $f(\alpha)$ singularity spectrum. Phys. Rev. Lett. 62, 1327–1330. https://doi.org/10.1103/PhysRevLett.62.1327

Collins, J.J., De Luca, C.J., 1995. Upright, correlated random walks: A statistical-biomechanics approach to the human postural control system. Chaos An Interdiscip. J. Nonlinear Sci. 5, 57–63. https://doi.org/10.1063/1.166086

Collins, J.J., De Luca, C.J., 1993. Open-loop and closed-loop control of posture: A random-walk analysis of center-of-pressure trajectories. Exp. Brain Res. 95, 308–318. https://doi.org/10.1007/BF00229788

Doyon, J.K., Hajnal, A., Surber, T., Clark, J.D., Kelty-Stephen, D.G., 2019. Multifractality of posture modulates multisensory perception of stand-on-ability. PLoS One 14, e0212220. https://doi.org/10.1371/journal.pone.0212220

Duarte, M., Sternad, D., 2008. Complexity of human postural control in young and older adults during prolonged standing. Exp. Brain Res. 191, 265–276. https://doi.org/10.1007/s00221-008-1521-7

Duarte, M., Zatsiorsky, V.M., 2001. Long-range correlations in human standing. Phys. Lett. A 283, 124–128. https://doi.org/10.1016/S0375-9601(01)00188-8

Ducharme, S.W., van Emmerik, R.E.A., 2018. Fractal dynamics, variability, and coordination in human locomotion. Kinesiol. Rev. 7, 26–35. https://doi.org/10.1123/kr.2017-0054

Eddy, C.L., Kelty-Stephen, D.G., 2015. Nesting of focal within peripheral vision promotes interactions across nested time scales in head sway: Multifractal evidence from accelerometry during manual and walking-based fitts tasks. Ecol. Psychol. 27, 43–67. https://doi.org/10.1080/10407413.2015.991663

Ferronato, P.A.M., Barela, J.A., 2011. Age-related changes in postural control: Rambling and trembling trajectories. Motor Control 15, 481–493. https://doi.org/10.1123/mcj.15.4.481

Fiorelli, C.M., Polastri, P.F., Rodrigues, S.T., Baptista, A.M., Penedo, T., Pereira, V.A.I., Simieli, L., Barbieri, F.A., 2017. Gaze position interferes in body sway in young adults. Neurosci. Lett. 660, 130–134. https://doi.org/10.1016/j.neulet.2017.09.008

Gibson, E.J., 1988. Exploratory behavior in the development of perceiving, acting, and the acquiring of knowledge. Annu. Rev. Psychol. 39, 1–42. https://doi.org/0.1146/annurev.ps.39.020188.000245

Gibson, J.J., 1979. The Ecological Approach to Visual Perception. Houghton Mifflin, Boston, MA.

Gibson, J.J., 1966. The Senses Considered as Perceptual Systems. Houghton Mifflin, Boston, MA.

Gilden, D.L., 2009. Global model analysis of cognitive variability. Cogn. Sci. 33, 1441–1467. https://doi.org/10.1111/j.1551-6709.2009.01060.x

Gilfriche, P., Deschodt-Arsac, V., Blons, E., Arsac, L.M., 2018. Frequency-specific fractal analysis of postural control accounts for control strategies. Front. Physiol. 9, 293. https://doi.org/10.3389/fphys.2018.00293

Granger, C.W.J., Joyeux, R., 1980. An introduction to long-memory time series models and fractional
differencing. J. Time Ser. Anal. 1, 15–29. https://doi.org/10.1111/j.1467-9892.1980.tb00297.x

Haddad, J.M., Ryu, J.H., Seaman, J.M., Ponto, K.C., 2010. Time-to-contact measures capture modulations in
posture based on the precision demands of a manual task. Gait Posture 32, 592–596.
https://doi.org/10.1016/j.gaitpost.2010.08.008

Haddad, J.M., Van Emmerik, R.E.A., Wheat, J.S., Hamill, J., 2008. Developmental changes in the dynamical
structure of postural sway during a precision fitting task. Exp. Brain Res. 190, 431–441.
https://doi.org/10.1007/s00221-008-1483-9

Hagio, K., Obata, H., Nakazawa, K., 2018. Effects of breathing movement on the reduction of postural sway
during   postural-cognitive   dual   tasking.   PLoS   One   13,   e0197385.
https://doi.org/10.1371/journal.pone.0197385

Hajnal, A., Clark, J.D., Doyon, J.K., Kelty-Stephen, D.G., 2018. Fractality of body movements predicts
perception of affordances: Evidence from stand-on-ability judgments about slopes. J. Exp. Psychol. Hum.
Percept. Perform. 44, 836–841. https://doi.org/10.1037/xhp0000510

Halsey, T.C., Jensen, M.H., Kadanoff, L.P., Procaccia, I., Shraiman, B.I., 1986. Fractal measures and their
singularities: The   characterization   of   strange   sets.   Phys.   Rev.   A   33,   1141–1151.
https://doi.org/10.1103/PhysRevA.33.1141

Hatemi-J, A., 2004. Multivariate tests for autocorrelation in the stable and unstable VAR models. Econ. Model.
21, 661–683. https://doi.org/10.1016/j.econmod.2003.09.005

Hayhoe, M.M., 2017. Vision and action. Annu. Rev. Vis. Sci. 3, 389–413. https://doi.org/10.1146/annurev-
vision-102016-061437

Hove, M.J., Suzuki, K., Uchitomi, H., Orimo, S., Miyake, Y., 2012. Interactive rhythmic auditory stimulation
reinstates   natural $1/f$ timing   in   gait   of   Parkinson's   patients.   PLoS   One   7,   e32600.
https://doi.org/10.1371/journal.pone.0032600

Hunt, N., McGrath, D., Stergiou, N., 2014. The influence of auditory-motor coupling on fractal dynamics in
human gait. Sci. Rep. 4, 5879. https://doi.org/10.1038/srep05879

Ignaccolo, M., Latka, M., West, B.J., 2010. Detrended fluctuation analysis of scaling crossover effects.
Europhys. Lett. 90, 10009. https://doi.org/10.1209/0295-5075/90/10009

Ihlen, E., 2012. Introduction to multifractal detrended fluctuation analysis in Matlab. Front. Physiol. 3, 141.
https://doi.org/10.3389/fphys.2012.00

Ihlen, E.A.F., Vereijken, B., 2010. Interaction-dominant dynamics in human cognition: Beyond $1/f$ fluctuation. J.
Exp. Psychol. Gen. 139, 436–463. https://doi.org/10.1037/a0019098

Ingber, D.E., 2010. From cellular mechanotransduction to biologically inspired engineering. Ann. Biomed. Eng.
38, 1148–1161. https://doi.org/10.1007/s10439-010-9946-0

Ingber, D.E., 2008. Tensegrity-based mechanosensing from macro to micro. Prog. Biophys. Mol. Biol. 97, 163–
179. https://doi.org/10.1016/j.pbiomolbio.2008.02.005

Ingber, D.E., 2006. Cellular mechanotransduction: Putting all the pieces together again. FASEB J. 20, 811–827.
https://doi.org/10.1096/fj.05-5424rev

Ivanov, P.C., Nunes Amaral, L.A., Goldberger, A.L., Havlin, S., Rosenblum, M.G., Stanley, H.E., Struzik, Z.R., 2001. From 1/f noise to multifractal cascades in heartbeat dynamics. Chaos An Interdiscip. J. Nonlinear Sci. 11, 641–652. https://doi.org/10.1063/1.1395631

Iwasaki, T., Tawara, A., 2002. Effects of viewing distance on accommodative and pupillary responses following a three-dimensional task. Ophthalmic Physiol. Opt. 22, 113–118. https://doi.org/10.1046/j.1475-1313.2002.00014.x

Jacobson, N., Berleman-Paul, Q., Mangalam, M., Kelty-Stephen, D.G., 2020. Multifractality in postural sway supports quiet eye training in aiming tasks: A study of golf putting. bioRxiv 258053. https://doi.org/10.1101/2020.08.19.258053

Jaschinski-Kruza, W., 1991. Eyestrain in VDU users: Viewing distance and the resting position of ocular muscles. Hum. Factors 33, 69–83. https://doi.org/10.1177/001872089103300106

Jaschinski, W., 2002. The proximity-fixation-disparity curve and the preferred viewing distance at a visual display as an indicator of near vision fatigue. Optom. Vis. Sci. 79, 158–168.

Jensen, H.J., 1998. Self-Organized Criticality: Emergent Complex Behavior in Physical and Biological Systems. Cambridge university press, Cambridge, UK.

Johnston, T.D., Turvey, M.T., 1980. A sketch of an ecological metatheory for theories of earning. Psychol. Learn. Motiv. 14, 147–205. https://doi.org/10.1016/S0079-7421(08)60161-9

Kaldy, Z., Blaser, E., 2020. Putting effort into infant cognition. Curr. Dir. Psychol. Sci. 29, 180–185. https://doi.org/10.1177/0963721420903015

Kantelhardt, J.W., Zschiegner, S.A., Koscielny-Bunde, E., Havlin, S., Bunde, A., Stanley, H.E., 2002. Multifractal detrended fluctuation analysis of nonstationary time series. Phys. A Stat. Mech. its Appl. 316, 87–114. https://doi.org/10.1016/S0378-4371(02)01383-3

Kelso, S.J.A., 1995. Dynamic Patterns: The Self-Organization of Brain and Behavior. MIT press, Cambridge, MA.

Kelty-Stephen, D.G., 2018. Multifractal evidence of nonlinear interactions stabilizing posture for phasmids in windy conditions: A reanalysis of insect postural-sway data. PLoS One 13, e0202367. https://doi.org/10.1371/journal.pone.0202367

Kelty-Stephen, D.G., 2017. Threading a multifractal social psychology through within-organism coordination to within-group interactions: A tale of coordination in three acts. Chaos, Solitons & Fractals 104, 363–370. https://doi.org/10.1016/j.chaos.2017.08.037

Kelty-Stephen, D.G., Dixon, J.A., 2014. Interwoven fluctuations during intermodal perception: Fractality in head sway supports the use of visual feedback in haptic perceptual judgments by manual wielding. J. Exp. Psychol. Hum. Percept. Perform. 40, 2289–2309. https://doi.org/10.1037/a0038159

Kelty-Stephen, D.G., Dixon, J.A., 2013. Temporal correlations in postural sway moderate effects of stochastic resonance on postural stability. Hum. Mov. Sci. 32, 91–105. https://doi.org/10.1016/j.humov.2012.08.006

Kelty-Stephen, D.G., Furmanek, M.P., Mangalam, M., 2020. Multifractality distinguishes reactive from proactive cascades in postural control. Chaos, Solitons & Fractals 110471. https://doi.org/10.1016/j.chaos.2020.110471

Kelty-Stephen, D.G., Wallot, S., 2017. Multifractality versus (mono-) fractality as evidence of nonlinear interactions across timescales: Disentangling the belief in nonlinearity from the diagnosis of nonlinearity in empirical data. Ecol. Psychol. 29, 259–299. https://doi.org/10.1080/10407413.2017.1368355

Kim, N.-G., Turvey, M.T., 1999. Eye movements and a rule for perceiving direction of heading. Ecol. Psychol. 11, 233–248. https://doi.org/10.1207/s15326969eco1103_3

Ko, J.-H., Newell, K.M., 2016. Aging and the complexity of center of pressure in static and dynamic postural tasks. Neurosci. Lett. 610, 104–109. https://doi.org/10.1016/j.neulet.2015.10.069

Korge, A., Krueger, H., 1984. Influence of edge sharpness on the accommodation of the human eye. Graefe's Arch. Clin. Exp. Ophthalmol. 222, 26–28. https://doi.org/10.1007/BF02133773

Koslucher, F., Munafo, J., Stoffregen, T.A., 2016. Postural sway in men and women during nauseogenic motion of the illuminated environment. Exp. Brain Res. 234, 2709–2720. https://doi.org/10.1007/s00221-016-4675-8

Kuznetsov, N., Wallot, S., 2011. Effects of accuracy feedback on fractal characteristics of time estimation. Front. Integr. Neurosci. 5, 62. https://doi.org/10.3389/fnint.2011.00062

Latash, M.L., 2020. On primitives in motor control. Motor Control 24, 318–346. https://doi.org/10.1123/mc.2019-0099

Latash, M.L., 2012. The bliss (not the problem) of motor abundance (not redundancy). Exp. Brain Res. 217, 1–5. https://doi.org/10.1007/s00221-012-3000-4

Lê, T.-T., Kapoula, Z., 2008. Role of ocular convergence in the Romberg quotient. Gait Posture 27, 493–500. https://doi.org/10.1016/j.gaitpost.2007.06.003

Lee, D.N., Lishman, J.R., 1975. Visual proprioceptive control of stance. J. Hum. Mov. Stud. 1, 87–95.

Lee, I.-C., Pacheco, M.M., Newell, K.M., 2019a. The precision demands of viewing distance modulate postural coordination and control. Hum. Mov. Sci. 66, 425–439. https://doi.org/10.1016/j.humov.2019.05.019

Lee, I.-C., Pacheco, M.M., Newell, K.M., 2019b. Postural coordination and control to the precision demands of light finger touch. Exp. Brain Res. 237, 1339–1346. https://doi.org/10.1007/s00221-019-05513-2

Lee, I.-C., Pacheco, M.M., Newell, K.M., 2018. Constraints specific influences of vision, touch and surface compliance in postural dynamics. Gait Posture 59, 117–121. https://doi.org/10.1016/j.gaitpost.2017.09.014

Lei, C., Sunzi, K., Dai, F., Liu, X., Wang, Y., Zhang, B., He, L., Ju, M., 2019. Effects of virtual reality rehabilitation training on gait and balance in patients with Parkinson's disease: A systematic review. PLoS One 14, e0224819. https://doi.org/10.1371/journal.pone.0224819

Leibowitz, H.W., Owens, D.A., 1978. New evidence for the intermediate position of relaxed accommodation. Doc. Ophthalmol. 46, 133–147. https://doi.org/10.1007/BF00174103

Likens, A.D., Amazeen, P.G., West, S.G., Gibbons, C.T., 2019. Statistical properties of Multiscale Regression Analysis: Simulation and application to human postural control. Phys. A Stat. Mech. its Appl. 532, 121580. https://doi.org/10.1016/j.physa.2019.121580

Likens, A.D., Fine, J.M., Amazeen, E.L., Amazeen, P.G., 2015. Experimental control of scaling behavior: What is not fractal? Exp. Brain Res. 233, 2813–2821. https://doi.org/10.1007/s00221-015-4351-4

Lipsitz, L.A., 2002. Dynamics of stability: The physiologic basis of functional health and frailty. Journals Gerontol. Ser. A 57, B115–B125. https://doi.org/10.1093/gerona/57.3.B115

Lovejoy, S., 2019. Weather, Macroweather, and the Climate: Our Random Yet Predictable Atmosphere. Oxford University Press, Oxford, UK.

Lutkepohl, H., 2007. New Introduction to Multiple Time Series Analysis. Springer, New York, NY.

Mandelbrot, B.B., 1999. Multifractals and $1/f$ Noise. Springer-Verlag, New York, NY.

Mandelbrot, B.B., 1982. The Fractal Geometry of Nature. W H Freeman, New York, NY.

Mandic, D.P., Chen, M., Gautama, T., Van Hulle, M.M., Constantinides, A., 2008. On the characterization of the deterministic/stochastic and linear/nonlinear nature of time series. Proc. R. Soc. A Math. Phys. Eng. Sci. 464, 1141–1160. https://doi.org/10.1098/rspa.2007.0154

Mangalam, M., Carver, N.S., Kelty-Stephen, D.G., 2020a. Global broadcasting of local fractal fluctuations in a bodywide distributed system supports perception via effortful touch. Chaos, Solitons & Fractals 135, 109740. https://doi.org/10.1016/j.chaos.2020.109740

Mangalam, M., Carver, N.S., Kelty-Stephen, D.G., 2020b. Multifractal signatures of perceptual processing on anatomical sleeves of the human body. J. R. Soc. Interface 17, 20200328. https://doi.org/10.1098/rsif.2020.0328

Mangalam, M., Chen, R., McHugh, T.R., Singh, T., Kelty-Stephen, D.G., 2020c. Bodywide fluctuations support manual exploration: Fractal fluctuations in posture predict perception of heaviness and length via effortful touch by the hand. Hum. Mov. Sci. 69, 102543. https://doi.org/10.1016/j.humov.2019.102543

Mangalam, M., Kelty-Stephen, D.G., 2020. Multiplicative-cascade dynamics supports whole-body coordination for perception via effortful touch. Hum. Mov. Sci. 70, 102595. https://doi.org/10.1016/j.humov.2020.102595

Martinez-Conde, S., Otero-Millan, J., Macknik, S.L., 2013. The impact of microsaccades on vision: Towards a unified theory of saccadic function. Nat. Rev. Neurosci. 14, 83–96. https://doi.org/10.1038/nrn3405

Mayo, A.M., Wade, M.G., Stoffregen, T.A., 2010. Postural effects of the horizon on land and at sea. Psychol. Sci. 22, 118–124. https://doi.org/10.1177/0956797610392927

Meyberg, S., Sinn, P., Engbert, R., Sommer, W., 2017. Revising the link between microsaccades and the spatial cueing of voluntary attention. Vision Res. 133, 47–60. https://doi.org/10.1016/j.visres.2017.01.001

Montesano, G., Crabb, D.P., Jones, P.R., Fogagnolo, P., Digiuni, M., Rossetti, L.M., 2018. Evidence for alterations in fixational eye movements in glaucoma. BMC Ophthalmol. 18, 191. https://doi.org/10.1186/s12886-018-0870-7

Munafo, J., Curry, C., Wade, M.G., Stoffregen, T.A., 2016a. The distance of visual targets affects the spatial magnitude and multifractal scaling of standing body sway in younger and older adults. Exp. Brain Res. 234, 2721–2730. https://doi.org/10.1007/s00221-016-4676-7

Munafo, J., Wade, M.G., Stergiou, N., Stoffregen, T.A., 2016b. The rim and the ancient mariner: The nautical horizon affects postural sway in older adults. PLoS One 11, e0166900. https://doi.org/10.1371/journal.pone.0166900

Muñoz-Diosdado, A., 2005. A non linear analysis of human gait time series based on multifractal analysis and cross correlations. J. Phys. Conf. Ser. 23, 87–95. https://doi.org/10.1088/1742-6596/23/1/010

Murnaghan, C.D., Carpenter, M.G., Chua, R., Inglis, J.T., 2016. Keeping still doesn't "make sense": Examining a role for movement variability by stabilizing the arm during a postural control task. J. Neurophysiol. 117, 846–852. https://doi.org/10.1152/jn.01150.2015

Nelson, C.M., Jean, R.P., Tan, J.L., Liu, W.F., Sniadecki, N.J., Spector, A.A., Chen, C.S., 2005. Emergent patterns of growth controlled by multicellular form and mechanics. Proc. Natl. Acad. Sci. U. S. A. 102, 11594–11599. https://doi.org/10.1073/pnas.0502575102

Newell, K.M., Corcos, Daniel M, 1993. Issues in variability and motor control, in: Newell, K.M., Corcos, D M (Eds.), Variability and Motor Control. Human Kinetics, Champaign, IL, pp. 1–12.

Newell, K.M., Jordan, K., 2007. Task constraints and movement organization: A common language, in: Broadhead, G.D., Davis, W.E. (Eds.), Ecological Task Analysis and Movement. Human Kinetics, Champaign, IL, pp. 5–23.

Palatinus, Z., Dixon, J.A., Kelty-Stephen, D.G., 2013. Fractal fluctuations in quiet standing predict the use of mechanical information for haptic perception. Ann. Biomed. Eng. 41, 1625–1634. https://doi.org/10.1007/s10439-012-0706-1

Palatinus, Z., Kelty-Stephen, D.G., Kinsella-Shaw, J., Carello, C., Turvey, M.T., 2014. Haptic perceptual intent in quiet standing affects multifractal scaling of postural fluctuations. J. Exp. Psychol. Hum. Percept. Perform. 40, 1808–1818. https://doi.org/10.1037/a0037247

Párraga, C.A., Troscianko, T., Tolhurst, D.J., 2002. Spatiochromatic properties of natural images and human vision. Curr. Biol. 12, 483–487. https://doi.org/10.1016/S0960-9822(02)00718-2

Peng, C.-K., Buldyrev, S. V, Havlin, S., Simons, M., Stanley, H.E., Goldberger, A.L., 1994. Mosaic organization of DNA nucleotides. Phys. Rev. E 49, 1685–1689. https://doi.org/10.1103/PhysRevE.49.1685

Peng, C.-K., Havlin, S., Stanley, H.E., Goldberger, A.L., 1995. Quantification of scaling exponents and crossover phenomena in nonstationary heartbeat time series. Chaos An Interdiscip. J. Nonlinear Sci. 5, 82–87. https://doi.org/10.1063/1.166141

Pfaff, B., Stigler, M., Pfaff, M.B., 2018. Package 'vars.' R Packag. version 1.5-3.

Priplata, A., Niemi, J., Salen, M., Harry, J., Lipsitz, L.A., Collins, J.J., 2002. Noise-enhanced human balance control. Phys. Rev. Lett. 89, 238101. https://doi.org/10.1103/PhysRevLett.89.238101

Priplata, A.A., Niemi, J.B., Harry, J.D., Lipsitz, L.A., Collins, J.J., 2003. Vibrating insoles and balance control in elderly people. Lancet 362, 1123–1124. https://doi.org/10.1016/S0140-6736(03)14470-4

Rajachandrakumar, R., Mann, J., Schinkel-Ivy, A., Mansfield, A., 2018. Exploring the relationship between stability and variability of the centre of mass and centre of pressure. Gait Posture 63, 254–259. https://doi.org/10.1016/j.gaitpost.2018.05.008

Rempel, D., Willms, K., Anshel, J., Jaschinski, W., Sheedy, J., 2007. The effects of visual display distance on eye accommodation, head posture, and vision and neck symptoms. Hum. Factors 49, 830–838. https://doi.org/10.1518/001872007X230208

Rhea, C.K., Kiefer, A.W., D'Andrea, S.E., Warren, W.H., Aaron, R.K., 2014. Entrainment to a real time fractal
visual stimulus modulates fractal gait dynamics. Hum. Mov. Sci. 36, 20–34.
https://doi.org/10.1016/j.humov.2014.04.006

Riccio, G.E., 1993. Information in movement variability about the qualitative dynamics of posture and
orientation, in: Variability and Motor Control. Human Kinetics, Champaign, IL, pp. 317–357.

Rodger, M.W.M., Craig, C.M., 2016. Beyond the metronome: Auditory events and music may afford more than
just interval durations as gait cues in Parkinson's disease. Front. Neurosci. 10, 272. https://doi.org/10.3389/
fnins.2016.00272

Sakamoto, K., Asahara, S., Yamashita, K., Okada, A., 2012. Influence of viewing distance and size of TV on
visual fatigue and feeling of involvement. J. Hum. Ergol. (Tokyo). 41, 17–30.
https://doi.org/10.11183/jhe.41.17

Saltzman, E.L., Munhall, K.G., 1992. Skill acquisition and development: The roles of state-, parameter-, and
graph-dynamics. J. Mot. Behav. 24, 49–57. https://doi.org/10.1080/00222895.1992.9941600

Schertzer, D., Lovejoy, S., 2004. Uncertainty and predictability in geophysics: Chaos and multifractal insights.
State Planet Front. Challenges Geophys., Geophysical Monograph Series 317–334.
https://doi.org/10.1029/150GM25

Schleip, R., Mechsner, F., Zorn, A., Klingler, W., 2014. The bodywide fascial network as a sensory organ for
haptic perception. J. Mot. Behav. 46, 191–193. https://doi.org/10.1080/00222895.2014.880306

Shinbrot, T., Muzzio, F.J., 2001. Noise to order. Nature 410, 251–258. https://doi.org/10.1038/35065689

Sims, C.A., 1980. Macroeconomics and reality. Econometrica 48, 1–48. https://doi.org/10.2307/1912017

Singer, J.D., Willett, J.B., 2003. Applied Longitudinal Analysis: Modeling Change and Event Occurrence.
Oxford University Press, New York, NY.

Slifkin, A.B., Newell, K.M., 1999. Noise, information transmission, and force variability. J. Exp. Psychol. Hum.
Percept. Perform. 25, 837–851. https://doi.org/10.1037/0096-1523.25.3.837

Slifkin, A.B., Newell, K.M., 1998. Is variability in human performance a reflection of system noise? Curr. Dir.
Psychol. Sci. 7, 170–177. https://doi.org/10.1111/1467-8721.ep10836906

Sotirakis, H., Patikas, D., Stergiou, N., Hatzitaki, V., 2020. Swaying to the complex motion of a visual target
affects postural sway variability. Gait Posture 77, 125–131. https://doi.org/10.1016/j.gaitpost.2020.01.026

Stephen, D., Anastas, J., Dixon, J., 2012. Scaling in cognitive performance reflects multiplicative multifractal
cascade dynamics. Front. Physiol. 3, 102. https://doi.org/10.3389/fphys.2012.00102

Stephen, D.G., Dixon, J.A., 2011. Strong anticipation: Multifractal cascade dynamics modulate scaling in
synchronization behaviors. Chaos, Solitons & Fractals 44, 160–168.
https://doi.org/10.1016/j.chaos.2011.01.005

Stephen, D.G., Hajnal, A., 2011. Transfer of calibration between hand and foot: Functional equivalence and
fractal fluctuations. Attention, Perception, Psychophys. 73, 1302–1328. https://doi.org/10.3758/s13414-
011-0142-6

Stephen, D.G., Hsu, W.-H., Young, D., Saltzman, E.L., Holt, K.G., Newman, D.J., Weinberg, M., Wood, R.J., Nagpal, R., Goldfield, E.C., 2012. Multifractal fluctuations in joint angles during infant spontaneous kicking reveal multiplicativity-driven coordination. Chaos, Solitons & Fractals 45, 1201–1219. https://doi.org/10.1016/j.chaos.2012.06.005

Stergiou, N., Harbourne, R.T., Cavanaugh, J.T., 2006. Optimal movement variability: A new theoretical perspective for neurologic physical therapy. J. Neurol. Phys. Ther. 30. https://doi.org/10.1097/01.NPT.0000281949.48193.d9

Stoffregen, T.A., 2010. Affordances as properties of the animal-environment system. Ecol. Psychol. 15, 115–134. https://doi.org/10.1207/S15326969ECO1502_2

Stoffregen, T.A., 1986. The role of optical velocity in the control of stance. Percept. Psychophys. 39, 355–360. https://doi.org/10.3758/BF03203004

Stoffregen, T.A., Mantel, B., Bardy, B.G., 2017. The senses considered as one perceptual system. Ecol. Psychol. 29, 165–197. https://doi.org/10.1080/10407413.2017.1331116

Stoffregen, T.A., Pagulayan, R.J., Bardy, B.G., Hettinger, L.J., 2000. Modulating postural control to facilitate visual performance. Hum. Mov. Sci. 19, 203–220. https://doi.org/10.1016/S0167-9457(00)00009-9

Stoffregen, T.A., Smart, L.J., Bardy, B.G., Pagulayan, R.J., 1999. Postural stabilization of looking. J. Exp. Psychol. Hum. Percept. Perform. 25, 1641–1658. https://doi.org/10.1037/0096-1523.25.6.1641

Team, R.C., 2013. R: A language and environment for statistical computing. https://doi.org/URL https://www.R-project.org/.

Telesca, L., Lovallo, M., 2011. Revealing competitive behaviours in music by means of the multifractal detrended fluctuation analysis: Application to Bach's Sinfonias. Proc. R. Soc. A Math. Phys. Eng. Sci. 467, 3022–3032. https://doi.org/10.1098/rspa.2011.0118

Teng, D.W., Eddy, C.L., Kelty-Stephen, D.G., 2016. Non-visually-guided distance perception depends on matching torso fluctuations between training and test. Attention, Perception, Psychophys. 78, 2320–2328. https://doi.org/10.3758/s13414-016-1213-5

Thurner, S., Mittermaier, C., Ehrenberger, K., 2002. Change of complexity patterns in human posture during aging. Audiol. Neurotol. 7, 240–248. https://doi.org/10.1159/000063740

Turing, A.M., 1952. The chemical basis of morphogenesis. Philos. Trans. R. Soc. B Biol. Sci. 237, 37–72.

Turvey, M.T., 2009. On the notion and implications of organism-environment system. Ecol. Psychol. 21, 97–111. https://doi.org/10.1080/10407410902877041

Turvey, M.T., Fitzpatrick, P., 1993. Commentary: Development of perception-action systems and general principles of pattern formation. Child Dev. 64, 1175–1190. https://doi.org/10.1111/j.1467-8624.1993.tb04194.x

Turvey, M.T., Fonseca, S.T., 2014. The medium of haptic perception: A tensegrity hypothesis. J. Mot. Behav. 46, 143–187. https://doi.org/10.1080/00222895.2013.798252

Uchitomi, H., Ota, L., Ogawa, K., Orimo, S., Miyake, Y., 2013. Interactive rhythmic cue facilitates gait relearning in patients with Parkinson's disease. PLoS One 8, e72176. https://doi.org/10.1371/journal.pone.0072176

Vaillancourt, D.E., Newell, K.M., 2002. Changing complexity in human behavior and physiology through aging and disease. Neurobiol. Aging 23, 1–11. https://doi.org/10.1016/S0197-4580(01)00247-0

Vaz, J.R., Rand, T., Fujan-Hansen, J., Mukherjee, M., Stergiou, N., 2020. Auditory and visual external cues have different effects on spatial but similar effects on temporal measures of gait variability. Front. Physiol. 11, 67. https://doi.org/10.3389/fphys.2020.00067

Vecera, S.P., Cosman, J.D., Vatterott, D.B., Roper, Z.J.J., 2014. The control of visual attention: Toward a unified account, in: Ross, B.H. (Ed.), The Psychology of Learning and Motivation. Academic Press, Burlington, MA, pp. 303–347.

Wachtel, P.L., 1973. Psychodynamics, behavior therapy, and the implacable experimenter: An inquiry into the consistency of personality. J. Abnorm. Psychol. 82, 324–334. https://doi.org/10.1037/h0035132

Wagenmakers, E.-J., Farrell, S., Ratcliff, R., 2004. Estimation and interpretation of $1/f^{\alpha}$ noise in human cognition. Psychon. Bull. Rev. 11, 579–615. https://doi.org/10.3758/BF03196615

Wagman, J.B., Caputo, S.E., Stoffregen, T.A., 2016a. Hierarchical nesting of affordances in a tool use task. J. Exp. Psychol. Hum. Percept. Perform. 42, 1627–1642. https://doi.org/10.1037/xhp0000251

Wagman, J.B., Caputo, S.E., Stoffregen, T.A., 2016b. Sensitivity to hierarchical relations among affordances in the assembly of asymmetric tools. Exp. Brain Res. 234, 2923–2933. https://doi.org/10.1007/s00221-016-4695-4

Wagman, J.B., Miller, D.B., 2003. Nested reciprocities: The organism–environment system in perception–action and development. Dev. Psychobiol. 42, 317–334. https://doi.org/10.1002/dev.10114

Wagman, J.B., Stoffregen, T.A., 2020. It doesn't add up: Nested affordances for reaching are perceived as a complex particular. Attention, Perception, Psychophys. 82, 3832–3841. https://doi.org/10.3758/s13414-020-02108-w

Wagman, J.B., Stoffregen, T.A., Bai, J., Schloesser, D.S., 2017. Perceiving nested affordances for another person's actions. Q. J. Exp. Psychol. 71, 790–799. https://doi.org/10.1080/17470218.2016.1277249

Wallot, S., O'Brien, B., Coey, C.A., Kelty-Stephen, D., 2015. Power-law fluctuations in eye movements predict text comprehension during connected text reading, in: Noelle, D.C., Dale, R., Warlaumont, A.S., Yoshimi, J., Matlock, T., Jennings, C.D. (Eds.), Proceedings of the 37th Annual Meeting of the Cognitive Science Society. pp. 2583–2588.

Wang, Y., Bao, J., Ou, L., Thorn, F., Lu, F., 2013. Reading behavior of emmetropic schoolchildren in China. Vision Res. 86, 43–51. https://doi.org/10.1016/j.visres.2013.03.007

Williams, J.D., 1971. A multiple regression approach to multiple comparisons for comparing several treatments with a control. J. Exp. Educ. 39, 93–96.

Wimsatt, W.C., 1994. The ontology of complex systems: Levels of organization, perspectives, and causal thickets. Can. J. Philos. 20, 207–274. https://doi.org/10.1080/00455091.1994.10717400

Winter, D.A., 1995. Human balance and posture control during standing and walking. Gait Posture 3, 193–214. https://doi.org/10.1016/0966-6362(96)82849-9

Yamagata, M., Popow, M., Latash, M.L., 2019. Beyond rambling and trembling: Effects of visual feedback on slow postural drift. Exp. Brain Res. 237, 865–871. https://doi.org/10.1007/s00221-019-05470-w

Zamir, M., 2003. Critique of the test of multifractality as applied to biological data. J. Theor. Biol. 225, 407–412.
https://doi.org/10.1016/S0022-5193(03)00261-3

Zatsiorsky, V., Seluyanov, V., 1985. Estimation of the mass and inertia characteristics of the human body by
means of the best predictive regression equations. Biomechanics IX-B, 233–239.

Zatsiorsky, V.M., Duarte, M., 2000. Rambling and trembling in quiet standing. Motor Control 4, 185–200.
https://doi.org/10.1123/mcj.4.2.185

Zatsiorsky, V.M., Duarte, M., 1999. Instant equilibrium point and its   migration in standing tasks: Rambling
and trembling components of the   stabilogram.   Motor   Control   3,   28–38.
https://doi.org/10.1123/mcj.3.1.28
1605

1606 **Table 1.** Significant baseline effects common to all conditions from mixed-effect regression modeling[a] of
1607 impulse-response functions relating CoM-H_{fGn}, CoP-SD, and CoP-H_{fGn} (see Section 3.1.1.).

Effect	B	SE	t	P[b]
poly(Segment,2)1	-5.25×10^{-1}	2.41×10^{-1}	-2.18	$\mathbf{2.92 \times 10^{-2}}$
Impulse(CoP-H_{fGn})	-6.84×10^{-3}	2.18×10^{-3}	-3.13	$\mathbf{1.73 \times 10^{-3}}$
Trial × poly(Segment,2)1	2.96×10^{-1}	1.11×10^{-1}	2.66	$\mathbf{7.84 \times 10^{-3}}$
poly(Segment,2)1 × Impulse(CoP-H_{fGn})	1.25×10^{0}	3.40×10^{-1}	3.67	$\mathbf{2.43 \times 10^{-4}}$
poly(Segment,2)1 × Response(CoP-H_{fGn})	8.67×10^{-1}	3.40×10^{-1}	2.55	$\mathbf{1.08 \times 10^{-2}}$
Impulse(CoP-H_{fGn}) × Response(CoP-SD)	7.24×10^{-3}	3.10×10^{-3}	2.34	$\mathbf{1.91 \times 10^{-2}}$
Trial × poly(Segment,2)1 × Impulse(CoP-H_{fGn})	-5.47×10^{-1}	1.58×10^{-1}	-3.47	$\mathbf{5.23 \times 10^{-4}}$
Trial × poly(Segment,2)1 × Response(CoP-H_{fGn})	-4.95×10^{-1}	1.58×10^{-1}	-3.14	$\mathbf{1.68 \times 10^{-3}}$
poly(Segment,2)1 × Impulse(CoP-H_{fGn}) × Response(CoP-H_{fGn})	-1.15×10^{0}	4.81×10^{-1}	-2.40	$\mathbf{1.65 \times 10^{-2}}$
poly(Segment,2)1 × Impulse(CoP-H_{fGn}) × Response(CoP-SD)	-1.25×10^{0}	4.81×10^{-1}	-2.60	$\mathbf{9.23 \times 10^{-3}}$
Trial × poly(Segment,2)1 × Impulse(CoP-H_{fGn}) × Response(CoP-H_{fGn})	5.73×10^{-1}	2.23×10^{-1}	2.57	$\mathbf{1.02 \times 10^{-2}}$
Trial × poly(Segment,2)1 × Impulse(CoP-SD) × Response(CoP-H_{fGn})	4.85×10^{-1}	2.23×10^{-1}	2.18	$\mathbf{2.94 \times 10^{-2}}$
Trial × poly(Segment,2)1 × Impulse(CoP-H_{fGn}) × Response(CoP-SD)	5.43×10^{-1}	2.23×10^{-1}	2.44	$\mathbf{1.48 \times 10^{-2}}$

1608 [a]Fitted model: IRF ~ (EC + CVD + DF_CVD)*Trial*poly(segment,2)*Impulse*Response + (1|Participant)
1609 [b]Boldfaced values indicate statistical significance at the the alpha level of 0.05.

1610 **Table 2.** Significant effects of Eyes Closed (EC) from mixed-effect regression modeling[a] of impulse-response
1611 functions relating CoM-H_{fGn}, CoP-SD, and CoP-H_{fGn} (see Section 3.1.2.).

Effect	*B*	*SE*	*t*	*P*[b]
EC × poly(Segment,2)2	-9.57×10^{-1}	3.56×10^{-1}	-2.69	**7.16×10^{-3}**
EC × Impulse(CoP-H_{fGn})	7.06×10^{-3}	3.23×10^{-3}	2.19	**2.88×10^{-2}**
EC × Trial × poly(Segment,2)2	5.46×10^{-1}	1.65×10^{-1}	3.31	**9.23×10^{-4}**
EC × poly(Segment,2)1 × Impulse(CoP-H_{fGn})	-1.95×10^{0}	5.03×10^{-1}	-3.88	**1.04×10^{-4}**
EC × poly(Segment,2)2 × Impulse(CoP-H_{fGn})	1.63×10^{0}	5.03×10^{-1}	3.24	**1.21×10^{-3}**
EC × poly(Segment,2)2 × Impulse(CoP-SD)	1.15×10^{0}	5.03×10^{-1}	2.28	**2.27×10^{-2}**
EC × Trial × poly(Segment,2)1 × Impulse(CoP-H_{fGn})	1.03×10^{0}	2.33×10^{-1}	4.43	**9.37×10^{-6}**
EC × Trial × poly(Segment,2)2 × Impulse(CoP-H_{fGn})	-1.05×10^{0}	2.33×10^{-1}	-4.51	**6.51×10^{-6}**
EC × Trial × poly(Segment,2)2 × Impulse(CoP-SD)	-6.69×10^{-1}	2.33×10^{-1}	-2.87	**4.10×10^{-3}**
EC × Trial × poly(Segment,2)2 × Response(CoP-H_{fGn})	-5.25×10^{-1}	2.33×10^{-1}	-2.25	**2.42×10^{-2}**
EC × Trial × poly(Segment,2)2 × Response(CoP-SD)	-5.50×10^{-1}	2.33×10^{-1}	-2.36	**1.83×10^{-2}**
EC × poly(Segment,2)1 × Impulse(CoP-H_{fGn}) × Response(CoP-H_{fGn})	1.72×10^{0}	7.12×10^{-1}	2.41	**1.58×10^{-2}**
EC × poly(Segment,2)2 × Impulse(CoP-H_{fGn}) × Response(CoP-H_{fGn})	-1.74×10^{0}	7.12×10^{-1}	-2.45	**1.45×10^{-2}**
EC × poly(Segment,2)1 × Impulse(CoP-H_{fGn}) × Response(CoP-SD)	1.97×10^{0}	7.12×10^{-1}	2.76	**5.72×10^{-3}**
EC × poly(Segment,2)2 × Impulse(CoP-H_{fGn}) × Response(CoP-SD)	-1.66×10^{0}	7.12×10^{-1}	-2.33	**2.00×10^{-2}**
EC × Trial × poly(Segment,2)1 × Impulse(CoP-H_{fGn}) × Response(CoP-H_{fGn})	-1.13×10^{0}	3.30×10^{-1}	-3.42	**6.31×10^{-4}**
EC × Trial × poly(Segment,2)2 × Impulse(CoP-H_{fGn}) × Response(CoP-H_{fGn})	1.29×10^{0}	3.30×10^{-1}	3.90	**9.60×10^{-5}**
EC × Trial × poly(Segment,2)2 × Impulse(CoP-SD) × Response(CoP-H_{fGn})	7.26×10^{-3}	3.30×10^{-1}	2.20	**2.76×10^{-2}**
EC × Trial × poly(Segment,2)1 × Impulse(CoP-H_{fGn}) × Response(CoP-SD)	-1.02×10^{0}	3.30×10^{-1}	-3.11	**1.89×10^{-3}**
EC × Trial × poly(Segment,2)2 × Impulse(CoP-H_{fGn}) × Response(CoP-SD)	1.06×10^{0}	3.30×10^{-1}	3.22	**1.30×10^{-3}**
EC × Trial × poly(Segment,2)2 × Impulse(CoP-SD) × Response(CoP-SD)	6.77×10^{-1}	3.30×10^{-1}	2.05	**3.99×10^{-2}**

1612 [a]Fitted model: IRF ~ (EC + CVD + DF_CVD)*Trial*poly(segment,2)*Impulse*Response + (1|Participant)
1613 [b]Boldfaced values indicate statistical significance at the the alpha level of 0.05.

1614 **Table 3.** Significant effects of Comfortable Viewing Distance (CVD) from mixed-effect regression modeling[a] of
1615 impulse-response functions relating CoM-H_{fGn}, CoP-SD, and CoP-H_{fGn} (see Section 3.1.3.).

Effect	B	SE	t	P[b]
CVD × poly(Segment,2)1	$1.38×10^0$	$3.56×10^{-1}$	3.88	**$1.05×10^{-4}$**
CVD × poly(Segment,2)2	$-9.42×10^{-1}$	$3.56×10^{-1}$	-2.65	**$8.14×10^{-3}$**
CVD × Impulse(CoP-SD)	$9.30×10^{-3}$	$3.23×10^{-3}$	2.88	**$3.98×10^{-3}$**
CVD × Trial × poly(Segment,2)1	$-7.14×10^{-1}$	$1.65×10^{-1}$	-4.33	**$1.47×10^{-5}$**
CVD × Trial × poly(Segment,2)2	$4.68×10^{-1}$	$1.65×10^{-1}$	2.84	**$4.51×10^{-3}$**
CVD × Trial × Impulse(CoP-H_{fGn})	$4.45×10^{-3}$	$1.49×10^{-3}$	2.98	**$2.91×10^{-3}$**
CVD × poly(Segment,2)1 × Impulse(CoP-H_{fGn})	$-2.76×10^0$	$5.03×10^{-1}$	-5.49	**$4.03×10^{-8}$**
CVD × poly(Segment,2)2 × Impulse(CoP-H_{fGn})	$1.02×10^0$	$5.03×10^{-1}$	2.02	**$4.33×10^{-2}$**
CVD × poly(Segment,2)1 × Impulse(CoP-SD)	$-1.92×10^0$	$5.03×10^{-1}$	-3.81	**$1.38×10^{-4}$**
CVD × poly(Segment,2)2 × Impulse(CoP-SD)	$1.13×10^0$	$5.03×10^{-1}$	2.24	**$2.49×10^{-2}$**
CVD × poly(Segment,2)1 × Response(CoP-H_{fGn})	$-1.37×10^0$	$5.03×10^{-1}$	-2.71	**$6.69×10^{-3}$**
CVD × poly(Segment,2)1 × Response(CoP-SD)	$-1.36×10^0$	$5.03×10^{-1}$	-2.70	**$6.98×10^{-3}$**
CVD × Impulse(CoP-SD) × Response(CoP-H_{fGn})	$-1.03×10^{-2}$	$4.57×10^{-3}$	-2.26	**$2.36×10^{-2}$**
CVD × Impulse(CoP-SD) × Response(CoP-SD)	$-1.00×10^{-2}$	$4.57×10^{-3}$	-2.19	**$2.85×10^{-2}$**
CVD × Trial × poly(Segment,2)1 × Impulse(CoP-H_{fGn})	$1.52×10^0$	$2.33×10^{-1}$	6.52	**$7.05×10^{-11}$**
CVD × Trial × poly(Segment,2)2 × Impulse(CoP-H_{fGn})	$-4.78×10^{-1}$	$2.33×10^{-1}$	-2.05	**$4.01×10^{-2}$**
CVD × Trial × poly(Segment,2)1 × Impulse(CoP-SD)	$8.64×10^{-1}$	$2.33×10^{-1}$	3.71	**$2.09×10^{-4}$**
CVD × Trial × poly(Segment,2)2 × Impulse(CoP-SD)	$-4.81×10^{-1}$	$2.33×10^{-1}$	-2.07	**$3.89×10^{-2}$**
CVD × Trial × poly(Segment,2)1 × Response(CoP-H_{fGn})	$6.67×10^{-1}$	$2.33×10^{-1}$	2.86	**$4.20×10^{-3}$**
CVD × Trial × poly(Segment,2)1 × Response(CoP-SD)	$6.93×10^{-1}$	$2.33×10^{-1}$	2.97	**$2.95×10^{-3}$**
CVD × Trial × poly(Segment,2)2 × Response(CoP-SD)	$-4.61×10^{-1}$	$2.33×10^{-1}$	-1.98	**$4.81×10^{-2}$**
CVD × Trial × Impulse(CoP-H_{fGn}) × Response(CoP-SD)	$-4.22×10^{-3}$	$2.11×10^{-3}$	-2.00	**$4.58×10^{-2}$**
CVD × poly(Segment,2)1 × Impulse(CoP-H_{fGn}) × Response(CoP-H_{fGn})	$1.69×10^0$	$7.12×10^{-1}$	2.38	**$1.75×10^{-2}$**
CVD × poly(Segment,2)1 × Impulse(CoP-SD) × Response(CoP-H_{fGn})	$2.08×10^0$	$7.12×10^{-1}$	2.93	**$3.44×10^{-3}$**
CVD × poly(Segment,2)1 × Impulse(CoP-H_{fGn}) × Response(CoP-SD)	$2.76×10^0$	$7.12×10^{-1}$	3.88	**$1.06×10^{-4}$**
CVD × poly(Segment,2)1 × Impulse(CoP-SD) × Response(CoP-SD)	$1.92×10^0$	$7.12×10^{-1}$	2.69	**$7.14×10^{-3}$**
CVD × Trial × poly(Segment,2)1 × Impulse(CoP-H_{fGn}) × Response(CoP-H_{fGn})	$-8.49×10^{-1}$	$3.29×10^{-1}$	-2.58	**$9.97×10^{-3}$**
CVD × Trial × poly(Segment,2)1 × Impulse(CoP-SD) × Response(CoP-H_{fGn})	$-8.88×10^{-1}$	$3.29×10^{-1}$	-2.69	**$7.06×10^{-3}$**
CVD × Trial × poly(Segment,2)1 × Impulse(CoP-H_{fGn}) × Response(CoP-SD)	$-1.50×10^0$	$3.29×10^{-1}$	-4.55	**$5.43×10^{-3}$**
CVD × Trial × poly(Segment,2)1 × Impulse(CoP-SD) × Response(CoP-H_{fGn})	$-8.56×10^{-1}$	$3.29×10^{-1}$	-2.60	**$9.41×10^{-3}$**

1616 [a]Fitted model: IRF ~ (EC + CVD + DF_CVD)*Trial*poly(segment,2)*Impulse*Response + (1|Participant)
1617 [b]Boldfaced values indicate statistical significance at the the alpha level of 0.05.

1618 **Table 4.** Significant effects of Distance from Comfortable Viewing Distance (DF_CVD) from mixed-effect
1619 regression modeling[a] of impulse-response functions relating CoM-H_{fGn}, CoP-SD, and CoP-H_{fGn} (see Section
1620 3.1.4.).

Effect	B	SE	t	P[b]
Effects of Distance from Comfortable Viewing Distance (DF_CVD)				
DF_CVD × Impulse(CoP-H_{fGn})	5.03×10^{-5}	1.37×10^{-5}	3.68	**2.36×10^{-4}**
DF_CVD × Trial × Impulse(CoP-H_{fGn})	-1.43×10^{-5}	6.34×10^{-6}	−2.26	**2.36×10^{-2}**
DF_CVD × poly(Segment,2)1 × Impulse(CoP-H_{fGn})	-4.91×10^{-3}	2.13×10^{-3}	−2.30	**2.15×10^{-2}**
DF_CVD × Impulse(CoP-H_{fGn}) × Response(CoP-H_{fGn})	-5.21×10^{-5}	1.94×10^{-5}	−2.69	**7.14×10^{-3}**
DF_CVD × Impulse(CoP-H_{fGn}) × Response(CoP-SD)	-5.33×10^{-5}	1.94×10^{-5}	−2.75	**5.88×10^{-3}**
DF_CVD × Trial × poly(Segment,2)1 × Response(CoP-H_{fGn})	1.97×10^{-3}	9.88×10^{-4}	2.00	**4.57×10^{-2}**
DF_CVD × poly(Segment,2)1 × Impulse(CoP-SD) × Response(CoP-H_{fGn})	6.09×10^{-3}	3.02×10^{-3}	2.02	**4.35×10^{-2}**
DF_CVD × Trial × poly(Segment,2)1 × Impulse(CoP-SD) × Response(CoP-H_{fGn})	-2.83×10^{-3}	1.40×10^{-3}	−2.03	**4.28×10^{-2}**

1621 [a]Fitted model: IRF ~ (EC + CVD + DF_CVD)*Trial*poly(segment,2)*Impulse*Response + (1|Participant)
1622 [b]Boldfaced values indicate statistical significance at the the alpha level of 0.05.

Table 5. Significant baseline effects common to all conditions from mixed-effect regression modeling[a] of impulse-response functions incorporating CoP-t_{MF} along with the prior set of CoM-H_{fGn}, CoP-SD, and CoP-H_{fGn} (see Section 3.2.1.). '*' after H_{fGn} and SD indicate the impulse-response functions from the second vector autoregression.

Effect	B	SE	t	P[b]
Impulse(CoP-H_{fGn}*)	1.24×10^0	3.84×10^{-1}	3.22	**1.28×10^{-3}**
poly(Segment, 9)1 × Impulse(CoP-H_{fGn}*)	-1.77×10^2	8.05×10^1	−2.20	**2.80×10^{-2}**
poly(Segment, 9)3 × Impulse(CoP-H_{fGn}*)	-1.68×10^2	8.05×10^1	−2.09	**3.70×10^{-2}**
poly(Segment, 9)5 × Impulse(CoP-H_{fGn}*)	-2.17×10^2	8.05×10^1	−2.70	**7.03×10^{-3}**
poly(Segment, 9)7 × Impulse(CoP-H_{fGn}*)	-2.22×10^2	8.05×10^1	−2.75	**5.92×10^{-3}**
poly(Segment, 9)9 × Impulse(CoP-H_{fGn}*)	-2.72×10^2	8.05×10^1	−3.38	**7.33×10^{-4}**
Impulse(CoP-H_{fGn}*) × Response(CoP_H_{fGn})	-1.25×10^0	4.52×10^{-1}	−2.77	**5.58×10^{-3}**
poly(Segment, 9)5 × Impulse(CoP-H_{fGn}*) × Response(CoP-H_{fGn})	2.30×10^2	9.86×10^1	2.33	**1.99×10^{-2}**
poly(Segment, 9)7 × Impulse(CoP-H_{fGn}*) × Response(CoP-H_{fGn})	2.29×10^2	9.86×10^1	2.33	**2.01×10^{-2}**
poly(Segment, 9)9 × Impulse(CoP-H_{fGn}*) × Response(CoP-H_{fGn})	2.76×10^2	9.86×10^1	2.80	**5.14×10^{-3}**
Impulse(CoP-H_{fGn}*) × Response(CoP-SD)	-1.24×10^0	4.52×10^{-1}	−2.74	**6.16×10^{-3}**
poly(Segment, 9)5 × Impulse(CoP-H_{fGn}*) × Response(CoP-SD)	2.17×10^2	9.86×10^1	2.20	**2.78×10^{-2}**
poly(Segment, 9)7 × Impulse(CoP-H_{fGn}*) × Response(CoP-SD)	2.22×10^2	9.86×10^1	2.25	**2.47×10^{-2}**
poly(Segment, 9)9 × Impulse(CoP-H_{fGn}*) × Response(CoP-SD)	2.72×10^2	9.86×10^1	2.76	**5.83×10^{-3}**
Impulse(CoP-t_{MF})	9.75×10^{-1}	3.84×10^{-1}	2.54	**1.12×10^{-2}**
poly(Segment, 9)1 × Impulse(CoP-t_{MF})	1.73×10^2	8.05×10^2	2.15	**3.17×10^{-2}**
poly(Segment, 9)2 × Impulse(CoP-t_{MF})	2.03×10^2	8.05×10^2	2.52	**1.19×10^{-2}**
poly(Segment, 9)3 × Impulse(CoP-t_{MF})	4.14×10^2	8.05×10^2	5.14	**2.71×10^{-7}**
poly(Segment, 9)4 × Impulse(CoP-t_{MF})	2.38×10^2	8.05×10^2	2.95	**3.17×10^{-3}**
poly(Segment, 9)5 × Impulse(CoP-t_{MF})	4.73×10^2	8.05×10^2	5.87	**4.41×10^{-9}**
poly(Segment, 9)6 × Impulse(CoP-t_{MF})	2.64×10^2	8.05×10^2	3.28	**1.05×10^{-3}**
poly(Segment, 9)7 × Impulse(CoP-t_{MF})	4.91×10^2	8.05×10^2	6.10	**1.08×10^{-9}**
poly(Segment, 9)8 × Impulse(CoP-t_{MF})	2.53×10^2	8.05×10^2	3.14	**1.68×10^{-3}**
poly(Segment, 9)9 × Impulse(CoP-t_{MF})	5.72×10^2	8.05×10^2	7.11	**1.21×10^{-12}**
Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	-9.74×10^{-1}	4.52×10^{-1}	−2.15	**3.11×10^{-2}**
poly(Segment, 9)2 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	-2.02×10^2	9.86×10^1	−2.05	**4.01×10^{-2}**
poly(Segment, 9)3 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	-4.14×10^2	9.86×10^1	−4.20	**2.70×10^{-5}**
poly(Segment, 9)4 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	-2.37×10^2	9.86×10^1	−2.41	**1.61×10^{-2}**
poly(Segment, 9)5 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	-4.72×10^2	9.86×10^1	−4.79	**1.68×10^{-6}**
poly(Segment, 9)6 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	-2.64×10^2	9.86×10^1	−2.67	**7.51×10^{-3}**
poly(Segment, 9)7 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	-4.91×10^2	9.86×10^1	−4.98	**6.51×10^{-7}**
poly(Segment, 9)8 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	-2.53×10^2	9.86×10^1	−2.56	**1.04×10^{-2}**
poly(Segment, 9)9 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	-5.72×10^2	9.86×10^1	−5.80	**6.78×10^{-9}**
Response(CoP-SD*)	-9.78×10^{-1}	3.63×10^{-1}	−2.70	**7.00×10^{-3}**
poly(Segment, 9)1 × Response(CoP-SD*)	-1.73×10^2	7.87×10^1	−2.20	**2.78×10^{-2}**
poly(Segment, 9)2 × Response(CoP-SD*)	-2.02×10^2	7.87×10^1	−2.58	**9.93×10^{-3}**
poly(Segment, 9)3 × Response(CoP-SD*)	-4.14×10^2	7.87×10^1	−5.27	**1.38×10^{-7}**

poly(Segment, 9)4 × Response(CoP-*SD**)	-2.37×10^{2}	7.87×10^{1}	-3.02	**2.54×10^{-3}**
poly(Segment, 9)5 × Response(CoP-*SD**)	-4.73×10^{2}	7.87×10^{1}	-6.01	**1.83×10^{-9}**
poly(Segment, 9)6 × Response(CoP-*SD**)	-2.64×10^{2}	7.87×10^{1}	-3.35	**7.97×10^{-4}**
poly(Segment, 9)7 × Response(CoP-*SD**)	-4.91×10^{2}	7.87×10^{1}	-6.25	**4.20×10^{-10}**
poly(Segment, 9)8 × Response(CoP-*SD**)	-2.53×10^{2}	7.87×10^{1}	-3.22	**1.30×10^{-3}**
poly(Segment, 9)9 × Response(CoP-*SD**)	-5.73×10^{2}	7.87×10^{1}	-7.28	**3.39×10^{-13}**
Response(CoP-t_{MF})	-5.57×10^{-1}	2.66×10^{-1}	-2.09	**3.64×10^{-2}**
poly(Segment, 9)9 × Response(CoP-t_{MF})	1.15×10^{2}	5.69×10^{1}	2.02	**4.33×10^{-2}**

[a]Fitted model: IRF ~ (EC + CVD + DF_CVD)*poly(segment,9)*Impulse*Response + (1|Participant)

[b]Boldfaced values indicate statistical significance at the the alpha level of 0.05.

Table 6. Significant effects of Distance from Comfortable Viewing Distance (CVD) from mixed-effect
regression modeling[a] of impulse-response functions incorporating CoP-t_{MF} along with the prior set of CoM-H_{fGn},
CoP-SD, and CoP-H_{fGn} (see Section 3.2.4.). '*' after H_{fGn} and SD indicate the impulse-response functions from
the second vector autoregression.

Effect	B	SE	t	P[b]
CVD × poly(Segment, 9)9 × Impulse(CoP-H_{fGn}*)	2.32×10^2	1.15×10^2	2.02	**4.30×10^{-1}**
CVD × poly(Segment, 9)3 × Impulse(CoP-t_{MF})	-3.53×10^2	1.15×10^2	−3.07	**2.12×10^{-2}**
CVD × poly(Segment, 9)4 × Impulse(CoP-t_{MF})	-2.47×10^2	1.15×10^2	−2.15	**3.17×10^{-1}**
CVD × poly(Segment, 9)5 × Impulse(CoP-t_{MF})	-3.70×10^2	1.15×10^2	−3.23	**1.26×10^{-2}**
CVD × poly(Segment, 9)6 × Impulse(CoP-t_{MF})	-2.63×10^2	1.15×10^2	−2.29	**2.22×10^{-1}**
CVD × poly(Segment, 9)7 × Impulse(CoP-t_{MF})	-4.10×10^2	1.15×10^2	−3.57	**3.52×10^{-4}**
CVD × poly(Segment, 9)8 × Impulse(CoP-t_{MF})	-2.33×10^2	1.15×10^2	−2.03	**4.27×10^{-2}**
CVD × poly(Segment, 9)9 × Impulse(CoP-t_{MF})	-4.93×10^2	1.15×10^2	−4.30	**1.74×10^{-5}**
CVD × poly(Segment, 9)3 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	3.53×10^2	1.41×10^2	2.51	**1.21×10^{-2}**
CVD × poly(Segment, 9)5 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	3.70×10^2	1.41×10^2	2.63	**8.50×10^{-3}**
CVD × poly(Segment, 9)7 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	4.10×10^2	1.41×10^2	2.92	**3.55×10^{-3}**
CVD × poly(Segment, 9)9 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	4.93×10^2	1.41×10^2	3.50	**4.58×10^{-4}**
CVD × poly(Segment, 9)3 × Response(CoP-SD*)	3.53×10^2	1.10×10^2	3.22	**1.29×10^{-3}**
CVD × poly(Segment, 9)4 × Response(CoP-SD*)	2.47×10^2	1.10×10^2	2.25	**2.44×10^{-2}**
CVD × poly(Segment, 9)5 × Response(CoP-SD*)	3.70×10^2	1.10×10^2	3.38	**7.34×10^{-4}**
CVD × poly(Segment, 9)6 × Response(CoP-SD*)	2.63×10^2	1.10×10^2	2.40	**1.65×10^{-2}**
CVD × poly(Segment, 9)7 × Response(CoP-SD*)	4.10×10^2	1.10×10^2	3.74	**1.84×10^{-4}**
CVD × poly(Segment, 9)8 × Response(CoP-SD*)	2.33×10^2	1.10×10^2	2.12	**3.37×10^{-2}**
CVD × poly(Segment, 9)9 × Response(CoP-SD*)	4.93×10^2	1.10×10^2	4.50	**6.89×10^{-6}**

1633 [a]Fitted model: IRF ~ (EC + CVD + DF_CVD)*poly(segment,9)*Impulse*Response + (1|Participant)
1634 [b]Boldfaced values indicate statistical significance at the the alpha level of 0.05.

Table 7. Significant effects of Eyes Closed (EC) from mixed-effect regression modeling[a] of impulse-response functions incorporating CoP-t_{MF} along with the prior set of CoM-H_{fGn}, CoP-SD, and CoP-H_{fGn} (see Section 3.2.2.). '*' after H_{fGn} and SD indicate the impulse-response functions from the second vector autoregression.

Effect	B	SE	t	P[b]
DF_CVD × Impulse(CoP-H_{fGn}*)	$7.10×10^{-3}$	$2.29×10^{-3}$	3.09	**$1.99×10^{-3}$**
DF_CVD × poly(Segment, 9)5 × Impulse(CoP-H_{fGn}*)	$1.18×10^{0}$	$5.06×10^{-1}$	2.33	**$1.99×10^{-2}$**
DF_CVD × poly(Segment, 9)7 × Impulse(CoP-H_{fGn}*)	$1.12×10^{0}$	$5.06×10^{-1}$	2.22	**$2.63×10^{-2}$**
DF_CVD × poly(Segment, 9)9 × Impulse(CoP-H_{fGn}*)	$1.40×10^{0}$	$5.06×10^{-1}$	2.77	**$5.63×10^{-3}$**
DF_CVD × Impulse(CoP-H_{fGn}*) × Response(CoP-H_{fGn})	$-7.20×10^{-3}$	$2.81×10^{-3}$	-2.56	**$1.04×10^{-2}$**
DF_CVD × poly(Segment, 9)9 × Impulse(CoP-H_{fGn}*) × Response(CoP-H_{fGn})	$-1.38×10^{0}$	$6.19×10^{-1}$	-2.23	**$2.60×10^{-2}$**
DF_CVD × Impulse(CoP-H_{fGn}*) × Response(CoP-SD)	$-7.09×10^{-3}$	$2.81×10^{-3}$	-2.52	**$1.16×10^{-2}$**
DF_CVD × poly(Segment, 9)9 × Impulse(CoP-H_{fGn}*) × Response(CoP-SD)	$-1.40×10^{0}$	$6.19×10^{-1}$	-2.26	**$2.38×10^{-2}$**
DF_CVD × poly(Segment, 9)1 × Impulse(CoP-t_{MF})	$-1.01×10^{0}$	$5.06×10^{-1}$	-2.00	**$4.60×10^{-2}$**
DF_CVD × poly(Segment, 9)2 × Impulse(CoP-t_{MF})	$-1.20×10^{0}$	$5.06×10^{-1}$	-2.37	**$1.78×10^{-2}$**
DF_CVD × poly(Segment, 9)3 × Impulse(CoP-t_{MF})	$-1.68×10^{0}$	$5.06×10^{-1}$	-3.33	**$8.64×10^{-4}$**
DF_CVD × poly(Segment, 9)4 × Impulse(CoP-t_{MF})	$-1.51×100$	$5.06×10^{-1}$	-2.99	**$2.80×10^{-3}$**
DF_CVD × poly(Segment, 9)5 × Impulse(CoP-t_{MF})	$-2.03×10^{0}$	$5.06×10^{-1}$	-4.02	**$5.79×10^{-5}$**
DF_CVD × poly(Segment, 9)6 × Impulse(CoP-t_{MF})	$-1.57×10^{0}$	$5.06×10^{-1}$	-3.11	**$1.90×10^{-3}$**
DF_CVD × poly(Segment, 9)7 × Impulse(CoP-t_{MF})	$-2.19×10^{0}$	$5.06×10^{-1}$	-4.33	**$1.49×10^{-5}$**
DF_CVD × poly(Segment, 9)8 × Impulse(CoP-t_{MF})	$-1.39×10^{0}$	$5.06×10^{-1}$	-2.75	**$5.87×10^{-3}$**
DF_CVD × poly(Segment, 9)9 × Impulse(CoP-t_{MF})	$-2.64×10^{0}$	$5.06×10^{-1}$	-5.22	**$1.84×10^{-7}$**
DF_CVD × poly(Segment, 9)3 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	$1.68×10^{0}$	$6.19×10^{-1}$	2.72	**$6.56×10^{-3}$**
DF_CVD × poly(Segment, 9)4 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	$1.51×10^{0}$	$6.19×10^{-1}$	2.44	**$1.47×10^{-2}$**
DF_CVD × poly(Segment, 9)5 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	$2.03×10^{0}$	$6.19×10^{-1}$	3.28	**$1.03×10^{-3}$**
DF_CVD × poly(Segment, 9)6 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	$1.57×10^{0}$	$6.19×10^{-1}$	2.53	**$1.13×10^{-2}$**
DF_CVD × poly(Segment, 9)7 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	$2.19×10^{0}$	$6.19×10^{-1}$	3.53	**$4.10×10^{-4}$**
DF_CVD × poly(Segment, 9)8 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	$1.39×10^{0}$	$6.19×10^{-1}$	2.25	**$2.46×10^{-2}$**
DF_CVD × poly(Segment, 9)9 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	$2.64×10^{0}$	$6.19×10^{-1}$	4.26	**$2.09×10^{-5}$**
DF_CVD × poly(Segment, 9)1 × Response(CoP-SD*)	$1.01×10^{0}$	$4.94×10^{-1}$	2.04	**$4.13×10^{-2}$**
DF_CVD × poly(Segment, 9)2 × Response(CoP-SD*)	$1.20×10^{0}$	$4.94×10^{-1}$	2.43	**$1.53×10^{-2}$**
DF_CVD × poly(Segment, 9)3 × Response(CoP-SD*)	$1.68×10^{0}$	$4.94×10^{-1}$	3.41	**$6.56×10^{-4}$**
DF_CVD × poly(Segment, 9)4 × Response(CoP-SD*)	$1.51×10^{0}$	$4.94×10^{-1}$	3.06	**$2.22×10^{-3}$**
DF_CVD × poly(Segment, 9)5 × Response(CoP-SD*)	$2.03×10^{0}$	$4.94×10^{-1}$	4.11	**$3.90×10^{-5}$**
DF_CVD × poly(Segment, 9)6 × Response(CoP-SD*)	$1.57×10^{0}$	$4.94×10^{-1}$	3.18	**$1.48×10^{-3}$**
DF_CVD × poly(Segment, 9)7 × Response(CoP-SD*)	$2.19×10^{0}$	$4.94×10^{-1}$	4.43	**$9.42×10^{-6}$**
DF_CVD × poly(Segment, 9)8 × Response(CoP-SD*)	$1.39×10^{0}$	$4.94×10^{-1}$	2.82	**$4.82×10^{-3}$**
DF_CVD × poly(Segment, 9)9 × Response(CoP-SD*)	$2.64×10^{0}$	$4.94×10^{-1}$	5.34	**$9.52×10^{-8}$**
DF_CVD × Response(CoP-t_{MF})	$-7.30×10^{-3}$	$1.62×10^{-3}$	-4.50	**$6.82×10^{-6}$**

[a]Fitted model: IRF ~ (EC + CVD + DF_CVD)*poly(segment,9)*Impulse*Response + (1|Participant)
[b]Boldfaced values indicate statistical significance at the the alpha level of 0.05.

1640 **Table 8.** Significant effects of Distance from Comfortable Viewing Distance (DF_CVD) from mixed-effect
1641 regression modeling[a] of impulse-response functions incorporating CoP-t_{MF} along with the prior set of CoM-H_{fGn},
1642 CoP-SD, and CoP-H_{fGn} (see Section 3.2.3.). '*' after H_{fGn} and SD indicate the impulse-response functions from
1643 the second vector autoregression.

Effect	B	SE	t	P[b]
EC × Impulse(CoP-H_{fGn}*)	1.75×10^{0}	5.91×10^{-1}	2.95	**3.15×10^{-3}**
EC × poly(Segment, 9)9 × Impulse(CoP-H_{fGn}*)	2.84×10^{2}	1.30×10^{2}	2.18	**2.92×10^{-2}**
EC × Impulse(CoP-H_{fGn}*) × Response(CoP-H_{fGn})	-1.67×10^{0}	7.24×10^{-1}	−2.31	**2.11×10^{-2}**
EC × Impulse(CoP-H_{fGn}*) × Response(CoP-SD)	-1.75×10^{0}	7.24×10^{-1}	−2.41	**1.59×10^{-2}**
EC × Impulse(CoP-t_{MF})	2.26×10^{0}	5.91×10^{-1}	3.83	**1.31×10^{-4}**
EC × poly(Segment, 9)1 × Impulse(CoP-t_{MF})	-3.52×10^{2}	1.30×10^{2}	−2.70	**7.00×10^{-3}**
EC × poly(Segment, 9)2 × Impulse(CoP-t_{MF})	-2.59×10^{2}	1.30×10^{2}	−1.98	**4.74×10^{-2}**
EC × poly(Segment, 9)3 × Impulse(CoP-t_{MF})	-2.82×10^{2}	1.30×10^{2}	−2.17	**3.03×10^{-2}**
EC × poly(Segment, 9)4 × Impulse(CoP-t_{MF})	-3.84×10^{2}	1.30×10^{2}	−2.94	**3.24×10^{-3}**
EC × poly(Segment, 9)5 × Impulse(CoP-t_{MF})	-3.39×10^{2}	1.30×10^{2}	−2.60	**9.34×10^{-3}**
EC × poly(Segment, 9)6 × Impulse(CoP-t_{MF})	-3.59×10^{2}	1.30×10^{2}	−2.75	**5.92×10^{-3}**
EC × poly(Segment, 9)7 × Impulse(CoP-t_{MF})	-4.44×10^{2}	1.30×10^{2}	−3.41	**6.58×10^{-4}**
EC × poly(Segment, 9)8 × Impulse(CoP-t_{MF})	-2.59×10^{2}	1.30×10^{2}	−1.99	**4.71×10^{-2}**
EC × poly(Segment, 9)9 × Impulse(CoP-t_{MF})	-5.62×10^{2}	1.30×10^{2}	−4.31	**1.62×10^{-5}**
EC × Impulse(CoP-t_{MF}) × Response(CoP-t_{MF})	-2.26×10^{0}	7.24×10^{-1}	−3.12	**1.80×10^{-3}**
EC × poly(Segment, 9)1 × Impulse(CoP-t_{MF}) × Response(CoP-t_{MF})	3.51×10^{2}	1.60×10^{2}	2.20	**2.78×10^{-2}**
EC × poly(Segment, 9)4 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	3.83×10^{2}	1.60×10^{2}	2.40	**1.63×10^{-2}**
EC × poly(Segment, 9)5 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	3.39×10^{2}	1.60×10^{2}	1.12	**3.39×10^{-2}**
EC × poly(Segment, 9)6 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	3.59×10^{2}	1.60×10^{2}	2.25	**2.48×10^{-2}**
EC × poly(Segment, 9)7 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	4.44×10^{2}	1.60×10^{2}	2.78	**5.44×10^{-3}**
EC × poly(Segment, 9)9 × Impulse(CoP-t_{MF}) × Response(CoP-H_{fGn})	5.62×10^{2}	1.60×10^{2}	3.52	**4.35×10^{-4}**
EC × Response(CoP-SD*)	-2.26×10^{0}	6.04×10^{-1}	−3.74	**1.86×10^{-4}**
EC × poly(Segment, 9)1 × Response(CoP-SD*)	3.51×10^{2}	1.33×10^{2}	2.64	**8.35×10^{-3}**
EC × poly(Segment, 9)3 × Response(CoP-SD*)	2.83×10^{2}	1.33×10^{2}	2.12	**3.38×10^{-2}**
EC × poly(Segment, 9)4 × Response(CoP-SD*)	3.84×10^{2}	1.33×10^{2}	2.88	**3.96×10^{-3}**
EC × poly(Segment, 9)5 × Response(CoP-SD*)	3.39×10^{2}	1.33×10^{2}	2.54	**1.10×10^{-2}**
EC × poly(Segment, 9)6 × Response(CoP-SD*)	3.59×10^{2}	1.33×10^{2}	2.69	**7.07×10^{-3}**
EC × poly(Segment, 9)7 × Response(CoP-SD*)	4.44×10^{2}	1.33×10^{2}	3.33	**8.54×10^{-4}**
EC × poly(Segment, 9)9 × Response(CoP-SD*)	5.63×10^{2}	1.33×10^{2}	4.22	**2.43×10^{-5}**
EC × Response(CoP-t_{MF})	-8.56×10^{-1}	4.18×10^{-1}	−5.12	**3.13×10^{-7}**

[a]Fitted model: IRF ~ (EC + CVD + DF_CVD)*poly(segment,9)*Impulse*Response + (1|Participant)
[b]Boldfaced values indicate statistical significance at the the alpha level of 0.05.

Figure 1. Schematic of the task and effects of eye-to-target distance on postural sway. (**A**) The suprapostural
viewing task of standing quietly with the eyes fixated at a distant visual element. (**B**, **C**) Visual angle gain for
short vs. long eye-to-target distances along the anterior-posterior (*AP*) and medial-lateral (*ML*) axes. (**D**, **E**)
Visual angle gain as a function of eye-to-target distance for different sway magnitudes. Closer targets increase
AP sway, and farther targets increase *ML* sway.

Figure 2. An overview of detrended fluctuation analysis (DFA). (A, B) CoM and CoP in a representative 10-s segment. (C to F) CoM displacement along the medial-lateral (ML), anterior-posterior (AP), and superior-inferior (SI) axes; CoM spatial Euclidean series; CoP displacement along the ML, AP, and SI axes; CoP planar Euclidean series. (G, H) Log-log plots of fluctuation function, f(N), vs. bin size (N), reflecting the fractal evidence, H_{fGn}, yielded by DFA. Solid circles and solid trend lines represent f(N) for the original (unshuffled) series; open circles and dashed trend lines represent f(N) for a shuffled version of the original series.

Figure 3. Multifractal spectra $(\alpha, f(\alpha))$ for the representative CoM spatial Euclidean series (**A**), and CoP planar
Euclidean series (**B**), from Fig. 2.

Figure 4. An overview of vector autoregressive (VAR) analysis. VAR analysis was used to model the diffusion
of fractal fluctuations across the body, as a time series of segment-by-segment values of CoM-H_{fGn}, CoP-SD, and
CoP-H_{fGn}. (Hypotheses-1–4), as well as CoP-SD, CoP-H_{fGn}, and CoP-t_{MF} (Hypothesis-5). Black arrows indicate
the effects of H_{fGn} in the previous segment on H_{fGn} in the current segment.

Figure 5. Impulse-response functions predicting the responses over ten segments ahead to an impulse in the current segment for each viewing condition. (**A**) CoM-H_{fGn} on CoM-H_{fGn}. (**B**) CoP-H_{fGn} on CoP-H_{fGn}. (**C**) CoM-H_{fGn} on CoP-H_{fGn}.(**D**) CoP-H_{fGn} on CoM-H_{fGn}. Shaded areas indicate $M\pm1SEM$ of all trials across all participants ($N = 15$). Solid circles indicate statistically significant ($P < 0.01$) responses to an impulse in the i^{th} segment. The curves eventually approach zero, indicating that impulse-responses weakened over subsequent segments and eventually diminished completely.

Figure 6. Impulse-response functions predicting the responses over ten segments ahead to an impulse in the current segment for each trial for each viewing condition. (**A**) CoP-H_{fGn} on CoP-SD. (**B**) CoP-SD on CoP-H_{fGn}. Line widths encode trial order (thin: trial-1; medium: trial-2; thick: trial-3). Shaded areas indicate $M\pm1SEM$ of trial averages across all participants ($N = 15$). Solid circles indicate statistically significant ($P < 0.01$) responses to an impulse in the i^{th} segment. The curves eventually approach zero, indicating that impulse-responses weakened over subsequent segments and eventually diminished completely.

Figure 7. Impulse-response functions predicting the responses over ten segments ahead to an impulse in the current segment averaged across all trials for each condition. (**A**) CoP-t_{MF} on CoP-*SD*. In 50-cm viewing condition, a trial with outlier values was substituted by another trial for the same condition, for one participant. (**B**) CoP-*SD* on CoP-t_{MF}. Shaded areas indicate $M \pm 1 SEM$ of all trial across all participants ($N = 45$). Solid circles indicate statistically significant ($P < 0.01$) responses to an impulse in the i^{th} segment. The curves eventually approach zero, indicating that impulse-responses weakened over subsequent segments and eventually diminished completely.

Figure 8. Impulse-response functions predicting the responses in CoP-H_{fGn} over ten segments ahead to an impulse in CoP-t_{MF} the current segment averaged across all trials for each condition. (**A**) CoP-t_{MF} on CoP-*SD*. (**B**) CoP-*SD* on CoP-t_{MF}. In 305-cm viewing condition, a trial with outlier values was substituted by another trial for the same condition, for two participants. Shaded areas indicate $M\pm1SEM$ of all trial across all participants ($N =$ 45). Solid circles indicate statistically significant ($P < 0.01$) responses to an impulse in the i^{th} segment. The curves eventually approach zero, indicating that impulse-responses weakened over subsequent segments and eventually diminished completely.

Figure 9. Impulse-response functions predicting the responses over ten segments ahead to an impulse in the current segment for each viewing condition. (**A**) CoP-t_{MF} on CoP-SD^*. (**B**) CoP-t_{MF} on on CoP-H_{fGn}. '*' after CoP-SD indicates the impulse-response functions from the second vector autoregression (see Section 2.7.1.).

1692 **Figure 10.** Impulse-response functions predicting the responses over ten segments ahead to an impulse in the
1693 current segment for each viewing condition. (**A**) CoP-H_{fGn}* on CoP-t_{MF}. (**B**) CoP-H_{fGn}* on CoP-SD. '*' after
1694 CoP-H_{fGn} indicates the impulse-response functions from the second vector autoregression (see Section 2.7.1.).

1695   **Figure 11.** Impulse-response functions predicting the responses in CoP-t_{MF} over ten segments ahead to an
1696   impulse in CoP-*SD*.