Research article

How many sirtuin genes are out there? evolution of sirtuin genes in vertebrates with a description of a new family member

Juan C. Opazo[1,2,3], Michael W. Vandewege[4], Federico G. Hoffmann[5,6], Kattina Zavala[2], Catalina Meléndez[7], Charlotte Luchsinger[8], Viviana A. Cavieres[7], Luis Vargas-Chacoff[1,9,10,11], Francisco J. Morera[1,12], Patricia V. Burgos[7,13,14], Cheril Tapia-Rojas[7,13], Gonzalo A. Mardones[1,8,15]

[1]Integrative Biology Group, Universidad Austral de Chile, Valdivia, Chile.
[2]Instituto de Ciencias Ambientales y Evolutivas, Facultad de Ciencias, Universidad Austral de Chile, Valdivia, Chile.
[3]Millennium Nucleus of Ion Channel-Associated Diseases (MiNICAD), Valdivia, Chile.
[4]Department of Biology, Eastern New Mexico University, Portales, New Mexico, USA.
[5]Department of Biochemistry, Molecular Biology, Entomology, and Plant Pathology, Mississippi State University, Starkville, Mississippi, USA.
[6]Institute for Genomics, Biocomputing and Biotechnology, Mississippi State University, Starkville, Mississippi, USA.
[7]Centro de Biología Celular y Biomedicina (CEBICEM), Facultad de Medicina y Ciencia, Universidad San Sebastián, Santiago, Chile.
[8]Department of Physiology, School of Medicine, Universidad Austral de Chile, Valdivia, Chile.
[9]Instituto de Ciencias Marinas y Limnológicas, Universidad Austral de Chile, Valdivia, Chile.
[10]Centro Fondap de Investigación de Altas Latitudes (IDEAL), Universidad Austral de Chile, Valdivia, Chile.
[11]Millennium Institute Biodiversity of Antarctic and Subantarctic Ecosystems, BASE, Universidad Austral de Chile, Valdivia, Chile.
[12]Applied Biochemistry Laboratory, Facultad de Ciencias Veterinarias, Instituto de Farmacología y Morfofisiología, Universidad Austral de Chile, Valdivia, Chile.
[13]Centro Ciencia & Vida, Fundación Ciencia & Vida, Santiago, Chile.
[14]Centro de Envejecimiento y Regeneración (CARE-UC), Facultad de Ciencias Biológicas, Pontificia Universidad Católica, Santiago, Chile.
[15]Center for Interdisciplinary Studies of the Nervous System (CISNe), Universidad Austral de Chile, Valdivia, Chile.

Corresponding authors:
Juan C. Opazo, jopazo@gmail.com
Gonzalo A. Mardones, gonzalo.mardones@uach.cl

Abstract

Studying the evolutionary history of gene families is a challenging and exciting task with

a wide range of implications. In addition to exploring fundamental questions about the

origin and evolution of genes, disentangling their evolution is also critical to those who do

functional/structural studies to allow a deeper and more precise interpretation of their

results in an evolutionary context. The sirtuin gene family is a group of genes that are

involved in a variety of biological functions mostly related to aging. Their duplicative

history is an open question, as well as the definition of the repertoire of sirtuin genes

among vertebrates. Our results show a well-resolved phylogeny that represents an

improvement in our understanding of the duplicative history of the sirtuin gene family. We

identified a new sirtuin gene family member (*SIRT8*) that was apparently lost in the last

common ancestor of amniotes, but retained in all other groups of jawed vertebrates.

According to our experimental analyses, elephant shark SIRT8 protein is located in

mitochondria, the overexpression of which leads to an increase in cellular levels of ATP.

Moreover, *in vitro* analysis demonstrated it has deacetylase activity being modulated in a

similar way to mammalian SIRT3. Our results indicate that there are at least eight sirtuin

paralogs among vertebrates and that all of them can be traced back to the last common

ancestor of the group that existed between 676 and 615 millions of years ago.

Keywords: aging, deacetylase, gene family evolution, gene duplication, mitochondria,

SIRT.

Introduction

The availability of whole-genome sequences in species of all main groups of vertebrates represents an opportunity to unravel the evolution of gene families. The number of genomes and their phylogenetic representativeness in the vertebrate tree of life allows performing robust inferences regarding how gene family members are related to each other and their modes of evolution (Nei and Rooney 2005). The available genomes also open an opportunity to discover new gene lineages that are not currently described maybe because they are not present in model species and/or to the absence of appropriate evolutionary analyses (Wichmann et al. 2016; Céspedes et al. 2017; Himmel et al. 2020). Gene copy number variation could be seen as a natural experiment (Albertson et al. 2009) that could help understand the evolutionary fate of duplicated genes, as individuals with different repertoires can, in principle, fulfill the biological functions with a different combination of paralogs (Gitelman 2007).

The sirtuin gene family, class III of histone deacetylase enzymes (HDACs), is a group of genes that, in mammals, is composed of seven paralogs (*SIRT1-7*) grouped into four classes (Fig. 1)(Frye 2000; Frye 2006). Sirtuin genes are involved in a variety of biological functions mostly related to aging, metabolic regulation, stress response, cell cycle among others (Fig. 1) (Michan and Sinclair 2007; Greiss and Gartner 2009; Haigis and Sinclair 2010; Zhao et al. 2019; Zhang et al. 2020). All sirtuin genes have a conserved catalytic domain and variable carboxy- and amino terminal domains. Most family members possess deacetylase activity, while others have, in addition to weak deacetylase activity, other enzymatic activities like ADP-ribosyltransferase, desuccinylase and demalonylase (Fig. 1). Further, they are located in different subcellular compartments and associated to different biological functions (Fig. 1).

The evolution of the sirtuin genes is an open question. There are multiple phylogenetic hypotheses describing evolutionary relationships among the paralogs in the sirtuin gene family (Frye 2000; North and Verdin 2004; Frye 2006; Greiss and Gartner 2009; Pereira et al. 2011; Slade et al. 2011; Vassilopoulos et al. 2011; Costantini et al. 2013; Scholte et al. 2017; Simó-Mirabet et al. 2017; Yang et al. 2017; Rajabi et al. 2018; Kabiljo et al. 2019; Zhao et al. 2019). Differences in the taxonomic sampling, number of paralogs included, and the inclusion of outgroups, are likely contributors to this variation.

Additionally, studies that are focused on resolving evolutionary relationships are scarce,
and in fact, most phylogenetic analyses for sirtuin genes are part of studies where the
sirtuin duplicative history is a secondary goal. Additionally, and probably for similar
reasons, there are no systematic efforts to characterize the full complement of sirtuin
genes among vertebrates. Thus, unraveling the evolutionary history of sirtuin genes
represents a challenging and exciting task with a wide range of implications. Because of
their role in the aging process, these genes are of great interest. In addition to exploring
fundamental questions about the origin and evolution of sirtuin genes, disentangling their
evolution is also critical to understanding the diversification of functional and structural
phenotypes present in the sirtuin gene family.

This study aims to take advantage of the genomic data available in public
databases to advance our understanding of the diversity of vertebrate sirtuin genes and
to infer its duplicative history. We also aim to characterize the subcellular localization,
enzymatic activity, and mitochondrial activity of the newly discovered *SIRT8* gene lineage
in the elephant shark (*Callorhinchus milii*) as a representative species. Our phylogenetic
tree is in general well resolved, recovering vertebrate sirtuin genes into three clades: 1)
SIRT4, *SIRT5*, 2) *SIRT6* and *SIRT7*, and 3) *SIRT1*, *SIRT2*, *SIRT3*, and *SIRT8*. The new
sirtuin family member (*SIRT8*) is retained in all vertebrates other than amniotes (the group
that includes mammals, birds, and reptiles). Based on how sirtuin genes are related to
each other and the information is already known for the other sirtuin family members, we
can predict that SIRT8 belongs to the class I, has a deacetylase activity, and is mainly
located on mitochondria. Our experimental analyses confirmed our evolutionary guided
inferences.

Gene phylogeny	Synteny	Protein Size (AA)	MW (KD)	Enzymatic activity	Subcellular localization	Function	Disease
SIRT1 (Class I)	Human Chr10 DNAJC12 SIRT1 HERC4	747	81.7	Deacetylase	Nucleus & Cytosol	Metabolism regulation, Regulation of chromatin and transcription, DNA repair & Inflammation suppression	Aging, Neurodegenration, Metabolic disease, Cardiovascular disease & Cancer
SIRT2 (Class I)	Human Chr19 RINL SIRT2 NFKBIB	389	41.5	Deacetylase	Nucleus & Cytosol	Cell differentiation, Metabolism regulation, Cell cycle regulation, Microtubule dynamics & Inflammation regulation	Cancer & Neurodegeneration
SIRT3 (Class I)	Human Chr11 RIC8A SIRT3 PSMD13	399	43.6	Deacetylase	Nucleus & Mitochondria	Metabolism regulation, Inflammation suppression, Inhibition of oxidative stress, Apoptosis regulation & Autophagy regulation	Aging, Neurodegenration, Metabolic disease, Cardiovascular disease & Cancer
SIRT8 (Class I)	Elephant shark NW_024704746.1* RIC8B SIRT8 TMEM263	401	44.2	Deacetylase	Mitochondria	?	?
SIRT6 (Class IV)	Human Chr19 CREB3L3 SIRT6 ANKRD24	355	39.1	Deacetylase & ADP-ribosyltransferase	Nucleus	Metabolism regulation & Chromatin and DNA repair	Aging, Metabolic disease, Cardiovascular disease & Cancer
SIRT7 (Class IV)	Human Chr17 PCYT2 SIRT7 MAFG	400	44.9	Deacetylase	Nucleus	Transcription regulation, Chromatin remodeling & Metabolism regulation	Metabolic disease, Cardiovascular disease & Cancer
SIRT4 (Class II)	Human Chr12 PXN SIRT4 PLA2G1B	314	35.2	ADP-ribosyltransferase, Lipoamidase & Deacetylase	Mitochondria	Metabolism regulation	Neurodegenration, Metabolic disease & Cancer
SIRT5 (Class III)	Human Chr6 GFOD1 SIRT5 NOL7	310	33.9	Deacetylase, Desuccinylase, Deglutarylase & Demalonylase	Mitochondria	Metabolism regulation & Immune response modulation	Metabolic disease, & Cancer

Figure 1. Gene phylogeny, synteny, protein size, molecular weight, enzymatic activity, subcellular localization, functions and diseases associated with sirtuin genes. Information regarding the sister group relationships among sirtuin genes was obtained from this study, synteny from ENSEMBL v.106 (Howe et al. 2021), protein size from Fujita and Yamashita (2018)(Fujita and Yamashita 2018), molecular weight from Vassilopoulos et al. (2011)(Vassilopoulos et al. 2011) while enzymatic activity, subcellular localization, functions and diseases from Zhang et al. (2020)(Zhang et al. 2020). In the case of the *SIRT8* of the elephant shark, synteny information was obtained from the National Center for Biotechnology Information (NCBI) (Sharma et al. 2019), protein size, molecular weight, enzymatic activity and subcellular localization from this study.

Results and Discussion

Vertebrate sirtuin paralogs are recovered into two main clades

We recovered the monophyly of all sirtuin family members with strong support (Fig. 2). The diversity of sirtuin genes was arranged into three main clades (Fig. 2). The first clade contains the SIRT4 and SIRT5 paralogs, the second clade contains the SIRT6 and SIRT7 paralogs, and the third clade includes the SIRT1, SIRT2, SIRT3 and SIRT8 gene lineages (Fig. 2). A diversity of phylogenetic arrangements for sirtuin genes have been

proposed in the past (Frye 2000; Frye 2006; Greiss and Gartner 2009; Slade et al. 2011;

Vassilopoulos et al. 2011; Costantini et al. 2013; Scholte et al. 2017; Simó-Mirabet et al.

2017; Yang et al. 2017; Rajabi et al. 2018; Kabiljo et al. 2019; Zhao et al. 2019), and our

results largely support the relationships proposed by Frye (2006).

In the first clade we recovered the sister group relationships between SIRT4 and

SIRT5 with strong support (Fig. 2). Evolutionary relationships between these two family

members are still a matter of debate, as a variety of phylogenetic positions have been

suggested for these paralogs (Slade et al. 2011; Vassilopoulos et al. 2011; Costantini et

al. 2013; Yang et al. 2017; Rajabi et al. 2018), and only in a fraction of the studies their

sister group relationship is supported (Frye 2000; North and Verdin 2004; Frye 2006;

Greiss and Gartner 2009; Hirschey 2011; Scholte et al. 2017; Simó-Mirabet et al. 2017;

Kabiljo et al. 2019; Zhao et al. 2019). In the second clade, we recovered SIRT6 sister to

the clade containing SIRT7 sequences with strong support (Fig. 2). The sister group

relationship between SIRT6 and SIRT7 has been recovered in all examined studies (Frye

2000; North and Verdin 2004; Greiss and Gartner 2009; Hirschey 2011; Slade et al.

2011; Vassilopoulos et al. 2011; Costantini et al. 2013; Scholte et al. 2017; Simó-Mirabet

et al. 2017; Yang et al. 2017; Rajabi et al. 2018; Kabiljo et al. 2019; Zhao et al. 2019),

suggesting there is robust support for the sister relationship between these sirtuin family

members. In the third clade, there is a broad consensus in the literature that SIRT2

shares a common ancestor more recently in time with SIRT3 than with any other sirtuin

paralog, and that the clade containing SIRT1 sequences is sister to the SIRT2/SIRT3

clade (Frye 2000; North and Verdin 2004; Frye 2006; Greiss and Gartner 2009; Hirschey

2011; Slade et al. 2011; Vassilopoulos et al. 2011; Costantini et al. 2013; Scholte et al.

2017; Simó-Mirabet et al. 2017; Yang et al. 2017; Rajabi et al. 2018; Zhao et al. 2019).

We recovered the same evolutionary relationships with strong support (Fig. 2). The

sister-group relationship between the SIRT6/SIRT7 and SIRT2/SIRT3/SIRT8/SIRT1

received moderate support (Fig. 2). The sister-group relationships among the three main

sirtuin clades is something that has been difficult to resolve (Simó-Mirabet et al. 2017;

Zhao et al. 2019), as it appears divergences were close in time, as evidence by the short

length of the corresponding branch (Fig. 2).

In summary, we present a phylogenetic analysis based on a taxonomic sampling that

included representative species from all main groups of vertebrates for all sirtuin family

members. Our phylogenetic tree is in general well resolved (Fig. 2), representing an

advance in our understanding of the duplicative history of the sirtuin gene family. In

comparison to the phylogenetic trees currently available in the literature only one study

shows the same topology as our study (Frye 2006).

Figure 2. Maximum likelihood tree showing sister group relationships among sirtuin genes of vertebrates. Numbers on the nodes correspond to support from the aBayes and ultrafast bootstrap values. Nicotinamide Nucleotide Transhydrogenase (NNT) sequences from the human (*Homo sapiens*) mouse (*Mus musculus*), spotted gar (*Lepisosteus oculatus*) and zebrafish (*Danio rerio*) were used as outgroups (not shown). The scale denotes substitutions per site and colors represent vertebrate sirtuin lineages.

Identification of a new sirtuin gene family member, *SIRT8*

In our analyses, we identified a new sirtuin family member (Fig. 2 and 3), *SIRT8*, which is present in a fraction of the vertebrate tree of life (Fig. 4) that was recovered sister to the *SIRT3* clade with strong support (Fig. 2 and 3). Synteny conservation provides further support to the monophyly of the *SIRT8* gene lineage in gnathostomes (Fig. 5a), as genes found at the 5` side (*POLR3B*, *RFX4* and *RIC8B*) and 3` side (*TMEM263*, *MTERF2*, and *CRY1*) of the *SIRT8* gene are well conserved (Fig. 5a). Synteny is also conserved in species in which the *SIRT8* gene was lost (Fig. 5a). Among vertebrates, we found orthologs of the *SIRT8* gene in representative species of cartilaginous fish, bony fish, coelacanth, and amphibians (Fig. 4). A comparison of the genomic region of the tropical clawed frog (*Xenopus tropicalis*), which possesses the *SIRT8* gene, with the corresponding region in the human (*Homo sapiens*), opossum (*Monodelphis domestica*), chicken (*Gallus gallus*), gharial (*Gavialis gangeticus*), red-eared slider (*Trachemys scripta*) and green anole (*Anolis carolinensis*) strongly suggests that the *SIRT8* gene is not present in the vertebrate lineages that these species represent (Fig. 5b). The absence of the *SIRT8* gene in mammals, birds and reptiles indicates that it was probably lost in the common ancestor of the group, between 352 and 312 millions of years ago (Kumar et al. 2017). In the case of cyclostomes, we found in the sea lamprey (*Petromyzon marinus*) a chromosomal region (Chr3) that possess *TMEM263* and *POLR3B* separated by 44 base pairs, suggesting that all the genes that should be in between (*RFX4*, *RIC8B*, *SIRT8*) were lost in cyclostomes. Alternatively, it is possible that *SIRT8*, and all the other genes are present in cyclostome genomes, but not in the current genome assembly. In the past, Pereira et al. (2011), with a limited taxonomic sampling, identified a clade sister to a group containing *SIRT3* sequences. This gene lineage included sequences from three species with an evolutionary history of whole-genome duplications, two teleost fish (zebrafish, *Danio rerio*, and green spotted puffer, *Tetraodon nigroviridis*), and one

 amphibian (African clawed frog, *Xenopus laevis*), complicating the definition of their

210 duplicative history. Thus, our results, including a balanced taxonomic sampling of all main

211 vertebrate groups and appropriate phylogenetic searches, define the existence of the

212 *SIRT8* gene lineage and its phyletic distribution.

Figure 3. Maximum likelihood tree showing sister group relationships among *SIRT3* and *SIRT8* genes in vertebrates. Numbers on the nodes correspond to support from the aBayes and ultrafast bootstrap values. The scale denotes substitutions per site and colors represent gene lineages. This tree does not represent a novel phylogenetic analysis, it is the *SIRT3*/*SIRT8* clade that was recovered from figure 2.

The lack of the *SIRT8* gene can be interpreted in different ways. We can think that the loss of *SIRT8* could have no physiological impact if the functional role of the lost gene is assumed by other family members, consistent with the idea that gene families possess functional redundancy (Ohno 1985; Félix and Barkoulas 2015; Albalat and Cañestro 2016). Alternatively, we can also think of gene loss as a source of adaptive evolution (Olson 1999; Albalat and Cañestro 2016; Helsen et al. 2020). The description of new gene lineages is not uncommon in the literature (Wichmann et al. 2016; Himmel et al. 2020; Opazo et al. 2021). It could be mainly attributed to the presence in non-model species and/or the absence of proper evolutionary analyses. The existence of species with different gene repertoires is an opportunity to better understand the evolutionary fate of duplicated genes (Lynch and Conery 2000) and the biological functions associated to a group of genes.

Figure 4. Phyletic distribution of sirtuin genes in vertebrates. The sister group relationships among sirtuin genes was obtained from this study, whereas, organismal phylogeny was obtained from the most updated phylogenetic hypotheses available in the literature (Iwabe et al. 2005; Delsuc et al. 2006; Delsuc et al. 2008; Hara et al. 2018). The colors of the circles are those used in figure 2 to define sirtuin gene lineages, whereas the classes to which the different sirtuins belong are shown with shading.

Figure 5. a Patterns of conserved synteny in the chromosomal region that harbor SIRT3-
like genes in gnathostomes. Asterisks denote that the orientation of the genomic piece is
from 3´ to 5´, gray lines represent intervening genes that do not contribute to conserved
synteny. **b** Pairwise dot-plot comparison of the genomic region of the tropical clawed frog
(*Xenopus tropicalis*) with the corresponding region in the human (*Homo sapiens*),
opossum (*Monodelphis domestica*), chicken (*Gallus gallus*), gharial (*Gavialis
gangeticus*), red-eared slider (*Trachemys scripta*) and green anole (*Anolis carolinensis*).
Light green vertical lines denote exons and regions in between are introns. Dot plots were
based on the complete coding region in addition to 6.7 kb of upstream and downstream
flanking sequence. The genome assembly version of the species depicted in this figure
is the following: Human: GRCh38.p13; Opossum: ASM229v1; Chicken: GRCg6a;
Gharial: GavGan_comp1; red-eared slider: CAS_Tse_1.0; Green anole: AnoCar2.0v2;
Tropical clawed frog: UCB_Xtro_10.0; Coelacanth: LatCha1; Spotted gar: LepOcu1 and
Elephant shark: Callorhinchus_milii-6.1.3.

An amino acid alignment of the SIRT8 and SIRT3 sequences shows that the
catalytic domain of SIRT8 is well conserved, where six amino acid positions were
identified as diagnostic characters (Fig. 6). The amino acid divergence of the SIRT8

catalytic domain varies from 32.2% (spotted gar vs coelacanth) to 33.6% (spotted gar vs

elephant shark). The same comparisons for the SIRT3 catalytic domain show slightly

lower amino acid divergence values, between 24.8% (spotted gar vs coelacanth) and

31.0% (coelacanth vs elephant shark). As expected, the interparalog distance (SIRT3 vs

SIRT8) of the catalytic domains shows higher divergence values, ranging from 40.7%

(spotted gar SIRT3 vs spotted gar SIRT8) and 43.6% (elephant shark SIRT3 vs elephant

shark SIRT8). Although nothing is known about the protein encoded by the SIRT8 gene,

based on how sirtuin genes are evolutionarily related and the information already known

for the other family members (Fig. 1), we can speculate that SIRT8 belongs to the class

I, is mainly located on the mitochondria/nucleus, and has deacetylase activity. From a

functional perspective it should perform functions similar to what has been described for

SIRT3 (Fig. 1) (Brown et al. 2013; McDonnell et al. 2015; Zhang et al. 2020). It is important

to highlight that the inferences regarding a newly discovered gene are better performed

if they are phylogenetically guided.

Figure 6. Alignment of the catalytic domain of SIRT3 in humans (*Homo sapiens*), and
SIRT3 and SIRT8 of spotted gar (*Lepisosteus oculatus*), coelacanth (*Latimeria*
chalumnae) and elephant shark (*Callorhinchus milii*). The shaded region denotes the
catalytic domain. Diagnostic characters -i.e. amino acids positions that distinguish
between SIRT3 and SIRT8 gene lineages - are indicated with a rectangle and green
colors.

Expression pattern of sirtuin gene family members

Our next step in characterizing the *SIRT8* gene was to investigate whether it is
transcribed and if it was, to characterize the transcription pattern. To do this, we mapped
RNASeq reads to reference gene sequences of representative species of vertebrates and
examined transcript abundance. In agreement with the transcription pattern reported for
sirtuin genes (Stelzer et al. 2016; Kabiljo et al. 2019), our results show that although they
exhibited wide variance in tissue expression, sirtuin genes are expressed in almost all
tissues, including the novel *SIRT8* gene lineage (Fig. 7). In the case of the tropical clawed
frog, the *SIRT8* gene is transcribed at a similar level in all tissues, other than the ovary
where high transcription levels were estimated (Fig. 7). Coincident with the pattern
observed in the tropical clawed frog, the elephant shark *SIRT8* gene is mostly transcribed
in the ovary (Fig. 7). These results should be taken with caution given the lack of biological
replicates, at least within species. In the case of the zebrafish, although our transcription
level estimations did not follow the trend previously mentioned, an extensive study
examining the expression of all sirtuin genes in zebrafish showed that the *SIRT8* gene in
this species is mainly expressed in the ovary (Pereira et al. 2011). Thus, in addition to
showing that the *SIRT8* gene is transcribed, our analyses allow us to suggest that this
gene could be involved in biological processes associated with reproduction. This
observation agrees with the literature as for SIRT3 there are many studies in which its
functions related to reproduction have been reported (Zhao et al. 2016; Tatone et al. 2018;
Vazquez et al. 2020; Di Emidio et al. 2021).

Figure 7. Heatmap representation of within-species relative transcriptional levels of sirtuin paralogs between 7 chosen tissues. Transcription values were calculated independently for each species and normalized over all tissues.

Elephant shark SIRT8 protein localizes to mitochondria

To further explore the structural relationship between SIRT3 and SIRT8 proteins, we

performed an unbiased protein structure modeling of SIRT8 proteins from elephant shark

(*Callorhinchus milii*), spotted gar (*Lepisosteus oculatus*), salmon (*Salmo salar*) and

coelacanth (*Latimeria chalumnae*). Models of SIRT8 proteins were generated by the

SWISS-MODEL workspace that used as template crystal structure models of human

SIRT3: 4bvh.1.A for elephant shark, 5z94.1.A for spotted gar, 5d7n.3.A for salmon and

5bwo.1.B for coelacanth (data not shown). All models generated comprise only the central

portion of the SIRT8 proteins where the catalytic and its regulatory regions are expected

to be: amino acids G95 to Q365 for elephant shark, L115 to V388 for spotted gar, P114

to G387 for salmon, and K113 to P386 for coelacanth. The overall root mean square

deviation for 272 superimposable Cα coordinates for the models of human SIRT3 (pdb:

4bvh.1.A; amino acids G121 to G392) and elephant shark SIRT8 (amino acids G95 to

Q365) is 0.101 Å (Fig. 8a), suggesting extensive structural match. Some SIRT proteins,

such as SIRT1, SIRT2 and SIRT3, possess evolutionarily conserved, distinct cellular

localizations and functions, as evidenced by their analysis in several model species that

include Drosophila melanogaster, Mus musculus and Homo sapiens (McBurney et al.

2003; Blander and Guarente 2004; North and Verdin 2004; Michishita et al. 2005; Haigis

et al. 2006) (Fig. 1). Some sirtuins localize in more than one compartment, like SIRT1

and SIRT2 that localize to the nucleus and cytosol, and SIRT3 that localizes mainly to the

mitochondrial matrix, but with a small fraction localizing to the nucleus (Scher et al. 2007).

In contrast, SIRT4 and SIRT5 localize only to mitochondria, and SIRT6 and SIRT7

localize only to the nucleus. The sequence and structural comparison between SIRT8

and SIRT3 proteins, as well as subcellular localization prediction by LocTree3 (Goldberg

et al. 2014), suggest that the localization of SIRT8 proteins is also in the mitochondria.

To determine the subcellular localization of SIRT8 proteins, we chose to analyze SIRT8

from elephant shark. To this end, we expressed in human cultured cells full-length

elephant shark SIRT8 tagged with three copies of the c-Myc epitope at the C-terminus

(SIRT8-3myc). To obtain biochemical evidence of the expression of this protein, we

performed immunoblot analysis with monoclonal antibody 9E10 against the c-Myc
epitope, which showed the detection of an expected 48.3 kDa protein in all cell lines
analyzed (Fig. 8b). Fluorescence microscopy analysis of human H4 neuroglioma cells
expressing SIRT8-3myc showed decoration by the anti-c-Myc antibody of structures
highly reminiscent of mitochondria, in addition to a faint fluorescent signal in the rest of
the cytoplasm (Fig. 8c, d). Mitochondria localization of SIRT8-3myc was confirmed by co-
localization with the fluorescent signal of MitoTracker Orange (Pearson's correlation
coefficient [r] = 0.94 ± 0.02; n = 20; Fig. 8c and Supplementary Fig. 1a). Importantly,
SIRT8-3myc showed no localization in any other major compartment, as shown by lack
of localization in the nucleus (detected with the nuclear stain DAPI; r = 0.21 ± 0.03; n =
20; Fig. 8c, d and Supplementary Fig. 1b), the Golgi apparatus (detected with antibody
against Trans-Golgi network integral membrane protein 2 (TGN46) (Ponnambalam et al.
1996) (r = 0.34 ± 0.09; n = 20; Fig. 8c, d and Supplementary Fig. 1c), or the endoplasmic
reticulum (detected with antibody against Cytoskeleton-associated protein 4 (P63)
(Schweizer et al. 1993) (r = 0.47 ± 0.15; n = 20; Fig. 8d and Supplementary Fig. 1d). The
same pattern of expression of SIRT8-3myc was obtained using other human cell lines,
such as HeLa or MDA-MB-231 cells (data not shown). Altogether, these results indicate
that SIRT8 proteins are targeted to the mitochondria.

Figure 8. Structural model of elephant shark SIRT8 and expression of SIRT8-like-3myc in mammalian cells. **a** Superposition of the ribbon representations of elephant shark SIRT8 (green model; amino acids G95 to Q365) and human SIRT3 (magenta model; pdb: 4bvh.1.A; amino acids G121 to G392). Depicted in ellipses are the indicated functional/structural domains of SIRT3 and in dotted lines the binding sites for peptide substrates and NAD$^+$. **b** The indicated cells were left untreated (Control, lanes 1, 3 and 5) or transfected to express SIRT8-3myc (lanes 2, 4 and 6), followed by immunoblot analysis with antibody against the c-Myc epitope or antibody against β-ACTIN used as loading control. The position of molecular mass markers is indicated on the left. SIRT8-3myc exhibits an electrophoretic mobility corresponding to a protein of 48.3 kDa, which is expected considering the extra amino acids of the three copies of the c-Myc epitope at the C-terminus. **c** and **d** H4 cells grown in glass coverslips were transfected to express SIRT8-3myc (green channel) and incubated with the mitochondrial probe MitoTracker Orange (red channel in c) or mock incubated (d). Cells were fixed, permeabilized, and double- (c) or triple-labeled (d) with mouse monoclonal antibody against the c-Myc epitope (c and d), rabbit polyclonal antibody against the endoplasmic reticulum protein Cytoskeleton-associated protein 4 (P63; d) and sheep polyclonal antibody against the Golgi apparatus protein Trans-Golgi network integral membrane protein 2 (TGN46; c and d). Secondary antibodies were Alexa-Fluor-488-conjugated donkey anti-mouse IgG (green channel), Alexa-Fluor-594-conjugated donkey anti-rabbit IgG (red channel), and Alexa-Fluor-647-conjugated donkey anti-sheep IgG (blue channel), and nuclei were

stained with the DNA probe DAPI (white channel). Stained cells were examined by fluorescence microscopy. Insets: X3 magnification with yellow arrows indicating colocalization (c) and green and red arrows indicating lack of colocalization (d). Bar, 10 μm.

Supplementary Figure 1. Colocalization analysis of the immunofluorescence detection of SIRT8-3myc. H4 cells were processed as indicated in Figure legend 8c and 8d. The figure shows a representative colocalization analysis between the fluorescent signals of SIRT8-3myc and the mitochondrial probe MitoTracker Orange (a), SIRT8-3myc and the nuclear probe DAPI (b), SIRT8-3myc and antibody against the Golgi apparatus protein Trans-Golgi network integral membrane protein 2 (TGN46; c), and SIRT8-3myc and

antibody against the endoplasmic reticulum protein Cytoskeleton-associated protein 4
(P63; d). The first column shows the overlapping of corresponding second and third
images in each row. The fourth column shows the corresponding two-dimensional
cytofluorograms of the colocalization analysis using the plugin JACoP implemented in the
software ImageJ (version 1.47h). The corresponding Pearson's correlation coefficient is
indicated in the bottom left corner of the cytofluorograms. Diagonal in a cytofluorogram
indicates high degree of colocalization. Bar, 10 μm.

Elephant shark SIRT8 protein has deacetylase activity

Reversible lysine acetylation is one of the most common post-translational modifications

on proteins that regulate a variety of physiological processes including gene expression,

enzymatic activity, protein-protein interactions, and subcellular localization (Glozak et al.

2005). Sirtuin proteins share a catalytic core domain (North and Verdin 2004) that have

NAD^+-dependent deacetylase activity (Shahgaldi and Kahmini 2021). After translocation

into the mitochondrial matrix, human SIRT3 is inactive, but after proteolytic processing of

its N-terminus signal peptide it becomes active as deacetylase (Onyango et al. 2002;

Schwer et al. 2002). Similar to human SIRT3, secondary structure prediction of

representative SIRT8 proteins indicates that the first ~100 N-terminal amino acids are in

a disordered domain (data not shown). Therefore, to determine if SIRT8 proteins also

exhibit deacetylase activity, we produced in bacteria N-terminal truncated elephant shark

SIRT8 protein (amino acids 95-372; ΔNT-SIRT8). To analyze the enzymatic activity of

purified ΔNT-SIRT8, we used a commercial fluorometric deacetylation activity assay that

contains an acetylated substrate peptide for recombinant human SIRT3 (Abcam). An

initial analysis of activity at 37ºC showed that ΔNT-SIRT8 possesses a deacetylase

activity that is similar to that of human SIRT3, being also NAD^+-dependent (Fig. 9a, e, f).

Because the enzymatic activity of elephant shark SIRT8 could be influenced by the

environmental conditions, we next evaluated the optimal temperature of ΔNT-SIRT8

compared to that of human SIRT3. As expected, the optimal temperature of the human

SIRT3 was ~37ºC (Fig. 9b). In contrast, ΔNT-SIRT8 exhibited an optimal temperature at

~24ºC (Fig. 9b). The optimal temperature value for the SIRT8 deacetylase activity of the

elephant shark could represent an evolutionary adjustment to the temperature regime of

the habitat in which this species lives, as it has been demonstrated for other enzymes in

other species (McCormick 1993; Somero 2004; Bilyk et al. 2021; Saravia et al. 2021).

Thus, all subsequent enzymatic activity analyses for ΔNT-SIRT8 were performed at 24ºC.

The apparent specific activity of ΔNT-SIRT8 was 260.7 ± 9.9 fluorescence units (FU) min^{-1}

mg^{-1}, which was significantly lower than that of SIRT3 that was 1143.3 ± 46.3 FU min^{-1}

mg^{-1}. Apparent K_m of ΔNT-SIRT8 and SIRT3 were 5.7 and 5.5, respectively, which

resulted not significantly different (Fig. 9c, d). In contrast, apparent V_{max} of ΔNT-SIRT3-

like was 0.322 FU sec^{-1}, which was significantly different to that of SIRT3 that was 3.908

FU sec^{-1} (Fig. 9c, d). We next compared the effect of the sirtuin inhibitors nicotinamide

(NAM) (Sauve and Schramm 2003) and quercetin (QUE) (You et al. 2019) on the activities

of SIRT3 and ΔNT-SIRT8. As expected, SIRT3 activity was significantly inhibited by NAM

and QUE (Fig. 9e). We also found a significant inhibitory effect of both NAM and QUE on

the activity of ΔNT-SIRT8 (Fig. 9f). We also compared the effect of the polyphenolic

antioxidant resveratrol (REV), which is an activator of sirtuins (Wood et al. 2004). As

expected, we found a significant concentration-dependent activation of SIRT3

deacetylase activity up to a 1.6 ± 0.1-fold increase in the presence of 100 μM REV (Fig.

9e). Strikingly, we found a much more potent concentration dependent activation effect

of REV on ΔNT-SIRT8, showing up to a 11.1 ± 0.6 fold increase in the presence of 100

μM REV (Fig. 9f). However, because the specific activity of ΔNT-SIRT8 was ~23% that of

SIRT3 in the assay conditions, the activation of ΔNT-SIRT8 by REV in fact represents a

value only ~1.6 fold higher compared to the activation of SIRT3 by REV. Because different

sirtuins have activity on different proteins, as well as remove with different specificity a

variety of acyl groups other than the acetyl (Fig. 1) (Anderson et al. 2014), the modulation

of ΔNT-SIRT8 activity by REV suggests that it has distinct functional substrates. Together,

these results demonstrate that elephant shark ΔNT-SIRT8 has deacetylase activity and

indicate that the activity of SIRT8 proteins is modulated in a similar manner to that of

mammalian sirtuins.

Figure 9. Comparison of deacetylase activity between human SIRT3 and elephant shark ΔNT-SIRT8. a-f) Results obtained with a fluorometric deacetylation assay of an acetylated substrate peptide. a) Time course of deacetylase activity at the indicated concentration of SIRT3 and concentrations of ΔNT-SIRT8. b) Temperature dependence of deacetylase activity. c-d) Michaelis-Menten plot and Lineweaver-Burk plot of SIRT3 (c) and ΔNT-SIRT8 (d) deacetylase activity. e-f) NAD$^+$ dependence and concentration dependent effects of the SIRT3 inhibitors nicotinamide (NAM) and quercetin (QUE) and of the SIRT3 activator resveratrol (REV) on the deacetylase activity of SIRT3 (e) and ΔNT-SIRT8 (f). In a-d, graphs depict the mean ± standard deviation (n = 3). In e-f, bars represent the mean ± standard deviation. Statistical analysis was performed using two-tailed unpaired Student's t-test (n = 3; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; ns, not statistically significant).

Elephant shark SIRT8 protein increases ATP cellular content

Considering that SIRT3 plays an important role in maintaining basal ATP levels by deacetylation of mitochondrial proteins involved in electron transport chain (ETC), the tricarboxylic acid cycle, and fatty-acid oxidation (Hirschey et al. 2010), we investigated the impact of transient SIRT8-3myc expression on ATP cellular content in H4 and HeLa cells (Fig. 10a). Similar to the well-known effect of mammalian SIRT3, SIRT8-3myc expression increased ATP cellular content, suggesting that it shares with SIRT3 the ability to enhance the activity of the mitochondrial ETC. Because an enhancement in the activity of ETC can cause an increase in reactive oxygen species (ROS), we also evaluated the levels of ROS in cells expressing SIRT8-3myc (Fig. 10b). We found similar ROS levels in cells expressing SIRT3-like-3myc and untransfected cells (Fig. 10b). This finding suggests that SIRT8 proteins share the mammalian SIRT3 characteristic of protection against oxidative damage (Bause and Haigis 2013), likely by deacetylation of enzymes involved in ROS clearance to protect mitochondrial membranes from oxidative stress (Tseng et al. 2013). A candidate that could be activated by the deacetylation activity of SIRT3-like proteins is SOD2 that in human cells reduces cellular ROS content promoting oxidative stress resistance (Qiu et al. 2010; Tao et al. 2010). Alternatively, SIRT8 proteins could activate the last complex of the ETC by deacetylation, enhancing ATP-synthase activity without affecting ROS production, similar to the effect that SIRT3 has in response to exercise-induced stress (Vassilopoulos et al. 2014). Together, our results show that the SIRT8 protein is a new sirtuin family member with deacetylase activity that could have important roles in the homeostasis of oxidative species and mitochondrial metabolism.

Figure 10. SIRT8-3myc expression causes an increment on ATP cellular content with no difference in ROS levels. (a) ATP levels were measured by a luciferin/luciferase bioluminescence assay in H4 and HeLa cells either left untreated (Control) or transiently expressing SIRT8-3myc. (b) ROS content was measured by the intensity fluorescence of the dye CM-H2DCFDA in H4 and HeLa cells either left untreated (Control) or transiently expressing SIRT8-3myc. Bars represent the mean ± standard deviation. Statistical analysis was performed using two-tailed unpaired Student's t-test ($n = 4$; ** $p < 0.01$; ns, not statistically significant).

Conclusions

We performed an evolutionary analysis of the sirtuin gene family in vertebrates. In addition to describing the duplicative history of sirtuins, our phylogenetic analyses revealed the presence of a new gene family member, *SIRT8*. This new vertebrate gene lineage is present in all vertebrates other than mammals, birds, and reptiles. Furthermore, our transcriptomic analyses show that *SIRT8* is transcribed in almost all examined tissues, showing a tendency to be more expressed in the ovary, suggesting biological

functions related to reproduction. According to our analyses, SIRT8 localized in the

mitochondria, had deacetylase activity and increased the ATP cellular content. The

finding of a new sirtuin gene lineage highlights the need for more detailed assessments

of orthology with a more broad taxonomic sampling to better define the membership

composition of gene families (Glover et al. 2019). In the literature, there are examples in

which more comprehensive analyses provide a better description of gene families,

including previously unknown family members (Castro et al. 2012; Wichmann et al. 2016;

Céspedes et al. 2017; Ramos-Vicente et al. 2018; Opazo, Kuraku, et al. 2019; Opazo,

Hoffmann, et al. 2019). The availability of species with different gene repertoires could be

seen as a natural experiment (Albertson et al. 2009) that helps us understand the

evolutionary fate of duplicated genes, as they can fulfill the biological functions associated

with the gene family but with a different combination of paralogs.

Methods

Protein sequences and phylogenetic analyses

We retrieved sirtuin amino acid sequences in representative species of all main groups

of vertebrates. Our sampling included mammals, birds, reptiles, amphibians, lungfish,

coelacanth, bony fish, cartilaginous fish, and cyclostomes (Supplementary Table S1).

Sequences were obtained from the Orthologous MAtrix project (OMA), December 2021

release (Altenhoff et al. 2021)last accessed on April 8th, 2022). In addition, we

complemented our sampling by retrieving sequences from vertebrate groups that were

not present or poorly represented in the Orthologous MAtrix project (e.g., crocodiles) from

the National Center for Biotechnology Information (NCBI) (Sharma et al. 2019) using the

program blast (tblastn) with default parameters (Altschul et al. 1990) Supplementary

Table S1). Protein sequences were aligned using the software MAFFT v.7 (Katoh and

Standley 2013), allowing the program to choose the alignment strategy (FFT-NS-i,

alignment length 2490 amino acids). To select the best-fitting model of molecular

evolution we used the proposed model tool in the program IQ-Tree v1.6.12

(Kalyaanamoorthy et al. 2017), which selected JTT+F+I+G4. We used a maximum-

likelihood approach to obtain the best tree using the program IQ-Tree v1.6.12

(Trifinopoulos et al. 2016). We carried out 20 independent runs to explore the tree space

changing the value of the strength of the perturbation (-pers) parameter. We conducted

the following analyses:

1. Five runs modifying the strength of the perturbation parameter from 0.5 (default

537   value) to 0.3.

538   2. Five runs using the default value (0.5) for the strength of the perturbation

539   parameter.

540   3. Five runs modifying the strength of the perturbation parameter from 0.5 (default

541   value) to 0.7.

542   4. Five runs modifying the strength of the perturbation parameter from 0.5 (default

543   value) to 0.9.

In all cases, the number of unsuccessful iterations to stop parameter (-nstop) value was

changed from 100 (default value) to 500. The tree with the highest likelihood score was

chosen (log-likelihood: -127052.963, -pers 0.5 and -nstop 500). Support for the nodes

was evaluated using two approaches: the aBayes test (Anisimova et al. 2011) and the

ultrafast bootstrap procedure using 1000 replicates (Hoang et al. 2018). Nicotinamide

Nucleotide Transhydrogenase (NNT) protein sequences, a member of the DHS-like

NAD/FAD-binding domain superfamily, from the human (*Homo sapiens*), mouse (*Mus*

musculus), zebrafish (*Danio rerio*) and spotted gar (*Lepisosteus oculatus*) were used as

outgroups.

Assessment of conserved synteny

We examined genes found upstream and downstream of the sirtuin genes. For

comparative purposes, we used the estimates of orthology and paralogy derived from the

Ensembl Compara database (Herrero et al. 2016); these estimates are obtained from a

pipeline that considers both synteny and phylogeny to generate orthology mappings.

These predictions were visualized using the program Genomicus v100.01 (Nguyen et al.

2018). Our assessments were performed in humans (*Homo sapiens*), chicken (*Gallus*

gallus), gharial (*Gavialis gangeticus*), red-eared slider (*Trachemys scripta*), green anole

(*Anolis carolinensis*), tropical clawed frog (*Xenopus tropicalis*), coelacanth (*Latimeria*

chalumnae), spotted gar (*Lepisosteus oculatus*), and elephant shark (*Callorhinchus milii*).

Dot-plots

We retrieved the chromosomal region containing the *SIRT8* gene of the tropical clawed frog (*Xenopus tropicalis*, Ch3) including 6.7 kb up- and downstream, and the corresponding syntenic region in the human (*Homo sapiens*, Chr12), opossum (*Monodelphis domestica*, Chr8), chicken (*Gallus gallus*, Chr1), gharial (*Gavialis gangeticus*, NW_017729022.1), red-eared slider (*Trachemys scripta*, Chr1), and green anole (*Anolis carolinensis*, Chr5) based on the location of the flanking genes (*RIC8B* and *TMEM263*). We aligned SIRT3-like syntenic regions using Advanced PipMaker (Schwartz et al. 2000).

Transcript abundance analyses

Sirtuin transcript abundance was measured from a representative sample of vertebrates including the elephant shark (*Callorhinchus milii*), zebrafish (*Danio rerio*), tropical clawed frog (*Xenopus tropicalis*), anole lizard (*Anolis carolinensis*), and human (*Homo sapiens*). RNASeq libraries from brain, heart, kidney, liver, muscle, ovary, and testis from each species were gathered from the NCBI Short Read Archive (SRA; (Leinonen et al. 2011). Accession numbers for species and tissue specific libraries can be found in Supplemental Table S2. Reference transcript sequences were collected from Ensembl v.100 (Yates et al. 2020) and we removed sequences that were shorter than 100 bp. For each library adapters were removed using Trimmomatic 0.38 (Bolger et al. 2014) and reads were filtered for quality using the parameters HEADCROP:5, SLIDINGWINDOW:5:30, and MINLEN:50. We mapped quality filtered paired-end RNAseq reads back to reference sequences using Bowtie 1.2.2 (Langmead et al. 2009) and default parameters of RSEM (Li and Dewey 2011). Transcripts with < 10 mapped reads across all seven tissues per species were removed prior to normalization. Normalization of raw read counts for each species was performed using the median of ratios (Anders and Huber 2010) method implemented by the estimateSizeFactors function in DESeq2 v1.26 (Love et al. 2014). Briefly, for all samples within a species, the geometric mean is calculated for the read counts of each gene. Read counts are then divided by the geometric mean and the median ratio is determined for each sample. Normalized read counts are calculated by dividing raw read counts by the sample-specific median ratio. If multiple SIRT transcripts

596   were present, we presented the expression data from the transcript with the most mapped
597   reads.
598

Protein structure homology modeling

600   Protein structure homology modeling was performed using the SWISS-MODEL server
601   (https://swissmodel.expasy.org/) (Waterhouse et al. 2018). Structural figures were
602   prepared with PyMOL Molecular Graphics System, Version 2.0.6 Schrödinger, LLC.
603

Recombinant cDNA Constructs

For expression in mammalian cells, a codon-optimized mammalian expression construct
encoding full-length (amino acids 1-372) elephant shark (*Callorhinchus milii*) SIRT8
(Supplementary table S1), cloned in-frame into the *Eco*RI and *Sal*I sites of the pCMV-
3Tag-4a vector followed by vector encoding sequence of three successive Myc epitope
tags before a stop codon, was acquired from GenScript (Piscataway, NJ). For the
generation of a codon-optimized bacterial expression construct, a cDNA encoding N-
terminal truncated (amino acids 95-372) elephant shark SIRT8, cloned in-frame into the
*Eco*RI and *Sal*I sites of the pGEX-4T-1 vector, was also acquired from GenScript. This
cDNA was subsequently cloned in-frame into the *Eco*RI and *Sal*I sites of the pGST-
Parallel-1 vector (Sheffield et al. 1999). The nucleotide sequence of all recombinant
constructs was confirmed by dideoxy sequencing using the AUSTRAL-omics core facility
at Universidad Austral de Chile (https://australomics.cl/).
617

Cell culture, cell transfection and preparation of protein extracts

H4 human neuroglioma cells, HeLa human cervix adenocarcinoma cells and MDA-MB-
231 human mammary gland adenocarcinoma cells were obtained from the American
Type Culture Collection (Manassas, VA). H4 and HeLa cells were maintained in
Dulbecco's modified Eagle's medium (DMEM), and MDA-MB-231 in DMEM-F12
(ThermoFisher, Waltham, MA). For all cell lines, media were supplemented with 10%
heat-inactivated fetal bovine serum, 100 U/ml penicillin, 100 μg/ml streptomycin
(ThermoFisher), and 5 μg/ml plasmocin (InvivoGen, San Diego, CA), and cells were
cultured in a humidified incubator with 5% CO_2 at 37 °C. For transient transfections, cells

were either seeded on top of glass coverslips on 24-well plates or on 6-well plates. When
cells were ~60% confluent, transfections were performed with Lipofectamine 2000
(ThermoFisher), according to the manufacturer's instructions. Preparation of protein
extracts from cultured, transfected or non-transfected cells was performed using methods
described elsewhere (Tenorio et al. 2016).

Antibodies and cell reagents

We used the following antibodies: mouse clone 9E10 against the c-Myc epitope
(Covance, Princeton, NJ), mouse clone BA3R against β-ACTIN (ThermoFisher), rabbit
polyclonal against Cytoskeleton-associated protein 4 (P63; Merck, Germany; cat #
HPA001225), and sheep polyclonal against Trans-Golgi network integral membrane
protein 2 (TGN46; Bio-Rad Laboratories, Hercules, CA; cat # AHP500G). The following
fluorochrome-conjugated antibodies were from ThermoFisher: Alexa Fluor-488–
conjugated donkey anti mouse IgG, Alexa Fluor-594–conjugated donkey anti rabbit IgG
and Alexa Fluor-647-conjugated donkey anti sheep IgG. HRP-conjugated secondary
antibodies were from Jackson ImmunoResearch (West Grove, PA). Depending on their
reactivity, primary antibodies were used at a dilution 1/200 to 1/2000. HRP-conjugated
secondary antibodies were used at dilutions 1/5000 to 1/20000 also depending on their
reactivity. All Alexa Fluor-conjugated secondary antibodies were used at a dilution
1/1000. The fluorescent nuclear stain 4',6-diamidino-2-phenylindole (DAPI) was also from
ThermoFisher.

Protein Electrophoresis, Immunoblotting and Immunofluorescence
Microscopy

SDS-PAGE and immunoblotting were performed as described (Bustamante et al. 2020).
For fluorescence microscopy, cells grown on glass coverslips were transfected, and after
16-h cells were left untreated or treated for 30 min with MitoTracker™ Orange CMTMRos
(ThermoFisher) according to the manufacturer's instructions. After washing with
phosphate buffered saline (PBS) supplemented with 0.1 mM $CaCl_2$ and 1 mM $MgCl_2$
(PBS-CM), cells were fixed in 4% paraformaldehyde for 30 min at room temperature,
permeabilized with 0.2% Triton X-100 in PBS-CM for 15 min at room temperature, and

incubated with blocking solution (0.2% gelatin in PBS-CM) for 10 min at room

temperature. Cells were incubated simultaneously either with antibodies to the c-Myc

epitope (1/200) and to TGN46 (1/1000) or to the c-Myc epitope, to TGN46 and to P63

(1/600) for 30 min at 37 ºC in a humidified chamber. After washing coverslips with PBS,

cells were incubated as before, but with fluorochrome-conjugated antibodies, followed by

PBS-washing, and coverslips were mounted onto glass slides with Fluoromount-G

mounting medium (ThermoFisher). Fluorescence microscopy images were acquired with

an AxioObserver.D1 microscope equipped with a PlanApo 63x oil immersion objective

(NA 1.4), and an AxioCam MRm digital camera (Carl Zeiss). To quantitatively evaluate

colocalization of fluorescence signals, we obtained the Pearson's correlation coefficient

and cytofluorograms of pairwise comparisons from images acquired under identical

settings, avoiding signal saturation and corrected for background, crosstalk and noise

signals on each set of images, using the plugin JACoP (Bolte and Cordelières 2006)

implemented in the software ImageJ (version 1.47h)(Schneider et al. 2012), and the

plugin Colocalization Finder implemented in the software ImageJ2 (version 2.3.0/1.53)

(Rueden et al. 2017). To prepare figures, 12-bit images were processed with ImageJ

software and Adobe Photoshop CS3 software (Adobe Systems, Mountain View, CA).

Expression and Purification of Recombinant N-terminal truncated elephant

shark SIRT8 protein

Recombinant, N-terminal truncated elephant shark SIRT8 (ΔNT-SIRT8; amino acids 95-

372) tagged with an N-terminal glutathione S-transferase (GST) followed by a tobacco

etch virus (TEV) protease cleavage site was expressed and purified using a method

described previously (Ross et al. 2014), with some modifications. Briefly, expression in

E. coli B834(DE3) (Novagen, Madison, WI) was induced with 0.25 mM IPTG at 25 °C for

16 h. Pellets of bacteria were resuspended in homogenization buffer (50 mM Tris HCl,

0.5 M NaCl, 5 mM EDTA, 5 mM β-mercaptoethanol, and 2 mM phenylmethylsulfonyl

fluoride, pH 8.0) and lysed by sonication. The clarified supernatant was purified on

glutathione-Sepharose 4B (GE Healthcare). After removal of the GST moiety by TEV

cleavage, and sequential further passage through glutathione-Sepharose 4B and Ni-NTA

(QIAGEN) resins, ΔNT-SIRT8 was further purified on a Superdex 200 pg column (GE

Healthcare) equilibrated in storage buffer (50 mM Tris HCl, 150 mM NaCl, 1 mM DTT).

Aliquots of purified protein were kept at -80 ºC until use.

Deacetylation activity

To characterize deacetylation activity of recombinant, N-terminal truncated elephant

shark SIRT8 (ΔNT-SIRT8), we used a fluorometric SIRT3 activity assay kit (Abcam, cat

ab156067; Cambridge, UK), according to the manufacturer's instructions. This assay

allows detection of a fluorescent signal upon deacetylation of an acetylated substrate

peptide for recombinant human SIRT3. The intensity of fluorescence was measured on a

fluorometric microplate reader (Varioskan Flash, ThermoFisher) with excitation set at 350

nm and emission detection set at 450 nm. To analyze the deacetylation activity of N-

terminal truncated elephant shark SIRT8 protein, first we determined its optimum

temperature that resulted to be ~24 ºC. Subsequent analyses were performed by

comparing the activity of N-terminal truncated elephant shark SIRT8 protein at 24 ºC to

that of human SIRT3, which is provided by the assay kit, at 37 ºC. K_m and V_{max} were

obtained by varying the concentration of the fluorogenic acetylated substrate peptide

provided by the assay kit. This assay was also used to determine the effect of

nicotinamide (NAM), quercetin (QUE) and resveratrol (REV; Sigma-Aldrich) on the

deacetylation activity of ΔNT-SIRT8.

ATP and ROS content

ATP levels were quantified in cell lysates using a luciferin/luciferase bioluminescence

assay (ATP determination kit, Molecular Probes cat # A22066, Thermo Fisher Scientific)

as previously described (Jara et al. 2018). Briefly, cells were lysed in HEPES buffer (125

mM NaCl, 25 mM NaF, 1 mM EDTA, 1 mM EGTA, 1% NP-40, 25 mM HEPES, pH 7.4)

supplemented with a protease inhibitor mixture (cat # 78429, Thermo Fisher Scientific)

and centrifuged at 12000 rpm using a refrigerated centrifuge (Eppendorf model 5424R)

for 15 min at 4 ºC. Proteins in the supernatant were quantified, and 10 µl of lysate were

used to determine ATP levels according to the kit manufacturer's instructions. The ATP

level in each sample was calculated using standard curves and normalized to the protein

concentration in each sample. ROS content was determined in cell lysates using 25 µM

CM-H2DCFDA (DCF; a fluorogenic indicator of ROS), as previously described (Jara et
al. 2018; Olesen et al. 2020). The intensity of fluorescence was measured on a
fluorescence plate reader (Biotek Synergy HT, Agilent, Santa Clara, CA) with excitation
set at 485 nm and emission detection set at 530 nm. Briefly, in a dark 96-well plate, 25
µg of cell lysate proteins were incubated with CM-H2DCFDA (DCF) dye with shaking for
5 min at room temperature, followed by determination of the fluorescence of each sample
that was subtracted by the fluorescence of the respective blank.

Statistical Analysis

In the case of the deacetylation, ATP content and ROS content assays, quantification
was performed from at least three independent experiments/measurements. Statistical
analysis was performed using Microsoft Excel for Mac 2011 (Microsoft Corporation).
When appropriate, results are represented in graphs depicting the mean ± standard
deviation. Statistical significance was determined by two-tailed, paired t-test. P-values >
0.05 or ≤ 0.05 were regarded as not statistically significant or statistically significant,
respectively. In the figures, P-values between 0.01 and 0.05 are indicated with one
asterisk, P-values between 0.001 and 0.01 are indicated with two asterisks, and P-values
less than 0.001 are indicated with three asterisks.

Data accessibility

Data and supplementary material are available online:

Acknowledgements

This work was supported by the Fondo Nacional de Desarrollo Científico y Tecnológico
from Chile (FONDECYT 1210471) and The Millennium Nucleus of Ion Channel-
Associated Diseases is a Millennium Nucleus of the Millennium Scientific Initiative,
National Agency of Research and Development (ANID, NCN19_168), Ministry of Science,
Technology, Knowledge and Innovation, Chile to JCO, Fondo Nacional de Desarrollo
Científico y Tecnológico from Chile (FONDECYT 1211481) to GAM, National Science

Foundation (EPS-0903787, DBI-1262901 and DEB-1354147) to F.G.H., Financiamiento
Basal No. ANID/BASAL/ACE210009 to PVB and No. ANID/BASAL/FB210008 to PVB
and CTR, Fondap-Ideal Grant N° 15150003 to LV-C, ANID-Millennium Science Initiative
Program-Center ICM-ANID ICN2021_002 to LV-C, Fondo Nacional de Desarrollo
Científico y Tecnológico from Chile (FONDECYT 1180957) to FJM, Vicerrectoría de
Investigación, Desarrollo y Creación Artística, Universidad Austral de Chile to JCO, FJM,
LVC, GAM. We thank Stefanie Teuber, John Quiroga and Rafael Burgos, Instituto de
Farmacología y Morfofisiología, Facultad de Ciencias Veterinarias, Universidad Austral
de Chile, for providing assistance in the use of the Varioskan Flash microplate reader;
Hianara Bustamante, Instituto de Microbiología Clínica, Facultad de Medicina,
Universidad Austral de Chile, for providing quercetin and resveratrol; Víctor Castro,
Departamento de Biología, Facultad de Ciencias, Universidad de Chile, for providing
NAD^+ and nicotinamide; and Gonzalo Astroza, and Andrés Rivera-Dictter for technical
assistance.

Contributions

765 JCO and GAM designed the study. JCO, MWV, FGH, KZ, CM, CL, VC, FJM, PVB, CTR,
766 GAM, collected and/or analyzed data. JCO, PVB and GAM wrote the manuscript. JCO,
767 MWV, FGH, KZ, CM, CL, VC, LV-C, FJM, PVB, CTR and GAM, reviewed and edited the
768 manuscript. All authors contributed to the article and approved the submitted version.

Ethics declarations

Conflict of interest disclosure

772 The authors declare they have no conflict of interest relating to the content of this
773 article. Juan C. Opazo is a recommender for PCI Evol Biol and PCI Genomics.

Data availability

776 Data and supplementary material are available online at zenodo.org

References

Albalat R, Cañestro C. 2016. Evolution by gene loss. *Nat. Rev. Genet.* 17:379–391.

Albertson RC, Craig Albertson R, Cresko W, William Detrich H, Postlethwait JH. 2009.
Evolutionary mutant models for human disease. *Trends in Genetics* [Internet]
25:74–81. Available from: http://dx.doi.org/10.1016/j.tig.2008.11.006

Altenhoff AM, Train C-M, Gilbert KJ, Mediratta I, Mendes de Farias T, Moi D, Nevers Y,
Radoykova H-S, Rossier V, Warwick Vesztrocy A, et al. 2021. OMA orthology in
2021: website overhaul, conserved isoforms, ancestral gene order and more.
Nucleic Acids Res. 49:D373–D379.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment
search tool. *J. Mol. Biol.* 215:403–410.

Anderson KA, Green MF, Huynh FK, Wagner GR, Hirschey MD. 2014. SnapShot:
Mammalian Sirtuins. *Cell* 159:956–956.e1.

Anders S, Huber W. 2010. Differential expression analysis for sequence count data.
Genome Biol. 11:R106.

Anisimova M, Gil M, Dufayard J-F, Dessimoz C, Gascuel O. 2011. Survey of branch
support methods demonstrates accuracy, power, and robustness of fast likelihood-
based approximation schemes. *Syst. Biol.* 60:685–699.

Bause AS, Haigis MC. 2013. SIRT3 regulation of mitochondrial oxidative stress. *Exp.*
Gerontol. 48:634–639.

Bilyk KT, Zhuang X, Vargas-Chacoff L, Cheng C-HC. 2021. Evolution of chaperome
gene expression and regulatory elements in the antarctic notothenioid fishes.
Heredity 126:424–441.

Blander G, Guarente L. 2004. The Sir2 family of protein deacetylases. *Annu. Rev.*
Biochem. 73:417–435.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* 30:2114–2120.

Bolte S, Cordelières FP. 2006. A guided tour into subcellular colocalization analysis in
light microscopy. *J. Microsc.* 224:213–232.

Brown K, Xie S, Qiu X, Mohrin M, Shin J, Liu Y, Zhang D, Scadden DT, Chen D. 2013.
SIRT3 reverses aging-associated degeneration. *Cell Rep.* 3:319–327.

Bustamante HA, Cereceda K, González AE, Valenzuela GE, Cheuquemilla Y,
Hernández S, Arias-Muñoz E, Cerda-Troncoso C, Bandau S, Soza A, et al. 2020.
The Proteasomal Deubiquitinating Enzyme PSMD14 Regulates Macroautophagy

by Controlling Golgi-to-ER Retrograde Transport. *Cells* [Internet] 9. Available from:
http://dx.doi.org/10.3390/cells9030777

Castro LFC, Lopes-Marques M, Gonçalves O, Wilson JM. 2012. The evolution of
pepsinogen C genes in vertebrates: duplication, loss and functional diversification.
PLoS One 7:e32852.

Céspedes HA, Zavala K, Vandewege MW, Opazo JC. 2017. Evolution of the α-
adrenoreceptors in vertebrates: ADRA2D is absent in mammals and crocodiles.
Gen. Comp. Endocrinol. 250:85–94.

Costantini S, Sharma A, Raucci R, Costantini M, Autiero I, Colonna G. 2013. Genealogy
of an ancient protein family: the Sirtuins, a family of disordered members. *BMC*
Evol. Biol. 13:60.

Delsuc F, Brinkmann H, Chourrout D, Philippe H. 2006. Tunicates and not
cephalochordates are the closest living relatives of vertebrates. *Nature* 439:965–
968.

Delsuc F, Tsagkogeorga G, Lartillot N, Philippe H. 2008. Additional molecular support
for the new chordate phylogeny. *Genesis* 46:592–604.

Di Emidio G, Falone S, Artini PG, Amicarelli F, D'Alessandro AM, Tatone C. 2021.
Mitochondrial Sirtuins in Reproduction. *Antioxidants (Basel)* [Internet] 10. Available
from: http://dx.doi.org/10.3390/antiox10071047

Félix M-A, Barkoulas M. 2015. Pervasive robustness in biological systems. *Nat. Rev.*
Genet. 16:483–496.

Frye RA. 2000. Phylogenetic classification of prokaryotic and eukaryotic Sir2-like
proteins. *Biochem. Biophys. Res. Commun.* 273:793–798.

Frye RA. 2006. Evolution of Sirtuins From Archaea to Vertebrates. *Histone*
Deacetylases [Internet]:183–202. Available from: http://dx.doi.org/10.1385/1-59745-
024-3:183

Fujita Y, Yamashita T. 2018. Sirtuins in Neuroendocrine Regulation and Neurological
Diseases. *Front. Neurosci.* 12:778.

Gitelman I. 2007. Evolution of the vertebrate twist family and synfunctionalization: a
mechanism for differential gene loss through merging of expression domains. *Mol.*
Biol. Evol. 24:1912–1925.

Glover N, Dessimoz C, Ebersberger I, Forslund SK, Gabaldón T, Huerta-Cepas J,
Martin M-J, Muffato M, Patricio M, Pereira C, et al. 2019. Advances and
Applications in the Quest for Orthologs. *Mol. Biol. Evol.* 36:2157–2164.

Glozak MA, Sengupta N, Zhang X, Seto E. 2005. Acetylation and deacetylation of non-

histone proteins. *Gene* 363:15–23.

Goldberg T, Hecht M, Hamp T, Karl T, Yachdav G, Ahmed N, Altermann U, Angerer P,
Ansorge S, Balasz K, et al. 2014. LocTree3 prediction of localization. *Nucleic Acids*
Res. 42:W350–W355.

Greiss S, Gartner A. 2009. Sirtuin/Sir2 phylogeny, evolutionary considerations and
structural conservation. *Mol. Cells* 28:407–415.

Haigis MC, Mostoslavsky R, Haigis KM, Fahie K, Christodoulou DC, Murphy AJ,
Valenzuela DM, Yancopoulos GD, Karow M, Blander G, et al. 2006. SIRT4 inhibits
glutamate dehydrogenase and opposes the effects of calorie restriction in
pancreatic beta cells. *Cell* 126:941–954.

Haigis MC, Sinclair DA. 2010. Mammalian sirtuins: biological insights and disease
relevance. *Annu. Rev. Pathol.* 5:253–295.

Hara Y, Yamaguchi K, Onimaru K, Kadota M, Koyanagi M, Keeley SD, Tatsumi K,
Tanaka K, Motone F, Kageyama Y, et al. 2018. Shark genomes provide insights
into elasmobranch evolution and the origin of vertebrates. *Nat Ecol Evol* 2:1761–
1771.

Helsen J, Voordeckers K, Vanderwaeren L, Santermans T, Tsontaki M, Verstrepen KJ,
Jelier R. 2020. Gene Loss Predictably Drives Evolutionary Adaptation. *Mol. Biol.*
Evol. 37:2989–3002.

Herrero J, Muffato M, Beal K, Fitzgerald S, Gordon L, Pignatelli M, Vilella AJ, Searle
SMJ, Amode R, Brent S, et al. 2016. Ensembl comparative genomics resources.
Database [Internet] 2016. Available from:
http://dx.doi.org/10.1093/database/bav096

Himmel NJ, Gray TR, Cox DN. 2020. Phylogenetics Identifies Two Eumetazoan TRPM
Clades and an Eighth TRP Family, TRP Soromelastatin (TRPS). *Mol. Biol. Evol.*
37:2034–2044.

Hirschey MD. 2011. Old enzymes, new tricks: sirtuins are NAD(+)-dependent de-
acylases. *Cell Metab.* 14:718–719.

Hirschey MD, Shimazu T, Goetzman E, Jing E, Schwer B, Lombard DB, Grueter CA,
Harris C, Biddinger S, Ilkayeva OR, et al. 2010. SIRT3 regulates mitochondrial
fatty-acid oxidation by reversible enzyme deacetylation. *Nature* 464:121–125.

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2:
Improving the Ultrafast Bootstrap Approximation. *Mol. Biol. Evol.* 35:518–522.

Howe KL, Achuthan P, Allen J, Allen J, Alvarez-Jarreta J, Amode MR, Armean IM, Azov
AG, Bennett R, Bhai J, et al. 2021. Ensembl 2021. *Nucleic Acids Res.* 49:D884–
D891.

Iwabe N, Hara Y, Kumazawa Y, Shibamoto K, Saito Y, Miyata T, Katoh K. 2005. Sister
group relationship of turtles to the bird-crocodilian clade revealed by nuclear DNA-
coded proteins. *Mol. Biol. Evol.* 22:810–813.

Jara C, Aránguiz A, Cerpa W, Tapia-Rojas C, Quintanilla RA. 2018. Genetic ablation of
tau improves mitochondrial function and cognitive abilities in the hippocampus.
Redox Biol 18:279–294.

Kabiljo J, Murko C, Pusch O, Zupkovitz G. 2019. Spatio-temporal expression profile of
sirtuins during aging of the annual fish Nothobranchius furzeri. *Gene Expression*
Patterns [Internet] 33:11–19. Available from:
http://dx.doi.org/10.1016/j.gep.2019.05.001

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017.
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat.*
Methods 14:587–589.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol. Biol. Evol.* 30:772–780.

Kumar S, Stecher G, Suleski M, Hedges SB. 2017. TimeTree: A Resource for
Timelines, Timetrees, and Divergence Times. *Mol. Biol. Evol.* 34:1812–1819.

Kuraku S. 2013. Impact of asymmetric gene repertoire between cyclostomes and
gnathostomes. *Semin. Cell Dev. Biol.* 24:119–127.

Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biol.* 10:R25.

Leinonen R, Sugawara H, Shumway M, International Nucleotide Sequence Database
Collaboration. 2011. The sequence read archive. *Nucleic Acids Res.* 39:D19–D21.

Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. *BMC Bioinformatics* [Internet] 12. Available
from: http://dx.doi.org/10.1186/1471-2105-12-323

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15:550.

Lynch M, Conery JS. 2000. The Evolutionary Fate and Consequences of Duplicate
Genes. *Science* [Internet] 290:1151–1155. Available from:
http://dx.doi.org/10.1126/science.290.5494.1151

McBurney MW, Yang X, Jardine K, Hixon M, Boekelheide K, Webb JR, Lansdorp PM,
Lemieux M. 2003. The mammalian SIR2alpha protein has a role in embryogenesis
and gametogenesis. *Mol. Cell. Biol.* 23:38–54.

McCormick SD. 1993. Methods for nonlethal gill biopsy and measurement of Na+, K+-

920   ATPase activity. *Can. J. Fish. Aquat. Sci.* 50:656–658.

921 McDonnell E, Peterson BS, Bomze HM, Hirschey MD. 2015. SIRT3 regulates
922   progression and development of diseases of aging. *Trends Endocrinol. Metab.*
923   26:486–492.

924 Michan S, Sinclair D. 2007. Sirtuins in mammals: insights into their biological function.
925   *Biochem. J* 404:1–13.

926 Michishita E, Park JY, Burneskis JM, Barrett JC, Horikawa I. 2005. Evolutionarily
927   conserved and nonconserved cellular localizations and functions of human SIRT
928   proteins. *Mol. Biol. Cell* 16:4623–4635.

929 Nei M, Rooney AP. 2005. Concerted and birth-and-death evolution of multigene
930   families. *Annu. Rev. Genet.* 39:121–152.

931 Nguyen NTT, Vincens P, Roest Crollius H, Louis A. 2018. Genomicus 2018: karyotype
932   evolutionary trees and on-the-fly synteny computing. *Nucleic Acids Res.* 46:D816–
933   D822.

934 North BJ, Verdin E. 2004. Sirtuins: Sir2-related NAD-dependent protein deacetylases.
935   *Genome Biol.* 5:224.

936 Ohno S. 1985. Dispensable genes. *Trends Genet.* 1:160–164.

937 Olesen MA, Torres AK, Jara C, Murphy MP, Tapia-Rojas C. 2020. Premature synaptic
938   mitochondrial dysfunction in the hippocampus during aging contributes to memory
939   loss. *Redox Biol* 34:101558.

940 Olson MV. 1999. When less is more: gene loss as an engine of evolutionary change.
941   *Am. J. Hum. Genet.* 64:18–23.

942 Onyango P, Celic I, McCaffery JM, Boeke JD, Feinberg AP. 2002. SIRT3, a human
943   SIR2 homologue, is an NAD-dependent deacetylase localized to mitochondria.
944   *Proc. Natl. Acad. Sci. U. S. A.* 99:13653–13658.

945 Opazo JC, Hoffmann FG, Zavala K, Edwards SV. 2019. Evolution of the DAN gene
946   family in vertebrates. *Evolutionary Biology*:185.

947 Opazo JC, Kuraku S, Zavala K, Toloza-Villalobos J, Hoffmann FG. 2019. Evolution of
948   nodal and nodal-related genes and the putative composition of the heterodimers
949   that trigger the nodal pathway in vertebrates. *Evolution & Development* [Internet]
950   21:205–217. Available from: http://dx.doi.org/10.1111/ede.12292

951 Opazo JC, Vandewege MW, Gutierrez J, Zavala K, Vargas-Chacoff L, Morera FJ,
952   Mardones GA. 2021. Independent duplications of the Golgi phosphoprotein 3
953   oncogene in birds. *Sci. Rep.* 11:12483.

Pereira TCB, Rico EP, Rosemberg DB, Schirmer H, Dias RD, Souto AA, Bonan CD,
Bogo MR. 2011. Zebrafish as a model organism to evaluate drugs potentially able
to modulate sirtuin expression. *Zebrafish* 8:9–16.

Ponnambalam S, Girotti M, Yaspo ML, Owen CE, Perry AC, Suganuma T, Nilsson T,
Fried M, Banting G, Warren G. 1996. Primate homologues of rat TGN38: primary
structure, expression and functional implications. *J. Cell Sci.* 109 (Pt 3):675–685.

Qiu H, Hildebrand F, Kuraku S, Meyer A. 2011. Unresolved orthology and peculiar
coding sequence properties of lamprey genes: the KCNA gene family as test case.
BMC Genomics 12:325.

Qiu X, Brown K, Hirschey MD, Verdin E, Chen D. 2010. Calorie restriction reduces
oxidative stress by SIRT3-mediated SOD2 activation. *Cell Metab.* 12:662–667.

Rajabi N, Galleano I, Madsen AS, Olsen CA. 2018. Targeting Sirtuins: Substrate
Specificity and Inhibitor Design. *Prog. Mol. Biol. Transl. Sci.* 154:25–69.

Ramos-Vicente D, Ji J, Gratacòs-Batlle E, Gou G, Reig-Viader R, Luís J, Burguera D,
Navas-Perez E, García-Fernández J, Fuentes-Prior P, et al. 2018. Metazoan
evolution of glutamate receptors reveals unreported phylogenetic groups and
divergent lineage-specific events. *Elife* [Internet] 7. Available from:
http://dx.doi.org/10.7554/eLife.35774

Ross BH, Lin Y, Corales EA, Burgos PV, Mardones GA. 2014. Structural and functional
characterization of cargo-binding sites on the μ4-subunit of adaptor protein complex
4. *PLoS One* 9:e88147.

Rueden CT, Schindelin J, Hiner MC, DeZonia BE, Walter AE, Arena ET, Eliceiri KW.
2017. ImageJ2: ImageJ for the next generation of scientific image data. *BMC*
Bioinformatics 18:529.

Saravia J, Paschke K, Oyarzún-Salazar R, Cheng C-HC, Navarro JM, Vargas-Chacoff
L. 2021. Effects of warming rates on physiological and molecular components of
response to CTMax heat stress in the Antarctic fish Harpagifer antarcticus. *J.*
Therm. Biol. 99:103021.

Sauve AA, Schramm VL. 2003. Sir2 regulation by nicotinamide results from switching
between base exchange and deacetylation chemistry. *Biochemistry* 42:9249–9256.

Scher MB, Vaquero A, Reinberg D. 2007. SirT3 is a nuclear NAD+-dependent histone
deacetylase that translocates to the mitochondria upon cellular stress. *Genes Dev.*
21:920–928.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of
image analysis. *Nat. Methods* 9:671–675.

Scholte LLS, Mourão MM, Pais FS-M, Melesina J, Robaa D, Volpini AC, Sippl W, Pierce

RJ, Oliveira G, Nahum LA. 2017. Evolutionary relationships among protein lysine
deacetylases of parasites causing neglected diseases. *Infect. Genet. Evol.* 53:175–
188.

Schwartz S, Zhang Z, Frazer KA, Smit A, Riemer C, Bouck J, Gibbs R, Hardison R,
Miller W. 2000. PipMaker--a web server for aligning two genomic DNA sequences.
Genome Res. 10:577–586.

Schweizer A, Ericsson M, Bächi T, Griffiths G, Hauri HP. 1993. Characterization of a
novel 63 kDa membrane protein. Implications for the organization of the ER-to-
Golgi pathway. *J. Cell Sci.* 104 (Pt 3):671–683.

Schwer B, North BJ, Frye RA, Ott M, Verdin E. 2002. The human silent information
regulator (Sir)2 homologue hSIRT3 is a mitochondrial nicotinamide adenine
dinucleotide-dependent deacetylase. *J. Cell Biol.* 158:647–657.

Shahgaldi S, Kahmini FR. 2021. A comprehensive review of Sirtuins: With a major focus
on redox homeostasis and metabolism. *Life Sci.* 282:119803.

Sharma S, Ciufo S, Starchenko E, Darji D, Chlumsky L, Karsch-Mizrachi I, Schoch CL.
2019. The NCBI BioCollections Database. *Database* [Internet] 2019. Available
from: http://dx.doi.org/10.1093/database/baz057

Sheffield P, Garrard S, Derewenda Z. 1999. Overcoming expression and purification
problems of RhoGDI using a family of "parallel" expression vectors. *Protein Expr.*
Purif. 15:34–39.

Simó-Mirabet P, Bermejo-Nogales A, Calduch-Giner JA, Pérez-Sánchez J. 2017.
Tissue-specific gene expression and fasting regulation of sirtuin family in gilthead
sea bream (Sparus aurata). *Journal of Comparative Physiology B* [Internet]
187:153–163. Available from: http://dx.doi.org/10.1007/s00360-016-1014-0

Slade KM, Freggiaro S, Cottrell KA, Smith JJ, Wiley EA. 2011. Sirtuin-mediated nuclear
differentiation and programmed degradation in Tetrahymena. *BMC Cell Biol.* 12:40.

Smith JJ, Kuraku S, Holt C, Sauka-Spengler T, Jiang N, Campbell MS, Yandell MD,
Manousaki T, Meyer A, Bloom OE, et al. 2013. Sequencing of the sea lamprey
(Petromyzon marinus) genome provides insights into vertebrate evolution. *Nat.*
Genet. 45:415–421, 421e1–e2.

Somero GN. 2004. Adaptation of enzymes to temperature: searching for basic
"strategies." *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 139:321–333.

Stelzer G, Rosen N, Plaschkes I, Zimmerman S, Twik M, Fishilevich S, Stein TI, Nudel
R, Lieder I, Mazor Y, et al. 2016. The GeneCards Suite: From Gene Data Mining to
Disease Genome Sequence Analyses. *Curr. Protoc. Bioinformatics* 54:1.30.1–
1.30.33.

Tao R, Coleman MC, Pennington JD, Ozden O, Park S-H, Jiang H, Kim H-S, Flynn CR,
Hill S, Hayes McDonald W, et al. 2010. Sirt3-mediated deacetylation of
evolutionarily conserved lysine 122 regulates MnSOD activity in response to stress.
Mol. Cell 40:893–904.

Tatone C, Di Emidio G, Barbonetti A, Carta G, Luciano AM, Falone S, Amicarelli F.
2018. Sirtuins in gamete biology and reproductive physiology: emerging roles and
therapeutic potential in female and male infertility. *Hum. Reprod. Update* 24:267–
289.

Tenorio MJ, Ross BH, Luchsinger C, Rivera-Dictter A, Arriagada C, Acuña D, Aguilar M,
Cavieres V, Burgos PV, Ehrenfeld P, et al. 2016. Distinct Biochemical Pools of
Golgi Phosphoprotein 3 in the Human Breast Cancer Cell Lines MCF7 and MDA-
MB-231. *PLoS One* 11:e0154719.

Trifinopoulos J, Nguyen L-T, von Haeseler A, Minh BQ. 2016. W-IQ-TREE: a fast online
phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res.* 44:W232–
W235.

Tseng AHH, Shieh S-S, Wang DL. 2013. SIRT3 deacetylates FOXO3 to protect
mitochondria against oxidative damage. *Free Radic. Biol. Med.* 63:222–234.

Vassilopoulos A, Fritz KS, Petersen DR, Gius D. 2011. The human sirtuin family:
evolutionary divergences and functions. *Hum. Genomics* 5:485–496.

Vassilopoulos A, Pennington JD, Andresson T, Rees DM, Bosley AD, Fearnley IM, Ham
A, Flynn CR, Hill S, Rose KL, et al. 2014. SIRT3 deacetylates ATP synthase F1
complex proteins in response to nutrient- and exercise-induced stress. *Antioxid.*
Redox Signal. 21:551–564.

Vazquez BN, Vaquero A, Schindler K. 2020. Sirtuins in female meiosis and in
reproductive longevity. *Mol. Reprod. Dev.* 87:1175–1187.

Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, Heer FT, de
Beer TAP, Rempfer C, Bordoli L, et al. 2018. SWISS-MODEL: homology modelling
of protein structures and complexes. *Nucleic Acids Res.* 46:W296–W303.

Wichmann IA, Zavala K, Hoffmann FG, Vandewege MW, Corvalán AH, Amigo JD,
Owen GI, Opazo JC. 2016. Evolutionary history of the reprimo tumor suppressor
gene family in vertebrates with a description of a new reprimo gene lineage. *Gene*
591:245–254.

Wood JG, Rogina B, Lavu S, Howitz K, Helfand SL, Tatar M, Sinclair D. 2004. Sirtuin
activators mimic caloric restriction and delay ageing in metazoans. *Nature*
430:686–689.

Yang L, Ma X, He Y, Yuan C, Chen Q, Li G, Chen X. 2017. Sirtuin 5: a review of
structure, known inhibitors and clues for developing new inhibitors. *Sci. China Life*

Sci. 60:249–256.

Yates AD, Achuthan P, Akanni W, Allen J, Allen J, Alvarez-Jarreta J, Amode MR,
Armean IM, Azov AG, Bennett R, et al. 2020. Ensembl 2020. *Nucleic Acids Res.*
48:D682–D688.

You W, Zheng W, Weiss S, Chua KF, Steegborn C. 2019. Structural basis for the
activation and inhibition of Sirtuin 6 by quercetin and its derivatives. *Sci. Rep.*
9:19176.

Zhang J, Xiang H, Liu J, Chen Y, He R-R, Liu B. 2020. Mitochondrial Sirtuin 3: New
emerging biological function and therapeutic target. *Theranostics* 10:8315–8342.

Zhao E, Hou J, Ke X, Abbas MN, Kausar S, Zhang L, Cui H. 2019. The Roles of Sirtuin
Family Proteins in Cancer Progression. *Cancers* [Internet] 11. Available from:
http://dx.doi.org/10.3390/cancers11121949

Zhao H-C, Ding T, Ren Y, Li T-J, Li R, Fan Y, Yan J, Zhao Y, Li M, Yu Y, et al. 2016.
Role of Sirt3 in mitochondrial biogenesis and developmental competence of human
in vitro matured oocytes. *Hum. Reprod.* 31:607–622.