Research article

How many sirtuin genes are out there? evolution of sirtuin genes in vertebrates with a description of a new family member

Juan C. Opazo[1,2,3], Michael W. Vandewege[4], Federico G. Hoffmann[5,6], Kattina Zavala[2], Catalina Meléndez[7], Charlotte Luchsinger[8], Viviana A. Cavieres[7], Luis Vargas-Chacoff[1,9,10,11], Francisco J. Morera[1,12], Patricia V. Burgos[7,13,14], Cheril Tapia-Rojas[7,13], Gonzalo A. Mardones[1,8,15]

[1]Integrative Biology Group, Universidad Austral de Chile, Valdivia, Chile.
[2]Instituto de Ciencias Ambientales y Evolutivas, Facultad de Ciencias, Universidad Austral de Chile, Valdivia, Chile.
[3]Millennium Nucleus of Ion Channel-Associated Diseases (MiNICAD), Valdivia, Chile.
[4]Department of Biology, Eastern New Mexico University, Portales, New Mexico, USA.
[5]Department of Biochemistry, Molecular Biology, Entomology, and Plant Pathology, Mississippi State University, Starkville, Mississippi, USA.
[6]Institute for Genomics, Biocomputing and Biotechnology, Mississippi State University, Starkville, Mississippi, USA.
[7]Centro de Biología Celular y Biomedicina (CEBICEM), Facultad de Medicina y Ciencia, Universidad San Sebastián, Santiago, Chile.
[8]Department of Physiology, School of Medicine, Universidad Austral de Chile, Valdivia, Chile.
[9]Instituto de Ciencias Marinas y Limnológicas, Universidad Austral de Chile, Valdivia, Chile.
[10]Centro Fondap de Investigación de Altas Latitudes (IDEAL), Universidad Austral de Chile, Valdivia, Chile.
[11]Millennium Institute Biodiversity of Antarctic and Subantarctic Ecosystems, BASE, Universidad Austral de Chile, Valdivia, Chile.
[12]Applied Biochemistry Laboratory, Facultad de Ciencias Veterinarias, Instituto de Farmacología y Morfofisiología, Universidad Austral de Chile, Valdivia, Chile.
[13]Centro Ciencia & Vida, Fundación Ciencia & Vida, Santiago, Chile.
[14]Centro de Envejecimiento y Regeneración (CARE-UC), Facultad de Ciencias Biológicas, Pontificia Universidad Católica, Santiago, Chile.
[15]Center for Interdisciplinary Studies of the Nervous System (CISNe), Universidad Austral de Chile, Valdivia, Chile.

Corresponding authors:
Juan C. Opazo, jopazo@gmail.com
Gonzalo A. Mardones, gonzalo.mardones@uach.cl

Abstract

Studying the evolutionary history of gene families is a challenging and exciting task with

a wide range of implications. In addition to exploring fundamental questions about the

origin and evolution of genes, disentangling their evolution is also critical to those who do

functional/structural studies to allow a deeper and more precise interpretation of their

results in an evolutionary context. The sirtuin gene family is a group of genes that are

involved in a variety of biological functions mostly related to aging. Their duplicative

history is an open question, as well as the definition of the repertoire of sirtuin genes

among vertebrates. Our results show a well-resolved phylogeny that represents an

improvement in our understanding of the duplicative history of the sirtuin gene family. We

identified a new sirtuin family member (SIRT3-like) that was apparently lost in the last

common ancestor of amniotes, but retained in all other groups of jawed vertebrates.

According to our experimental analyses, elephant shark SIRT3-like protein is located in

mitochondria, the overexpression of which leads to an increase in cellular levels of ATP.

Moreover, *in vitro* analysis demonstrated it has deacetylase activity being modulated in a

similar way to mammalian SIRT3. Our results indicate that there are at least eight sirtuin

paralogs among vertebrates and that all of them can be traced back to the last common

ancestor of the group that existed between 676 and 615 millions of years ago.

Keywords: aging, deacetylase, gene family evolution, gene duplication, mitochondria,

SIRT.

Introduction

The availability of whole-genome sequences in species of all main groups of vertebrates represents an opportunity to unravel the evolution of gene families. The number of genomes and their phylogenetic representativeness in the vertebrate tree of life allows performing robust inferences regarding how gene family members are related to each other and their modes of evolution (Nei and Rooney 2005). The available genomes also open an opportunity to discover new gene lineages that are not currently described maybe because they are not present in model species and/or to the absence of appropriate evolutionary analyses (Wichmann et al. 2016; Céspedes et al. 2017; Himmel et al. 2020). Gene copy number variation could be seen as a natural experiment (Albertson et al. 2009) that could help understand the evolutionary fate of duplicated genes, as individuals with different repertoires can, in principle, fulfill the biological functions with a different combination of paralogs.

The sirtuin gene family, class III of histone deacetylase enzymes (HDACs), is an ancient group of genes that, in mammals, is composed of seven paralogs (SIRT1-7) grouped into four classes (Fig. 1)(Frye 2000; Frye 2006). Sirtuin genes are involved in a variety of biological functions mostly related to aging, metabolic regulation, stress response, cell cycle among others (Fig. 1) (Michan and Sinclair 2007; Greiss and Gartner 2009; Haigis and Sinclair 2010; Zhao et al. 2019; Zhang et al. 2020). All sirtuin genes have a conserved catalytic domain and variable carboxy- and amino terminal domains. Most family members possess deacetylase activity, while others have, in addition to weak deacetylase activity, other enzymatic activities like ADP-ribosyltransferase, desuccinylase and demalonylase (Fig. 1). Further, they are located in different subcellular compartments and associated to different biological functions (Fig. 1).

The evolution of the sirtuin genes is an open question. There are multiple phylogenetic hypotheses describing evolutionary relationships among the paralogs in the sirtuin gene family (Frye 2000; North and Verdin 2004; Frye 2006; Greiss and Gartner 2009; Pereira et al. 2011; Slade et al. 2011; Vassilopoulos et al. 2011; Costantini et al. 2013; Scholte et al. 2017; Simó-Mirabet et al. 2017; Yang et al. 2017; Rajabi et al. 2018; Kabiljo et al. 2019; Zhao et al. 2019). Differences in the taxonomic sampling, number of paralogs included, and the inclusion of outgroups, are likely contributors to this variation.

Additionally, studies that are focused on resolving evolutionary relationships are scarce,
and in fact, most phylogenetic analyses for sirtuin genes are part of studies where the
sirtuin duplicative history is a secondary goal. Additionally, and probably for similar
reasons, there are no systematic efforts to characterize the full complement of sirtuin
genes among vertebrates. Thus, unraveling the evolutionary history of sirtuin genes
represents a challenging and exciting task with a wide range of implications. Because of
their role in the aging process, these genes are of great interest. In addition to exploring
fundamental questions about the origin and evolution of sirtuin genes, disentangling their
evolution is also critical to understanding the diversification of functional and structural
phenotypes present in the sirtuin gene family.

This study aims to take advantage of the genomic data available in public
databases to advance our understanding of the diversity of vertebrate sirtuin genes and
to infer its duplicative history. We also aim to characterize the subcellular localization,
enzymatic activity, and mitochondrial activity of the newly discovered SIRT3-like gene
lineage in the elephant shark (*Callorhinchus milii*) as a representative species. Our
phylogenetic tree is in general well resolved, recovering sirtuin genes into two clades: 1)
SIRT4, SIRT5, SIRT6 and SIRT7, and 2) SIRT1, SIRT2, SIRT3, and SIRT3-like. The new
sirtuin family member (SIRT3-like) is retained in all vertebrates other than amniotes (the
group that includes mammals, birds, and reptiles). Based on how sirtuin genes are related
to each other and the information is already known for the other sirtuin family members,
we can predict that SIRT3-like belongs to the class I, has a deacetylase activity, and is
mainly located on mitochondria. Our experimental analyses confirmed our evolutionary
guided inferences.

Gene phylogeny	Synteny	Protein Size	MW (KD)	Enzymatic activity	Subcellular localization	Function	Disease
SIRT1 (Class I)	Human Chr10 — DNAJC12 SIRT1 HERC4	747	81.7	Deacetylase	Nucleus & Cytosol	Metabolism regulation, Regulation of chromatin and transcription, DNA repair & Inflammation suppression	Aging, Neurodegenration, Metabolic disease, Cardiovascular disease & Cancer
SIRT2 (Class I)	Human Chr19 — RINL SIRT2 NFKBIB	389	41.5	Deacetylase	Nucleus & Cytosol	Cell differentiation, Metabolism regulation, Cell cycle regulation, Microtubule dynamics & Inflammation regulation	Cancer & Neurodegeneration
SIRT3 (Class I)	Human Chr11 — RIC8A SIRT3 PSMD13	399	43.6	Deacetylase	Nucleus & Mitochondria	Metabolism regulation, Inflammation suppression, Inhibition of oxidative stress, Apoptosis regulation & Autophagy regulation	Aging, Neurodegenration, Metabolic disease, Cardiovascular disease & Cancer
SIRT3-like (Class I)	Elephant shark NW_024704746.1* — RIC8B SIRT3l TMEM263	401	44.2	Deacetylase	Mitochondria	?	?
SIRT4 (Class II)	Human Chr12 — PXN SIRT4 PLA2G1B	314	35.2	ADP-ribosyltransferase, Lipoamidase & Deacetylase	Mitochondria	Metabolism regulation	Neurodegenration, Metabolic disease & Cancer
SIRT5 (Class III)	Human Chr6 — GFOD1 SIRT5 NOL7	310	33.9	Deacetylase, Desuccinylase, Deglutarylase & Demalonylase	Mitochondria	Metabolism regulation & Immune response modulation	Metabolic disease, & Cancer
SIRT6 (Class IV)	Human Chr19 — CREB3L3 SIRT6 ANKRD24	355	39.1	Deacetylase & ADP-ribosyltransferase	Nucleus	Metabolism regulation & Chromatin and DNA repair	Aging, Metabolic disease, Cardiovascular disease & Cancer
SIRT7 (Class IV)	Human Chr17 — PCYT2 SIRT7 MAFG	400	44.9	Deacetylase	Nucleus	Transcription regulation, Chromatin remodeling & Metabolism regulation	Metabolic disease, Cardiovascular disease & Cancer

Figure 1. Gene phylogeny, synteny, protein size, molecular weight, enzymatic activity, subcellular localization, functions and diseases associated with sirtuin genes. Information regarding the sister group relationships among sirtuin genes was obtained from this study, synteny from ENSEMBL v.106 (Howe et al. 2021), protein size from Fujita and Yamashita (2018)(Fujita and Yamashita 2018), molecular weight from Vassilopoulos et al. (2011)(Vassilopoulos et al. 2011) while enzymatic activity, subcellular localization, functions and diseases from Zhang et al. (2020)(Zhang et al. 2020). In the case of the SIRT3-like of the elephant shark, synteny information was obtained from the National Center for Biotechnology Information (NCBI) (Sharma et al. 2019), protein size, molecular weight, enzymatic activity and subcellular localization from this study.

Results and Discussion

Sirtuin paralogs are recovered into two main clades

We recovered the monophyly of all sirtuin family members with strong support (Fig. 2), except for the SIRT3 clade which received moderate support (Fig. 3). In this case, the phylogenetic position of the sea lamprey (*Petromyzon marinus*) could be considered ambiguous (Fig. 3). This situation could be explained by the sequence itself, instead of the well-known compositional bias of the cyclostome genomes (Qiu et al. 2011; Kuraku

2013; Smith et al. 2013), given that the phylogenetic position of the other SIRT3
cyclostome sequence (inshore hagfish, *Eptatretus burgeri*) was recovered, as expected,
sister to all other vertebrate SIRT3 with strong support (Fig. 3). Additionally, our
phylogenetic reconstructions did not have difficulties inferring orthology for all other
cyclostome sirtuin genes. Further, in support of our tree topology, the sea lamprey
chromosomal region that harbors SIRT3 shares one syntenic gene, Bet1 Golgi vesicular
membrane trafficking protein-like (Bet1l), with humans.

According to our results, the diversity of sirtuin genes was arranged into two main
clades (Fig. 2). The first includes the SIRT4, SIRT5, SIRT6 and SIRT7, whereas the
second includes the SIRT1, SIRT2, SIRT3 and SIRT3-like lineages (Fig. 2). In the
literature, a diversity of phylogenetic arrangements have been proposed (Frye 2000;
North and Verdin 2004; Frye 2006; Greiss and Gartner 2009; Hirschey 2011; Pereira et
al. 2011; Slade et al. 2011; Vassilopoulos et al. 2011; Costantini et al. 2013; Scholte et
al. 2017; Simó-Mirabet et al. 2017; Yang et al. 2017; Rajabi et al. 2018; Kabiljo et al.
2019; Zhao et al. 2019), not being directly comparable as different studies implemented
different taxonomic samplings and not all of them include outgroups. One of the above
mentioned studies recovered the diversity of sirtuin genes into two main clades
(Vassilopoulos et al. 2011), however they did not recover the sister group relationship
between SIRT4 and SIRT5.

In the first clade we recovered the sister group relationships between SIRT4 and
SIRT5 with strong support (Fig. 2). Evolutionary relationships between these two family
members are still a matter of debate, as a variety of phylogenetic positions have been
suggested for them (Slade et al. 2011; Vassilopoulos et al. 2011; Costantini et al. 2013;
Yang et al. 2017; Rajabi et al. 2018), and only in a fraction of the studies their sister group
relationship is supported (Frye 2000; North and Verdin 2004; Frye 2006; Greiss and
Gartner 2009; Hirschey 2011; Pereira et al. 2011; Scholte et al. 2017; Simó-Mirabet et al.
2017; Kabiljo et al. 2019; Zhao et al. 2019). In the same clade, we also recovered SIRT6
sister to the clade containing SIRT7 sequences with strong support (Fig. 2). The sister
group relationship between SIRT6 and SIRT7 has been recovered in all examined studies
(Frye 2000; North and Verdin 2004; Frye 2006; Greiss and Gartner 2009; Hirschey 2011;
Pereira et al. 2011; Scholte et al. 2017; Simó-Mirabet et al. 2017; Kabiljo et al. 2019; Zhao

et al. 2019), suggesting a robust support for the sister group relationship between these

sirtuin lineages. The sister-group relationship between the SIRT6/SIRT7 and

SIRT4/SIRT5 clades is moderately supported (Fig. 2), being consistent with the diversity

of topologies reported in the literature, and suggesting that more work is necessary (Frye

2000; North and Verdin 2004; Frye 2006; Greiss and Gartner 2009; Hirschey 2011; Slade

et al. 2011; Vassilopoulos et al. 2011; Costantini et al. 2013; Scholte et al. 2017; Simó-

Mirabet et al. 2017; Yang et al. 2017; Rajabi et al. 2018; Kabiljo et al. 2019; Zhao et al.

2019). In the second clade, there is a broad consensus in the literature that SIRT2 shares

a common ancestor more recently in time with SIRT3 than with any other sirtuin family

member, and that the clade containing SIRT1 sequences is sister to the SIRT2/SIRT3

clade (Frye 2000; North and Verdin 2004; Frye 2006; Greiss and Gartner 2009; Hirschey

2011; Pereira et al. 2011; Slade et al. 2011; Vassilopoulos et al. 2011; Costantini et al.

2013; Scholte et al. 2017; Simó-Mirabet et al. 2017; Yang et al. 2017; Rajabi et al. 2018;

Zhao et al. 2019). Accordingly, we recovered the same evolutionary relationships with

strong support (Fig. 2).

In summary, we present a phylogenetic analysis based on a taxonomic sampling

that included representative species from all main groups of vertebrates for all sirtuin

family members. Our phylogenetic tree is in general well resolved (Fig. 2), representing

an advance in our understanding of the duplicative history of the sirtuin gene family. In

comparison to the gene trees currently available in the literature our tree topology

represents a new hypothesis for their duplicative history.

 Figure 2. Maximum likelihood tree showing sister group relationships among sirtuin
193 genes of vertebrates. Numbers on the nodes correspond to support from the aBayes and
194 ultrafast bootstrap values. Nicotinamide Nucleotide Transhydrogenase (NNT) sequences
195 from the human (*Homo sapiens*) mouse (*Mus musculus*), spotted gar (*Lepisosteus*
196 *oculatus*) and zebrafish (*Danio rerio*) were used as outgroups (not shown). The scale
197 denotes substitutions per site and colors represent sirtuin lineages.
198

Identification of a new sirtuin gene family member, SIRT3-like

In our analyses, we identified a new sirtuin family member (Fig. 2 and 3), SIRT3-like,

which is present in a fraction of the vertebrate tree of life (Fig. 4) that was recovered sister

to the SIRT3 clade with strong support (Fig. 2 and 3). Synteny conservation provides

further support to the monophyly of the SIRT3-like gene lineage in gnathostomes (Fig.

5a), as genes found at the 5` side (POLR3B, RFX4 and RIC8B) and 3` side (TMEM263,

MTERF2, and CRY1) of the SIRT3-like gene are well conserved (Fig. 5a). Synteny is also

conserved in species in which the SIRT3-like gene was lost (Fig. 5a). Among vertebrates,

we found orthologs of the SIRT3-like gene in representative species of cartilaginous fish,

bony fish, coelacanth, and amphibians (Fig. 4). A comparison of the genomic region of

the tropical clawed frog (*Xenopus tropicalis*), which possesses the SIRT3-like gene, with

the corresponding region in the human (*Homo sapiens*), opossum (*Monodelphis*

domestica), chicken (*Gallus gallus*), gharial (*Gavialis gangeticus*), red-eared slider

(*Trachemys scripta*) and green anole (*Anolis carolinensis*) strongly suggests that the

SIRT3-like gene is not present in the vertebrate lineages that these species represent

(Fig. 5b). The absence of the SIRT3-like gene in mammals, birds and reptiles indicates

that it was probably lost in the common ancestor of the group, between 352 and 312

millions of years ago (Kumar et al. 2017). In the case of cyclostomes, we found in the sea

lamprey (*Petromyzon marinus*) a chromosomal region (Chr3) that possess TMEM263 and

POLR3B separated by 44 base pairs, suggesting that all the genes that should be in

between (RFX4, RIC8B, SIRT3-like) were lost in cyclostomes. Alternatively, it is possible

that SIRT3-like, and all the other genes are present in cyclostome genomes, but not in

the current genome assembly. In the past, Pereira et al. (2011), with a limited taxonomic

sampling, identified a clade sister to a group containing SIRT3 sequences. This gene

lineage included sequences from three species with an evolutionary history of whole-

genome duplications, two teleost fish (zebrafish, *Danio rerio*, and green spotted puffer,

Tetraodon nigroviridis), and one amphibian (African clawed frog, *Xenopus laevis*),

complicating the definition of their duplicative history. Thus, our results, including a

balanced taxonomic sampling of all main vertebrate groups and appropriate phylogenetic

searches, define the existence of the SIRT3-like gene lineage and its phyletic distribution.

Figure 3. Maximum likelihood tree showing sister group relationships among SIRT3 and
SIRT3-like genes in vertebrates. Numbers on the nodes correspond to support from the
aBayes and ultrafast bootstrap values. The scale denotes substitutions per site and colors
represent gene lineages. This tree does not represent a novel phylogenetic analysis, it is

the SIRT3/SIRT3-like clade that was recovered from figure 2. The scale denotes
substitutions per site and colors represent sirtuin lineages.
236

The lack of the SIRT3-like gene can be interpreted in different ways. We can think

that the loss of SIRT3-like could have no physiological impact if the functional role of the

lost gene is assumed by other family members, consistent with the idea that gene families

possess functional redundancy (Ohno 1985; Félix and Barkoulas 2015; Albalat and

Cañestro 2016). Alternatively, we can also think of gene loss as a source of adaptive

evolution (Olson 1999; Albalat and Cañestro 2016; Helsen et al. 2020). The description

of new gene lineages is not uncommon in the literature (Wichmann et al. 2016; Himmel

et al. 2020; Opazo et al. 2021). It could be mainly attributed to the presence in non-model

species and/or the absence of proper evolutionary analyses. The existence of species

with different gene repertoires is an opportunity to better understand the evolutionary fate

of duplicated genes (Lynch and Conery 2000) and the biological functions associated to

a group of genes.

Figure 4. Phyletic distribution of sirtuin genes in vertebrates. The sister group
relationships among sirtuin genes was obtained from this study, whereas, organismal
phylogeny was obtained from the most updated phylogenetic hypotheses available in the
literature (Iwabe et al. 2005; Delsuc et al. 2006; Delsuc et al. 2008; Hara et al. 2018). The

colors of the circles are those used in figure 2 to define sirtuin gene lineages, whereas
the classes to which the different sirtuins belong are shown with shading. The question
mark denotes that we cannot determine whether or not the sirtuin3-like gene is present
in cyclostomes.
259

260
261

Figure 5. a Patterns of conserved synteny in the chromosomal region that harbor SIRT3-
like genes in gnathostomes. Asterisks denote that the orientation of the genomic piece is
from 3´ to 5´, gray lines represent intervening genes that do not contribute to conserved
synteny. **b** Pairwise dot-plot comparison of the genomic region of the tropical clawed frog
(*Xenopus tropicalis*) with the corresponding region in the human (*Homo sapiens*),
opossum (*Monodelphis domestica*), chicken (*Gallus gallus*), gharial (*Gavialis*
gangeticus), red-eared slider (*Trachemys scripta*) and green anole (*Anolis carolinensis*).
Light green vertical lines denote exons and regions in between are introns. Dot plots were
based on the complete coding region in addition to 6.7 kb of upstream and downstream
flanking sequence.
272

An amino acid alignment of the SIRT3-like and SIRT3 sequences shows that the

catalytic domain of SIRT3-like is well conserved, where six amino acid positions were

identified as diagnostic characters (Fig. 6). The amino acid divergence of the SIRT3-like

catalytic domain varies from 32.2% (spotted gar vs coelacanth) to 33.6% (spotted gar vs

elephant shark). The same comparisons for the SIRT3 catalytic domain show slightly

lower amino acid divergence values, between 24.8% (spotted gar vs coelacanth) and

31.0% (coelacanth vs elephant shark). As expected, the interparalog distance (SIRT3 vs

SIRT3-like) of the catalytic domains shows higher divergence values, ranging from 40.7%

(spotted gar SIRT3 vs spotted gar SIRT3-like) and 43.6% (elephant shark SIRT3 vs

elephant shark SIRT3-like). Although nothing is known about the protein encoded by the

SIRT3-like gene, based on how sirtuin genes are evolutionarily related and the

information already known for the other family members (Fig. 1), we can speculate that

SIRT3-like belongs to the class I, is mainly located on the mitochondria/nucleus, and has

deacetylase activity. From a functional perspective it should perform functions similar to

what has been described for SIRT3 (Fig. 1) (Brown et al. 2013; McDonnell et al. 2015;

Zhang et al. 2020). It is important to highlight that the inferences regarding a newly

discovered gene are better performed if they are phylogenetically guided.

Figure 6. Alignment of the catalytic domain of SIRT3 in humans (*Homo sapiens*), and
SIRT3 and SIRT3-like of spotted gar (*Lepisosteus oculatus*), coelacanth (*Latimeria*
chalumnae) and elephant shark (*Callorhinchus milii*). The shaded region denotes the
catalytic domain. Diagnostic characters -i.e. amino acids positions that distinguish

between SIRT3 and SIRT3-like gene lineages - are indicated with a rectangle and green
colors.

Expression pattern of sirtuin gene family members

Our next step in characterizing the SIRT3-like gene was to investigate whether it
is transcribed and if it was, to characterize the transcription pattern. To do this, we
mapped RNASeq reads to reference gene sequences of representative species of
vertebrates and examined transcript abundance. In agreement with the transcription
pattern reported for sirtuin genes (Stelzer et al. 2016; Kabiljo et al. 2019), our results show
that although they exhibited wide variance in tissue expression, sirtuin genes are
expressed in almost all tissues, including the novel SIRT3-like gene lineage (Fig. 7). In
the case of the tropical clawed frog, the SIRT3-like gene is transcribed at a similar level
in all tissues, other than the ovary where high transcription levels were estimated (Fig. 7).
Coincident with the pattern observed in the tropical clawed frog, the elephant shark
SIRT3-like gene is mostly transcribed in the ovary (Fig. 7). In the case of the zebrafish,
although our transcription level estimations did not follow the trend previously mentioned,
an extensive study examining the expression of all sirtuin genes in zebrafish showed that
the SIRT3-like gene in this species is mainly expressed in the ovary (Pereira et al. 2011).
Thus, in addition to showing that the SIRT3-like gene is transcribed, our analyses allow
us to suggest that this gene could be involved in biological processes associated with
reproduction. This observation seems not to be unexpected as for SIRT3 there are many
studies in which its functions related to reproduction have been reported (Zhao et al. 2016;
Tatone et al. 2018; Vazquez et al. 2020; Di Emidio et al. 2021).

Figure 7. Heatmap representation of transcription levels of sirtuin genes among vertebrates. Transcription values were calculated independently for each species and normalized over all tissues.

Elephant shark SIRT3-like protein localizes to mitochondria

To further explore the structural relationship between SIRT3 and SIRT3-like proteins, we performed an unbiased protein structure homology modeling of SIRT3-like proteins from elephant shark (*Callorhinchus milii*), spotted gar (*Lepisosteus oculatus*), salmon (*Salmo salar*) and coelacanth (*Latimeria chalumnae*). Models of SIRT3-like proteins were generated by the SWISS-MODEL workspace that used as template crystal structure models of human SIRT3: 4bvh.1.A for elephant shark, 5z94.1.A for spotted gar, 5d7n.3.A for salmon and 5bwo.1.B for coelacanth (data not shown). All models generated comprise only the central portion of the SIRT3-like proteins where the catalytic and its regulatory regions are expected to be. The overall root mean square deviation for 272 superimposable Cα coordinates for the models of human SIRT3 (pdb: 4bvh.1.A; amino acids G121 to G392) and elephant shark SIRT3-like (amino acids G95 to Q365) is 0.101 Å (Fig. 8a), suggesting extensive structural match. SIRT proteins possess evolutionary conserved, distinct cellular localizations and functions (Fig. 1). Some sirtuins localize in more than one compartment, like SIRT1 and SIRT2 that localize to the nucleus and cytosol, and SIRT3 that localizes mainly to the mitochondrial matrix, but with a small fraction localizing to the nucleus (Scher et al. 2007). In contrast, SIRT4 and SIRT5 localize only to mitochondria, and SIRT6 and SIRT7 localize only to the nucleus. The sequence and structural comparison between SIRT3-like and SIRT3 proteins, as well as subcellular localization prediction by LocTree3 (Goldberg et al. 2014), suggest that the localization of SIRT3-like proteins is also in the mitochondria. To determine the subcellular localization of SIRT3-like proteins, we chose to analyze SIRT3-like from elephant shark. To this end, we expressed in human cultured cells full-length elephant shark SIRT3-like tagged with three copies of the c-Myc epitope at the C-terminus (SIRT3-like-3myc). To obtain biochemical evidence of the expression of this protein, we performed immunoblot analysis with monoclonal antibody 9E10 against the c-Myc epitope, which showed the

detection of an expected 48.3 kDa protein in all cell lines analyzed (Fig. 8b). Fluorescence
microscopy analysis of human H4 neuroglioma cells expressing SIRT3-like-3myc showed
decoration by the anti-c-Myc antibody of structures highly reminiscent of mitochondria, in
addition to a faint fluorescent signal in the rest of the cytoplasm (Fig. 8c, d). Mitochondria
localization of SIRT3-like-3myc was confirmed by co-localization with the fluorescent
signal of MitoTracker Orange (Fig. 8c). Importantly, SIRT3-like-3myc showed no
localization in any other major compartment, as shown by lack of localization in the
nucleus (detected with the nuclear stain DAPI; Fig. 8c, d), the Golgi apparatus (detected
with antibody against TGN46; Fig. 8c, d), or the endoplasmic reticulum (detected with
antibody against P63; Fig. 8d). The same pattern of expression of SIRT3-like-3myc was
obtained using other human cell lines, such as HeLa or MDA-MB-231 cells (data not
shown). Altogether, these results indicate that SIRT3-like proteins are targeted to the
mitochondria.

Figure 8. Structural model of elephant shark SIRT3-like and expression in mammalian cells of SIRT3-like-3myc. **a** Superposition of the ribbon representations of elephant shark SIRT3-like (green model; amino acids G95 to Q365) and human SIRT3 (magenta model; pdb: 4bvh.1.A; amino acids G121 to G392). Depicted in ellipses are the indicated functional/structural domains of SIRT3 and in dotted lines the binding sites for peptide substrates and NAD$^+$. **b** The indicated cells were left untreated (Control, lanes 1, 3 and 5) or transfected to express SIRT3-like-3myc (lanes 2, 4 and 6), followed by immunoblot analysis with antibody against the c-Myc epitope or antibody against β-ACTIN used as loading control. The position of molecular mass markers is indicated on the left. SIRT3-like-3myc exhibits an electrophoretic mobility corresponding to a protein of 48.3 kDa, which is expected considering the extra amino acids of the three copies of the c-Myc epitope at the C-terminus. **c** and **d** H4 cells grown in glass coverslips were transfected to express SIRT3-like-3myc (green channel) and incubated with MitoTracker Orange (red channel in c) or mock incubated (d). Cells were fixed, permeabilized, and double- (c) or triple-labeled (d) with mouse monoclonal antibody against the c-Myc epitope (c and d), rabbit polyclonal antibody against the endoplasmic reticulum protein P63 (d) and sheep polyclonal antibody against the Golgi apparatus protein TGN46 (c and d). Secondary antibodies were Alexa-Fluor-488-conjugated donkey anti-mouse IgG (green channel), Alexa-Fluor-594-conjugated donkey anti-rabbit IgG (red channel), and Alexa-Fluor-647-conjugated donkey anti-sheep IgG (blue channel), and nuclei were stained with DAPI (white channel). Stained cells were examined by fluorescence microscopy. Insets: X3 magnification with yellow arrows indicating colocalization (c) and green and red arrows indicating lack of colocalization (d). Bar, 10 μm.

Elephant shark SIRT3-like protein has deacetylase activity

Reversible lysine acetylation is one of the most common post-translational modifications on proteins that regulate a variety of physiological processes including gene expression, enzymatic activity, protein-protein interactions, and subcellular localization (Glozak et al. 2005). Sirtuin proteins share a catalytic core domain (North and Verdin 2004) that have NAD$^+$-dependent deacetylase activity (Shahgaldi and Kahmini 2021). After translocation into the mitochondrial matrix, human SIRT3 is inactive, but after proteolytic processing of its N-terminus signal peptide it becomes active as deacetylase (Onyango et al. 2002; Schwer et al. 2002). Similar to human SIRT3, secondary structure prediction of representative SIRT3-like proteins indicates that the first ~100 N-terminal amino acids are in a disordered domain (data not shown). Therefore, to determine if SIRT3-like proteins also exhibit deacetylase activity, we produced in bacteria N-terminal truncated elephant shark SIRT3-like protein (amino acids 95-372; ΔNT-SIRT3-like). To analyze the enzymatic activity of purified ΔNT-SIRT3-like, we used a commercial fluorometric

deacetylation activity assay that contains an acetylated substrate peptide for recombinant

human SIRT3 (Abcam). An initial analysis of activity at 37ºC showed that ΔNT-SIRT3-like

possesses a deacetylase activity that is similar to that of human SIRT3, being also NAD$^+$-

dependent (Fig. 9a, e, f). Because the enzymatic activity of elephant shark SIRT3-like

could be influenced by the environmental conditions, we next evaluated the optimal

temperature of ΔNT-SIRT3-like compared to that of human SIRT3. As expected, the

optimal temperature of the human SIRT3 was ~37ºC (Fig. 9b). In contrast, ΔNT-SIRT3-

like exhibited an optimal temperature at ~24ºC (Fig. 9b). The optimal temperature value

for the SIRT3-like deacetylase activity of the elephant shark could represent an

evolutionary adjustment to the temperature regime of the habitat in which this species

lives, as it has been demonstrated for other enzymes in other species (McCormick 1993;

Somero 2004; Bilyk et al. 2021; Saravia et al. 2021). Thus, all subsequent enzymatic

activity analyses for ΔNT-SIRT3-like were performed at 24ºC. The apparent specific

activity of ΔNT-SIRT3-like was 260.7 ± 9.9 fluorescence units (FU) min^{-1} mg^{-1}, which was

significantly lower than that of SIRT3 that was 1143.3 ± 46.3 FU min^{-1} mg^{-1}. Apparent K_m

of ΔNT-SIRT3-like and SIRT3 were 5.7 and 5.5, respectively, which resulted not

significantly different (Fig. 9c, d). In contrast, apparent V_{max} of ΔNT-SIRT3-like was 0.322

FU sec^{-1}, which was significantly different to that of SIRT3 that was 3.908 FU sec^{-1} (Fig.

9c, d). We next compared the effect of the sirtuin inhibitors nicotinamide (NAM) (Sauve

and Schramm 2003) and quercetin (QUE) (You et al. 2019) on the activities of SIRT3 and

ΔNT-SIRT3-like. As expected, SIRT3 activity was significantly inhibited by NAM and QUE

(Fig. 9e). We also found a significant inhibitory effect of both NAM and QUE on the activity

of ΔNT-SIRT3-like (Fig. 9f). We also compared the effect of the polyphenolic antioxidant

resveratrol (REV), which is an activator of sirtuins (Wood et al. 2004). As expected, we

found a significant concentration-dependent activation of SIRT3 deacetylase activity up

to a 1.6 ± 0.1-fold increase in the presence of 100 μM REV (Fig. 9e). Strikingly, we found

a much potent concentration dependent activation effect of REV on ΔNT-SIRT3-like,

showing up to a 11.1 ± 0.6 fold increase in the presence of 100 μM REV (Fig. 9f).

However, because the specific activity of ΔNT-SIRT3-like was ~23% that of SIRT3 in the

assay conditions, the activation of ΔNT-SIRT3-like by REV in fact represents a value only

~1.6 fold higher compared to the activation of SIRT3 by REV. Because different sirtuins

have activity on different proteins, as well as remove with different specificity a variety of
acyl groups other than the acetyl (Fig. 1) (Anderson et al. 2014), the modulation of ΔNT-
SIRT3-like activity by REV suggests that it has distinct functional substrates. Together,
these results demonstrate that elephant shark ΔNT-SIRT3-like has deacetylase activity
and indicate that the activity of SIRT3-like proteins is modulated in a similar manner to
that of mammalian sirtuins.

443
444
445

Figure 9. Comparison of deacetylase activity between human SIRT3 and elephant shark ΔNT-SIRT3-like. a-f) Results obtained with a fluorometric deacetylation assay of an acetylated substrate peptide. a) Time course of deacetylase activity at the indicated concentration of SIRT3 and concentrations of ΔNT-SIRT3-like. b) Temperature dependence of deacetylase activity. c-d) Michaelis-Menten plot and Lineweaver-Burk plot of SIRT3 (c) and ΔNT-SIRT3-like (d) deacetylase activity. e-f) NAD$^+$ dependence and concentration dependent effects of the SIRT3 inhibitors nicotinamide (NAM) and quercetin (QUE) and of the SIRT3 activator resveratrol (REV) on the deacetylase activity of SIRT3 (e) and ΔNT-SIRT3-like (f). In a-d, graphs depict the mean ± standard deviation (n = 3). In e-f, bars represent the mean ± standard deviation. Statistical analysis was performed using two-tailed unpaired Student's t-test (n = 3; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; ns, not statistically significant).

Elephant shark SIRT3-like protein increases ATP cellular content

Considering that SIRT3 plays an important role in maintaining basal ATP levels by deacetylation of mitochondrial proteins involved in electron transport chain (ETC), the tricarboxylic acid cycle, and fatty-acid oxidation (Hirschey et al. 2010), we investigated the impact of transient SIRT3-like-3myc expression on ATP cellular content in H4 and HeLa cells (Fig. 10a). Similar to the well-known effect of mammalian SIRT3, SIRT3-like-3myc expression increased ATP cellular content, suggesting that it shares with SIRT3 the ability to enhance the activity of the mitochondrial ETC. Because an enhancement in the activity of ETC can cause an increase in reactive oxygen species (ROS), we also evaluated the levels of ROS in cells expressing SIRT3-like-3myc (Fig. 10b). We found similar ROS levels in cells expressing SIRT3-like-3myc and untransfected cells (Fig. 10b). This finding suggests that SIRT3-like proteins share the mammalian SIRT3 characteristic of protection against oxidative damage (Bause and Haigis 2013), likely by deacetylation of enzymes involved in ROS clearance to protect mitochondrial membranes from oxidative stress (Tseng et al. 2013). A candidate that could be activated by the deacetylation activity of SIRT3-like proteins is SOD2 that in human cells reduces cellular ROS content promoting oxidative stress resistance (Qiu et al. 2010; Tao et al. 2010). Alternatively, SIRT3-like proteins could activate the last complex of the ETC by

deacetylation, enhancing ATP-synthase activity without affecting ROS production, similar

to the effect that SIRT3 has in response to exercise-induced stress (Vassilopoulos et al.

2014). Together, our results show that the SIRT3-like protein is a new sirtuin family

member with deacetylase activity that could have important roles in the homeostasis of

oxidative species and mitochondrial metabolism.

Figure 10. SIRT3-like-3myc expression causes an increment on ATP cellular
content with no difference in ROS levels. (a) ATP levels were measured by a
luciferin/luciferase bioluminescence assay in H4 and HeLa cells either left untreated
(Control) or transiently expressing SIRT3-like-3myc. (b) ROS content was measured by
the intensity fluorescence of the dye CM-H2DCFDA in H4 and HeLa cells either left
untreated (Control) or transiently expressing SIRT3-like-3myc. Bars represent the mean
± standard deviation. Statistical analysis was performed using two-tailed unpaired
Student's t-test (n = 4; ** $p < 0.01$; ns, not statistically significant).

Conclusions

We performed an evolutionary analysis of the sirtuin gene family in vertebrates. In
addition to describing the duplicative history of sirtuins, our phylogenetic analyses
revealed the presence of a new gene family member, SIRT3-like. This new gene lineage
is present in all vertebrates other than mammals, birds, and reptiles. Furthermore, our
transcriptomic analyses show that SIRT3-like is transcribed in almost all examined
tissues, showing a tendency to be more expressed in the ovary, suggesting biological
functions related to reproduction. According to our analyses, SIRT3-like, localized in the
mitochondria, had deacetylase activity and increased the ATP cellular content. The
finding of a new sirtuin gene lineage highlights the need for more detailed assessments
of orthology with a more broad taxonomic sampling to better define the membership
composition of gene families (Glover et al. 2019). In the literature, there are examples in
which more comprehensive analyses provide a better description of gene families,
including previously unknown family members (Castro et al. 2012; Wichmann et al. 2016;
Céspedes et al. 2017; Ramos-Vicente et al. 2018; Opazo, Kuraku, et al. 2019; Opazo,
Hoffmann, et al. 2019). The availability of species with different gene repertoires could be
seen as a natural experiment (Albertson et al. 2009) that helps us understand the
evolutionary fate of duplicated genes, as they can fulfill the biological functions associated
with the gene family but with a different combination of paralogs.

Methods

Protein sequences and phylogenetic analyses

We retrieved sirtuin amino acid sequences in representative species of all main groups
of vertebrates. Our sampling included mammals, birds, reptiles, amphibians, coelacanth,
bony fish, cartilaginous fish, and cyclostomes (Supplementary Table S1). Sequences
were obtained from the Orthologous MAtrix project (OMA), December 2021 release
(Altenhoff et al. 2021)(last accessed on April 8th, 2022). In addition, we complemented
our sampling by retrieving sequences from vertebrate groups that were not present or
poorly represented in the Orthologous MAtrix project (e.g., crocodiles) from the National
Center for Biotechnology Information (NCBI) (Sharma et al. 2019) using the program blast
(tblastn) with default parameters (Altschul et al. 1990)(Supplementary Table S1). Protein
sequences were aligned using the software MAFFT v.7 (Katoh and Standley 2013),

allowing the program to choose the alignment strategy (FFT-NS-i). To select the best-
fitting model of molecular evolution we used the proposed model tool in the program IQ-
Tree v1.6.12 (Kalyaanamoorthy et al. 2017), which selected JTT+F+I+G4. We used a
maximum-likelihood approach to obtain the best tree using the program IQ-Tree v1.6.12
(Trifinopoulos et al. 2016). We carried out 20 independent runs to explore the tree space
changing the value of the strength of the perturbation (-pers) parameter. We conducted
the following analyses:

1. Five runs modifying the strength of the perturbation parameter from 0.5 (default
value) to 0.3.
2. Five runs using the default value (0.5) for the strength of the perturbation
parameter.
3. Five runs modifying the strength of the perturbation parameter from 0.5 (default
value) to 0.7.
4. Five runs modifying the strength of the perturbation parameter from 0.5 (default
value) to 0.9.

In all cases, the number of unsuccessful iterations to stop parameter (-nstop) value was
changed from 100 (default value) to 500. The tree with the highest likelihood score was
chosen (log-likelihood: -127052.963, -pers 0.5 and -nstop 500). Support for the nodes
was evaluated using two approaches: the aBayes test (Anisimova et al. 2011) and the
ultrafast bootstrap procedure (Hoang et al. 2018). Nicotinamide Nucleotide
Transhydrogenase (NNT) protein sequences from the human (*Homo sapiens*), mouse
(*Mus musculus*), zebrafish (*Danio rerio*) and spotted gar (*Lepisosteus oculatus*) were
used as outgroups.

Assessment of conserved synteny
We examined genes found upstream and downstream of the sirtuin genes. For
comparative purposes, we used the estimates of orthology and paralogy derived from the
Ensembl Compara database (Herrero et al. 2016); these estimates are obtained from a
pipeline that considers both synteny and phylogeny to generate orthology mappings.
These predictions were visualized using the program Genomicus v100.01 (Nguyen et al.
2018). Our assessments were performed in humans (*Homo sapiens*), chicken (*Gallus*

gallus), gharial (*Gavialis gangeticus*), red-eared slider (*Trachemys scripta*), green anole

(*Anolis carolinensis*), tropical clawed frog (*Xenopus tropicalis*), coelacanth (*Latimeria*

chalumnae), spotted gar (*Lepisosteus oculatus*), and elephant shark (*Callorhinchus milii*).

Dot-plots

We retrieved the chromosomal region containing the SIRT3-like gene of the tropical

clawed frog (*Xenopus tropicalis*, Ch3) including 6.7 kb up- and downstream, and the

corresponding syntenic region in the human (*Homo sapiens*, Chr12), opossum

(*Monodelphis domestica*, Chr8), chicken (*Gallus gallus*, Chr1), gharial (*Gavialis*

gangeticus, NW_017729022.1), red-eared slider (*Trachemys scripta*, Chr1), and green

anole (*Anolis carolinensis*, Chr5) based on the location of the flanking genes (RIC8B and

TMEM263). We aligned SIRT3-like syntenic regions using Advanced PipMaker (Schwartz

et al. 2000).

Transcript abundance analyses

Sirtuin transcript abundance was measured from a representative sample of vertebrates

including the elephant shark (*Callorhinchus milii*), zebrafish (*Danio rerio*), tropical clawed

frog (*Xenopus tropicalis*), anole lizard (*Anolis carolinensis*), and human (*Homo sapiens*).

RNASeq libraries from brain, heart, kidney, liver, muscle, ovary, and testis from each

species were gathered from the NCBI Short Read Archive (SRA)(Leinonen et al. 2011).

Accession numbers for species and tissue specific libraries can be found in Supplemental

Table S2. Reference transcript sequences were collected from Ensembl v.100 (Yates et

al. 2020) and we removed sequences that were shorter than 100 bp. For each library

adapters were removed using Trimmomatic 0.38 (Bolger et al. 2014) and reads were

filtered for quality using the parameters HEADCROP:5, SLIDINGWINDOW:5:30, and

MINLEN:50. We mapped quality filtered paired-end RNAseq reads back to reference

sequences using Bowtie 1.2.2 (Langmead et al. 2009) and default parameters of RSEM

(Li and Dewey 2011). Transcripts with < 10 mapped reads across all seven tissues per

species were removed prior to normalization. Normalization of raw read counts for each

species was performed using the estimateSizeFactors and estimateDispersions functions

in DESeq2 v1.26 (Love et al. 2014). If multiple SIRT transcripts were present, we

presented the expression data from the transcript with the most mapped reads.

Protein structure homology modeling

Protein structure homology modeling was performed using the SWISS-MODEL server

(https://swissmodel.expasy.org/) (Waterhouse et al. 2018). Structural figures were

prepared with PyMOL Molecular Graphics System, Version 2.0.6 Schrödinger, LLC.

Recombinant cDNA Constructs

For expression in mammalian cells, a codon-optimized mammalian expression construct

encoding full-length (amino acids 1-372) elephant shark (*Callorhinchus milii*) SIRT3-like

(Supplementary table S1), cloned in-frame into the *Eco*RI and *Sal*I sites of the pCMV-

3Tag-4a vector followed by vector encoding sequence of three successive Myc epitope

tags before a stop codon, was acquired from GenScript (Piscataway, NJ). For the

generation of a codon-optimized bacterial expression construct, a cDNA encoding N-

terminal truncated (amino acids 95-372) elephant shark SIRT3-like, cloned in-frame into

the *Eco*RI and *Sal*I sites of the pGEX-4T-1 vector, was also acquired from GenScript.

This cDNA was subsequently cloned in-frame into the *Eco*RI and *Sal*I sites of the pGST-

Parallel-1 vector (Sheffield et al. 1999). The nucleotide sequence of all recombinant

constructs was confirmed by dideoxy sequencing using the AUSTRAL-omics core facility

at Universidad Austral de Chile (https://australomics.cl/).

Cell culture, cell transfection and preparation of protein extracts

H4 human neuroglioma cells, HeLa human cervix adenocarcinoma cells and MDA-MB-

231 human mammary gland adenocarcinoma cells were obtained from the American

Type Culture Collection (Manassas, VA). H4 and HeLa cells were maintained in

Dulbecco's modified Eagle's medium (DMEM), and MDA-MB-231 in DMEM-F12

(ThermoFisher, Waltham, MA). For all cell lines, media were supplemented with 10%

heat-inactivated fetal bovine serum, 100 U/ml penicillin, 100 µg/ml streptomycin

(ThermoFisher), and 5 µg/ml plasmocin (InvivoGen, San Diego, CA), and cells were

cultured in a humidified incubator with 5% CO_2 at 37 °C. For transient transfections, cells

were either seeded on top of glass coverslips on 24-well plates or on 6-well plates. When

cells were ~60% confluent, transfections were performed with Lipofectamine 2000

(ThermoFisher), according to the manufacturer's instructions. Preparation of protein

extracts from cultured, transfected or non-transfected cells was performed using methods

described elsewhere (Tenorio et al. 2016).

Antibodies and cell reagents

We used the following antibodies: mouse clone 9E10 to the c-Myc epitope (Covance,

Princeton, NJ), mouse clone BA3R to β-actin (ThermoFisher), rabbit polyclonal to p63

(Merck, Germany; cat # HPA001225), and sheep polyclonal to TGN46 (Bio-Rad

Laboratories, Hercules, CA; cat # AHP500G). The following fluorochrome-conjugated

antibodies were from ThermoFisher: Alexa Fluor-488–conjugated donkey anti mouse

IgG, Alexa Fluor-594–conjugated donkey anti rabbit IgG and Alexa Fluor-647-conjugated

donkey anti sheep IgG. HRP-conjugated secondary antibodies were from Jackson

ImmunoResearch (West Grove, PA). Depending on their reactivity, primary antibodies

were used at a dilution 1/200 to 1/2000. HRP-conjugated secondary antibodies were used

at dilutions 1/5000 to 1/20000 also depending on their reactivity. All Alexa Fluor-

conjugated secondary antibodies were used at a dilution 1/1000. The fluorescent nuclear

stain 4',6-diamidino-2-phenylindole (DAPI) was also from ThermoFisher.

Protein Electrophoresis, Immunoblotting and Immunofluorescence Microscopy

SDS-PAGE and immunoblotting were performed as described (Bustamante et al. 2020).

For fluorescence microscopy, cells grown on glass coverslips were transfected, and after

16-h cells were left untreated or treated for 30 min with MitoTracker™ Orange CMTMRos

(ThermoFisher) according to the manufacturer's instructions. After washing with

phosphate buffered saline (PBS) supplemented with 0.1 mM $CaCl_2$ and 1 mM $MgCl_2$

(PBS-CM), cells were fixed in 4% paraformaldehyde for 30 min at room temperature,

permeabilized with 0.2% Triton X-100 in PBS-CM for 15 min at room temperature, and

incubated with blocking solution (0.2% gelatin in PBS-CM) for 10 min at room

temperature. Cells were incubated simultaneously either with antibodies to the c-Myc

epitope (1/200) and to TGN46 (1/1000) or to the c-Myc epitope, to TGN46 and to p63

(1/600) for 30 min at 37 ºC in a humidified chamber. After washing coverslips with PBS,

cells were incubated as before, but with fluorochrome-conjugated antibodies, followed by

PBS-washing, and coverslips were mounted onto glass slides with Fluoromount-G

mounting medium (ThermoFisher). Fluorescence microscopy images were acquired with

an AxioObserver.D1 microscope equipped with a PlanApo 63x oil immersion objective

(NA 1.4), and an AxioCam MRm digital camera (Carl Zeiss). To prepare figures, 12-bit

images were processed with ImageJ software (version 1.44o; Wayne Rasband, NIH,

http://imagej.nih.gov) and Adobe Photoshop CS3 software (Adobe Systems, Mountain

View, CA).

Expression and Purification of Recombinant N-terminal truncated elephant shark SIRT3-like protein

659
660

Recombinant, N-terminal truncated elephant shark SIRT3-like (ΔNT-SIRT3-like; amino

acids 95-372) tagged with an N-terminal glutathione S-transferase (GST) followed by a

tobacco etch virus (TEV) protease cleavage site was expressed and purified using a

method described previously (Ross et al. 2014), with some modifications. Briefly,

expression in *E. coli* B834(DE3) (Novagen, Madison, WI) was induced with 0.25 mM IPTG

at 25 °C for 16 h. Pellets of bacteria were resuspended in homogenization buffer (50 mM

Tris HCl, 0.5 M NaCl, 5 mM EDTA, 5 mM β-mercaptoethanol, and 2 mM

phenylmethylsulfonyl fluoride, pH 8.0) and lysed by sonication. The clarified supernatant

was purified on glutathione-Sepharose 4B (GE Healthcare). After removal of the GST

moiety by TEV cleavage, and sequential further passage through glutathione-Sepharose

4B and Ni-NTA (QIAGEN) resins, ΔNT-SIRT3-like was further purified on a Superdex 200

column (GE Healthcare) equilibrated in storage buffer (50 mM Tris HCl, 150 mM NaCl, 1

mM DTT). Aliquots of purified protein were kept at -80 ºC until use.

Deacetylation activity

To characterize deacetylation activity of recombinant, N-terminal truncated elephant

shark SIRT3-like protein (ΔNT-SIRT3-like), we used a fluorometric SIRT3 activity assay

kit (Abcam, cat # ab156067; Cambridge, UK), according to the manufacturer's

instructions. This assay allows detection of a fluorescent signal upon deacetylation of an
acetylated substrate peptide for recombinant human SIRT3. The intensity of fluorescence
was measured on a fluorometric microplate reader (Varioskan Flash, ThermoFisher) with
excitation set at 350 nm and emission detection set at 450 nm. To analyze the
deacetylation activity of N-terminal truncated elephant shark SIRT3-like protein, first we
determined its optimum temperature that resulted to be ~24 ºC. Subsequent analyses
were performed by comparing the activity of N-terminal truncated elephant shark SIRT3-
like protein at 24 ºC to that of human SIRT3, which is provided by the assay kit, at 37 ºC.
K_m and V_{max} were obtained by varying the concentration of the fluorogenic acetylated
substrate peptide provided by the assay kit. This assay was also used to determine the
effect of nicotinamide (NAM), quercetin (QUE) and resveratrol (REV; Sigma-Aldrich) on
the deacetylation activity of ΔNT-SIRT3-like.

ATP and ROS content

ATP levels were quantified in cell lysates using a luciferin/luciferase bioluminescence
assay (ATP determination kit, Molecular Probes cat # A22066, Thermo Fisher Scientific)
as previously described (Jara et al. 2018). Briefly, cells were lysed in HEPES buffer (125
mM NaCl, 25 mM NaF, 1 mM EDTA, 1 mM EGTA, 1% NP-40, 25 mM HEPES, pH 7.4)
supplemented with a protease inhibitor mixture (cat # 78429, Thermo Fisher Scientific)
and centrifuged at 12000 rpm using a refrigerated centrifuge (Eppendorf model 5424R)
for 15 min at 4 ºC. Proteins in the supernatant were quantified, and 10 µl of lysate were
used to determine ATP levels according to the kit manufacturer's instructions. The ATP
level in each sample was calculated using standard curves and normalized to the protein
concentration in each sample. ROS content was determined in cell lysates using 25 µM
CM-H2DCFDA (DCF; a fluorogenic indicator of ROS), as previously described (Jara et
al. 2018; Olesen et al. 2020). The intensity of fluorescence was measured on a
fluorescence plate reader (Biotek Synergy HT, Agilent, Santa Clara, CA) with excitation
set at 485 nm and emission detection set at 530 nm. Briefly, in a dark 96-well plate, 25
µg of cell lysate proteins were incubated with CM-H2DCFDA (DCF) dye with shaking for
5 min at room temperature, followed by determination of the fluorescence of each sample
that was subtracted by the fluorescence of the respective blank.

Statistical Analysis

In the case of the deacetylation, ATP content and ROS content assays, quantification

was performed from at least three independent experiments/measurements. Statistical

analysis was performed using Microsoft Excel for Mac 2011 (Microsoft Corporation).

When appropriate, results are represented in graphs depicting the mean ± standard

deviation. Statistical significance was determined by two-tailed, paired t-test. P-values >

0.05 or ≤ 0.05 were regarded as not statistically significant or statistically significant,

respectively. In the figures, P-values between 0.01 and 0.05 are indicated with one

asterisk, P-values between 0.001 and 0.01 are indicated with two asterisks, and P-values

less than 0.001 are indicated with three asterisks.

Data accessibility

723 Data and supplementary material are available online:

Acknowledgements

This work was supported by the Fondo Nacional de Desarrollo Científico y Tecnológico

from Chile (FONDECYT 1210471) and Millennium Nucleus of Ion Channels Associated

Diseases (MiNICAD), Iniciativa Científica Milenio, Ministry of Economy, Development and

Tourism from Chile to JCO, Fondo Nacional de Desarrollo Científico y Tecnológico from

Chile (FONDECYT 1211481) to GAM, National Science Foundation (EPS-0903787, DBI-

1262901 and DEB-1354147) to F.G.H., Financiamiento Basal No.

ANID/BASAL/ACE210009 to PVB and No. ANID/BASAL/FB210008 to PVB and CTR,

Fondap-Ideal Grant N° 15150003 to LV-C, ANID-Millennium Science Initiative Program-

Center ICM-ANID ICN2021_002 to LV-C, Fondo Nacional de Desarrollo Científico y

Tecnológico from Chile (FONDECYT 1180957) to FJM, Vicerrectoría de Investigación,

Desarrollo y Creación Artística, Universidad Austral de Chile to JCO, FJM, LVC, GAM.

We thank Stefanie Teuber, John Quiroga and Rafael Burgos, Instituto de Farmacología

y Morfofisiología, Facultad de Ciencias Veterinarias, Universidad Austral de Chile, for

Contributions

JCO and GAM designed the study. JCO, MWV, FGH, KZ, CM, CL, VC, FJM, PVB, CTR, GAM, collected and/or analyzed data. JCO, PVB and GAM wrote the manuscript. JCO, MWV, FGH, KZ, CM, CL, VC, LV-C, FJM, PVB, CTR and GAM, reviewed and edited the manuscript. All authors contributed to the article and approved the submitted version.

Ethics declarations

Conflict of interest disclosure

The authors declare they have no conflict of interest relating to the content of this article. Juan C. Opazo is a recommender for PCI Evol Biol and PCI Genomics.

Data availability

Data and supplementary material are available online at zenodo.org

References

Albalat R, Cañestro C. 2016. Evolution by gene loss. *Nat. Rev. Genet.* 17:379–391.

Albertson RC, Craig Albertson R, Cresko W, William Detrich H, Postlethwait JH. 2009. Evolutionary mutant models for human disease. *Trends in Genetics* [Internet] 25:74–81. Available from: http://dx.doi.org/10.1016/j.tig.2008.11.006

Altenhoff AM, Train C-M, Gilbert KJ, Mediratta I, Mendes de Farias T, Moi D, Nevers Y, Radoykova H-S, Rossier V, Warwick Vesztrocy A, et al. 2021. OMA orthology in 2021: website overhaul, conserved isoforms, ancestral gene order and more. *Nucleic Acids Res.* 49:D373–D379.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J. Mol. Biol.* 215:403–410.

Anderson KA, Green MF, Huynh FK, Wagner GR, Hirschey MD. 2014. SnapShot: Mammalian Sirtuins. *Cell* 159:956–956.e1.

Anisimova M, Gil M, Dufayard J-F, Dessimoz C, Gascuel O. 2011. Survey of branch support methods demonstrates accuracy, power, and robustness of fast likelihood-based approximation schemes. *Syst. Biol.* 60:685–699.

Bause AS, Haigis MC. 2013. SIRT3 regulation of mitochondrial oxidative stress. *Exp. Gerontol.* 48:634–639.

Bilyk KT, Zhuang X, Vargas-Chacoff L, Cheng C-HC. 2021. Evolution of chaperome gene expression and regulatory elements in the antarctic notothenioid fishes. *Heredity* 126:424–441.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120.

Brown K, Xie S, Qiu X, Mohrin M, Shin J, Liu Y, Zhang D, Scadden DT, Chen D. 2013. SIRT3 reverses aging-associated degeneration. *Cell Rep.* 3:319–327.

Bustamante HA, Cereceda K, González AE, Valenzuela GE, Cheuquemilla Y, Hernández S, Arias-Muñoz E, Cerda-Troncoso C, Bandau S, Soza A, et al. 2020. The Proteasomal Deubiquitinating Enzyme PSMD14 Regulates Macroautophagy by Controlling Golgi-to-ER Retrograde Transport. *Cells* [Internet] 9. Available from: http://dx.doi.org/10.3390/cells9030777

Castro LFC, Lopes-Marques M, Gonçalves O, Wilson JM. 2012. The evolution of pepsinogen C genes in vertebrates: duplication, loss and functional diversification. *PLoS One* 7:e32852.

Céspedes HA, Zavala K, Vandewege MW, Opazo JC. 2017. Evolution of the α-adrenoreceptors in vertebrates: ADRA2D is absent in mammals and crocodiles.

Gen. Comp. Endocrinol. 250:85–94.

Costantini S, Sharma A, Raucci R, Costantini M, Autiero I, Colonna G. 2013. Genealogy
of an ancient protein family: the Sirtuins, a family of disordered members. *BMC*
Evol. Biol. 13:60.

Delsuc F, Brinkmann H, Chourrout D, Philippe H. 2006. Tunicates and not
cephalochordates are the closest living relatives of vertebrates. *Nature* 439:965–
968.

Delsuc F, Tsagkogeorga G, Lartillot N, Philippe H. 2008. Additional molecular support
for the new chordate phylogeny. *Genesis* 46:592–604.

Di Emidio G, Falone S, Artini PG, Amicarelli F, D'Alessandro AM, Tatone C. 2021.
Mitochondrial Sirtuins in Reproduction. *Antioxidants (Basel)* [Internet] 10. Available
from: http://dx.doi.org/10.3390/antiox10071047

Félix M-A, Barkoulas M. 2015. Pervasive robustness in biological systems. *Nat. Rev.*
Genet. 16:483–496.

Frye RA. 2000. Phylogenetic classification of prokaryotic and eukaryotic Sir2-like
proteins. *Biochem. Biophys. Res. Commun.* 273:793–798.

Frye RA. 2006. Evolution of Sirtuins From Archaea to Vertebrates. *Histone*
Deacetylases [Internet]:183–202. Available from: http://dx.doi.org/10.1385/1-59745-
024-3:183

Fujita Y, Yamashita T. 2018. Sirtuins in Neuroendocrine Regulation and Neurological
Diseases. *Front. Neurosci.* 12:778.

Glover N, Dessimoz C, Ebersberger I, Forslund SK, Gabaldón T, Huerta-Cepas J,
Martin M-J, Muffato M, Patricio M, Pereira C, et al. 2019. Advances and
Applications in the Quest for Orthologs. *Mol. Biol. Evol.* 36:2157–2164.

Glozak MA, Sengupta N, Zhang X, Seto E. 2005. Acetylation and deacetylation of non-
histone proteins. *Gene* 363:15–23.

Goldberg T, Hecht M, Hamp T, Karl T, Yachdav G, Ahmed N, Altermann U, Angerer P,
Ansorge S, Balasz K, et al. 2014. LocTree3 prediction of localization. *Nucleic Acids*
Res. 42:W350–W355.

Greiss S, Gartner A. 2009. Sirtuin/Sir2 phylogeny, evolutionary considerations and
structural conservation. *Mol. Cells* 28:407–415.

Haigis MC, Sinclair DA. 2010. Mammalian sirtuins: biological insights and disease
relevance. *Annu. Rev. Pathol.* 5:253–295.

Hara Y, Yamaguchi K, Onimaru K, Kadota M, Koyanagi M, Keeley SD, Tatsumi K,

Tanaka K, Motone F, Kageyama Y, et al. 2018. Shark genomes provide insights
into elasmobranch evolution and the origin of vertebrates. *Nat Ecol Evol* 2:1761–
1771.

Helsen J, Voordeckers K, Vanderwaeren L, Santermans T, Tsontaki M, Verstrepen KJ,
Jelier R. 2020. Gene Loss Predictably Drives Evolutionary Adaptation. *Mol. Biol.*
Evol. 37:2989–3002.

Herrero J, Muffato M, Beal K, Fitzgerald S, Gordon L, Pignatelli M, Vilella AJ, Searle
SMJ, Amode R, Brent S, et al. 2016. Ensembl comparative genomics resources.
Database [Internet] 2016. Available from:
http://dx.doi.org/10.1093/database/bav096

Himmel NJ, Gray TR, Cox DN. 2020. Phylogenetics Identifies Two Eumetazoan TRPM
Clades and an Eighth TRP Family, TRP Soromelastatin (TRPS). *Mol. Biol. Evol.*
37:2034–2044.

Hirschey MD. 2011. Old enzymes, new tricks: sirtuins are NAD(+)-dependent de-
acylases. *Cell Metab.* 14:718–719.

Hirschey MD, Shimazu T, Goetzman E, Jing E, Schwer B, Lombard DB, Grueter CA,
Harris C, Biddinger S, Ilkayeva OR, et al. 2010. SIRT3 regulates mitochondrial
fatty-acid oxidation by reversible enzyme deacetylation. *Nature* 464:121–125.

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2:
Improving the Ultrafast Bootstrap Approximation. *Mol. Biol. Evol.* 35:518–522.

Howe KL, Achuthan P, Allen J, Allen J, Alvarez-Jarreta J, Amode MR, Armean IM, Azov
AG, Bennett R, Bhai J, et al. 2021. Ensembl 2021. *Nucleic Acids Res.* 49:D884–
D891.

Iwabe N, Hara Y, Kumazawa Y, Shibamoto K, Saito Y, Miyata T, Katoh K. 2005. Sister
group relationship of turtles to the bird-crocodilian clade revealed by nuclear DNA-
coded proteins. *Mol. Biol. Evol.* 22:810–813.

Jara C, Aránguiz A, Cerpa W, Tapia-Rojas C, Quintanilla RA. 2018. Genetic ablation of
tau improves mitochondrial function and cognitive abilities in the hippocampus.
Redox Biol 18:279–294.

Kabiljo J, Murko C, Pusch O, Zupkovitz G. 2019. Spatio-temporal expression profile of
sirtuins during aging of the annual fish Nothobranchius furzeri. *Gene Expression*
Patterns [Internet] 33:11–19. Available from:
http://dx.doi.org/10.1016/j.gep.2019.05.001

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017.
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat.*
Methods 14:587–589.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol. Biol. Evol.* 30:772–780.

Kumar S, Stecher G, Suleski M, Hedges SB. 2017. TimeTree: A Resource for
Timelines, Timetrees, and Divergence Times. *Mol. Biol. Evol.* 34:1812–1819.

Kuraku S. 2013. Impact of asymmetric gene repertoire between cyclostomes and
gnathostomes. *Semin. Cell Dev. Biol.* 24:119–127.

Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biol.* 10:R25.

Leinonen R, Sugawara H, Shumway M, International Nucleotide Sequence Database
Collaboration. 2011. The sequence read archive. *Nucleic Acids Res.* 39:D19–D21.

Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. *BMC Bioinformatics* [Internet] 12. Available
from: http://dx.doi.org/10.1186/1471-2105-12-323

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15:550.

Lynch M, Conery JS. 2000. The Evolutionary Fate and Consequences of Duplicate
Genes. *Science* [Internet] 290:1151–1155. Available from:
http://dx.doi.org/10.1126/science.290.5494.1151

McCormick SD. 1993. Methods for nonlethal gill biopsy and measurement of Na+, K+-
ATPase activity. *Can. J. Fish. Aquat. Sci.* 50:656–658.

McDonnell E, Peterson BS, Bomze HM, Hirschey MD. 2015. SIRT3 regulates
progression and development of diseases of aging. *Trends Endocrinol. Metab.*
26:486–492.

Michan S, Sinclair D. 2007. Sirtuins in mammals: insights into their biological function.
Biochem. J 404:1–13.

Nei M, Rooney AP. 2005. Concerted and birth-and-death evolution of multigene
families. *Annu. Rev. Genet.* 39:121–152.

Nguyen NTT, Vincens P, Roest Crollius H, Louis A. 2018. Genomicus 2018: karyotype
evolutionary trees and on-the-fly synteny computing. *Nucleic Acids Res.* 46:D816–
D822.

North BJ, Verdin E. 2004. Sirtuins: Sir2-related NAD-dependent protein deacetylases.
Genome Biol. 5:224.

Ohno S. 1985. Dispensable genes. *Trends Genet.* 1:160–164.

Olesen MA, Torres AK, Jara C, Murphy MP, Tapia-Rojas C. 2020. Premature synaptic
mitochondrial dysfunction in the hippocampus during aging contributes to memory
loss. *Redox Biol* 34:101558.

Olson MV. 1999. When less is more: gene loss as an engine of evolutionary change.
Am. J. Hum. Genet. 64:18–23.

Onyango P, Celic I, McCaffery JM, Boeke JD, Feinberg AP. 2002. SIRT3, a human
SIR2 homologue, is an NAD-dependent deacetylase localized to mitochondria.
Proc. Natl. Acad. Sci. U. S. A. 99:13653–13658.

Opazo JC, Hoffmann FG, Zavala K, Edwards SV. 2019. Evolution of the DAN gene
family in vertebrates. *Evolutionary Biology*:185.

Opazo JC, Kuraku S, Zavala K, Toloza-Villalobos J, Hoffmann FG. 2019. Evolution of
nodal and nodal-related genes and the putative composition of the heterodimers
that trigger the nodal pathway in vertebrates. *Evolution & Development* [Internet]
21:205–217. Available from: http://dx.doi.org/10.1111/ede.12292

Opazo JC, Vandewege MW, Gutierrez J, Zavala K, Vargas-Chacoff L, Morera FJ,
Mardones GA. 2021. Independent duplications of the Golgi phosphoprotein 3
oncogene in birds. *Sci. Rep.* 11:12483.

Pereira TCB, Rico EP, Rosemberg DB, Schirmer H, Dias RD, Souto AA, Bonan CD,
Bogo MR. 2011. Zebrafish as a model organism to evaluate drugs potentially able
to modulate sirtuin expression. *Zebrafish* 8:9–16.

Qiu H, Hildebrand F, Kuraku S, Meyer A. 2011. Unresolved orthology and peculiar
coding sequence properties of lamprey genes: the KCNA gene family as test case.
BMC Genomics 12:325.

Qiu X, Brown K, Hirschey MD, Verdin E, Chen D. 2010. Calorie restriction reduces
oxidative stress by SIRT3-mediated SOD2 activation. *Cell Metab.* 12:662–667.

Rajabi N, Galleano I, Madsen AS, Olsen CA. 2018. Targeting Sirtuins: Substrate
Specificity and Inhibitor Design. *Prog. Mol. Biol. Transl. Sci.* 154:25–69.

Ramos-Vicente D, Ji J, Gratacòs-Batlle E, Gou G, Reig-Viader R, Luís J, Burguera D,
Navas-Perez E, García-Fernández J, Fuentes-Prior P, et al. 2018. Metazoan
evolution of glutamate receptors reveals unreported phylogenetic groups and
divergent lineage-specific events. *Elife* [Internet] 7. Available from:
http://dx.doi.org/10.7554/eLife.35774

Ross BH, Lin Y, Corales EA, Burgos PV, Mardones GA. 2014. Structural and functional
characterization of cargo-binding sites on the μ4-subunit of adaptor protein complex
4. *PLoS One* 9:e88147.

Saravia J, Paschke K, Oyarzún-Salazar R, Cheng C-HC, Navarro JM, Vargas-Chacoff

L. 2021. Effects of warming rates on physiological and molecular components of
response to CTMax heat stress in the Antarctic fish Harpagifer antarcticus. *J.*
Therm. Biol. 99:103021.

Sauve AA, Schramm VL. 2003. Sir2 regulation by nicotinamide results from switching
between base exchange and deacetylation chemistry. *Biochemistry* 42:9249–9256.

Scher MB, Vaquero A, Reinberg D. 2007. SirT3 is a nuclear NAD+-dependent histone
deacetylase that translocates to the mitochondria upon cellular stress. *Genes Dev.*
21:920–928.

Scholte LLS, Mourão MM, Pais FS-M, Melesina J, Robaa D, Volpini AC, Sippl W, Pierce
RJ, Oliveira G, Nahum LA. 2017. Evolutionary relationships among protein lysine
deacetylases of parasites causing neglected diseases. *Infect. Genet. Evol.* 53:175–
188.

Schwartz S, Zhang Z, Frazer KA, Smit A, Riemer C, Bouck J, Gibbs R, Hardison R,
Miller W. 2000. PipMaker--a web server for aligning two genomic DNA sequences.
Genome Res. 10:577–586.

Schwer B, North BJ, Frye RA, Ott M, Verdin E. 2002. The human silent information
regulator (Sir)2 homologue hSIRT3 is a mitochondrial nicotinamide adenine
dinucleotide-dependent deacetylase. *J. Cell Biol.* 158:647–657.

Shahgaldi S, Kahmini FR. 2021. A comprehensive review of Sirtuins: With a major focus
on redox homeostasis and metabolism. *Life Sci.* 282:119803.

Sharma S, Ciufo S, Starchenko E, Darji D, Chlumsky L, Karsch-Mizrachi I, Schoch CL.
2019. The NCBI BioCollections Database. *Database* [Internet] 2019. Available
from: http://dx.doi.org/10.1093/database/baz057

Sheffield P, Garrard S, Derewenda Z. 1999. Overcoming expression and purification
problems of RhoGDI using a family of "parallel" expression vectors. *Protein Expr.*
Purif. 15:34–39.

Simó-Mirabet P, Bermejo-Nogales A, Calduch-Giner JA, Pérez-Sánchez J. 2017.
Tissue-specific gene expression and fasting regulation of sirtuin family in gilthead
sea bream (Sparus aurata). *Journal of Comparative Physiology B* [Internet]
187:153–163. Available from: http://dx.doi.org/10.1007/s00360-016-1014-0

Slade KM, Freggiaro S, Cottrell KA, Smith JJ, Wiley EA. 2011. Sirtuin-mediated nuclear
differentiation and programmed degradation in Tetrahymena. *BMC Cell Biol.* 12:40.

Smith JJ, Kuraku S, Holt C, Sauka-Spengler T, Jiang N, Campbell MS, Yandell MD,
Manousaki T, Meyer A, Bloom OE, et al. 2013. Sequencing of the sea lamprey
(Petromyzon marinus) genome provides insights into vertebrate evolution. *Nat.*
Genet. 45:415–421, 421e1–e2.

Somero GN. 2004. Adaptation of enzymes to temperature: searching for basic
"strategies." *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 139:321–333.

Stelzer G, Rosen N, Plaschkes I, Zimmerman S, Twik M, Fishilevich S, Stein TI, Nudel
R, Lieder I, Mazor Y, et al. 2016. The GeneCards Suite: From Gene Data Mining to
Disease Genome Sequence Analyses. *Curr. Protoc. Bioinformatics* 54:1.30.1–
1.30.33.

Tao R, Coleman MC, Pennington JD, Ozden O, Park S-H, Jiang H, Kim H-S, Flynn CR,
Hill S, Hayes McDonald W, et al. 2010. Sirt3-mediated deacetylation of
evolutionarily conserved lysine 122 regulates MnSOD activity in response to stress.
Mol. Cell 40:893–904.

Tatone C, Di Emidio G, Barbonetti A, Carta G, Luciano AM, Falone S, Amicarelli F.
2018. Sirtuins in gamete biology and reproductive physiology: emerging roles and
therapeutic potential in female and male infertility. *Hum. Reprod. Update* 24:267–
289.

Tenorio MJ, Ross BH, Luchsinger C, Rivera-Dictter A, Arriagada C, Acuña D, Aguilar M,
Cavieres V, Burgos PV, Ehrenfeld P, et al. 2016. Distinct Biochemical Pools of
Golgi Phosphoprotein 3 in the Human Breast Cancer Cell Lines MCF7 and MDA-
MB-231. *PLoS One* 11:e0154719.

Trifinopoulos J, Nguyen L-T, von Haeseler A, Minh BQ. 2016. W-IQ-TREE: a fast online
phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res.* 44:W232–
W235.

Tseng AHH, Shieh S-S, Wang DL. 2013. SIRT3 deacetylates FOXO3 to protect
mitochondria against oxidative damage. *Free Radic. Biol. Med.* 63:222–234.

Vassilopoulos A, Fritz KS, Petersen DR, Gius D. 2011. The human sirtuin family:
evolutionary divergences and functions. *Hum. Genomics* 5:485–496.

Vassilopoulos A, Pennington JD, Andresson T, Rees DM, Bosley AD, Fearnley IM, Ham
A, Flynn CR, Hill S, Rose KL, et al. 2014. SIRT3 deacetylates ATP synthase F1
complex proteins in response to nutrient- and exercise-induced stress. *Antioxid.*
Redox Signal. 21:551–564.

Vazquez BN, Vaquero A, Schindler K. 2020. Sirtuins in female meiosis and in
reproductive longevity. *Mol. Reprod. Dev.* 87:1175–1187.

Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, Heer FT, de
Beer TAP, Rempfer C, Bordoli L, et al. 2018. SWISS-MODEL: homology modelling
of protein structures and complexes. *Nucleic Acids Res.* 46:W296–W303.

Wichmann IA, Zavala K, Hoffmann FG, Vandewege MW, Corvalán AH, Amigo JD,
Owen GI, Opazo JC. 2016. Evolutionary history of the reprimo tumor suppressor
gene family in vertebrates with a description of a new reprimo gene lineage. *Gene*

591:245–254.

Wood JG, Rogina B, Lavu S, Howitz K, Helfand SL, Tatar M, Sinclair D. 2004. Sirtuin
activators mimic caloric restriction and delay ageing in metazoans. *Nature*
430:686–689.

Yang L, Ma X, He Y, Yuan C, Chen Q, Li G, Chen X. 2017. Sirtuin 5: a review of
structure, known inhibitors and clues for developing new inhibitors. *Sci. China Life*
Sci. 60:249–256.

Yates AD, Achuthan P, Akanni W, Allen J, Allen J, Alvarez-Jarreta J, Amode MR,
Armean IM, Azov AG, Bennett R, et al. 2020. Ensembl 2020. *Nucleic Acids Res.*
48:D682–D688.

You W, Zheng W, Weiss S, Chua KF, Steegborn C. 2019. Structural basis for the
activation and inhibition of Sirtuin 6 by quercetin and its derivatives. *Sci. Rep.*
9:19176.

Zhang J, Xiang H, Liu J, Chen Y, He R-R, Liu B. 2020. Mitochondrial Sirtuin 3: New
emerging biological function and therapeutic target. *Theranostics* 10:8315–8342.

Zhao E, Hou J, Ke X, Abbas MN, Kausar S, Zhang L, Cui H. 2019. The Roles of Sirtuin
Family Proteins in Cancer Progression. *Cancers* [Internet] 11. Available from:
http://dx.doi.org/10.3390/cancers11121949

Zhao H-C, Ding T, Ren Y, Li T-J, Li R, Fan Y, Yan J, Zhao Y, Li M, Yu Y, et al. 2016.
Role of Sirt3 in mitochondrial biogenesis and developmental competence of human
in vitro matured oocytes. *Hum. Reprod.* 31:607–622.

1038