Quantifying Evolutionary Importance of Protein Sites:

A Tale of Two Measures

Avital Sharir-Ivry[1], Yu Xia[1*]

[1]Department of Bioengineering, McGill University, Montreal, QC, Canada

Corresponding author:

Yu Xia

Department of Bioengineering

McGill University

Montreal, Quebec H3A 0E9, Canada

Email: brandon.xia@mcgill.ca

Office: McConnell Engineering Building, Room 360

Phone: 514-398-5026

Abstract

A key challenge in evolutionary biology is the accurate quantification of selective pressure on proteins and other biological macromolecules at single-site resolution. The evolutionary importance of a protein site under purifying selection is typically measured by the degree of conservation of the protein site itself. A possible alternative measure is the strength of the site-induced conservation gradient in the rest of the protein structure. However, the quantitative relationship between these two measures remains unknown. Here, we show that despite major differences, there is a strong linear relationship between the two measures such that more conserved protein sites also induce stronger conservation gradient in the rest of the protein. This linear relationship is universal as it holds for different types of proteins and functional sites in proteins. Our results show that the strong selective pressure acting on the functional site in general percolates through the rest of the protein via residue-residue contacts. Surprisingly however, catalytic sites in enzymes are the principal exception to this rule. Catalytic sites induce significantly stronger conservation gradients in the rest of the protein than expected from the degree of conservation of the site alone. The unique requirement for the active site to selectively stabilize the transition state of the catalyzed chemical reaction imposes additional selective constraints on the rest of the enzyme.

Introduction

The evolutionary importance of protein sites under purifying selection can be quantified in two very different ways. The classical, "intrinsic" measure for the evolutionary importance of a protein site is the degree of conservation or evolutionary rate of the protein site itself. Protein residues experience different degrees of selective pressure as a result of the different roles they play in protein stability and function(Worth et al. 2009; Echave et al. 2016). For example, residues in a protein core are generally under stronger selective pressure than surface residues due to their importance in stabilizing the protein. Indeed, structural determinants such as solvent exposure(Overington et al. 1992; Conant and Stadler 2009; Franzosa and Xia 2009; Franzosa et al. 2013; Sharir-Ivry and Xia 2017) and degree of packing (Hamelryck 2005; Yeh et al. 2014; Marcos and Echave 2015) were shown to explain a large portion of the variability in the observed site-specific evolutionary rates. In addition, residues in functional sites such as catalytic sites(Bartlett et al. 2002) and ligand-binding sites are also under stronger selective pressure than non-functional residues.

An alternative, "extrinsic" measure for the evolutionary importance of a protein site is the conservation gradient the site exerts on the rest of the protein. Rather than quantifying evolutionary conservation of the protein site itself, this measure captures how the evolutionary conservation of other residues surrounding the site gradually decreases with distance from the site in the tertiary structure. Several studies have indicated the possibility for selective pressure to propagate from the functional site to the rest of the protein via physical interactions between neighboring residues in the three-dimensional structure (Tóth-Petróczy and Tawfik 2011; Jack et al. 2016; Nelson and Grishin 2016; Sharir-Ivry and Xia 2019a). While it is clear that the two

measures of the evolutionary importance of protein sites are substantially different, it remains

unknown how the two measures relate to each other.

In this paper, we addressed the fundamental question whether there is a direct relationship

between the intrinsic and extrinsic measures of the evolutionary importance of protein sites. We

have based our study on a dataset of homology-based structural models of the yeast

proteome(Franzosa and Xia 2009; Sharir-Ivry and Xia 2017). Despite their major differences, we

show here that there is a strong linear relationship between the degree of conservation of a

protein site and the conservation gradient induced from it. In other words, more conserved

protein sites tend to induce stronger conservation gradient in the rest of the protein, as selective

pressure acting on the protein site percolates via residue-residue interactions. This linear

evolutionary conservation-percolation relationship is universal in that it holds for different types

of proteins as well as for different types of functional sites in proteins. Remarkably however,

catalytic sites in enzymes are the only exception to this universal rule, as catalytic sites induce

significantly stronger long-range conservation gradient than other types of functional sites with

similar degrees of conservation. We conclude that for many different types of functional sites,

site-induced conservation gradient can be explained by the percolation of site-specific selective

pressure through the rest of the protein via residue-residue contacts. However, catalytic sites in

enzymes induce significantly stronger long-range conservation gradient in the rest of the protein

than expected from the percolation theory. This is likely due to the unique requirement for the

enzyme active site to selectively bind to and stabilize the transition state of the catalyzed

chemical reaction(Warshel et al. 2006).

Overall, we show that a more complete understanding of the selective pressure on protein sites

can be achieved by integrating the intrinsic measure of site-specific evolutionary conservation

with the extrinsic measure of site-induced conservation gradient, with potential implications in

protein design, study of disease mutations and better phylogenetic inference(Yang 1996).

Results

Evolutionary conservation gradient induced from a protein residue is linearly correlated with

the conservation rank of the residue

We based our study on a dataset of homology-based structural models of the *S. cerevisiae*

proteome. This basic dataset contains structural templates from the Protein Data Bank

(PDB)(Berman et al. 2000) mapped to ORFs of *S. cerevisiae* via sequence alignment (see Methods).

We had 1274 yeast ORFs with structural models from the PDB for which residue conservation

scores are available in ConSurf-DB (Goldenberg et al. 2009; Celniker et al. 2013). Each residue in

the dataset was ranked according to its relative conservation within the protein (the residue's

rank of conservation divided by the total number of residues). This normalized conservation rank

of a residue ranges from 0 to 1, with higher rank corresponding to lower conservation within the

protein. We have also calculated for each residue the Pearson correlation between the

conservation scores of all other residues in the protein and their distance from that reference

residue. This Pearson correlation between conservation and distance describes the degree of

percolation of the evolutionary conservation from a residue throughout the protein tertiary

structure, i.e., the 'long-range conservation gradient'. All residues were then binned into 20

equally spaced bins of conservation rank and the average conservation gradient for each bin was

calculated.

A clear negative linear trend is observed between the conservation rank of a residue and the

strength of the evolutionary conservation gradient induced from it (Fig 1a). The more a residue

is conserved within the protein (lower conservation rank), the stronger the evolutionary

conservation gradient it induces. The distribution of overall per-protein Pearson correlations

between conservation ranks and conservation gradients is mainly between -0.3 and -0.7, and the

average correlation in the dataset is -0.5 (Fig 1b), indicating that the linear relationship between

site-specific conservation and site-induced conservation gradient is high for all types of proteins.

We have also examined the relationship between site-specific conservation and site-induced

conservation gradient specifically for functional sites. We identified different functional sites in

the dataset: catalytic sites, non-catalytic ligand-binding sites (on enzymes and on nonenzymatic

proteins), protein-protein interaction sites, and allosteric sites. We identified catalytic sites using

the Catalytic Site Atlas(Furnham et al. 2014), ligand-binding sites using BioLip(Yang et al. 2013),

allosteric sites using the Allosteric Database (ASD)(Huang et al. 2011; Shen et al. 2016) and

protein-protein interaction sites using our previous protocol of identifying protein-protein

interfaces in the yeast proteome(Franzosa and Xia 2009). Relative solvent accessibility (RSA) of

residues was calculated and residues were classified as buried if RSA=0.0, exposed if RSA>0.8 and

middle for 0.0<RSA≤0.8. The linear relationship between site-specific conservation rank and site-

induced conservation gradient holds regardless of the residue's location within the protein or its

functional role (Fig 2). Overall, our results reveal the existence of a conservation–percolation

relationship in which higher residue conservation leads to stronger long-range percolation of

selective pressure to adjacent sites in the tertiary structure. Furthermore, this relationship holds

for residues in different types of functional sites.

Conservation gradient induced from a protein site is linearly correlated with the evolutionary

rate of the site

The linear trend between conservation and percolation shown above is based on conservation

ranks of residues, which are relative within a protein. In these calculations, we have lumped

together residues from different proteins having similar conservation rank, however these

residues, being from different proteins, can be under different selective pressure. To address this

caveat, we examined whether the conservation-percolation linear trend still holds when

conservation is measured in terms of absolute evolutionary rate (dN/dS). The evolutionary rate

is calculated for the yeast proteins in *S. cerevisiae* compared with its orthologs in four closely

related yeast species (see Methods). Conservation score annotations are transferred from

structural homologs onto yeast proteins using sequence alignment. Then, we binned all residues

into 100 equally spaced bins of conservation rank and calculated their average evolutionary rate

(dN/dS). The average evolutionary rate increases for residues with increasing conservation rank

(decreasing conservation) (S1 Fig), showing high correlation between the evolutionary rate of

yeast protein sites and the conservation scores of their structural homologs. The conservation-

percolation linear trend is shown to still hold here, where conservation is measured in terms of

absolute evolutionary rate (Fig 3). The lower the average evolutionary rate of a protein site is,

the stronger percolation of evolutionary conservation is induced from it.

Catalytic sites induce stronger conservation gradients than predicted by the conservation-

percolation trend

The annotation of functional and structural site residues was transferred to yeast proteins from

their structural models using sequence alignment. The average conservation gradient induced

from them is shown in Fig 3. As expected, functional residues evolve more slowly (low dN/dS)

than other residues. The average conservation gradients induced from ligand binding sites and

protein-protein interaction sites can all be predicted reasonably well from the linear

conservation-percolation trend. Protein-protein interaction sites induce slightly weaker

conservation gradient than expected, possibly due to their tendency to rewire during evolution.

Exposed non-functional residues induce slightly stronger conservation gradient than expected,

possibly due to hidden functional sites on protein surfaces. Remarkably, catalytic sites are the

only functional sites with the most significant deviation from the linear conservation-percolation

trend. The average conservation gradient induced from catalytic site residues is significantly

stronger than expected by the linear conservation-percolation trend. This result suggests that

the strong long-range conservation gradient induced from catalytic residues cannot be solely

attributed to the percolation of their low evolutionary rate.

Catalytic site residues induce stronger conservation gradients than non-catalytic functional site

residues with similar evolutionary rates

We have shown that on average, catalytic site residues induce stronger conservation gradients

than expected by their average evolutionary rate. Next, we further reinforce this result by

comparing the conservation gradient induced from subsets of catalytic and non-catalytic site

residues with similar evolutionary rates. We have sampled 1,000 random subsets of residues

from each type of functional site. For each such subset (represented by a circle in Fig 4), we

calculated the average evolutionary rate and average conservation gradient. As expected,

catalytic site residue subsets span the lowest dN/dS values (x-axis), followed by non-catalytic

ligand binding sites and allosteric sites, protein-protein interaction sites, ligand binding sites in

non-enzymes and finally buried residues. The linear trend between evolutionary rate and

conservation gradient holds for each of these functional site types (Fig 4). This result indicates

the robustness of the conservation-percolation linear trend regardless of the functional or

structural role of the residues. Interestingly, residue subsets from different functional sites with

similar evolutionary rates induce conservation gradients with different magnitudes. In particular,

catalytic sites induce significantly stronger long-range conservation gradients than all other non-

catalytic sites with similar evolutionary rates, including non-catalytic ligand binding sites (Fig 4).

Finally, protein-protein interaction site residues induce lower conservation gradients than most

other functional site residues with similar evolutionary rates, possibly due to the tendency for

protein-protein interactions to rewire during evolution.

Catalytic site residues often induce stronger conservation gradients than more conserved non-

catalytic functional site residues within the same protein

We have shown that conservation gradients induced by functional site residues in general

correlate linearly with the evolutionary rates of these functional site residues. We have also

shown that catalytic site residues are special in that they induce the strongest conservation

gradients among subsets of different functional site residues with similar average evolutionary

rates. However, our analyses have so far been carried out by grouping together functional site

residues from different proteins. In this section, we perform a stringent, per-protein analysis by

focusing on multi-functional proteins with at least two distinct functional sites and comparing

the evolutionary properties of different functional site residues within the same protein. Some

proteins in our dataset contain both catalytic sites as well as non-catalytic functional sites,

whereas other proteins in our dataset contain two distinct non-catalytic functional sites. We

found that for the majority of the functional site residue pairs within the same multi-functional

protein, the more conserved functional site residue indeed induces stronger conservation

gradient (86% if the more conserved residue is catalytic, Fig 5A; 67% if both residues are non-

catalytic, Fig 5B). These results agree with the hypothesis that induced conservation gradients

are largely driven by the percolation of selective pressure acting on functional sites.

Remarkably, in cases where the catalytic site residue is less conserved than the non-catalytic

functional site residue within the same multi-functional protein, the catalytic site residue still

induces stronger conservation gradient than the non-catalytic functional site for many of these

cases (discordance of 49%, Fig 5C). This large discordance shown in cases where the lower

conservation residue is a catalytic site residue, is significantly higher compared to the cases

where the lower conservation residue is a non-catalytic site residue (binomial test $P<<0.01$).

These results clearly show that within the same protein, less conserved catalytic site residues

often induce stronger conservation gradient than the more conserved non-catalytic site.

Therefore, the strong conservation gradients from catalytic sites cannot be entirely explained by

the percolation of the strong selective pressure acting on the catalytic sites.

Discussion

In this paper we have shown a linear relationship between two measures of evolutionary importance of protein sites under purifying selection. These are the degree of evolutionary conservation of the site itself, as well as the percolation of evolutionary conservation induced from the site via neighboring residues in the protein tertiary structure. Despite major differences between these two measures, we have shown that the linear relationship between the two measures holds for different types of proteins as well as for different types of functional sites in proteins. However, catalytic sites in enzymes are the principal exception to this rule. We have shown here that catalytic sites in enzymes induce significantly stronger conservation gradients in the rest of the protein than expected from the degree of conservation of the site alone. Catalytic sites have a unique and complex functionality as they both bind a substrate as well as reduce the free energy barrier required for a chemical reaction to occur. These catalytic sites were shown to be under stronger selective pressure compared with other functional sites such as protein-protein interaction sites and ligand binding sites(Jack et al. 2016; Sharir-Ivry and Xia 2019a). It was also shown that they induce a significantly stronger long-range evolutionary rate gradient than other functional sites. One hypothesis regarding the origin of the strong conservation gradient from catalytic sites is that it is simply due to the strong selective pressure acting on these sites percolating through the rest of the protein via residue-residue contacts. However, we have shown here that the strong selective pressure acting on catalytic sites cannot entirely explain the strong conservation gradients induced from catalytic sites.

The main determinant of the long-range conservation gradient from catalytic sites is still not completely understood. Local structural constraints (such as residue burial and packing) are

usually potential contributors as they are known to generally have a significant effect on residue

evolutionary rate(Franzosa and Xia 2009; Worth et al. 2009; Marcos and Echave 2015; Echave et

al. 2016). However, it was shown that local structural constraints are not the main determinants

of long-range conservation gradients in enzymes(Jack et al. 2016; Sharir-Ivry and Xia 2018; Sharir-

Ivry and Xia 2019b). Moreover, the fact that non-catalytic ligand binding sites and allosteric

binding sites induce significantly weaker evolutionary rate gradients implies that the ligand

binding and allosteric function are not the main determinants of the long-range conservation

gradients in enzymes either(Sharir-Ivry and Xia 2019a). We are therefore left with the hypothesis

that the uniquely strong conservation gradient in enzymes is imposed by the special requirement

for catalytic sites to differentially bind the transition state of a chemical reaction rather than the

reactants or products with very similar properties (Warshel et al. 2006), a function which is

unique to catalytic sites compared with other functional binding sites. A recent physical model of

residue evolutionary rates in enzymes introduced an activation term in addition to a stability

term and showed improved predictive ability(Echave 2019). The model attributed the activity to

the free energy required to transform a distorted catalytic site upon mutation to its native

conformation. The improved ability of the model supports the hypothesis that the main

determinant of the observed long-range conservation gradient in enzymes is a functional rather

than structural constraint. Overall, our results suggest that the stringent requirement for the

catalytic site to differentially bind to and stabilize the transition state of the catalyzed chemical

reaction imposes extensive evolutionary constraints on a large portion of the enzyme beyond

just the catalytic site, all of which play key roles in maintaining the catalytic function.

The results in this paper provide a complete, quantitative picture of evolutionary conservation

patterns within proteins induced by functional sites. The linear relationship between the degree

of conservation of a protein functional site and the induced conservation gradient in the rest of

the protein suggests that the strong selective pressure acting on the functional site percolates

through the rest of the protein via residue-residue contacts. Our results also clearly show the

unique evolutionary behavior of enzymes in which the catalytic site induces significantly stronger

evolutionary constraints on their surroundings than can be explained by the percolation theory

alone. Finally, our proposed measure of the evolutionary importance of protein sites can assist

in better phylogenetic inference, functional site prediction and protein design.

Methods

Protein dataset collection and functional site annotations

The current dataset is based on a dataset of structural homologs of yeast proteins(Sharir-Ivry and Xia 2019a). The dataset was created first by using gapped BLAST(Altschul et al. 1997) searches between protein subunit sequences with solved structure from the Protein Data Bank(Berman et al. 2000) and 5,861 translated open reading frames (ORFs) of the yeast *Saccharomyces cerevisiae*(Cherry et al. 1998). The ORF–subunit pairs were chosen such that both the subunit sequence and the ORF sequence had coverage of ≥50% in the alignment and E-value <10^{-5} and could be paired with their orthologs in four other closely-related yeast species *S. paradoxus*, *S. mikatae*, *S. bayanus and S. pombe*. This way, 1,555 yeast ORFs were mapped to homologs in the PDB. The procedure included the following steps:

First, if one of the homology-based structural models of a yeast ORF had an annotated allosteric site, this model was chosen. For 115 yeast proteins the structural model was identified with a known allosteric site as well as pre-calculated conservation scores in ConSurf-DB(Goldenberg et al. 2009). For all other yeast ORFs, if they had structural models with known ligand binding sites that do not overlap with catalytic sites and the closest catalytic site residue is at least 6Å away (non-catalytic ligand binding site), the model with the lowest E-value out of them was chosen. Overall, 54 non-enzymes with 57 ligand binding sites and 41 enzymes with 47 non-catalytic ligand-binding sites were part of the dataset. For all other yeast ORFs, the structural model with the lowest E-value was chosen. In this manner, 1064 more ORFs for which the best structural model had pre-calculated conservation scores in ConSurf-DB were added to the dataset. Overall, 1,274 ORF-subunit pairs were included in the study. Out of them, 148 yeast proteins were

identified with 280 protein-protein interactions sites and 348 proteins were identified with

catalytic sites. Full list of yeast proteins, their structural models along with identified functional

sites can be found in Supplementary Table S1.

Functional site annotations

Allosteric sites within the structural subunits were found and their residues annotated using the

Allosteric Database (ASD)(Huang et al. 2011; Shen et al. 2016). Biologically-relevant ligand-

binding sites were found using Binding MOAD (Mother of all Databases)(Hu et al. 2005; Ahmed

et al. 2015), a database of biologically significant protein-ligand binding in the PDB. Using MOAD,

sites with bound crystallographic additives, buffers, salts, metals and sites with covalently linked

ligands are excluded. Ligand-binding residues were identified using the BioLip(Yang et al. 2013)

database. Catalytic sites within the protein structural subunits were found using the Catalytic Site

Atlas(Furnham et al. 2014). In order to find proteins that participate in protein-protein

interactions in our dataset, we identified structural subunits where each subunit is both in

physical contact with another subunit and the corresponding modelled ORFs are reported as

interacting by at least one physical experiment in the BioGRID (Stark et al. 2006; Chatr-

Aryamontri et al. 2017). Our dataset contains 148 proteins with 280 protein-protein interaction

interfaces. Interfacial residues were identified as residues with different solvent accessibility

values when in complex compared to when the interacting partner is manually deleted from the

tertiary structure. Distances between residues were calculated as distances between their

respective Cα atoms. All functional site annotations of the chosen structural subunits were

transferred to the yeast ORF sequence according to the sequence alignment.

Evolutionary conservation and rate calculations

In this study we calculated both evolutionary conservation scores for the residues of the

structural subunits as well as average absolute evolutionary rates (dN/dS) for the yeast ORFs.

Evolutionary conservation scores were taken from ConSurf-DB(Goldenberg et al. 2009), which is

a database of pre-calculated conservation scores of residues in proteins with known structures

in the Protein Data Bank (PDB). ConSurf-DB conservation scores are based on collected sequence

homologs of the PDB structure and using the Rate4Site algorithm(Pupko et al. 2002).

To calculate the average evolutionary rates (dN/dS) for residues of *S. cerevisiae*, we first used the

orthology assignment of the protein-coding genes of *S. cerevisiae* with four other closely-related

yeast species (*S. paradoxus*, *S. mikatae*, *S. bayanus, and S. pombe*)*,* according to the Fungal

Orthogroup Repository(Wapinski et al. 2007). We then aligned the ORFs using MAFFT(Katoh and

Standley 2013). Then, evolutionary rates were calculated using the program codeml within the

PAML software package(Yang 2007). The tree was specified as ((((*S. cerevisiae*, *S. paradoxus*), *S.*

mikatae), *S. bayanus*), *S. pombe*). Codon frequencies were assumed equal (CodonFreq=0) and

other parameters in codeml were left to their default values. The codon alignments can be found

in Supplementary S2 File.

Statistical analysis

1000 random sample of 250 residues were collected from each type of functional site residues

for Fig 4.

Estimated standard errors in our measurements of conservation gradients (Pearson correlations)

and of dN/dS values were done using 50 rounds of bootstrap resampling.

Acknowledgments

This work was supported by Natural Sciences and Engineering Research Council of Canada (grant numbers RGPIN-2019-05952, RGPAS-2019-00012), Canada Foundation for Innovation (grant number JELF-33732), and Canada Research Chairs program.

Figure Legends

Figure 1. **Long-range conservation gradient induced from a protein residue is linearly correlated with its conservation rank within the protein**. (a) Average long-range conservation gradient from a residue as a function of conservation rank for all residues in the dataset binned into 20 equally spaced bins of conservation rank. (b) Distribution of per-protein Pearson correlation between residues' conservation ranks and conservation gradients.

Figure 2. Residues in functional and structural sites exhibit linear correlation between conservation and conservation gradient. Conservation gradient of residues as a function of their conservation rank within a protein along with the corresponding Pearson correlation for residues in different types of structural and functional sites.

Figure 3. **Conservation gradient induced from residues is linearly (negatively) correlated with their evolutionary rate (dN/dS), with catalytic site residues inducing stronger conservation gradient than expected by the linear trend**. Average conservation gradient (calculated as the average Pearson correlation between conservation of residues and their distance from a site) as a function of average evolutionary rate (dN/dS) for all yeast protein residues binned according to their annotated conservation rank into 100 equally spaced bins.

Figure 4. **Catalytic site residues induce stronger conservation gradients than non-catalytic functional site residues with similar evolutionary rates.** Average conservation gradients of subsets of functional residues. Each circle represents a subset of residues, colored by different types of functional sites.

Figure 5. Catalytic site residues often induce stronger conservation gradients than more conserved non-catalytic functional site residues within the same protein. Within the same protein, (A) more conserved catalytic site residues tend to induce stronger conservation gradient than less conserved non-catalytic site residues (binomial test, P<<0.001); (B) more conserved non-catalytic site residues tend to induce stronger conservation gradient than less conserved non-catalytic site residues (binomial test, P <<0.001); (C) less conserved catalytic site residues often induce stronger conservation gradient than more conserved non-catalytic site residues (discordance for half of the cases, binomial test, P=0.13). Functional site residue pairs for which the ordering of residue conservation agrees with the ordering of induced conservation gradient (concordance) are marked in blue. Functional site residue pairs for which the ordering of residue conservation disagrees with the ordering of induced conservation gradient (discordance) are marked in orange.

References

Ahmed A, Smith RD, Clark JJ, Dunbar JB, Carlson HA. 2015. Recent improvements to Binding MOAD: a resource for protein–ligand binding affinities and structures. *Nucleic Acids Res.* 43:D465–D469.

Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* 25:3389–3402.

Bartlett GJ, Porter CT, Borkakoti N, Thornton JM. 2002. Analysis of catalytic residues in enzyme active sites. *J. Mol. Biol.* 324:105–121.

Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat TN, Weissig H, Shindyalov IN, Bourne PE. 2000. The Protein Data Bank. *Nucleic Acids Res.* 28:235–242.

Celniker G, Nimrod G, Ashkenazy H, Glaser F, Martz E, Mayrose I, Pupko T, Ben-Tal N. 2013. ConSurf: Using Evolutionary Data to Raise Testable Hypotheses about Protein Function. *Isr. J. Chem.* 53:199–206.

Chatr-Aryamontri A, Oughtred R, Boucher L, Rust J, Chang C, Kolas NK, O'Donnell L, Oster S, Theesfeld C, Sellam A, et al. 2017. The BioGRID interaction database: 2017 update. *Nucleic Acids Res.* 45:D369–D379.

Cherry JM, Adler C, Ball C, Chervitz SA, Dwight SS, Hester ET, Jia Y, Juvik G, Roe T, Schroeder M, et al. 1998. SGD: Saccharomyces Genome Database. *Nucleic Acids Res.* 26:73–79.

Conant GC, Stadler PF. 2009. Solvent exposure imparts similar selective pressures across a range of yeast proteins. *Mol. Biol. Evol.* 26:1155–1161.

Echave J. 2019. Beyond Stability Constraints: A Biophysical Model of Enzyme Evolution with Selection on Stability and Activity. *Mol. Biol. Evol.* 36:613–620.

Echave J, Spielman SJ, Wilke CO. 2016. Causes of evolutionary rate variation among protein sites. *Nat. Rev. Genet.* 17:109–121.

Franzosa EA, Xia Y. 2009. Structural determinants of protein evolution are context-sensitive at the residue level. *Mol. Biol. Evol.* 26:2387–2395.

Franzosa EA, Xue R, Xia Y. 2013. Quantitative residue-level structure-evolution relationships in the yeast membrane proteome. *Genome Biol. Evol.* 5:734–744.

Furnham N, Holliday GL, De Beer TAP, Jacobsen JOB, Pearson WR, Thornton JM. 2014. The Catalytic Site Atlas 2.0: Cataloging catalytic sites and residues identified in enzymes. *Nucleic Acids Res.* 42:D485–D489.

Goldenberg O, Erez E, Nimrod G, Ben-Tal N. 2009. The ConSurf-DB: pre-calculated evolutionary conservation profiles of protein structures. *Nucleic Acids Res.* 37:D323-7.

Hamelryck T. 2005. An amino acid has two sides: A new 2D measure provides a different view
of solvent exposure. *Proteins* 59:38–48.

Hu L, Benson ML, Smith RD, Lerner MG, Carlson HA. 2005. Binding MOAD (Mother Of All
Databases). *Proteins Struct. Funct. Bioinforma.* 60:333–340.

Huang Z, Zhu L, Cao Y, Wu G, Liu X, Chen Y, Wang Q, Shi T, Zhao Y, Wang Y, et al. 2011. ASD: a
comprehensive database of allosteric proteins and modulators. *Nucleic Acids Res.*
39:D663–D669.

Jack BR, Meyer AG, Echave J, Wilke CO. 2016. Functional sites induce long-range evolutionary
constraints in enzymes. *PLOS Biol.* 14:e1002452.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
Improvements in performance and usability. *Mol. Biol. Evol.* 30:772–780.

Marcos ML, Echave J. 2015. Too packed to change: side-chain packing and site-specific
substitution rates in protein evolution. *PeerJ* 3:e911.

Nelson ED, Grishin N V. 2016. Evolution of off-lattice model proteins under ligand binding
constraints. *Phys. Rev. E* 94:022410.

Overington J, Donnelly D, Johnson MS, Sali A, Blundell TL. 1992. Environment-specific amino
acid substitution tables: tertiary templates and prediction of protein folds. *Protein Sci.*
1:216–226.

Pupko T, Bell RE, Mayrose I, Glaser F, Ben-Tal N. 2002. Rate4Site: an algorithmic tool for the
identification of functional regions in proteins by surface mapping of evolutionary
determinants within their homologues. *Bioinformatics* 18:S71–S77.

Sharir-Ivry A, Xia Y. 2017. The impact of native state switching on protein sequence evolution.
Mol. Biol. Evol. 34:1378–1390.

Sharir-Ivry A, Xia Y. 2018. Nature of Long-Range Evolutionary Constraint in Enzymes: Insights
from Comparison to Pseudoenzymes with Similar Structures. *Mol. Biol. Evol.* 35:2597–
2606.

Sharir-Ivry A, Xia Y. 2019a. Non-catalytic Binding Sites Induce Weaker Long-Range Evolutionary
Rate Gradients than Catalytic Sites in Enzymes. *J. Mol. Biol.* 431:3860–3870.

Sharir-Ivry A, Xia Y. 2019b. Using Pseudoenzymes to Probe Evolutionary Design Principles of
Enzymes. *Evol. Bioinforma.* 15:117693431985593.

Shen Q, Wang G, Li S, Liu X, Lu S, Chen Z, Song K, Yan J, Geng L, Huang Z, et al. 2016. ASD v3.0:
unraveling allosteric regulation with structural mechanisms and biological networks.
Nucleic Acids Res. 44:D527–D535.

Stark C, Breitkreutz B-J, Reguly T, Boucher L, Breitkreutz A, Tyers M. 2006. BioGRID: a general
repository for interaction datasets. *Nucleic Acids Res.* 34:D535-9.

Tóth-Petróczy A, Tawfik DS. 2011. Slow protein evolutionary rates are dictated by surface-core

association. *Proc. Natl. Acad. Sci. U. S. A.* 108:11151–11156.

Wapinski I, Pfeffer A, Friedman N, Regev A. 2007. Natural history and evolutionary principles of
gene duplication in fungi. *Nature* 449:54–61.

Warshel A, Sharma PK, Kato M, Xiang Y, Liu H, Olsson MHM. 2006. Electrostatic basis for
enzyme catalysis. *Chem. Rev.* 106:3210–3235.

Worth CL, Gong S, Blundell TL. 2009. Structural and functional constraints in the evolution of
protein families. *Nat. Rev. Mol. Cell Biol.* 10:709–720.

Yang J, Roy A, Zhang Y. 2013. BioLiP: A semi-manually curated database for biologically relevant
ligand-protein interactions. *Nucleic Acids Res.* 41:D1096-1103.

Yang Z. 1996. Among-site rate variation and its impact on phylogenetic analyses. *Trends Ecol.*
Evol. 11:367–372.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* 24:1586–
1591.

Yeh SW, Huang TT, Liu JW, Yu SH, Shih CH, Hwang JK, Echave J. 2014. Local packing density is
the main structural determinant of the rate of protein sequence evolution at site level.
Biomed Res. Int.:572409.

Figures

Figure 1. **Long-range conservation gradient induced from a protein residue is linearly correlated with its conservation rank within the protein**. (a) Average long-range conservation gradient from a residue as a function of conservation rank for all residues in the dataset binned into 20 equally spaced bins of conservation rank. (b) Distribution of per-protein Pearson correlation between residues' conservation ranks and conservation gradients.

Fig 2. Residues in functional and structural sites exhibit linear correlation between conservation and conservation gradient. Conservation gradient of residues as a function of their conservation rank within a protein along with the corresponding Pearson correlation for residues in different types of structural and functional sites.

Fig 3. **Conservation gradient induced from residues is linearly (negatively) correlated with their evolutionary rate (dN/dS), with catalytic site residues inducing stronger conservation gradient than expected by the linear trend**. Average conservation gradient (calculated as the average Pearson correlation between conservation of residues and their distance from a site) as a function of average evolutionary rate (dN/dS) for all yeast protein residues binned according to their annotated conservation rank into 100 equally spaced bins.

520 **Figure 4**. **Catalytic site residues induce stronger conservation gradients than non-catalytic functional**
521 **site residues with similar evolutionary rates.** Average conservation gradients of subsets of functional
522 residues. Each circle represents a subset of residues, colored by different types of functional sites.

Fig 5. Catalytic site residues often induce stronger conservation gradients than more conserved non-catalytic functional site residues within the same protein. Within the same protein, (A) more conserved catalytic site residues tend to induce stronger conservation gradient than less conserved non-catalytic site residues (binomial test, P<<0.001); (B) more conserved non-catalytic site residues tend to induce stronger conservation gradient than less conserved non-catalytic site residues (binomial test, P <<0.001); (C) less conserved catalytic site residues often induce stronger conservation gradient than more conserved non-catalytic site residues (discordance for half of the cases, binomial test, P=0.13). Functional site residue pairs for which the ordering of residue conservation agrees with the ordering of induced conservation gradient (concordance) are marked in blue. Functional site residue pairs for which the ordering of residue conservation disagrees with the ordering of induced conservation gradient (discordance) are marked in orange.

Pearson correlation between conservation rank and
conservation gradient in a protein

(A) Catalytic site / Non-catalytic site

(B) Non-catalytic site / Non-catalytic site

(C) Non-catalytic site / Catalytic site

Conservation

■ Discordance - Lower conservation induces stronger conservation gradient
■ Concordance – Lower conservation residue induces weaker conservation gradient