Confirmation of interpersonal expectations is intrinsically rewarding

Niv Reggev* [1,2,3], Anoushka Chowdhary[1], and Jason P. Mitchell[1]

[1] Department of Psychology, Harvard University, 33 Kirkland St, Cambridge, MA, US, 0218

[2] Department of Psychology and [3] Zlotowski Center for Neuroscience, Ben-Gurion University of the Negev, P.O. Box 653, Be'er-Sheva, Israel, 84105

ORCiD identifiers:

Niv Reggev: 0000-0002-5734-7457

Corresponding author:

Niv Reggev, Department of Psychology, Ben Gurion University of the Negev, P.O. Box 653,

Be'er-Sheva, Israel, 84105

Phone number: +972-8-67472030

Email address: reggevn@bgu.ac.il

This article is a preprint, not yet published in a peer-reviewed journal. We welcome any comments or feedback - please send these to the corresponding author

1   **Abstract**

People want to interact successfully with other individuals, and they invest significant efforts

in attempting to do so. Decades of research have demonstrated that to simplify the dauntingly

complex task of interpersonal communication, perceivers use stereotypes and other sources of

prior knowledge to predict the responses of individuals in their environment. Here, we show that

these top-down expectations can also shape the subjective value of expectation-consistent and

expectation-violating targets. Specifically, in two neuroimaging experiments ($n = 58$), we

observed increased activation in brain regions associated with reward processing—including the

nucleus accumbens—when perceivers observed information consistent with their social

expectations. In two additional behavioral experiments ($n = 704$), we observed that perceivers

were willing to forgo money to encounter an expectation-consistent target and avoid an

expectation-violating target. Together, these findings suggest that perceivers value having their

social expectations confirmed, much like food or monetary rewards.

Keywords: Stereotypes, Expectations, Reward, fMRI, NAcc, Value, Motivation

People dedicate a substantial portion of their time to interacting with other individuals. When we succeed in doing so, we feel better physically and psychologically [1–4]. At the same time, other people also present one of the most complicated challenges we have to face. Understanding another person involves inferring hidden states based on fragmentary sensory, verbal and visceral cues, each of which conveys only a small amount of information. To simplify the highly demanding challenge of social cognition, perceivers use top-down predictions (e.g., stereotypes) that help make sense of others in a rapid fashion [5–8]. Perceivers can thus seamlessly interact with their environment while refraining from the effortful construction of elaborative representations for each individual they encounter [9,10]. For example, on her first day of school, a freshman might assume that her female peers may be interested in conversing about shopping. Utilizing these predictions, the freshman could effortlessly engage in spontaneous conversations with her new female peers, potentially facilitating her social bonds.

However, although they often facilitate interpersonal interaction, social predictions also impose a cost on perceivers. Individuals tend to adhere to predictions they have previously formed and fail to modify them even in the face of contradictory evidence [11–15]. For instance, when perceivers first learn that someone is 'intelligent' and then subsequently learn that he is also 'envious', they form a favorably-skewed impression of that person; on the other hand, when perceivers learn about these same two character traits in reverse order, they form an unfavorable impression of the target [16,17]. To date, research has not been able to identify the sources that support the persistence of these initial predictions about other individuals. Here, we integrate insights from social psychology and neuroscience to explore the idea that perceivers prefer to 'stick' with their initial predictions because they attribute subjective value to the confirmation of these predictions. That is, we posit that targets who confirm our expectations about them (such as

stereotype-consistent targets) will trigger a reward-like response similar to food, sex, or chocolate.

Several lines of research already hint at such an effect. Perceivers generally like individuals who conform to expectations more than individuals who violate them [18–20]; for example, observers typically prefer female teachers to male teachers, but like male leaders better than their equally competent female peers [21,22]. Similarly, participants express greater trust in targets that fit gender-based predictions [23,24]. Moreover, perceivers demonstrate similar effects for emotion-based expectations, regardless of the valence of the emotion [25]. Several theorists have suggested that perceivers may gradually develop a habitual hedonic response for targets conforming to normative expectations [26–29]. These suggestions dovetail with the well-documented aversive reactions people experience when confronted with violations of predictions and the uncertainty associated with such violations [15,29–32]. Put together, these positive and aversive responses motivate perceivers to seek expectation-consistent information.

In spite of ample evidence for perceivers' motivation to confirm their social expectations, scholars are still debating the mechanisms supporting the persistence of this motivation. Here we suggest that neural activity can offer a novel insight on this topic. In recent years, scholars have identified the involuntary effects of motivation and expectation in several neural systems, most notably in the mesolimbic dopaminergic system [33]. Animal models suggest that midbrain dopaminergic activity signals one's internal desire to obtain a goal [34]. In humans, dopamine modulates the motivation to experience positive effects by adjusting midbrain and striatal responses to better- or worse-than-expected information [35–37]. Likewise, participants expecting a painful stimulus demonstrate increased striatal activity while experiencing pain, compared with participants who do not expect to feel pain [38,39]. Similarly, stigmatized participants demonstrate

different striatal activity compared to non-stigmatized participants [40]. Finally, a recent meta-analysis reported that when perceivers agreed with expected group opinions, they demonstrated robust striatal activity, as compared with times in which they deviated from the group consensus [41]. These studies suggest that the striatum responds to events that align with perceivers' motivation.

Notably, the involvement of the striatum hints at a potential mechanism mediating the effects of expectation and motivation. Researchers repeatedly identify striatal activity, and most prominently activity in its ventral portion, in anticipation and receipt of various types of reward [42–44]. For example, the nucleus accumbens (NAcc), located at the ventral-rostral tip of the striatum, responds both to primary rewards (e.g., food or erotic) and secondary rewards (e.g., money or positive feedback) [45–47]. The NAcc also responds to social experiences, such as engagement with attractive or smiling faces, prosocial actions, or placing one's trust in peers [48–51].

Together, these studies suggest that consistency with stereotypes and other forms of interpersonal predictions is intrinsically rewarding. To test this hypothesis, we first measured NAcc activity in response to information consistent or inconsistent with social expectations. Using functional magnetic resonance imaging (fMRI), we scanned participants while they judged target individuals on characteristics that either conformed to or violated interpersonal expectations. As the NAcc is consistently involved in rewarding experiences, activity in this region can serve as a marker of a neural reward response. If perceivers value having their social expectations confirmed, we should observe increased NAcc activity for trials in which targets were associated with characteristics that confirm expectations compared to trials in which targets violate them.

In addition, we assessed whether perceivers actively prefer expectancy-confirming social

information by creating experimental situations in which participants could trade money for the

chance to view targets with expectation-consistent characteristics. To do so, we relied on a

modified version of a "pay-per-view" task, previously used with human and non-human

primates, to measure the monetary value associated with expectancy-confirming and expectancy-

violating stimuli [52,53]. If perceivers experience expectancy-confirming information as

intrinsically more valuable, we expected participants to forgo money to interact with individuals

associated with expectancy-consistent rather than expectancy-violating information.

Interestingly, social expectations can take multiple forms. For example, stereotypes relate

specific attributes to social groups regardless of personal knowledge about group members.

Conversely, we can construct detailed individuated expectations about familiar individuals, such

as personally familiar others or famous people [10,14]. Accordingly, we also probed whether

predictions from these two distinct sources evoked qualitatively different reward responses.

Together, the results of four studies support the hypothesis that perceivers are motivated to

reaffirm their interpersonal forecasts, in part because they experience the confirmation of social

expectations as a powerful form of subjective reward.

17   **Results**

Study 1: Neural response to stereotype confirmation vs. violation

In Study 1 we observed greater NAcc activity when participants saw targets associated with

gender-stereotype-consistent information than when they saw targets associated with stereotype-

violating information. We scanned participants ($n = 28$) while they formed impressions about

targets that varied in the degree to which they confirmed gender stereotypes. Gender stereotypes

included various characteristics typically associated with men (e.g., "emotionally closed" or

"CEO of a big company") or women (e.g., "loves children" or "an admired preschool teacher").

In each of 204 trials, participants first read a short description for 1.5 seconds and then saw the

face of a target man or woman (see Fig S1A). Participants rated how likely the target was to have

the presented characteristic (see SI results and Table S1 for behavioral results). We conducted

three parallel analyses to examine whether the neural region most associated with reward—

namely, the NAcc—was more engaged when the target was associated with stereotype-derived

expectations compared to when the target violated them. First, a whole-brain random-effects

contrast identified regions that were more active for *stereotype-consistent > stereotype-violating*

trials ($p < 0.05$, corrected; see Table S2 for full results). This analysis indicated a significantly

greater response in the NAcc when the presented target matched the stereotypical expectation set

by the preceding statement than when the target violated that expectation (Fig. 1A). Parallel

results were obtained when we modeled expectation consistency as a continuous rather than

dichotomized predictor (see Table S2 and SI Materials and Methods for full details).

Second, to confirm that this region overlapped with those responsive to rewards, we

independently defined a neural regions of interest (ROI) based on spheres around peak voxels

identified in a comprehensive meta-analysis [47]. In this independently defined region consistency

with stereotypes resulted in significantly greater activity compared to their violation [one-sided

test: $t_{(54)} = 3.53$, $p = 0.0004$, *Hedges's g* = 0.37 [95% confidence intervals: 0.1–0.64], Fig. 1B].

Third, we corroborated this finding by defining ROIs from a task in which participants received

monetary rewards based on their performance [the Monetary Incentive Delay (MID) task; see

Materials and Methods] [54]. This analysis yielded similar results [$t_{(54)} = 2.87$, $p = 0.0029$, $g = 0.30$

[0.03–0.59]]. Together, these patterns suggest that seeing a person associated with a stereotypical

characteristic triggers activation in the very same region that responds to primary and secondary

reinforcers, highlighting the intrinsic value of stereotype confirmation.

3
Figure 1. Neural responses associated with the confirmation of expectations about other
individuals. (a) Whole-brain random-effects contrasts comparing expectation-consistent >
expectation-violating *trials revealed activity in the nucleus accumbens (NAcc) in (A) Study 1 and*
(B) Study 3. (C) We independently defined a region of interest in the NAcc using a
comprehensive meta-analysis (MNI coordinates: -6, 10, -6; 10, 12, -6). Analysis of parameter
estimates in this region confirmed that the bilateral NAcc showed a stronger response during
consistent than during violating trials in Study 1 and (D) Study 3. Upper panel: Across all
figures, individual dots represent parameter estimates for individual participants. Each figure
also visualizes the mean of each condition (as a red dot), the median (solid horizontal line), and
first and third quartiles (boxplot). Lower panel: effect size (the mean difference between
respective conditions, indicated by the black circles), the bootstrapped 95% confidence intervals
(illustrated by the vertical lines), and the resampled distribution of the effect size given the
observed data, indicated by the curve (see Methods section).

Additionally, to investigate whether this NAcc activation is limited to stereotype-derived

expectations, we included a third type of statements in our study: stereotype-neutral statements

(e.g., "drinks coffee every morning"). Interestingly, we found that the overall NAcc response to

targets associated with stereotype-neutral information was higher than the response to

stereotype-violating information and not different from stereotype-consistent information [Fig.

1C; Bonferroni corrected comparisons: neutral versus violating: Meta-analysis ROIs: $t_{(54)}$ = 2.83,

p = 0.0195, g = 0.29 [0.09–0.51]; MID ROIs: $t_{(54)}$ = 2.38, p = 0.062, g = 0.25 [0.04–0.47]; neutral

versus consistent: Meta-analysis ROIs: $t_{(54)}$ = 0.7, p = 0.76, g = 0.07 [-0.19–0.34]; MID ROIs:

$t_{(54)}$ = 0.49, p = 0.63, g = 0.05 [-0.18–0.28]]. At first blush, this finding suggests that NAcc

activation was modulated only by stereotype-inconsistent information, not that it was especially

driven by stereotype-consistent information which, on average, did not differ from stereotype-

neutral trials. However, additional analyses belie this interpretation. On each trial, participants

rated the likelihood that the target could be described by the accompanying characteristic (e.g.,

"enjoys drinking coffee in the morning"). Importantly, the NAcc was activated during

stereotype-neutral trials only when participants endorsed those characteristics as descriptive of

the target and not when they rejected its applicability to a target (Meta-analysis ROIs: $F_{(1,76.18)}$ =

15.67, p = 0.0002; η_p^2 = 0.17 [0.06, 0.29]; MID ROIs: $F_{(1,76.5)}$ = 12.66, p = 0.0006; η_p^2 = 0.14

[0.04, 0.26]). In contrast, the preferential NAcc activation observed for stereotype-consistent

trials was unaffected by participants' likelihood ratings (interaction between response and

condition: Meta-analysis ROIs: $F_{(1.77,47.68)}$ = 5.4, p = 0.01, η_p^2 = 0.17 [0.02, 0.3]; MID ROIs:

$F_{(1.87,50.4)}$ = 4.01, p = 0.03, η_p^2 = 0.13 [0.01, 0.26]; see Fig. 2. Stereotype-consistent trials: meta-

analysis ROIs: $F_{(1,76.18)}$ = 0.2, p > 0.5; η_p^2 = 0.003 [0, 0.05]; MID ROIs: $F_{(1,76.5)}$ = 0.63, p > 0.5;

η_p^2 = 0.005 [0, 0.06]). Together, these data points suggest that NAcc is activated specifically

when one's expectations are confirmed, regardless of whether those expectations derive from

culturally-salient stereotypes or from more personal and idiosyncratic sources. Finally, as for

stereotype-neutral trials, stereotype-violating trials (e.g., a man who "enjoys shopping for

shoes") were associated with greater NAcc activation when participants endorsed the likelihood

of such characteristics (Meta-analysis ROIs: $F_{(1,76.18)} = 17.84$, p = 0.0001; $\eta_p^2 = 0.19$ [0.07, 0.31];

MID ROIs: $F_{(1,76.5)} = 15.76$, p = 0.0002; $\eta_p^2 = 0.17$ [0.06, 0.29]).

a

x = -6   y = 10

b

Condition: Consistent

Parameter estimates

$\widehat{\mu}=0.35$
$\widehat{\mu}=0.25$

Likely   Unlikely

Condition: Neutral

Parameter estimates

$\widehat{\mu}=0.58$
$\widehat{\mu}=-0.3$

***

Likely   Unlikely

Condition: Violate

Parameter estimates

$\widehat{\mu}=0.46$
$\widehat{\mu}=-0.48$

***

Likely   Unlikely

Mean diff. and 95% CI

Likely
minus Unlikely

Mean diff. and 95% CI

Likely
minus Unlikely

Mean diff. and 95% CI

Likely
minus Unlikely

Figure 2. The effect of behavioral ratings of trials on neural responses. In each trial,
participants indicated whether they thought the presented target was likely or unlikely to be
associated with the presented statement. (A) The independently defined regions of interest in the
NAcc. (B) We observed a significant interaction in the independently defined NAcc in Study 1:
Participants' ratings modulated the neural response only for stereotype-neutral and stereotype-
violating targets, suggesting that stereotype-confirming targets are involuntarily rewarding.

Study 2: The monetary value of stereotype confirmation

Although activation of the NAcc often reflects the presence of rewarding stimuli [55], this

region can also respond to non-value related processes including information coding or salience

effects [56]. To complement our initial neural results, in the preregistered Study 2 we examined a

behavioral measure of the value associated with rating expectancy-confirming targets.

Specifically, we tested how much money participants were willing to forgo to view stereotype-

consistent instead of stereotype-violating targets.

Participants on Amazon Mechanical Turk (*n*s = 174 and 169 in Study 2a and 2b, respectively)

made a series of choices to rate one of two target types: a stereotype-consistent target (e.g., a

man who enjoys riding motorcycles) or a stereotype-violating target (e.g., a man who enjoys

shopping for shoes), designated as "typical" and "atypical" respectively. After each choice,

participants saw a target accompanied by a statement and rated the likelihood that the target

would be associated with the statement on a 0–100 scale. Participants had up to 5 seconds for

each phase of the task (see Fig S2). To avoid potentially different responses to male and female

targets, Study 2a included only male faces and Study 2b included only female faces. On each of

the 25 trials, small monetary payoffs ($0.03–$0.09 in increments of 2 cents) were associated with

each target choice. Participants received a subset of these payoffs as a monetary bonus for the

task. Payoff amounts for each target choice varied across trials (and were occasionally equal), as

did the location of the option for which participants received the larger amount. If consistency

with stereotypes is intrinsically rewarding, participants should be willing to forgo money—to

choose the lower-paying option—to see stereotype-consistent individuals. On the other hand, a

participant seeking to maximize monetary payoff should consistently choose the higher paying

option regardless of the stereotypicality of the information that follows.

We modeled the relative value of each target type by calculating the point of subjective

equivalence (PSE) between the two options. This value was derived by fitting a cumulative

normal distribution curve to participants' choices (Fig. 3A) and finding the monetary value at

which participants effectively chose arbitrarily between the two target types [57]. Thus, the PSE

represents the relative monetary value of one target type over another.

As predicted, participants demonstrated a significant preference for seeing stereotype-

consistent targets over stereotype-violating targets. When the two trial types shared the same

payoff amounts, participants chose the stereotypical targets 58% and 57% of the time for male

and female targets, respectively (significantly more than chance, as indicated in a generalized

mixed model analysis by an odds ratio of 1.4, $Z = 3.11$, $p = 0.0019$ and an odds ratio of 1.41,

$Z = 2.91$, $p = 0.00369$ for the zero-centered intercept in Study 2a and 2b, respectively).

Moreover, the calculated PSE indicated that participants forewent an average of 0.34 and 0.335

cents per trial to rate a stereotype-consistent over a stereotype-violating target in Study 2a and

2b, respectively (95% Confidence Interval [CI]: 0.21–0.47, $t_{(173)} = 5.17$, $p < 0.0001$, *Cohen's*

d $= 0.39$ [0.24–0.55] and CI: 0.25–0.43, $t_{(168)} = 7.3$, $p < 0.0001$, $d = 0.56$ [0.4–0.72] for the two

studies, respectively; Fig 3b). This PSE resulted in an average loss of 10% of potential earnings,

as participants chose lower monetary amounts to view stereotype-consistent targets. Just as non-

human primates prefer to view dominant groupmates over receiving juice [52] and students are

willing to forgo money to talk about themselves [53] or to view attractive members of the opposite

sex [58], our participants gave up money to view information that was in line with their

stereotypical expectations.

Figure 3. The monetary values of consistency with stereotype and person-specific expectations.
(A) Visualization of the cumulative distribution function we used to calculate the Point of
Subjective Equivalence (PSE), illustrated by group data from Study 2a. The x-axis represents the
difference between the monetary values associated with the two target types presented in each
trial. Each dot indicates the proportion of trials in which participants chose to rate a stereotype-
confirming over a stereotype-violating target. The PSE was calculated as the point at which a
cumulative normal distribution function, fit to these responses, passes 50%. This point represents
the relative monetary value associated with one target type over the other. Negative values
indicate that participants preferred to incur a relative monetary loss to rate a confirming target.
Error bars depict 95% confidence intervals. (B) Distribution of individual PSE values for Study
2. Rating stereotype-consistent targets in Studies 2a and 2b was associated with significantly
higher subjective value than rating stereotype-violating targets. Each gray dot depicts PSE for a
specific participant. Red dots indicate the sample mean. Error bars depict 95% confidence
intervals. (C) Distribution of individual PSE values for Study 4. Rating Obama-consistent trials
was associated with significantly higher subjective value than rating Obama-violating trials.
However, rating Trump-consistent trials did not significantly differ from rating Trump-violating
trials.

Study 3: Neural response to interpersonal expectations

Together, Studies 1 and 2 suggest that perceivers experience consistency with social

expectations as intrinsically rewarding. However, we designed these studies primarily to test the

effects of a specific type of social expectations – gender-based stereotypes. Although stereotypes

are a significant source of interpersonal expectations, perceivers routinely make use of

additional, idiosyncratic sources of information, especially for individuals with whom they are

highly familiar. Does the reward value of expectancy-consistent information extend to person-

specific predictions?

To examine this question, Study 3 assessed the responses of the neural reward system to the

consistency with and violation of expectations regarding two highly familiar targets, the current

and previous presidents of the United States at the time of the study: Donald Trump and Barack

Obama, respectively. To facilitate comparison to Study 1, the preregistered Study 3 also included

stereotype-derived expectations about unfamiliar targets. As in Study 1, participants ($n = 30$)

rated how likely each of 240 specific statements described a specific man or woman or described

Donald Trump or Barack Obama. We presented stereotype-related and person-specific

statements in blocks of 15 trials per content domain for a total of 60 trials per condition

(consistent/violating targets in the stereotype/person-specific domain). Person-specific

statements included various characteristics typically associated with one – but not the other –

leader (e.g., "Supports a wall along the borders" versus "Acts to support women's rights").

Participants first read a short statement for 1.5 seconds and then saw the face of a target (a man,

a woman, Trump or Obama; see Fig S1C). Participants rated how likely the statement was to

describe the target (see Table S3 and SI Results for full details) .

We conducted two parallel analyses to examine whether the NAcc was more engaged when

presented with information that was consistent expectations than with information that violated

them. First, a whole-brain random-effects contrast identified regions that were more active for

expectation-consistent > expectation-violating trials (p < 0.05, corrected; see Table S3 for full

results). This analysis indicated significantly greater response in several regions, including the

NAcc, when the target was consistent with the expectation set by the preceding statement than

when the target violated that expectation (Fig. 1B).

In a second analysis we defined the NAcc via two independent procedures, first as bilateral

spheres around peak voxels identified in a meta-analysis, and then by examining participants'

neural responses during the MID task. Across both procedures the NAcc demonstrated a robust

difference in activity between expectation-consistent and expectation-violating targets (Fig. 1D).

However, we did not observe any difference between stereotypes and person-specific trials in

any of the analyses. Specifically, a 2 X 2 repeated measures ANOVA over activity in bilateral

NAcc revealed a main effect of expectation-consistency, with higher activation associated with

expectation-consistent compared to expectation-violation (meta-analysis ROIs: $F_{(1,29)} = 21.89$,

$p < 0.0001$, $\eta^2_p = 0.43$ [0.19–0.58]; MID ROIs: $F_{(1,29)} = 11.2$, $p = 0.002$, $\eta^2_p = 0.28$ [0.07–0.46]).

Activation did not significantly differ between stereotype and person-specific content (meta-

analysis ROIs: $F_{(1,29)} = 3.64$, $p = 0.07$, $\eta^2_p = 0.11$ [0–0.29]; MID ROIs: $F_{(1,29)} = 1.41$, $p = 0.24$,

$\eta^2_p = 0.05$ [0–0.2]) and no interaction was observed between the factors (meta-analysis ROIs:

$F_{(1,29)} = 0.31$, $p = 0.58$, $\eta^2_p = 0.01$ [0–0.13]; MID ROIs: $F_{(1,29)} < 0.01$, $p = 0.99$, $\eta^2_p < 0.0001$).

Accordingly, expectation-consistency yielded more neural activity for each of the two content

domains when examined separately (meta-analysis ROIs: $F_{(1,54.8)} = 5.96$, $p = 0.0179$,

$\eta^2_p = 0.1$ [0.01–0.23] and $F_{(1,54.8)} = 11.03$, $p = 0.0016$, $\eta^2_p = 0.17$ [0.04–0.31] for gender

stereotypes and person-specific expectations, respectively; MID ROIs: $F_{(1,55.47)} = 4.36$, $p = 0.04$,

$\eta^2_p = 0.07$ [0.001–0.2] and $F_{(1,55.47)} = 4.45$, $p = 0.04$, $\eta^2_p = 0.07$ [0.002–0.2], respectively). Thus,

consistency with social expectations triggered more activation in the NAcc than violation of such

expectations, regardless of whether source of the expectation was general social knowledge

(stereotypes) or person-specific knowledge.

Finally, similar to Study 1, variability in behavioral responses in Study 3 enabled us to test the

effects of subjective endorsement of characteristics for stereotype-based characteristics (but not

for person-specific-expectations; see SI results). As in Study 1, NAcc activity was not modulated

by subjective endorsement of characteristics for stereotype-consistent targets (simple effects

analyses: meta-analysis ROIs: $F_{(1,58)} = 3.99$, $p = 0.0503$; $\eta_p^2 = 0.06$ [0, 0.18]; MID ROIs:

$F_{(1,56.5)} = 3.20$, $p = 0.08$; $\eta_p^2 = 0.05$ [0, 0.17]). However, we did not observe an interaction

between response and condition (meta-analysis ROIs: $F_{(1,29)} = 0.08$, $p > 0.5$,

$\eta_p^2 = 0.003$ [0, 0.09]; MID ROIs: $F_{(1,29)} = 0.40$, $p > 0.5$, $\eta_p^2 = 0.01$ [0, 0.14]). In other words,

unlike in Study 1, NAcc activation was *in*sensitive to the degree to which a participant judges

information to apply to a target, regardless of the stereotypicality of the target. Together, these

results suggest that consistency with expectations about other targets is valuable for the two most

dominant sources of interpersonal expectations – group-based stereotypes and person-specific

knowledge.

Study 4: The monetary value of person-specific expectation-confirmation

In Study 2, we observed that perceivers are willing to forgo money to view stereotype-

consistent (rather than stereotype-violating) information. To examine whether this behavioral

effect extends to expectations about specific individuals, Study 4 replicated the procedure from

Study 2 using familiar individuals (Obama and Trump). On each of 32 trials, participants on

Prolific Academic (n = 189 and 172 in Study 4a and 4b, respectively) first chose between seeing

either an expectation-consistent or an expectation-violating target and then rated the target. Study

4a included only Obama as the target of statements and Study 4b included only Trump. After

choosing the type of content they would like to see, participants rated the likelihood that a

specific statement would be associated with the target on a 0–100 scale. Payoff amounts for each

choice varied across trials (and were occasionally equal), as did the option for which participants

received the larger amount. As in Study 2, we quantified the subjective monetary value of each

option by calculating the PSE between the two display types by fitting a cumulative normal

distribution curve to participants' choices and finding the monetary value at which participants

were indifferent to the two options. If participants experience consistency with expectation as

equally rewarding regardless of the target of expectations, then they should choose to forgo

money to rate statements consistent with Obama and Trump in Studies 4a and 4b, respectively.

The results from Study 4a indicate that participants preferred to see expectation-consistent

statements of Barack Obama. When the payoff amounts were equal for confirming and violating

statements, participants chose the consistent statements 60% of the time (significantly more than

chance, as indicated in a generalized mixed model analysis by an odds ratio of 1.42, $Z = 4.74$,

$p < 0.0001$ for the zero-centered intercept in Study 4a). Moreover, the calculated PSE indicated

that, on average, participants gave up 0.39 cents per trial to rate an expectancy-consistent over an

expectancy-violating statement about Obama (95% [CI]: 0.28–0.49, $t_{(188)} = 7.21$, $p < 0.0001$,

$d = 0.52$ [0.37–0.68]; Fig 3C). However, the same was not true for Trump-related statements in

Study 4b. At equal payoff amounts, participants had no preference between

expectancy-consistent and expectation-violating statements (choosing the consistent option 49%

of the time; odds ratio of 1.01, $z = 0.13$, $p = 0.9$), and the calculated PSE was not different from

zero (0.07 cents; [-0.05–0.19], $t_{(173)} = 1.17$, $p = 0.12$, $d = 0.09$ [-0.06–0.24]; Fig 3C). The

difference between the studies was significant ($t_{(328.29)} = 2.28$, $p = 0.023$, $d = 0.25$ [0.03–0.46]).

However, this finding should be interpreted with caution, as our sample was demographically

skewed towards liberals, potentially diluting the effect of consistency with expectations from

Trump with additional factors (see Fig S4). Together, these findings suggest that the rewarding

effect of expectation-consistency is not limited to stereotypes, but also applies to knowledge

about specific individuals. Notably, however, not all sources of knowledge equally contribute to

the reward value; people were not willing to forgo monetary amounts to interact with content

consistent with their expectations about Donald Trump (see SI results for potentially related

findings in Study 3).

Discussion

The human preference for consistent and predictable social interactions has long been

acknowledged as a core motivational component driving everyday behavior [31,32]. To predict the

behavior of others, perceivers regularly employ biased strategies to collect and interpret

information that corresponds to their expectations [59–63]. Here we provide evidence to suggest that

humans associate expectation-consistent information with intrinsic value, much like other forms

of reward such as food or money. Our findings suggest that this reward value is generated

regardless of the source of the social expectation. Participants were willing to forgo money to

rate an expectation-consistent target rather than its expectancy-violating counterpart. Moreover,

doing so was associated with increased activity in a brain region in the neural reward circuitry,

regardless of whether the target was consistent with gender stereotypes or knowledge about US

presidents. Put simply: people find it rewarding to have their expectations (stereotypical or

idiosyncratic) confirmed.

This line of research coincides with emerging theories that highlight the instrumental value of

behaviors and perceptions that fall in line with our expectations. In an early example of this

value, Allport [64] described a mental process in which 'A Scotsman who is penurious delights us

because he vindicates our prejudgment' (p.22). Some recent theories suggest that, because most

of our social expectations are anchored in our social environment, repeated interaction with

expectancy-confirming information leads to continuous reinforcement of our expectations [27,63,65].

Once established, these expectations induce motivations and cognitive representations that

persist even in the face of disconfirmation [28,66,67]. One theory further suggests that the metabolic

costs associated with the violation of expectations increase the desirability of expectation-

consistent behavior from an evolutionary standpoint [29]. In line with these theories, the current

studies demonstrate that information consistent with social expectations is indeed associated with

subjective value.

The current findings provide a neural extension to prominent accounts of implicit (i.e.,

involuntary) stereotyping and prejudice [68,69]. Group-based stereotypes typically draw on

categorical distinctions to facilitate easier decision making by enabling faster and more efficient

processing of stereotype-confirming information [15]. Downstream, perceivers evaluate

expectation-confirming individuals more positively, allocate more economic resources to them,

and judge them as more hirable [18,19,23]. Complementarily, violations of social expectations pose a

threat to individuals and social structures alike, which, in turn, often try to eliminate the threat

and reinforce the original expectation [70]. Our results provide a candidate mechanism for these

effects, whereby the preference of expectation-consistent information translates into a subjective

value that shapes how we evaluate specific individuals [cf., 71]. Furthermore, our results hint at

when perceivers can assign value to expectation-violating information. In our task, behavior

asymmetrically affected the neural response in the nucleus accumbens. Whereas stereotype-

consistent targets always evoked the same level of neural activity regardless of participants'

responses, in Study 1 stereotype-violating targets elicited enhanced NAcc activity only if

perceivers judged them as likely to be associated with the expectancy. This pattern suggests that

participants can assign value to stereotype-violating targets, perhaps depending on the

believability of the counter-stereotypical judgment.

The expectation-consistency account of ventral striatum activity we put forward complements

earlier hypotheses that activity in the mesolimbic circuit reflects prediction errors [72]. Prediction

errors refer to discrepancies between actual and expected outcomes, typically in the context of

learning [73]. Numerous studies have found that the ventral striatum increases its activity as the gap

between the expected and the received outcome grow [e.g., 74,75]. However, the present results

diverge from this prediction error pattern. We observed increased striatal activity when an

outcome—the target face—was <u>not</u> different from the expected outcome; a prediction error

account would suggest that such activity should accompany unexpected information. Notably,

unlike the vast majority of studies exploring the prediction error account, the paradigms we used

in the current investigation did not involve learning. Participants saw each target only once

during the entire study and formed their expectations based solely on previously established

knowledge. Future studies could formally test whether striatal activity in response to expectancy-

consistent information also emerges when participants need to learn the new information.

An additional and important future route of investigation should explore the generalizability

of our findings to non-social contexts. Here we focused on expectations concerning social

targets—unfamiliar men and women as well as familiar targets. We cannot ascertain whether

similar effects would emerge for non-social expectations, such as about inanimate objects or the

weather. Given some accounts suggesting that social information processing has a preferred

status in the primate brain [76–78], we might expect that the subjective value attributed to

consistency should be greater for the kinds of social expectations studied here.

Altogether, these findings join a growing body of literature that characterizes how our prior

beliefs modulate information-processing to fortify a world view and protect established

expectations [36,39,66,79,80]. We suggest that the subjective value imbued upon targets who conform

to societal expectations may serve to sustain multiple stereotype- and expectation-induced biases.

To mitigate the negative implications associated with these expectations, society will need to

acknowledge the subjective value associated with their confirmation.

10   **Materials and Method**

Materials. To create expectation-setting statements we generated a list of verbal statements

that described relevant individual preferences, traits, behaviors or professions. Studies 1 and 2

included 136 gender-related and 68 gender-neutral statements (see SI Materials and Methods).

We verified the stereotypicality of these statements in a pilot study ($n = 78$) in which participants

from the local community indicated how typical the characteristic was for a specific gender on a

visual scale of 0 ("very untypical") to 100 ("very typical"; the scale had no other tick marks).

Each participant was randomly assigned to rate each statement either for men or for women.

Participants were instructed to base their ratings on how they thought the average person would

respond. This verification procedure was successful; men were associated with men-stereotypic

statements more than women (mean difference: 25.7) and women were associated with women-

stereotypic statements more than men (mean difference: 25.1). Overall, the statements contained

2-9 words (mean: 4.69, S.D: 1.44; no difference between experimental conditions, $p > 0.2$) and

9-45 characters (mean: 26.68, S.D.: 7.69; $p > 0.18$). Studies 3 and 4 further included 120 person-

specific statements pertaining to Barack Obama and Donald Trump. A total of 243 participants

from Amazon Mechanical Turk rated a sample of 60 of these statements, randomly determined

per participant. On each trial participants indicated how typical the presented characteristic was

for the two targets (a separate scale for each target; the two scales were presented simultaneously

with a randomly determined order). Obama was associated with Obama-related statements more

than Trump (mean difference: 56.1) and Trump was associated with Trump-related statements

more than Obama (mean difference: 54.5). Overall, the statements contained 2-9 words (mean:

4.65, S.D: 1.52; no difference between experimental conditions, $p > 0.5$) and 11-45 characters

(mean: 28.33, S.D.: 8.43; $p > 0.5$).

Studies 1 and 3. Twenty-eight individuals participated in Study 1 and 30 individuals

participated in Study 3. Additional participants were excluded due to excessive motion, technical

issues or lack of response to more than 20% of trials (3 and 6 participants from Studies 1 and 3,

respectively). All participants provided informed consent in a manner approved by the

Committee on the Use of Human Subjects in Research at Harvard University. Study 3 was pre-

registered (https://osf.io/h9c6x/?view_only=6fa03fc0ceb04e2083db9e485dbe6615). See SI for

demographic information. The current sample size allowed a power of 0.8 to detect a medium

effect size (*Cohen's d* = 0.5) in the planned one-tailed contrast between expectation-consistent

and expectation-violating targets at the region of interest. In both studies participants formed

impressions about target individuals. On each trial, participants first saw a statement for 1.5

seconds. The statements in Study 1 described a stereotypically neutral, stereotypically male, or

stereotypically female characteristic; statements in Study 3 described a characteristic which was

either stereotypically male or female or closely associated with Barack Obama or Donald Trump

(see https://osf.io/tgja3/?view_only=d849bad1b606474f80c1a3e40e740875 for open materials).

Next, participants saw the statement with a face of a man or a woman (in Study 1; Study 3 also

included face images of the relevant leaders). The statement-face pair appeared on screen for 4

additional seconds (3.5 seconds in Study 3) for a total of 5.5 seconds (5 seconds) per trial. Each

trial ended with a 0.5 second fixation crosshair. Participants used their left hand to indicate how

likely the presented target was to be described by the specific characteristic using a 4-point scale

(1- 'very unlikely'; 4 -'very likely'). Participants responded while the pair appeared on screen.

Trials in both studies were separated by variable intertrial intervals of 0-9s [81] optimized for our

contrast of interest (see SI Materials and Methods for details).

To localize brain regions associated with the processing of rewarding stimuli, we defined

8-mm spheres around peak coordinates drawn from a comprehensive meta-analysis [47]. To

functionally identify these brain regions, participants in both studies completed a Monetary

Incentive Delay (MID) task [54] immediately after the impression formation task (see Fig. S1B and

SI Materials and Methods for details). This task allows the identification of monetary-reward-

sensitive ROIs by comparing trials in which participants won money to trials in which

participants could not earn any reward. We extracted and averaged parameter estimates across

voxels in each ROI per condition of interest and analyzed them using within-participant

ANOVAs as implemented by afex package [82] for R, version 0.22-1. We plotted the results using

the package ggstatsplot [83], version 0.2.0. Additionally, we plotted the mean effect size and the

bootstrapped 95% confidence intervals using the package dabestr [84], version 0.2.2.

We collected neuroimaging data with a 3T Siemens Prisma scanner system (Siemens Medical

Systems, Erlangen, Germany). First, we acquired high-resolution anatomical images using a T1-

weighted 3D MPRAGE sequence. Next, whole brain functional images were collected using a

simultaneous multi-slice (multiband) T2*-weighted gradient echo sequence (TR = 2000 msec,

TE = 30 msec, voxel size = 2 × 2 × 2 mm3, 75 slices auto-aligned to -25 degrees of the AC-PC

line). Participants completed four impression formation task runs consisting of 229 volumes each

(245 volumes in Study 3). Finally, participants completed the MID task in a single run consisting

of 110 volumes using identical parameters to those mentioned above. We used SPM12 version

6225 (Wellcome Department of Cognitive Neurology, London, UK) to process and analyze the

fMRI data. Data were corrected for differences in acquisition time between slices, corrected for

inhomogeneities in the magnetic field using fieldmap [85], realigned to the first image to correct for

head movement, unwarped to account for residual movement-related variance and co-registered

with each participant's anatomical data. Functional data were then transformed into a standard

anatomical space (2 mm isotropic voxels) based on the ICBM152 brain template (Montreal

Neurological Institute). Normalized data were then spatially smoothed (6 mm full-width at half-

maximum, FWHM) using a Gaussian Kernel (see SI Materials and Methods for full details of

scanning and analysis procedures). We analyzed preprocessed data using a general linear model

in which we modeled trials as boxcar functions with an onset at face presentation (1.5s after

statement presentation) and with variable duration determined per trial by reaction time to

control for effects of reaction time on the neural response [86]. Our main analysis included a model

in which we conditionalized trials based on trial type (stereotype-consistent, stereotype-neutral or

stereotype-violating trials in Study 1; stereotype-consistent, stereotype-violating, person-

specific-confirming and person-specific-violating trials in Study 3). In our secondary analysis

(Fig 2) we split each of the trial types included in the main analysis with two regressors, one in

which participants provided a "Likely" rating and one in which they provided an "Unlikely"

rating. We convolved events with a canonical hemodynamic response function and its temporal

derivative and included additional covariates of no interest (session mean, no response trials, six

motion parameters and their temporal derivative). The final first-level GLM was high-pass

filtered at 128 s. Analyses were performed individually for each participant, and contrast images

were subsequently entered into a second-level analysis treating participants as a random effect.

We report activations that survived a threshold of p<0.001 (uncorrected) at the voxel level and

(cluster-size) corrected to p<0.05 at the cluster level using Monte Carlo simulations (1,000

iterations) with the current imaging and analysis parameters [87].

Studies 2 and 4. A total of 343 participants were included in Study 2 (174 in Study 2a) and

361 in Study 4 (189 in Study 4a). Additional participants were excluded by criteria set in the pre-

registered protocols for each study (see SI Materials and Methods for details; see also

https://aspredicted.org/blind.php?x=rt3wi6 (Study 2a),

https://aspredicted.org/blind.php?x=db3gz4 (Study 2b), and

https://aspredicted.org/blind.php?x=9z83yb (Studies 4a and 4b)). Sample size was set to allow

sufficient power (0.8) to detect a small effect size (*Cohen's d* = 0.2) in a one-sample t-test for

each study. Informed consent was obtained from all participants in a manner approved by the

Committee on the Use of Human Subjects at Harvard University. We analyzed choice data with

logit generalized linear mixed models as implemented in the lme4 package version 1.1-14 [88] for

R version 3.4.2 (R Core Team, 2017). We used the probit link and included fixed effects for the

intercept and for the value difference between the two target types, as well as random effects for

the intercepts for participants for the by-participant random slopes for the fixed effect of value

difference. We calculated the PSE (and the related SEs) for the difference between the two target

types for each study by the Delta Method as implemented in the MixedPsy package [57,89] for R.

The Delta Method relies on responses to <u>all trials</u> aggregated across participants in a generalized

linear model to approximate the PSE with a Gaussian distribution [57] and to plot the cumulative

distribution function. The model included the difference between the two target types on each

trial as the predictor value and a binary outcome (stereotypical/knowledge-consistent option

chosen) as the predicted value, with the probit link. To present individual-level data (Figure 3B

and 3C), we calculated the PSE for each participant. This resulted in some PSE values that

exceeded the possible values in the current studies, as PSE models do not accurately reflect

behavior when one choice option is rarely selected, as was the case for some participants.

Therefore, we excluded participants for whom the calculated PSE value exceeded ± 6 ($n = 4\text{--}14$

across studies).

Acknowledgements

This work was partially supported by the Israeli Science Foundation (No. 79/18, to N.R.).

N.R was supported by Yad Ha'Nadiv (Rothschild) Foundation and the Mind, Brain and Behavior

interfaculty program at Harvard University. Parts of this work were conducted at the Harvard

Center for Brain Science, which is supported by the NIH Shared Instrumentation Grant Program

(No. S10OD020039). We thank Kirstan Brodie for her assistance with data collection for Study

3.

Data availability

The data that support the findings of this study are available from the corresponding author

(N.R.) upon reasonable request.

References

1. Tay, L., Tan, K., Diener, E. & Gonzalez, E. Social relations, health behaviors, and health outcomes: A survey and synthesis. *Appl. Psychol. Heal. Well-Being* **5**, 28–78 (2013).

2. Baumeister, R. F. & Leary, M. R. The need to belong: Desire for interpersonal attachments as a fundamental human motivation. *Psychol. Bull.* **117**, 497–529 (1995).

3. Cacioppo, J. T., Cacioppo, S. & Boomsma, D. I. Evolutionary mechanisms for loneliness. *Cogn. Emot.* **28**, 1–22 (2014).

4. Matthews, G. A. & Tye, K. M. Neural mechanisms of social homeostasis. *Ann. N. Y. Acad. Sci.* 1–21 (2019) doi:10.1111/nyas.14016.

5. Otten, M., Seth, A. K. & Pinto, Y. A social Bayesian brain: How social knowledge can shape visual perception. *Brain Cogn.* **112**, 69–77 (2017).

6. Hutchinson, J. B. & Barrett, L. F. The power of predictions: An emerging paradigm for psychological research. *Curr. Dir. Psychol. Sci.* **28**, 280–291 (2019).

7. Tamir, D. I. & Thornton, M. A. Modeling the predictive social mind. *Trends Cogn. Sci.* **22**, 201–212 (2018).

8. Freeman, J. B. & Johnson, K. L. More than meets the eye: Split-second social perception. *Trends Cogn. Sci.* **20**, 362–374 (2016).

9. Macrae, C. N. & Bodenhausen, G. V. Social cognition: Thinking categorically about others. *Annu. Rev. Psychol.* **51**, 93–120 (2000).

10. Fiske, S. T. & Neuberg, S. L. A continuum of impression formation, from category-based to individuating processes: Influences of information and motivation on attention and interpretation. in *Advances in Experimental Social Psychology* (ed. Zanna, M. P.) vol. 23 1–74 (Academic Press, 1990).

11. Gregg, A. P., Seibt, B. & Banaji, M. R. Easier done than undone: Asymmetry in the malleability of implicit ireferences. *J. Pers. Soc. Psychol.* **90**, 1–20 (2006).

12. Dunsmoor, J. E., Kubota, J. T., Li, J., Coelho, C. A. O. & Phelps, E. A. Racial stereotypes impair flexibility of emotional learning. *Soc. Cogn. Affect. Neurosci.* **11**, 1363–1373 (2016).

13. Wyer, N. A. You never get a second chance to make a first (implicit) impression: The role of elaboration in the formation and revision of implicit impressions. *Soc. Cogn.* **28**, 1–19 (2010).

14. Hamilton, D. L. & Sherman, S. J. Perceiving persons and groups. *Psychol. Rev.* **103**, 336–355 (1996).

15. Roese, N. J. & Sherman, J. W. Expectancy. in *Social Psychology: handbook of basic principles* (eds. Higgins, E. T. & Kruglanski, A. W.) 91–115 (The Guilford Press, 2007). doi:10.1017/S0030605315000393.

16. Asch, S. E. Forming impressions of personality. *J. Abnorm. Soc. Psychol.* **41**, 258–290 (1946).

17. Sullivan, J. The primacy effect in impression formation: Some replications and extensions. *Soc. Psychol. Personal. Sci.* **10**, 432–439 (2019).

18. Phelan, J. E. & Rudman, L. A. Reactions to ethnic deviance: The role of backlash in racial stereotype maintenance. *J. Pers. Soc. Psychol.* **99**, 265–281 (2010).

19. Stern, C., West, T. V. & Rule, N. O. Conservatives negatively evaluate counterstereotypical people to maintain a sense of certainty. *Proc. Natl. Acad. Sci.* **112**, 15337–15342 (2015).

20. Eagly, A. H. & Karau, S. J. Role congruity theory of prejudice toward female leaders. *Psychol. Rev.* **109**, 573–598 (2002).

21. Rudman, L. A., Moss-Racusin, C. A., Phelan, J. E. & Nauts, S. Status incongruity and backlash effects: Defending the gender hierarchy motivates prejudice against female leaders. *J. Exp. Soc. Psychol.* **48**, 165–179 (2012).

22. Moss-Racusin, C. A. & Johnson, E. R. Backlash against male elementary educators. *Journal of Applied Social Psychology* vol. 46 379–393 (2016).

23. Stern, C. & Rule, N. O. Physical androgyny and categorization difficulty shape political conservatives' attitudes toward transgender people. *Soc. Psychol. Personal. Sci.* **9**, 24–31 (2018).

24. Olszanowski, M., Kaminska, O. K. & Winkielman, P. Mixed matters: fluency impacts trust ratings when faces range on valence but not on motivational implications. *Cogn. Emot.* **32**, 1032–1051 (2018).

25. Chanes, L., Wormwood, J. B., Betz, N. & Barrett, L. F. Facial expression predictions as drivers of social perception. *J. Pers. Soc. Psychol.* **114**, 380–396 (2018).

26. Jost, J. & Hunyady, O. The psychology of system justification and the palliative function of ideology. *Eur. Rev. Soc. Psychol.* **13**, 111–153 (2003).

27. Huebner, B. Implicit bias, reinforcement learning, and scaffolded moral cognition. in *Implicit Bias and Philosophy, Volume 1: Metaphysics and Epistemology* (eds. Brownstein, M. & Saul, J.) 47–79 (Oxford University Press, 2016). doi:10.1093/acprof:oso/9780198713241.001.0001.

28. Berridge, K. C. From prediction error to incentive salience: Mesolimbic computation of reward motivation. *Eur. J. Neurosci.* **35**, 1124–1143 (2012).

29. Theriault, J. E., Young, L. & Barrett, L. F. The sense of should: A biologically-based framework for modeling social pressure. *Phys. Life Rev.* (2020) doi:10.1016/j.plrev.2020.01.004.

30. Feldmanhall, O. & Shenhav, A. Resolving uncertainty in a social world. *Nat. Hum. Behav.* **3**, 426–435 (2019).

31. Festinger, L. *A theory of cognitive dissonance (Vol. 2)*. (Stanford university press, 1957).

32. Gawronski, B. Back to the future of dissonance theory: Cognitive consistency as a core motive. *Soc. Cogn.* **30**, 652–668 (2012).

33. Kohli, A. *et al.* Using Expectancy Theory to quantitatively dissociate the neural representation of motivation from its influential factors in the human brain: An fMRI study. *Neuroimage* **178**, 552–561 (2018).

34. Berridge, K. C. From prediction error to incentive salience□: mesolimbic computation of reward motivation. *Eur. J. Neurosci.* **35**, 1124–1143 (2012).

35. Sharot, T., Guitart-masip, M., Korn, C. W., Chowdhury, R. & Dolan, R. J. How Dopamine enhances an optimism bias in humans. *Curr. Biol.* **22**, 1477–1481 (2012).

36. Charpentier, C. J., Bromberg-Martin, E. S. & Sharot, T. Valuation of knowledge and ignorance in mesolimbic reward circuitry. *Proc. Natl. Acad. Sci.* **115**, E7255–E7264 (2018).

37. Lefebvre, G., Lebreton, M., Meyniel, F., Bourgeois-Gironde, S. & Palminteri, S. Behavioural and neural characterization of optimistic reinforcement learning. *Nat. Hum. Behav.* **1**, 1–9 (2017).

38. Schwarz, K. A. *et al.* How stereotypes affect pain. *Sci. Rep.* **9**, 8626 (2019).

39. Jepma, M., Koban, L., Doorn, J., Jones, M. & Wager, T. D. Behavioural and neural evidence for self-reinforcing expectancy effects on pain. *Nat. Hum. Behav.* **2**, 838–855 (2018).

40. Welborn, B. L., Hong, Y. & Ratner, K. G. Exposure to negative stereotypes influences representations of monetary incentives in the nucleus accumbens. *Soc. Cogn. Affect. Neurosci.* 261–271 (2020) doi:10.1093/scan/nsaa041.

41. Wu, H., Luo, Y. & Feng, C. Neural signatures of social conformity: A coordinate-based activation likelihood estimation meta-analysis of functional brain imaging studies. *Neurosci. Biobehav. Rev.* **71**, 101–111 (2016).

42. Schultz, W. Multiple reward signals in the brain. *Nat. Rev. Neurosci.* **1**, 199–207 (2000).

43. Haber, S. N. & Knutson, B. The reward circuit: Linking primate anatomy and human imaging. *Neuropsychopharmacology* **35**, 4–26 (2010).

44. Hare, T. A., O'Doherty, J. P., Camerer, C. F., Schultz, W. & Rangel, A. Dissociating the role of the orbitofrontal cortex and the striatum in the computation of goal values and prediction errors. *J. Neurosci.* **28**, 5623–5630 (2008).

45. Sescousse, G., Caldú, X., Segura, B. & Dreher, J.-C. Processing of primary and secondary rewards: A quantitative meta-analysis and review of human functional neuroimaging studies. *Neurosci. Biobehav. Rev.* **37**, 681–696 (2013).

46. Peters, J. & Büchel, C. Neural representations of subjective reward value. *Behav. Bran Res.* **213**, 135–141 (2010).

47. Bartra, O., McGuire, J. T. & Kable, J. W. The valuation system: A coordinate-based meta-analysis of BOLD fMRI experiments examining neural correlates of subjective value.

Neuroimage **76**, 412–427 (2013).

48. Lin, A., Adolphs, R. & Rangel, A. Social and monetary reward learning engage overlapping neural substrates. *Soc. Cogn. Affect. Neurosci.* **7**, 274–281 (2012).

49. Hackel, L. M., Doll, B. B. & Amodio, D. M. Instrumental learning of traits versus rewards: dissociable neural correlates and effects on choice. *Nat. Neurosci.* **18**, 1233–1240 (2015).

50. Harris, L. T. & Fiske, S. T. Neural regions that underlie reinforcement learning are also active for social expectancy violations. *Soc. Neurosci.* **5**, 76–91 (2010).

51. Krosch, A. R. & Amodio, D. M. Scarcity disrupts the neural encoding of Black faces: A aocioperceptual pathway to discrimination. *J. Pers. Soc. Psychol.* **117**, 859–875 (2019).

52. Deaner, R. O., Khera, A. V. & Platt, M. L. Monkeys pay per view: Adaptive valuation of social images by rhesus macaques. *Curr. Biol.* **15**, 543–548 (2005).

53. Tamir, D. I. & Mitchell, J. P. Disclosing information about the self is intrinsically rewarding. *Proc. Natl. Acad. Sci.* **109**, 8038–8043 (2012).

54. Knutson, B., Westdorp, A., Kaiser, E. & Hommer, D. FMRI visualization of brain activity during a monetary incentive delay task. *Neuroimage* **12**, 20–27 (2000).

55. Bhanji, J. P. & Delgado, M. R. The social brain and reward: Social information processing in the human striatum. *Wiley Interdiscip. Rev. Cogn. Sci.* **5**, 61–73 (2014).

56. O'Doherty, J. P. The problem with value. *Neurosci. Biobehav. Rev.* **43**, 259–268 (2014).

57. Moscatelli, A., Mezzetti, M. & Lacquaniti, F. Modeling psychophysical data at the population-level: The generalized linear mixed model. *J. Vis.* **12**, 26–26 (2012).

58. Hayden, B. Y., Parikh, P. C., Deaner, R. O. & Platt, M. L. Economic principles motivating social attention in humans. *Proc. R. Soc. B* **274**, 1751–1756 (2007).

59. Johnston, L. C. & Macrae, C. N. Changing social stereotypes: The case of the information seeker. *European Journal of Social Psychology* vol. 24 581–592 (1994).

60. Frimer, J. A., Skitka, L. J. & Motyl, M. Liberals and conservatives are similarly motivated to avoid exposure to one another's opinions. *J. Exp. Soc. Psychol.* **72**, 1–12 (2017).

61. Falben, J. *et al.* Predictably confirmatory: The influence of stereotypes during decisional processing. *Q. J. Exp. Psychol.* **72**, 2437–2451 (2019).

62. Lord, C. G., Ross, L. & Lepper, M. R. Biased assimilation and attitude polarization: The effects of prior theories on subsequently considered evidence. *J. Pers. Soc. Psychol.* **37**, 2098–2109 (1979).

63. Oyserman, D. & Yan, V. X. Making meaning: A culture-as-situated-cognition approach to the consequences of cultural fluency and disfluency. in *Handbook of Cultural Psychology* (eds. Cohen, D. & Kitayama, S.) 536–565 (Guilford Press, 2019). doi:10.5840/philstudies197120057.

64. Allport, G. W. *The nature of prejudice*. (Addison–Wesley, 1954).

65. Peters, U. What is the function of confirmation bias? *Erkenntnis* (2020) doi:10.1007/s10670-020-00252-1.

66. Yon, D., Lange, F. P. De & Press, C. The predictive brain as a stubborn scientist. *Trends Cogn. Sci.* **23**, 6–8 (2019).

67. Uusberg, A., Suri, G., Dweck, C. S. & Gross, J. J. Motivation: A valuation systems perspective. in *Emotion in the mind and body. Nebraska symposium on motivation. Vol. 66* (eds. Neta, M. & Haas, I. J.) 161–192 (Springer, Cham, 2019).

68. Tibboel, H., Houwer, J. De & Bockstaele, B. Van. Implicit measures of "wanting" and "liking" in humans. *Neurosci. Biobehav. Rev.* **57**, 350–364 (2015).

69. Greenwald, A. G. & Lai, C. K. Implicit social cognition. *Annu. Rev. Psychol.* **71**, (2020).

70. Morgenroth, T. & Ryan, M. K. The Effects of Gender Trouble: An Integrative Theoretical Framework of the Perpetuation and Disruption of the Gender/Sex Binary. *Perspect. Psychol. Sci.* (2020) doi:10.1177/1745691620902442.

71. Amodio, D. M. & Devine, P. G. Stereotyping and evaluation in implicit race bias: Evidence for independent constructs and unique effects on behavior. *J. Pers. Soc. Psychol.* **91**, 652–661 (2006).

72. Daw, N. D., Gershman, S. J., Seymour, B., Dayan, P. & Dolan, R. J. Model-based influences on humans' choices and striatal prediction errors. *Neuron* **69**, 1204–1215 (2011).

73. Niv, Y. & Schoenbaum, G. Dialogues on prediction errors. *Trends Cogn. Sci.* **12**, 265–272 (2008).

74. Glimcher, P. W. Understanding dopamine and reinforcement learning: The dopamine reward prediction error hypothesis. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 15647–15654 (2011).

75. Ballard, I., Miller, E. M., Piantadosi, S. T., Goodman, N. D. & McClure, S. M. Beyond Reward Prediction Errors: Human Striatum Updates Rule Values during Learning. *Cereb. Cortex* **28**, 3965–3975 (2018).

76. Adolphs, R. The social brain: Neural basis of social knowledge. *Annu. Rev. Psychol.* **60**, 693–716 (2009).

77. Atzil, S., Gao, W., Fradkin, I. & Barrett, L. F. Growing a social brain. *Nat. Hum. Behav.* **2**, (2018).

78. Lockwood, P. L., Apps, M. A. J. & Chang, S. W. C. Is there a 'social' brain? Implementations and algorithms. *Trends Cogn. Sci.* **24**, 802–813 (2020).

79. Gershman, S. J. How to never be wrong. *Psychon. Bull. Rev.* **26**, 13–28 (2019).

80. Golman, R., Hagmann, D. & Loewenstein, G. Information avoidance. *J. Econ. Lit.* **55**, 96–135 (2017).

81. Dale, A. M. Optimal experimental design for event-related fMRI. *Hum. Brain Mapp.* **8**,

1   109–114 (1999).

2   82.   Singmann, H., Bolker, B., Westfall, J. & Aust, F. afex: Analysis of factorial experiments.
3   (2018).

4   83.   Patil, I. ggstatsplot: 'ggplot2' Based Plots with Statistical Details. (2018)
5   doi:10.5281/zenodo.2074621.

6   84.   Ho, J., Tumkaya, T., Aryal, S., Choi, H. & Claridge-Chang, A. Moving beyond P values:
7   data analysis with estimation graphics. *Nat. Methods* **16**, 565–566 (2019).

8   85.   Cusack, R. & Papadakis, N. New robust 3-D phase unwrapping algorithms: Application to
9   magnetic field mapping and undistorting echoplanar images. *Neuroimage* **16**, 754–764
(2002).

86.   Grinband, J., Wager, T. D., Lindquist, M., Ferrera, V. P. & Hirsch, J. Detection of time-
varying signals in event-related fMRI designs. *Neuroimage* **43**, 509–520 (2008).

87.   Slotnick, S. D. Cluster success: fMRI inferences for spatial extent have acceptable false-
positive rates. *Cogn. Neurosci.* **8**, 150–155 (2017).

88.   Bates, D. M., Maechler, M., Bolker, B. & Walker, S. lme4: Linear mixed-effects models
using Eigen and S4. (2014).

89.   Moscatelli, A. & Balestrucci, P. Psychophysics with R: The R package MixedPsy. (2017).

Supplementary Information for

Confirmation of interpersonal expectations is intrinsically rewarding

Niv Reggev, Anoushka Chowdhary and Jason P. Mitchell, Harvard University.

Corresponding author: Niv Reggev

Email: reggevn@bgu.ac.il

This PDF file includes:

Supplementary text

Figures S1 to S4

Tables S1 to S5

SI References

SI Materials and Methods

Below we provide a detailed description of all the materials and methods used including all

measures and data exclusions.

Participants

Studies 1 and 3. Twenty-eight individuals participated in Study 1 (mean age: 21.85, S.D.:

2.95, range: 18–30, 21 females, 15 Caucasian, 4 Hispanic, 4 mixed, 3 Asian, 2 African

American) and 30 individuals participated in Study 3 (mean age: 22.37, S.D.: 2.62, range: 18–30,

17 females, 14 Caucasian, 7 Asian, 4 African American, 3 mixed, 1 Hispanic, 1 did not self-

identify). Participants were recruited from Harvard University and its surroundings using

Harvard's Psychology Study Pool website. Additional participants were excluded due to

technical issues (1 participant from Study 3), lack of response to more than 20% of trials (2

participants from each study) or excessive movement (more than 1mm – half-width of the

acquired voxel size) in more than one functional run in the scanner (1 participant in Study 1, 3

participants in Study 3). All participants were healthy, right-handed, native English speakers

with normal or corrected-to-normal vision and no history of neurological or psychiatric

conditions. Participants were compensated with $65 in Study 1 and $55 in Study 3. We

determined supplemental compensation (up to $30) based on participants' performance in the

monetary incentive delay task (see below). Pilot versions of the imaging tasks were conducted

outside the scanner to ensure the functionality of the task.

Studies 2 and 4. Three-hundred-ninety-six individuals from Amazon Mechanical Turk

completed Study 2 (in Study 2a, 198 out of 216 individuals who started the study; in Study 2b,

198 out of 223 individuals). Study 2a was conducted in March 2018, and Study 2b was

conducted in April 2018. Four-hundred-seventy individuals from Prolific Academic completed

Studies 4a and 4b in January 2019 (235 in each study; a total of 503 participants started the

study). All participants were 18 years old or older, had an approval rate of 95% or higher, held a

US nationality and participated only in one of the studies. In addition, participants from Amazon

Mechanical Turk were required to have completed at least 100 tasks to be eligible for Study 2.

The final sample size we report in the main text incorporates the following pre-registered

exclusion criteria for participants who (1) answered incorrectly at least two of the manipulation

or attention checks, (2) completed the survey in more than 20 minutes or less than 2.5 standard

deviations below the sample mean duration, (3) failed to answer more than 15% of the survey,

(4) provided similar ratings (on the 0–100 scale) on all trials; specifically, participants whose

standard variation of the rating was 2 standard deviations below the sample mean standard

deviation (not applied in Study 2a), (5) provided an identical response in all 2AFC trials (applied

only in Study 4) and (6) indicated in their debriefing that they had understood the goal of the task

and had acted per this understanding at any point during the task. Participants in all studies were

similarly distributed on gender (Study 2a: 51.72% males, 45.98% females; Study 2b: 50.3%

males, 48.52% females; Study 4a: 51.32% males, 46.56% females; Study 4b: 41.86% males,

56.98% females; across studies 1%–3% of participants self-identified as gender-nonconforming

or other identification), age (Study 2a: Mean [S.D.]: 37.47 [11.75]; Study 2b: 35.13 [10.04];

Study 4a: 31.22 [10.32]; Study 4b: 31.99 [10.97]) and ethnicity (Study 2a: 78.16%

White/European American; Study 2b: 74.56% White/European American; Study 4a: 69.84%

White/European American; Study 4b: 69.19% White/European American). Study 4 measured

political orientation on a 1–9 scale (see below). We observed no differences on this measure

between Study 4a and Study 4b (Study 4a: 3.79 [2.13], Study 4b: 3.45 [1.96]).

Stimuli

All studies included face-statement pairs. We selected faces from the 10k US Adult Faces

Database (a large-scale database of natural face photographs of the U.S. adult population (1)).

The current investigation focused on Caucasian faces to avoid intrusion of racial or intersectional

stereotypes. We further restricted the faces to be moderately memorable (0.4 – 0.6 hit rate in the

10k database, computed over an average of 81.7 people per photo) and excluded faces with any

kind of distinguishable accessories (e.g. hats, big necklaces, etc.). This resulted in a total of 204

faces, 102 per gender. Three additional photos per gender were used in the practice sessions. All

face pictures had neutral to mildly positive expressions. We resized all pictures to 240 by 256

pixels and presented them in color on a gray background (see Fig S1). For Studies 3 and 4 we

also included images portraying the faces of Barack Obama and Donald Trump (a single image

per leader). These images were taken from open access sources and cropped to be identical in

dimensions to the rest of the face stimuli.

We generated gender-stereotype-relevant statements in the form of individual preference,

trait, behavior or profession (e.g., "Loves taking risks", "Doesn't cry", "Swears a lot", "Is a truck

driver", etc.; see https://osf.io/tgja3/?view_only=d849bad1b606474f80c1a3e40e740875). We

piloted the verbal statements in a two-phase procedure using non-overlapping groups of

participants that did not participate in the reported studies. In the first session, 23 participants

rated 160 gender-related statements generated by the authors based on various sources (e.g., 2).

Participants saw each statement once and indicated how typical the characteristic was for a

specific gender on a visual scale of 0 (= "very untypical") to 100 (= "very typical"; the scale had

no other tick marks). Each participant rated each statement either for men or for women, never

for both, randomly determined per participant. Participants were instructed to base their ratings

on how they thought the average person would respond. Based on the ratings from this first

phase, we selected 136 gender-related statements (68 per gender) for which the average rating

was at least 65 for one of the genders and less than 35 for the other gender. In a second session of

the pilot testing, we verified the stereotypicality of these statements with a total of 78 participants

who used the same rating procedure to rate the selected 136 statements and an additional 68

gender-neutral statements (e.g., "Exercises regularly", "Thinks positively", "Is a TV reporter").

The statements included in the studies contained 2–9 words (mean: 4.69, S.D: 1.44; no difference

between experimental conditions, $p > 0.2$) and 9–45 characters (mean: 26.68, S.D.: 7.69; $p >$

0.18).

We utilized a similar procedure for leader-relevant statements. First, we generated 160

statements based on common knowledge about the selected leaders. We constructed the

statements such that a statement consistent with knowledge about one leader would violate

knowledge about the other leader. We piloted these statements with 243 Amazon Mechanical

Turk participants. Each participant saw a randomly selected subset of 40 or 60 statements and

rated them on two scales: "how likely is the statement to be attributed to Barack Obama" and

"how likely is the statement to be attributed to Donald Trump" on a visual scale of 0 (= "very

unlikely") to 100 (= "very likely"). Participants were instructed to base their ratings on how they

thought the average person would respond. Each statement had an average of 81.75 raters (S.D.:

6.23; range: 67–100). We selected the 120 statements that differed the most between ratings for

the two leaders. Overall, the statements contained 2–9 words (mean: 4.65, S.D: 1.52; no

difference between experimental conditions, $p > 0.5$) and 11–45 characters (mean: 28.33, S.D.:

8.43; $p > 0.5$).

Behavioral procedure

Studies 1 and 3. The impression formation task included 204 face-statement pairs in Study 1 and

240 face-statement pairs in Study 3. To create stereotype-consistent and stereotype-violating

pairs, we yoked half of the 136 stereotypical statements (120 statements in Study 3) with faces

from the gender corresponding to the stereotype and the other half with faces from the

mismatching gender (yoking of statements was randomized across participants; e.g., the

statement "Doesn't cry" could be paired with a male face for one participant and with a female

face for another participant to create a stereotype-consistent or a stereotype–violating pair,

respectively). Gender-neutral statements (presented only in Study 1; 68 statements) were

randomly and evenly yoked with male and female faces. We randomized the specific face

identity paired with each statement across participants. We used a similar procedure to create

leader-specific pairs (60 statements per leader), with the exception that a single photo was used

per leader.

Impression Formation Task. Each trial in the Impression Formation task (see Fig S1) started

with a statement (describing a neutral, stereotypically male or stereotypically female

characteristic in Study 1; stereotypically male, stereotypically female, Obama-specific or Trump-

specific in Study 3) presented for 1.5 seconds. Then, a face joined the statement to form a face-

statement pair that was either consistent or inconsistent (or, in Study 1, neutral) with gender

stereotypes or (in Study 3) person-specific knowledge. The pair appeared on screen for 4 seconds

for a total of 5.5 seconds per trial in Study 1, and 3.5 seconds for a total of 5 seconds in Study 3.

Each trial ended with a 0.5-second fixation crosshair. For each pair, participants used their left

hand to indicate how likely the presented target was to be described by the specific characteristic

using a 4-point scale (1- 'very unlikely'; 4 -'very likely'). Participants were to provide a response

while the pair appeared on screen.

Each of four functional runs included 51 (in Study 1) or 60 (in Study 3) unique face-

statement pairs (17 per condition in Study 1, 15 per condition in Study 3). In Study 3 we

presented statements in mini-blocks by knowledge domain (stereotypes versus person-specific),

2 blocks per type per run; each mini-block contained 15 trials. Order of the mini-blocks was

randomly determined within each run with a limitation that consecutive blocks never displayed

the same type of content. To optimize estimation of the event-related fMRI response, we

intermixed conditions in a pseudorandom order and separated trials by a variable interstimulus

interval (Study1: 0–9 seconds, mean: 3.03, S.D.: 2.25; Study 3: 0–7 seconds, mean: 1.24, S.D:

1.84). We used OptSeq2 (3) to generate sequences optimized for the efficiency of a 3-conditions

design for a first-order counterbalanced event sequence in Study 1 and for the efficiency of the

(expectancy-consistent versus expectancy-violating) contrast for a first-order counterbalanced

event sequence in Study 3. Of these sequences, we selected 6 sequences that contained no more

than 5 consecutive events of the same condition (separate sequences were generated and selected

per Study). We randomly assigned (with replacement) an event sequence for each functional run

to avoid spurious results attributable to differences between conditions in one specific event

sequence (4). Within conditions, trials were presented in random order. To facilitate

familiarization with the task, participants completed a brief practice session before entering the

MRI machine. This practice session included six statement-face pairs in Study 1 and 12 pairs in

Study 3; the stimuli used in the practice were not used in any other phase of the experiment.

Monetary Incentive Delay (MID) Task. After completing the impression formation task,

participants in Studies 1 and 3 completed the Monetary-Incentive Delay (MID) task (5) to allow

us to localize brain regions associated with the processing of rewarding stimuli in a non-social

context. Participants were not informed about this task before its execution to prevent them from

forming an association between the main task and reward processing.

The MID task included a series of trials in which participants attempted to respond, via a

button press, to a briefly presented target (a white rectangle) (see Fig S1B). Each trial started

with a cue (a blue circle or a green circle) that was presented for 0.5 seconds. The green circle

predicted a modest monetary reward ($1) upon a successful response to the target, whereas the

blue circle predicted no reward. Nevertheless, participants were instructed to respond to both cue

types. Cues were followed by a delay interval randomly varying in duration between 2 and 2.5

seconds. The target was then briefly presented for a duration varied between 130 and 350

milliseconds. Duration varied as a function of the participants' performance. Specifically, we

implemented a 2-down 1-up staircase procedure to create a level of difficulty that would allow

participants to successfully respond to the target on two-thirds of the trials. This algorithm

succeeded; on average, participants were rewarded on approximately 20 of the 30 trials (Study 1

mean: 20.26; Study 3: 20.21). At the end of each trial, participants saw the amount of money

they had earned on that trial along with the total amount they had earned during the task up to

that point (presented for 0.5 seconds). The task included 45 trials (with 30 green cue trials). We

added participants' gains in this task to their overall compensation.

Memory Test. Once outside the scanner, participants completed a surprise associative

memory task that will be reported elsewhere (Reggev & Mitchell, in preparation). Briefly, none

of the results reported in the current manuscript were affected by including memory in the

analyses.

Additional measures. Next, participants completed several individual differences and

explicit attitudes scales measuring beliefs about sexism (Ambivalent Sexism Inventory – ASI)

(6), social dominance orientation (SDO) (7), motivation to control sexism (MCS) (8) and need

for cognitive closure (NFC) (9). Order of the scales was randomized between participants. We

included these scales to facilitate future individual differences analyses. Individual differences

scores were not used in the current manuscript in any of the analyses due to insufficient power

and are reported solely for full disclosure's sake.

Finally, participants in Study 1 indicated the extent to which they thought the different

statements presented during the impression formation task were associated with women and men

using the procedure used for piloting the statements. We presented each statement with the

gender to which it was yoked in the impression formation task. Participants had up to 10 seconds

per trial and were told that they should base their judgments on their own beliefs about men and

women, rather than on what the "average" person in the population thinks.

After completing these tasks, participants provided demographic details (age, self-identified

gender and self-identified race in an open response format, and in Study 3 political affiliation on

a 1–9 scale, 1 = "extremely liberal" and 9 = "extremely conservative"). Then, we probed

participants for their understanding of the goal of the study and asked whether they had

suspected a memory test. Lastly, we paid and fully debriefed them.

Imaging procedure

Images were collected with a 3T Siemens Prisma scanner system (Siemens Medical

Systems, Erlangen, Germany) using a 64-channel radiofrequency head coil. Stimuli were

projected onto a screen at the end of the magnet bore that participants viewed via a mirror

mounted on the head coil. Stimulus presentation was controlled by PsychoPy v1.84.2(10)

running under Windows 7. Prior to entering the scanner participants were extensively briefed by

one of the authors about potential movements that can occur in the scanner and ways to mitigate

them. Participants were then set up in the scanner, head first and supine in the scanner bore, with

a response box in their left hand. Foam cushions were placed within the head coil to minimize

head movements. First, high-resolution anatomical images were acquired using a T1-weighted

3D MPRAGE sequence (TR = 2200 msec, TI = 1100 msec, acquisition matrix = $256 \times 256 \times$

176, flip angle = 7, voxel size = $1 \times 1 \times 1$ mm^3). Second, a fieldmap was acquired in the same

plane as the functional images (see below) to correct for inhomogeneities in the magnetic field

(11). Next, whole-brain functional images were collected using a simultaneous multi-slice

(multiband) T2*-weighted gradient echo sequence, sensitive to BOLD contrast, developed at the

Center for Magnetic Resonance Research (CMRR) at University of Minnesota (12–14) (TR =

2000 msec, TE = 30 msec, voxel size = $2 \times 2 \times 2$ mm^3, 75 slices auto-aligned to -25 degrees of

the AC-PC line, image matrix = 104×104, FOV = 208 * 208 mm^2, flip angle = 75º, GRAPPA

acceleration factor = 2, multiband factor = 3, phase encoding direction = A -> P). After a brief

practice run (identical in content to the practice session completed before entering the scanner),

participants completed four impression formation task runs consisting of 229 volumes each in

Study 1 and 245 volumes each in Study 3; all runs were complemented by two additional dummy

scans and an initial period of approximately 26 s dedicated to references for the GRAPPA

procedure. The first four volumes from each run (i.e., in addition to dummy scans) were

discarded to ensure T1 equilibrium. The last 5 volumes from each run always included a

crosshair fixation to ensure the appropriate estimation of the hemodynamic function for the last

events in the run. Finally, participants completed the MID task in a single run consisting of 110

volumes using identical parameters to those mentioned above.

Imaging analysis

We processed and analyzed the fMRI data using SPM12 version 6225 (Wellcome

Department of Cognitive Neurology, London, UK) on a 2015b MATLAB platform (Mathworks,

Natick, MA, USA). Functional data were corrected for differences in acquisition time between

slices, corrected for inhomogeneities in the magnetic field using the fieldmap (11), realigned to

the first image to correct for head movement using a 2^{nd} degree B-spline interpolation, unwarped

to account for residual movement-related variance using a 4^{th} degree B-spline interpolation and

co-registered with each participant's anatomical data. Then, the functional data were transformed

into standard anatomical space (2 mm isotropic voxels) based on the ICBM152 brain template

(Montreal Neurological Institute). Normalized data were spatially smoothed (6 mm full-width at

half-maximum, FWHM) using a Gaussian Kernel. In addition to the GLM models reported in the

main text, in Study 1 we also examined a model in which the stereotypicality of trials was

modeled continuously rather than with the dichotomous binning approach. Specifically, this

additional model included two regressors – one for trials including a woman's face and another

for trials including a man's face. We included a separate parametric modulator for each regressor

to model the extent of the stereotypicality of the statement included in that trial based on our

pilot ratings. For example, the statement "Can lift heavy things" was rated as related more to

men than to women in our pilot studies with a 28-points difference on the 0–100 scale.

Consequently, the parametric modulation value was 0.28 for trials in which this statement was

presented with a man's face, and -0.28 for trials in which this statement was presented with a

woman's face. Similar to the models reported in the main text, in this additional model we

convolved events with a canonical hemodynamic response function and its temporal derivative

and included additional covariates of no interest (session mean, no response trials, six motion

parameters, and their temporal derivative).

Regions of interest (ROIs)

We used two complementary approaches to localize regions involved in the processing of

rewards. For independently defined ROIs, we defined 8-mm spheres around peak coordinates

drawn from a recent meta-analysis (15). Specifically, we utilized the peaks of the region

identified as supporting the processing of both monetary and primary incentives: bilateral ventral

striatum (x=-6, y=10, z=-6 and x=10, y=12, z=-6).

To functionally locate these regions, we examined the MID task to identify voxels that

responded more to rewarded trials (i.e., trials in which participants successfully responded to the

target) than to no-reward trials (i.e., trials in which no reward was available). Whole-brain

corrected clusters (using the procedure described in the main text) were defined as independent

ROIs.

As we had no prediction about laterality of the hypothesized effects, we collapsed across

hemispheres to create a single ROI. We extracted and averaged parameter estimates across

voxels and analyzed them with planned contrasts in a repeated-measures ANOVA context using

$p < 0.05$ as a threshold.

Studies 2 and 4. In each trial, participants chose one of two decks of cards (see Fig S2). We

instructed participants to choose based on their preferences regarding the information they had

available for each trial. Participants had two sources of information to rely on. First, each deck

was associated with a small monetary payoff (ranging from $0.03 to $0.09 in 2 cents

increments). Participants were told that a subset of their choices (5 trials in Study 2, 7 trials in

Study 4) would be added to their final compensation for the study. Payoff amounts for each

choice varied across trials (and were occasionally equal). Second, each deck was associated with

a specific label ("Typical" versus "Atypical" in Study 2, "Common" versus "Uncommon" in

Study 4) that determined the content presented when that deck is selected (stereotypical or

counter-stereotypical targets in Study 2, knowledge-consistent or knowledge-violating targets in

Study 4). After making their selection, participants saw a face-statement pair corresponding to

their choice. For example, if a participant in Study 2 selected the "Typical" deck, they would be

presented with a stereotypical target (e.g., in Study 2a a male associated with the statement

"CEO of a big company"). Participants then indicated how likely that target was to possess that

characteristic on a visual scale of 0 (= "Not at all likely") to 100 (= "Very likely"; the scale had

no other tick marks). The location of the specific labels (right deck versus left deck) was

counterbalanced between participants. Study 2a included only faces of men, Study 2b included

only faces of women, Study 4a included only the face of Barack Obama and Study 4b included

only the face of Donald Trump.

All studies included a demo trial, 4 practice trials, and 25 deck selection trials (32 trials in

Study 4). We also included several catch trials to detect if participants were responding without

considering the statement presented. The catch trials prompted the participants to slide the bar to

the right tick mark or to the left tick mark. Participants had up to 4s to select a card and up to 5s

to rate an individual. Participants that did not respond fast enough to more than 20% of the trials

got their survey terminated midway through the task. The specific amounts associated with each

trial and the face-statement pairs presented per trial were randomized. Following these trials,

participants were presented with 4 final manipulation-check trials in which they were asked to

select the card with the higher value.

After completing the main task, participants responded to the individual difference scales

mentioned above – ASI, SDO, MCP, and NFC. Order of the scales was randomized between

participants. No significant correlations between Point of Subjective Equivalence (PSE) and the

individual scores on these scales were consistently detected across studies. Finally, participants

supplied demographic information (including age, self-reported gender identity, self-reported

race with multiple answers enabled and whether they were born in the US). Study 4 also probed

participants' political affiliation (as in Study 3), how much they liked Barack Obama and how

much they liked Donald Trump on two separate 0–100 scales. Then, we probed for participants'

intuitions about the goal of the task and fully debriefed them.

SI Results

Behavioral analyses: Neuroimaging studies

Table S1 presents the summaries of participants' ratings and reaction times in the impression

formation task in Study 1. We analyzed rating data with mixed models as implemented in the

lme4 package version 1.1-14 (16) and the 'ordinal' package version 2018.4-19 (17) for R version

3.4.2 (R Core Team, 2017). As the behavioral data obtained in the impression formation task

were ordinal, we analyzed them using cumulative link mixed models (CLMM) with the logit

link. To avoid the transformation of raw reaction time data, we used generalized linear models

(gLMMs) with the inverse Gaussian identity link (18). We included random effects for the

intercepts for participants and statements, as well as by-participant random slopes for the fixed

effect of stereotypicality. Trials that elicited no response (<1.5% of all trials; no difference

between conditions) were excluded from all analyses.

To examine whether our stereotype-consistent and stereotype-violating targets were

indeed perceived differently by our participants in Study 1, we tested the effect of our a-priori

categorization on behavioral ratings in a CLMM. Stereotypicality was dummy coded with

stereotype-neutral trials as the intercept and behavioral ratings were centered on 0. Overall

stereotypicality affected ratings, as indicated by leave-one-out model comparison, comparing our

model to an intercept only model ($\chi^2_{(2)} = 44.45$, $p < 0.001$). The manipulation worked as

anticipated: Stereotype-consistent targets received higher ratings than stereotype-neutral targets

($\beta \pm SE = 0.42 \pm 0.14$, $Z = 2.99$, $p = 0.003$), whereas stereotype-violating targets received lower

ratings ($\beta \pm SE = -1.22 \pm 0.15$, $Z = -8.31$, $p < 0.001$).

Stereotypicality did not have a main effect on reaction time for the behavioral ratings

($\chi^2_{(2)} = 0.35$, $p > 0.8$). However, reaction time did vary by the interaction of stereotypicality and

behavioral ratings, as indicated by comparing a model with an interaction term to a model

without it, $\chi^2_{(2)} = 133.88$, $p < 0.001$); participants were slower to provide responses that were not

in line with our a-priori definitions (e.g., indicating that a woman is very likely to be a firefighter

or saying that a man is very unlikely to be a CEO); see Table S1 for full descriptive results.

Behavioral results in Study 3 generally replicated the behavioral results we obtained in

Study 1 (see Table S3). The outcome of the expectation (whether the target were consistent with

or violated the expectation) significantly affected participants' ratings ($\chi^2_{(1)} = 73.61$, $p < 0.001$).

Domain (stereotype-based or person-specific-based knowledge) also affected participants'

ratings ($\chi^2_{(1)} = 27.58$, $p < 0.001$). Interestingly, participants distributed their expectation-based

responses differently between the domains (outcome by domain interaction: $\chi^2_{(1)} = 862.52$, $p <$

0.001). Participants used more extreme outcome-congruent ratings for person-specific

expectations compared to stereotype-based expectations (see Table S3).

Similar to Study 1, participants' reaction time did not differ between expectation-

consistent or expectation-violating trials ($t = -0.29$, $p = 0.77$), nor between person-specific or

stereotype-based expectations ($t = 1.89$, $p = 0.059$). Comparable to Study 1, we observed a

significant interaction between outcome and behavioral ratings ($t = -15.11$, $p < 0.001$), an

interaction that was qualified by a 3-way interaction with domain ($t = 6.36$, $p < 0.001$). To

interpret this interaction, we analyzed responses separately for the two knowledge domains. In

both knowledge domains, participants responded faster when their responses were in line with

our a-priori definitions ($t = -9.72$ and $t = -13.81$ for stereotype- and person-specific-based

expectations, respectively; $p's < 0.001$).

Neuroimaging

In Study 3, in addition to the main findings (see Fig 1 and Table S4), the design allowed us to

compare the effects of expectation-consistency with different specific targets, namely, men,

women, Trump or Obama. Analyzing activity in the NAcc in a 2 (expectation result) X 2

(content domain) X 2 (specific target) ANOVA yielded, in addition to the main effect of

expectation consistency ($F_{(1,29)} = 21.41$, $p < 0.0001$, $\eta^2_p = 0.42$ [0.19–0.58], a main effect of

content domain ($F_{(1,29)} = 4.56$, $p = 0.04$, $\eta^2_p = 0.14$ [0.003–0.32]; all other effects p>0.1) such that

gender-related trials were associated with increased activity. To complement the findings, we

conducted a whole-brain interaction analysis to examine whether any neural regions responded

differently to the confirmation or violation of expectations between specific targets. Given the

different experimental context between the two content domains, we first examined the

interaction of expectations and specific targets within each content domain separately. This

analysis yielded 3 regions (p<0.05, FWE-corrected; Table S5 and Figure S3), including the left

inferior frontal gyrus (LIFG) and left superior temporal gyrus (LSTG), in which consistency with

expectations yielded more activity than their violation only when the statements pertained to

Trump (LIFG results: $F_{(1,114.24)} = 40.12$, $p < 0.0001$, $\eta^2_p = 0.26$ [0.15–0.36]) and not to any of the

other targets (triple interaction: $F_{(1,29)} = 33.32$, $p < 0.0001$, $\eta^2_p = 0.53$ [0.30–0.66; similar patterns

were observed in LSTG]. Disambiguating this triple interaction, we verified the interaction

between expectation results and specific targets for leaders: $F_{(1,29)} = 42.73$, $p < 0.0001$, $\eta^2_p = 0.6$

[0.37–0.71]; the parallel interaction within gender content was not significant: $F_{(1,29)} = 0.55$, $p >$

0.4, $\eta^2_p = 0.02$ [0–0.03]). Notably, no main effect of expectation results was observed in these

regions (all *p's* > 0.05). Together, these findings suggest that the consistency with expectations

pertaining to Donald Trump evokes an additional process which is not triggered for confirmation

of other types of expectations.

Behavioral analysis: Online studies

In addition to the analyses reported in the main manuscript, in Study 4 we also performed an

exploratory pre-registered analysis to examine the relationship between PSEs and support for the

specific leader. To that end, we collected from each participant how much they liked Barack

Obama and how much they liked Donald Trump on two separate 0–100 scales. Unsurprisingly

for online data collection, our participants demonstrated a skewed preference toward Barack

Obama (mean Obama liking for participants in Study 4a: 71.89, *S.D.* = 28.68; mean Trump

liking for participants in Study 4b: 17.1, *S.D.* = 27.33). We correlated these data with PSEs for

the corresponding leader in each study. As can be seen in Figure S4, the data are extremely

skewed, with many participants indicating the maximum liking rating for Obama and the

minimum liking rating for Trump. As such, we do not provide inferential statistics as these data

violate the statistical assumptions. Descriptively, we can see a small positive slope between PSE

and liking for each leader. These descriptive trends can be taken to cautiously hint that the more

one likes a leader, the more they would be willing to forgo money to see information consistent

with the expectation from that leader.

1   **SI References**

3   1.   W. A. Bainbridge, P. Isola, A. Oliva, The intrinsic memorability of face photographs. *J. Exp.*

4   *Psychol. Gen.* **142**, 1323–1334 (2013).

5   2.   D. A. Prentice, E. Carranza, What women and men should be, shouldn't be, are allowed to be, and

6   don't have to be: the contents of prescriptive gender stereotypes. *Psychol. Women Q.* **26**, 269–281

7   (2002).

8   3.   A. M. Dale, Optimal experimental design for event-related fMRI. *Hum. Brain Mapp.* **8**, 109–114

9   (1999).

10   4.   J. A. Mumford, T. Davis, R. A. Poldrack, The impact of study design on pattern estimation for

11   single-trial multivariate pattern analysis. *Neuroimage* **103**, 130–138 (2014).

12   5.   B. Knutson, A. Westdorp, E. Kaiser, D. Hommer, FMRI visualization of brain activity during a

13   monetary incentive delay task. *Neuroimage* **12**, 20–27 (2000).

14   6.   P. Glick, S. T. Fiske, The Ambivalent Sexism Inventory: Differentiating hostile and benevolent

15   sexism. *J. Pers. Soc. Psychol.* **70**, 491–512 (1996).

16   7.   A. K. Ho, *et al.*, The nature of social dominance orientation: Theorizing and measuring

17   preferences for intergroup inequality using the new SDO☐ scale. *J. Pers. Soc. Psychol.* **109**,

18   1003–1028 (2015).

19   8.   S. C. Klonis, E. A. Plant, P. G. Devine, Internal and external motivation to respond without

20   sexism. *Personal. Soc. Psychol. Bull.* **31**, 1237–1249 (2005).

21   9.   A. W. Kruglanski, D. M. Webster, A. Klem, Motivated resistance and openness to persuasion in

22   the presence or absence of prior information. *J. Pers. Soc. Psychol.* **65**, 861–876 (1993).

1   10.   J. W. Peirce, PsychoPy-Psychophysics software in Python. *J. Neurosci. Methods* **162**, 8–13

2   (2007).

3   11.   R. Cusack, N. Papadakis, New robust 3-D phase unwrapping algorithms: Application to magnetic

4   field mapping and undistorting echoplanar images. *Neuroimage* **16**, 754–764 (2002).

5   12.   S. Moeller, *et al.*, Multiband multislice GE-EPI at 7 tesla, with 16-fold acceleration using partial

6   parallel imaging with application to high spatial and temporal whole-brain fMRI. *Magn. Reson.*

7   *Med.* **63**, 1144–1153 (2010).

8   13.   D. A. Feinberg, *et al.*, Multiplexed echo planar imaging for sub-second whole brain FMRI and fast

9   diffusion imaging. *PLoS One* **5**, e15710 (2010).

14.   J. Xu, *et al.*, Evaluation of slice accelerations using multiband echo planar imaging at 3 T.

Neuroimage **83**, 991–1001 (2013).

15.   O. Bartra, J. T. McGuire, J. W. Kable, The valuation system: A coordinate-based meta-analysis of

BOLD fMRI experiments examining neural correlates of subjective value. *Neuroimage* **76**, 412–

427 (2013).

16.   D. M. Bates, M. Maechler, B. Bolker, S. Walker, lme4: Linear mixed-effects models using Eigen

and S4 (2014).

17.   R. H. B. Christensen, ordinal---Regression Models for Ordinal Data (2018).

18.   S. Lo, S. Andrews, To transform or not to transform□: Using Generalized Linear Mixed Models to

analyse reaction time data. *Front. Psychol.* **6**, 1–16 (2015).

2   **Fig S1.**

Experimental designs of Studies 1 and 3. (A) In Study 1, participants saw 204 unique trials. Each

started with a gender-relevant or irrelevant statement, followed by a target face consistent with,

violating or neutral in respect to the displayed statement. Participants indicated how likely the

presented person was to have the characteristic described in the statement on a 1 ("very

unlikely") to 4 ("very likely") scale. (B) Following the impression formation task, participants

completed the monetary incentive delay (MID) task. In each trial, participants saw a cue

predicting the outcome of a successful response to the target. A green cue always indicated

monetary reward, a blue cue always indicated no reward. Participants saw 30 reward cues and 15

no-reward cues. After a randomly jittered delay a target appeared on screen for a brief duration

(determined by a 2-up-1-down staircase procedure). Participants received feedback about their

performance in each trial and across the entire task. (C) In Study 3, participants rated 240 trials,

including 120 stereotype-related targets and 120 person-specific trials. Each content domain

(stereotypes versus person-specific) was presented in separate blocks of 15 consecutive trials per

type.

3   **Fig S2.** Experimental design of Study 2a. Participants chose which target type to rate (typical,

4   denoting stereotype-confirming targets, versus atypical, denoting stereotype-violating targets).

5   Each target type was associated with a variable amount of money. After deciding which target

6   type to rate, participants then rated a target of the chosen type. The design of studies 2b, 4a and

7   4b followed this procedure.

2
3   **Fig S3.** Neural responses as a function of expectancy-outcome and type of target. (A) results of a

4   whole-brain analysis examining the interaction of outcome and type of target (see main text and

5   Table S5 for details). (B) Parameter estimates drawn from the Left Inferior Frontal Gyrus ROI.

Fig S4. The correlations between the subjective value of expectation-consistent information and the liking

of a specific leader in Study 4. The subjective value was calculated as the point of subjective equivalence

(PSE) per participant, as detailed in the main text. (a) Results from Study 4a in which participants chose

between seeing trials with expectation-consistent and expectation-inconsistent information about Barack

Obama. (b) Results from Study 4b in which participants made equivalent choices about trials including

Donald Trump. We do not provide inferential statistics as these data are heavily skewed.

1   *Table S1.* Distribution of participants' ratings and reaction time (RT, in milliseconds) as a

2   function of stereotypicality in the impression formation task in Study 1. Parentheses indicate

3   standard error of the mean.

	Very Likely	Somewhat Likely	Somewhat Unlikely	Very Unlikely	No Response
Stereotype-confirming (68)					
Proportion	0.27 (0.03)	0.39 (0.03)	0.23 (0.02)	0.09 (0.01)	0.01 (0.005)
RT	1676 (28)	1845 (27)	2100 (30)	1751 (54)	
Stereotype-neutral (68)					
Proportion	0.18 (0.02)	0.42 (0.03)	0.26 (0.0.2)	0.12 (0.02)	0.01 (0.004)
RT	1676 (34)	1855 (31)	2047 (35)	1814 (69)	
Stereotype-violating (68)					
Proportion	0.07 (0.01)	0.27 (0.02)	0.36 (0.02)	0.29 (0.02)	0.01 (0.005)
RT	1828 (77)	1946 (26)	2015 (31)	1640 (32)	

1 *Table S2.* Study 1: Gray matter regions showing differences in activity between stereotypical

2 (stereotype-consistent) and counter-stereotypical (stereotype-violating) targets in model 1 (with

3 no specification of behavioral response; p<0.05, corrected).

Region	MNI coordinates			Z value	# voxels
	X	Y	Z		
(a) Stereotypical > Counter-Stereotypical					
Nucleus Accumbens	-2	4	-6	4.98	30
L Inferior Temporal	-50	-56	-12	4.03	108
(b) Counter-Stereotypical > Stereotypical					
R Motor	34	-22	58	4.38	145
R Inferior Frontal	46	32	-10	4.07	18
Dorsomedial Prefrontal	8	36	50	3.8	50
(c) Parametric Modulation (Counter-Stereotypical -> Stereotypical)					
L inferior Parietal Sulcus	-30	-82	36	4.22	17
L Inferior Temporal	-48	-58	-12	4.19	60
Extrastriate Cortex	14	-84	-12	3.72	22
Nucleus Accumbens	0	6	-6	4.15	20
R Nucleus Accumbens	-8	8	-4	3.91	13

(d) Parametric Modulation (Stereotypical -> Counter-Stereotypical)

R Motor	34	-20	58	4.41	128
R Medial prefrontal	12	48	12	4.33	18
R Inferior Parietal	58	-50	38	3.94	18
Dorsomedial Prefrontal	8	36	50	3.85	16

1 *Table S3.* Distribution of participants' ratings and reaction time (RT, in milliseconds) as a

2 function of expectation domain and expectation-consistency in the impression formation task in

3 Study 3. Parentheses indicate standard error of the mean.

		Very Likely	Somewhat Likely	Somewhat Unlikely	Very Unlikely	No Response
Stereotype	**Confirming (60)**					
	Proportion	0.30 (0.03)	0.38 (0.03)	0.21 (0.02)	0.09 (0.01)	0.02 (0.01)
	RT	1635 (33)	1805 (27)	1957 (34)	1703 (46)	
	Violating (60)					
	Proportion	0.08 (0.01)	0.27 (0.02)	0.33 (0.03)	0.30 (0.03)	0.02 (0.01)
	RT	1770 (49)	1892 (38)	1954 (30)	1583 (34)	
Person-specific	**Confirming (60)**					
	Proportion	0.73 (0.03)	0.18 (0.02)	0.05 (0.01)	0.03 (0.01)	0.02 (0.01)
	RT	1297 (40)	1793 (42)	2080 (79)	1694 (118)	
	Violating (60)					
	Proportion	0.06 (0.01)	0.13 (0.01)	0.28 (0.03)	0.51 (0.03)	0.02 (0.01)
	RT	1459 (60)	1854 (45)	1925 (41)	1437 (36)	

1 *Table S4.* Study 3: Gray matter regions showing differences in activity between expectation-

2 consistent and expectation-violating targets. (a) and (b) portray the results collapsed across

3 content domain, (c) through (f) include the results per content domain (p<0.05, corrected).

Region	MNI coordinates			Z value	# voxels
	X	Y	Z		
(a) Main effect: Confirming > Violating					
Extrastriate Cortex	12	-80	-4	5.63	2800
R Motor	46	-14	52	5.43	298
L Ventral Striatum	-12	20	-2	4.93	89
L Insula	-48	4	8	4.87	59
Cerebellum	10	-74	-38	4.82	56
Cerebellum	-18	-76	-44	4.58	35
L Occipitotemporal Cortex	-44	-68	0	4.57	463
R Ventral Striatum	6	6	-8	4.52	49
R Insula	36	6	10	4.51	40
L Parietal Operculum	-44	-32	20	4.34	166
Occipital Cortex	24	-96	20	4.29	266
Posterior Cingulate	-2	-34	44	4.26	240
R Fusiform Gyrus	48	-42	-16	4.25	57

R Occipitotemporal Cortex	54	-60	0	4.19	221
L Superior Temporal	-66	-36	20	4.08	62
L Insula	-36	-12	14	4.06	42
L Parahippocampus	-24	-40	-6	4.06	50
L Inferior Frontal	-42	38	12	4.01	34
Precuneus	-16	-62	16	3.98	103
Ventromedial Prefrontal	6	42	-16	3.86	28
L Motor	-50	-16	54	3.77	112
L Temporal Pole	-48	14	-14	3.72	20
R Orbitofrontal	26	38	-18	3.69	26
R Inferior Frontal	52	8	-4	3.64	39

(b) Main effect: Violating > Confirming

Dorsomedial Prefrontal	12	34	52	4.22	23
R Motor	36	-24	54	3.99	50
Dorsomedial Prefrontal	-8	42	50	3.92	62
L Middle Frontal	-38	20	46	3.88	32

(c) Gender: Confirming > Violating

Extrastriate Cortex	10	-78	-6	4.48	146
R Occipitotemporal Cortex	54	-62	2	3.80	47
Ventral Striatum	4	10	-10	3.78	21
L Motor	52	-12	54	3.66	64
R Lateral Occipital	30	-84	38	3.55	28

(d) Gender: Violating > Confirming

Dorsomedial Prefrontal	12	34	52	3.92	33
L Middle Frontal	-34	20	40	3.55	19

(e) Leaders: Confirming > Violating

Extrastriate Cortex	8	-76	-6	5.56	1854
R Motor	48	-14	52	5.42	204
Posterior Cingualte	-8	-40	54	4.34	57
L Parietal Operculum	-42	-36	26	4.31	248
R Insula	36	6	10	4.23	24
L Temporal Pole	-50	12	-12	4.19	80
Cerebellum	24	-66	-44	4.18	24
R Supramarginal gyrus	50	-36	8	4.06	50

L Inferior Frontal / Insula	-48	4	6	4.01	35
L Occipitotemporal Cortex	-44	-68	0	3.89	21
Cuneus	-2	-94	16	3.83	21
R Fusiform Gyrus	48	-44	-16	3.78	30
Cerebellum	8	-74	-38	3.76	20
L Occipitotemporal Cortex	-48	-58	-2	3.71	111
Precuneus	-14	-66	28	3.71	55
L Inferior Frontal	-60	6	10	3.68	22
Precuneus	-14	-78	42	3.62	74
Ventral Striatum	-10	20	-2	3.57	21
Anterior Cingualte	-2	14	32	3.49	31
R Inferior Frontal / Insula	50	4	-2	3.48	22
L Motor	-50	-16	54	3.45	22

(f) Leaders: Violating > Confirming

R Motor	40	-24	60	4.17	90
Dorsomedial Prefrontal	-8	42	50	3.77	22

1 *Table S5.* Study 3: Results of a set of two whole-brain analyses examining the interactive effects

2 of expectation-consistency and specific target, done separately for gender and leaders. (p<0.05,

3 corrected).

Region	MNI coordinates			Z value	# voxels
	X	Y	Z		
(a) Gender: Confirmation effect in men > Confirmation effect in women					
None					
(b) Gender: Confirmation effect in women > Confirmation effect in men					
None					
(c) Leaders: Confirmation effect in Obama > Confirmation effect in Trump					
None					
(d) Leaders: Confirmation effect in Trump > Confirmation effect in Obama					
Cuneus	12	-88	24	5.16	435
L Inferior Frontal	-50	34	2	4.83	473
L Superior Temporal	-62	-36	4	4.32	352
Extrastriate Cortex	-6	-80	6	3.67	75