Title page

Once upon a time in the south: local drivers of plant invasion in the harsh sub-Antarctic islands

Bazzichetto[1]*, M., Massol[2], F., Carboni[3], M., Lenoir[4], J., Lembrechts[5], J. J., Joly[1], R. & Renault[1,6], D.

[1] Université de Rennes, CNRS, EcoBio (Ecosystèmes, biodiversité, évolution) - UMR 6553, F-35000 Rennes, France

[2] Université de Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 - UMR 9017 - CIIL - Center for Infection and Immunity of Lille, F-59000 Lille, France

[3] Department of Sciences, University Roma Tre, Rome, Italy

[4] UR 'Ecologie et Dynamique des Systèmes Anthropisés' (EDYSAN, UMR 7058 CNRS-UPJV), Université de Picardie Jules Verne, Amiens, France

[5] Research Group PLECO (Plants and Ecosystems), University of Antwerp, 2610 Wilrijk, Belgium

[6] Institut Universitaire de France, 1 rue Descartes, 75231 Paris Cedex 05, France

*corresponding author

Authors information:

Massol François: Univ. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 - UMR 9017 - CIIL - Center for Infection and Immunity of Lille, F-59000 Lille, France

francois.massol@univ-lille.fr

ORCID: https://orcid.org/0000-0002-4098-955X

Carboni Marta: Department of Sciences, University Roma Tre, Rome, Italy

marta.carboni@uniroma3.it

ORCID: https://orcid.org/0000-0002-9348-4758

Lenoir Jonathan: UR 'Ecologie et Dynamique des Systèmes Anthropisés' (EDYSAN, UMR 7058 CNRS-UPJV), Université de Picardie Jules Verne, Amiens, France

jonathan.lenoir@u-picardie.fr

ORCID: https://orcid.org/0000-0003-0638-9582

Lembrechts Jonas Johan: Research Group PLECO (Plants and Ecosystems), University of Antwerp, 2610 Wilrijk, Belgium

jonas.lembrechts@uantwerpen.be

ORCID: https://orcid.org/0000-0002-1933-0750

Joly Rémi: Université de Rennes, CNRS, EcoBio (Ecosystèmes, biodiversité, évolution) - UMR 6553, F-35000 Rennes, France

remi.joly88@hotmail.fr

Renault David: Université de Rennes, CNRS, EcoBio (Ecosystèmes, biodiversité, évolution) - UMR 6553, F-35000 Rennes, France; Institut Universitaire de France (IUF), 1 rue Descartes, 75231 Paris Cedex 05, France

david.renault@univ-rennes1.fr

ORCID: https://orcid.org/0000-0003-3644-1759

Bazzichetto Manuele (corresponding author): Université de Rennes, CNRS, EcoBio (Ecosystèmes, biodiversité, évolution) - UMR 6553, F-35000 Rennes, France

manuele.bazzichetto@univ-rennes1.fr

ORCID: https://orcid.org/0000-0002-9874-5064

Once upon a time in the south: local drivers of plant invasion in the harsh sub-Antarctic islands

Manuele Bazzichetto[1*], François Massol[2], Marta Carboni[3], Jonathan Lenoir[4], Jonas J Lembrechts[5], Rémi Joly[1], & David Renault[1,6]

[1] Université de Rennes, CNRS, EcoBio (Ecosystèmes, biodiversité, évolution) - UMR 6553, F-35000 Rennes, France

[2] Université de Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 - UMR 9017 - CIIL - Center for Infection and Immunity of Lille, F-59000 Lille, France

[3] Department of Sciences, University Roma Tre, Rome, Italy

[4] UR 'Ecologie et Dynamique des Systèmes Anthropisés' (EDYSAN, UMR 7058 CNRS-UPJV), Université de Picardie Jules Verne, Amiens, France

[5] Research Group PLECO (Plants and Ecosystems), University of Antwerp, 2610 Wilrijk, Belgium

[6] Institut Universitaire de France, 1 rue Descartes, 75231 Paris Cedex 05, France

*corresponding author

Abstract

Aim

To investigate the local effect of environmental and human-related factors on alien plant invasion in
sub-Antarctic islands. To explore the relationship between alien species features and dependence on
anthropogenic propagule pressure to unravel key traits conferring invasiveness in the sub-Antarctic.

Location

Possession Island, Crozet archipelago (French sub-Antarctic islands).

Taxon

Non-native vascular plants (Poaceae, Caryophyllaceae, Juncaceae).

Methods

Single-species distribution models were used to explore the effect of high-resolution topoclimatic and
human-related variables on the occurrence of six of the most aggressive alien plants colonizing
French sub-Antarctic islands. Furthermore, the interaction between alien species traits and their
response to anthropogenic propagule pressure was analysed by means of a multi-species distribution
model. This allowed identifying the features of species that were associated to low dependence on
human-assisted introductions, and were thus potentially more invasive.

Results

We observed two main invasion patterns: low-spread species strongly dependent on anthropogenic
propagule pressure and high-spread species limited mainly by harsh climatic conditions. Differences
in invasiveness across species mostly related to their residence time, life history and plant height,
with older introductions, perennial and low-stature species being most invasive.

Main conclusions

The availability of high-resolution data allowed for a fine understanding of the role of environmental and human-related factors in driving alien species distribution on sub-Antarctic islands. At the same time, the identification of alien species features conferring invasiveness may help anticipating future problematic invasions.

Keywords: alien plants, anthropogenic propagule pressure, invasiveness, plant invasion, species distribution models, sub-Antarctic islands, topoclimate

Running title: plant invasion in sub-Antarctic islands

1. Introduction

Sub-Antarctic islands are extremely remote territories which harbour a unique biodiversity with a high degree of endemism (Shaw, 2013). As a consequence of their relatively recent discovery and environmental harshness, these islands have long remained pristine and largely free of human disturbances. Yet, due to the gradual relaxation of these natural barriers, sub-Antarctic islands are now counted among the most threatened environments on Earth. In particular, invasion by alien plants, boosted by ongoing climate changes and increasing human disturbances (Duffy & Lee, 2019; Hughes et al., 2019), has become one of the main threats to the endemic biodiversity of these territories, and is bound to rise in the next decades (Lebouvier et al., 2011; Hughes, Pertierra, Molina-Montenegro, & Convey, 2015). Over the past century, alien plants have been increasingly introduced in the sub-Antarctic region (Frenot et al., 2005; Huiskes et al., 2014). European whalers and scientific activities, respectively in the 19th and 20th century, determined the first main introduction events (Convey & Lebouvier, 2009; Shaw, 2013), while, since the mid-twentieth century, climate modifications have progressively made sub-Antarctic islands more suitable to cold-tolerant alien species (Shaw, 2013; Pertierra et al., 2017; Duffy & Lee, 2019). Nevertheless, despite their demonstrated impacts on native biodiversity, little attention has been given to plant compared to animal invasions in these islands (Molina-Montenegro et al., 2012; le Roux et al., 2013), leaving a knowledge gap in the mechanisms underpinning plant invasion processes in these unique environments (Greve, Mathakutha, Steyn, & Chown, 2017).

The outcome of any biological invasion is jointly determined by propagule pressure (i.e. frequency of propagules introduction), abiotic conditions (i.e. physico-chemical features of the invaded environment) and biotic features (i.e. alien species characteristics and interactions with the recipient community), with anthropogenic disturbances affecting all three (Richardson & Pyšek, 2006; Catford, Jansson, & Nilsson, 2009; Lembrechts et al., 2016). The relative importance of these factors is, however, context-dependent and species-specific (Catford et al., 2009). In sub-Antarctic islands, due to the high specialization but low diversity of the native flora, biotic interactions are thought to play a minor role (le Roux et al., 2013; Duffy et al., 2017; Moser et al., 2018), so it is mainly the first two factors that determine the distribution and spread of alien plants. First, invasions depend on human-induced propagule pressure: the frequency of propagule introduction correlates with the number of ship landings and is highest in the vicinity of human facilities (Huiskes et al., 2014). Second, local abiotic conditions are strongly limiting, and particularly the climatic mismatch between the alien species' native range and the sub-Antarctic can strongly constrain invasions (Frenot et al., 2005). Some alien plants are more limited during the introduction phase, while others quickly become relatively independent of human-related propagule-pressure and are only climatically limited. Once, established, the species which are least dependent on continuous introductions are the most likely to spread widely and become invasive (Richardson & Pyšek, 2006; Catford et al., 2009). Therefore, quantifying the degree of alien species dependence on propagule pressure might aid in identifying potentially invasive species.

A lower dependence on human-related propagule pressure is potentially related to certain species features which are more generally known to affect invasiveness. First of all, alien species with longer residence times are more likely to become independent of anthropogenic propagule pressure (Wilson et al., 2007; Pyšek et al., 2015). Second, certain plant traits are considered key for profiling successful invaders (Pyšek & Richardson, 2008): invasive alien plants across most environments are growing faster and taller than non-invasive alien species, and typically produce resource-acquisitive leaves and many small seeds (van Kleunen, Weber, & Fischer, 2010; van Kleunen, Dawson, & Maurel, 2015). More specifically, Mathakutha et al. (2019) performed a first functional comparison between

invasive and non-invasive alien species colonizing the sub-Antarctic Marion Island, reporting that
species generally considered invasive had lower plant height, smaller leaf area, lower frost tolerance
and higher specific leaf area than other alien species. Nevertheless, it is still unclear which traits can
actually make some alien plants less dependent on human-related propagule pressure, and thus more
likely to become invasive, especially in the sub-Antarctic islands. This knowledge could facilitate the
early screening of highly invasive alien species in these environments (Frenot et al., 2005;
Mathakutha et al., 2019).

Correlative species distribution models (SDMs) are statistical tools that model the species-
environment relationship relying on geo-referenced occurrence data and spatial environmental layers
(Guisan, Thuiller, & Zimmermann, 2017). SDMs already proved to be valuable tools for analysing
alien plant invasion in Antarctica and the sub-Antarctic regions. For instance, Pertierra et al. (2017)
modelled the distribution of *Poa annua* and *Poa pratensis* in the Antarctic peninsula as a function of
bioclimatic variables, while Duffy et al. (2017) generated future scenarios of invasion across
Antarctica and the sub-Antarctic regions using climate-based SDMs. Whilst these previous SDM
applications have revealed large-scale determinants of alien plant invasion in the Antarctic
biogeographic region, they have up till now failed to account for how environmental and
anthropogenic factors regulate alien plant distributions at the fine spatial resolutions meaningful for
local management. This is chiefly due to the lack of high-resolution environmental (e.g. climatic,
topographic) and human-related data layers, which limits the implementation of SDMs at fine spatial
resolutions in remote areas (Gutt et al., 2012).

The sub-Antarctic Possession Island constitutes an ideal arena to analyse alien plant invasions in the
sub-Antarctic region. The availability of historical vegetation observations allows retracing the
invasion history of most alien species. Moreover, this island witnessed past human colonization and
climate changes comparable to the other sub-Antarctic islands, allowing inference on the mechanisms
underpinning alien plant invasion in these unique areas. Previous work showed that there is
considerable variation in the spread of alien plants established on Possession Island, with some
species clustering close to their introduction locations and others spreading widely and far from the
initial introduction sites (Frenot et al., 2005), which allows testing for differences in the dependence
on human introductions. In the present study, we model the distribution of the most relevant alien
plant species colonizing Possession Island using a combination of environmental and human-related
spatial data derived at an unprecedented high spatial resolution (i.e. 30-m) for these latitudes, and
long-term monitoring observations of plant occurrences. Our aim is to test the local effect of
environmental and anthropogenic factors on alien plant invasion in sub-Antarctic ecosystems. We
hypothesize that both abiotic and human-related factors jointly define the local occurrence of alien
plant species, but that these two factors will not be equally important among species. Furthermore, to
identify plant characteristics conferring high invasiveness in sub-Antarctic ecosystems, we
investigate how plant functional traits affect species dependence on anthropogenic propagule
pressures. In this regard, our working hypothesis is that the more invasive species share specific
functional characteristics allowing them to become independent of human-assisted introductions and
spread widely once established.

2. Materials and methods

2.1 *Study area*

The study was carried out on Possession Island in the Crozet archipelago, which is included in the *Réserve Naturelle Terres Australes Françaises* (RN-TAF) and listed as UNESCO World Heritage site since 2019. Possession Island (Longitude: 51.7469, Latitude: -46.4046; EPSG: 4326) is characterized by a complex topography, with an altitudinal gradient ranging from 0 to 934 m above the sea level (*Pic du Mascarin*) over a relatively short spatial extent (147 km^2). The island is characterized by a typical sub-Antarctic climate, with mean annual temperature of 5.6 °C and annual precipitation of 2300 mm (Météo France, data: 1960-2019). Frequent and strong western winds occur throughout most of the year.

The first human settlements date back to the 19[th] century, when whalers and sealers established on the north-east side of the island during the hunting season, facilitating a first series of alien species introductions. In 1963, a permanent research station (*Alfred Faure*, hereafter the 'research station') was built on the easternmost area of Possession Island, fostering a new invasion front. Beyond the research station, other shelters (inhabited for short periods) are currently present on each side of the island: north (*Pointe Basse*); south-west (*Baie du La Pérouse*); and north-east (*Baie Américaine*). Among these, the research station is by far the biggest human settlement and main hub of propagule introduction. The vegetation at Possession Island has experienced relatively low grazing pressure from large herbivores in the past (Convey & Lebouvier, 2009), in comparison to other sub-Antarctic islands where introduced large herbivores still strongly affect the distribution of alien plants (Shaw, 2013).

2.2 *Study species*

Despite the 68 alien species recorded on Possession Island (RN-TAF management plan 2018-2027: https://taaf.fr//content/uploads/sites/2/2019/09/180607-Volet-A_pour-CNPN.pdf), only few have established persistent populations (Frenot, Gloaguen, Massé, & Lebouvier, 2001). In this study, we restricted our analysis to those alien plants that are either known to be generally widespread on sub-Antarctic islands or are particularly widespread on Possession Island, and for which sufficient occurrence data were available (total number of presences > 100). Specifically, we selected the following species: *Poa annua* and *Poa pratensis* (Poaceae); *Cerastium fontanum*, *Sagina procumbens* and *Stellaria alsine* (Caryophyllaceae); and *Juncus bufonius* (Juncaceae). The two grasses, *P. annua* and *P. pratensis,* have colonized most of the sub-Antarctic islands (Shaw, 2013), and are the longest-established alien plants in the Antarctic Peninsula (Pertierra et al., 2017). *Cerastium fontanum* and *S. procumbens* are currently widely distributed in this environment (Frenot et al., 2005; Shaw, 2013) with, in particular, *S. procumbens* exhibiting the highest rate of spread among the alien plants of Marion and Prince Edward Islands (le Roux et al., 2013). Finally, both *J. bufonius* and *S. alsine* currently occur at significant distances from the research station on Possession Island (Frenot el al., 2001). While the former has been recently observed up to the Maritime Antarctica latitudes (Cuba-Díaz, Fuentes, & Rondanelli-Reyes, 2015), the latter has been singled out by some authors as the potentially most problematic future invasive plant species on Possession Island (Frenot et al., 2001; Convey, Key, & Key, 2010).

2.2.1 *Species distribution data*

We analysed the invasion patterns of the six selected alien plant species relying on georeferenced occurrence (presence/absence) data collected within the context of a yearly vegetation monitoring survey carried out by the RN-TAF since 2010. The vegetation sampling is implemented within a system of 675 squared cells 100x100-m, where floristic data (presence and abundance of vascular plant species) are collected along with habitat characteristics through phytosociological *relevés*

165 (Dengler, 2016). In this study, we used data collected from 2010 to 2017 (3354 occurrences for the
166 selected species across 1572 sampled plot).

2.2.2 *Species features and functional trait data*

To inform species features potentially related to invasiveness, we collected data on plants residence
time and functional traits. Residence time positively interacts with propagule pressure in determining
plant invasion success (Richardson & Pyšek, 2006; Lockwood, Cassey, & Blackburn, 2005; Pyšek et
al., 2015), and this relationship was also observed on sub-Antarctic islands (le Roux et al., 2013;
Shaw, 2013; Mathakutha et al., 2019). To test how residence time influences alien species'
dependence on propagule pressure, we considered the introduction date of the selected plants on
Possession Island (Frenot et al., 2001) and used this information to assign them to two groups: old *vs*
new resident species (Appendix S1, Table S1.1). In particular, we considered as old resident species
those which were firstly observed on Possession Island before the research station was built (1963),
while referring to the others as new resident species.

We then collated data on seven plant traits commonly used to synthesize species strategies known to
be related to invasiveness (van Kleunen et al., 2010; van Kleunen et al., 2015): (1) life history (annual
vs perennial); (2) plant height; (3) leaf area; (4) specific leaf area (SLA); (5) vegetative reproduction
(present *vs* absent, i.e. sexual and vegetative *vs* only sexual reproduction); (6) seed dry mass; and (7)
number of seeds/plant. We excluded traits related to flowering since pollinating insects are absent
from almost all sub-Antarctic islands (Convey et al., 2010). Life history, plant height and leaf area
relate to plant persistence and tolerance to environmental stress (Cornelissen et al., 2003; Pérez-
Harguindeguy et al., 2013). In addition, life history is used to assess maximum lifespan and plant
height is associated with competitiveness for light and whole plant fecundity (Pérez-Harguindeguy et
al., 2013). Specific leaf area is the one-sided leaf area per leaf mass and is associated with resource
acquisition and photosynthetic rate (Pérez-Harguindeguy et al., 2013). Reproduction strategy, seed
dry mass and number of seeds per plant do not only relate with species persistence, but also with
dispersal capacity (Ottaviani et al., 2020). In particular, alien species reproducing predominantly
sexually may benefit from lower dispersal limitation and greater genetic diversity (van Kleunen et
al., 2015). At the same time, while small and light seeds are better dispersed at longer distances, large-
seeded plants may benefit from more stored resources (van Kleunen et al., 2015).

Functional trait data collected in areas environmentally analogous to sub-Antarctic islands were
compiled from the literature (Frenot et al., 2005; Mathakutha et al., 2019). Whenever we could not
find information collected in comparable environments, we relied on functional trait data included in
the TRY database (Kattge et al., 2020). For each alien species, the dominant reproduction strategy in
the study area was assessed relying on expert-based knowledge (personal communication, Lebouvier,
M., & Bittebiere, A.K.).

2.3 *Topoclimatic layers*

To model the species-environment relationship at fine spatial resolution, we first downloaded coarse-
grained temperature (BIO1, BIO5 and BIO6 – annual mean temperature, max temperature warmest
month and min temperature coldest month) and annual precipitation (BIO12) grid layers at 1-km
resolution (at the equator) from the CHELSA database (Karger et al., 2017) and then disaggregated
their spatial resolution using physiographically informed models fitted through geographically
weighted regression (GWR; Fotheringham & Rogerson, 2008). This downscaling technique allows
statistically predicting the local value of the coarse-grain CHELSA climatic variables as a function
of environmental grid layers available at finer spatial resolution (30-m) and known to drive

microclimate heterogeneity (Lenoir, Hattab, & Pierre 2017; Lembrechts et al., 2019). GWR-derived
topoclimatic layers, beyond allowing to model the species-environment relationship at a more
meaningful spatial resolution, have already proved to account better for the complex interactions
between macroclimate and topography (Lenoir et al., 2017; Lembrechts et al., 2019).

As using BIO5 (max temperature warmest month) and BIO6 (min temperature coldest month) in
place of BIO1 did not improve species distribution models, we ultimately used BIO1 (hereafter mean
temperature) and BIO12 as topoclimatic predictors. A full description of the downscaling procedure
is reported in Appendix S2 along with the results of the GWR models.

2.4 *Human-related layers*

As human disturbances are known to favour the establishment of alien plants through propagule
introduction and alteration of habitat conditions, we generated a 30-m resolution layer reporting the
distance between each human settlement (the research station, *Baie du La Pérouse*, *Pointe Basse* and
Baie Américaine) and any location on the island. Specifically, assuming that human disturbance is
stronger in most accessible areas, we derived a least cost distance grid layer providing a measure of
accessibility. Terrain slope changes between both orthogonally and diagonally neighbouring raster
cells were used to compute the cost of reaching any location on Possession Island starting from any
human settlement and following all potential paths of raster cells (function "accCost", "gdistance" R
package; Etten, 2018). High costs were thus associated with locations not easily reachable from
human settlements due to high topographic roughness (Appendix S3, Figure S3.2).

A network of hiking paths has been designed by the RN-TAF to restrict human movements for
wildlife conservation purposes, and walking these paths currently constitutes the only authorized way
to move across the island. As humans are a critical vector of propagule introduction and dispersal on
sub-Antarctic islands, we derived a 30-m resolution raster layer reporting the distance between any
location on Possession Island and the closest hiking path using the function "distance" from the
"raster" R package (Hijmans, 2019) (Appendix S3, Figure S3.2).

2.5 *Alien species distribution modelling*

The occurrence probability of the six alien plant species was separately modelled as a function of the
topoclimatic (mean temperature and annual precipitation) and human-related variables (path distance
and least cost) by means of logit binary generalized linear models (GLM). The single-species
distribution models (single-SDMs) were trained and tested on datasets obtained through a re-sampling
procedure of the presence/absence data performed in the environmental space to reflect all available
environmental conditions on Possession Island (see Appendix S4). All four topoclimatic and human-
related predictors were retained to fit the single-SDMs as the relative variance inflation factor
(function "vif", R package "car"; Fox & Weisberg, 2019) was always below a threshold of 3. Second-
order polynomial terms were included in the model to allow for intermediate niche optima of the
species or in case lack-of-fit tests detected consistent departure from linearity in the profile of Pearson
residuals (function "residualPlots", R package "car"; Fox & Weisberg, 2019). The statistical
significance of each predictor was tested using type II analysis of deviance (function "Anova", R
package "car"; Fox & Weisberg, 2019). We then computed the likelihood profile-based 95%
confidence intervals of the regression parameters.

Single-SDMs predictive performance was measured using the true skill statistic (TSS, equal to
sensitivity + specificity – 1; function "ecospat.max.tss", R package "ecospat"; Broennimann, Di Cola,
& Guisan, 2018) computed on the testing datasets obtained through the environmental matching
described in Appendix S4. We used the TSS as it has desirable properties of other accuracy measures

(e.g. Kappa and AUC), while being unaffected by prevalence (Allouche, Tsoar, & Kadmon, 2006).
Also, we computed the deviance-based R² value as a measure of goodness-of-fit of each single-SDM.

The occurrence probability estimated by the full single-SDMs (including both topoclimatic and
human-related predictors) for each alien plant species was mapped on a 30-m raster grid layer to
visualize their predicted distribution across Possession Island.

2.6 *Relationship between plant traits and alien species dependence on propagule pressure*

As preliminary analyses, we measured the relative importance of human-related variables in
determining alien species occurrence in the single-SDMs. To this aim, we used the sum of Akaike
weights (*w*), which provides an easily interpretable measure of variable importance (it ranges from 0
to 1, with a high value for a given variable indicating its high importance relative to the others;
Burnham & Anderson, 2002). Then we graphically related the species-specific values of the
functional traits to the sum of weights to look for relationships between plant traits and the importance
of human-related variables (see Appendix S7).

Secondly, we investigated how the interaction between human-related variables and plant traits
affected alien species occurrence in a multi-species distribution model (multi-SDM), focusing on
those functional traits that showed some relationship with the dependence on human-related variables
in the single-SDMs. To this aim, we modelled the occurrence of all alien species together as a function
of topoclimatic and human-related variables by means of a logit binary GLM, including the
interaction between species identity and topoclimatic variables on the one hand and the interaction
between species functional traits and human-related variables on the other hand. This allowed
exploring how the effect of human-related variables on alien species occurrence varied according to
plant traits, while controlling for species-specific responses to topoclimate. To select the most
parsimonious model, we fitted all possible sub-models including different combinations of the
functional traits-anthropogenic variables interaction terms (function "dredge", R package "MuMIn";
Barton, 2019), always retaining the species-topoclimate interaction terms and the main effect of path
distance and least cost in each candidate sub-model. Then, we computed the sum of Akaike weights
for each model term and used the evidence ratio as a measure of the relative importance of variables
(Massol et al., 2007; Burnham & Anderson, 2002). Specifically, we computed the evidence ratio of
the *i-th* variable (ER_i) as the odds of its sum of Akaike weights:

$$ER_i = \frac{w_i}{1 - w_i}$$

The evidence ratio was then compared with its expected value (ER_{null}) under the "null hypothesis"
that the variable explained as much deviance as a randomly generated explanatory variable, and
would thus be as likely as not to be incorporated in the best models. As all the variables were tested
in a balanced design, $ER_{null} = 1$ in all tested cases. Following Massol et al. (2007), the effect of a
variable *i* was deemed unlikely if $ER_i < 0.37\ ER_{null}$, implausible when $0.37\ ER_{null} < ER_i < ER_{null}$,
plausible when $ER_{null} < ER_i < 2.72\ ER_{null}$, and likely when $ER_i > 2.72\ ER_{null}$. These thresholds
correspond to differences in Akaike information criterion equal to +2 or -2, which are commonly
admitted as a good rule-of-thumb gap to compare model performance.

3. Results

3.1 *Effect of topoclimatic and human-related variables on single species occurrence*

Predictive performances of the single-SDMs varied greatly across species (Table 1): high values of
TSS were observed for *P. pratensis*, *S. alsine* and *J. bufonius* (0.80-0.82), while low values were

obtained for the remaining species (from 0.09 to 0.29). The R^2 values showed a similar trend, with
the highest value obtained for *J. bufonius* (0.48) and the lowest for *C. fontanum* (0.02).

Overall, the occurrence of *P. pratensis*, *S. alsine* and *J. bufonius* appeared to be strongly conditioned
by both topoclimatic and human-related variables, while *C. fontanum*, *P. annua* and *S. procumbens*
were less affected by human-related variables (Table 1, Figure 1, 2).

All alien species, except *S. alsine*, exhibited a significant positive or humped-shaped relationship with
mean temperature (Table 1), meaning that their occurrence probability increased with increasing
temperature (Figure 1, Appendix S5). More specifically, the occurrence probability of *J. bufonius*, *S.*
procumbens and *C. fontanum* peaked at mean temperature values around 4.5 °C, while the presence
of *P. pratensis* and *P. annua* "linearly" increased with temperature.

Annual precipitation significantly affected the presence of *P. pratensis*, *S. alsine* and *J. bufonius*,
while it had a minor influence on the occurrence of the other species (Table 1). In particular, the odds
of finding *P. pratensis* and *J. bufonius* decreased approximately by 90% for each 500 mm increment
in annual precipitation, while the occurrence probability of *S. alsine* sharply decreased for annual
precipitation values above 1500 mm (Figure 1).

All species except *C. fontanum* exhibited a significant negative relationship with path distance (i.e.
the occurrence probability of the species decreased at increasing distances from the hiking paths),
though its influence varied among species (Table 1, Figure 1, Appendix S5). In this regard, the odds
of finding *P. pratensis*, *S. alsine* and *J. bufonius* decreased respectively by 20%, 16% and 19%
moving 100 m away from the paths, while the odds of finding *P. annua* and *S. procumbens* decreased
by 4% and 6%, respectively.

Least cost distance to settlements appeared to influence the occurrence of all analysed species except
S. procumbens (Table 1). In particular, the odds of finding *P. pratensis*, *S. alsine* and *J. bufonius*
decreased, respectively, by 17%, 13% and 44% for each increment of 5000 units of cost of travelling
a given path from a human settlement (Figure 1, Appendix S5). On the contrary, *C. fontanum* and *P.*
annua showed a positive relationship with least cost, with their odds of occurring increasing
respectively by 9% and 5% for each increment of 5000 units of cost of travelling a given path from a
human settlement (Figure 1 and Appendix S5).

Figure 1 – Response curves of the analysed alien species in the single-SDMs.

Table 1 – Single-SDM type II analysis of deviance tables and performance measures (R^2 and TSS). LR: Likelihood Ratio statistic; Df: degrees of freedom; p-val: p-value (*** p <*
0.001; ** p < 0.01; * p < 0.05).

	Poa pratensis			*Juncus bufonius*			*Stellaria alsine*			*Poa annua*			*Sagina procumbens*			*Cerastium fontanum*		
Predictors	LR	Df	p-val	LR	Df	p-val	LR	Df	p-val	LR	Df	p-val	LR	Df	p-val	LR	Df	p-val
Mean temperature	5.083	1	*	32.516	2	***	0.312	1	= 0.576	55.538	1	***	62.554	2	***	18.325	2	***
Annual precipitation	106.443	1	***	73.647	1	***	68.406	2	***	4.759	1	*	3.681	1	= 0.055	0.041	1	= 0.840
Least cost	10.837	1	***	40.250	1	***	4.420	1	*	5.778	1	*	1.118	1	= 0.290	12.483	1	***
Path distance	15.623	1	***	8.740	1	**	7.524	1	**	3.927	1	*	6.782	1	**	2.877	1	= 0.089
R^2		0.46			0.48			0.36			0.06			0.10			0.02	
TSS		0.80			0.81			0.82			0.19			0.29			0.09	

329 *Figure 2 – Predicted occurrence of Sagina procumbens and Juncus bufonius. Pocc: occurrence probability. Dashed lines represent hiking paths, while black dots represent human*
330 *settlements. Occurrence maps of the other alien species are reported in Appendix S6 (Figure S6.4 and S6.5).*

3.2 *Plant traits and species dependence on propagule pressure*

In the preliminary analyses, residence time, life history, vegetative reproduction and plant height showed some relationship with the sum of weights of the human-related variables in the single-SDMs (Appendix S7, Figure S7.6 and S7.8), while seed- and leaf-related traits clearly showed no relationship (Appendix S7, Figure S7.7 and S7.9).

Then, the multi-SDM confirmed significant interactions of residence time, life history and plant height with the human-related variables (Appendix S7, Figure S7.10). Residence time and plant height appeared to interact with both human-related variables, while life history seemed to interact only with least cost in determining alien species occurrence. In particular, the effect of human-related variables on alien species occurrence varied with plant height, and while the occurrence probability of taller plants sharply decreased moving far from human facilities, a weaker and sometimes opposite trend was observed for shorter species (Figure 3a,b and Appendix S7, Figure S7.11). In addition, old residents were on average less affected by the human-related variables than new residents (Figure 3a,b and Appendix S7, Figure S7.11a). Finally, perennials appeared to be on average less negatively affected by least cost distance to human settlements than annuals (Figure 3c and Appendix S7, Figure S7.11b,c).

Figure 3 – Effect of the interaction between human-related variables and plant features on alien species occurrence probability (logit scale). Panel (a): effect of the path distance-plant height interaction on old and new resident occurrence probability. Panel (b): effect of the least cost-plant height interaction on old and new (annual) resident occurrence probability. Panel (c): effect of least cost distance to human settlements on annual and perennial (100 mm height) alien species occurrence (for both old and new residents). All plots are reported on a logit scale.

4. Discussion

As hypothesized, both environmental and human-related variables locally affected alien species occurrence on Possession Island, though with differences among species. Overall, results confirmed the key role of human-related propagule pressure in favouring alien species establishment and spread on sub-Antarctic islands (Frenot et al., 2005; le Roux et al., 2013; Shaw, 2013), though we also observed a significant effect of abiotic conditions. Indeed, climate barriers seemed to prevent alien species occurrence in the most environmentally stressful areas of the island (Lembrechts et al., 2016). In particular, our results suggested the existence of two main invasion patterns arising from the species-specific dependence on human-related propagule pressure (Frenot et al., 2005; Shaw, 2013): low-spread species (*P. pratensis*, *S. alsine* and *J. bufonius*) strongly relying on human-assisted dispersal along hiking trails and in the vicinity of human settlements; and high-spread species (*C. fontanum*, *P. annua* and *S. procumbens*) mostly limited by harsh climatic conditions at high altitudes.

Due to their dependence on human-related variables, low-spread species were predicted to occur mainly close to hiking paths and human settlements, pointing to the importance of anthropogenic activities as key drivers of continuous propagule pressure favouring species establishment (Whinam et al., 2005; Pickering & Mount, 2010). Once introduced through ship-to-shore transport, propagules are then likely to be dispersed on hiking paths through trampling (Whinam et al., 2005). However, the harsher environmental conditions characterizing the west side of Possession Island also limit the occurrence of low-spread species to the east. In particular, the west-east gradient of annual precipitation (Appendix S2, Figure S2.1) clearly overlaps with the low-spread species distribution (Figure 2 and Appendix S6), suggesting that their establishment might be prevented in areas with abundant precipitation. Nevertheless, the precipitation gradient is also connected to human presence, so that in the west side of the island (less inhabited and more preserved) anthropogenic propagule pressure is weaker. In any case, our results evidenced that low-spread species may lack important adaptations to successfully colonize less disturbed areas with limiting abiotic conditions, while remaining relegated to areas of high human presence (Lembrechts et al., 2018).

On the contrary, high-spread species appeared weakly (yet positively) influenced by human-related variables, suggesting that, in spite of the undisputed importance of anthropogenic activities in promoting alien plants establishment (Whinam et al., 2005; Huiskes et al., 2014), high-spread species may possess key traits releasing them from direct dependence on anthropogenic propagule pressure. Consequently, these species appeared to be mostly limited by the extreme climatic conditions of the high and cold inner sectors of Possession Island. Furthermore, the low predictive performance of high-spread SDMs indicate that the occurrence of these species can only be partly explained by the influence of topoclimatic and human-related variables, so that other factors not considered here (e.g. soil properties, plant-soil microbiota interactions, snow cover) may also play an important role in driving their distribution at even finer spatial resolutions.

Critically, although we managed to obtain relatively high-resolution topoclimatic data, it is important to realize that the CHELSA climate for the island 1) might lack the accuracy it has at temperature latitudes, being based on extrapolations from a single weather station only, and 2) represents air temperature only, while short plants as those analysed here relate more strongly to soil and near-surface temperatures (Lembrechts et al., 2019). This highlights the need for *in-situ* soil- and near-surface temperature measurements in remote locations to get more ecologically meaningful climate data.

Although the small set of analysed alien plant species calls for caution in interpretation, we confirmed here that certain plant traits confer greater invasiveness in sub-Antarctic environments. By relating

plant traits to species responses to human-related variables and analysing the effect of their interaction
on alien species occurrence, we found evidence that low-stature was a key feature that discriminated
invasive from non-invasive alien species on Possession Island. However, residence time and life
history also appeared to affect species invasiveness.

High-spread species were shorter than low-spread species (Mathakutha et al., 2019). Consistently, we
observed a sharper decrease in the occurrence probability of taller plants moving away from both
hiking paths and human settlements. As plant height is generally associated with species adaptations
to harsh environments (Cornelissen et al., 2003) and, specifically, low-stature has been attributed to
frost avoidance mechanisms in high mountains (Márquez, Rada, & Fariñas, 2006; Ladinig, Hacker,
Neuner, & Wagner, 2013), shorter species may be reasonably favoured in windy and cold sub-
Antarctic environments (Mathakutha et al., 2019) and therefore be more easily released of human
dependence. Indeed, the importance of functional traits providing tolerance to abiotic stress increases
with environmental harshness, even under strong anthropogenic disturbance (Zefferman et al., 2015).
Further, our results supported the hypothesis that residence time positively affects invasiveness
(Lockwood et al., 2005; Pyšek et al., 2015), though with some exceptions. Generally, old residents
(e.g. *C. fontanum* and *P. annua*) were less dependent on human-related propagule pressure and more
widely spread than new resident species. Nevertheless, among the old residents, *P. pratensis* was
strongly dependent on human-related variables and was still mostly restricted to the original
introduction sites. On the other hand, *S. procumbens*, a new resident, has been able to spread
extensively and quicker than the other (old residents) high-spread species. However, this might be
due to the difference in plant height of the two species: while *P. pratensis* is among the tallest analysed
species, *S. procumbens* is the shortest. The multi-SDM showed that perennials were slightly less
dependent on human presence than annuals (Figure 3c). Although annuals might benefit from high
dispersal abilities (e.g. abundant light seeds) and usually spread quicker and wider (Pertierra et al.,
2017), perennials can sustain short growing seasons (Frenot et al., 2001; Shaw, 2013) and potentially
colonize harsher environments (Dietz & Edwards 2006). In our case, short perennials (e.g. *C.
fontanum*) might be favoured over tall annuals (e.g. *J. bufonius*) due to the interaction between stress-
tolerant traits, such as plant height, and high abiotic tolerance.

Though the interaction between vegetative reproduction and human-related variables was not
included in the most parsimonious multi-SDM, alien species may still benefit from sexual
reproduction, as suggested from the lower importance of human-related variables for the occurrence
of alien plants reproducing sexually in the single-SDMs (Appendix S7). As discussed above, the high
dispersal potential of sexually reproducing alien species may foster their extensive spread as, for
instance, observed for *Poa annua* in the Antarctic Peninsula (Pertierra et al., 2017). Nonetheless, by
reproducing vegetatively, perennials might outcompete annuals in maintaining viable and persistent
populations during unfavourable seasons. Finally, in spite of their acknowledged importance in
conferring invasiveness in sub-Antarctic islands (Mathakutha et al., 2019), we found no evidence of
the role of seed and leaf traits in affecting species dependence on human-related variables. This is
possibly due to the small set of analysed alien species, which might have prevented the emergence of
further functional traits-anthropogenic variables relationships.

Overall, our approach allowed identifying fine-scale drivers of alien species distribution on sub-
Antarctic islands, along with the most likely features that favour their spread beyond sources of
continuous human-assisted introductions. Despite some limitations (e.g. limited number of species),
our study opens avenues for a more targeted management of alien plant invasions in these unique
environments, potentially allowing to anticipate future problematic invasions.

Acknowledgements

We would like to thank Marc Lebouvier (Ing. CNRS) for his suggestions which strongly improved the manuscript. We acknowledge the contribution of Lise Chambrin (Ing. RN-TAF) and all civil volunteers who collected data used in this research.

Funding: Project Granted by the French Polar Institute Paul-Emile Victor 'IPEV 136 Subanteco', Institut Universitaire de France (IUF) 'ENVIE', Stratégie d'attractivité durable (SAD) Région Bretagne 'INFLICT'. JJL is funded by the Research Foundation Flanders (FWO, project OZ7828), MC by a Rita Levi-Montalcini grant.

Authors' contribution

DR, MB, FM and JL conceived the idea; MB analyzed the data with FM and MC; MB led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

Data availability statement

Data and R code available on Zenodo: https://doi.org/10.5281/zenodo.3951465

Conflict of interest disclosure

The authors of this article declare that they have no financial conflict of interest with the content of this article.

References

Allouche, O., Tsoar, A., & Kadmon, R. (2006). Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS). *Journal of applied ecology*, *43*(6), 1223-1232.

Barton, K. (2019). MuMIn: Multi-Model Inference. R package version 1.43.6. https://CRAN.R-project.org/package=MuMIn

Broennimann, O., Di Cola, V., & Guisan, A. (2018). ecospat: Spatial Ecology Miscellaneous Methods. R package version 3.0. https://CRAN.R-project.org/package=ecospat

Burnham, K.P. & Anderson, D.R. (2002) *Model Selection and Multimodel Inference: a Practical Information‐Theoretic Approach*, 2nd edn. Springer‐Verlag, New York.

Catford, J.A., Jansson, R., & Nilsson, C. (2009). Reducing redundancy in invasion ecology by integrating hypotheses into a single theoretical framework. *Divers. Distrib.*, *15*(1), 22-40.

Convey, P., Key, R.S., & Key, R.J.D. (2010). The establishment of a new ecological guild of pollinating insects on sub-Antarctic South Georgia. *Antarct. Sci.*, *22*(5), 508-512.

Convey, P., & Lebouvier, M. (2009). Environmental change and human impacts on terrestrial ecosystems of the sub-Antarctic islands between their discovery and the mid-twentieth century. In *Papers and proceedings of the royal society of Tasmania* (Vol. 143, No. 1, pp. 33-44).

Cornelissen, J.H.C., Lavorel, S., Garnier, E., Diaz, S., Buchmann, N., Gurvich, D.E., ... Pausas, J.G. (2003). A handbook of protocols for standardised and easy measurement of plant functional traits worldwide. *Aust. J. Bot.*, 51(4), 335-380.

Cuba-Díaz, M., Fuentes, E., & Rondanelli-Reyes, M. (2015). Experimental culture of non-indigenous *Juncus bufonius* from King George Island, South Shetland Island, Antarctica. 极地研究, *26*(1-English), 24-29.

Dengler, J. (2016). Phytosociology. *International Encyclopedia of Geography: People, the Earth, Environment and Technology: People, the Earth, Environment and Technology*, 1-6.

Dietz, H., & Edwards, P. J. (2006). Recognition that causal processes change during plant invasion helps explain conflicts in evidence. *Ecology*, *87*(6), 1359-1367.

Duffy, G.A., & Lee, J.R. (2019). Ice-free area expansion compounds the non-native species threat to Antarctic terrestrial biodiversity. *Biol. Cons.*, *232*, 253-257.

Duffy, G.A., Coetzee, B.W., Latombe, G., Akerman, A.H., McGeoch, M.A., & Chown, S.L. (2017). Barriers to globally invasive species are weakening across the Antarctic. *Divers. Distrib.*, *23*(9), 982-996.

Etten, J.V. (2018). gdistance: Distances and Routes on Geographical Grids. R package version 1.2-2. https://CRAN.R-project.org/package=gdistance

Fotheringham, A.S., & Rogerson, P.A. (Eds.). (2008). *The SAGE handbook of spatial analysis*. Sage.

Fox, J., & Weisberg, S. (2019). An R Companion to Applied Regression (Third). Thousand Oaks CA: Sage. https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Frenot, Y., Chown, S.L., Whinam, J., Selkirk, P.M., Convey, P., Skotnicki, M., & Bergstrom, D.M. (2005). Biological invasions in the Antarctic: extent, impacts and implications. *Biol. Rev.*, *80*(1), 45-72.

Frenot, Y., Gloaguen, J.C., Massé, L., & Lebouvier, M. (2001). Human activities, ecosystem disturbance and plant invasions in subantarctic Crozet, Kerguelen and Amsterdam Islands. *Biol. Cons.*, *101*(1), 33-50.

Greve, M., Mathakutha, R., Steyn, C., & Chown, S.L. (2017). Terrestrial invasions on sub-Antarctic Marion and Prince Edward Islands. *Bothalia-African Biodiversity & Conservation*, *47*(2), 1-21.

Guisan, A., Thuiller, W., & Zimmermann, N.E. (2017). *Habitat suitability and distribution models: with applications in R*. Cambridge University Press.

Gutt, J., Zurell, D., Bracegridle, T., Cheung, W., Clark, M., Convey, P., ... Griffiths, H. (2012). Correlative and dynamic species distribution modelling for ecological predictions in the Antarctic: a cross-disciplinary concept. *Polar Res.*, *31*(1), 11091.

Hijmans, R.J. (2019). raster: Geographic Data Analysis and Modeling. R package version 3.0-2. https://CRAN.R-project.org/package=raster

Hughes, K.A., Convey, P., Pertierra, L.R., Vega, G.C., Aragón, P., & Olalla-Tárraga, M.Á. (2019). Human-mediated dispersal of terrestrial species between Antarctic biogeographic regions: A preliminary risk assessment. *J. Environ. Manage.*, *232*, 73-89.

Hughes, K.A., Pertierra, L.R., Molina-Montenegro, M.A., & Convey, P. (2015). Biological invasions in terrestrial Antarctica: what is the current status and can we respond?. *Biodiversity and Conservation*, *24*(5), 1031-1055.

Huiskes, A.H., Gremmen, N.J., Bergstrom, D.M., Frenot, Y., Hughes, K.A., Imura, S., ... Ware, C. (2014). Aliens in Antarctica: assessing transfer of plant propagules by human visitors to reduce invasion risk. *Biol. Cons.*, *171*, 278-284.

Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., ... Kessler, M. (2017). Climatologies at high resolution for the earth's land surface areas. *Sci. data*, *4*, 170122.

Kattge, J., Bönisch, G., Díaz, S., Lavorel, S., Prentice, I.C., Leadley, P., ... Acosta, A.T. (2020). TRY plant trait database–enhanced coverage and open access. *Global Change Biol*.

Ladinig U, Hacker J, Neuner G, Wagner J (2013) How endangered is sexual reproduction of high-mountain plants by summer frosts? Frost resistance, frequency of frost events and risk assessment. *Oecologia* 171:743–760

Lebouvier, M., Laparie, M., Hulle, M., Marais, A., Cozic, Y., Lalouette, L., ... Renault, D. (2011). The significance of the sub-Antarctic Kerguelen Islands for the assessment of the vulnerability of native communities to climate change, alien insect invasions and plant viruses. *Biol. Invasions*, *13*(5), 1195-1208.

Lembrechts, J.J., Lenoir, J., Roth, N., Hattab, T., Milbau, A., Haider, S., ... Nuñez, M.A. (2019). Comparing temperature data sources for use in species distribution models: From in-situ logging to remote sensing. *Global Ecol. Biogeogr.*, *28*(11), 1578-1596.

Lembrechts, J.J., Lenoir, J., Nuñez, M.A., Pauchard, A., Geron, C., Bussé, G., ... Nijs, I. (2018). Microclimate variability in alpine ecosystems as stepping stones for non-native plant establishment above their current elevational limit. *Ecography*, *41*(6), 900-909.

Lembrechts, J.J., Pauchard, A., Lenoir, J., Nuñez, M.A., Geron, C., Ven, A., ... Milbau, A. (2016). Disturbance is the key to plant invasions in cold environments. *PNAS*, *113*(49), 14061-14066.

Lenoir, J., Hattab, T., & Pierre, G. (2017). Climatic microrefugia under anthropogenic climate change: implications for species redistribution. *Ecography*, *40*(2), 253-266.

le Roux, P.C., Ramaswiela, T., Kalwij, J.M., Shaw, J.D., Ryan, P.G., Treasure, A.M., ... Chown, S.L. (2013). Human activities, propagule pressure and alien plants in the sub-Antarctic: Tests of generalities and evidence in support of management. *Biol. Cons.*, 161, 18-27.

Lockwood, J.L., Cassey, P., & Blackburn, T. (2005). The role of propagule pressure in explaining species invasions. *Trends Eco. Evo.*, 20(5), 223-228.

Márquez, E.J., Rada, F., & Fariñas, M.R. (2006). Freezing tolerance in grasses along an altitudinal gradient in the Venezuelan Andes. *Oecologia*, *150*(3), 393-397.

Massol, F., David, P., Gerdeaux, D., & Jarne, P. (2007). The influence of trophic status and large-scale climatic change on the structure of fish communities in Perialpine lakes. *J. Anim. Ecol.*, *76*(3), 538-551.

Mathakutha, R., Steyn, C., le Roux, P.C., Blom, I.J., Chown, S.L., Daru, B.H., ... Greve, M. (2019). Invasive species differ in key functional traits from native and non-invasive alien plant species. *J. Veg. Sci.*, *30*(5), 994-1006.

Molina-Montenegro, M.A., Carrasco-Urra, F., Rodrigo, C., Convey, P., Valladares, F., & Gianoli, E. (2012). Occurrence of the non-native annual bluegrass on the Antarctic mainland and its negative effects on native plants. *Cons. Biol.*, *26*(4), 717-723.

Moser, D., Lenzner, B., Weigelt, P., Dawson, W., Kreft, H., Pergl, J., ... Cassey, P. (2018). Remoteness promotes biological invasions on islands worldwide. *PNAS*, *115*(37), 9270-9275.

Ottaviani, G., Keppel, G., Götzenberger, L., Harrison, S., Opedal, Ø. H., Conti, L., ... Negoita, L. (2020). Linking Plant Functional Ecology to Island Biogeography. *Trends Plant Sci*.

Pérez-Harguindeguy, N., Diaz, S., Gamier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., ... Urcelay,
C. (2013). New handbook for stand-ardised measurement of plant functional traits worldwide. *Aust.*
J. Bot. 61: 167-234.

Pertierra, L.R., Aragón, P., Shaw, J.D., Bergstrom, D.M., Terauds, A., & Olalla-Tárraga, M.Á.
(2017). Global thermal niche models of two European grasses show high invasion risks in
Antarctica. *Global Change Biol.*, *23*(7), 2863-2873.

Pickering, C., & Mount, A. (2010). Do tourists disperse weed seed? A global review of unintentional
human-mediated terrestrial seed dispersal on clothing, vehicles and horses. *Journal of Sustainable*
Tourism, *18*(2), 239-256.

Pyšek, P., Manceur, A.M., Alba, C., McGregor, K.F., Pergl, J., Štajerová, K., ... Lučanová, M. (2015).
Naturalization of central European plants in North America: species traits, habitats, propagule
pressure, residence time. *Ecology*, *96*(3), 762-774.

Pyšek, P., & Richardson, D.M. (2008). Traits associated with invasiveness in alien plants: where do
we stand?. In *Biol. Invasions* (pp. 97-125). Springer, Berlin, Heidelberg.

Richardson, D.M., & Pyšek, P. (2006). Plant invasions: merging the concepts of species invasiveness
and community invasibility. *Prog. Phys. Geogr.*, *30*(3), 409-431.

Shaw, J.D. (2013). Southern Ocean Islands invaded: conserving biodiversity in the world's last
wilderness. In *Plant Invasions in Protected Areas* (pp. 449-470). Springer, Dordrecht.

van Kleunen, M., Dawson, W., & Maurel, N. (2015). Characteristics of successful alien
plants. *Molecular Ecology*, *24*(9), 1954-1968.

van Kleunen, M., Weber, E., & Fischer, M. (2010). A meta-analysis of trait differences between
invasive and non-invasive plant species. *Ecology letters*, *13*(2), 235-245.

Whinam, J., Chilcott, N., & Bergstrom, D.M. (2005). Subantarctic hitchhikers: expeditioners as
vectors for the introduction of alien organisms. *Biol. Cons.*, *121*(2), 207-219.

Wilson, J.R.U., Richardson, D.M., Rouget, M., Proches, S., Amis, M.A., Henderson, L. … Thuiller,
W. (2007). Residence time and potential range: crucial considerations in modelling plant invasions.
Divers. Distrib., 13, 11–22.

Zefferman, E., Stevens, J. T., Charles, G. K., Dunbar-Irwin, M., Emam, T., Fick, S., ... Young, T. P.
(2015). Plant communities in harsh sites are less invaded: a summary of observations and proposed
explanations. *AoB Plants*, *7*.