Title: **Species-Specific Responses of Bird Song Output in the Presence of Drones**

Authors: Andrew M. Wilson[1], Kenneth S. Boyle[1], Jennifer L. Gilmore[1], Cody J. Kiefer[1],

Matthew F. Walker[1]

Affiliations: [1] Environmental Studies Department, Gettysburg College, 300 North Washington

Street, Gettysburg, PA, 17325, United States

Correspondence author: Andrew M. Wilson, awilson@gettysburg.edu

Key words: drone; UAV; UAS; songbird; bioacoustics; noise pollution

Abstract

Drones are now widely used to study wildlife, but applications for studying bioacoustics have

been limited. Drones can be used to collect data on bird vocalizations, but an ongoing concern is

that noise from the drones could change bird vocalization behavior. To test this we conducted an

experiment using 30 sound localization arrays to track the song output of seven songbird species

before, during, and after a 3-minute flight of a small quadcopter drone hovering at 50 m above

ground level. We analyzed 8,303 song bouts, of which 2,285 song bouts of 184 individual birds

were within 50 meters of the array centers. We used GLMMs to assess patterns in song output

showed patterns that could be attributed to the drone's presence. We found no evidence of any

effect of the drone for five species: American Robin *Turdus migratorius*, Common Yellowthroat

Geothlypis trichas, Field Sparrow *Spizella pusilla*, Song Sparrow *Melospiza melodia*, and Indigo

Bunting *Passerina cyanea*. However, we found a substantial decrease in Yellow Warbler

Setophaga petechia song detections during the 3-minute drone hover, such that there was an 81%

drop in detections after 2 minutes (Wald-test, $p<0.001$), compared with before the drone's

introduction. In contrast, the number of singing Northern Cardinal *Cardinalis cardinalis*

increased after the drone was introduced, and remained almost five-fold higher for 4-minutes

after the drone departed ($p<0.001$). Further, we found an increase in cardinal contact/alarm calls

when the drone was overhead, with the elevated calling-rate sustaining for 2 minutes after the

drone had departed ($p<0.001$). Our study suggests that responses of songbirds to drones may be

species-specific, an important consideration when proposing the use of drones in avian studies.

We note that recent advances in drone technology have resulted in much quieter drones, which

makes us hopeful that the impacts that we detected could be greatly reduced.

Introduction

Drones or Unoccupied Aircraft Systems (UASs) are now well-established tools in field ecology, especially for purposes of mapping habitats or counting and tracking megafauna (Vermeulen et al. 2013, Christiansen et al. 2016, Witczuk et al. 2017, Kelaher et al. 2020). Drones have been proven to be effective tools for counting and monitoring large birds, especially those found in open habitats (Ratcliffe et al. 2015, Barnas et al. 2019) or that nest or roost in tree canopies (Weissensteiner et al. 2015, Afán et al. 2018), but there have been few studies that show drones to be useful for monitoring small birds, or those that inhabit dense vegetation, where the reach of aerial imagery is limited. Despite some of the potential advantages of drones, for example their efficiency, ease of access, and potential to reduce disturbance (Borrelle and Fletcher 2017), drones are not widely used for monitoring smaller birds, such as songbirds, which constitute the majority of birds species and abundance in most terrestrial environments (Musgrove et al. 2013, Inger et al. 2015, Rosenberg et al. 2019). Recent advances in using drones coupled with audio-recorders for bioacoustics monitoring (Wilson et al. 2017, Kloepper and Kinniry 2018), or infrared cameras to detect passerine nests (Scholten et al. 2019), suggest that the utility of using drones for monitoring smaller birds could grow.

An important drawback of drones is that they emit noise, which, in addition to their visual impacts, can cause disturbance to wildlife. There have now been many studies that investigate the impacts of drones on wildlife. A review of these concluded that smaller drones, flown at greater heights, and spending reduced time overhead, were less likely to elicit a response in study organisms (Mulero-Pázmány et al. 2017). However, there are still many unknown impacts of drones that have yet to be studied, especially for non-target organisms and their responses.

In development protocols for using drones for bioacoustics monitoring, we are interested

in the possible impacts of drones on songbirds for two important reasons: firstly, if drones do

impact songbird behavior, it introduces an ethical decision into whether they should be used at

all, and secondly, if drones cause songbirds to move or change vocalization behavior, data

derived from airborne bioacoustics could be biased. In this study we aim to assess the impacts of

hovering a drone over a study area for the intention of airborne bioacoustics monitoring, to test

the hypothesis that drones initiate vocal behavior responses in songbirds. While the bounds of

our study are narrow--we focus on the impacts of a single drone model, flown at a specific

height--we hope that this study sheds light on potential impacts of drones on what to most

ecologists would be non-target species. Further, there is a wider concern that as an additional

source of anthropogenic noise pollution, drones flown by recreationists as well as professionals

may be adding to growing noise pollution problems. Hence, we place our study within a broader

context of anthropogenic noise disturbance.

Methods

To test to see whether there was a change in songbird vocalizing behavior that could be

attributed to the drone's presence, we set up an experiment whereby ambient bird sound in the

area under the drone was recorded before, during and after a drone flight. Following a previous

study (Wilson et al. 2017), we simulated an airborne attempt at conducting a bird point count by

hovering a small quadcopter (DJI Mavic Pro) over a pre-determined count location for 3-

minutes. Traditional point count duration is typically in the range of 3 to 10 minutes, and we

chose the lower end of that range because this minimizes flight time, thereby allowing for greater

survey efficiency—potentially allowing for multiple points to be surveyed on each drone battery.

Additionally, shorter flight duration is less likely to cause excessive disturbance (Mulero-

Pázmány et al. 2017).

Field Study

We used sound localization to locate singing birds in the vicinity of a ground-based

Automated Recording Unit (ARU) array to track bird song output before, during, and after our

drone flights (Figure 1i). By arranging four ARUs in a 50 x 50 m array, we were able to measure

the differences in the time taken for bird vocalizations to reach the recorders, a method called

"time difference of arrival" (TDOA). The differences in the time of arrival allowed us to

triangulate birds in two dimensional space inside and beyond the array using the Sound Finder

Microsoft Excel application (Wilson et al. 2014). The Sound Finder method has previously been

found to provide accurate localizations, with a mean error of 4.3 m, and 74% to within 10 m

(Wilson et al. 2014).

We conducted the experiments in June and early July of 2017 at Pennsylvania State

Game Lands 249, Heidlersburg, Adams County, Pennsylvania (39°56'14.6"N 77°10'38.6"W).

The experiment had 30 replicates at locations on a 200 m grid to ensure a representative sample

of the habitat and to reduce the likelihood of resampling birds between replicates. At each

location, we deployed four SM3 ARUs (Wildlife Acoustics) on tripods 1.5 m off the ground and

in a 50 m x 50 m array centered on the predetermined grid points. The recorders were fitted with

Garmin GPS time-synch receivers to synchronize their clocks. Once the recorders were set up,

we retreated to at least 150 m from the center of the array to allow normal bird activity to resume

following possible human disturbance. After at least 10 minutes, we deployed the drone from a

base station at least 150 m away and programmed it to autonomously hover 48 m above ground

level at the center of the array for three minutes, to simulate an aerial point count as described in

Wilson et al. (2017). After returning the drone to the base station, we continued recording

ambient noise in the study array for an additional five minutes. All flights were preprogrammed

using the Litchi App (VC Technology Ltd.) to ensure high degrees of spatial precision.

Bioacoustic Analysis and Sound Localization

We used Audacity to clip the recordings from each of the four ARUs to the exact same

11-minute time-span (to nearest millisecond), which included 4-minutes before the drone flight,

3-minutes while the drone was overhead, and 4-minutes after the drone flight. We used 4-

minutes before and after the drone flight so that we could exclude the one-minute immediately

before and after the drone-flight, during which the drone was audible on the recordings as it

approached and departed the array. This allowed us to isolate 3 x 3-minute blocks for "before"

(minutes 1-3) "during" (minutes 5-7) and "after" (minutes 9-11) the drone flight.

We focused our data analysis on seven species that were known to be present on the site

in moderate to high densities: American Robin *Turdus migratorius*, Yellow Warbler *Setophaga*

petechia, Common Yellowthroat *Geothlypis trichas*, Field Sparrow *Spizella pusilla*, Song

Sparrow *Melospiza melodia*, Indigo Bunting *Passerina cyanea*, and Northern Cardinal

Cardinalis cardinalis. Some other abundant species were not included in the study because we

could not reliably identify individual birds in the audio recordings due to incessant song (Gray

Catbird *Dumetella carolinensis*) and localized high densities (Red-winged Blackbird *Agelaius*

phoeniceus). For our seven study species, we identified all song bouts that were clearly visible in

spectrograms from all four ARUs (Figure 1ii), and assigned a unique identification code to

identify each individual bird based on location, song volume, song-bout spacing, and unique

characteristics of song.

Time differences of arrival were measured manually from spectrograms in program

Raven Pro 1.5 (Figure 1iii). Spectrogram settings were Hanning window with 512 samples and

89% overlap. The time differences and air temperature at the time of the experiment (logged by

the ARUs) were then input into the Sound Finder spreadsheet to localize the location of each

bird, with locations given in UTM coordinates. Each songbird location was added to a scatterplot

in the Excel spreadsheet relative to the coordinates of the ARUs, to manually verify that the

calculated coordinates were logical (Figure 1iv), for example, if the vocalization reached ARU C

first, then the bird should be closer to that recorder than the others. The location of each

individual bird was localized for every song bout throughout the 11-minute experiment, so that

locations and song output could be tracked through space and time.

While listening to the recordings to locate song bouts of our target species, we observed

an apparent increase in Northern Cardinal "tik" contact/alarm calls when the drone was

overhead. Because these calls are quieter than songs, and therefore difficult to localize because

many are not detected on all four ARUs, we used a different approach to see whether the rates of

these calls changed in the presence of the drone. We used Kaleidoscope software (Wildlife

Acoustics 2019) to automatically detect Northern Cardinal "tik" calls in recordings from a single

ARU (A) from each array. To train Kaleidoscope we developed a training set of 10 recordings of

"tik" calls. All identified calls were verified aurally, to exclude calls of other species.

 Statistical Analysis

To test whether birdsong detections varied by time, we constructed linear mixed-effect

models (GLMMs) using the 'glmer' function in R package lme4 (Bates et al. 2015). The

response variable was whether or not an individual bird was detected, modelled with a binomial

error distribution. We constructed eight models: a null model with individual bird as a random

effect (model 1), a model with distance to bird as a covariate (model 2), three models with time

effects (models 3-5), and three models with time effects plus distance (models 6-8). The three

time effects were: individual minutes (models 3 and 6); the 3-minute periods "before", "during"

and "after" (models 4 and 7); and "before" and "after" but with effects varying by minute during

the drone flight (models 5 and 8; Figure 2). All eight models included individual bird as a

random effect. Models were compared using AIC, the size of effects determined using odd-

ratios, and significance determined using Wald tests.

We used the same eight model structures illustrated in Figure 2 to test whether the

singing rate (number of song bouts per minute) varied according to time period, this time with a

Poisson error distribution. The same Poisson model was used to test whether the Northern

Cardinal's "tik" contact/alarm call showed time dependent patterns consistent with a response to

the drone flight.

Results

We identified 8,303 song bouts across the seven study species, which were initially

estimated to belong to 467 different individual birds (Table 1). However, ˜50% of the song bouts

were localized to distances greater than 100 m from the array centers, and hence duplication of

individuals between adjacent arrays was likely. Further, many song bout detections were either

not localizable (typically because they were not clear enough in all 4 spectrograms), or had large

localization errors (greater than 5 ms) and were excluded from data analysis.

Song bout detections showed a decrease in the three minutes when the drone was

overhead, but most of this decline was of more distant birds, while the number of detections

within 50 m was less variable over time (Figure 3). However, there were clear differences among

the seven study species. For three species: Common Yellowthroat, Song Sparrow, and Indigo

Burning, the patterns did not suggest a significant response to the drone. Detections of American

Robin, Yellow Warbler, and Field Sparrow all showed a decline when the drone was overhead,

while Northern Cardinal detections increased during the drone flight, and the increased song

output persisted 4-minutes after the drone had departed (Figure 3).

Because most of the decline in song detections during the drone flight was of more

distant birds (50-100 m from array centers; Figure 3), we assume that this was due to masking of

songs by the drone noise on the recordings; it is unlikely that distant birds responded to the drone

more than those closer to it. For this reason, we restricted our statistical analysis to the 2,285

song bouts of birds localized to within 50 m of the array centers (mean distance across bouts),

which we attributed to 184 individual birds (Table 1). Yellow Warbler and Song Sparrow were

the most abundant species, followed by Field Sparrow; all three species averaged more than one

singing bird per array (within 50 m radius).

For five of the seven species, the null model (no time effects) GLMM had the best fit

(lowest AIC), indicating no evidence for an effect of the drone on song detection within a 50 m

radius (Table S1). The null model was also the best model when data for all seven species were

combined. The two exceptions were the Yellow Warbler and the Northern Cardinal. For the

Yellow Warbler, model 5 was the best fit, which indicates that song output was depressed when

the drone was overhead and that the effect increased over time, such that there was a non-

significant (Wald-test; p=0.311) 34% decrease in detections in the first minute, a non-significant

54% (p=0.035) decrease in the second minute, and a highly significant 81% (p=0.001) decrease

in the third minute, when compared with the period before the drone flight (Table 2). Song

detections remained 51% lower after the drone had departed (p=0.013). For the Northern

Cardinal, the drone effect was also significant, but the results were the opposite of those for the

Yellow Warbler--cardinal song detections increased by a non-significant 46% (p=0.45) when the

drone was overhead, and increased almost five-fold (p<0.001) after the drone had departed

(Table 2).

The null model (no time effects) was the best model for singing-rate for four species:

Yellow Warbler, Field Sparrow, Song Sparrow, and Common Yellowthroat (Table S2). For the

American Robin, model 7 was the best fit, but none of the estimates for time periods were

significant (Table 3). For the Indigo Burning, model 4 was the best fit, but the AIC was only

marginally lower than that of the null model, and none of the estimates for time periods were

significant (Table 3). We conclude that there was no evidence that singing-rates changed in

response to the drone for six of the seven species. Once again, the results for Northern Cardinal

were anomalous with evidence of a significant drop in singing-rate in minute 6 (p=0.019; Table

3), although once again, the time-effect model (5) did not appreciably improve model fit over the

null model (Table S2).

A total of 1,148 Northern Cardinal "tik" calls were detected on recordings from ARU

"A", these detections coming from 23 of the 30 arrays. The calling rate increase from an average

of 3.21/minute (se 1.39) during minutes 1 to 4 to 5.96/minute (se 1.11) in minutes 4 to 9, then

declined to 2.60/minute (se 0.79) in the last two minutes of the experiment (Figure 4). Calling

rates were significantly higher in minutes 5 thru 9 compared with minute 1 (Wald Tests,

$p<0.001$). It should be noted that there appeared to be an inflection point in call rate at about 3

minute 30 second, which is around the time that the drone was first audible in the recording as it

approached the array (Figure 4).

Discussion

Drawing on multiple lines of evidence, we conclude that overall, a drone hovering at 48

m a.g.l. for a 3-minute duration, causes little change in songbird output in the area below the

drone, within a 50-m radius. However, there were significant impacts of the drone on the

detection of songs of two species: the Yellow Warbler, which saw a substantial drop in song

detection while the drone was overhead, and the Northern Cardinal, which saw a significant

increase in song detections, especially after the drone's departure. The pronounced increase in

Northern Cardinal contact/alarm calls during and following the drone flights provides further

evidence that the vocal behavior of that species was affected by the drone's presence.

The fact that we found that responses to the drone were varied among a small group of

passerines is intriguing. Differential responses to anthropogenic noise disturbance have been

detected by others (Francis et al. 2009, McClure et al. 2013, Roca et al. 2016). The differences

have, in part, been attributed to whether the vocalization frequency of a bird species overlaps

with the frequency of the noise disturbance (Francis et al. 2009), presumably because birds

would not waste energy vocalizing if it was likely to be masked by other noise. However, of our

seven study species, two have relative low frequency songs (American Robin and Northern

Cardinal; peak sound pressure 2 to 4 kHz), Field Sparrows are slightly higher (peak sound

pressure 3 to 5 kHz) and the others have higher frequency songs (peak sound pressure 4 to 8

kHz) (Pieplow 2017), so we find no discernible association between sound frequency and effects

of the drone. Being a small vehicle, the noise emitted by a DJI Mavic Pro is of higher

frequencies than larger drones, and with a broad spectrum (Miljkovic 2018), which overlaps with

the songs of all species included in our study. The increased singing and calling of cardinals

indicates agitation at the drone noise. The Northern Cardinal is a species that has been show to

increase its song frequency in response to anthropogenic noise (Dowling et al. 2012, Seger-

Fullam et al. 2012).

It should be noted that we chose the DJI Mavic Pro for this study because of its suitability

for conducting aerial bioacoustics surveys: it is small and therefore quieter than the larger

vehicles that are typically used in wildlife research. Although we don't have our own

measurement of the drone noise from the ground, the manufacturer reports sound pressure

(SPLa) of 70 dB at 1 meter (per Valle and Scarton 2019), which would attenuate to around 36 dB

at ground-level directly below a drone hovering at 48 m, and around 33 dB at ground-level at the

edge of the 50 m radius. In a review of anthropogenic disturbance affects, it was found that

impacts on bird song, abundance, reproduction, and stress hormone levels were typically found

when anthropogenic noise levels exceeded 45 dB SPL (Shannon et al. 2016). Therefore, while

not wishing to downplay the noise pollution emitted from small drones, the noise levels that

birds were subjected to in our experiment were modest in comparison to some other sources of

anthropogenic noise. However, the relatively high frequency of drone noise compared to most

forms of anthropogenic noise pollution, which is loudest below 2 kHz (Kight et al. 2012), results

in more overlap with the frequency of most bird song, and hence it could be that drone noise

disturbance to birds is a function of its frequency, rather than loudness.

It has previously been shown that prolonged exposure to drones has an increased

disturbance effect (Mulero-Pázmány et al. 2017). Our results support this with impacts most

evident after 2 minutes of exposure for both the Yellow Warbler and Northern Cardinal. If

drones are to be used for bioacoustics surveys for birds, we recommend that very short duration

counts are used. Not only does this reduce disturbance, but it also allows for multiple counts to

be made on a single drone battery. If short duration counts are considered inadequate, for

example if the probability of detecting a bird is too low, an alternative approach would be to do

repeated counts over the duration of a day, or perhaps over several days. This would then provide

a sampling framework that was amenable to estimating species presence or abundance using

occupancy models (Royle and Nichols 2003, Hayes and Monfils 2015) or spatially replicated N-

mixture models (Royle 2004). An added advantage of using drones for aerial bioacoustics

surveys under those circumstances is that they are highly replicable, both in terms of location and

duration (due the ability to precisely program missions), and because audio recordings provide a

permanent record of bird song that can be analyzed by multiple observers after the fact

(Shonfield and Bayne 2017).

A limiting factor in our study is that despite the very large number of song bouts

analyzed, sample sizes for the number of individual birds for some of our study species were

small, which may have reduced the power to detect more modest responses. However given the

fact that many of the birds tracked were found to sing throughout the time that drone was

overhead, and that there was no evidence that singing-rates declined for six of our seven species,

we can be certain that effects were very limited for five of the seven species. It should also be

noted that most of the species in our study sing from relatively low vegetation (at least at our

study site, pers. obs.); the Indigo Bunting being an exception. Therefore, we can't generalize our

results to species that sing from more elevated perches, which would be closer to the drone and

perhaps more susceptible to disturbance. Clearly, further studies of the impacts of drone on bird

singing behavior are merited to see whether our results hold for a wider range of species, and

different habitat types.

Measurement error, especially with respect to the sound localization process could also

have impacted our results. Limiting the statistical analysis to within 50 m of the array center will

have reduced localization errors because errors increase with distance from the array (Campbell

and Francis 2013, Wilson et al. 2014), but even so, we should expect that errors of 5 to 10 m

were commonplace, even if the mean error was small (Wilson et al. 2014). Given that our study

did not aim to estimate populations densities—a common application of sound localization—the

effects of these errors on our findings may be limited to whether or not individual birds were

included in the statistical analysis (i.e. whether were they <50 m from array center). We do not

think that erroneous inclusion or exclusion of a few individuals would be sufficient to change our

findings, especially as those marginal birds are furthest from the drone, and hence the least likely

to be affected by its presence.

It is possible that our ability to individually identify birds based on their song pattern and

location was not always accurate. Anecdotally, we observed that Yellow Warblers may have

shown song-pattern plasticity (Lowther et al. 1999), which could potentially have resulted in us

over-estimating the number of different birds. However, because we were using sound

localization to track birds in space and time, we think cases of confusion due to individual bird

song-switching would be limited to a few individuals at the most.

Our study focused on assessing the effects of drone noise on bird song for a specific

vehicles type and altitude, which was due to our interest in using drones for aerial bioacoustics.

However, our results have more general interest for those concerned about disturbance effects of

drones on wildlife, and to our knowledge, ours is the first study to focus specifically on how

drones affect bird vocalizing behavior. A survey of UAS survey altitudes, focusing on wildlife

applications, shows that 50 m is frequently used for small UASs (Sherman and Chatowsky in.

prep.).

That fact that we did find some bird behavioral responses to the drone in our study gives

us reason for pause with respect to using drones for airborne bioacoustics. However, there has

been a rapid improvement in technology to reduce unwanted drone noise, even in the time since

our study. More recent DJI Mavic models are substantially quieter, with refined propeller shapes

reportedly reducing sound pressure by 4 dB, which represent about 60% reduction in noise

(Valle and Scarton 2019). With the advent of quieter drones, we hope that concerns about noise

pollution impacts on wildlife will diminish, but we urge for continued vigilance with regards to

the behavioral impacts of using drones to study wildlife.

Ethics Statement

Our experiments were approved by the Gettysburg College Institutional Animal Care and Use Committee (IACUC). The drone was operated by the US Federal Aviation Authority (FAA) Part 107 license holder A.M. Wilson, and in accordance with FAA guidelines.

Acknowledgements

We thank the Pennsylvania Game Commission for permitting us to use State Game Lands 249 for the experiments. State Game Lands 249 is on the lands of the Susquehannock people. Gettysburg College students Taylor Derick, Arthur Garst, and Samantha Trueman helped with data analysis. The research was funding by the Gettysburg College Cross-Disciplinary Science Institute (X-SIG).

Literature Cited

Afán, I., M. Máñez, and R. Díaz-Delgado (2018). Drone Monitoring of Breeding Waterbird Populations: The Case of the Glossy Ibis. Drones 2:42.

Barnas, A. F., B. J. Darby, G. S. Vandeberg, R. F. Rockwell, and S. N. Ellis-Felege (2019). A comparison of drone imagery and ground-based methods for estimating the extent of habitat destruction by lesser snow geese (Anser caerulescens caerulescens) in La Pérouse Bay. PloS one 14:e0217049–e0217049.

Bates, D., M. Mächler, B. M. Bolker, and S. C. Walker (2015). Fitting linear mixed-effects models using lme4. Journal of Statistical Software 67:0–48.

Borrelle, S. B., and A. T. Fletcher (2017). Will drones reduce investigator disturbance to surface-nesting birds? Marine Ornithology 45:89–94.

Campbell, M., and C. M. Francis (2013). Using Stereo-Microphones to Evaluate Observer Variation in North American Breeding Bird Survey Point Counts. http://dx.doi.org/10.1525/auk.2011.10005.

Christiansen, F., A. M. Dujon, K. R. Sprogis, J. P. Y. Arnould, and L. Bejder (2016). Noninvasive unmanned aerial vehicle provides estimates of the energetic cost of reproduction in humpback whales. Ecosphere 7:e01468.

Dowling, J. L., D. A. Luther, and P. P. Marra (2012). Comparative effects of urban development and anthropogenic noise on bird songs. Behavioral Ecology 23:201–209.

Francis, C. D., C. P. Ortega, and A. Cruz (2009). Noise Pollution Changes Avian Communities and Species Interactions. Current Biology 19:1415–1419.

Hayes, D. B., and M. J. Monfils (2015). Occupancy modeling of bird point counts: Implications of mobile animals. Journal of Wildlife Management 79:1361–1368.

Inger, R., R. Gregory, J. P. Duffy, I. Stott, P. Voříšek, and K. J. Gaston (2015). Common European birds are declining rapidly while less abundant species' numbers are rising. Ecology Letters 18:28–36.

Kelaher, B. P., A. P. Colefax, A. Tagliafico, M. J. Bishop, A. Giles, and P. A. Butcher (2020). Assessing variation in assemblages of large marine fauna off ocean beaches using drones. Marine and Freshwater Research 71:68–77.

Kight, C. R., M. S. Saha, and J. P. Swaddle (2012). Anthropogenic noise is associated with reductions in the productivity of breeding Eastern Bluebirds (Sialia sialis). Ecological Applications 22:1989–1996.

Kloepper, L. N., and M. Kinniry (2018). Recording animal vocalizations from a UAV: Bat echolocation during roost re-entry. Scientific Reports 8:8–13.

Lowther, P. E., C. Celada, N. K. Klein, C. C. Rimmer, and D. A. Spector (1999). Yellow Warbler (Dendroica petechia). The Birds of North America Online.

https://doi.org/10.2173/bna.454

McClure, C. J. W., H. E. Ware, J. Carlisle, G. Kaltenecker, J. R. Barber, K. Riitters, J. Wickham,
L. Fahrig, T. Rytwinski, A. Benítez-López, R. Alkemade, et al. (2013). An experimental
investigation into the effects of traffic noise on distributions of birds: avoiding the phantom
road. Proceedings. Biological sciences / The Royal Society 280:20132290.

Miljkovic, D. (2018). Methods for attenuation of unmanned aerial vehicle noise. 2018 41st
International Convention on Information and Communication Technology, Electronics and
Microelectronics, MIPRO 2018 - Proceedings:914–919.

Mulero-Pázmány, M., S. Jenni-Eiermann, N. Strebel, T. Sattler, J. J. Negro, and Z. Tablado
(2017). Unmanned aircraft systems as a new source of disturbance for wildlife: A
systematic review. PLOS ONE 12:e0178448.

Musgrove, A., N. Aebischer, M. Eaton, R. Hearn, S. Newson, D. Noble, M. Parsons, K. Risely,
and D. Stroud (2013). Population estimates of birds in Great Britain and the UK. British
Birds 106:231–232.

Pieplow, N. (2017). Peterson Field Guide to Bird Sounds of Eastern North Americae. Houghton
Mifflin Harcour, New York, New York, USA.

Ratcliffe, N., D. Guihen, J. Robst, S. Crofts, A. Stanworth, and P. Enderlein (2015). A protocol
for the aerial survey of penguin colonies using UAVs1. http://dx.doi.org/10.1139/juvs-
2015-0006.

Roca, I. T., L. Desrochers, M. Giacomazzo, A. Bertolo, P. Bolduc, R. Deschesnes, C. A. Martin,
V. Rainville, G. Rheault, and R. Proulx (2016). Shifting song frequencies in response to
anthropogenic noise: a meta-analysis on birds and anurans. Behavioral Ecology 27:1269–
1274.

Rosenberg, K. V, A. M. Dokter, P. J. Blancher, J. R. Sauer, A. C. Smith, P. A. Smith, J. C.
Stanton, A. Panjabi, L. Helft, M. Parr, and P. P. Marra (2019). Decline of the North
American avifauna. Science 366:120 LP – 124.

Royle, J. A. (2004). N-mixture models for estimating population size from spatially replicated
counts. Biometrics 60:108–115.

Royle, J. A., and J. D. Nichols (2003). Estimating abundance from repeated presence-absence
data or point counts. Ecology 84:777–790.

Scholten, C. N., A. J. Kamphuis, K. J. Vredevoogd, K. G. Lee-Strydhorst, J. L. Atma, C. B.
Shea, O. N. Lamberg, and D. S. Proppe (2019). Real-time thermal imagery from an
unmanned aerial vehicle can locate ground nests of a grassland songbird at rates similar to
traditional methods. Biological Conservation 233:241–246.

Seger-Fullam, K. D., A. D. Rodewald, and Ji. A. Soha (2012). Urban noise predicts song
frequency in Northern Cardinals and American Robins. Bioacoustics 20:267–276.

Shannon, G., M. F. McKenna, L. M. Angeloni, K. R. Crooks, K. M. Fristrup, E. Brown, K. A.

Warner, M. D. Nelson, C. White, J. Briggs, S. McFarland, and G. Wittemyer (2016). A
synthesis of two decades of research documenting the effects of noise on wildlife.
Biological Reviews 91:982–1005.

Shonfield, J., and E. M. Bayne (2017). Autonomous recording units in avian ecological research:
current use and future applications. Avian Conservation and Ecology 12.

Valle, R. G., and F. Scarton (2019). Effectiveness, efficiency, and safety of censusing eurasian
oystercatchers haematopus ostralegus by unmanned aircraft. Marine Ornithology 47:81–87.

Vermeulen, C., P. Lejeune, J. Lisein, P. Sawadogo, and P. Bouché (2013). Unmanned aerial
survey of elephants. PloS one 8:e54700–e54700.

Weissensteiner, M. H., J. W. Poelstra, and J. B. W. Wolf (2015). Low-budget ready-to-fly
unmanned aerial vehicles: an effective tool for evaluating the nesting status of canopy-
breeding bird species. Journal of Avian Biology 46:425–430.

Wildlife Acoustics (2019). Kaleidoscope Pro Analysis Software. [Online.] Available at
https://www.wildlifeacoustics.com/products/kaleidoscope-pro/overview.

Wilson, A. M., J. Barr, and M. Zagorski (2017). The feasibility of counting songbirds using
unmanned aerial vehicles. The Auk 134:350–362.

Wilson, D. R., M. Battiston, J. Brzustowski, and D. J. Mennill (2014). Sound Finder: a new
software approach for localizing animals recorded with a microphone array. Bioacoustics
23:99–112.

Witczuk, J., S. Pagacz, A. Zmarz, and M. Cypel (2017). Exploring the feasibility of unmanned
aerial vehicles and thermal imaging for ungulate surveys in forests - preliminary results.
International Journal of Remote Sensing:1–18.

Figure 1. Methods. Pane **i** shows the setup of each experimental array, **ii** shows the spectrogram of a Yellow Warbler song from recorder C with a black rectangle showing inset in pane **iii,** which shows the measurement of time difference of arrival in Program Raven, and **iv** the estimated positions of the Yellow Warbler over the entire 11-minute experiment, relative to ARUs A, B, C and D.

428 **Figure 2**. Gantt chart showing models used to test whether bird song output is related to the
429 drone's presence. All models include individual bird as a random effect. The null model and
430 model 2 do not include time effects. Time effects of other models shown by numbered
431 sequences. Note that minutes 4 (drone approaching) and 8 (drone departing) of the experiment
432 were not included in the statistical analysis.

Figure 3. Number of individual signing birds detected by minutes and within four radial 25 meter distance bands from the array center.

Figure 4. Calling rate of Northern Cardinals from a single ARU across all 30 arrays. Black line shows the cumulative total of calls and blue circles show GLMM coefficients for each 1-minute interval, relative to minute 1, with 95% confidence limits.

440   **Table 1**. Total number of song bouts detected, and number of bouts localized within 100 m and
441   50 m of the array centers, and number of individual birds within the 50 m radius.

	Total song bout detected	Localized song bouts		Birds within 50-m of array centers
		Within 100-m of array centers	Within 50-m of array centers	
American Robin	1,597	858	331	18
Yellow Warbler	1,643	759	529	42
Common Yellowthroat	734	382	137	13
Field Sparrow	1,388	532	388	31
Song Sparrow	1,661	963	470	42
Indigo Bunting	611	357	241	14
Northern Cardinal	669	294	189	24
all seven species	8,303	4,145	2,285	184

443 **Table 2**. Results of GLMMs for models showing significant effects of the drone on detections of
444 birds within 50 m of the array centers. Estimates (Est) are odd-ratios relative the first time period
445 (either minute 1, or period 1). UL and UL are lower and upper confidence limits, and p-values
446 are from Wald tests.

Yellow Warbler

		Est	LL	UL	P
Intercept		0.665	0.407	1.087	0.103
	Minute 5	0.662	0.297	1.472	0.311
During {	Minute 6	0.438	0.189	1.017	0.054
	Minute 7	0.189	0.066	0.495	0.001
After	Minutes 9-11	0.483	0.271	0.860	0.013

Northern Cardinal

		Est	LL	UL	p
Intercept		0.111	0.050	0.248	<0.001
During	Minutes 5-6	1.462	0.544	3.925	0.450
After	Minutes 9-11	4.888	1.977	12.084	<0.001

448 **Table 3**. Results of GLMMs for models showing significant effects of the drone on singing rates
449 of birds within 50 m of the array centers. Estimates (Est) are odd-ratios relative the first time
450 period (either minute 1, or period 1). UL and UL are lower and upper confidence limits, and p-
451 values are from Wald tests.

American Robin

		Est	LL	UL	P
Intercept		6.124	4.274	8.774	<0.001
During	Minutes 5-7	0.847	0.605	1.188	0.337
After	Minutes 9-11	1.240	0.922	1.669	0.155
distance		0.984	0.972	0.995	0.005

Indigo Bunting

		Est	LL	Est	LL
Intercept		4.846	3.786	6.204	<0.001
During	Minutes 5-7	0.711	0.501	1.009	0.056
After	Minutes 9-11	0.728	0.511	1.037	0.079

Northern Cardinal

			Est	LL	Est	LL
Intercept			3.792	2.494	5.764	<0.001
During	{	Minute 5	0.759	0.351	1.640	0.483
		Minute 6	0.231	0.068	0.783	0.019
		Minute 7	0.942	0.506	1.754	0.851
After		Minutes 9-11	0.782	0.491	1.247	0.302

all seven species

		Est	LL	Est	LL
Intercept		3.736	3.200	4.363	<0.001
During	Minutes 5-7	0.907	0.804	1.024	0.115
After	Minutes 9-11	0.968	0.862	1.087	0.586
distance		0.996	0.991	1.000	0.055

Supplementary Materials

Table S1. AIC values for the eight models for each species, for probability of detection. ΔAIC is compared with the null model (model 1). Gray shading indicates the lowest AIC (or joint lowest if ΔAIC less than 2).

Species	1 (null) AIC	2	3	4	5	6	7	8
				ΔAIC				
American Robin	194.8	1.8	11.1	2.6	6.4	13.0	4.5	8.3
Yellow Warbler	451.5	1.9	-5.3	-6.8	-8.0	-3.4	-4.9	-6.1
Common Yellowthroat	131.8	0.5	11.8	3.0	6.7	12.8	4.0	7.7
Field Sparrow	316.5	1.9	6.6	3.2	6.2	8.5	5.2	8.1
Song Sparrow	400.6	0.2	14.0	3.0	6.9	14.1	3.2	7.1
Indigo Bunting	157.6	1.4	7.9	2.2	5.4	9.3	3.6	6.8
Northern Cardinal	224.1	2.0	-2.7	-11.7	-8.6	-0.7	-9.8	-6.7
all seven species	1898.6	0.7	11.4	1.3	4.1	12.2	2.0	4.8

(Header "model (see figure 2)" spans columns 2–8.)

Table S2. AIC values for the eight models for each species, for singing-rate. ΔAIC is compared with the null model (model 1). Gray shading indicates the lowest AIC (or joint lowest if ΔAIC less than 2).

Species	1 (null) AIC	2	3	4	5	6	7	8
				ΔAIC				
American Robin	264.1	-4.0	7.2	-0.4	1.8	2.5	-5.5	-4.0
Yellow Warbler	492.2	2.5	10.8	2.7	6.5	12.7	4.6	8.5
Common Yellowthroat	146.4	1.6	3.6	3.8	6.0	5.6	5.3	7.7
Field Sparrow	383.3	1.6	15.1	3.6	7.3	16.7	5.2	9.0
Song Sparrow	468.1	2.0	13.7	3.3	6.7	15.7	5.2	8.6
Indigo Bunting	202.5	1.7	10.3	-0.1	3.4	12.2	1.8	5.4
Northern Cardinal	193.0	1.7	6.6	2.3	-0.3	8.5	4.1	1.6
all seven species	2152.2	-1.8	7.7	1.1	4.0	6.0	-0.4	2.5

(The top of the table has the heading "model (see figure 2)" spanning columns 1–8.)