1 **Power analysis of transcriptome-wide association study: implications for**

2 **practical protocol choice**

3 Bowei Ding[1,¶] , Chen Cao[2,¶], Qing Li[2], Devin Kwok[1], Jingjing Wu[1,*], Quan Long[1,2,3,4,*]

4

5 [1]Department of Mathematics & Statistics, University of Calgary, Calgary, Canada.

6 [2]Department of Biochemistry & Molecular Biology, Alberta Children's Hospital Research

7 Institute, University of Calgary, Calgary, Canada.

8 [3]Department of Medical Genetics, University of Calgary, Calgary, Canada.

9 [4]Hotchkiss Brain Institute, O'Brien Institute for Public Health, University of Calgary, Calgary,

10 Canada.

11

12 ¶ = Joint first authors: B.D. and C.C.

13 * = Joint corresponding authors: J.W. (jinwu@ucalgary.ca) and Q.L. (quan.long@ucalgary.ca)

14

15 **Abstract**

16

17 The transcriptome-wide association study (TWAS) has emerged as one of several promising

18 techniques for integrating multi-scale 'omics' data into traditional genome-wide association

19 studies (GWAS). Unlike GWAS, which associates phenotypic variance directly with genetic

20 variants, TWAS uses a reference dataset to train a predictive model for gene expressions, which

21 allows it to associate phenotype with variants through the mediating effect of expressions.

22 Although effective, this core innovation of TWAS is poorly understood, since the predictive

23 accuracy of the genotype-expression model is generally low and further bounded by expression

24 heritability. This raises the question: to what degree does the accuracy of the expression model

25 affect the power of TWAS? Furthermore, would replacing predictions with actual,

26 experimentally determined expressions improve power? To answer these questions, we

27 compared the power of GWAS, TWAS, and a hypothetical protocol utilizing real expression

28 data. We derived non-centrality parameters (NCPs) for linear mixed models (LMMs) to enable

29 closed-form calculations of statistical power that do not rely on specific protocol

30 implementations. We examined two representative scenarios: causality (genotype contributes to

phenotype through expression) and pleiotropy (genotype contributes directly to both phenotype

and expression), and also tested the effects of various properties including expression

heritability. Our analysis reveals two main outcomes: (1) Under pleiotropy, the use of predicted

expressions in TWAS is superior to actual expressions. This explains why TWAS can function

with weak expression models, and shows that TWAS remains relevant even when real

expressions are available. (2) GWAS outperforms TWAS when expression heritability is below a

threshold of 0.04 under causality, or 0.06 under pleiotropy. Analysis of existing publications

suggests that TWAS has been misapplied in place of GWAS, in situations where expression

heritability is low.

Keywords: Power analysis, GWAS, TWAS, Non-centrality parameter, Expression heritability

Author Summary

We compared the effectiveness of three methods for finding genetic effects on disease in order to quantify their strengths and help researchers choose the best protocol for their data. The genome-wide association study (GWAS) is the standard method for identifying how the genetic differences between individuals relate to disease. Recently, the transcriptome-wide association study (TWAS) has improved GWAS by also estimating the effect of each genetic variant on the activity level (or expression) of genes related to disease. The effectiveness of TWAS is surprising because its estimates of gene expressions are very inaccurate, so we ask if a method using real expression data instead of estimates would perform better. Unlike past studies, which only use simulation to compare these methods, we incorporate novel statistical calculations to make our comparisons more accurate and universally applicable. We discover that depending on the type of relationship between genetics, gene expression, and disease, the estimates used by TWAS could be actually more relevant than real gene expressions. We also find that TWAS is not always better than GWAS when the relationship between genetics and expression is weak and identify specific turning points where past studies have incorrectly used TWAS instead of GWAS.

Introduction

High-throughput sequencing instruments have enabled the rapid profiling of transcriptomes (RNA expression of genes) [1-4], proteomes (proteins) [5-7] and other 'omics' data [8-10]. These 'omics' provide insight into the intermediary effects of genotypes on endophenotypes, and can improve the ability of genome-wide association studies (GWAS) to find associations between genetic variants and disease phenotypes. [11-13]. The integration of diverse 'omics' data sources remains a challenging and active field of research [14-17].

One approach to integrating 'omics' and GWAS is the transcriptome-wide association study (TWAS), which quantitatively aggregates multiple genetic variants into a single test using transcriptome data. Pioneered by Gamazon *et al* [18], the TWAS protocol typically has two steps. First, a model is trained to predict gene expressions from local genetic variants near the focal genes, using a reference dataset containing both genotype and expression data. Second, the pretrained model is used to predict expressions from genotypes in the association mapping dataset under study, which contains genotypes and phenotypes (but not expression). The

predicted expressions are then associated to the phenotype of interest. TWAS can also be

conducted with summary statistics from GWAS datasets (i.e. meta-analysis) as first

demonstrated by Gusev *et al.* [19] [20]. TWAS has since achieved significant popularity and

success in identifying the genetic basis of complex traits [21-27], inspiring similar protocols for

other endophenotypes such as IWAS for images [28] and PWAS for proteins [29].

Despite its demonstrated effectiveness, important questions remain regarding the

theoretical conditions under which TWAS is superior to GWAS. **First**: TWAS mapping relies

entirely on predicted expressions, but as shown by many methodological papers, the mean R^2

between predicted and actual expressions is very low (around $0.02 \sim 0.05$). This is in part due to

low expression heritability [18], which bounds the maximum predictive accuracy attainable by

the genotype-expression model. Naturally, one can ask: given sufficiently low expression

heritability, is there is a point at which TWAS performs worse than GWAS? Indeed in real data,

genes discovered with significant TWAS *p*-values tend to have a higher R^2, and thus expression

heritability, than on average [18,19,30-32]. We therefore investigate the effect of expression

heritability on the power of TWAS, as well as its interactions with trait heritability, phenotypic

variance from expressions, number of causal genes, and genetic architecture. **Second**: as

described by Gamazon et al. [18], the key insight of TWAS is that it aggregates sensible genetic

variants to estimate "genetically regulated gene expression", or GReX [18], for use in

downstream GWAS. Given this hypothesis, one may ask if actual expression data would further

improve the power of downstream GWAS over predicted expressions. This is not a trivial

question, as although actual expressions do not suffer from prediction errors, they also include

experimental or environmental noise which masks the genetic component of expression. To test

this problem, we invent a hypothetical protocol associating real expressions to phenotype, which

we call "expression mediated GWAS" or emGWAS. While emGWAS is not in practical use due

to the difficulties of accessing relevant tissues (e.g., in the studies of brain diseases), it can

potentially be applied to future analyses of diseases where tissues are routinely available (e.g.,

blood or cancerous tissues). More importantly, emGWAS serves as a useful benchmark for

evaluating the theoretical properties of TWAS-predicted expressions against ground truth

expression data. By analyzing the power of TWAS, GWAS, and emGWAS, we develop practical

guidelines for choosing each protocol given different expression heritability and genetic

architectures.

While there has been an existing study comparing the power of GWAS, TWAS, and a protocol which integrates eQTLs with GWAS [33], the existing study is purely simulation-based, whereas we determine power directly using traditional closed-form analysis. We derive non-centrality parameters (NCPs) for the relevant statistical tests and the linear mixed model (LMM) in particular (**Methods**). Our derivation uses a novel method to convert an LMM into a linear regression by decorrelating the covariance structure of the LMM response variable (**Methods**). To our best knowledge, this is the first closed-form derivation of the NCP for LMMs in current literature, with potential for broad applications as LMMs are the dominant models used in GWAS and portions of the TWAS pipeline.

Unlike pure simulations, which stochastically resample the alternative hypothesis to estimate statistical power, our closed-form derivation directly calculates power from a particular configuration of association mapping data. As a result, our method saves computational resources, yields more accurate power estimations, and adapts easily to similar protocols such as IWAS [28] and PWAS [29,34]. Moreover, as the closed-form derivation avoids conducting the actual regression, our power calculations do not depend on specific implementations of GWAS and TWAS, which could otherwise cause our results to vary due to differences in filtering inputs or parameter optimizations. Our work therefore characterizes the theoretical power of the protocols across all LMM-based implementations and datasets, although we are unable to account for power losses due to practical implementation issues.

In the following section we describe our novel derivation of NCPs for LMMs and our power analyses of GWAS, TWAS, and emGWAS. We present guidelines on the applicability of each protocol under different input conditions and discuss potential limitations of our approach as well as areas for future research.

Materials & Methods

Mathematical definitions of GWAS, TWAS, and emGWAS protocols

While there are many variations of GWAS and TWAS [18,19,35-39], in this work we assume that multiple genes contribute to phenotypic variation, and for each causal gene, multiple single nucleotide polymorphisms (SNPs) contribute to both gene expression and phenotype. This setting is motivated by the fact that most complex traits are known to have multiple contributing loci, and TWAS fundamentally assumes that genes have multiple local causal variants. To ensure

135 consistency, we apply the same assumptions in the design of the hypothetical protocol

136 emGWAS. Specifically, we define the following models:

137 **GWAS**. For GWAS, we adopted a standard LMM similar to EMMAX [35]

$$Y = \beta_{j0}\mathbf{1} + X_j\beta_{j1} + u + \varepsilon, \qquad j = 1,2,\dots,n_x, \tag{1}$$

where n is the number of individuals, n_x is the total number of genetic variants, Y is an $n \times 1$

vector of phenotypes, $\mathbf{1}$ is an $n \times 1$ vector of ones, X_j is an $n \times 1$ genotype vector with $X_{ij} \in$

$\{0,1,2\}$ representing the number of minor allele copies for the i^{th} individual and j^{th} genetic

variant, β_{j0} and β_{j1} are the intercept and effect size of the genetic variant, u is an $n \times 1$ vector of

random effects following the multivariate normal distribution, i.e. $u \sim N(0, \sigma_g^2 K_x)$, and ε is an

$n \times 1$ vector of errors with $\varepsilon \sim N(0, \sigma_e^2 I)$. In the distributions of u and ε, σ_g^2 and σ_e^2 are their

respective variance components, I is an $n \times n$ identity matrix, and K_x is the genomic relationship

matrix (GRM), which is a known $n \times n$ real symmetric matrix. Following Patterson *et al* [40],

K_x is calculated by

$$K_x = \frac{1}{n_x}\tilde{X}\tilde{X}^T, \tag{2}$$

149 where n_x is the total number of genetic variants and $\tilde{X}$ is a standardized $n \times n_x$ matrix. For

150 example, an element $\tilde{X}_{ij}$ in the j^{th} genetic variant column is calculated as

$$\tilde{X}_{ij} = \frac{X_{ij} - \bar{X}_{.j}}{S_{Xj}}, \tag{3}$$

where $\bar{X}_{.j} = \frac{1}{n}\sum_{i=1}^{n} X_{ij}$ and $S_{Xj}^2 = \frac{1}{n-1}\sum_{i=1}^{n}(X_{ij} - \bar{X}_{.j})^2$ are the sample mean and sample variance

of the j^{th} variant, respectively.

emGWAS. For emGWAS, we first regress the phenotype on the actual (not predicted)

expressions, and then regress the expressions on individual local genetic variants in a similar

manner as a cis-eQTL analysis. We chose the LMM to associate phenotype with expression,

since under the assumption that multiple genes contribute to phenotype, we expect that the

random term of the LMM can capture the effects of non-focal genes. We calculate the GRM

from DNA instead of expressions using the assumption that the ultimate goal is to identify

genetic variants underlying expressions. We chose to use linear regression (LM) to model the

association between expression and local genetic variants (which correspond to cis-eQTLs), as it

is the most common model used in cis-eQTL analyses.

Specifically, the phenotype-expression model is

164
$$Y = \beta_{l0}\mathbf{1} + \beta_{l1}Z_l + u + \varepsilon, \qquad l = 1,2,\dots,n_z, \tag{4}$$

165   where n, Y, $\mathbf{1}$, u and ε have identical interpretations as in the GWAS model from (1), n_z is the

166   total number of genes, Z_l is an $n \times 1$ gene expression vector for the l^{th} gene, and β_{l0} and β_{l1} are

167   the intercept and effect size of the gene.

168   The linear regression associating gene expression with local genetic variants is

169
$$Z_l = \beta_{lk0}\mathbf{1} + \beta_{lk1}X_{lk} + \varepsilon_{el}, \qquad l = 1,2,\dots,n_z, k = 1,2,\dots,n_{el}, \tag{5}$$

170   where X_{lk} is an $n_{el} \times 1$ vector of the k^{th} local genetic variants for the l^{th} gene, $\varepsilon_{el} \sim N(0, \sigma_{el}^2 I)$

171   is a $n \times 1$ vector of errors with variance component σ_{el}^2, n_{el} is the total number of local genetic

172   variants in the l^{th} gene, and β_{lk0} and β_{lk1} are the intercept and effect size of the variant.

173   **TWAS**. For TWAS, we apply an analysis similar to emGWAS, except that gene

174   expressions are predicted using a pretrained elastic-net model. Specifically,

175
$$Y = \beta_{Pl0}\mathbf{1} + \beta_{Pl1}\hat{Z}_l + u + \varepsilon, \qquad l = 1,2,\dots,n_z, \tag{6}$$

where $\hat{Z}_l$ is the altered notation representing an $n \times 1$ vector of predicted gene expressions for

the l^{th} gene, and β_{Pl0} and β_{Pl1} are the intercept and effect size of the predicted gene expression.

There are several methods to estimate gene expression including least absolute shrinkage

and selection operator (LASSO) and elastic-net. Gamazon *et al.* has shown that elastic-net has

good performance and is more robust to minor changes in the input variants [18]. We therefore

use the "glmnet" package in R to train a predictive model using elastic-net. The objective

function in "glmnet" is

183
$$L_{enet}(\beta) = \frac{1}{2n}\|Z - X\beta\|^2 + \lambda\left(\frac{1-\alpha}{2}\|\beta\|^2 + \alpha\|\beta\|_1\right) \tag{7}$$

where λ and α are tuning parameters. The penalty term is a convex (linear) combination of

LASSO and ridge penalties, where $\alpha = 1$ is equivalent to the LASSO objective function, and

$\alpha = 0$ is equivalent to ridge regression. Optimal values of λ and α were chosen by minimizing

the cross-validated squared-error. Readers are referred to **S1 Appendix** for details.

In practice, the specific regression model varies depending on the tool in use. For

example, the leading TWAS tool PrediXcan [18] does not include the random effects of a mixed

model, and many TWAS tools can also analyze summary statistics instead of subject-level

genotypes [19]. The motivation of this work is to reveal the key issues of using gene expressions

as mediations, therefore has to adapt comparable framework. In other works, we do not intend to

compare LMM against linear regression, which will mislead the comparison between GWAS

and TWAS. Since LMMs are dominant in GWAS, we chose LMMs as the underlying model for

all of the protocols we analyze, which allows us to compare them under an equivalent statistical

framework. We believe that LMMs are a sensible approach for TWAS, since the random term

can capture the genetic contributions of non-focal genes.

Closed-form derivation of NCP and power calculation

The non-centrality parameter (NCP) measures the distance between a non-central

distribution and a central distribution under a specific alternative hypothesis. The NCP enables

calculation of the probability of rejecting the null hypothesis, assuming the central distribution,

when the alternative hypothesis is correct. As such, the NCP naturally allows the power of a

statistical test to be determined in a closed form. We have developed the following method to

derive the NCP for LMMs, which we believe is new to the literature.

For a standard simple linear regression, the NCP of a *t*-test of the coefficient of the

predictor variable can be derived similarly to a one-sample *t*-test statistic as follows: if

$X_1, \ldots, X_n \sim N(\mu, \sigma)$ is a simple random sample, then the one-sample *t*-test statistic for evaluating

the null hypothesis $H_0: \mu = \mu_0$ is

$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} = \frac{\sqrt{n}(\bar{X} - \mu_0)/\sigma}{\sqrt{\frac{(n-1)S^2/\sigma^2}{n-1}}} \sim t_{n-1}, \tag{8}$$

211 where $\bar{X}$ and S are the sample mean and (unbiased) sample standard deviation respectively.

212 Under H_0, $\sqrt{n}(\bar{X} - \mu_0)/\sigma \sim N(0,1)$ and $(n-1)S^2/\sigma^2 \sim \chi^2_{n-1}$, and thus $T \sim t_{n-1}$. Under the

213 alternative hypothesis $H_a: \mu = \mu_a$, the test statistic $T = \frac{\sqrt{n}[(\bar{X}-\mu_a)+(\mu_a-\mu_0)]/\sigma}{\sqrt{\frac{(n-1)S^2/\sigma^2}{n-1}}}$ follows a non-

214 central *t* distribution with NCP given by

$$v = \frac{\mu_a - \mu_0}{\sigma/\sqrt{n}} \tag{9}$$

To derive a closed-form NCP for LMMs, we convert the LMM to a linear regression

without intercept by decorrelating the response variable and the predictors, a technique that has

previously been applied to mixed models [41,42]. The procedure is as follows: we first fit the

null model $Y_c = u + \varepsilon$ with no genetic variants, following an existing innovation for reducing the

computational cost of repeatedly factorizing the GRM when analyzing many variants [35,42].

We then estimate σ_g^2 using the Newton-Raphson method detailed in **S2 Appendix**. Denoting the

eigen decomposition of the GRM as $K_x = U_x \Lambda_x U_x^{-1}$, we construct the de-correlation matrix as

$$D_x = \left(\sigma_g^2 \Lambda_x + \sigma_e^2 I\right)^{-\frac{1}{2}} U_x^T. \tag{10}$$

By left multiplying both X and Y by D_x, and denoting $X^* = D_x X = (X_1^*, X_2^*, \ldots, X_n^*)^T$ and

$Y^* = D_x Y = (Y_1^*, Y_2^*, \ldots, Y_n^*)^T$, the covariance structure in Y^* is thus removed and a linear

regression of Y^* on X^* is equivalent to the original LMM model. A proof of the validity of this

decorrelation structure is presented in **S3 Appendix**.

Based on the closed-form NCP for linear regression, we derive the estimated NCP of the

LMM from (1), which is given by

$$\hat{v}_{Gj} = \frac{\sum_{i=1}^n \hat{X}_{ij}^* \hat{Y}_i^* \sum_{i=1}^n \hat{D}_{xi\cdot}^2 - \sum_{i=1}^n \hat{Y}_i^* \hat{D}_{xi\cdot} \sum_{i=1}^n \hat{X}_{ij}^* \hat{D}_{xi\cdot}}{\sqrt{\sum_{i=1}^n (\hat{X}_{ij}^*)^2 \left(\sum_{i=1}^n \hat{D}_{xi\cdot}\right)^2 - \left(\sum_{i=1}^n \hat{D}_{xi\cdot} \hat{X}_{ij}^*\right)^2 \sum_{i=1}^n \hat{D}_{xi\cdot}^2}}, \tag{11}$$

where $\hat{X}_j^* = \hat{D}_x X_j = (\hat{X}_{1j}^*, \hat{X}_{2j}^*, \ldots, \hat{X}_{nj}^*)^T$, $\hat{Y}^* = \hat{D}_x Y = (\hat{Y}_1^*, \hat{Y}_2^*, \ldots, \hat{Y}_n^*)^T$, and $\hat{D}_{xi\cdot} = \sum_{j=1}^n \hat{D}_{xij}$. A

proof of this expression of the NCP for LMMs is in **S4 Appendix.**

The above result allows us to derive the statistical power of the GWAS, emGWAS, and

TWAS protocols. For GWAS, we use the Bonferroni-corrected significance level $\alpha_x = \frac{0.05}{n_x}$ to

account for multiple testing [43], where n_x is the total number of SNPs. Throughout this paper,

we use $f(t; v)$ to denote the probability density function of the non-central t distribution with n-2

degrees of freedom and NCP v. The statistical power of the j^{th} SNP can then be estimated by

$P_{Gj} = \int_{F_0^{-1}(1-\alpha_x)}^{+\infty} f\left(t; \hat{v}_{Gj}\right) dt$ using the estimated NCP $\hat{v}_{Gj}$, where $F_0(t)$ is the cumulative

distribution function of the central t distribution with n-2 degrees of freedom, and $F_0^{-1}(1 - \alpha_x)$

gives the critical value for the central distribution. We directly implement this power

computation in R via the function "pt", which takes the critical value, NCP, and degrees of

freedom as parameters.

For emGWAS, we assume that the two regression models (4) and (5) are independent of

each other. For the model $Y = \beta_{l0} \mathbf{1} + Z_l \beta_{l1} + u + \varepsilon$ from (4), we left multiply the estimated $\hat{D}_x$

to both sides of the equation so that the estimated NCP for the l^{th} gene expression is given by

$$\hat{v}_{eZl} = \frac{\sum_{i=1}^n \hat{Z}_{il}^* \hat{Y}_i^* \sum_{i=1}^n \hat{D}_{xi\cdot}^2 - \sum_{i=1}^n \hat{Y}_i^* \hat{D}_{xi\cdot} \sum_{i=1}^n \hat{Z}_{il}^* \hat{D}_{xi\cdot}}{\sqrt{\sum_{i=1}^n (\hat{Z}_{il}^*)^2 \left(\sum_{i=1}^n \hat{D}_{xi\cdot}^2\right)^2 - \left(\sum_{i=1}^n \hat{D}_{xi\cdot} \hat{Z}_{il}^*\right)^2 \sum_{i=1}^n \hat{D}_{xi\cdot}^2}}, \tag{12}$$

247 where $\hat{Z}_l^* = \widehat{D}_x Z_l = (\hat{Z}_{1l}^*, \hat{Z}_{2l}^*, \dots, \hat{Z}_{nl}^*)^T$. We use the significance level $\alpha_z = \frac{0.05}{n_z}$ for each

248 individual test, where n_z is the total number of genes. The statistical power of detecting the l^{th}

249 gene expression is then estimated by $P_{eZl} = \int_{F_0^{-1}(1-\alpha_z)}^{+\infty} f(t; \hat{v}_{eZl}) dt$. For the model from (5), we

250 simply calculate the estimated NCP of the standard linear regression, which is

$$\hat{v}_{eXlk} = \frac{\sum_{i=1}^n (X_{ilk} - \bar{X}_{\cdot lk}) Z_{il}}{\sqrt{\sum_{i=1}^n (X_{ilk} - \bar{X}_{\cdot lk})^2}\, \hat{\sigma}_{el}}, \tag{13}$$

252 where

$$\hat{\sigma}_{el} = \frac{1}{n-2} \sum_{i=1}^n \left(Z_{il} - \bar{Z}_{\cdot l} + \hat{\beta}_{lk}(X_{ilk} - \bar{X}_{\cdot lk}) \right)^2. \tag{14}$$

254 Again, we use the significance level $\alpha_{el} = \frac{0.05}{n_{el}}$, where n_{el} is the total number of local genetic

255 variants in the l^{th} gene, so that the power of detecting X_{lk} is estimated by $P_{eXlk} =$

256 $\int_{F_0^{-1}(1-\alpha_{el})}^{+\infty} f(t; \hat{v}_{eXlk}) dt$. Since we assume (4) and (5) are independent, the power of detecting

257 the l^{th} gene and the k^{th} variants in the l^{th} gene simultaneously is given by $P_{eZl} P_{eXlk}$.

258   For TWAS, the NCP is estimated in a similar manner as the first step of emGWAS, i.e.

$$\hat{v}_{Tl} = \frac{\sum_{i=1}^n \hat{\hat{Z}}_{il}^* \hat{Y}_i^* \sum_{i=1}^n \widehat{D}_{xi\cdot}^2 - \sum_{i=1}^n \hat{Y}_i^* \widehat{D}_{xi\cdot} \sum_{i=1}^n \hat{\hat{Z}}_{il}^* \widehat{D}_{xi\cdot}}{\sqrt{\sum_{i=1}^n \left(\hat{\hat{Z}}_{il}^*\right)^2 \left(\sum_{i=1}^n \widehat{D}_{xi\cdot}^2\right)^2 - \left(\sum_{i=1}^n \widehat{D}_{xi\cdot} \hat{\hat{Z}}_{il}^*\right)^2 \sum_{i=1}^n \widehat{D}_{xi\cdot}^2}}, \tag{15}$$

where the only difference between (12) and (15) is that $\hat{Z}_{il}^* = \widehat{D}_x Z_{il}$ in (15) is replaced by $\hat{\hat{Z}}_{il}^* =$

$\widehat{D}_x \hat{Z}_{il}$ in (15). The significance level is again $\alpha_z = \frac{0.05}{n_z}$ and the power is estimated by $P_{Tl} =$

$\int_{F_0^{-1}(1-\alpha_z)}^{+\infty} f(t; \hat{v}_{Tl}) dt$.

Simulation of phenotype and expression

As the statistical power of each protocol depends on the magnitude of the genetic effect,

we simulated input data at various effect sizes. While effect size depends on a combination of

many factors, we chose to focus on the following three aspects. 1) We considered two genetic

architectures: causality and pleiotropy (**Fig 1**). In the causality scenario, the contribution of

genotype to phenotype is mediated through expression (**Fig 1a**), whereas in the pleiotropy

scenario, genotype contributes to both expression and phenotype directly (**Fig 1b**). We did not

consider the scenario where phenotype is causal to expression. 2) We considered the strength of

three different variant components: trait heritability (the variance component of phenotype

explained by genotype, denoted $h^2_{x=>y}$), expression heritability (the variance component of

expression explained by genotype, denoted $h^2_{x=>z}$), and the phenotypic variance component

explained by expression, denoted $h^2_{z=>y}$ and abbreviated as *PVX*. 3) We also considered the

number of genes contributing to phenotype and the number of local genetic variants contributing

to expression.

In all our simulations, we use real genotypes from the 1000 Genomes Project (*N* = 2504).

Although there are multiple existing datasets containing both expressions and genotype, we

chose to use simulated expressions instead as it is difficult to match real data exactly to desired

properties such as expression heritability or the number of contributing genetic variants. By

simulating expressions, we can perform a consistent power analysis across a comprehensive

range of prespecified input conditions.

In the causality scenario, phenotypes were simulated with the following procedure. First,

several genes (n_{z-sig}= 4, 9, or 13) were selected as causal genes. For each gene (indexed by $l =$

$1,2,...,n_{z-sig}$), several common and independent genetic variants were selected as causal

variants ($n_{z(l)-sig} = 4 \sim 9$, $MAF > 0.05$, and $R^2 < 0.01$). A linear combination of local variants

in the l^{th} gene is generated to produce the expression values $Z_{(l)}$, and a linear combination of

these gene expressions **Z** is generated as the genomic contribution to phenotype. Note that at

each step, we ensure the simulated linear combinations of variants and expressions match our

desired values for expression heritability $h^2_{x=>z}$ and PVX $h^2_{z=>y}$ (**S5 Appendix**).

In the pleiotropy scenario, we followed a similar procedure except that the phenotype *Y*

was directly generated from a linear combination of genotypes, instead of expressions (**S6**

Appendix). Note that although the expressions *Z* and phenotype *Y* are unrelated by genuine

biological causality, they are generated from the same genetic variants and are therefore

statistically correlated. Therefore, if the trait heritability and expression heritability are

sufficiently large, TWAS can still identify causal genes using the statistical correlation between

genetic variants and expression.

We simulated both scenarios with expression heritability $h^2_{x=>z}$ from the values (2.5%,

3%, 4%, 6%, 8%, 10%, 30%), and with trait heritability $h^2_{x=>y}$ in the pleiotropy scenario or PVX

$h^2_{z=>y}$ in the causality scenario from the values (0.5%, 1%, 2.5%, 5%, 10%). Although we

initially tested more extreme values, our **Results** show that the turning points where TWAS

outperforms GWAS are well within the range of values presented here, and the relative

performance of the protocols remains consistent under more extreme conditions. We therefore

chose to restrict our discussion to the most relevant values for protocol selection, noting that the

expression heritability values we examine are at the high-end of real observed values [18], while

the trait heritability values are lower than typically found in GWAS.

Finally, as each simulation involves multiple variants and genes, the overall power of

each protocol is defined as follows: the power of GWAS is the probability of detecting at least

one causal variant in any causal gene, the power of emGWAS is the probability of detecting at

least one gene and one local SNP of that gene simultaneously, and the power of TWAS is the

probability that at least one predicted gene expression is significant. Specifically,

313
$$P_{GWAS} = 1 - \prod_{j=1}^{n_{x-sig}} \left(1 - P_{G(j)}\right), \tag{16}$$

314
$$P_{emGWAS} = 1 - \prod_{l=1}^{n_{z-sig}} \left(1 - P_{eZ(l)}P_{eX(l)}\right), \; where \; P_{eX(l)} = 1 - \prod_{k=1}^{n_{z(l)-sig}} \left(1 - P_{eX(l)(k)}\right), \tag{17}$$

315
$$P_{TWAS} = 1 - \prod_{l=1}^{n_{z-sig}} \left(1 - P_{T(l)}\right), \tag{18}$$

316 where n_{x-sig}, n_{z-sig} and $n_{z(l)-sig}$ denote the numbers of significant SNPs, genes, and SNPs in

317 the l^{th} significant gene respectively, $G(j)$ denotes the j^{th} significant SNP identified by GWAS,

318 $Z(l)$ and $X(l)(k)$ denote the l^{th} significant gene and the k^{th} significant SNP of the l^{th}

319 significant gene identified by emGWAS, and $T(l)$ denotes the l^{th} significant gene identified by

320 TWAS.

Results

As a quality control measure, we compared the actual expression heritability and the mean R^2 of the predicted expressions (**Table 1**). As expected, the mean R^2 grows closer to the actual heritability value as expression heritability increases.

Table 1: Comparisons of R^2 of imputed gene expression under different levels of expression heritability and number of genetic variants.

	Mean of R^2	Sample Standard Deviation of R^2
$h_1^2 = 0.025$	0.007847616	0.007415877
$h_1^2 = 0.03$	0.01259302	0.008410582
$h_1^2 = 0.04$	0.02319834	0.009481371
$h_1^2 = 0.06$	0.04415579	0.01083593
$h_1^2 = 0.08$	0.06465895	0.01175991
$h_1^2 = 0.1$	0.08518152	0.01264175
$h_1^2 = 0.3$	0.2886779	0.01514781

Causality scenario

We first analyzed cases where expression heritability is high ($h_{x=>z}^2 = 0.1$ or 0.3) but the PVX is low (**Fig 2**). Overall, emGWAS clearly outperforms both GWAS and TWAS by a large margin, and TWAS also generally outperforms GWAS. Note that although the PVX is low and favors GWAS, TWAS is still more powerful due to the high expression heritability, which shows that expression heritability affects the performance of TWAS more than the PVX. Consistent with intuition, we observed that GWAS and TWAS have higher power as expression heritability increases, whereas this increase is much smaller for emGWAS. The power of GWAS and emGWAS reduces as the number of causal genes grows, whereas TWAS is largely unaffected by the number of causal genes. This is also consistent with intuition since TWAS uses GReX ($\hat{Z}$) to aggregate genetic effects, avoiding the burden of multiple-testing correction.

We then analyzed cases where the PVX is high, but expression heritability is relatively

low ($h^2_{x=>z}$= 0.025, 0.03, 0.04 or 0.08). Evidently, emGWAS performs best with powers

consistently at 1.0. The comparison between TWAS and GWAS is more nuanced, as TWAS is

suboptimal to GWAS when the expression heritability is 0.025 or 0.03 (**Figs 3a and 3b**), begins

to outperform GWAS when the expression heritability is 0.04 (**Fig 3c**), and clearly outperforms

GWAS when the expression heritability is 0.08 (**Fig 3d**). This quantifies an important turning

point in that GWAS is superior to TWAS when expression heritability is less than 0.04, even if

PVX is high (favoring TWAS).

Pleiotropy scenario

Again, we first analyze cases where expression heritability is high and trait heritability is

low (**Fig 4**). Unlike in the causality scenario, the power of emGWAS is very low compared to

TWAS and GWAS. A potential explanation is that when the effect of genetic variants on

phenotype is not mediated through expressions, the non-genetic effects within the actual

expressions add noise to emGWAS predictions. In contrast, the elastic-net model in TWAS

captures only the genetic component of expressions, meaning the predicted expressions are a

more accurate model of the direct genetic effect on phenotype. While errors are unavoidable in

the elastic-net training process (as revealed in **Table 1**), our results show that the loss of power

due to non-genetic effects is overwhelmingly greater than the loss due to training errors. As in

the casualty scenario, TWAS generally outperforms GWAS except in the case where trait

heritability is extremely low and the number of contributing genes is large, which is rare in

practice. We therefore conclude that in both scenarios, TWAS has better power than GWAS

when expression heritability is high.

We finally analyze cases where expression heritability is low but trait heritability is high.

Here, emGWAS continues to be the least powerful of the three protocols. As in the causality

scenario, we again observe a turning point where TWAS outperforms GWAS: TWAS has lower

power than GWAS when the expression heritability is 0.025 or 0.04 (**Figs 5a and 5b**), TWAS

has comparable power when the expression heritability is 0.06 (**Fig 5c**), and TWAS outperforms

GWAS when the expression heritability is 0.08 (**Fig 5d**).

Our results can be summarized in two observations (**Fig 6**). First, emGWAS outperforms

TWAS and GWAS in the casualty scenario, but is less powerful in the pleiotropy scenario

regardless of the accuracy of the predicted expressions (**Table 1**). This demonstrates that when

non-genetic components in expression do not contribute to phenotype (i.e. pleiotropy scenario),

predicted expressions capture genetic contributions better than actual expressions (which include

non-genetic components). Second, the turning point at which traditional GWAS outperforms

TWAS is an expression heritability of less than 0.04 in the causality scenario, or 0.06 in the

pleiotropy scenario.

These turning points are immediately relevant to the practical conduct of association

mapping studies, as shown by the following analysis of expression heritability in existing TWAS

publications. As few publications disclose their estimated expression heritability, we use

published R^2 values of the correlation between predicted and actual expressions to approximate

the underlying expression heritability. We use the difference between expression heritability and

R^2 as calculated from our simulations (**Table 1**) to map these R^2 values to an estimated

expression heritability (i.e. R^2 of 0.023 and 0.044 give expression heritability values 0.04 and

0.06, respectively), although in practice the true difference may vary depending on the predictive

model used in each study. Table 1 of the PrediXcan publication lists significant results from their

paper, in which 14 out of 41 discovered genes have R^2 values less than 0.044, with 2 values less

than 0.023. Additionally, our review of recent TWAS publications shows that most of the genes

presented have mean R^2 values less than 0.044 or 0.023 (**Table 2**). As our power analysis

indicated, GWAS may have better power than TWAS given these low expression heritability

conditions. Although we are unable to determine if the genes discovered by these publications

follow the causality or pleiotropy scenario, other advanced statistical models [44] may be used to

determine appropriate thresholds to distinguish between pleiotropy and causality.

In summary, we suggest the following modifications to the TWAS protocol. First, one

may estimate expression heritability in the reference panel and filter out genes with expression

heritability less than 0.04. Second, after conducting TWAS association mapping, determine the

underlying causality scenario (causality or pleiotropy) in order to choose an appropriate

expression heritability threshold (0.04 or 0.06). Finally, conduct GWAS for each gene with an

expression heritability below the given threshold.

Table 2: Mean R^2 in published TWAS projects.

Title of the publication	Description of prediction accuracy
Large-scale transcriptome-wide association study identifies new prostate cancer risk regions [22]	The mean R^2 = 0.07 for measured and predicted gene expression for TCGA normal prostate samples using models fitted in GTEx normal prostate.
A framework for transcriptome-wide association studies in breast cancer in diverse study populations [45]	The median CV R^2 for the 153 genes is 0.011 in both African American and white women.
Evaluation of PrediXcan for prioritizing GWAS associations and predicting gene Expression [46]	The average of prediction accuracy (R^2) is 0.023 for the DGN model and 0.02 for the GTEx model, with both using whole blood model.
A gene-based association method for mapping traits using reference transcriptome data [18]	The average prediction R^2 value is 0.0197 for GEUVADIS LCLs. For GTEx tissues, the prediction R^2 values are 0.0367 (adipose), 0.0358 (tibial artery), 0.0356 (left-ventricular heart), 0.0359 (lung), 0.0269 (muscle), 0.0422 (tibial nerve), 0.0374 (sun-exposed skin), 0.0398 (thyroid) and 0.0458 (whole blood).

Discussion

In this work, we produced a novel derivation of the NCP for LMMs based on the decorrelation procedure, allowing us to calculate closed-form estimates of statistical power for three protocols: GWAS, emGWAS, and TWAS. Our power analysis revealed two practical insights. **First**, in the pleiotropy scenario, the use of predicted expressions in TWAS is overwhelmingly more powerful than the use of actual expressions in emGWAS, regardless of the accuracy of the predicted expressions *per se* (**Table 1**). This suggests that even if real expressions can be experimentally determined, TWAS is still superior for the analysis of some genes. While this appears counterintuitive, in statistical terms it is a direct result of the lack of a causal relationship between expression and phenotype under pleiotropy. This result reinforces

the key insight, as presented by some publications [18], that TWAS uses expression as an

objective function to select a linear combination of genetic variants, rather than attempting to

accurately predict expressions. We note that this is equivalent to denoising in the field of

machine learning [47]. **Second**, expression heritability determines the relative power of TWAS

and GWAS. When the expression heritability is lower than 0.04 (in the casualty scenario) or 0.06

(in the pleiotropy scenario), GWAS outperforms TWAS despite not utilizing gene expression

information. This suggests that in practice, TWAS may often be suboptimal when expression

heritability is low (**Table 2** & Table 1 in [18]), which can be mitigated by choosing the optimal

association mapping protocol according to this work's quantitative guidelines.

A recent publication has also compared the statistical powers of GWAS and TWAS using

pure simulations [33]. However, since we calculate power from a closed-form NCP derivation,

our work establishes theoretical benchmarks for the performance of each protocol, independent

of their implementations. Our work also has a different focus: rather than comparing techniques

for training the genotype-expression predictive model and the impact of the actual number of

causal genetic variants, we rank the effectiveness of GWAS, TWAS and emGWAS to better

guide the practical application of TWAS. We analyze the theoretical effectiveness of real

expressions as utilized by emGWAS, but exclude the protocol eGWAS as analyzed in [33],

which uses eQTLs to assist association mapping. Our conclusions also differ slightly, as while

the previous publication highlighted the importance of expression heritability, they concluded

that expression heritability affects power only under the causality scenario, and not pleiotropy. In

contrast, we concluded that expression heritability affects both scenarios.

Finally, our closed-form derivation is readily adaptable to other methods utilizing middle

'omics' (endophenotypes) such as IWAS [28] and PWAS [29,34]. In fact, the variable Z in

formula (15) can already represent such data as images or proteins, and thus no further

modifications of the NCPs are necessary to adapt this work.

There are several limitations in the present study. Although our closed-form derivation is

easily adaptable and works independently of specific implementations, it is unable to capture

power loss due to implementation limitations or bias in specific datasets. Additionally, closed-

form derivations are more sensitive to model assumptions than simulation-based methods. Our

calculation of the NCP also requires the variance component σ_g^2 to be estimated from data, in

order to form the decorrelation matrix D_x. Although this approximation introduces extra

variability and may therefore cause a decrease in power, we have omitted this variability from

our analyses as the estimation of σ_g^2 is generally well-established, and has high accuracy in

practice when given thousands of samples. Finally, we only compared linear models for GWAS

and TWAS. As a future work, we may explore kernel-based nonparametric and semiparametric

methods for conducting both GWAS [48,49] and TWAS [50].

Acknowledgements

J.W. is supported by an NSERC Discovery Grant (RGPIN-2018-04328). Q.L. is supported by an

NSERC Discovery Grant (RGPIN-2017-04860), a Canada Foundation for Innovation JELF grant

(36605), a New Frontiers in Research Fund (NFRFE-2018-00748), a Clinical Research Fund and

a Startup grant supported by Alberta Children's Hospital Research Institute (ACHRI). C.C. is

supported by ACHRI postdoctoral scholarship.

References

1. Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, et al. (2010) Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. Nat Biotechnol 28: 511-515.
2. Wang Z, Gerstein M, Snyder M (2009) RNA-Seq: a revolutionary tool for transcriptomics. Nat Rev Genet 10: 57-63.
3. McGettigan PA (2013) Transcriptomics in the RNA-seq era. Curr Opin Chem Biol 17: 4-11.
4. Ozsolak F, Milos PM (2011) RNA sequencing: advances, challenges and opportunities. Nat Rev Genet 12: 87-98.
5. Selevsek N, Chang CY, Gillet LC, Navarro P, Bernhardt OM, et al. (2015) Reproducible and consistent quantification of the Saccharomyces cerevisiae proteome by SWATH-mass spectrometry. Mol Cell Proteomics 14: 739-749.
6. Pible O, Armengaud J (2015) Improving the quality of genome, protein sequence, and taxonomy databases: a prerequisite for microbiome meta-omics 2.0. Proteomics 15: 3418-3423.
7. Bell AW, Deutsch EW, Au CE, Kearney RE, Beavis R, et al. (2009) A HUPO test sample study reveals common problems in mass spectrometry-based proteomics. Nat Methods 6: 423-430.
8. Zhang A, Sun H, Wang P, Han Y, Wang X (2012) Modern analytical techniques in metabolomics analysis. Analyst 137: 293-300.
9. Coats VC, Rumpho ME (2014) The rhizosphere microbiota of plant invaders: an overview of recent advances in the microbiomics of invasive plants. Front Microbiol 5: 368.
10. Teperino R, Lempradl A, Pospisilik JA (2013) Bridging epigenomics and complex disease: the basics. Cell Mol Life Sci 70: 1609-1621.
11. Klein RJ, Zeiss C, Chew EY, Tsai JY, Sackler RS, et al. (2005) Complement factor H polymorphism in age-related macular degeneration. Science 308: 385-389.
12. Ozaki K, Ohnishi Y, Iida A, Sekine A, Yamada R, et al. (2002) Functional SNPs in the lymphotoxin-alpha gene that are associated with susceptibility to myocardial infarction. Nat Genet 32: 650-654.
13. Mills MC, Rahal C (2019) A scientometric review of genome-wide association studies. Commun Biol 2: 9.
14. Eddy S, Mariani LH, Kretzler M (2020) Integrated multi-omics approaches to improve classification of chronic kidney disease. Nat Rev Nephrol.
15. Hasin Y, Seldin M, Lusis A (2017) Multi-omics approaches to disease. Genome Biol 18: 83.
16. Yan J, Risacher SL, Shen L, Saykin AJ (2018) Network approaches to systems biology analysis of complex disease: integrative methods for multi-omics data. Brief Bioinform 19: 1370-1381.
17. Fukushima A, Kusano M, Redestig H, Arita M, Saito K (2009) Integrated omics approaches in plant systems biology. Curr Opin Chem Biol 13: 532-538.
18. Gamazon ER, Wheeler HE, Shah KP, Mozaffari SV, Aquino-Michaels K, et al. (2015) A gene-based association method for mapping traits using reference transcriptome data. Nat Genet 47: 1091-1098.
19. Gusev A, Ko A, Shi H, Bhatia G, Chung W, et al. (2016) Integrative approaches for large-scale transcriptome-wide association studies. Nat Genet 48: 245-252.
20. Barbeira AN, Dickinson SP, Bonazzola R, Zheng J, Wheeler HE, et al. (2018) Exploring the phenotypic consequences of tissue specific gene expression variation inferred from GWAS summary statistics. Nat Commun 9: 1825.
21. Gusev A, Mancuso N, Won H, Kousi M, Finucane HK, et al. (2018) Transcriptome-wide association study of schizophrenia and chromatin activity yields mechanistic disease insights. Nat Genet 50: 538-548.

22. Mancuso N, Gayther S, Gusev A, Zheng W, Penney KL, et al. (2018) Large-scale
transcriptome-wide association study identifies new prostate cancer risk regions. Nat
Commun 9: 4079.
23. Theriault S, Gaudreault N, Lamontagne M, Rosa M, Boulanger MC, et al. (2018) A
transcriptome-wide association study identifies PALMD as a susceptibility gene for
calcific aortic valve stenosis. Nat Commun 9: 988.
24. Gong L, Zhang D, Lei Y, Qian Y, Tan X, et al. (2018) Transcriptome-wide association study
identifies multiple genes and pathways associated with pancreatic cancer. Cancer Med
7: 5727-5732.
25. Ratnapriya R, Sosina OA, Starostik MR, Kwicklis M, Kapphahn RJ, et al. (2019) Retinal
transcriptome and eQTL analyses identify genes associated with age-related macular
degeneration. Nat Genet 51: 606-610.
26. Atkins I, Kinnersley B, Ostrom QT, Labreche K, Il'yasova D, et al. (2019) Transcriptome-
Wide Association Study Identifies New Candidate Susceptibility Genes for Glioma.
Cancer Res 79: 2065-2071.
27. Zhang W, Voloudakis G, Rajagopal VM, Readhead B, Dudley JT, et al. (2019) Integrative
transcriptome imputation reveals tissue-specific and shared biological mechanisms
mediating susceptibility to complex traits. Nat Commun 10: 3834.
28. Xu Z, Wu C, Pan W, Alzheimer's Disease Neuroimaging I (2017) Imaging-wide association
study: Integrating imaging endophenotypes in GWAS. Neuroimage 159: 159-169.
29. Brandes N, Linial N, Linial M. PWAS: Proteome-Wide Association Study. Research in
Computational Molecular Biology; 2020; Cham. Springer International Publishing. pp.
237-239.
30. Mancuso N, Shi H, Goddard P, Kichaev G, Gusev A, et al. (2017) Integrating Gene
Expression with Summary Association Statistics to Identify Genes Associated with 30
Complex Traits. Am J Hum Genet 100: 473-487.
31. Wainberg M, Sinnott-Armstrong N, Mancuso N, Barbeira AN, Knowles DA, et al. (2019)
Opportunities and challenges for transcriptome-wide association studies. Nat Genet 51:
592-599.
32. Mogil LS, Andaleon A, Badalamenti A, Dickinson SP, Guo X, et al. (2018) Genetic
architecture of gene expression traits across diverse populations. PLoS Genet 14:
e1007586.
33. Veturi Y, Ritchie MD (2018) How powerful are summary-based methods for identifying
expression-trait associations under different genetic architectures? Pac Symp Biocomput
23: 228-239.
34. Okada H, Ebhardt HA, Vonesch SC, Aebersold R, Hafen E (2016) Proteome-wide
association studies identify biochemical modules associated with a wing-size phenotype
in Drosophila melanogaster. Nat Commun 7: 12649.
35. Kang HM, Sul JH, Service SK, Zaitlen NA, Kong SY, et al. (2010) Variance component
model to account for sample structure in genome-wide association studies. Nat Genet
42: 348-354.
36. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, et al. (2007) PLINK: a tool set
for whole-genome association and population-based linkage analyses. Am J Hum Genet
81: 559-575.
37. Aulchenko YS, Ripke S, Isaacs A, van Duijn CM (2007) GenABEL: an R library for genome-
wide association analysis. Bioinformatics 23: 1294-1296.
38. Gogarten SM, Bhangale T, Conomos MP, Laurie CA, McHugh CP, et al. (2012)
GWASTools: an R/Bioconductor package for quality control and analysis of genome-
wide association studies. Bioinformatics 28: 3329-3331.
39. Zhou X, Stephens M (2012) Genome-wide efficient mixed-model analysis for association
studies. Nat Genet 44: 821-824.

40. Patterson N, Price AL, Reich D (2006) Population structure and eigenanalysis. PLoS Genet
562   2: e190.
563   41. Rakitsch B, Lippert C, Stegle O, Borgwardt K (2013) A Lasso multi-marker mixed model for
564   association mapping with population structure correction. Bioinformatics 29: 206-214.
565   42. Long Q, Zhang Q, Vilhjalmsson BJ, Forai P, Seren U, et al. (2013) JAWAMix5: an out-of-
566   core HDF5-based java implementation of whole-genome association studies using
567   mixed models. Bioinformatics 29: 1220-1222.
568   43. Shaffer JP (1995) Multiple hypothesis testing. Annual review of psychology 46: 561-584.
569   44. Schadt EE, Lamb J, Yang X, Zhu J, Edwards S, et al. (2005) An integrative genomics
570   approach to infer causal associations between gene expression and disease. Nat Genet
571   37: 710-717.
572   45. Bhattacharya A, Garcia-Closas M, Olshan AF, Perou CM, Troester MA, et al. (2020) A
573   framework for transcriptome-wide association studies in breast cancer in diverse study
574   populations. Genome Biol 21: 42.
575   46. Li B, Verma SS, Veturi YC, Verma A, Bradford Y, et al. (2018) Evaluation of PrediXcan for
576   prioritizing GWAS associations and predicting gene expression. Pac Symp Biocomput
577   23: 448-459.
578   47. Tian C, Fei L, Zheng W, Xu Y, Zuo W, et al. (2019) Deep learning on image denoising: An
579   overview. arXiv preprint arXiv:191213171.
580   48. Wu MC, Lee S, Cai T, Li Y, Boehnke M, et al. (2011) Rare-variant association testing for
581   sequencing data with the sequence kernel association test. Am J Hum Genet 89: 82-93.
582   49. Wu MC, Kraft P, Epstein MP, Taylor DM, Chanock SJ, et al. (2010) Powerful SNP-set
583   analysis for case-control genome-wide association studies. Am J Hum Genet 86: 929-
584   942.
585   50. Cao C, Kwok D, Edie S, Li Q, Ding B, et al. (2020) kTWAS: integrating kernel-machine with
586   transcriptome-wide association studies improves statistical power and reveals novel
587   genes. bioRxiv.
588

589 **Figures**

591 **Fig 1: Causality (a) and Pleiotropy (b) scenarios for genotype (X), expression (Z) and**
592 **phenotype (Y).**

Fig 2: Causality scenario when expression heritability is high and PVX is low.

The PVX is 0.005, 0.01, 0.025, and 0.05 in the four columns as indicated by the X-axis labels.

The number of genes contributing to phenotype for (a), (b) and (c) are 4, 9, and 13 respectively.

The expression heritability for the top and bottom rows of (a), (b) and (c) are 0.1 and 0.3

respectively. The number of causal variants per gene is randomly sampled from the interval

[4,9].

Fig 3: Causality scenario when expression heritability is low and PVX is high.

The PVX is 0.05 and 0.1 in the two columns as indicated by the X-axis labels. The numbers of genes contributing to phenotype in the left, middle and right panels are 4, 9, and 13 respectively. The expression heritability levels in (a), (b), (c) and (d) are 0.025, 0.03, 0.04, and 0.08 respectively. The number of causal variants per gene is randomly sampled from the interval [4,9].

Fig 4: Pleiotropy scenario when expression heritability is high and trait heritability is low.

The trait heritability is 0.005, 0.01, 0.025, and 0.05 in the four columns as indicated by the X-axis labels. The numbers of genes contributing to phenotype for (a), (b) and (c) are 4, 9, and 13 respectively. The expression heritability for the top and bottom rows of (a), (b) and (c) are 0.1 and 0.3 respectively. The number of causal variants per gene is randomly sampled from the interval [4,9].

Fig 5: Pleiotropy scenario when expression heritability is low and trait heritability is high.

616 The PVX is 0.05 and 0.1 in the two columns as indicated by the X-axis labels. The numbers of

617 genes contributing to phenotype for the left, middle and right panels are 4, 9, and 13 respectively.

618 The expression heritability levels in (a), (b), (c) and (d) are 0.025, 0.04, 0.06, and 0.08

619 respectively. The number of causal variants per gene is randomly sampled from the interval

620 [4,9].

Fig 6: Overall comparison between emGWAS, GWAS and TWAS.