Phylogenetic diversity analysis of shotgun metagenomic reads describes gut microbiome development and treatment effects in the post-weaned pig

Daniela Gaio[1], Matthew Z. DeMaere[1], Kay Anantanawat[1], Graeme J Eamens[2], Tiziana Zingali[1], Linda Falconer[2], Toni A. Chapman[2], Steven Djordjevic[1], Aaron E. Darling[1]*

[1]The iThree institute, University of Technology Sydney, Sydney, NSW, Australia

[2]NSW Department of Primary Industries, Elizabeth Macarthur Agricultural Institute, Menangle, NSW, Australia

Corresponding author:

Daniela Gaio[1]

The iThree institute, University of Technology Sydney, P.O. Box 123, Broadway, NSW, 2007, Australia

d.gaio@outlook.com

Abstract

Background: Intensive farming practices can increase exposure of animals to infectious agents against which antibiotics are used. Besides leading to antimicrobial resistance (AMR), orally administered antibiotics are well known to cause dysbiosis. To counteract dysbiotic effects, numerous studies in the past two decades sought to understand whether probiotics are a valid tool to help re-establish a healthy gut microbial community after antibiotic treatment. However, although dysbiotic effects of antibiotics are well investigated, little is known about the effects of intramuscular antibiotic treatment on the gut microbiome and a few studies attempted to study treatment effects using phylogenetic diversity analysis techniques. In this study we sought to determine the effects of two probiotic- and one intramuscularly administered antibiotic treatment on the developing gut microbiome of post-weaning piglets between their 3rd and 9th week of life.

Methods: Shotgun metagenomic sequences from over 800 faecal time-series samples derived

from 126 piglets and 42 sows were analysed in a phylogenetic framework to characterise the

developing gut microbial community composition of post-weaning piglets. We assessed the

effects of intramuscular antibiotic treatment and probiotic oral treatment on the diversity of these

gut microbial communities using alpha and beta diversity measures.

Results: Differences between individual hosts such as breed, litter, and age, were found to be

important contributors to variation in the community composition. Host age was the dominant

factor in shaping the gut microbiota of piglets after weaning. The post-weaning pig gut

microbiome appeared to follow a highly structured developmental program with characteristic

post-weaning changes that can distinguish hosts that were born as little as two days apart in the

second month of life. Treatment effects of the antibiotic and probiotic treatments were found but

were subtle and included a higher representation of *Mollicutes* associated with intramuscular

antibiotic treatment, and an increase of *Lactobacillus* associated with probiotic treatment.

Discussion: The discovery of correlations between experimental factors and microbial

community composition is more commonly addressed with OTU-based methods and rarely

analysed via phylogenetic diversity measures. The latter method, although less intuitive than the

former, suffers less from library size normalization biases, and it proved to be instrumental in

this study for the discovery of correlations between microbiome composition and host-, and

treatment factors.

Introduction

As the world population grows, there is an accompanying demand for animal-derived products.

Intensive animal husbandry and early weaning practices are commonly used to maximize

production rates while minimizing costs. In a semi-natural environment, pig weaning occurs

between the 12th and the 17th week following birth whereas in intensively farmed pigs it typically

occurs at 3-4 weeks of age [1,2]. In intensive pig production early weaning practices increase the

risk of enteric infections [3–6], and thereby the need for antimicrobial strategies, which has

included the common metaphylactic use of antibiotics [7,8]. Besides contributing to the concerning

issue of antimicrobial resistance (AMR) build up, antibiotic usage causes dysbiosis [9–11], a

disruption of a balanced state within a gut microbial community, which increases the chance of

pathogens gaining a foothold (*i.e.*: colonization of the host or overgrowth) [12,13]. Researchers

sought to determine the gut microbial modulatory effects of oral and in-feed antibiotic treatment

in the pig [10,11], mouse [14–16], and human [17], and only a few studies report the effects of

intramuscular (IM) antibiotic treatment on the gut microbial community of the pig [18,19].

Dysbiosis following antibiotic treatment of the young pig via intramuscular (IM) [20] and

subcutaneous [9] routes has been documented. It has been reported that a single dose of

intramuscularly administered amoxicillin in pigs, immediately after birth, causes long lasting

dysbiosis [21], but no studies, to our knowledge, assessed the effects of IM neomycin treatment on

the post-weaning pig gut microbiota. IM antibiotic use has a lower risk for inducing AMR than

oral antibiotics in livestock production [22–25], and according to a national survey of antibiotic use

in 197 large Australian commercial pig herds, 9.6% reported the use of injectable

apramycin/neomycin [7]. For these reasons it is relevant to the industry to assess if, and to what

extent, IM neomycin causes dysbiosis.

Probiotics may play a role in counteracting the dysbiotic effects of antibiotics. Probiotic

treatment has been reported to affect the physiology of the host by improving mucosal integrity

[26–29], inducing competitive exclusion against pathogenic species [30–35], reducing intestinal

inflammation [26,36,37], and pathogen translocation [26,38,39]. In terms of microbial community

modulating potential of probiotics, studies in swine using culture based methods reported the

reduction of pathogenic species [32,34,40–48] and an increase of *Lactobacillus* species [43,47–49]

following the administration of probiotic formulas. However, only in the past decade have

advances in and higher accessibility to high-throughput sequencing techniques enabled

researchers to better characterize microbial dynamics following probiotic treatment. Several of

these studies used the amplification of the bacterial 16S rDNA V4 region, determining the

effects of probiotics on the gut microbial community of swine [50–53]. The choice of shotgun

metagenomic sequencing over amplicon sequencing, in this study, was dictated by the use of this

dataset outside of the scope described in this manuscript. In this study we assessed the effects of

two probiotic formulas on the gut microbial community of post-weaning piglets undergoing a

two-weeks probiotic treatment immediately after weaning. To assess the effects on the microbial

community of the administration of probiotic following antibiotic treatment, two treatment

groups were administered IM neomycin treatment followed by two weeks of either of the two

probiotic formulae.

The data was analysed from the perspective of the phylogenetic diversity of microbial

communities. Firstly, we describe the phylogenetic diversity of microbial communities, and

secondly, the role of age, breed and litter in microbial community composition. Finally, we

describe the temporal effects on microbial communities, and the associated effects of IM

neomycin and probiotic treatments.

Materials & Methods

Animal trial

Metagenomic samples were derived from a pig study conducted at the Elizabeth Macarthur

Agricultural Institute (EMAI) NSW, Australia, and approved by the EMAI Ethics Committee

(Approval M16/04). Below, we briefly summarise the origin of the samples, with comprehensive

details on the animal trial and sample workflow being described previously [54].

Post-weaning piglets (*n=126*) derived from a large commercial pig herd were transferred to

the study facility in January 2017. Piglets, aged 22.5±2.5 days at the start of the trial, consisted of

4 main cross-breed types: "Duroc × Landrace" (*n=46*), "Duroc × Large White" (*n=59*),

"Landrace × cross bred (LW×D)" (*n=9*), "Large White × Duroc" (*n=12*); and three pig lines

(line 319: *n=9*; line 316: *n=46*; line 326: *n=71*) [54]. The study facility consisted of 4

environmentally controlled rooms (Rooms 1–4) with air conditioning, concrete slatted block

flooring with underground drainage, and open rung steel pens. The floor was swept daily and the

under-floor drainage was flushed twice weekly. A rubber ball was added to each pen for

environmental enrichment. Piglets were fed *ad libitum,* a commercial pig grower mix of 17.95%

protein, free of antibiotics.

The piglets were distributed over 6 treatment cohorts: a placebo group (Control *n=29*); two

probiotic groups (D-Scour *n=18*; ColiGuard *n=18*); one antibiotic group (Neomycin *n=24*) and

two antibiotic-then-probiotic treatment groups (Neomycin+D-Scour *n=18*;

Neomycin+ColiGuard *n=18*). The commercial probiotic paste preparation D-Scour™ was

sourced from International Animal Health, Australasia, and is composed of 1.8×10^8 CFU/g of

each of the following: *Lactobacillus acidophilus, Lactobacillus*

delbrueckii subspecies *bulgaricus*, *Lactobacillus plantarum*, *Lactobacillus rhamnosus*,

Bifidobacterium bifidum, *E. faecium*, and *Streptococcus salivarius* subspecies *thermophilus*, with

an additional 20mg/g of garlic extract (*Allium sativum*). ColiGuard® is a probiotic formulation

developed for the treatment of entero-toxigenic *Escherichia coli* (ETEC) in weaner pigs,

developed in collaboration between the NSW DPI and International Animal Health Product, and

contains undefined concentrations of: *L. plantarum* and *Lactobacillus salivarius*. In addition, an

in-house developed mock community was used as a positive control, composed of *Bacillus*

subtilis strain 168, *Enterococcus faecium*, *Staphylococcus aureus* ATCC25923, *Staphylococcus*

epidermidis ATCC35983, *Enterobacter hormaechei* CP_032842, *Escherichia coli* K-12

MG1655, and *Pseudomonas aeruginosa* PAO1, in the following proportions:

8.7:13.0:7.7:16.7:38.9:14.5:0.4.

Groups of piglets were euthanized at the start (t0: *n*=6), a week after (t2: *n*=12), two weeks

after the start of the trial (t4: *n*=36) and at the end of the trial (t10: *n*=72) to obtain biopsy

samples that were used in another study. All surviving animals were euthanized at the end of the

animal trial. Euthanasia occurred as follows: the piglets were restrained in dorsal recumbency in

a bleeding cradle and pentobartitone euthanasia solution (Lethabarb, Virbac Australia, 325 mg

pentobarbitone sodium/mL) diluted 1:1 in sterile normal saline was administered via the precava

using a 20-21G needle depending on pig weight. The piglets received a dose of approximately 30

mg pentobarbitone/Kg to achieve deep anaesthesia, and then immediately exsanguinated.

Exsanguination after a non-lethal dose of barbiturate was undertaken to reduce excessive

congestion of the visceral blood vessels at the time of intestinal specimen collection (which is

likely to occur if euthanasia is undertaken using an overdose of barbiturate alone).

Faecal sampling occurred twice weekly for a subset of the piglets (8 per cohort; *n*=48), while

it occurred weekly for all the living piglets (*n*=126; see euthanasia groups described above). In

addition, 42 faecal samples derived from the piglets' mothers (*n*=42), 18 samples derived from

three distinct positive controls (D-Scour™, ColiGuard®, and mock community), and 20 negative

controls were included.

A thorough description of the animal trial, and the metadata containing behaviour, weight,

and faecal consistency scores recorded over the 6-week period of the trial, is available in our

previous work [54].

Metagenomic samples processing

Samples underwent homogenization and storage at -80 ºC. After sample thawing, DNA extraction was performed with the PowerMicrobiome DNA/RNA EP kit (Qiagen), and libraries were prepared using the Hackflex method [55]. Sequencing was performed on three Illumina NovaSeq S4 flow cells, after library normalization and pooling. The data is deposited to the NCBI Short Read Archive under project PRJNA526405 and http://dx.doi.org/10.5524/100890 [54]. Samples were assessed for quality using FASTQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and a combined report of all samples was obtained with MULTIQC [56] (available in our GitHub repository https://github.com/GaioTransposon/metapigs_phylodiv).

Determination of microbial diversity among samples

Phylogenetic diversity of all samples was assessed with PhyloSift [57] using the first 1M read pairs of each sample (parameters: `--chunk-size 1000000 --paired`) (script: *phylosift.nf*). In addition, a separate analysis with PhyloSift [57] was performed using a smaller downsampling, by including only the first 100K read pairs of each sample (parameters: `--chunk-size 100000 --paired`). In order to test for associations of phylogenetic diversity with treatment, time of sampling, and differences among hosts at the start of the trial (first post-weaning sample collection time point), analysis of the unrooted phylogenetic diversity (PD) [58], the balance weighted phylogenetic diversity (BWPD) [59] and principal component analysis (PCA) of the Kantorovich-Rubinstein distances [60] (beta diversity analysis) were performed. Collections of phylogenetic placements produced by PhyloSift are compared using guppy [60] to produce the Kantorovich-Rubenstein distances output. The collections of phylogenetic placements are grouped based on the variable of interest (*i.e.*: treatment, time of sampling, and differences among hosts at the start of the trial), and consequently the selection is fed to guppy. Guppy is used in two modalities: 1. `guppy epca` to obtain edge principal components; 2. `guppy fat` to annotate the edges of the phylogenetic tree using the relative abundance of reads placed on each lineage. Alpha-diversity and beta-diversity were analysed and the results were visualized with R [61] and R packages [62–76]. Scripts for the analysis are available at

https://github.com/GaioTransposon/metapigs_phylodiv. The data analysis workflow is
schematically represented in **Figure 1**.

All samples were included in the analysis, except for the analysis of treatment effect, where,
in order to minimize the effect of age as a confounding factor, we excluded animals born on
2017-01-06 and on 2017-01-07. In this manner the largest age gap between animals was 3 days,
with animals born on 2017-01-08 ($n=24$), 2017-01-09 ($n=24$), 2017-01-10 ($n=23$), and 2017-01-
11 ($n=39$). As a result, sample sizes at each time point were the following: Control (t0: 26; t2:
20; t4: 15; t6: 12; t8: 12; t10: 12), D-Scour™ (t0: 15; t2: 15; t4: 15; t6: 10; t8: 12; t10: 10),
ColiGuard® (t0: 16; t2: 16; t4: 16; t6: 12; t8: 12; t10: 12), neomycin (t0: 22; t2: 22; t4: 17; t6: 12;
t8: 12; t10: 12), neomycin+D-Scour™ (t0: 17; t2: 17; t4: 17; t6: 12; t8: 12; t10: 12), and
neomycin+ColiGuard® (t0: 14; t2: 13; t4: 14; t6: 10; t8: 10; t10: 10).

Analysis of raw reads with SortMeRNA[77] (version 4.0.0) was performed as described in
Gaio *et al* (2021) [54]. Briefly, raw reads were mapped against the rRNA reference database silva-
bac-16s-id90.fasta (parameters: `--fastx --blast 1 --num_alignments 1`) (script:
sortmerna.sh) and filtered based on e-value (e-value <= 1e-30), identity (identity ≥ 80%), and
alignment length (length ≥ 100 bp) (script: *sortmerna_filter.sh*). [54] The output from
SortMeRNA[77] was used to compute Principal Component Analysis (PCA) with R [61] (script:
08_sortmerna.R). Sample counts were normalized for library size by proportions and were tested
with the Spearman's Rank correlation coefficient method to find lineages correlating with the
weight of the piglets across the trial (script: *08_sortmerna.R*).

Batch effects
A randomized block design was adopted to mitigate batch effects. Because samples were
distributed across ten 96-well plates during DNA extraction and library preparation, plate effects
were expected. Although samples did not visibly cluster by DNA extraction plate across the first
five principal components, a batch effect was found by multiple comparison analysis with
ANOVA (alpha diversity: *p* range=0.0001-1; beta diversity: *p* value<0.0001) (**Supplementary**
Figure 1). Significance of pairwise comparisons was obtained by running Tukey's *post-hoc*
analysis (**Supplementary Table 1**). Batch effects were removed with ComBat [78] (script:

02_phylodiv.R). ComBat relies on robust empirical Bayes regression to remove heterogeneity

due to the use of batches (in this study a batch is a 96-well plate) while maintaining the

biological variance among samples.[78] The input data used for ComBat in this study is not subject

to normalization as it consists of phylogenetic diversity values obtained from a defined number

of read-pairs per sample. A parametric adjustment was used in the batch effect correction

(`par.prior=TRUE`) and both means and variances were adjusted (`mean.only=FALSE`).

Data availability

217   The raw sequencing data has been submitted to the NCBI Short Read Archive under project

218   PRJNA526405 and is also available through GigaDB at http://dx.doi.org/10.5524/100890. A

219   thorough description of this dataset and the metadata has been reported previously [54]. Scripts

220   used for the data analysis in this study are available in our GitHub repository

221   https://github.com/GaioTransposon/metapigs_phylodiv.

Results

PhyloSift [57] was employed as a means to study microbial community diversity among the

samples, and to test for associations with treatment, time of sampling, and differences among

hosts during the first week post-weaning. To this end, analysis of the unrooted phylogenetic

diversity (PD) [58], the balance weighted phylogenetic diversity (BWPD) [59], and principal

component analysis (PCA) of the Kantorovic-Rubinstein distances [60] (beta diversity analysis)

were performed.

Phylogenetic diversity of positive controls

Unrooted PD was highest for positive control D-Scour™ (mean±SD=162.2±28.3), slightly lower

for positive control ColiGuard® (mean±SD=129.9±50.0) and lowest for the mock community

(mean±SD=123.6±28.4). BWPD was highest for positive control D-Scour™ (mean±SD:

1.9±0.2), lower for the mock community (mean±SD: 1.6±0.1), and lowest for positive control

ColiGuard® (mean±SD: 0.9±0.1). (**Supplementary Table 1**; **Supplementary Figure 2**).

Principal component analysis (PCA) of the Kantorovich-Rubenstein distances (beta

diversity analysis) was performed on positive control samples. Samples clearly separated in PC1

(accounting for 83.62% of the variation), where mock community samples clustered in the lower

end of principal component 1 (PC1) showing a higher representation of *Enterobacteriaceae* and

Pseudomonadaceae, while ColiGuard® and D-Scour™ samples clustered in the higher end of

PC1, showing a higher representation of *Lactobacillales*. D-Scour™ samples clustered in the

higher end of principal component 2 (PC2) (accounting for 14.52% of the variation), showing a

higher representation of *Lactobacillus delbrueckii*, *Enterococcus faecium*, and *Bifidobacterium*,

while in the lower end of PC2, mock community samples and ColiGuard® samples, forming two

separate clusters, showed a higher representation of *Bacilli*, *Enterobacteriaceae*, and

Gammaproteobacteria. (**Supplementary Figure 3**).

Phylogenetic diversity of piglet gut communities: host factors

Based on Kruskal-Wallis one-way analysis of variance (Hommel adjusted *p* values to correct for

multiple hypotheses testing), alpha diversity of the piglet samples did not cluster significantly by

cross-breed type ($p > 0.05$) or by pig line ($p > 0.05$) in the first week post-weaning (**Figure 2;**

Supplementary Table 1). A correlation of cross-breed type was found with beta diversity

(principal component 3) at t2 ($p=0.024$). (**Figure 2**; **Supplementary Table 1**)

The piglets differed slightly by age, being born between 1 and 5 days apart. Notably, we

found a significant correlation between alpha diversity and the age of the piglets at the first

sampling time point (unrooted PD: $p < 0.0001$; BWPD: $p=0.011$) (**Figure 2**; **Supplementary**

Table 1) and between age and beta diversity at t2 (PC3: $p=0.047$) and t7 (PC3: $p=0.018$) (**Figure**

2; **Supplementary Table 1**). As age groups were confounded with cross-breed types (*i.e.* not all

age groups are represented by each of the four cross-breed types), we compared the phylogenetic

diversity of age groups within each breed. As cross-breed types "Landrace × cross bred (LW ×

D)" and "Large White × Duroc" had only a small number of piglets in each age group ($n=9$ and

$n=12$, respectively), we tested for an association between phylogenetic diversity and age in

cross-breeds "Duroc × Landrace" and "Duroc × Large White" ($n=46$ and $n=59$, respectively).

Among these cross-breeds, age in the "Duroc × Landrace" piglets ($n=46$) correlates with alpha

diversity during the first week post-weaning (unrooted PD: $p=0.006$; BWPD: $p=0.047$) and with

beta diversity at t2 (PC2: *p*=0.048) (**Figure 2**; **Supplementary Figure 4**; **Supplementary Table**
1).

As piglets were derived from 42 distinct sows (maternal sows) and nursed by either the
same or a different sow (a nurse sow), a litter effect was expected and determined. Based on
Hommel adjusted *p* values, a similarity of alpha phylogenetic diversity can be seen among
piglets from the same maternal sow (unrooted PD: *p*=0.001; BWPD: *p*=0.017) and in piglets
from the same nurse sow (unrooted PD: *p*=0.002; BWPD: *p*=0.027) (**Figure 2**; **Supplementary**
Table 1). The litter effect described was found significant only in samples from the first week
post-weaning as the significance of the correlations did not persist thereafter (**Figure 2**;
Supplementary Table 1).

A strong effect of aging on phylogenetic diversity
Beta diversity analysis of all samples revealed a distinct and consistent change of the microbial
community over time in all piglets, regardless of the treatment. Beta diversity analysis was
performed from all reads and from the analysis of 16S rDNA V4 region-containing reads.
Samples collected immediately after weaning (t0) were characterized by a higher representation
of *Bacteroidetes* chlorobi group and *Clostridia* (PC1, 47.68%), particularly of *Sedimibacterium*
and *Desulfosporosinus* (PC2 17.2%), respectively. Between day 0 (t0; immediately after
weaning) and day 14 (t4), samples shifted towards a higher representation of *Bifidobacterium*
and *Lactobacillus*, as measured from beta diversity analysis (PC1; 47.68% var. explained) and
from analysis of 16S rRNA reads (PC1 and PC2; 23.9% and 17.2% var. explained, respectively).
During the last two weeks (t6-t10), corresponding to day 21 and 35 of post-weaning, samples
shifted towards a higher representation of *Actinobacteria*, (PC2 21.79% var. explained),
particularly of *Collinsella* (PC1 23.9%, PC2 17.2%). Lineages of the *Erysipelotrichales*
Catenibacterium and *Solobacterium* were also found to be representative of samples of this time
interval (t6-t10). (**Figure 3**).

Beta diversity analysis was performed separately for samples within each time point in order
to find lineages associated with variation within each time point. Extent of variation was derived
from the product of branch width by the variation explained by the principal component (**Figure**

4). The lineages *Enterobacteriaceae* (t0=0.05) and *Bacteroides* (t0=0.06) were responsible for

variation only during the first week after weaning, and *Methanobrevibacter smithii* during the

first and the second week (t0=0.05; t2=0.03). The *Bifidobacterium* lineage was responsible for

variation in the second week (t2=0.09). The following lineages were responsible for variation

throughout the 6 weeks after weaning: *Bacteroidales* (min=0.04; max=0.20), *Prevotellaceae*

(min=0.02; max=0.13), *Coriobacteriaceae* (min=0.01; max=0.09). *Lactobacillus* became

variable after the first week post-weaning and remained highly variable throughout the rest of the

trial (min=0.03; max=0.30) (**Figure 4**).

Taxonomic representation in terms of relative abundance was derived from the branch width

of the phylogenetic tree (**Supplementary Figure 5**) and combined with PhyloSift's taxonomic

annotation of the marker gene phylogeny. *Lactobacillus acidophilus* increased at the start

(t0=0.00; t2=0.07; t4=0.17) then decreased (t6=0.11; t8=0.06; t10=0.08). Among other prevalent

lineages during the week after weaning, up to the next week and dropping at later time points, we

found *Methanobrevibacter smithii* (t0=0.05; t2=0.03) and *Bacteroidales* (t0=0.01; t2=0.01).

Following an opposite trend we found *Ruminococcus* sp. JC304 (t0=0.00; t2=0.00; t4=0.01;

t6=0.03; t8=0.03; t10=0.03), *Solobacterium moorei* (t0=0.00; t2=0.01; t4=0.02; t6=0.03;

t8=0.02; t10=0.03) and *Prevotella copri* (t0=0.00; t2=0.00; t4=0.05; t6=0.03; t8=0.06; t10=0.03).

In modest and stable abundance across the post-weaning period were the following lineages of

the order of the *Clostridiales*: *Mogibacterium* sp. CM50 (t0=0.03; t2=0.05; t4=0.03; t6=0.04;

t8=0.03; t10=0.04), *Oscillibacter* (t0=0.08; t2=0.06; t4=0.04; t6=0.03; t8=0.04; t10=0.03),

Subdoligranulum variabile (t0=0.07; t2=0.03; t4=0.05; t6=0.07; t8=0.07; t10=0.08), and

Ruminococcus bromii (t0=0.03; t2=0.02; t4=0.01; t6=0.01; t8=0.1; t10=0.01). In transient

abundance we found *Bifidobacterium thermophilum* RBL67 (t2=0.02; t4=0.03). Gradually

increasing from the second week we found *Eubacterium biforme* DSM3989 (t2=0.02; t4=0.02;

t6=0.03; t8=0.03; t10=0.02), *Eubacterium rectale* (t4=0.03; t6=0.03; t8=0.03; t10=0.01) and,

after the third week, *Faecalibacterium prausnitsii* (t6=0.01; t8=0.01; t10=0.01).

(**Supplementary Figure 5**)

The effect of time was also clear in alpha diversity, where all the piglet cohorts followed a

similar trend over time, independent of the treatment (**Supplementary Figure 6**;

Supplementary Table 1). Immediately after weaning (t0), the unrooted PD was lower for the

piglets than the sows (sows: 328.5±24.0; piglets: 296.5±34.7) and reached a higher unrooted PD

to the sows' in the following week (t2) (piglets: 336.18±33.0) (**Supplementary Table 1**). In

comparing four timepoints at one-week intervals from the start of the trial, changes in alpha

diversity among all the piglets were tested for and significance was determined using the

Bonferroni correction. Unrooted phylogenetic diversity increased in the first week following

weaning (t0-t2: +9.15%; $p<0.001$) and decreased in the following week (t2-t4: -4.54%; $p<0.001$)

amongst the piglets. In contrast, BWPD decreased in the week after weaning (t0-t2: -6.07%;

$p<0.001$), to increase in the following week (t2-t4: +4.77%; $p<0.001$) and decrease in the fourth

week (t4-t6: -3.67%; $p=0.002$). (**Supplementary Figure 6**; **Supplementary Table 1**).

Effect of antibiotic and probiotic treatment on alpha diversity

We hypothesized that the probiotic treatments, whether alone (D-Scour™ and ColiGuard®) or

administered after neomycin (neomycin+D-Scour™ and neomycin+ColiGuard®) would cause a

change in the microbial community composition that would be measurable via phylogenetic

diversity. We tested whether the treatments correlated with a change in phylogenetic diversity

independently of the changes occurring with time. Given the differences in alpha and beta

diversity detected among the subjects after weaning, we analyzed the deltas of phylogenetic

diversity instead of relying on the absolute means, similar to the procedure applied by Kembel *et*

al (2012) [79]. Time-point measurements of alpha diversity were taken, and deltas were computed

for each piglet. Delta means were compared between cohorts, where the control cohort would

serve as a control group for neomycin, D-Scour™ and ColiGuard® cohorts, whereas the

neomycin cohort would serve as a control group for the neomycin+D-Scour™ and

neomycin+ColiGuard® cohorts.

One week after weaning, 90% of the piglets displayed an increase in unrooted PD and 71%

displayed a decrease of BWPD. The following week the trend was opposite: 72% of the piglets

displayed an increase of BWPD and 76% displayed a decrease of unrooted PD (**Supplementary**

Table 1). However, the neomycin cohort displayed the smallest BWPD drop in the week

following weaning, and the overall trend of neomycin in unrooted PD was the most different

from the other cohorts (**Supplementary Figure 6**). Due to the lower drop in BWPD of

neomycin, significance was found in BWPD during the first week between neomycin and

Neomycin+ColiGuard® (t0-t2; Tukey adjusted *p* value=0.031), and between neomycin and

ColiGuard® (t0-t2; Tukey adjusted *p* value=0.041). No other significant differences were found

in changes in alpha diversity between time points among the cohorts. (**Supplementary Table 1**)

Effect of antibiotic and probiotic treatment on beta diversity

To investigate the treatment effect on beta diversity, principal component analysis (PCA) of the

Kantorovich-Rubenstein distances (beta diversity analysis) was performed on all samples and,

additionally, on samples within individual time points. This analysis is conceptually similar to

the weighted Unifrac approach for beta diversity analysis, but is designed to work with

phylogenetic placement data [60]. When examining all samples together, there was no clear

separation of cohorts on any of the five principal component axes. When individual time points

were analysed, some clustering by cohort was observed (**Supplementary Figure 7**). D-Scour™

separated from the control cohort in PC3 (10.97%) during the first week of probiotic treatment

(t2) (**Supplementary Figure 7**). ColiGuard® separated from control during the first week of

probiotic treatment (t2) in PC5 (2.82%), and two weeks after the end of probiotic treatment in

PC3 (7.87%), showing a higher representation of *Lactobacillus*.

Two weeks after the end of neomycin treatment (t6), neomycin separated from control in

PC4 (5.65%), with a smaller clustering and a higher representation of *Mollicutes*. Neomycin+D-

Scour™ separated from neomycin in PC2 (17.80%) and PC4 (5.65%) at the end of probiotic

treatment (t6), as well as a week later (t9) in PC5 (3.93%). In these instances, neomycin clustered

towards a higher representation of *Mollicutes*, while Neomycin+D-Scour™ showed a higher

representation of *Lactobacillus*. (**Supplementary Figure 7**)

Association between weight and community composition

Weight correlated with the abundance of certain taxa at each time point as it resulted from

principal component analysis of 16S rRNA reads. Positively correlating with weight we found

among others: *Blautia* (t0), *Cetobacterium* (t0), *Lactobacillus* (t6), *Mycoplasma* (t6),

Anaerostipes (t8), *Ruminococcus* (t8), *Cerasibacillus* (t10). Negatively correlating with weight

we found among others: *Pyramidobacter* (t0), *Odoribacter* (t2), *Schwartzia* (t6), *Streptococcus*

(t6), *Dokdonella* (t8). Correlations remained significant after *p* value correction for multiple

hypotheses testing (Hommel, and Benjamini & Yekutieli), but only one taxa

(*Coprothermobacter*; t0) remained significantly, positively correlated with host weight after

applying the more stringent Bonferroni correction (**Supplementary Table 1**).

Effect of treatments on weight gain

Overall weight gain from initial to final weight (**Supplementary Table 1**) was not significantly

affected by any treatment. However, the probiotic ColiGuard® was found to have a partial effect

on piglet weight gain (**Supplementary Figure 8**; **Supplementary Table 1**). Weight was

measured weekly for a total of six measurements. To minimize age as a confounding factor from

this analysis, we kept samples from piglets that were born a max of 3 days apart. Based on Tukey

adjusted *p* values, a lower weight gain was detected in the ColiGuard® cohort compared to the

control cohort between the last day of probiotic treatment and a week after treatment (t4-t6)

(*p*=0.015) (**Supplementary Figure 8**; **Supplementary Table 1**). Similarly, a lower weight gain

was detected in the neomycin+ColiGuard® cohort between the 9th day of treatment and the 9th

day after probiotic treatment (t4-t8) compared to the neomycin cohort (*p*=0.011)

(**Supplementary Figure 8**; **Supplementary Table 1**). Age and cross-breed differences among

piglets were not associated with weight gain (**Supplementary Table 1**).

Estimation of phylogenetic diversity using sample downsampling sizes: 100K versus 1M

We compared unrooted PD and BWPD values obtained using either 100,000 or 1,000,000 reads

per sample (corresponding to 0.612% or 6.12% of the average sample). The median unrooted PD

was 2.6x higher, and variance improved when the analysis was run from 1M reads per sample,

compared to the analysis run using 100K reads per sample (100K: 121±18.67; 1M: 317±50.67;

Pearson's r=0.8965), while the median BWPD remained nearly unchanged (100K: 2.11±0.203;

1M: 2.11±0.193; Pearson's r=0.9747). Comparing beta diversity values obtained using either

100,000 or 1,000,000 reads per sample, we found a high correlation for all five principal

components (Pearson's r range=0.9809-0.9956). (**Supplementary Figure 9**)

Discussion

The microbial composition of positive control samples was analysed from a taxonomic
perspective in our previous study [54], and it was here compared to the phylogenetic diversity
obtained for these samples. The unrooted PD reflects the absolute diversity, independently of the
relative abundance of each species, within a sample. In fact, low-level contamination (<0.1%)
detected in each of the positive controls (mock community: 1 taxon; ColiGuard®: 20 taxa; D-
Scour™: 25 taxa) [54] contributes toward the absolute diversity in the unrooted PD, inflating this
value. Unrooted PD values obtained for the positive controls were directly proportional to their
respective count of taxa, which include contaminants (D-Scour™: $n=33$; ColiGuard®: $n=22$;
mock community: $n=8$) [54]. On the other hand, the contribution to total diversity of phylogenetic
tree edges with uneven quantities of reads placed on either side is down-weighted in BWPD. The
fact that ColiGuard®, mock community, and D-Scour™ are mainly composed of 2, 7, and 8 taxa
taking up even proportions in the samples, respectively (hence excluding contaminants) [54], was
reflected by their lowest, higher, and highest BWPD measured from these samples.

The consistent trend in community composition over time, across all the cohorts,
indicates that an age-related process of ecological succession is the largest factor shaping the
microbial community of post-weaning piglets, as found in this study where animals aged 20-63
days were fed the same diet. A peak in unrooted phylogenetic diversity and drop in balance
weighted phylogenetic diversity (BWPD) reflects the acquisition of new species with the loss of
dominating species. Similarly to Pollock *et al* (2018) [80], we found the relative abundance of
Clostridiales, and *Lachnospiraceae* to decrease from the start of weaning to 2 weeks into
weaning, while the relative abundance of *Lactobacillus*, *Prevotella copri*, *Faecalibacterium*
prausnitsii and *Erysipelotrichaceae* increased. The increase of *Prevotella* with weaning [80–82] is
well documented and it is associated with the increased polysaccharide consumption associated
with the start of solid food consumption [81,83]. The relative abundance of *Lactobacillus* goes up
with weaning, as other studies suggested [80,81]. However, according to our analysis, this increase
concerns *Lactobacillus acidophilus*, while other *Lactobacillus* species, such as *Lactobacillus*
vaginalis ATCC follow the opposite trend. We also found a clear gradual increase in abundance

of *Solobacterium moorei* and *Eubacterium biforme* DSM3989, a transient increase of

Bifidobacterium thermophilum RBL67 during the second and the third week post weaning, and a

sharp decrease in the abundance of *Methanobrevibacter smithii* after the second week post-

weaning.

The change in phylogenetic diversity detected in the week following the piglets' arrival at

the trial site irrespective of the cohort, could be linked to the piglets being subjected to microbial

interchange (*e.g.*: new pen mates) and/or to diet transition (peri-weaning transition) to solid food

leading to the reshaping of the gut microbial community. [84,85] The week following the drop of

BWPD, a significant increase of BWPD was recorded, reflecting the acquisition of a larger

proportion of the community by the newly introduced species. The strong changes in

phylogenetic diversity detected in the first and the second week could as well be attributable to

other post-weaning related physiological changes, as previous studies report [84–87].

The highest inter-individual differences among piglets are seen in the first week of life,

irrespective of maternal or environmental effects. The microbiota of three week old piglets is still

very dynamic, but environmental factors become evident [85]. At six weeks of age, CD8+ T cells

infiltrate the intestinal tissue and the mucosa and intestinal lining resemble that of an adult pig [79].

The increase of alpha diversity associated with weaning has been measured before [81,82,88], but

rarely using metrics that allow the distinction of absolute diversity from evenness. In this study,

one week after weaning, piglets reached a comparable absolute diversity to the sows, at which

time the piglets were aged between 3.8 and 4.6 weeks. Unrooted PD did not reach higher levels

at later sampling time points. The highest BWPD accompanied by a high unrooted PD was

reached after the second week post-weaning when piglets were aged between 4.9 and 5.6 weeks.

Age-dependent physiological changes could explain i) the major shifts we detected in alpha

diversity during the first two weeks after weaning and, ii) the distinct differences in community

composition with age, even with a narrow age difference between piglets (1-5 days). We found a

significant difference in microbial composition in both absolute diversity as in balance-weighted

PD, only in the first week after weaning (piglets aged between 3.8 and 4.6 weeks), between

groups of piglets that were separated by up to five days maximum by day of birth. Since age

groups were confounded with breeds in our study, we attempted to determine the correlation

within single cross-breeds. Unfortunately, although the correlations with age could still be

detected, we could not determine the association at later time points due to the introduction of

treatment effects.

Animal trials are often conducted in controlled environments to minimize environmental

effects. However, individual variations such as breed and age are often unavoidable in large

animal trials, especially involving animals derived from commercial herds. Previously reported

confounding factors include: individual variation, cohabitation, age, maternal effects, hormones,

behavioural differences between breeds (*e.g.* coprophagy, mouth to mouth contact) and extent of

long-term behavioural adaptation, which can differ between breeds for reasons not attributable to

genetics [84–86,89]. A litter effect was found in piglets at the start of the trial (piglets aged between

3.8 and 4.6 weeks; samples collected immediately after weaning) and was lost at later time

points during the trial. This could be due to either of the aforementioned factors. Co-housing,

aging and the splitting of the piglets in separate rooms to receive a different treatment, are

possible causes for loss of the litter effect. In this study we confirm the importance of these

factors in the contribution to inter-individual variability of gut microbial composition. Motta *et al*

(2019) report a correlation of beta diversity with age and no correlation of genotype and litter

effect with either alpha or beta diversity [90]. On the contrary, we found the piglet samples to

significantly cluster by age and litter immediately after weaning in alpha diversity, and by age

and cross-breed in beta diversity during the second week after weaning. Samples also clustered

by age groups in beta diversity four weeks after weaning. The groups we tested, based on age,

cross-breed, and litter, differed in sample sizes, therefore a non-parametric test was used to test

for associations of host factors with phylogenetic diversity. Based on our results we conclude

that even small age differences among post- weaning piglets, down to the day, must be

accounted for in an experimental set up, but we acknowledge that the method we used may have

less power to detect small effects than methods which make assumptions of balance in group

sizes.

Three groups of piglets (cohorts neomycin, neomycin+D-Scour™ and

neomycin+ColiGuard®) underwent five days of treatment with neomycin, via intramuscular

administration. Intramuscular neomycin poorly diffuses (<10%) into a healthy gastrointestinal

tract [91], therefore a direct effect of neomycin on the gut microbiome may not be expected.

However, neomycin showed a different trend in unrooted PD between the second and the third

week post-weaning, corresponding to the week following the neomycin treatment period for the

neomycin cohort. Taking this time frame into consideration, the neomycin cohort did not

increase in BWPD to the extent of the Control cohort. Although statistically significant

differences between neomycin and Control in alpha diversity were not reached, possibly due to

the effect being too small to detect within a relatively small sample size [92–94], BWPD of the

neomycin cohort appears to follow a different trend to the Control cohort from the first week

(during neomycin treatment) where neomycin treated piglets show the lowest decrease of BWPD

compared to the control cohort and all other cohorts. While all cohorts show an increase in

absolute phylogenetic diversity accompanied by a decrease of diversity evenness during this time

frame, the neomycin cohort piglets show a lower drop in BWPD, suggesting an increase of

species richness, without a corresponding loss of species evenness. Furthermore the neomycin

cohort significantly separated from the control cohort in beta diversity two weeks after neomycin

treatment, showing a higher representation of *Mollicutes*. Numerous studies report the link of

oral antibiotic use with dysbiosis [10,14–17,86,95], as well as with host physiology changes [15]. On the

contrary, the effect of intramuscular antibiotic administration on the microbiome is less well

investigated. Correlation between intramuscular antibiotic use and dysbiosis has been reported in

fish [96], gorillas [19], and pigs [21,97]. In one day old piglets, a single IM injection of amoxicillin

(penicillin class) is reported to have an effect on the intestinal microbiota, detectable 40 days

post treatment [21]. Zeineldin *et al* (2018) tested the effects of IM administration of several

antibiotics of various classes (penicillin, macrolide, cephalosporin and tetracycline), in 8-week

old piglets, reporting shifts of the *Firmicutes*/*Bacteroidetes* ratio following treatment (length of

the treatment not reported) [97]. The effects of intramuscular administration of neomycin

(aminoglycoside class) on the gut microbiota have to our knowledge not been investigated.

Based on our results we conclude that a mild effect on phylogenetic diversity is appreciable post

IM neomycin treatment, up to two weeks after termination of the treatment. Additional

compositional and functional analysis is necessary to determine the source of this mild variation.

Differences were not detected at later time points, based on our phylogenetic diversity analyses,

suggesting a full recovery of the microbial communities after two weeks from the end of the

treatment.

It is possible that the large shifts in phylogenetic diversity taking place in the first two weeks

irrespective of the treatment (an increase, then decrease of unrooted PD, and an opposite trend of

BWPD) have masked the milder effects of the treatment, despite our efforts to control for the

effects of aging. This could be the reason why a significantly distinct alpha diversity trend was

found in the neomycin+D-Scour™ cohort compared to the neomycin cohort, but not in the D-

Scour™ cohort compared to the Control cohort. There are multiple studies reporting beneficial

effects of probiotic treatment in sucker and weaner piglets in terms of improved gut mucosal

integrity[27,98], growth rate[98–100,] digestibility of proteins and water absorption [98,101] , reduction of

pathogen invasion efficiency [38,45,98], and decreased mortality [98,100]. Although the assessment of

physiologic changes from probiotic treatments was outside the scope of this study, we found

significant separation of neomycin+D-Scour™ cohort samples to neomycin cohort samples in

beta diversity 2 and 12 days after D-Scour™ treatment, where neomycin+D-Scour™ samples

showed a higher representation of *Lactobacillales* compared to neomycin samples, suggesting a

transient establishment of the probiotic strains in the piglet guts.

The second probiotic in this study, ColiGuard®, did not have an effect on alpha diversity,

but clustering was detected in beta diversity, where ColiGuard® samples separated from Control

cohort samples in the first week of probiotic treatment and two weeks after probiotic treatment.

Additionally, the ColiGuard® treatment correlated with a lower weight gain, whether or not it

was preceded by the antibiotic treatment. However, when comparing the overall weight gain

(from the start to the end of the trial, corresponding to the six weeks after weaning) the weight

gain in the cohorts receiving ColiGuard® did not differ from the other cohorts.

We extracted the 16S rRNA gene hypervariable regions from our dataset, obtained the

counts, and ran a correlation analysis to discover lineages that correlated with the weight of the

piglets. As a consequence of the library size normalization step, the use of correlation with

compositional data can inflate the false discovery rate [102,103]. For this reason it can be expected

that some of the lineages we found to correlate with the weight of the piglets (eighty-three

distinct species) could be spurious while other correlations may have been missed.

As our initial analysis of phylogenetic diversity was obtained using a 100K reads

downsampling size per sample [104], we wanted to test to what extent the use of a larger sampling

size would affect the diversity obtained from these samples. A 100K downsampling size

corresponds to a 0.6% of reads of an average sample. Increasing the sampling size from 0.6% to

6%, affected BWPD only slightly, while it increased the unrooted PD and decreased its variance.

As expected, a larger sampling size enriches the absolute diversity by increasing the chance to

make new read placements, and it lessens the variance in estimates, but it does not affect the core

microbiome. However, the small difference between sampling sizes could 1. demonstrate how

powerful the method is in describing diversity using a small sampling size, or 2. reflect the limits

imposed by the phylogenetic markers database, as it was pointed out by Darling *et al* (2014) [57].

Conclusions

571
Our findings stress the importance of confounding factors such as breed, age and maternal

effects when assessing the effect of treatment on the gut microbiome. We found that age, even

within a narrow age span (1-5 days) can have an impact on microbial shifts and should be

accounted for in microbiome studies, either (i). by accounting for it as a confounding variable in

the hypothesis-testing model used, or (ii). by avoiding, where possible, the inclusion of subjects

of different ages, or (iii). by allowing a sufficiently long period of time prior to the start of the

treatment. The latter allows animals to become accustomed to the new environment and

researchers to perform additional sampling. Allowing animals a sufficiently long period of time

to become accustomed to the new environment (*e.g.*: temperature, humidity, new microbes, etc)

is meant to reduce noise derived from external factors, while repeated sampling, likewise, aims

to increase the confidence in the signal prior to treatment.

Intramuscular neomycin treatment correlated with a clustering in alpha diversity and a

higher representation of *Mollicutes* compared to control. D-Scour™ treated piglets displayed a

transient establishment of *Lactobacillales*. ColiGuard® treated piglets displayed a clustering in

beta diversity and a transient lower weight gain compared to control. Weight correlated with the

abundance of a number of lineages. Age was the strongest factor shaping phylogenetic diversity

of the piglets.

As previously mentioned, phylogenetic diversity is based on distinct lineages (richness) and
their collective structure (proportions reflected by BWPD) and not on a direct assessment of
composition and function. These types of analyses will be necessary to further describe the
effects of the treatments.

Acknowledgements
Shayne Fell is acknowledged for overseeing daily activities during the pig trial. Thanks to Amy
Bottomley, Giulia Ballerin, and Rosy Cavaliere, for providing the mock community strains, and
Akane Tanaka, Leigh Monahan, Michael Liu and Joyce To for the technical support. John
Webster is thanked an internal review of this manuscript and for the helpful suggestions
regarding its content. International Animal Health Products is acknowledged for providing
access to the probiotic supplements used in this study.

Conflicts of interest
D-Scour™ was sourced from International Animal Health Products (IAHP). ColiGuard® was
developed in a research project with NSW DPI, IAHP and AusIndustry Commonwealth
government funding.

Funding information
This work was supported by the Australian Research Council, linkage grant LP150100912. This
project was funded by the Australian Centre for Genomic Epidemiological Microbiology
(Ausgem), a collaborative partnership between the NSW Department of Primary Industries and
the University of Technology Sydney. TZ and DG are recipients of UTS International Research
and UTS President's Scholarships. NSW DPI approved the paper before submission for
publication.

Captions:

Figure 1. Workflow.

Schematic workflow from sample collection to sequencing (orange) and data analysis (green). Scripts (italic) are available in our GitHub repository.

Figure 2. Significant correlations of phylogenetic diversity with specified host factors.

The correlation of host factors birth day, cross-breed, birth day within the D × L cross-breed, birth day within the D × LW cross-breed, line, litter (maternal sow and nurse sow) with alpha phylogenetic diversity (**A**) and beta phylogenetic diversity (**B**) from each time point (x-axes) was obtained. Shown in A and B are the significance values of these correlations derived from Kruskal-Wallis analysis, where the symbol shape indicates the phylogenetic diversity measure of alpha (unrooted pd and BWPD) and beta (PC components 1 to 5), while the symbol color indicates the significance before (black) and after (red) Hommel *p* value adjustment. C) For each red symbol in A and B, describing a significant correlation after Hommel *p* value adjustment, a plot of the phylogenetic diversity estimates is shown. Phylogenetic diversity estimates of the litter effect are not shown. All p values are reported in Supplementary Table 1. Abbreviations: D × L = "Duroc × Landrace" cross breed; D × LW = "Duroc × Large White" cross breed.

Figure 3. Effect of time on beta diversity.

Principal component analysis (PCA) of samples. PCA from edge component analysis with PhyloSift (top) and PCA from 20 most abundant 16S reads extracted with SortMeRNA (bottom). In the top, distribution of samples on either side of the plot (left *versus* right; top *versus* bottom) reflect the lineages that were found to explain the variation. Samples are color coded by time post-weaning (days). In the lower plot, arrows indicate which of the 20 lineages contributed to the variation of samples across time, where arrows thickness represents a higher (thicker) or lower (thinner) contribution. Samples are color coded by time point during the trial (t0: start of the trial/first day after weaning; t10: last day of the trial/40th day after weaning).

Figure 4. Lineages displaying the highest variation in beta diversity across time.

Heatmap of lineages explaining the community composition of samples from separate time points

of the trial (1 week interval between time points) derived from edge principal component analysis.

Intensity is derived from branch width by the percentage of variability explained by the principal

components.

Supplementary Figure 1. Batch effect on alpha and beta diversity before batch effect removal.

Batch effect by alpha (top two plots) and beta diversity (bottom five plots) before batch effect

removal. Samples are grouped by DNA extraction plate. The p values are derived from multiple

comparison analysis with ANOVA, indicating equality of the means. *Post hoc* corrected p values

for pairwise comparisons are provided in Supplementary Table 1.

Supplementary Figure 2. Alpha phylogenetic diversity of cohorts.

Alpha phylogenetic diversity per cohort from samples across all time points. Balance-weighted

phylogenetic diversity (BWPD) (top) (mean±SD: Positive control Mock community: 1.59±0.07;

Positive control D-Scour™: 1.94±0.21; Positive control ColiGuard®: 0.86±0.14; Control:

2.13±0.13; D-Scour™: 2.16±0.12; ColiGuard®: 2.12±0.12; neomycin: 2.12±0.16; neomycin+D-

Scour™: 2.13±0.13; neomycin+ColiGuard®: 2.10±0.14; sows: 2.12±0.15; all piglet cohorts:

2.13±0.13); Unrooted phylogenetic diversity (bottom) (mean±SD: Positive control Mock

community: 123.62±24.41; Positive control D-Scour™: 162.14±28.27; Positive control

ColiGuard®: 129.88±50.00; Control: 311.23±29.23; D-Scour™: 316.35±24.99; ColiGuard®:

311.98±42.51;   neomycin: 314.81±44.01;   neomycin+D-Scour™:   312.66±40.18;

neomycin+ColiGuard®: 316.29±31.20; sows: 328.51±24.00; all piglet cohorts: 313.86±36.00).

Supplementary Figure 3. Beta diversity of positive controls.

Principal component analysis (PCA) of positive control samples. PCA from edge component
analysis with PhyloSift. Distribution of samples on either side of the plot (left *versus* right; top
versus bottom) reflect the lineages that were found to explain the variation.

Supplementary Figure 4. Alpha phylogenetic diversity by age breed.

Alpha diversity of samples from the start of the trial (immediately after weaning) grouped by breed
and by date of birth. Unrooted phylogenetic diversity (top) and balance-weighted phylogenetic
diversity (bottom). P values are derived from Kruskal-Wallis analysis of variance. Piglets of the
Duroc × Landrace breed (n=46) separated significantly by age in unrooted phylogenetic diversity
and in BWPD at the start of the trial (t0) (Hommel adjusted p value: unrooted pd=0.006;
BWPD=0.047). All post hoc corrected p values are provided in Supplementary Table 1.

Supplementary Figure 5. Relative abundance heatmap.

Most abundant lineages within each time point (columns) are obtained from analysis with guppy
fat. Guppy fat outputs trees with fattened edges in proportion to the relative abundance of reads
place in each lineage. The branch width of the trees, each corresponding to samples from distinct
time points, are the entries for this heat map. The distance between each time point is one week.

Supplementary Figure 6. Time trend of alpha diversity by cohort.

Unrooted phylogenetic diversity (**A**) and balance-weighted phylogenetic diversity (**B**) describe
richness and evenness, respectively, of alpha phylogenetic diversity for all samples across time,
grouped and color coded by cohort. The p values derive from pairwise comparisons of time points
of all treatment cohorts. The p values and *post hoc* corrected p values of time points comparisons
for each separate treatment cohort are provided in Supplementary Table 1.

Supplementary Figure 7. Significant differences in beta diversity between cohorts at

specific time points.

Significance was determined by comparing groups by pairwise t-test and the resulting p values

were adjusted with the Bonferroni method. Significance values are provided in Supplementary

Table 1. The x-axes represent the principal component. As plots are derived from distinct guppy

runs, each principal component explains variation to a different extent (percentage specified in

parentheses). The number of samples is specified on the y-axis. Distribution of the samples on

either side of a plot (left *versus* right) reflects the lineages that were found to explain the variation.

Distributions are color coded by cohort.

Supplementary Figure 8. Change in weight gain of piglets by cohort across the trial.

On the y-axis of plots the change in weight gain between time points is provided in percentage.

Letters on the top left of each plot indicate the time points compared with one week interval (t0-

t2, t2-t4, t6-t8, t8-t10) and with two weeks interval (t2-t6, t4-t8, t0-t8). Pairwise t-test comparisons

between cohorts were computed. A significant difference was found between Control and

ColiGuard (t4-t6, Tukey adjusted p value=0.0084), between neomycin and neomycin+ColiGuard

(t4-t6, Tukey adjusted p value=0.0152) and between neomycin and neomycin+ColiGuard (t4-t8,

Tukey adjusted p value=0.011).

Supplementary Figure 9. Correlation of diversity indices between the analysis run using 100,000

(100K) and 1,000,000 (1M) reads per sample.

Phylogenetic diversity analysis (alpha and beta) was run using either 100K or 1M reads per sample,

corresponding to a 0.6% or 6% of the average sample. We show the Pearson's correlation of these

analyses run using distinct downsampling sizes. Significance of correlations is reported within

each plot. All the diversity indices are provided in Supplementary Table 1.

Supplementary Table 1. Statistical analysis output.

Statistical analysis output is split across the following sheets. **all_padj_Hommel** : list of p values

obtained from running Kruskal-Wallis test, adjusted using the Hommel correction for

phylogenetic diversity values: unrooted pd, BWPD, and five principal components. Grouping of

samples is based on: cross-breed, line, date of birth, date of birth within distinct cross-breeds,

maternal sow and nurse sow. Sample size and sample collection date is reported. **all_pvalues** :

reports the *p* values of "all_padj_Hommel" sheet, prior to Hommel *p* value adjustment.

alpha_delta_cohorts : deltas between time points are obtained per cohort and these deltas are

compared using ANOVA, adjusting *p* values using the TukeyHSD method. **alpha_means** : the

means and standard deviations of alpha diversity (unrooted pd and BWPD) obtained per time

point and cohort. **alpha_time** : results of comparison between time points of alpha diversity

(unrooted pd and BWPD) using the t-test and Bonferroni *p* value correction.

batch_post_process : results of comparison of alpha diversity values by DNA extraction plate

(1-10) after batch effect removal, run using ANOVA and TukeyHSD *p* value adjustment method.

batch_pre_process : results of comparison of alpha diversity values by DNA extraction plate (1-

10) before batch effect removal, run using ANOVA and TukeyHSD *p* value adjustment method.

deltas_percent_change : comparison of alpha diversity values (unrooted pd and BWPD)

between time points within distinct cohorts and within all cohorts. Sample size (*n*), deltas

(percentage) are shown. **guppy_padj** : *p* values from "guppy_pvalues" are adjusted using the

Bonferroni method and results are reported. **guppy_pvalues** : output of single guppy runs is

analyzed by comparing beta diversity values by cohort at each time point using the pairwise t

test; results of the tests are here shown. **PD_100K_vs_1M_stats** : alpha and beta diversity values

obtained with PhyloSift using either 100,000 reads or 1,000,000 reads downsampling size.

weight_cohort_stats : piglets were weighted at each time point. Deltas are obtained per piglet

and means were compared between cohorts using ANOVA and TukeyHSD *p* value correction,

results are reported. **weight_taxa** : Spearman's rank correlation was assessed between pig weight

and abundance of lineages obtained from 16S rRNA containing reads. Significance values are

adjusted using the distinct methods listed.

References

1. Jensen, P. & Recén, B. When to wean — Observations from free-ranging domestic pigs. *Appl. Anim. Behav. Sci.* **23**, 49–60 (1989).

2. Jensen, P. Observations on the maternal behaviour of free-ranging domestic pigs. *Appl. Anim. Behav. Sci.* **16**, 131–142 (1986).

3. Lalles, J.-P., Bosi, P., Smidt, H. & Stokes, C. R. Nutritional management of gut health in pigs around weaning. *Proc. Nutr. Soc.* **66**, 260–268 (2007).

4. Van Heugten, E., Coffey, M. T. & Spears, J. W. Effects of immune challenge, dietary energy density, and source of energy on performance and immunity in weanling pigs. *J. Anim. Sci.* **74**, 2431–2440 (1996).

5. Nabuurs, M. J. A. Weaning piglets as a model for studying pathophysiology of diarrhea. *Vet. Q.* **20**, 42–45 (1998).

6. Heo, J. M. *et al.* Gastrointestinal health and function in weaned pigs: a review of feeding strategies to control post-weaning diarrhoea without using in-feed antimicrobial compounds. *J. Anim. Physiol. Anim. Nutr.* **97**, 207–237 (2013).

7. Jordan, D. *et al.* Antimicrobial use in the Australian pig industry: results of a national survey. *Aust. Vet. J.* **87**, 222–229 (2009).

8. Barton, M. D. Impact of antibiotic use in the swine industry. *Curr. Opin. Microbiol.* **19**, 9–15 (2014).

9. Schokker, D. *et al.* Long-lasting effects of early-life antibiotic treatment and routine animal handling on gut microbiota composition and immune system in pigs. *PloS One* **10**, e0116523 (2015).

10. Looft, T. P. *The swine intestinal microbiota: localized adaptations and responses to in-feed antibiotics*. (Iowa State University, 2012).

11.   Zhang, D. *et al.* Changes in the diversity and composition of gut microbiota of weaned

piglets after oral administration of Lactobacillus or an antibiotic. *Appl. Microbiol. Biotechnol.*

100, 10081–10093 (2016).

12.   Sachs, J. L., Mueller, U. G., Wilcox, T. P. & Bull, J. J. The evolution of cooperation. *Q. Rev.*

Biol. **79**, 135–160 (2004).

13.   Foster, K. R. & Wenseleers, T. A general model for the evolution of mutualisms. *J. Evol.*

Biol. **19**, 1283–1293 (2006).

14.   Cho, I. *et al.* Antibiotics in early life alter the murine colonic microbiome and adiposity.

Nature **488**, 621–626 (2012).

15.   Antunes, L. C. M. *et al.* Effect of Antibiotic Treatment on the Intestinal Metabolome.

Antimicrob. Agents Chemother. **55**, 1494–1503 (2011).

16.   Gonzalez-Perez, G. *et al.* Maternal Antibiotic Treatment Impacts Development of the

Neonatal Intestinal Microbiome and Antiviral Immunity. *J. Immunol.* **196**, 3768–3779

(2016).

17.   Dethlefsen, L. & Relman, D. A. Incomplete recovery and individualized responses of the

human distal gut microbiota to repeated antibiotic perturbation. *Proc. Natl. Acad. Sci.* **108**,

4554–4561 (2011).

18.   Kim, A. *et al.* Administration of antibiotics can cause dysbiosis in fish gut. *Aquaculture* **512**,

734330 (2019).

19.   Vlčková, K. *et al.* Effect of Antibiotic Treatment on the Gastrointestinal Microbiome of Free-

Ranging Western Lowland Gorillas (Gorilla g. gorilla). *Microb. Ecol.* **72**, 943–954 (2016).

20.   Thymann, T. *et al.* Antimicrobial treatment reduces intestinal microflora and improves

protein digestive capacity without changes in villous structure in weanling pigs. *Br. J. Nutr.*

97, 1128–1137 (2007).

21.   Janczyk, P. *et al.* Parenteral long-acting amoxicillin reduces intestinal bacterial community

diversity in piglets even 5 weeks after the administration. *ISME J.* **1**, 180–183 (2007).

22. Birck, M. M. *et al.* Enteral but not parenteral antibiotics enhance gut function and prevent

necrotizing enterocolitis in formula-fed newborn preterm pigs. *Am. J. Physiol.-Gastrointest.*

Liver Physiol. **310**, G323–G333 (2016).

23. Wiuff, C., Lykkesfeldt, J., Svendsen, O. & Aarestrup, F. M. The effects of oral and

intramuscular administration and dose escalation of enrofloxacin on the selection of

quinolone resistance among Salmonella and coliforms in pigs. *Res. Vet. Sci.* **75**, 185–193

(2003).

24. Chantziaras, I., Smet, A., Haesebrouck, F., Boyen, F. & Dewulf, J. Studying the effect of

administration route and treatment dose on the selection of enrofloxacin resistance in

commensal Escherichia coli in broilers. *J. Antimicrob. Chemother.* **72**, 1991–2001 (2017).

25. Ricker, N. *et al.* Toward antibiotic stewardship: route of antibiotic administration impacts

the microbiota and resistance gene diversity in swine feces. *Front. Vet. Sci.* **7**, 255 (2020).

26. Lessard, M. *et al.* Administration of Pediococcus acidilactici or Saccharomyces cerevisiae

boulardii modulates development of porcine mucosal immunity and reduces intestinal

bacterial translocation after Escherichia coli challenge1, 2. *J. Anim. Sci.* **87**, 922 (2009).

27. Deng, J., Li, Y., Zhang, J. & Yang, Q. Co-administration of Bacillus subtilis RJGP16 and

Lactobacillus salivarius B1 strongly enhances the intestinal mucosal immunity of piglets.

Res. Vet. Sci. **94**, 62–68 (2013).

28. Garrido-Mesa, N. *et al.* The association of minocycline and the probiotic Escherichia coli

Nissle 1917 results in an additive beneficial effect in a DSS model of reactivated colitis in

mice. *Biochem. Pharmacol.* **82**, 1891–1900 (2011).

29. Ukena, S. N. *et al.* Probiotic Escherichia coli Nissle 1917 Inhibits Leaky Gut by Enhancing

Mucosal Integrity. *PloS One* **2**, e1308 (2007).

30. Rangan, K. J. *et al.* A secreted bacterial peptidoglycan hydrolase enhances tolerance to

enteric pathogens. *science* **353**, 1434–1437 (2016).

31.   Deriu, E. *et al.* Probiotic bacteria reduce salmonella typhimurium intestinal colonization by

832   competing for iron. *Cell Host Microbe* **14**, 26–37 (2013).

833   32.   Hancock, V., Dahl, M. & Klemm, P. Probiotic Escherichia coli strain Nissle 1917

834   outcompetes intestinal pathogens during biofilm formation. *J. Med. Microbiol.* **59**, 392–399

835   (2010).

836   33.   Heyman, M. Effect of lactic acid bacteria on diarrheal diseases. *J. Am. Coll. Nutr.* **19**,

837   137S-146S (2000).

838   34.   Jin, L.-Z., Marquardt, R. R. & Baidoo, S. K. Inhibition of enterotoxigenic Escherichia coli

839   K88, K99 and 987P by the Lactobacillus isolates from porcine intestine. *J Sci Food Agric* 6

840   (2000).

841   35.   Maltby, R., Leatham-Jensen, M. P., Gibson, T., Cohen, P. S. & Conway, T. Nutritional

842   basis for colonization resistance by human commensal Escherichia coli strains HS and

843   Nissle 1917 against E. coli O157: H7 in the mouse intestine. *PloS One* **8**, e53957 (2013).

844   36.   Suda, Y. *et al.* Immunobiotic Lactobacillus jensenii as immune-health promoting factor to

845   improve growth performance and productivity in post-weaning pigs. *BMC Immunol.* **15**, 24

846   (2014).

847   37.   Shimazu, T. *et al.* Immunobiotic Lactobacillus jensenii Elicits Anti-Inflammatory Activity in

848   Porcine Intestinal Epithelial Cells by Modulating Negative Regulators of the Toll-Like

849   Receptor Signaling Pathway. *Infect. Immun.* **80**, 276–288 (2012).

850   38.   Splichalova, A. *et al.* Interference of Bifidobacterium choerinum or Escherichia coli Nissle

851   1917 with Salmonella Typhimurium in gnotobiotic piglets correlates with cytokine patterns

852   in blood and intestine. *Clin. Exp. Immunol.* **163**, 242–249 (2011).

853   39.   Roselli, M. *et al.* The novel porcine Lactobacillus sobrius strain protects intestinal cells from

854   enterotoxigenic Escherichia coli K88 infection and prevents membrane barrier damage. *J.*

855   *Nutr.* **137**, 2709–2716 (2007).

856   40.   Bomba, A., Nemcová, R., Gancarčíková, S., Herich, R. & Kaštel, R. Potentiation of the

857   effectiveness of Lactobacillus casei in the prevention of E. coli induced diarrhea in

858   conventional and gnotobiotic pigs. in *Mechanisms in the pathogenesis of enteric diseases*

859   *2* 185–190 (Springer, 1999).

860   41.   Nemcova, R., Bomba, A., Gancarcikova, S., Herich, R. & Guba, P. Study of the effect of

861   Lactobacillus paracasei and fructooligosaccharides on the faecal microflora in weanling

862   piglets. *Berl. Munch. Tierarztl. Wochenschr.* **112**, 225–228 (1999).

863   42.   Estrada, A., Drew, M. D. & Van Kessel, A. Effect of the dietary supplementation of

864   fructooligosaccharides and Bifidobacterium longum to early-weaned pigs on performance

865   and fecal bacterial populations. *Can. J. Anim. Sci.* **81**, 141–148 (2001).

866   43.   Huang, C., Qiao, S., Li, D., Piao, X. & Ren, J. Effects of Lactobacilli on the performance,

867   diarrhea incidence, VFA concentration and gastrointestinal microbial flora of weaning pigs.

868   *Asian-Australas. J. Anim. Sci.* **17**, 401–409 (2004).

869   44.   De Angelis, M. *et al.* Survival and persistence of Lactobacillus plantarum 4.1 and

870   Lactobacillus reuteri 3S7 in the gastrointestinal tract of pigs. *Vet. Microbiol.* **123**, 133–144

871   (2007).

872   45.   Šmajs, D. *et al.* Experimental Administration of the Probiotic Escherichia coli Strain Nissle

873   1917 Results in Decreased Diversity of E. coli Strains in Pigs. *Curr. Microbiol.* **64**, 205–210

874   (2012).

875   46.   Liu, H. *et al.* Oral administration of Lactobacillus fermentum I5007 favors intestinal

876   development and alters the intestinal microbiota in formula-fed piglets. *J. Agric. Food*

877   *Chem.* **62**, 860–866 (2014).

878   47.   Zhao, P. & Kim, I. Effect of direct-fed microbial on growth performance, nutrient

879   digestibility, fecal noxious gas emission, fecal microbial flora and diarrhea score in

880   weanling pigs. *Anim. Feed Sci. Technol.* **200**, 86–92 (2015).

881   48.   Dowarah, R., Verma, A. K. & Agarwal, N. The use of Lactobacillus as an alternative of

882   antibiotic growth promoters in pigs: A review. *Anim. Nutr.* **3**, 1–6 (2017).

883   49.   Pieper, R. *et al.* Effect of a single oral administration of Lactobacillus plantarum DSMZ

884   8862/8866 before and at the time point of weaning on intestinal microbial communities in

885   piglets. *Int. J. Food Microbiol.* **130**, 227–232 (2009).

886   50.   Chae, J. P., Pajarillo, E. A. B., Oh, J. K., Kim, H. & Kang, D.-K. Revealing the combined

887   effects of lactulose and probiotic enterococci on the swine faecal microbiota using 454

888   pyrosequencing. *Microb. Biotechnol.* **9**, 486–495 (2016).

889   51.   Xu, J. *et al.* Yeast probiotics shape the gut microbiome and improve the health of early-

890   weaned piglets. *Front. Microbiol.* **9**, 2011 (2018).

891   52.   Solano-Aguilar, G. I. *et al.* The effect of feeding cocoa powder and Lactobacillus

892   rhamnosus on the composition and function of pig intestinal microbiome. *Curr. Dev. Nutr.*

893   **2**, nzy011 (2018).

894   53.   Shin, D. *et al.* Beneficial roles of probiotics on the modulation of gut microbiota and

895   immune response in pigs. *PloS One* **14**, e0220843 (2019).

896   54.   Gaio, D. *et al.* A large-scale metagenomic survey dataset of the post-weaning piglet gut

897   lumen. *GigaScience* (2021).

898   55.   Gaio, D. *et al.* Hackflex: low cost Illumina sequencing library construction for high sample

899   counts. *bioRxiv* 779215 (2019).

900   56.   Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize analysis results for

901   multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048 (2016).

902   57.   Darling, A. E. *et al.* PhyloSift: phylogenetic analysis of genomes and metagenomes. *PeerJ*

903   **2**, e243 (2014).

904   58.   Faith, D. P. Conservation evaluation and phylogenetic diversity. *Biol. Conserv.* **61**, 1–10

905   (1992).

59.   McCoy, C. O. & Matsen, F. A. Abundance-weighted phylogenetic diversity measures
distinguish microbial community states and are robust to sampling depth. *PeerJ* **1**, e157
(2013).
60.   Evans, S. N. & Matsen, F. A. The phylogenetic Kantorovich-Rubinstein metric for
environmental sequence samples. *J. R. Stat. Soc. Ser. B Stat. Methodol.* **74**, 569–592
(2012).
61.   R Core Team, R. C. T. R: A language and environment for statistical computing. (2013).
62.   Wickham, H., Hester, J. & Francois, R. *readr: Read Rectangular Text Data*. (2018).
63.   Wickham, H. & Bryan, J. *readxl: Read Excel Files*. (2019).
64.   Wickham, H., François, R., Henry, L. & Müller, K. *dplyr: A Grammar of Data Manipulation*.
(2019).
65.   Wickham, H. *et al.* Welcome to the tidyverse. *J. Open Source Softw.* **4**, 1686 (2019).
66.   Robinson, D. & Hayes, A. *broom: Convert Statistical Analysis Objects into Tidy Tibbles*.
(2019).
67.   Vu, V. Q. *ggbiplot: A ggplot2 based biplot*. (2011).
68.   Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. & R. Core Team. *nlme: Linear and
Nonlinear Mixed Effects Models*. (2019).
69.   Kassambara, A. *ggpubr: 'ggplot2' Based Publication Ready Plots*. (2019).
70.   Mahto, A. *splitstackshape: Stack and Reshape Datasets After Splitting Concatenated
Values*. (2019).
71.   Kassambara, A. *rstatix: Pipe-Friendly Framework for Basic Statistical Tests*. (2019).
72.   Henry, L. & Wickham, H. *purrr: Functional Programming Tools*. (2019).
73.   Dinno, A. *dunn.test: Dunn's Test of Multiple Comparisons Using Rank Sums*. (2017).
74.   Auguie, B. *gridExtra: Miscellaneous Functions for 'Grid' Graphics*. (2017).
75.   Bache, S. M. & Wickham, H. *magrittr: A Forward-Pipe Operator for R*. (2014).
76.   Bates, D. & Maechler, M. *Matrix: Sparse and Dense Matrix Classes and Methods*. (2019).

77.   Kopylova, E., Noé, L. & Touzet, H. SortMeRNA: fast and accurate filtering of ribosomal

933   RNAs in metatranscriptomic data. *Bioinformatics* **28**, 3211–3217 (2012).

934   78.   Johnson, W. E., Li, C. & Rabinovic, A. Adjusting batch effects in microarray expression

935   data using empirical Bayes methods. *Biostatistics* **8**, 118–127 (2007).

936   79.   Kembel, S. W. *et al.* Architectural design influences the diversity and structure of the built

937   environment microbiome. *ISME J.* **6**, 1469–1479 (2012).

938   80.   Pollock, J. *et al.* Analysis of temporal fecal microbiota dynamics in weaner pigs with and

939   without exposure to enterotoxigenic Escherichia coli. *J. Anim. Sci.* **96**, 3777–3790 (2018).

940   81.   Frese, S. A., Parker, K., Calvert, C. C. & Mills, D. A. Diet shapes the gut microbiome of

941   pigs during nursing and weaning. *Microbiome* **3**, 28 (2015).

942   82.   Meng, Q. *et al.* Weaning alters intestinal gene expression involved in nutrient metabolism

943   by shaping gut microbiota in pigs. *Front. Microbiol.* **11**, 694 (2020).

944   83.   Guevarra, R. B. *et al.* Piglet gut microbial shifts early in life: causes and effects. *J. Anim.*

945   *Sci. Biotechnol.* **10**, 1 (2019).

946   84.   Pajarillo, E. A. B., Chae, J. P., Kim, H. B., Kim, I. H. & Kang, D.-K. Barcoded

947   pyrosequencing-based metagenomic analysis of the faecal microbiome of three purebred

948   pig lines after cohabitation. *Appl. Microbiol. Biotechnol.* **99**, 5647–5656 (2015).

949   85.   Thompson, C. L., Wang, B. & Holmes, A. J. The immediate environment during postnatal

950   development has long-term impact on gut community structure in pigs. *ISME J.* **2**, 739

951   (2008).

952   86.   Xiao, L. *et al.* A reference gene catalogue of the pig gut microbiome. *Nat. Microbiol.* **1**,

953   16161 (2016).

954   87.   Bailey, M. *et al.* Regulation of mucosal immune responses in effector sites. *Proc. Nutr.*

955   *Soc.* **60**, 427–435 (2001).

956   88.   Wang, W., Hu, H., Zijlstra, R. T., Zheng, J. & Gänzle, M. G. Metagenomic reconstructions

957   of gut microbial metabolism in weanling pigs. *Microbiome* **7**, 1–11 (2019).

958   89.   Funkhouser, L. J. & Bordenstein, S. R. Mom knows best: the universality of maternal

959   microbial transmission. *PLoS Biol* **11**, e1001631 (2013).

960   90.   Motta, V., Luise, D., Bosi, P. & Trevisi, P. Faecal microbiota shift during weaning transition

961   in piglets and evaluation of AO blood types as shaping factor for the bacterial community

962   profile. *PloS One* **14**, e0217001 (2019).

963   91.   Ludlow, K. Australian Pesticides and Veterinary Medicines Authority. in *Encyclopedia of*

964   *Nanoscience and Society* (SAGE Publications, Inc., 2010).

965   doi:10.4135/9781412972093.n22.

966   92.   Button, K. S. *et al.* Power failure: why small sample size undermines the reliability of

967   neuroscience. *Nat. Rev. Neurosci.* **14**, 365–376 (2013).

968   93.   Dirnagl, U. *The p value wars (again)*. (Springer, 2019).

969   94.   Anderson, S. F. Misinterpreting p: The discrepancy between p values and the probability

970   the null hypothesis is true, the influence of multiple testing, and implications for the

971   replication crisis. *Psychol. Methods* **25**, 596 (2020).

972   95.   Deshmukh, H. S. *et al.* The microbiota regulates neutrophil homeostasis and host

973   resistance to Escherichia coli K1 sepsis in neonatal mice. *Nat. Med.* **20**, 524–530 (2014).

974   96.   Kim, P. I. *et al.* Probiotic properties of Lactobacillus and Bifidobacterium strains isolated

975   from porcine gastrointestinal tract. *Appl. Microbiol. Biotechnol.* **74**, 1103–1111 (2007).

976   97.   Zeineldin, M., Aldridge, B., Blair, B., Kancer, K. & Lowe, J. Impact of parenteral

977   antimicrobial administration on the structure and diversity of the fecal microbiota of growing

978   pigs. *Microb. Pathog.* **118**, 220–229 (2018).

979   98.   Valeriano, V. D. V., Balolong, M. P. & Kang, D. K. Probiotic roles of *Lactobacillus* sp. in

980   swine: insights from gut microbiota. *J. Appl. Microbiol.* **122**, 554–567 (2017).

981   99.   Abe, F., Ishibashi, N. & Shimamura, S. Effect of administration of bifidobacteria and lactic

982   acid bacteria to newborn calves and piglets. *J. Dairy Sci.* **78**, 2838–2846 (1995).

100. Davis, M. E. *et al.* Effect of a Bacillus-based direct-fed microbial feed supplement on

growth performance and pen cleaning characteristics of growing-finishing pigs. *J. Anim.*

Sci. **86**, 1459–1467 (2008).

101. Chu, G. M., Lee, S. J., Jeong, H. S. & Lee, S. S. Efficacy of probiotics from anaerobic

microflora with prebiotics on growth performance and noxious gas emission in growing

pigs: SYNBIOTICS ON NOXIOUS GAS EMISSION IN PIGS. *Anim. Sci. J.* **82**, 282–290

(2011).

102. Friedman, J. & Alm, E. J. Inferring Correlation Networks from Genomic Survey Data. *PLoS*

Comput. Biol. **8**, e1002687 (2012).

103. Lovell, D., Pawlowsky-Glahn, V., Egozcue, J. J., Marguerat, S. & Bähler, J. Proportionality:

A Valid Alternative to Correlation for Relative Data. *PLOS Comput. Biol.* **11**, e1004075

(2015).

104. Gaio, D. *et al.* Community composition and development of the post-weaning piglet gut

microbiome. *bioRxiv* 211326 (2020).

997
998
999