Community composition and development of the post-weaning piglet gut microbiome

D. Gaio[1], Matthew Z. DeMaere[1], Kay Anantanawat[1], Graeme J. Eamens[2], Michael Liu[1], Tiziana Zingali[1], Linda Falconer[2], Toni A. Chapman[2], Steven P. Djordjevic[1], Aaron E. Darling[1]*

[1]The ithree institute, University of Technology Sydney, Sydney, NSW, Australia

[2]NSW Department of Primary Industries, Elizabeth Macarthur Agricultural Institute, Menangle, NSW, Australia

*Corresponding author: Aaron E. Darling, aaron.darling@uts.edu.au

Keywords: shotgun metagenomics; pig; post-weaning; gut microbiome; phylogenetic diversity

ABSTRACT

We report on the largest metagenomic analysis of the pig gut microbiome to date. By processing over 800 faecal time-series samples from 126 piglets and 42 sows, we generated over 8Tbp of metagenomic shotgun sequence data. Here we describe the animal trial procedures, the generation of our metagenomic dataset and the analysis of the microbial community composition using a phylogenetic framework. We assess the effects of intramuscular antibiotic treatment and probiotic oral treatment on the diversity of gut microbial communities. We found differences between individual hosts such as breed, litter, and age, to be important contributors to variation in the community composition. Treatment effects of the antibiotic and probiotic treatments were found but were subtle, while host age was the dominant factor in shaping the gut microbiota of piglets after weaning. The post-weaning pig gut microbiome appears to follow a highly structured developmental program with characteristic post-weaning changes that can distinguish hosts that were born as little as two days apart in the second month of life.

INTRODUCTION

As the world population grows, there is an accompanying demand for animal derived products. In a semi-natural environment pig weaning occurs between the 12th and the 17th weeks from birth, whereas in the farm this typically occurs at 4 weeks of age[1]. Intensive animal husbandry and early weaning practices are commonly used to maximise production rates while minimizing costs. Both practices increase the risk of infections with pathogenic organisms, and thereby the need for antimicrobial strategies, which has included the common use of antibiotics. [2–12] Although antibiotics may kill some pathogens, the surviving bacteria can develop antimicrobial resistance (AMR) against the class of antibiotic used [13–20], as well as against other antibiotic classes [21–31]. A retrospective U.S. livestock study found evidence of multidrug resistance (resistance to >3 antimicrobial drug classes) in *Escherichia coli* that increased from 7.2% to 63.6% between the 1950s and the 2000s [32]. As the incidence of multidrug resistant (MDR) pathogens expands, the World Health Organization (WHO) has recognized AMR as one of the top health challenges of the current century [33,34].

Besides leading to AMR development, antibiotic usage is known to cause dysbiosis [34–37], the disruption of a balanced state within a gut microbial community. A balanced state decreases the chance of pathogens gaining a foothold, as there is niche and nutrient competition in play [39,40]. In livestock production, intramuscular (IM) antibiotic treatment is preferred over oral antibiotic treatment, as it is thought to contribute less to AMR than oral antibiotic treatment [41–43]. Extensive evidence exists on dysbiosis as a consequence of oral antibiotic treatment on the gut microbiome [22,35–38,44,45], whereas only a handful of studies report the effects of intramuscular (IM) antibiotic treatment on the gut microbial community [46–48], and none, to our knowledge have studied the use of IM neomycin treatment.

A number of non-antibiotic strategies to increase resistance to disease have been studied [49]. Strategies for which beneficial effects have been reported consist of hydrolases [50], fiber intake [51], β-lactamase enzymes [52], non-resistance inducing antimicrobial drugs [53,54], vaccination [55–58], phage therapy [59], in-feed organic acids [60], starches [61], and liquid feed [62–64]. Extensive evidence exists on the use of probiotics. Probiotics are reported to promote intestinal health in multiple ways: improvement of mucosal integrity [65–68], competitive exclusion against pathogenic species [50,69–73], reduction of intestinal inflammation [65,74,75] and of pathogen translocation [65,76,77]. Their efficacy has been determined, among others, in cell cultures [70,74,75,78], mice [50,67,68,73], and swine [65,66,74,76,79–83]. However, while most evidence of beneficial effects from probiotic treatment is derived from host immunity responses [65–68,74–

[76,80], host physiology [74,80–82], and pathogen relocation [50,65,69,70,72,77–79,84], fewer studies exist on

the effects of probiotics on the whole gut microbial community [83,85].

This study was conducted to: 1) generate a public metagenomic databank of the gut

microbiome of weaner piglets aged between 3 and 9 weeks old; 2) assess the effects of IM

neomycin antibiotic use on the gut community; and 3) assess the effects of two probiotic

formulations on the gut community. The data we present here is analysed from the

perspective of the phylogenetic diversity of the microbial communities.

In the first results section we describe the phylogenetic diversity of the intestinal

microbiome of the piglets and how it compares to the known composition of positive

controls. Then, we describe evidence that highlights the importance of age, breed and litter

as strongly associated factors with changes in community composition. Lastly, we describe

the strong effect of time on community composition and the milder effect of antibiotic and

probiotic treatments on community composition.

METHODS

Pig trial

Animal studies were conducted at the Elizabeth Macarthur Agricultural Institute (EMAI)

NSW, Australia and were approved by the EMAI Ethics Committee (Approval M16/04). The

trial animals comprised 4-week old male weaner pigs (*n*=126) derived from a commercial

swine farm and transferred to the study facility in January 2017. These were cross-bred

animals of Landrace, Duroc and Large White breeds and had been weaned at approximately

3 weeks of age.

The pig facility consisted of four environmentally controlled rooms (Rooms 1-4) with

air conditioning, concrete slatted block flooring with underground drainage and open rung

steel pens (**S1**). Each room had nine pens, consisting of a set of six and a set of three pens,

designated a-f and g-i respectively, with the two sets of pens being physically separate, *i.e.*

animals could come in contact with each other through the pen's bars within each set of

pens, but not between sets. The rooms were physically separated by concrete walls and

contamination between rooms was minimized by using separate equipment (boots, gloves,

coveralls) for each room. In addition, under-floor drainage was flushed twice weekly and the

flushed faeces/urine was retained in under-floor channels that ran the length of the facility,
so that Rooms 1, 2 were separate from Rooms 3, 4 and flushing was in the direction 1 to 2
and 3 to 4.

The pigs were fed *ad libitum* a commercial pig grower mix of 17.95% protein free of
antibiotics, via self-feeders. On the day of arrival (day 1) 30, 18, 18, and 60 pigs were
allocated randomly to Rooms 1, 2, 3 and 4 respectively in groups of 6, 6, 6 and 6-7 pigs per
pen respectively (**S1A**). Pigs were initially weighed on day 2, and some pigs were moved
between pens to achieve an initial mean pig weight per treatment of approximately 6.5 kg
(range: 6.48-6.70; mean±SD: 6.53±0.08). Pigs were weighed weekly throughout the trial
(**Supplementary file 1**). Behaviour and faecal consistency scores were taken daily over the
6-week period of the trial (**Supplementary file 2**). Developmental and commercial probiotic
paste preparations ColiGuard® and D-Scour™ from International Animal Health,
Australasia, were used in some treatment groups.

The animals were acclimatised for 2 days before the following treatments were
administered: Room 1 - oral 1 g/pig of placebo paste daily for 14 d; Room 2 - oral 1 g/pig of
D-Scour™ paste daily for 14 d; Room 3 - oral 1 g/pig of ColiGuard® paste daily for 14 d;
Room 4 - intramuscular (IM) injection of antibiotic administered at 0.1 mL per pig daily
from a 200 mg/mL solution for a total treatment duration of 5 d.

On the day following the final neomycin treatment (day 8), 36 pigs were moved from
Room 4 to Room 2 (*n*=18, 6 in each pen, pens g-i), and to Room 3 (*n*=18, 6 in each pen, pens
g-i) (**S1**). The following day (day 9), oral administration of D-Scour™ (1 g/pig) and of
ColiGuard® (1 g/pig) commenced for pigs in Room 2 pens g-i and in Room 3 pens g-i,
respectively, and continued for a period of 14 days. Assignment of the 36 neomycin-treated
pigs to the treatment groups neomycin+D-Scour™ (*n*=18; Room 2 pens g-i) and
neomycin+ColiGuard® (*n*=18; Room 3 pens g-i), was carried out by distributing them so
that the mean weight of the animals distributed across pens and rooms was similar
(**Supplementary file 1**). By this time point, each occupied pen in the trial housed six pigs.
(**S1B**) From that time, twelve piglets from the original 126 were no longer present, as they
had been euthanised as pre-treatment controls at the start of the trial.

Faecal samples were collected from all piglets once per week and from a subset (*n*=48
pigs; 8 from each of the six cohorts) twice per week over the 6-week study period (**Figure

1). From each piglet, faeces were collected per rectum with new disposable gloves; where

minimal or no faeces could be collected on a collection day, sampling was performed the

following morning. Samples were placed in 50 mL Falcon tubes and stored at 4ºC within 30

mins of sample collection for a minimum of 30 mins and a maximum period of 6 h.

Faecal sample processing

Samples (3g/pig) were mixed with 15 mL PBS (200 mg/mL), in sterile stomacher bags

and homogenized with a Bio-Rad stomacher. The homogenised samples were divided in

replicates: one replicate was stored directly at -80ºC and one replicate was supplemented

with glycerol (20% v/v) (Sigma-Aldrich) then stored at -80ºC. In addition, single time-point

faecal samples from the dams of the trial pigs (n=42) were obtained from the commercial

facility of origin and were pre-processed at EMAI as described above. Thus, a total of 911

unique samples, between one and ten samples per subject (mean: 4.8; median: 3) (**S2**), were

obtained throughout this study. At the end of the trial period, all samples were transported

from EMAI to the University of Technology Sydney (UTS) for further processing.

Positive controls

As a positive control "mock community" for this study, four Gram positive (*Bacillus*

subtilis strain 168, *Enterococcus faecium*, *Staphylococcus aureus* ATCC25923,

Staphylococcus epidermidis ATCC35983) and three Gram negative (*Enterobacter*

hormaechei CP_032842, *Escherchia coli* K-12 MG1655, *Pseudomonas aeruginosa* PAO1)

bacterial strains from -80°C stocks were cultured at 37ºC for 16 h in LB (Luria-Bertani) then

centrifuged at 14,000 rpm for 10 mins. From the resulting pellets, 1 g was transferred to 1

mL of LB and homogenised and a 1:10 dilution of this was made for each bacterial culture.

Ten microliters of bacterial suspension from each of the cultures was used to determine the

number of colony forming units (CFU) in the original suspension in the following manner:

by further diluting tenfold in LB and by plating onto 1.6% LB agar plates and incubacted

overnight. The remaining suspensions (990 μL from each bacterial culture) were pooled into

a sterile tube, then aliquoted into Eppendorf tubes in 500 μL volumes/tube. As a washing

step, Eppendorf tubes were centrifuged at 14,000 rpm for 10 mins, 500 μL PBS was added to

the pellet and subsequently resuspended. These tubes constituted the mock community

samples and were stored at -80ºC. Expected proportions of the mock community members

were determined from the estimated colony forming units (CFU) multiplied by the genome

size and were as follows: 8.7:13.0:7.7:16.7:38.9:14.5:0.4 for *S. aureus*, *B. subtilis*, *E.*
faecium, S. epidermidis, P. aeruginosa, E. cloacae, and E.coli respectively.

The two probiotic formulations used in this study were used as two additional positive
controls. D-Scour™ is a commercially available probiotic formulation for livestock, with
each gram containing 180 million CFU of the following: *Lactobacillus acidophilus*,
Lactobacillus delbrueckii subspecies *bulgaricus*, *Lactobacillus plantarum*, *Lactobacillus*
rhamnosus, *Bifidobacterium bifidum*, *Enterococcus faecium*, *Streptococcus salivarius*
subspecies *thermophilus*, with an additional 20 mg of garlic extract (*Allium sativum*). The
probiotic ColiGuard is a probiotic formulation developed for the treatment of entero-
toxigenic *Escherichia coli* (ETEC) in weaner pigs, developed in collaboration between the
NSW DPI and International Animal Health Products, containing undefined concentrations of
Lactobacillus plantarum and *Lactobacillus salivarius*.

DNA extraction

Piglet and sow faecal samples, mock community samples, negative controls and
probiotic samples (D-Scour™ and ColiGuard® paste) were allocated to a randomized block
design to control for batch effects in DNA extraction and library preparation. The faecal
samples were thawed on ice first, followed by the probiotics and mock community samples.
MetaPolyzyme (Sigma-Aldrich) treatment was performed according to the manufacturer's
instructions except for the dilution factor, which we allowed to be 4.6 times higher.
Immediately after incubation, DNA extraction was performed with the MagAttract
PowerMicrobiome DNA/RNA EP kit (Qiagen) according to the manufacturer's instructions.
Quantification of DNA was performed using PicoGreen (Thermofisher) and measurements
were performed with a plate reader (Tecan, Life Sciences) using 50 and 80 gain settings. All
samples were diluted to 10 ng/μL.

Library preparation

Sample index barcode design using a previously introduced method [86] yielded a set of 96
x 8nt sequences with a 0.5 mean GC content and none of the barcodes containing 3 or more
identical bases in a row. Nine hundred-sixty different combinations of i5 and i7 primers were
used to create a uniquely barcoded library for each sample. The detailed sample-to-barcode
assignment is given in **Supplementary file 3**. Library preparation was carried out using a
modification of the Nextera Flex protocol to produce low bias, low cost shotgun libraries, as

described in a previous manuscript [86]. Following the amplification step, samples were
centrifuged at 280 x *g* for 1 min and stored between 1 and 5 days at 4°C.

Size selection and purification

Samples from the same 96-well plates were pooled into one tube by taking 5 μL from
each library. This generated 10 pooled samples, one for each plate. A master pool was
created by pooling 5 μL from the pool of each plate into a single pool. Forty microliters from
each of the 10 plate pools and 40 μL from the master pool underwent library size selection
and purification using equal volumes of SPRIselect beads (Beckman Coulter) and ultrapure
water (Invitrogen). Sample cleaning with SPRI-beads was performed as described previously
[86]. A purified master pool comprising samples from all plates, and purified pools of
individual plates to check for plate-specific anomalies, were diluted to 4 nM and fragment
size distribution was assessed using the High Sensitivity DNA kit on the Bioanalyzer
(Agilent Technologies, USA).

Normalisation and sequencing

The master pool was sequenced on an Illumina MiSeq v2 300 cycle nano flow cell
(Illumina, USA). Read counts were obtained and used to normalise libraries. The liquid
handling robot OT-One (Opentrons) was programmed to re-pool libraries based on read
counts obtained from the previous MiSeq run. The script used to achieve the normalization is
available through our Github repository.

The read count distribution after normalisation is displayed in supplementary figure
(**S3**). The normalized and purified pooled library was sequenced on an Illumina NovaSeq S4
flow cell at the Ramaciotti Centre for Genomics (Sydney, NSW, Australia), generating a
total of 27 billion read pairs from 911 samples.

Sequence data processing

Adapter trimming (parameters: k=23 hdist=1 tpe tbo mink=11), PhiX DNA removal
(parameters: k=31 hdist=1), and quality filtering (parameters: ftm=0 qtrim=r trimq=20), were
performed using bbduk.sh (http://jgi.doe.gov/data-and-tools/bbtools). Quality assessment of
raw   reads   was   carried   out   using   FASTQC
(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and a combined report of all
samples was obtained with MULTIQC [87] (**Supplementary File 4**). The presence of PCR

duplicates was assessed by feeding read pairs to dedupe.sh (http://jgi.doe.gov/data-and-

tools/bbtools) (parameters: ac=f). Nextflow [88] was used to manage processing of the data on

the HPC.

Determination of microbial diversity among samples

Phylogenetic diversity of all samples was assessed with PhyloSift [89] using the first 100k

read pairs of each sample (parameters: --chunk-size 100000 --paired). In order to test for

associations of phylogenetic diversity with treatment, time of sampling, and differences

among hosts at the start of the trial, analysis of the unrooted phylogenetic diversity (PD) [90],

the balance weighted phylogenetic diversity (BWPD) [91] and principal component analysis

(PCA) of the Kantorovich-Rubinstein distances [92] (beta diversity analysis) were performed.

Alpha-diversity and beta-diversity were analyzed and the results were visualized with R [93]

and R packages [91–107]. The data analysis is schematically represented in **Supplementary**

Figure 4 (**S4**).

Additionally, SortmeRNA [111] (version 2.1) was applied to extract 16S ribosomal RNA

genes from raw reads. The rRNA reference database silva-bac-16s-id90 was used. Hits were

filtered (e-value <= 1e-30) and PCA was computed with R [93]. Sample counts were

normalized for library size by proportions and were tested with the Spearman's Rank

correlation coefficient method to find taxa correlating with the weight of the piglets across

the trial.

Batch effects

A randomized block design was adopted to mitigate batch effects. Because samples

were distributed across ten 96-well plates during DNA extraction and library preparation,

plate effects were expected. Although samples did not visibly cluster by DNA extraction

plate across the first five principal components (**S5**), a batch effect was found by multiple

comparison analysis with ANOVA and by applying Tukey post-hoc correction to pairwise

comparisons. Batch effects were detected (ANOVA, alpha diversity: p range=4.8e-12-

0.00011; beta diversity: p value<0.0023) (**S6**; **Supplementary file 5**) and removed with

COMBAT[7] (ANOVA p value=1) (**S7**; **Supplementary file 5**).

Technical analysis of positive controls

As a quality control step, taxonomic analysis of the positive control samples and their

technical replicates was performed with MetaPhlAn2 [112]. The positive controls included the

mock community (*n*=8), D-Scour™ (*n*=8), and ColiGuard® (*n*=8) samples.

Data availability

The raw sequencing data has been submitted to the NCBI Short Read Archive under

project PRJNA526405. Metadata and scripts are available in this article, its Supplementary

material and in our Github repository https://github.com/GaioDany/metapigs_base

RESULTS

PhyloSift [89] was employed as a means to study microbial community diversity among

the samples, and to test for associations with treatment, time of sampling, and differences

among hosts at the start of the trial. To this end, analysis of the unrooted phylogenetic

diversity (PD) [90], the balance weighted phylogenetic diversity (BWPD) [91] and principal

component analysis (PCA) of the Kantorovich-Rubinstein distances [92] (beta diversity

analysis) were performed.

Comparison of the expected and the observed taxonomic profile of the positive controls

All the mock community members, in seven of the eight technical replicates, were

detected by MetaPhlAn2 (**S8**). One sample failed to sequence, reporting zero counts for any

species. The observed mean relative abundances were as follows: *B. subtilis* (mean±SD:

2.92±0.994), *E. cloacae* (mean±SD: 38.0±6.404), *E. faecium* (mean±SD: 0.97±0.081), *E.coli*

(mean±SD: 10.12±1.480), *E.coli* unclassified (mean±SD: 7.83± 1.755), *P. aeruginosa*

(mean±SD: 26.72±3.026), *S. aureus* (mean±SD: 9.90±3.613), *S. epidermidis* (mean±SD:

3.54±1.435). Isolate *E. cloacae* C15117, used in this study for the make-up of the mock

community, was recently found to be most closely related to the *Enterobacter hormaechei*

phylogenomic group C type strain DSM 16687 and therefore re-identified as *Enterobacter*

hormaechei subsp. *oharae*[113] . For this reason, taxonomic assignment by MetaPhlAn2

attributed the reads to *E. cloacae* instead. The expected proportions of the mock community

members were derived from the CFU by the genome size. Based on the expected and the

observed relative abundance, we found, with the exception of *S. aureus* (exp: 8.7% obs:

9.9%), three Gram positive members to be under-represented (*B. subtilis*: exp: 13.0% obs:

2.9%; *E. faecium:* exp: 7.7% obs: 1.0%; *S. epidermidis*: exp: 16.7% obs: 3.5%) and, with the

exception of *P. aeruginosa* (exp: 38.9% obs: 26.8%), two Gram negative members to be

over-represented (*E. cloacae:* exp: 14.5% obs: 38.0%; *E.coli*: exp: 0.4% obs: 7.8-10.1%)

(**S9**). Taxonomic assignment of the mock community samples reported one contaminating

species in one of the eight replicates: *Lactobacillus salivarius* (mean: 0.008) (**S10**).

The probiotic D-Scour™ is expected to contain, per gram, a total of 180 million CFU of

Lactobacillus acidophilus, *Lactobacillus delbrueckii* subspecies *bulgaricus*, *Lactobacillus*

plantarum, *Lactobacillus rhamnosus*, *Bifidobacterium bifidum*, *Enterococcus faecium*,

Streptococcus salivarius subspecies *thermophilus* in unknown proportions. From taxonomic

analysis with MetaPhlAn2, we can conclude that 6 of the 7 expected species were

determined to be present in the replicates in the following mean relative abundances:

Bifidobacterium bifidum: mean±SD: 40.01±12.558; *Enterococcus faecium*: mean±SD:

30.98±13.472; *Lactobacillus delbrueckii*: mean±SD: 11.56±7.148; *Lactobacillus plantarum*:

mean±SD: 6.23±7.863; *Lactobacillus rhamnosus*: mean±SD: 2.08±1.226; *Streptococcus*

thermophilus: mean±SD: 4.28±1.523. *Lactobacillus acidophilus* was not detected and

Lactobacillus helveticus was detected instead (*Lactobacillus helveticus*: mean±SD:

4.75±2.431) (**S8**). An additional 25 taxa were detected, of which 18 and 7 were identified at

the species and at the genus level, respectively. Contaminants were present at a higher

concentration in three technical replicates (R3, R7, R8) with the most frequent contaminant

(*Methanobrevibacter* spp.) being present in 5 of the 8 replicates (**S11**).

Taxonomic analysis of the technical replicates of the probiotic ColiGuard® also showed

a species profile consistent with the expected profile, with *Lactobacillus salivarius* and

Lactobacillus plantarum in a 9:1 ratio (*Lactobacillus salivarius*: mean±SD: 93.52±1.617;

Lactobacillus plantarum: mean±SD: 6.10±1.134) across the replicates (**S8**). ColiGuard®

contained a total of 20 contaminants, of which 16 and 4 were identified at the species and the

genus level, respectively. Contaminants were present at a higher level in two technical

replicates (R5, R7), with R7 displaying the most diverse and highest contamination rate (R7:

14 taxa; total contaminating reads: 2.67%; R5: 9 taxa; total contaminating reads: 0.30%).

(**S12**).

Phylogenetic diversity of positive controls and how it compares to the taxonomic profile

Alpha diversity of the positive controls reflects the expected alpha diversity, with the
mock community, ColiGuard® and D-Scour™ positive controls, displaying a progressively
higher unrooted PD (mean±SD: Mock community: 31.53±29.50; ColiGuard®: 58.52±21.70;
D-Scour™: 64.84±21.30). BWPD for the positive control ColiGuard® appears on the far left
of the plot (mean±SD: 0.82±0.15), distant from the mock community (mean±SD: 1.58±0.12)
and the positive control D-Scour™ (mean±SD: 1.89±0.26), which contains 2, 7 and 8 main
species, respectively (**S13**; **Supplementary file 1**). The unrooted PD of the positive control
ColiGuard® (mean±SD: 58.52±21.70) is closer to the unrooted PD of the Mock Community
(mean±SD: 31.53±29.50) and to that of the positive control D-Scour™ (mean±SD:
64.84±21.30), than the BWPD.

The contribution to total diversity of phylogenetic tree edges with uneven quantities of
reads placed on either side is down-weighted in BWPD, which is reflected in the low
reported BWPD of the positive control ColiGuard®. ColiGuard® is mainly composed of two
species in an uneven ratio (1:9) as by taxonomic analysis with MetaPhlAn2 (**S8**), in contrast
with the mock community (0.01:1:3:4:8:10:10:27:38) and the positive control D-Scour™
(2.08:4.28:4.75:6.23:11.56:30.98:40.01) (**S8**). On the other hand, the unrooted PD reflects
the absolute diversity, independently of the relative abundance of each species, within a
sample. In fact, low-level contamination (<0.1%) detected in each of the positive controls
(Mock Community: 1 taxon; ColiGuard®: 20 taxa; D-Scour™: 25 taxa) (**S11-13**) contributes
toward the absolute diversity in the unrooted PD, erroneously inflating this value (mean±SD:
Mock Community: 31.53±29.50; ColiGuard®: 58.52±21.70; D-Scour™: 64.84±21.30)
(**Figure 2**).

Phylogenetic diversity of piglet gut communities immediately after weaning

The 126 piglets used in this study were derived from 4 main breed cross types "Duroc ×
Landrace" (n=46), "Duroc × Large White" (n=59), "Landrace × cross bred (LWxD)" (n=9),
"Large White × Duroc" (n=12), and three pig lines (line 319: n=9; line 316: n=46; line 326:
n=71). The piglets also differed slightly by age, being born between 1 and 6 days apart.

Based on Kruskal-Wallis one-way analysis of variance (Hommel adjusted p values to
correct for multiple testing), phylogenetic diversity of the piglet samples did not cluster
significantly by breed cross type ($p > 0.05$) or by pig line ($p > 0.05$) at the start of the trial,
but a significant difference was found with breed and line in the fourth week of the trial

(breed: $p=0.04$; pig line: $p=0.02$) (**Figure 2**). Additionally a correlation of breed cross type

was found with beta diversity principal components (PC) in the second week of the trial

(PC3: $p=0.0357$) and in the fifth week of the trial (PC1: $p=0.027$). Alpha diversity

significantly correlated with pig line in the fourth week of the trial (BWPD: $p=0.017$).

(**Figure 2**; **Supplementary file 5**)

Notably, we found a significant correlation between alpha diversity and the age of the

piglets at the first sampling time point (unrooted PD: $p=0.0016$; BWPD: $p=0.0355$) (**Figure**

2; **Supplementary file 5**). While a correlation with age was found for alpha diversity only at

the start of the trial, age of the piglet was observed to be weakly associated with differences

in community composition in week two (PC3: $p=0.0507$) (**S14**; **Supplementary file 5**).

As age groups were confounded with cross-breed types (*i.e.* not all age groups are

represented by each of the four cross-breed types), we compared the phylogenetic diversity

of age groups within each breed. As cross-breed types "Landrace × cross bred (LW × D)"

and "Large White × Duroc" had only a small number of piglets in each age group, we tested

for an association between phylogenetic diversity and age in cross-breeds "Duroc ×

Landrace" and "Duroc × Large White". In these cross-breeds, alpha and beta diversity are

correlated with age in the "Duroc × Landrace" piglets (unrooted PD: $p=0.0059$; BWPD:

$p=0.0226$; PC2: $p=0.0263$; PC5: $p=0.0063$) and to a lesser extent in the "Duroc × Large

White" piglets (unrooted PD: $p=0.0529$; PC5: $p=0.0310$) during the first week of the trial. In

the "Duroc × Landrace" piglets, a correlation between age and beta diversity was detected in

week two (PC2: $p=0.0347$) (**Figure 2**; **S14**; **Supplementary file 5**). Differences in beta

diversity by date of birth, within the same cross-breed (Duroc × Large white), were found to

be significant (TukeyHSD adjusted p value: 1 day difference: 0.000782; 2 days difference:

0.018603) (**Supplementary file 5**).

Maternal effect on phylogenetic diversity

As piglets were derived from 42 distinct sows (maternal sows), and nursed by either the

same or a different sow (a nurse sow) (**Supplementary File 5**), a litter effect was expected

and determined. Based on Hommel adjusted p values, a similarity of alpha and beta

phylogenetic diversity can be seen among piglets from the same maternal sow (unrooted PD:

$p=0.0096$; BWPD: $p=0.0467$) and in piglets from the same nurse sow (unrooted PD:

$p=0.0320$; PC1: $p=0.0391$) (**Figure 2**; **S16-19**; **Supplementary file 5**). The litter effect

described was found in samples at the start of the trial. Significance of the correlations did

not persist thereafter (**Figure 2**; **Supplementary file 5**).

A strong effect of aging on phylogenetic diversity

Beta diversity analysis revealed a distinct and consistent change of the microbial

community over time in all piglets, regardless of the treatment. Samples clustered in PC2

(accounting for 21.68% of the variation), showing a higher representation of *Bacteroidetes*,

Gammaproteobacteria and *Prevotellaceae* from day 0-20, and a higher representation of

Firmicutes, *Mollicutes* and *Ruminococcaceae* from day 20-40 (**Figure 3**; **S20**). In PC1

(accounting for 46.99% of the variation) samples shifted towards a higher representation of

Lactobacilli from day 0 to day 20, and towards a higher representation of *Actinobacteria*,

Clostridiales and *Mollicutes* from day 20 to day 40 (**Figure 3**; **Supplementary file 5**). A

temporary shift towards a higher representation of *Bifidobacteriaceae* (*Bifidobacterium*

being a component of the probiotic D-Scour™) is seen in PC5 (3.04% of the variation

explained) one and two weeks from the start of the trial (**S21**).

Beta diversity analysis was performed separately for samples within each time point in

order to find taxa associated with variation within each time point. Extent of variation was

derived from the product of branch width by the variation explained by the principal

component (**S22**). The following taxa were responsible for variation only at the start of the

trial: *Gammaproteobacteria* (t0=0.12), *Enterobacteriaceae* (t0=0.08) and *Archaea* (t0=0.03).

The following taxa were responsible for variation throughout the trial: *Clostridiales*

(min=0.11; max=0.25), *Bacteroidetes* (min=0.06; max=0.11), *Bacteroidetes chlorobi* group

(min=0.05; max=0.10). *Lactobacillus* became variable after the first week of the trial and

remained highly variable throughout the trial (min=0.13; max=0.22). Some taxa contributed

less to the variability among the piglets' faecal population, but they nonetheless consistently

contributed to variability: *Bacteroidales* (min=0.01; max=0.04); *Firmicutes* (min=0.05;

max=0.08); *Bacilli* (min=0.06; max=0.09); *Lactobacillales* (min=0.04; max=0.08);

Actinobacteria (min=0.02; max=0.09); *Proteobacteria* (min=0.02; max=0.07). The

variability of *Prevotella* (min=0.01; max=0.04) increased from the second week of the trial.

(**S22**)

Taxonomic representation in terms of abundance was derived from analysis of samples

with guppy fat, where abundance was derived from the branch width (**Figure 4**) and

combined with PhyloSift's taxonomic annotation of the marker gene phylogeny. Overall,
Clostridiales were the most abundant throughout the trial (t0=0.18; t2=0.20; t4=0.11;
t6=0.09; t8=0.10; t10=0.08). *Lactobacillus acidophilus* increased at the start (t0=0.00;
t2=0.05; t4=0.13) then decreased (t6=0.09; t8=0.05; t10=0.07). *Subdoligranulum variabile*
decreased after the first week (t0=0.05; t2=0.02) then gradually increased (t4=0.04; t6=0.05;
t8=0.06; t10=0.07). Among other prevalent taxa at the start of the trial up to the second week
and dropping at later time points we found *Metahobrevibacter smithii* (t0=0.06; t2=0.03) and
Bacteroidales (t0=0.02; t2=0.01). Following an opposite trend we found *Ruminococcus* sp.
JC304 (t0=0.00; t2=0.00; t4=0.01; t6=0.02; t8=0.02; t10=0.03), *Solobacterium moorei*
(t0=0.00; t2=0.01; t4=0.02; t6=0.02; t8=0.02; t10=0.02) and *Prevotella copri* (t0=0.00;
t2=0.00; t4=0.04; t6=0.03; t8=0.05; t10=0.02). In modest and stable abundance across the
trial were *Mogibacterium* sp. CM50 (t0=0.02; t2=0.03; t4=0.02; t6=0.03; t8=0.02; t10=0.04)
and *Oscillibacter* (t0=0.06; t2=0.04; t4=0.03; t6=0.03; t8=0.03; t10=0.03). In transient
abundance we found *Bifidobacterium thermophilum* RBL67 (t2=0.01; t4=0.02). Gradually
increasing from the second week we found *Eubacterium biforme* DSM3989 (t2=0.01;
t4=0.02; t6=0.02; t8=0.02; t10=0.02), *Eubacterium rectale* (t4=0.02; t6=0.02; t8=0.03;
t10=0.01) and, after the third week, *Faecalibacterium prausnitsii* (t6=0.01; t8=0.02;
t10=0.02).

The effect of time was also clear in alpha diversity, where all the piglet cohorts followed
a similar trend over time, independent of the treatment (**S23**; **Supplementary file 5**). Upon
arrival on the trial site, the piglets' unrooted PD was lower than the sows (sows: 125.39;
piglets range: 104.6-112.7) and reached a similar unrooted PD to the sows' in the second
week (range: 123.4-130.7) (**Supplementary file 5**).

In comparing four timepoints at one week intervals from the start of the trial (intervals
labeled as A, B, C, D), changes in alpha diversity among all the piglets were tested for and
significance was determined using the Bonferroni correction. Unrooted phylogenetic
diversity increased after the start of the trial (A mean: 107.86; B mean: 127.21; $p<0.001$),
decreased in the following week (C mean: 122.92; p=ns) and the fourth week (D mean:
118.74; p=ns) amongst the piglets cohorts. In contrast, BWPD decreased after the start of the
trial (A mean: 2.17; B mean: 2.05; $p<0.0001$), to increase in the following week (C mean:
2.14; $p<0.0001$) and decrease in the fourth week (D mean: 2.07; $p<0.01$). (**S23**;
Supplementary file 5). The increase in unrooted PD in the first week was significant for all

treatment cohorts (*p* range: 0.0015-0.03) except ColiGuard® and neomycin (*p*=ns) cohorts.

The decrease in BWPD in the first week was only significant for the control (*p*=0.04),

ColiGuard® (*p*=0.0015) and neomycin+ColiGuard® (*p*=0.036) cohorts. In week two only

the increase of BWPD of the control cohort reached statistical significance (*p*=0.0121). (**S24**;

Supplementary file 5)

Effect of antibiotic and probiotic treatment on alpha diversity

We hypothesized that the probiotic treatments, whether alone (D-Scour™ and

ColiGuard®) or administered after neomycin (neomycin+D-Scour™ and

neomycin+ColiGuard®) would cause a change in the microbial community composition that

would be measurable via phylogenetic diversity. We tested whether the treatments correlated

with a change in phylogenetic diversity independently of the changes occurring with

time. Given the differences in alpha and beta diversity detected among the subjects at the

start of the trial, we analyzed the deltas of phylogenetic diversity instead of relying on the

absolute means, similar to the procedure applied by Kembel *et al* (2012) [114]. Time-point

measurements of alpha diversity were taken and deltas were computed for each piglet. Delta

means were compared between cohorts, where the control cohort would serve as a control

group for neomycin, D-Scour™ and ColiGuard® cohorts, whereas the neomycin cohort

would serve as a control group for the neomycin+D-Scour™ and neomycin+ColiGuard®

cohorts.

After the first week of the trial, the majority of the piglets displayed an increase in

unrooted PD (85%) and a decrease of BWPD (75%). The following week the trend was

opposite: an increase of BWPD (78%) and a decrease of unrooted PD (60%)

(**Supplementary file 5**). However, the neomycin cohort displayed the smallest BWPD drop

in the week following the start of the trial and the overall trend of neomycin in unrooted PD

was the most different from the other cohorts (**S25**). Due to the delayed rise in unrooted PD

of neomycin, significance was found in unrooted PD between neomycin and neomycin+D-

Scour™ during the first week of D-Scour™ treatment (neomycin mean: 1.59; neomycin+D-

Scour™ mean: -11.79; *p*= 0.012188) (**S25 B-C**) and in unrooted PD between neomycin and

neomycin+ColiGuard® (neomycin mean: -5.53; neomycin+ColiGuard®: -6.70; *p*=0.043138)

(**S25 C-D**). The change was opposite in BWPD (*p*=0.049263) (**S26 B-C**) and persisted

during the following week, up to 3 days post- D-Scour™ treatment (neomycin mean: -6.34;

neomycin+D-Scour™ mean: 4.52; BH adjusted *p*=0.024092) (**S26 B-D**).

The only significant difference between the neomycin and the control cohort was
detected in BWPD in the second week of the trial (control mean: 7.10; neomycin mean:
0.28) p=0.033751) (**S26 B-C**).

No significant differences were detected between the control and the D-Scour™ cohort
(p<0.05) (**Supplementary file 5**). Instead, a significant decrease in BWPD was detected in
the ColiGuard® cohort between the start of the trial and the end of the ColiGuard®
treatment, where nearly all piglets (88%) in the ColiGuard® cohort displayed a decrease in
BWPD, whereas the control cohort was split between piglets increasing (53%) and
decreasing (47%) in BWPD (control mean: 0.85; ColiGuard® mean: -5.49; p=0.040041)
(**S26 A-C**).

Effects of antibiotic and probiotic treatment on beta diversity

To investigate the treatment effect on beta diversity, principal component analysis
(PCA) of the Kantorovich-Rubenstein distances (beta diversity analysis) was performed on
all samples and, additionally, on samples within individual time points. This analysis is
conceptually similar to the weighted Unifrac approach for beta diversity analysis, but is
designed to work with phylogenetic placement data [92]. When examining all samples together,
there was no clear separation of cohorts on any of the five principal component axes. When
individual time points were analysed, some clustering by cohort was observed (**S27**). D-
Scour™ separated from the control cohort in PC2 (17.95%) and in PC3 (8.06%) one week
and two weeks after probiotic treatment, respectively (**S27**). ColiGuard® separated from
control two weeks post probiotic treatment in PC3 (7.88%). Two weeks after the end of
neomycin treatment, neomycin separated from control in PC4 (5.32%), with a smaller
clustering and a higher representation of *Mollicutes*. Two and four weeks after probiotic
treatment, neomycin+D-Scour™ separated from neomycin in PC2 (17.74%) and PC5
(3.81%), respectively. In both timepoints and principal components, neomycin+D-Scour™
showed a higher representation of *Lactobacillales*. (**S27**)

Association between weight and community composition

Weight correlated with the abundance of certain species (n=83) at each time point as it
resulted from principal component analysis of 16S rRNA reads. Positively correlating with
weight we found among others: *Blautia* (t0), *Cetobacterium* (t0), *Ruminibacter* (t0),
Rickettsia (t2), *Lactobacillus* (t6), *Mycoplasma* (t6), *Anaerostipes* (t8), *Ruminococcus* (t8-

t10), *Cerasibacillus* (t10). Negatively correlating with weight we found among others:
Pyramidobacter (t0), *Odoribacter* (t2), *Schwartzia* (t6), *Streptococcus* (t6), *Dokdonella* (t8).
(**S28-S33**) Significance values and correlation estimates are given in **Supplementary file 5**.

Effect of treatments on weight gain

Overall weight gain from initial to final weight (**S34 A-E**; **Supplementary file 5**) was
not significantly affected by any treatment. However, the probiotic ColiGuard® was found to
have a partial effect on piglet weight gain (**S34**; **Supplementary file 5**). Weight was
measured weekly for a total of six measurements. Based on Tukey adjusted *p* values, a lower
weight gain was detected in the ColiGuard® cohort in week 3 compared to the control cohort
(*p*=0.0008) (**S34 C-D**; **Supplementary file 5**). Similarly, a lower weight gain was detected
in the neomycin+ColiGuard® cohort in C-E compared to the neomycin cohort (*p*=0.0393)
(**S34 C-E**; **Supplementary file 5**). Breed and age differences among piglets were not
associated with weight gain (**Supplementary file 5**).

DISCUSSION

The consistent trend in community composition over time, across all the cohorts,
indicates that an age-related process of ecological succession is the largest factor shaping the
microbial community of post-weaning piglets, as found in this study where animals aged 20-
63 days and were fed the same diet. A peak in unrooted phylogenetic diversity and drop in
balance weighted phylogenetic diversity (BWPD) reflects the acquisition of new species
with the loss of dominating species. This change, detected in the week following the piglets'
arrival at the trial site irrespective of the cohort, could be linked to the piglets being
subjected to microbial interchange (e.g.: new pen mates [115,116]) and/or to diet transition (peri-
weaning transition to solid food [115,116]) leading to the reshaping of the gut microbial
community. The week following the drop of BWPD, a significant increase of BWPD was
recorded, reflecting the acquisition of a larger proportion of the community by the newly
introduced species. The strong changes in phylogenetic diversity detected in the first and the
second week could as well be attributable to other post-weaning related physiological
changes, as previous studies report [44,115–117].

The highest inter-individual differences among piglets are seen in the first week of life,
irrespective of maternal or environmental effects. The microbiota of 3 week old piglets is

still very dynamic, but environmental factors become evident [116]. At 6 weeks of age, CD8+ T
cells infiltrate the intestinal tissue and the mucosa and intestinal lining resemble that of an
adult pig [115] . In this study, piglets reached a comparable alpha diversity to the sows after the
first week of the trial, at which time the piglets were aged between 3.8 and 4.6 weeks.
Unrooted PD did not reach higher levels at later sampling time points. The highest BWPD
accompanied by a high unrooted PD was reached after the second week of the trial when
piglets were aged between 4.9 and 5.6 weeks. Age-dependent physiological changes could
explain i) the major shifts we detected in alpha diversity during the first two weeks of the
trial and, ii) the distinct differences in community composition with age, even with a narrow
age difference between piglets (1-6 days). We were able to appreciate a significant trend of
increasing unrooted PD and decreasing BWPD with age in piglets that separated a 6 days
maximum by day of birth from each other. Since age groups were confounded with breeds in
our study, we attempted to determine the correlation within single breeds. Unfortunately,
although the correlations with age could still be detected, we could not determine the
association at later time points due to the introduction of treatment effects.

Animal trials are often conducted in controlled environments so as to minimize
environmental effects. However, individual variations such as breed and age are often
unavoidable in large animal trials. Previously reported confounding factors include:
individual variation [44], cohabitation [115,116], age [44,115,116], maternal effects [115,116,118], hormones [44],
behavioural differences between breeds (e.g. coprophagy, mouth to mouth contact) and
extent of long-term behavioural adaptation, which can differ between breeds for reasons not
attributable to genetics [44,115,116]. A litter effect was found in piglets at the start of the trial and
was lost at later time points during the trial. This could be due to either of the
aforementioned factors. Co-housing, aging and the splitting of the piglets in separate rooms
to receive a different treatment, are possible causes for loss of the litter effect. In this study
we confirm the importance of these factors in the contribution to inter-individual variability
of gut microbial composition. Motta *et al* (2019) report a correlation of beta diversity with
age and no correlation of genotype and litter effect with either alpha or beta diversity [9] . On
the contrary, we found the piglet samples to significantly cluster by litter, breed and by age
up to the second and the fourth week post weaning, in alpha diversity and beta diversity,
respectively. We conclude that even small age differences among post- weaning piglets,
down to the day, must be accounted for in an experimental set up.

Three groups of piglets (cohorts neomycin, neomycin+D-Scour™ and

neomycin+ColiGuard®) underwent 5 days of treatment with the broad-spectrum antibiotic

neomycin, via intramuscular administration. Intramuscular neomycin poorly diffuses (<10%)

into a healthy gastrointestinal tract [119] , therefore a direct effect of neomycin on the gut

microbiome may not be expected. However, neomycin showed a different trend in unrooted

PD between the second and the third week of the trial, corresponding to the week following

the neomycin treatment period for the neomycin cohort. Taking this time frame into

consideration, the neomycin cohort did not increase in BWPD to the extent of the Control

cohort. Although statistically significant differences between neomycin and Control in alpha

diversity were not reached, the BWPD of the neomycin cohort appears to follow a different

trend to the Control from the first week (during neomycin treatment) where neomycin treated

piglets show the lowest decrease of BWPD compared to the control cohort and all other

cohorts. While all cohorts show an increase in absolute phylogenetic diversity accompanied

by a decrease of diversity evenness during this time frame, the neomycin cohort piglets show

a lower drop in BWPD, suggesting an increase of species richness, without a corresponding

loss of species evenness. Furthermore the neomycin cohort significantly separated from the

control cohort in beta diversity in the third week of the trial, showing a higher representation

of *Mollicutes*. Numerous studies report the link of oral antibiotic use with dysbiosis [22,35–

38,44,45], as well as with host physiology changes [37]. On the contrary, the effect of intramuscular

antibiotic administration on the microbiome is less well investigated. Correlation between

intramuscular antibiotic use and dysbiosis has been reported in fish [46], gorillas [47], humans [120],

and pigs [48,121]. In 1 day old piglets, a single IM injection of amoxicillin (penicillin class) is

reported to have an effect on the intestinal microbiota, detectable 40 days post treatment [48].

Zeineldin *et al* (2018) tested the effects of IM administration of several antibiotics of various

classes (penicillin, macrolide, cephalosporin and tetracycline), in 8-week old piglets,

reporting shifts of the *Firmicutes*/*Bacteroidetes* ratio following treatment (length of the

treatment not reported)[121]. The effects of intramuscular administration of neomycin

(aminoglycoside class) on the gut microbiota have to our knowledge not been investigated.

Based on our results we conclude that a mild effect on phylogenetic diversity is appreciable

post IM neomycin treatment, up to two weeks after termination of the treatment. Additional

compositional and functional analysis is necessary to determine the source of this mild

variation. Differences were not detected at later time points, based on our phylogenetic

diversity analyses, suggesting a full recovery of the microbial communities after two week
from the end of the treatment.

It is possible that the large shifts in phylogenetic diversity taking place in the first two
weeks irrespective of the treatment (an increase, then decrease of unrooted PD, and an
opposite trend of BWPD) have masked the milder effects of the treatment, despite our efforts
to control for the effects of aging. This could be the reason why a significantly distinct alpha
diversity trend was found in the neomycin+D-Scour™ cohort compared to the neomycin
cohort, but not in the D-Scour™ cohort compared to the Control cohort. The neomycin+D-
Scour™ cohort underwent 5 days of neomycin treatment followed by 2 weeks of D-Scour™
treatment. A significant increase of BWPD was detected in the two-week period of D-
Scour™ treatment, indicating a possible enhancement of microbiome evenness following
neomycin treatment. To our knowledge there are no studies reporting an increased evenness
in piglet gut community composition following a specific probiotic treatment. There are
instead multiple studies reporting beneficial effects of probiotic treatment in sucker and
weaner piglets in terms of improved gut mucosal integrity [66,80], growth rate [80–83], digestibility
of proteins and water absorption [80,83], reduction of pathogen invasion efficiency [76,79,80], and
decreased mortality [80,82]. Although the assessment of physiologic changes from probiotic
treatments was outside the scope of this study, we found significant separation of
neomycin+D-Scour™ cohort samples to neomycin cohort samples in beta diversity 3 and 10
days after D-Scour™ treatment, where neomycin+D-Scour™ samples showed a higher
representation of *Lactobacillales* compared to neomycin samples, suggesting a transient
establishment of the probiotic strains in the piglet guts.

The second probiotic in this study, ColiGuard®, did not have an effect on alpha
diversity, but clustering was detected in beta diversity, where ColiGuard® samples separated
from Control cohort samples in the third principal component (explaining 7.88% of the
variation) two weeks post probiotic treatment. Additionally, the ColiGuard® treatment
correlated with a lower weight gain, whether or not it was preceded by the antibiotic
treatment. However, when comparing the overall weight gain (from the start to the end of the
trial) the weight gain in the cohorts receiving ColiGuard® did not differ from the other
cohorts.

We extracted the 16S rRNA gene hypervariable regions from our dataset, obtained the
counts, and ran a correlation analysis to discover taxa that correlated with the weight of the

piglets. As a consequence of the library size normalization step, the use of correlation with
compositional data can inflate the false discovery rate [122,123]. For this reason it can be
expected that some of the taxa we found to correlate with the weight of the piglets (eighty-
three distinct species) could be spurious while other correlations may have been missed.

Technical controls in metagenomic studies and methodological limitations

Taxonomic assignment of the raw reads from the positive controls was performed with
MetaPhlAn2[112] which relies on a ca. 1M unique clade-specific markers derived from 17,000
reference genomes. Such a database to map against the positive controls suffices as these
organisms are cultivable, and for this reason they are widely studied hence the sequences are
known. This is not the case for real-world samples where mapping against a database (which
completeness relies on studied and often cultivable organisms) would narrow the view on the
true diversity within the sample.

Positive controls with well-studied members and known ratios within the samples, has
proven to be a valuable tool to assess consistency among technical replicates across batches
and to detect possible biases derived from the DNA extraction method.

Systematic taxonomic bias in microbiome studies, resulting from differences in cell wall
structures between Gram positive and Gram negative bacteria, have previously been
reported; sample treatment with enzymatic cocktails can modestly reduce this bias [124–126].
Although we implemented this step in our workflow, it seems that, from the read abundance
of our mock community, which contained three Gram negative and four Gram positive
strains, a bias towards Gram negative taxa may still be present.

In terms of contamination we concluded that: a) contamination in our study was not
batch specific; b) a problem of sample cross-contamination existed at the DNA extraction
step between neighbouring wells. During the bead beating step of DNA extraction, the deep-
well plate is sealed with a sealing mat, rotated and placed in a plate shaker for the bead
beating to take place. We consider that sample cross-contamination is most likely to occur
during this step.

Conclusion

Our study provides a publicly-available databank of the pig gut metagenome. Our findings
further stress the importance of confounding factors such as breed, age and maternal effects
when assessing the effect of treatment on the gut microbiome. We found that age, even
within a narrow age span (1-6 days) can have an impact on microbial shifts and should be
accounted for in microbiome studies. Intramuscular neomycin treatment correlated with a
clustering in alpha diversity and a higher representation of *Mollicutes* compared to control.
D-Scour™ treated piglets displayed a mild shift in alpha diversity compared to control, and a
transient establishment of *Lactobacillales*. ColiGuard® treated piglets displayed a clustering
in beta diversity and a transient lower weight gain compared to control. Weight correlated
with the abundance of a number of taxa. Age was the strongest factor shaping phylogenetic
diversity of the piglets.

As previously mentioned, phylogenetic diversity is based on distinct taxa (richness) and
their collective structure (proportions reflected by BWPD) and not on a direct assessment of
composition and function. These types of analyses will be necessary to further describe the
effects of the treatments.

Declarations

Competing Interests

D-Scour™ was sourced from International Animal Health Products (IAHP). ColiGuard®
was developed in a research project with NSW DPI, IAHP and AusIndustry Commonwealth
government funding.

Authors Contributions

Pig Trial: TC, LF, DG, TZ, GJE, AED, SPD

DNA extraction: DG, ML

Library prep, robot pooling: DG, ML, KA, AED

Sequencing data processing: MZD, DG, AED

Data analysis: DG, AED

Manuscript: DG

Manuscript editing: DG, AED, GJE, John Webster

Funding

This work was supported by the Australian Research Council, linkage grant LP150100912.
This project was funded by the Australian Centre for Genomic Epidemiological
Microbiology (Ausgem), a collaborative partnership between the NSW Department of
Primary Industries and the University of Technology Sydney. TZ and DG are recipients of
UTS International Research and UTS President's Scholarships. NSW DPI approved the
paper before submission for publication.

Acknowledgements

We would like to thank the on site team of the pig trial: Graeme Eamens, Linda Falconer,
Shayne Fell, Tiziana Zingali, Toni Chapman. Thank you Amy Bottomley, Giulia Ballerin,
Rosy Cavaliere, for providing the mock community strains. Thank you Akane Tanaka, Leigh
Monahan and Joyce To for the technical support. Thank you John Webster for providing DPI
internal review of this manuscript and for the helpful suggestion about its content.
International Animal Health Products for providing access to the probiotic supplements and
the Australian Centre for Genomic Epidemiological Microbiology (Ausgem) for financially
supporting this study.

Supplementary Files

Supplementary File 1. (animal details)

Supplementary File 2. (metadata)

Supplementary File 3. (barcodes)

Supplementary File 4. (multiqc report)

Supplementary file 5. (stats output)

REFERENCES

1.   Jensen, P. & Recén, B. When to wean — Observations from free-ranging domestic

708   pigs. *Appl. Anim. Behav. Sci.* **23**, 49–60 (1989).

709   2.   Van Heugten, E., Coffey, M. T. & Spears, J. W. Effects of immune challenge, dietary

710   energy density, and source of energy on performance and immunity in weanling pigs. *J.*

711   *Anim. Sci.* **74**, 2431–2440 (1996).

712   3.   Blecha, F. Immune system response to stress. *Biol. Anim. Stress Basic Princ. Implic.*

713   *Anim. Welf. Moberg GP Mench JA Eds* 111–123 (2000).

714   4.   Fairbrother, J. M., Nadeau, É. & Gyles, C. L. Escherichia coli in postweaning diarrhea

715   in pigs: an update on bacterial types, pathogenesis, and prevention strategies. *Anim.*

716   *Health Res. Rev.* **6**, 17–39 (2005).

717   5.   Faubert, C. & Drolet, R. Hemorrhagic gastroenteritis caused by Escherichia coli in

718   piglets: clinical, pathological and microbiological findings. *Can. Vet. J.* **33**, 251 (1992).

719   6.   Nabuurs, M. J. A. Weaning piglets as a model for studying pathophysiology of

720   diarrhea. *Vet. Q.* **20**, 42–45 (1998).

721   7.   Lallès, J.-P., Bosi, P., Smidt, H. & Stokes, C. R. Weaning — A challenge to gut

722   physiologists. *Livest. Sci.* **108**, 82–93 (2007).

723   8.   Heo, J. M. *et al.* Gastrointestinal health and function in weaned pigs: a review of

724   feeding strategies to control post-weaning diarrhoea without using in-feed antimicrobial

725   compounds. *J. Anim. Physiol. Anim. Nutr.* **97**, 207–237 (2013).

726   9.   Motta, V., Luise, D., Bosi, P. & Trevisi, P. Faecal microbiota shift during weaning

727   transition in piglets and evaluation of AO blood types as shaping factor for the bacterial

728   community profile. *PloS One* **14**, e0217001 (2019).

10. Alain B. Pajarillo, E., Chae, J.-P., P. Balolong, M., Bum Kim, H. & Kang, D.-K.

Assessment of fecal bacterial diversity among healthy piglets during the weaning

transition. *J. Gen. Appl. Microbiol.* **60**, 140–146 (2014).

11. Kaper, J. B., Nataro, J. P. & Mobley, H. L. T. Pathogenic Escherichia coli. *Nat. Rev.*

Microbiol. **2**, 123–140 (2004).

12. Jordan, D. *et al.* Antimicrobial use in the Australian pig industry: results of a national

survey. *Aust. Vet. J.* **87**, 222–229 (2009).

13. Zhu, Y.-G. *et al.* Diverse and abundant antibiotic resistance genes in Chinese swine

farms. *Proc. Natl. Acad. Sci.* **110**, 3435–3440 (2013).

14. Erik, K. & Knudsen, B. Development of antibiotic resistance and options to replace

antimicrobials in animal diets. *Proc. Nutr. Soc.* **60**, 291–299 (2001).

15. Levy, S. B. & Marshall, B. Antibacterial resistance worldwide: causes, challenges and

responses. *Nat. Med.* **10**, S122 (2004).

16. White, D. G. Antimicrobial resistance in pathogenic Escherichia coli from animals. in

Antimicrobial resistance in bacteria of animal origin 145–166 (American Society of

Microbiology, 2006).

17. Broens, E. M., Graat, E. A. M., Van der Wolf, P. J., Van de Giessen, A. W. & De Jong,

M. C. M. Prevalence and risk factor analysis of livestock associated MRSA-positive pig

herds in The Netherlands. *Prev. Vet. Med.* **102**, 41–49 (2011).

18. Modi, S. R., Lee, H. H., Spina, C. S. & Collins, J. J. Antibiotic treatment expands the

resistance reservoir and ecological network of the phage metagenome. *Nature* **499**,

219–222 (2013).

19. Prescott, J. F. History of antimicrobial usage in agriculture: an overview. in

Antimicrobial resistance in bacteria of animal origin 19–28 (American Society of

Microbiology, 2006).

20.   Marshall, B. M. & Levy, S. B. Food animals and antimicrobials: impacts on human

755   health. *Clin. Microbiol. Rev.* **24**, 718–733 (2011).

756   21.   Callens, B. *et al.* Presence of antimicrobial resistance and antimicrobial use in sows are

757   risk factors for antimicrobial resistance in their offspring. *Microb. Drug Resist.* **21**, 50–

758   58 (2015).

759   22.   Looft, T. P. *The swine intestinal microbiota: localized adaptations and responses to in-

760   feed antibiotics*. (Iowa State University, 2012).

761   23.   Salyers, A. A. & Shoemaker, N. B. Resistance gene transfer in anaerobes: new insights,

762   new problems. *Clin. Infect. Dis.* **23**, S36–S43 (1996).

763   24.   Salyers, A. A. & Amábile-Cuevas, C. F. Why are antibiotic resistance genes so resistant

764   to elimination? *Antimicrob. Agents Chemother.* **41**, 2321–2325 (1997).

765   25.   Dorado-García, A. *et al.* Dose-Response Relationship between Antimicrobial Drugs

766   and Livestock-Associated MRSA in Pig Farming1. *Emerg. Infect. Dis.* **21**, 950–959

767   (2015).

768   26.   Didelot, X. & Maiden, M. C. J. Impact of recombination on bacterial evolution. *Trends

769   Microbiol.* **18**, 315–322 (2010).

770   27.   Canchaya, C., Fournous, G., Chibani-Chennoufi, S., Dillmann, M.-L. & Brüssow, H.

771   Phage as agents of lateral gene transfer. *Curr. Opin. Microbiol.* **6**, 417–424 (2003).

772   28.   Claverys, J.-P., Martin, B. & Polard, P. The genetic transformation machinery:

773   composition, localization, and mechanism. *FEMS Microbiol. Rev.* **33**, 643–656 (2009).

774   29.   Ravi, A., Avershina, E., Ludvigsen, J., L'Abée-Lund, T. & Rudi, K. Integrons in the

775   Intestinal Microbiota as Reservoirs for Transmission of Antibiotic Resistance Genes.

776   *Pathogens* **3**, 238–248 (2014).

30.   Porse, A., Schønning, K., Munck, C. & Sommer, M. O. A. Survival and Evolution of a

Large Multidrug Resistance Plasmid in New Clinical Bacterial Hosts. *Mol. Biol. Evol.*

33, 2860–2873 (2016).

31.   Bednorz, C. *et al.* The broader context of antibiotic resistance: Zinc feed

supplementation of piglets increases the proportion of multi-resistant Escherichia coli in

vivo. *Int. J. Med. Microbiol.* **303**, 396–403 (2013).

32.   Tadesse, D. A. *et al.* Antimicrobial drug resistance in Escherichia coli from humans and

food animals, United States, 1950–2002. *Emerg. Infect. Dis.* **18**, 741 (2012).

33.   U. S. Food & Drug, A. FDA Task Force on Antimicrobial Resistance: Key

Recommendations and Report. (2000).

34.   Centers for Disease Control & Prevention. Get smart: know when antibiotics work.

Centers for Disease Control and Prevention, Atlanta, GA. (2010).

35.   Dethlefsen, L. & Relman, D. A. Incomplete recovery and individualized responses of

the human distal gut microbiota to repeated antibiotic perturbation. *Proc. Natl. Acad.*

Sci. **108**, 4554–4561 (2011).

36.   Cho, I. *et al.* Antibiotics in early life alter the murine colonic microbiome and adiposity.

Nature **488**, 621–626 (2012).

37.   Antunes, L. C. M. *et al.* Effect of Antibiotic Treatment on the Intestinal Metabolome.

Antimicrob. Agents Chemother. **55**, 1494–1503 (2011).

38.   Gonzalez-Perez, G. *et al.* Maternal Antibiotic Treatment Impacts Development of the

Neonatal Intestinal Microbiome and Antiviral Immunity. *J. Immunol.* **196**, 3768–3779

(2016).

39.   Sachs, J. L., Mueller, U. G., Wilcox, T. P. & Bull, J. J. The evolution of cooperation. *Q.*

Rev. Biol. **79**, 135–160 (2004).

40.   Foster, K. R. & Wenseleers, T. A general model for the evolution of mutualisms. *J.*
Evol. Biol. **19**, 1283–1293 (2006).

41.   Birck, M. M. *et al.* Enteral but not parenteral antibiotics enhance gut function and
prevent necrotizing enterocolitis in formula-fed newborn preterm pigs. *Am. J. Physiol.-*
Gastrointest. Liver Physiol. **310**, G323–G333 (2016).

42.   Wiuff, C., Lykkesfeldt, J., Svendsen, O. & Aarestrup, F. M. The effects of oral and
intramuscular administration and dose escalation of enrofloxacin on the selection of
quinolone resistance among Salmonella and coliforms in pigs. *Res. Vet. Sci.* **75**, 185–
193 (2003).

43.   Chantziaras, I., Smet, A., Haesebrouck, F., Boyen, F. & Dewulf, J. Studying the effect
of administration route and treatment dose on the selection of enrofloxacin resistance in
commensal Escherichia coli in broilers. *J. Antimicrob. Chemother.* **72**, 1991–2001
(2017).

44.   Xiao, L. *et al.* A reference gene catalogue of the pig gut microbiome. *Nat. Microbiol.* **1**,
16161 (2016).

45.   Deshmukh, H. S. *et al.* The microbiota regulates neutrophil homeostasis and host
resistance to Escherichia coli K1 sepsis in neonatal mice. *Nat. Med.* **20**, 524–530
(2014).

46.   Kim, A. *et al.* Administration of antibiotics can cause dysbiosis in fish gut. *Aquaculture*
512, 734330 (2019).

47.   Vlčková, K. *et al.* Effect of Antibiotic Treatment on the Gastrointestinal Microbiome of
Free-Ranging Western Lowland Gorillas (Gorilla g. gorilla). *Microb. Ecol.* **72**, 943–954
(2016).

48.   Janczyk, P. *et al*. Parenteral long-acting amoxicillin reduces intestinal bacterial

825   community diversity in piglets even 5 weeks after the administration. *ISME J*. **1**, 180–

826   183 (2007).

827   49.   Allen, H. K., Levine, U. Y., Looft, T., Bandrick, M. & Casey, T. A. Treatment,

828   promotion, commotion: antibiotic alternatives in food-producing animals. *Trends*

829   *Microbiol*. **21**, 114–119 (2013).

830   50.   Rangan, K. J. *et al*. A secreted bacterial peptidoglycan hydrolase enhances tolerance to

831   enteric pathogens. *science* **353**, 1434–1437 (2016).

832   51.   Tap, J. *et al*. Gut microbiota richness promotes its stability upon increased dietary fibre

833   intake in healthy adults. *Environ. Microbiol*. **17**, 4954–4964 (2015).

834   52.   Kaleko, M. *et al*. Development of SYN-004, an oral beta-lactamase treatment to protect

835   the gut microbiome from antibiotic-mediated damage and prevent Clostridium difficile

836   infection. *Anaerobe* (2016) doi:10.1016/j.anaerobe.2016.05.015.

837   53.   Song, J., Li, Y. li & Hu, C. hong. Effects of copper-exchanged montmorillonite, as

838   alternative to antibiotic, on diarrhea, intestinal permeability and proinflammatory

839   cytokine of weanling pigs. *Appl. Clay Sci*. **77–78**, 52–55 (2013).

840   54.   Holler, E. *et al*. Metagenomic Analysis of the Stool Microbiome in Patients Receiving

841   Allogeneic Stem Cell Transplantation: Loss of Diversity Is Associated with Use of

842   Systemic Antibiotics and More Pronounced in Gastrointestinal Graft-versus-Host

843   Disease. *Biol. Blood Marrow Transplant*. **20**, 640–645 (2014).

844   55.   Nogueira, M., Collins, A., Dunlop, R. & Emery, D. Effect of the route of administration

845   on the mucosal and systemic immune responses to *Lawsonia intracellularis* vaccine in

846   pigs. *Aust. Vet. J*. **93**, 124–126 (2015).

56. Bak, H. & Rathkjen, P. H. Reduced use of antimicrobials after vaccination of pigs

against porcine proliferative enteropathy in a Danish SPF herd. *Acta Vet. Scand.* **51**, 1

(2009).

57. Francis, D. H. & Willgohs, J. A. Evaluation of a live avirulent Escherichia coli vaccine

for K88+, LT+ enterotoxigenic colibacillosis in weaned pigs. *Am. J. Vet. Res.* **52**,

1051–1055 (1991).

58. Morgan, R. L., Isaacson, R. E., Moon, H. W., Brinton, C. C. & To, C. C. Immunization

of suckling pigs against enterotoxigenic Escherichia coli-induced diarrheal disease by

vaccinating dams with purified 987 or K99 pili: protection correlates with pilus

homology of vaccine and challenge. *Infect. Immun.* **22**, 771–777 (1978).

59. Callaway, T. R. *et al.* Evaluation of Phage Treatment as a Strategy to Reduce

Salmonella Populations in Growing Swine. *Foodborne Pathog. Dis.* **8**, 261–266 (2011).

60. Willamil, J., Creus, E., Francisco Pérez, J., Mateu, E. & Martín-Orúe, S. M. Effect of a

microencapsulated feed additive of lactic and formic acid on the prevalence of

Salmonella in pigs arriving at the abattoir. *Arch. Anim. Nutr.* **65**, 431–444 (2011).

61. Brouns, F., Kettlitz, B. & Arrigoni, E. Resistant starch and "the butyrate revolution".

Trends Food Sci. Technol. **13**, 251–261 (2002).

62. Missotten, J. A. M., Michiels, J., Ovyn, A., De Smet, S. & Dierick, N. A. Fermented

liquid feed for pigs. *Arch. Anim. Nutr.* **64**, 437–466 (2010).

63. Canibe, N. & Jensen, B. B. Fermented and nonfermented liquid feed to growing pigs:

effect on aspects of gastrointestinal ecology and growth performance. *J. Anim. Sci.* **81**,

2019–2031 (2003).

64. Moran, C. A. Development and benefits of liquid diets for newly weaned pigs. *PhD

Univ. Plymouth* (2001).

65. Lessard, M. *et al.* Administration of Pediococcus acidilactici or Saccharomyces

cerevisiae boulardii modulates development of porcine mucosal immunity and reduces

intestinal bacterial translocation after Escherichia coli challenge1, 2. *J. Anim. Sci.* **87**,

922 (2009).

66. Deng, J., Li, Y., Zhang, J. & Yang, Q. Co-administration of Bacillus subtilis RJGP16

and Lactobacillus salivarius B1 strongly enhances the intestinal mucosal immunity of

piglets. *Res. Vet. Sci.* **94**, 62–68 (2013).

67. Garrido-Mesa, N. *et al.* The association of minocycline and the probiotic Escherichia

coli Nissle 1917 results in an additive beneficial effect in a DSS model of reactivated

colitis in mice. *Biochem. Pharmacol.* **82**, 1891–1900 (2011).

68. Ukena, S. N. *et al.* Probiotic Escherichia coli Nissle 1917 Inhibits Leaky Gut by

Enhancing Mucosal Integrity. *PloS One* **2**, e1308 (2007).

69. Deriu, E. *et al.* Probiotic bacteria reduce salmonella typhimurium intestinal colonization

by competing for iron. *Cell Host Microbe* **14**, 26–37 (2013).

70. Hancock, V., Dahl, M. & Klemm, P. Probiotic Escherichia coli strain Nissle 1917

outcompetes intestinal pathogens during biofilm formation. *J. Med. Microbiol.* **59**, 392–

399 (2010).

71. Heyman, M. Effect of lactic acid bacteria on diarrheal diseases. *J. Am. Coll. Nutr.* **19**,

137S-146S (2000).

72. Jin, L.-Z., Marquardt, R. R. & Baidoo, S. K. Inhibition of enterotoxigenic Escherichia

coli K88, K99 and 987P by the Lactobacillus isolates from porcine intestine. *J Sci Food*

Agric 6 (2000).

73. Maltby, R., Leatham-Jensen, M. P., Gibson, T., Cohen, P. S. & Conway, T. Nutritional

basis for colonization resistance by human commensal Escherichia coli strains HS and

Nissle 1917 against E. coli O157: H7 in the mouse intestine. *PloS One* **8**, e53957

(2013).

74. Suda, Y. *et al*. Immunobiotic Lactobacillus jensenii as immune-health promoting factor

to improve growth performance and productivity in post-weaning pigs. *BMC Immunol*.

15, 24 (2014).

75. Shimazu, T. *et al*. Immunobiotic Lactobacillus jensenii Elicits Anti-Inflammatory

Activity in Porcine Intestinal Epithelial Cells by Modulating Negative Regulators of the

Toll-Like Receptor Signaling Pathway. *Infect. Immun*. **80**, 276–288 (2012).

76. Splichalova, A. *et al*. Interference of Bifidobacterium choerinum or Escherichia coli

Nissle 1917 with Salmonella Typhimurium in gnotobiotic piglets correlates with

cytokine patterns in blood and intestine. *Clin. Exp. Immunol*. **163**, 242–249 (2011).

77. Roselli, M. *et al*. The novel porcine Lactobacillus sobrius strain protects intestinal cells

from enterotoxigenic Escherichia coli K88 infection and prevents membrane barrier

damage. *J. Nutr*. **137**, 2709–2716 (2007).

78. Altenhoefer, A. *et al*. The probiotic Escherichia coli strain Nissle 1917 interferes with

invasion of human intestinal epithelial cells by different enteroinvasive bacterial

pathogens. *FEMS Immunol. Med. Microbiol*. **40**, 223–229 (2004).

79. Šmajs, D. *et al*. Experimental Administration of the Probiotic Escherichia coli Strain

Nissle 1917 Results in Decreased Diversity of E. coli Strains in Pigs. *Curr. Microbiol*.

64, 205–210 (2012).

80. Valeriano, V. D. V., Balolong, M. P. & Kang, D. K. Probiotic roles of Lactobacillus sp.

in swine: insights from gut microbiota. *J. Appl. Microbiol*. **122**, 554–567 (2017).

81. Abe, F., Ishibashi, N. & Shimamura, S. Effect of administration of bifidobacteria and

lactic acid bacteria to newborn calves and piglets. *J. Dairy Sci*. **78**, 2838–2846 (1995).

82.   Davis, M. E. *et al.* Effect of a Bacillus-based direct-fed microbial feed supplement on

growth performance and pen cleaning characteristics of growing-finishing pigs. *J.*

Anim. Sci. **86**, 1459–1467 (2008).

83.   Chu, G. M., Lee, S. J., Jeong, H. S. & Lee, S. S. Efficacy of probiotics from anaerobic

microflora with prebiotics on growth performance and noxious gas emission in growing

pigs: SYNBIOTICS ON NOXIOUS GAS EMISSION IN PIGS. *Anim. Sci. J.* **82**, 282–

290 (2011).

84.   Kim, P. I. *et al.* Probiotic properties of Lactobacillus and Bifidobacterium strains

isolated from porcine gastrointestinal tract. *Appl. Microbiol. Biotechnol.* **74**, 1103–1111

(2007).

85.   Takahashi, S., Egawa, Y., Simojo, N., Tsukahara, T. & Ushida, K. Oral administration

of Lactobacillus plantarum strain Lq80 to weaning piglets stimulates the growth of

indigenous lactobacilli to modify the lactobacillal population. *J. Gen. Appl. Microbiol.*

53, 325–332 (2007).

86.   Gaio, D. *et al.* Hackflex: low cost Illumina sequencing library construction for high

sample counts. *bioRxiv* 779215 (2019).

87.   Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize analysis

results for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048

(2016).

88.   Ewels, P. A. *et al.* nf-core *: Community curated bioinformatics pipelines.* (2019).

89.   Darling, A. E. *et al.* PhyloSift: phylogenetic analysis of genomes and metagenomes.

PeerJ **2**, e243 (2014).

90.   Faith, D. P. Conservation evaluation and phylogenetic diversity. *Biol. Conserv.* **61**, 1–

10 (1992).

91. McCoy, C. O. & Matsen, F. A. Abundance-weighted phylogenetic diversity measures

distinguish microbial community states and are robust to sampling depth. *PeerJ* **1**, e157

(2013).

92. Evans, S. N. & Matsen, F. A. The phylogenetic Kantorovich-Rubinstein metric for

environmental sequence samples. *J. R. Stat. Soc. Ser. B Stat. Methodol.* **74**, 569–592

(2012).

93. R Core Team, R. C. T. R: A language and environment for statistical computing.

(2013).

94. Wickham, H., Hester, J. & Francois, R. *readr: Read Rectangular Text Data*. (2018).

95. Wickham, H. & Bryan, J. *readxl: Read Excel Files*. (2019).

96. Wickham, H., François, R., Henry, L. & Müller, K. *dplyr: A Grammar of Data*

Manipulation. (2019).

97. Wickham, H. *et al*. Welcome to the tidyverse. *J. Open Source Softw.* **4**, 1686 (2019).

98. Robinson, D. & Hayes, A. *broom: Convert Statistical Analysis Objects into Tidy*

Tibbles. (2019).

99. Vu, V. Q. *ggbiplot: A ggplot2 based biplot*. (2011).

100. Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. & R. Core Team. *nlme: Linear and*

Nonlinear Mixed Effects Models. (2019).

101. Wilke, C. O. *cowplot: Streamlined Plot Theme and Plot Annotations for 'ggplot2'*.

(2019).

102. Kassambara, A. *ggpubr: 'ggplot2' Based Publication Ready Plots*. (2019).

103. Schauberger, P. & Walker, A. *openxlsx: Read, Write and Edit xlsx Files*. (2019).

104. Mahto, A. *splitstackshape: Stack and Reshape Datasets After Splitting Concatenated*

Values. (2019).

105. Kassambara, A. *rstatix: Pipe-Friendly Framework for Basic Statistical Tests*. (2019).

106. Henry, L. & Wickham, H. *purrr: Functional Programming Tools*. (2019).

107. Dinno, A. *dunn.test: Dunn's Test of Multiple Comparisons Using Rank Sums*. (2017).

108. Auguie, B. *gridExtra: Miscellaneous Functions for 'Grid' Graphics*. (2017).

109. Bache, S. M. & Wickham, H. *magrittr: A Forward-Pipe Operator for R*. (2014).

110. Bates, D. & Maechler, M. *Matrix: Sparse and Dense Matrix Classes and Methods*.

(2019).

111. Kopylova, E., Noé, L. & Touzet, H. SortMeRNA: fast and accurate filtering of

ribosomal RNAs in metatranscriptomic data. *Bioinformatics* **28**, 3211–3217 (2012).

112. Truong, D. T. *et al*. MetaPhlAn2 for enhanced metagenomic taxonomic profiling. *Nat*.

Methods **12**, 902–903 (2015).

113. Monahan, L. G. *et al*. High contiguity genome sequence of a multidrug-resistant

hospital isolate of Enterobacter hormaechei. *Gut Pathog*. **11**, 3 (2019).

114. Kembel, S. W. *et al*. Architectural design influences the diversity and structure of the

built environment microbiome. *ISME J*. **6**, 1469–1479 (2012).

115. Pajarillo, E. A. B., Chae, J. P., Kim, H. B., Kim, I. H. & Kang, D.-K. Barcoded

pyrosequencing-based metagenomic analysis of the faecal microbiome of three

purebred pig lines after cohabitation. *Appl*. *Microbiol*. *Biotechnol*. **99**, 5647–5656

(2015).

116. Thompson, C. L., Wang, B. & Holmes, A. J. The immediate environment during

postnatal development has long-term impact on gut community structure in pigs. *ISME*

J. **2**, 739 (2008).

117. Bailey, M. *et al*. Regulation of mucosal immune responses in effector sites. *Proc*. *Nutr*.

Soc. **60**, 427–435 (2001).

118. Funkhouser, L. J. & Bordenstein, S. R. Mom knows best: the universality of maternal

microbial transmission. *PLoS Biol* **11**, e1001631 (2013).

119. Ludlow, K. Australian Pesticides and Veterinary Medicines Authority. in *Encyclopedia*

of Nanoscience and Society (SAGE Publications, Inc., 2010).

120. Fouhy, F. *et al*. High-Throughput Sequencing Reveals the Incomplete, Short-Term

Recovery of Infant Gut Microbiota following Parenteral Antibiotic Treatment with

Ampicillin and Gentamicin. *Antimicrob. Agents Chemother*. **56**, 5811–5820 (2012).

121. Zeineldin, M., Aldridge, B., Blair, B., Kancer, K. & Lowe, J. Impact of parenteral

antimicrobial administration on the structure and diversity of the fecal microbiota of

growing pigs. *Microb. Pathog*. **118**, 220–229 (2018).

122. Friedman, J. & Alm, E. J. Inferring Correlation Networks from Genomic Survey Data.

PLoS Comput. Biol. **8**, e1002687 (2012).

123. Lovell, D., Pawlowsky-Glahn, V., Egozcue, J. J., Marguerat, S. & Bähler, J.

Proportionality: A Valid Alternative to Correlation for Relative Data. *PLOS Comput.*

Biol. **11**, e1004075 (2015).

124. von Wintzingerode, F., Göbel, U. B. & Stackebrandt, E. Determination of microbial

diversity in environmental samples: pitfalls of PCR-based rRNA analysis. *FEMS*

Microbiol. Rev. **21**, 213–229 (1997).

125. Han, Z., Sun, J., Lv, A. & Wang, A. Biases from different DNA extraction methods in

intestine microbiome research based on 16S rDNA sequencing: a case in the koi carp,

Cyprinus carpio var. Koi. *MicrobiologyOpen* **8**, e00626 (2019).

126. Moss, E. L., Maghini, D. G. & Bhatt, A. S. Complete, closed bacterial genomes from

microbiomes using nanopore sequencing. *Nat. Biotechnol*. (2020) doi:10.1038/s41587-

020-0422-6.

1015

Figure 1. Timeline.

Timeline of the trial indicating the start and the length of the treatment for each cohort. Piglets arrived on the site of the trial on January 30th and were allowed 2 days of acclimatisation before the start of the treatments (pink: placebo paste; yellow: probiotic D-Scour™ formulation; green: probiotic ColiGuard® formulation; blue: antibiotic neomycin intramuscular injection). Vertical lines indicate main days of sampling where all piglets where sampled ($n=126$). Small arrows indicate days of sampling for a subset of the piglets (8 per cohort; $n=48$).

Figure 2. Significance of correlations between phylogenetic diversity and specified factors.

Plots of *p* values derived from Kruskal-Wallis analysis of variance (black) and adjusted *p* values by Benjamini-Hochberg method (red). x-axis displays distinct sampling time points in chronological order. Correlation of specified factors of samples are shown with (**A**) alpha diversity and (**B**) beta diversity. Abbreviations: DxL = "Duroc x Landrace" cross breed; DxLW = "Duroc x Large White" cross breed.

Figure 3. Effect of time on beta diversity.

Principal component analysis (PCA) of samples. PCA from edge component analysis with PhyloSift (top) and PCA from 20 most abundant 16S reads extracted with SortMeRNA (bottom). In the top, distribution of samples on either side of the plot (left *versus* right; top *versus* bottom) reflect the taxa that were found to explain the variation. Samples are colour coded by days into the trial. In the lower plot, arrows indicate which of the 20 taxa read contributed to the variation of samples, where arrows thickness represents a higher (thicker) or lower (thinner) contribution. Samples are colour coded by time point during the trial.

Figure 4. Abundance heatmap.

Most abundant taxa within each time point are obtained from analysis with guppy fat. Abundance is derived from the branch width. The distance between each time point is of one week.

A. Allocation of piglets in rooms and pens on the day of arrival

B. Re-distribution of piglets on February 6th

Supplementary Figure 1. Piglets placements across rooms and pens.

Disposition of piglets in rooms (1-4) and pens (a-i) at the start of the trial (**A**) and re-arrangement of a subset of neomycin treated piglets from room 4 to room 2 and room 3, for D-Scour™ and ColiGuard® treatment, respectively (**B**).

Supplementary Figure 2. Sample collection.

Samples were collected from all piglets once weekly, and from a subset group (8 per cohort; *n*=48) twice weekly. Overall, piglets were sampled between 1 and 10 times.

Supplementary Figure 3. Read counts distribution.

Read counts distribution of all samples (main histogram) and low read counts samples (top right histogram).

Supplementary Figure 4. Summarised workflow.

Schematic workflow from sample collection to sequencing (orange) and data analysis (green). R scripts (italic) are available in our Github repository.

Supplementary Figure 5. Principal component analysis of batch effect before batch effect removal.

Principal components 1 to 4 from beta diversity analysis of all samples before batch effect removal. Samples are coloured by DNA extraction plate. No clustering of samples by DNA extraction plate is visible.

Batch effect by alpha and beta diversity

Supplementary Figure 6. Batch effect on alpha and beta diversity before batch effect removal.

Batch effect by alpha (top two plots) and beta diversity (bottom five plots) before batch effect removal. Samples are grouped by DNA extraction plate. The *p* values are derived from multiple comparison analysis with ANOVA, indicating equality of the means. *Post hoc* corrected *p* values for pairwise comparisons are provided in Supplementary File 5.

Supplementary Figure 7. Batch effect on alpha and beta diversity after batch effect removal.

Batch effect by alpha (top two plots) and beta diversity (bottom five plots) after batch effect removal. Samples are grouped by DNA extraction plate. The *p* values are derived from multiple comparison analysis with ANOVA, indicating equality of the means. *Post hoc* corrected *p* values for pairwise comparisons are provided in Supplementary File 5.

Supplementary Figure 8. Taxonomic assignment of reads from positive control samples.

Taxonomic profile of the positive controls used in this study obtained by mapping the reads against a ~1M bacterial genomes database with MetaPhlAn2. Each stacked barplot represents a technical replicate. Taxa of which reads are present in >0.1% are displayed. **A**. In-house made mock community; **B**. commercially available livestock probiotic D-Scour™ ; **C**. ColiGuard®.

Supplementary Figure 9. Expected and observed relative abundance of mock community members.

Expected relative abundance is derived by multiplying CFU count by genome size. Observed relative abundance is derived by multiplying reads mapping with MetaPhlAn2. A higher observed/expected ratio is observed in two of the three Gram negative species and a lower observed/expected ratio is observed in three of the four Gram negative species.

Supplementary Figure 10. Contamination rate in mock community samples.

Heatmap of contaminating species present in the positive control mock community technical replicates. *Lactobacillus salivarius* is found in one replicate at 0.01% of the total reads.

D–Scour contamination

	R1	R2	R3	R4	R5	R6	R7	R8	Species
	0.00	0.00	0.03	0.00	0.00	0.01	0.00	0.03	Methanobrevibacter_smithii
	0.00	0.00	0.01	0.00	0.01	0.01	0.01	0.05	Methanobrevibacter_unclassified
	0.00	0.00	0.00	0.00	0.00	0.00	0.03	0.00	Staphylococcus_aureus
	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	Staphylococcus_epidermidis
	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	Enterococcus_cecorum
	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	Enterococcus_faecalis
	0.00	0.00	0.02	0.00	0.00	0.00	0.01	0.00	Lactobacillus_amylovorus
	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	Lactobacillus_johnsonii
	0.00	0.00	0.02	0.00	0.00	0.00	0.00	0.07	Lactobacillus_reuteri
	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	Lactobacillus_salivarius
	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	Lactobacillus_vaginalis
	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	Peptostreptococcaceae_noname_unclassified
	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	Faecalibacterium_prausnitzii
	0.00	0.00	0.04	0.00	0.00	0.02	0.00	0.00	Subdoligranulum_unclassified
	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	Catenibacterium_mitsuokai
	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	Eubacterium_biforme
	0.00	0.00	0.02	0.00	0.00	0.00	0.00	0.00	Phascolarctobacterium_succinatutens
	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	Megasphaera_unclassified
	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	Mitsuokella_unclassified
	0.00	0.00	0.00	0.00	0.00	0.00	0.10	0.00	Enterobacter_cloacae
	0.00	0.00	0.00	0.00	0.00	0.01	0.01	0.00	Escherichia_coli
	0.00	0.00	0.00	0.01	0.01	0.00	0.05	0.01	Escherichia_unclassified
	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.00	Pseudomonas_aeruginosa
	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	Pseudomonas_unclassified
	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	Porcine_type_C_oncovirus

Supplementary Figure 11. Contamination rate in D-Scour™ samples.

Heatmap of contaminating species present in the positive control D-Scour™ technical replicates.
The D-Scour™ technical replicates contained 25 contaminants, of which 18 and 7 were identified at the species and at genus level, respectively. Contaminants are majorly present in three technical replicates (R3, R7, R8) and the most frequent contaminant (*Metahobrevibacter*) was present in 5 of the 8 replicates.

Supplementary Figure 12. Contamination rate in ColiGuard® samples.

Heatmap of contaminating species present in the positive control ColiGuard® technical replicates. ColiGuard® contained 20 contaminants, of which 16 and 4 were identified at the species and at genus level, respectively. Contaminants were present majorly in two technical replicates (R5, R7).

Supplementary Figure 13. Alpha phylogenetic diversity of cohorts.

Alpha phylogenetic diversity per cohort from samples across all time points. Phylogenetic entropy (top); Balance-weighted phylogenetic diversity (BWPD) (mean±SD: Positive control Mock community: 1.58±0.12; Positive control D-Scour™: 1.89±0.26; Positive control ColiGuard®: 0.82±0.15; Control: 2.12±0.14; D-Scour™: 2.14±0.14; ColiGuard®: 2.11±0.13; neomycin: 2.12±0.17; neomycin+D-Scour™: 2.12±0.14; neomycin+ColiGuard®: 2.10±0.15; sows: 2.13±0.14; all piglet cohorts: 2.12±0.14); (middle); Unrooted phylogenetic diversity (Positive control Mock community: 31.53±29.50; Positive control D-Scour™: 64.84±21.30; Positive control ColiGuard®: 58.52±21.70; Control: 117±16.8; D-Scour™: 121±14.7; ColiGuard®: 118±22.3; neomycin: 119±22.6; neomycin+D-Scour™: 116±23.4; neomycin+ColiGuard®: 121±15.8; sows: 125.39±7.56; all piglet cohorts: 118.54±19.70) (bottom).

Supplementary Figure 14. Alpha phylogenetic diversity by age group.

Alpha diversity of samples from the start of the trial grouped by the date of birth of the piglet. Unrooted phylogenetic diversity (top) and balance weighted phylogenetic diversity (bottom). *P* values are derived from Kruskal-Wallis analysis of variance. Pairwise comparisons and *post hoc* corrected *p* values are provided in Supplementary File 5.

Supplementary Figure 15. Alpha phylogenetic diversity by age breed.

Alpha diversity of samples from the start of the trial grouped by breed and by date of birth. Unrooted phylogenetic diversity (top) and balance weighted phylogenetic diversity (bottom). *P* values are derived from Kruskal-Wallis analysis of variance. Pairwise comparisons and *post hoc* corrected *p* values are provided in Supplementary File 5.

A Piglets alpha diversity (unrooted PD)
Grouped by maternal sow

B Piglets alpha diversity (unrooted)
Grouped by nurse sow

Supplementary Figure 16. Unrooted alpha phylogenetic diversity by maternal and nurse sow.

Alpha phylogenetic diversity (unrooted) of samples from piglets at the start of the trial (one time point -sample per piglet). The grouping is based on the maternal sow (**A**) or nurse sow (**B**). *P* values from multiple groups comparison by maternal sow: Kruskal-Wallis *p* values: 0.0011; BWPD: 0.0049. *P* values from multiple groups comparison by nurse sow: Kruskal-Wallis *p* values: 0.0036; BWPD: 0.0071. *Post-hoc* pairwise comparisons and adjusted *p* values are provided in Supplementary File 5.

Supplementary Figure 17. Balance weighted alpha phylogenetic diversity by maternal and nurse sow.

Alpha phylogenetic diversity (unrooted) of samples from piglets at the start of the trial (one time point -sample per piglet). The grouping is based on the maternal sow (**A**) or nurse sow (**B**). *P* values from multiple groups comparison by maternal sow: Kruskal-Wallis *p*=0.0049. *P* values from multiple groups comparison by nurse sow: Kruskal-Wallis *p*=0.0071. *Post-hoc* pairwise comparisons and adjusted *p* values are provided in Supplementary File 5.

Supplementary Figure 18. Principal component analysis of piglet samples by nurse sow.

First two principal components of beta phylogenetic diversity. Principal component 1 (PC1) and principal component 2 (PC2) explaining 46.99% and 21.68% of the variation, respectively.
Samples from piglets at the start of the trial (one time point -sample per piglet) coloured by nurse sow (n=30). Ellipse is drawn at 0.80 confidence level.

Supplementary Figure 19. Principal component analysis of piglet samples by maternal sow.

First two principal components of beta phylogenetic diversity. Principal component 1 (PC1) and principal component 2 (PC2) explaining 46.99% and 21.68% of the variation, respectively. Samples from piglets at the start of the trial (one time point -sample per piglet) colored by maternal sow (n=32). Ellipse is drawn at 0.80 confidence level.

Supplementary Figure 20. Time effect on beta diversity per cohort.

Edge principal component analysis of all samples coloured by sample collection time across the trial (day 0: red; day 10: yellow; day 20: green; day 30: light blue; day 40: dark blue) where principal component 1 (PC1) explains 46.99% of the variation and principal component 2 (PC2) explains 21.68% of the variation. Each plot displays all samples within each cohort.

Supplementary Figure 21. Time effect on beta diversity.

Each plot represents the samples distribution along one of the five principal components across time. Most variation is explained by PC1 (46.99%), followed by PC2 (21.68%), PC3 (8.37%), PC4 (5.1%) and PC5 (3.04%). Distribution of the samples on either side of a plot (left *versus* right) reflects the taxa that were found to explain the variation. The distributions are colour coded by time point during the trial.

Supplementary Figure 22. Taxa displaying the highest variation in beta diversity across time.

Heatmap of taxa explaining the community composition of samples from separate time points of the trial (1 week interval between time points) derived from edge principal component analysis. Intensity is derived from branch width by the percentage of variability explained by the principal components.

Supplementary Figure 23. Time trend of alpha diversity by cohort.

Unrooted PD (**A**) and BWPD (**B**) describe richness and evenness, respectively, of alpha phylogenetic diversity for all samples across time, grouped and colour coded by cohort. The *p* values derived from pairwise comparisons of time points within each of the cohorts. *P* values and *post-hoc* corrected *p* values of time points comparisons for each separate cohort are provided in Supplementary File 5.

Supplementary Figure 24 Alpha diversity by cohort across four time points.

Alpha phylogenetic diversity for each cohort for a subset of 4 time points. *P* values are derived from pairwise comparisons of the means of each time point of (**A**) unrooted phylogenetic diversity and (**B**) balance weighted phylogenetic diversity. Bonferroni correction of *p* values is applied. More time points were tested and significance was determined; reported in Supplementary File 5.

Supplementary Figure 25. Deltas of unrooted alpha phylogenetic diversity between time points.

Change of alpha unrooted phylogenetic diversity in samples between time points across the trial. On the y-axis of plots the percentage difference in unrooted PD between time points per cohort. Letters on the top left of each plot indicate the time points compared with one week interval (A-B, B-C, C-D, D-E) and two weeks interval (A-C, B-D). Pairwise t-test comparisons between cohorts were computed. Brackets in the timeline (top) indicate a significant *p* value (p<0.05) between cohorts (1-week interval); bold brackets indicate a significant *p* value (p<0.05) between cohorts (2-weeks interval). * indicate significance reached when applying Benjamini-Hochberg *post-hoc* correction.

Supplementary Figure 26. Deltas of balance weighted alpha phylogenetic diversity between time points.

Change of alpha balance weighted phylogenetic diversity in samples between time points across the trial. On the y-axis of plots the percentage difference in balance weighted phylogenetic diversity between time points per cohort. Letters on the top left of each plot indicate the time points compared with one week interval (A-B, B-C, C-D, D-E) and two weeks interval (A-C, B-D). Pairwise t-test comparisons between cohorts were computed. Brackets in the timeline (top) indicate a significant *p* value (p<0.05) between cohorts (1-week interval); bold brackets indicate a significant *p* value (p<0.05) between cohorts (2-weeks interval). * indicate significance reached when applying Benjamini-Hochberg *post-hoc* correction.

Supplementary Figure 27. Significant differences in beta diversity between cohorts at specific time points.

Significance was determined by comparing groups by pairwise t-test and the resulting p values where adjusted with the Bonferroni method. Significance values are provided in Supplementary File 5. The x-axes represent the principal component. As plots are derived from distinct guppy runs, each principal component explains variation to a different extent (percentage specified in parentheses). The number of samples is specified on the y-axis. Distribution of the samples on either side of a plot (left *versus* right) reflects the taxa that were found to explain the variation. Distributions are colour coded by cohort.

Supplementary Figure 28. Correlation of weight with taxonomy at time point 0.

16S rRNA reads were extracted with SortMeRNA. Abundance was estimated and correlation with weight was assessed with the Spearman method. Significance values are provided in Supplementary File 5. Plotted are the correlations with p value < 0.05. For visualization purposes, subjects are binned into weight groups (x-axis). Bins are formed with subjects whose weight falls below the first quartile (red), above the first quartile (green), below the third quartile (blue), above the third quartile (purple). At the bottom of each whisker plot, the samples size for each bin is provided.

t2

Supplementary Figure 29. Correlation of weight with taxonomy at time point 2.

16S rRNA reads were extracted with SortMeRNA. Abundance was estimated and correlation with weight was assessed with the Spearman method. Significance values are provided in Supplementary File 5. Plotted are the correlations with *p* value < 0.05. For visualization purposes, subjects are binned into weight groups (x-axis). Bins are formed with subjects whose weight falls below the first quartile (red), above the first quartile (green), below the third quartile (blue), above the third quartile (purple). At the bottom of each whisker plot, the samples size for each bin is provided.

t4

Supplementary Figure 30. Correlation of weight with taxonomy at time point 4.

16S rRNA reads were extracted with SortMeRNA. Abundance was estimated and correlation with weight was assessed with the Spearman method. Significance values are provided in Supplementary File 5. Plotted are the correlations with p value < 0.05. For visualization purposes, subjects are binned into weight groups (x-axis). Bins are formed with subjects whose weight falls below the first quartile (red), above the first quartile (green), below the third quartile (blue), above the third quartile (purple). At the bottom of each whisker plot, the samples size for each bin is provided.

Supplementary Figure 31. Correlation of weight with taxonomy at time point 6.

16S rRNA reads were extracted with SortMeRNA. Abundance was estimated and correlation with weight was assessed with the Spearman method. Significance values are provided in Supplementary File 5. Plotted are the correlations with *p* value < 0.05. For visualization purposes, subjects are binned into weight groups (x-axis). Bins are formed with subjects whose weight falls below the first quartile (red), above the first quartile (green), below the third quartile (blue), above the third quartile (purple). At the bottom of each whisker plot, the samples size for each bin is provided.

t8

Supplementary Figure 32. Correlation of weight with taxonomy at time point 8.

16S rRNA reads were extracted with SortMeRNA. Abundance was estimated and correlation with weight was assessed with the Spearman method. Significance values are provided in Supplementary File 5. Plotted are the correlations with p value < 0.05. For visualization purposes, subjects are binned into weight groups (x-axis). Bins are formed with subjects whose weight falls below the first quartile (red), above the first quartile (green), below the third quartile (blue), above the third quartile (purple). At the bottom of each whisker plot, the samples size for each bin is provided.

Supplementary Figure 33. Correlation of weight with taxonomy at time point 10.

16S rRNA reads were extracted with SortMeRNA. Abundance was estimated and correlation with weight was assessed with the Spearman method. Significance values are provided in Supplementary File 5. Plotted are the correlations with *p* value < 0.05. For visualization purposes, subjects are binned into weight groups (x-axis). Bins are formed with subjects whose weight falls below the first quartile (red), above the first quartile (green), below the third quartile (blue), above the third quartile (purple). At the bottom of each whisker plot, the samples size for each bin is provided.

Supplementary Figure 34. Weight gain of piglets by cohort across the trial.

On the y-axis of plots the weight gain between time points is provided in percentage. Letters on the top left of each plot indicate the time points compared with one week interval (A-B, B-C, C-D, D-E) and with two weeks interval (B-D, C-E, A-E). Pairwise t-test comparisons between cohorts were computed. Brackets in the timeline (top) indicate a significant *p* value between cohorts (1-week interval); bold brackets indicate a significant *p* value between cohorts (2-weeks interval). A significant difference was found between Control and ColiGuard® (C-D, Tukey adjusted *p* value=0.000747) and between neomycin and neomycin+ColiGuard® (C-E, Tukey adjusted *p* value=0.039286). *in the timeline indicate that the significance was maintained after applying Tukey HSD *post-hoc* correction.