Title: Warming indirectly increases invasion success in food webs

Authors: Arnaud Sentis[1,2], Jose M. Montoya[1], and Miguel Lurgi[1,3,*]

Affiliations:

[1] Centre for Biodiversity Theory and Modelling, Theoretical and Experimental Ecology Station,

CNRS, 2 route du CNRS, 09200, Moulis, France

[2] INRAE, Aix-Marseille University, UMR RECOVER, 3275 route Cézanne, 13182 Aix-en-

Provence, France.

[3] Department of Biosciences. Swansea University, Singleton Park, Swansea, SA2 8PP, UK.

[*]corresponding author: miguel.lurgi@swansea.ac.uk

Email addresses:

Arnaud Sentis: arnaud.sentis@inrae.fr

José M Montoya: Josemaria.MONTOYATERAN@sete.cnrs.fr

Miguel Lurgi: miguel.lurgi@swansea.ac.uk

Running title: Warming and invasions in food webs

Keywords: climate change; body size; allometric bio-energetic model; food web structure; food

web stability; temperature dependence; top-down control

Type of article: Letter

Statement of authorship: AS and ML conceived research. All authors designed research. ML

performed modelling work. ML and AS analysed output data. JMM financially managed the

project, and obtained the funding. AS and ML wrote the first draft of the manuscript, and all

authors contributed substantially to revisions.

Data accessibility statement: Source code and simulation output data supporting the results

presented in this manuscript will be archived in an appropriate public repository (e.g. Dryad) and

the data DOI will be included at the end of the article.

Number of words in abstract: 150

Number of words in main text: 4995

Number of references: 70

Number of figures: 2

Number of tables: 2

Corresponding author: Miguel Lurgi – Room 225BA. Wallace Building. College of Science.

Swansea University. SA2 8PP. Swansea, UK. – Phone: +44 (0) 1792 602157 – email:

miguel.lurgi@swansea.ac.uk

ABSTRACT

Climate warming and biological invasions are key drivers of biodiversity change. Their combined effects on ecological communities remain largely unexplored. We investigated the direct and indirect influences of warming on invasion success, and their synergistic effects on community structure and dynamics. Using size-structured food web models, we found that warming increased invasion success. The direct physiological effects of warming on invasions were minimal in comparison to indirect effects mediated by changes on food web structure and stability. Warmed communities with less connectivity, shortened food chains and reduced temporal variability were more susceptible to invasions. The directionality and magnitude of invasions effects on food webs varied across warming regimes. Warmer communities became smaller, more connected, and with more predator species when invaded than their colder counterparts. They were also less stable and their species more abundant. Considering food web structure is crucial to predict invasion success and its impacts under warming.

INTRODUCTION

Climate warming and biological invasions constitute two of the most pervasive drivers of global change (Nelson 2005; Díaz *et al.* 2020). Both drivers strongly impact ecosystems, causing not only species loss, but also affecting ecological interactions and the structure of interaction networks (Stachowicz *et al.* 2002; Holzapfel & Vinebrooke 2005; Romanuk *et al.* 2009; Britton *et al.* 2010; Huang *et al.* 2011; Lurgi *et al.* 2012b; Lu *et al.* 2013; Lurgi *et al.* 2014; Zhang *et al.* 2017). Climate warming and species invasions can act synergistically on ecosystems due to, for example, impacts of climate change on species range dynamics (Thuiller *et al.* 2008; Elith & Leathwick 2009; Tylianakis & Morris 2017), which differentially influence species' ability to colonise new habitats and thus invade new communities. Species range shifts affect not only species composition, but also the structure of species interaction networks creating novel communities. For example, by promoting species range shifts, climate warming can trigger the loss of specialised interactions and changes the body size ratio between predator and prey species, which in turn can influence predator control on prey populations (Lurgi *et al.* 2012b, a). Yet, we know surprisingly little about how invasions and climate change act together to affect species and links in ecosystems.

Previous studies have shown that warming can enhance invasions by increasing survival and reproduction of introduced species (Mandrak 1989; Johnson & Evans 1990; Stachowicz *et al.* 2002; Logan *et al.* 2003; Britton *et al.* 2010; Huang *et al.* 2011; Ricciardi *et al.* 2017). However, warming can also lead to the opposite effect by decreasing the potential for invaders to occupy new niches (Bradley *et al.* 2010; Bertelsmeier *et al.* 2013). Recent evidence suggests that warming influence on invasion success may depend on how warming influences trophic interaction strength and the persistence of native predators or competitors (Holzapfel &

Vinebrooke 2005; Lu *et al.* 2013; Seifert *et al.* 2015). On the one hand, warming can prevent

invasions by increasing top-down control on the invader prey (Bradley *et al.* 2010; Lu *et al.*

2013; Lu *et al.* 2016). On the other hand, warming can enhance invasions by releasing top-down

control following predator extinctions (Holzapfel & Vinebrooke 2005). Overall, previous studies

reported various outcomes on the effects of warming on invasions. Our limited understanding of

their causes poses the challenge of gaining a better understanding of the indirect effects of

warming on species and communities. Indirect effects of warming on community structure and

species interactions are often stronger than its direct effects on physiology and demography

(Ockendon *et al.* 2014). This suggests that investigating warming effects on communities and the

complex networks of interactions that structure them is a first step to address this challenge.

Ecologists have developed mechanistic frameworks to identify key processes underlying

temperature effects trophic interactions and networks (Binzer *et al.* 2012; Burnside *et al.* 2014;

Fussmann *et al.* 2014; Gilbert *et al.* 2014; Sentis *et al.* 2014). One first important finding is that,

since consumer metabolic rates often increase faster with temperature than their feeding rates,

most consumers become less efficient at processing matter and energy at warmer temperatures

(Vucic-Pestic *et al.* 2011; Fussmann *et al.* 2014; Iles 2014). This reduction of energetic efficiency

lessens energy flow between trophic levels and, if resulting in weakened interaction strengths, it

stabilizes food-web dynamics by reducing population fluctuations (Rip & McCann 2011; Binzer

et al. 2012; Gilbert *et al.* 2014). A second important finding is that elevated temperatures increase

consumer extinction risk when metabolic demands exceed ingestion rates, leading to consumer

starvation and extinction (Petchey *et al.* 1999; Rall *et al.* 2010; Sentis *et al.* 2020). Whether these

changes would favour invasion success is unclear, as previous studies exploring the role of

community structure and dynamics in preventing or facilitating invasions success have not

considered modifications in communities driven by climate change (e.g. Romanuk *et al.* 2009;

Lurgi *et al.* 2014).

In parallel to studies focusing on the effects of warming, much effort has been devoted to

understanding how invasions impact ecosystems (Hui & Richardson 2019). Several models have

unveiled the role of food web structure such as species richness, complexity or the heterogeneity

of distribution of interactions in preventing invasions. These models have also suggested that

invasions in food webs tend to decrease species richness and shorten food chains (e.g. Romanuk

et al. 2009; Lurgi et al. 2014). However, a more recent theoretical investigation suggest the

opposite, with invasions being instead beneficial for maintaining species richness and ecological

functions (Zhang et al. (2017). We need comprehensive mechanistic frameworks incorporating

both warming and invasion and their effects on complex communities to better understand and

predict their synergistic effects.

Here, we explored the combined effects of warming and invasions on food webs using a

theoretical model. Given our current understanding of the effects of warming on natural

communities on the one hand (Binzer *et al.* 2012; Sentis *et al.* 2017; Boukal *et al.* 2019), and of

the invasion process in complex food webs on the other (Romanuk et al. 2009; Lurgi et al. 2014;

Hui & Richardson 2019), along with previous studies showing a positive influence of weakened

top-down control on invasions (Holzapfel & Vinebrooke 2005), we hypothesise that warming

increases invasion success if it decreases top-down control or causes predator extinctions. We

further hypothesise warming to increase connectivity and shorten food chains in response to

species loss at higher trophic levels. These effects of warming should in turn destabilise

community dynamics (Boukal *et al.* 2019).

To test these hypotheses, we investigated how temperature can influence invasions on complex

food webs comprised of 30 species. Our model simulates population dynamics in food webs

following bio-energetic principles of species life histories and interactions. The model

incorporates the temperature dependency of biological rates allowing the exploration of a wide

range of temperature regimes. We aim at gaining a better understanding of (1) warming effects

on invasion success in food webs and (2) the ecological consequences of invasions on food web

structure in warmed communities.

MATERIAL AND METHODS

We modelled community dynamics in complex food webs using a size-structured bio-energetic

community model consisting of a set of ordinary differential equations (ODEs) incorporating the

effects of species growth and ecological interactions (Yodzis & Innes 1992; Brose *et al.* 2006).

Food webs were generated using the niche model (Williams and Martinez 2000). The effect of

temperature on population dynamics was incorporated into ODEs by introducing thermal

dependencies of relevant model parameters. A series of numerical simulations were then

computed and species invasions were modelled as the addition of a new species into the

community (Lurgi et al. 2014). Simulation results were analysed to assess the effects of

temperature on (1) food web proprieties (structure, stability and total biomass) before invasion,

(2) invasion success, and (3) the effects of invasions on community structure and stability.

Food web generation

Food webs were generated using the niche model (Williams & Martinez 2000). With only two

parameters (number of species (S) and network connectance (C), i.e. the fraction of links out of

all possible ones) this model generates networks that resemble real food web structure (Williams

& Martinez 2000). We generated food webs comprising 30 species and with 10% connectance.

These values are within the ranges reported for empirical food webs (Williams & Martinez 2000).

We kept these values fixed across experiments to avoid the confounding effects of variation in

species richness and connectance.

Non-linear model for population dynamics

To simulate network dynamics of species populations biomass we used an allometric bio-

energetic model adapted from its original formulation by Yodzis and Innes (1992). This model

defines interactions strengths between prey and predator according to their body mass ratios

(Brose *et al.* 2006) and has been used to investigate warming and invasion effects on complex

food webs (Binzer *et al.* 2011; Lurgi *et al.* 2014). Eqn. 1 gives dynamics of basal resource

species. These grow logistically with an intrinsic growth rate r_i and a carrying capacity K_i.

Consumers gain biomass according to Eqn. 2:

151
$$\frac{dB_i}{dt} = r_i B_i \left(1 - \frac{B_i}{K_i}\right) - \sum_m F_{im} B_i \qquad (1)$$

152
$$\frac{dB_i}{dt} = \sum_s e F_{si} B_i - \sum_m F_{im} B_m - x_i B_i \qquad (2)$$

153   where B_i is the biomass of species i; e is the assimilation efficiency of predators when ingesting

154   prey (kept constant across consumer-resource species pairs at a value $e = 0.85$ (following Yodzis

155   & Innes 1992)); x_i is the metabolic rate at which biomass of consumers is lost from the system

156   due to respiration and other metabolic processes. F_{ij} is a function that describes the feeding

157   relationship between prey i and predator j and is defined by the functional response:

158
$$F_{ij} = \frac{\alpha_{ij} B_i^{\,q}}{1 + \sum_k h_{kj} \alpha_{kj} B_k^{\,q}} \qquad (3)$$

where α_{ij}'s are the elements of a quantitative version of the adjacency matrix A, describing the

food web obtained according to the procedure explained above (*Food web generation*), and that

represent the attack rates of predator species j on prey species i (Eqn. 5a). h_{ij} is the handling

time, i.e. the average time spent by an individual of predator species i handling and digesting an

individual of prey species i. The shape of the functional response curve is controlled by the

parameter q (i.e. the Hill exponent). We kept q constant across interacting species pairs at a value

of 1.2 to simulate an intermediate response type between Type II (hyperbolic, $q = 1$) and Type III

(sigmoidal, $q = 2$), as in (Binzer et al. 2011, 2016).

Growth, metabolic, attack rates, and handling times are functions of species body masses and

temperature. Body mass of species i scales according to its position in the food web:

$$m_i = m_0 R^{L_i - 1} \qquad (4)$$

where m_0 is the body size of basal species in the food web and set here to $m_0 = 0.01$g, R is the

average predator-prey body mass ratio of all trophic interactions in the food web and was set here

to $R = 10^2$, L_i is the prey-averaged trophic level of species i (Williams & Martinez 2004).

Allometric and thermal dependencies of model parameters were incorporated as follows (Binzer

et al. 2016):

$$\alpha_{ij}, h_{ij} = dm_i^b m_j^c e^{E\frac{T_0 - T}{kTT_0}} \qquad (5a)$$

$$r_i, x_i = dm_i^b e^{E\frac{T_0 - T}{kTT_0}} \qquad (5b)$$

where $d = e^I$ is a rate-specific constant calculated for a species with body mass of 1g and at a

reference temperature $T_0 = 20°C$ (293.15K), m_i and m_j are the body masses of species i and j,

respectively, *b* and *c* are rate-specific allometric exponents. The temperature dependence term is

a version of the Arrhenius equation in which *E* is the rate-specific activation energy and *k* is the

Boltzmann constant. *T* is the current temperature of the system in Kelvin. T_0 is the reference

temperature at which the rate value is equal to the rate-specific constant *d*. Values and units for

the parameters in Eqns. 5a and 5b are presented in Table 1.

Species carrying capacity was assumed to be independent of temperature since empirical

evidence for the thermal dependency of carrying capacity is inconclusive (Fussmann *et al.* 2014;

Gilbert *et al.* 2014; Uszko *et al.* 2017; Bernhardt *et al.* 2018). Furthermore, we wanted to avoid

biases in the invasion experiments due to the intrinsic limit to community biomass caused by the

negative temperature dependence of carrying capacity, which would, in turn, influence invasion

success. We thus focused on the effects of temperature on species life-history traits such as

reproduction, death and species interactions (i.e. attack rates and handling times) and not on the

maximum population density of the basal resources.

Food web structure, community properties and ecological stability

To assess the synergistic effects of temperature and invasions on food webs, we measured a set of

statistical food web properties (Table 2), before and after invasions, across a range of

temperatures. In addition to changes in food web properties we also assessed community

properties such as total community biomass, the average species body size in the community

(*AvgBS*), and the average ratio of predator vs. prey body masses (*AvgPPMR*). Lastly, to assess

ecological stability, we focused on temporal variability of biomass both at the community and

population levels. We quantified stability using two measures: (1) community invariability,

measured as the inverse of total community biomass variability and (2) population invariability,

as the inverse of average population-level biomass variability (Haegeman et al. 2016).

Variabilities, both at the community (i.e. summing across the biomass of all species populations)

and at the population level, were calculated as the ratio of the standard deviation to the mean

biomass across the last 100 years of the simulations.

Numerical simulations

Using the food web model specified above (Eqns. 1-5) we simulated a range of temperature

regimes and the addition of new species (i.e. invasions) as follows:

1. 140 niche model food webs were generated ($S = 30$ and $C = 0.1$).

2. Initial biomass densities for basal species were set to their carrying capacity $K \approx 2.75$

following the allometric formulation in Eqn. 5b but omitting the temperature dependent

term and assuming an allometric scaling constant and exponents of 10 and 0.28

respectively (Binzer *et al.* 2016). Consumer species initial abundances were set to 1/8 of

this value as in Binzer *et al.* (2016).

3. Community dynamics were first simulated for an equivalent of 600 years (18.9216×10^9

seconds) to achieve persistence (i.e. no further species extinctions) after initial transient

dynamics. After these first 600 years, food web and community properties were

calculated. The objective here was to quantify the direct effects of temperature on

community structure before invasion. Resulting community features, and their

relationship to temperature values, can then be related to invasion success.

4. After the first 600 years, an invasion was simulated by introducing a new species into the

network. The introduced species was randomly drawn as an additional species from the

original niche model network (i.e. ensuring it was different from all species originally

present in the network). Since at this time point some species might have gone extinct

(see step 6), rendering the potential introduced species disconnected, we repeated the

drawing procedure if necessary until a connected species was found. This procedure

avoided introducing an invasive species with no interactions.

5. After the introduction, we simulated further 600 years of network/community dynamics.

We then recorded whether the invasion was successful (i.e. whether the introduced

species persisted). Structural network and community properties were calculated again to

assess community structure after the introduced species became invasive (i.e. established

itself successfully in the community) or went extinct.

6. A species was considered extinct if at any point during the simulations its biomass fell

below 10^{-9} g.m^{-2}, at which point its abundance was set to 0.

For each of the 140 food webs, this procedure (i.e. steps 2 to 6 above) was repeated for each of 41

constant temperature regimes ranging from 0 to 40ºC at 1ºC intervals, yielding a total of 5740

numerical simulations. We used the same unique food webs for each temperature treatment to

avoid confounding effects caused by initial differences in food web structure.

Statistical analyses

The relationship between temperature and food web structure, stability and community properties

- i.e. total biomass, *AvgBS*, and *AvgPPMR* -, and their corresponding effects on invasion success

were analysed using piecewise structural equation models (SEMs). We computed SEMs

considering invasion success (i.e. whether the invasive species established after introduction) as a

response variable, with temperature affecting it directly and indirectly via network and

community properties. The effect of species richness on food web and community properties was

also incorporated into the SEMs, accounting thus for the indirect effect of temperature on

network and community properties via *S* (see Appendix S2 for full SEM details).

To assess invasion effects on food web structure, stability and community properties, we

calculated the ratio of values after (i.e. at the end of the simulations) vs. before (i.e. after initial

transient dynamics) invasion for food web (Table 2) and community properties, as well as

stability measures. To disentangle the direct and indirect effects (manifested via species richness)

of temperature on these ratios, we performed another SEM following the same rationale as above

and using the ratio of the effects as a response variable. We additionally assessed the differences

between communities vulnerable vs. resistant to invasions in terms of these effects by comparing

after/before ratios of each property between invaded and non-invaded communities using Mann-

Whitney U tests.

All simulations and analyses were performed in R -language and environment for statistical

computing (R Development Core Team 2017)-. Numerical simulations of ODEs were computed

using the ode function of the deSolve package (Soetaert et al. 2010). Food web analyses were

conducted with cheddar (Hudson et al. 2013). Modularity (Q) was computed using the

cluster_louvain function from the igraph package (Csardi & Nepusz 2006). Piecewise SEMs and

models within were performed using the piecewiseSEM (Lefcheck 2016) and the lme4 (Bates et

al. 2015) packages respectively. Computer code developed to run model simulations and analyse

outputs is available from the following repository: https://github.com/mlurgi/temperature-

dependent-invasions.

RESULTS

We focus on (i) the influence of warming on invasion success and (ii) the community-wide

consequences of invasions. Effects of warming on food webs before invasion are detailed in

Appendix S3. In line with previous findings (Binzer *et al.* 2016), we found that warmer

communities harbour less species than their colder counterparts, particularly at high trophic

levels, which in turn translates into higher connectance. These structural changes prompt an

increase in both community biomass and stability in warmer environments (Appendix S3).

How does warming influence invasion success?

Communities exposed to warmer temperatures were more prompt to invasions. Temperature had

a direct positive effect on invasion success (Fig. 1 and Table S1). This result holds even after

accounting for temperature effects on species richness and other community and network

properties (Table S1). Temperature influenced invasion success indirectly by modifying

community properties, thus making communities more susceptible to invasions. Direct effects of

network properties on invasion success were about an order of magnitude larger than the direct

effect of temperature (compare arrow weights -i.e. standardised coefficients- on Fig. 1). In

particular, the number of links (L), the average number of links per species (L/S), mean food

chain length ($MFCL$) and the fraction of basal and intermediate species (B and I respectively) had

a strong and significant influence on invasion success (Fig. 1 and Table S1). Surprisingly, we

found no direct effect of species richness on invasion success (Table S1) which is likely due to

the small variability in species richness in our communities after the initial transient dynamics

(Fig. S1a).

Even though warming had a direct significant effect on most network properties (Table S1, Figs.

S1-S3), only a few of them affected invasion success. In particular, we found that communities

with longer food chains (*MFCL)* were more resistant to invasion. In addition, communities with

more links (*L*) and greater proportions of basal (*B*) and intermediate (*I*) species were more prompt

to invasion (Fig. 1). Communities harbouring more specialised species (i.e. small *L/S*) also were

more susceptible to invasion.

Changes in population stability and total community biomass also affected invasion success

under warming. Whereas larger total community biomass conferred resistance against invasions,

communities with higher population stability were easier to invade (Fig. 1). Overall our results

show that indirect effects of temperature on invasion success, mediated by changes in network

and community properties and dynamics, were stronger than direct ones.

Ecological consequences of invasions along the temperature gradient

Overall, invasions strongly decreased species richness (Fig. 2a), which, in turn, affected several

network properties. Moreover, we found that the magnitude of the change of food web and

community properties driven by invasions often depended on temperature (Fig. 2 & Table S2). In

particular, warmer communities loose more species and interactions when invaded than their

colder counterparts. This translates into more connected communities (Fig. 2a). Higher

connectance (*C*) was accompanied by a larger heterogeneity in the distribution of predators

among prey species (V*ulSD*) and a stronger increase in the fraction of top predators (*T*). On the

other hand, warming prompted a stronger decrease in the proportion of intermediate species (*I*)

when invaded. The proportion of basal species (B) was not influenced by invasion or temperature

(Table S2). This suggests that intermediate species became top predators when their predators

disappeared in invaded communities, yielding a lower fraction of intermediates while increasing

the fraction of top predator species. Warming also lead to a stronger decrease in modularity (*Q*)

which is explained by the presence of a larger fraction of top consumers and a higher

connectance.

Community properties and stability were affected more heterogeneously by invasions across

temperature regimes than network structure (Fig. 2). Total community biomass increased in

warm invaded communities relative to their pre-invasion state (positive effect sizes), whereas it

decreased under colder conditions (Fig. 2b, negative effect sizes). Similarly, population stability

increased in colder communities (at 0 and 5°C) but decreased in warmer ones (although the effect

was weak at 35°C). In addition, invasions always decreased community stability (Fig. 2b),

although this effect was weaker in colder environments compared to warmer ones. Overall, we

found that warm invaded communities have less species, with these species fluctuating more over

time (i.e. reduced stability) than cold invaded communities.

Lastly, the average body size (*AvgBS*) of species in the community remains mostly unaffected by

invasions except at very low temperatures where invasions decreased average body size. On the

other hand, the average predator:prey body size ratio (*AvgPPMR*) was negatively impacted by

invasion and this effect was more pronounced when temperature increased (Fig. 2b). The

decrease in body size ratio, along with the considerable increase in the fraction of top predator

species (Fig. 2a), while the fraction of basal species was unaffected by invasion or warming

reinforces the observation that top predators were lost and replaced by consumers further down

the food web. Interestingly, this replacement appeared to be stronger in warmer communities that

also lost more species than colder ones.

The effects of unsuccessful invasions (i.e. when the introduced species went extinct) on food

webs were more homogeneous across the temperature gradient than those caused by successful

invasions, mainly affecting species numbers, connectivity (*L* and *L/S*), and the fraction of

intermediate species (Appendix S4).

DISCUSSION

Global warming and biological invasions affect communities simultaneously. It is fundamental to

better understand their synergistic effects on biodiversity (Bradley *et al.* 2010). Using dynamical

models we have investigated the interactive effects of warming and invasions on the structure and

dynamics of complex food webs. We showed that warming has two overall effects on invasions

and on invaded communities. First, it modified key aspects of community structure and dynamics

that in turn facilitated invasions by introduced species. Secondly, warming mostly amplified the

impacts of invasions on the structure and organisation of communities. Importantly, the

directionality of the effects of invasions on recipient communities changed across the temperature

gradient for community biomass and population stability. Further, our results suggest that the

direct effects of temperature on invasion success are weaker to those mediated by changes in

community structure and stability.

Previous attempts to understand and predict the combined effects of warming and invasions have

been based mainly on bioclimatic envelop models, relying on temperature thresholds for survival

and reproduction of the invasive species (Stachowicz *et al.* 2002; Walther *et al.* 2009; Sorte *et al.*

2010; Kent *et al.* 2018). Such phenomenological approaches, even though informative, lack a

mechanistic understanding of how warming mediates invasions and how both can synergistically

affect ecological communities. Here, we have provided a first step towards a better understanding

of the synergistic effects of warming and invasions on complex ecological communities. Our

efforts complement recent attempts to better understand how warming modulates the effects of

invasions on natural communities (Zhang et al. 2017).

Warming effects on invasion success

We found that, before invasion, warming increased food web stability but decreased the

persistence of top predators, as reported in previous studies (Fussmann *et al.* 2014; Binzer *et al.*

2016; Sentis *et al.* 2017). This increased stability is likely explained by average trophic

interaction strength decreasing and by consumers being less efficient at feeding relative to their

metabolic losses (Boukal *et al.* 2019). Warmer communities thus contained less species, but

network connectance increased. Whether these warming-induced changes influence invasion

success remains an open question.

Previous theoretical models showed that more connected food webs are generally better at

repelling invaders (Romanuk *et al.* 2009). In contrast, Lurgi *et al.* (2014) showed that, when

controlling for the number of species, less connected food webs are more resistant to invasions.

Here, we found that invasions are more successful in warmer communities, which are more

connected but poorer in species richness than colder ones. Invasion success was strongly

mediated by the indirect effects of temperature on network and community properties and

stability, with these effects only weakly mediated by species richness. Temperature had an impact

on the composition of species across trophic levels, influencing greatly the proportion of top

predators. This, in turn, made communities more susceptible to invasions by reducing mean food

chain length. We also found that warming directly increased invasion success. Nevertheless, the

direct effect of warming on invasion success was much weaker than its indirect effects.

Our results are in line with empirical studies reporting that the influence of warming on invasion

success may depend on how warming affects the strength of trophic interactions and the

persistence of local predators or competitors (Holzapfel & Vinebrooke 2005; Lu *et al.* 2013;

Seifert *et al.* 2015). Warming can prevent invasions by increasing top-down control on

introduced species (Lu *et al.* 2013; Lu *et al.* 2016). On the other hand, Holzapfel and Vinebrooke

(2005) showed that warming can enhance invasions by removing top-down control on the

invader following predator extinctions. Our model results suggest that the loss of top predators,

which in turn reduces mean food chain length, is a driving mechanism of invasion success. This

highlights the importance of species loss at high trophic levels on facilitating invasions under

warming.

Ecological consequences of invasions along a temperature gradient

Our results are in line with previous studies showing that invasions decrease species richness and,

as a consequence, impact strongly community structure (Hui & Richardson 2019). However,

recent theoretical studies have suggested that warming is predicted to have an opposite effect on

invasions, by enhancing species richness and ecosystem functioning (Zhang et al. 2017). We

found that temperature can modulate the consequences of invasions for community structure.

Warmer communities tend to lose more species and interactions when invaded, which translates

into more connected communities compared to colder ones. This intensification in the effects of

invasions across the temperature range highlights the importance of considering a full spectrum

of temperature treatments to fully understand the effects of warming on invaded communities.

The proportion of top predator species in our model food webs increased as a consequence of

invasion, especially in warmer environments. This result apparently contradicts the loss of top

predators following invasions that we observed. Both observations are reconciled by a third

result: invasions shorten food chains. In short, when top predators go extinct they are replaced by

consumers further down the food web, which in turn become top predators. This switch decreases

the proportion of intermediate species while increasing the proportion of top species, ultimately

shortening food chain length. Invasions thus exacerbate the previously observed effect of

warming on top predator species, and corroborates previous empirical findings of higher trophic

levels being most vulnerable to climate change (Voigt *et al.* 2003). We should thus expect

warmer and invaded communities being even more susceptible to invasions, entering a positive

feedback loop via the loss of predator species.

As any modelling study, ours relies on a set of assumptions that can influence model predictions.

For instance, we did not account for temperature fluctuations, evolutionary change or differences

in the thermal traits of the invasive species (e.g. the invasive species may have thermal traits

better adapted to warmer climate) that can also influence invasion success and species persistence

(Vasseur *et al.* 2014; Zhang *et al.* 2017). The assumption of invasive species possessing similar

thermal performance than resident species is supported by the observation that most invasive

species are introduced by human transport and their invasion success is not proven to be strongly

related to the climate in their native environment (Gippet *et al.* 2019). Additionally, in our model,

introduced species, and their biotic interactions, have been defined following the same heuristics

as the rest of the species in the recipient community (i.e. they are drawn from the niche model),

whereas in natural communities invasive species tend to be generalists, both in terms of habitats

and interactions (Elton 1958). We decided to define introduced species in this way to be as

parsimonious as possible in our experiments and avoid the confounding effects of over-generalist

species in modelled communities. It would be interesting to extend the results presented here

using invasive species with more realistic traits. Despite these limitations, our model provides a

first step in the exploration of the consequences of warming and invasions in species-rich

communities.

CONCLUSION

Research on the effects of temperature on biological invasions has traditionally focused on the

species level and have not explicitly considered species interactions. Usually not taking into

account the way temperature can affect food web and community properties prior to invasions.

Understanding the joint effects of warming and invasions at the community level constitutes a

pressing challenge to unveil the full consequences of global change on natural communities. Here

we addressed this challenge and showed that temperature's direct effects on invasion success are

weaker than its indirect effects mediated by changes in food web structure, community properties

and stability. Moreover, we showed that the impact of invasions depend on the temperature

experienced by the invaded communities. Warmer food webs lose more species and interactions

when invaded than their colder counterparts. These changes are accompanied by an increase in

the fraction of top predator species, enhanced total community biomass and decreased stability.

Overall, our study suggest that both warming and invasion act synergistically to fasten species

loss creating smaller and more connected networks. It paves way for a better understanding of the

causes and consequences of invasions in a warming world.

ACKNOWLEDGEMENTS

This work was funded by the FRAGCLIM Consolidator Grant (number 726176) from the

European Research Council under the European Union's Horizon 2020 Research and Innovation

Program. ML was additionally supported by the French ANR through LabEx TULIP (ANR-10-

LABX-41; ANR-11-IDEX-002-02), and by a Region Midi-Pyrénées Project (CNRS 121090) to

JMM. AS was additionally supported by the ANR project EcoTeBo (ANR-19-CE02-0001-01)

from the French National Research Agency (ANR).

REFERENCES

Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015). Fitting linear mixed-effects models

using lme4. *Journal of Statistical Software*, 67, 1-48.

Bellard, C., Cassey, P. & Blackburn, T.M. (2016). Alien species as a driver of recent extinctions.

Biology letters, 12, 20150623.

Bernhardt, J.R., Sunday, J.M. & O'Connor, M.I. (2018). Metabolic theory and the temperature-

size rule Explain the temperature dependence of population carrying capacity. *The*

American Naturalist, 192, 687-697.

Bertelsmeier, C., Luque, G.M. & Courchamp, F. (2013). Global warming may freeze the invasion

of big-headed ants. *Biological Invasions*, 15, 1561-1572.

Binzer, A., Brose, U., Curtsdotter, A., Eklöf, A., Rall, B.C., Riede, J.O. *et al.* (2011). The

susceptibility of species to extinctions in model communities. *Basic and Applied Ecology*,

12, 590-599.

Binzer, A., Guill, C., Brose, U. & Rall, B.C. (2012). The dynamics of food chains under climate

change and nutrient enrichment. *Philosophical Transactions of the Royal Society B:*

Biological Sciences, 367, 2935-2944.

Binzer, A., Guill, C., Rall, B.C. & Brose, U. (2016). Interactive effects of warming,
eutrophication and size structure: impacts on biodiversity and food☐web structure.
Global Change Biology, 22, 220-227.

Blondel, V.D., Guillaume, J.-L., Lambiotte, R. & Lefebvre, E. (2008). Fast unfolding of
communities in large networks. *Journal of statistical mechanics: theory and experiment*,
2008, P10008.

Boukal, D.S., Bideault, A., Carreira, B.M. & Sentis, A. (2019). Species interactions under climate
change: connecting kinetic effects of temperature on individuals to community dynamics.
Current opinion in insect science, 35, 88-95.

Bradley, B.A., Blumenthal, D.M., Wilcove, D.S. & Ziska, L.H. (2010). Predicting plant invasions
in an era of global change. *Trends in Ecology & Evolution*, 25, 310-318.

Britton, J.R., Cucherousset, J., Davies, G.D., Godard, M.J. & Copp, G.H. (2010). Non-native
fishes and climate change: predicting species responses to warming temperatures in a
temperate region. *Freshwater Biology*, 55, 1130-1141.

Broennimann, O. & Guisan, A. (2008). Predicting current and future biological invasions: both
native and invaded ranges matter. *Biology letters*, 4, 585-589.

Brose, U., Williams, R.J. & Martinez, N.D. (2006). Allometric scaling enhances stability in
complex food webs. *Ecology Letters*, 9, 1228-1236.

Burnside, W.R., Erhardt, E.B., Hammond, S.T. & Brown, J.H. (2014). Rates of biotic interactions
scale predictably with temperature despite variation. *Oikos*, 123, 1449-1456.

Csardi, G. & Nepusz, T. (2006). The igraph software package for complex network research.
InterJournal, Complex Systems, 1695, 1-9.

Delmas, C., Delzon, S. & Lortie, C. (2011). A meta-analysis of the ecological significance of

density in tree invasions. *Community ecology*, 12, 171-178.

Díaz, S., Settele, J., Brondízio, E., Ngo, H., Guèze, M., Agard, J. *et al.* (2020). *Summary for*

policymakers of the global assessment report on biodiversity and ecosystem services of

the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.

IPBES secretariat, Bonn, Germany, 56 pages.

Elith, J. & Leathwick, J.R. (2009). Species distribution models: ecological explanation and

prediction across space and time. *Annual review of ecology, evolution, and systematics*,

40, 677-697.

Elton, C.S. (1958). *The ecology of invasions by animals and plants*. Methuen and Co. Ltd,

London.

Fussmann, K.E., Schwarzmüller, F., Brose, U., Jousset, A. & Rall, B.C. (2014). Ecological

stability in response to warming. *Nature Climate Change*, 4, 206-210.

Gilbert, B., Tunney, T.D., McCann, K.S., DeLong, J.P., Vasseur, D.A., Savage, V. *et al.* (2014).

A bioenergetic framework for the temperature dependence of trophic interactions.

Ecology Letters, 17, 902-9014.

Gippet, J., Liebhold, A., Fenn-Moltu, G. & Bertelsmeier, C. (2019). Human-mediated dispersal

in insects. *Current opinion in insect science*, 35, 96-102.

Haegeman, B., Arnoldi, J.-F., Wang, S., de Mazancourt, C., Montoya, J.M. & Loreau, M. (2016).

Resilience, Invariability, and Ecological Stability across Levels of Organization. *bioRxiv*,

085852.

Holzapfel, A.M. & Vinebrooke, R.D. (2005). Environmental warming increases invasion potential of alpine lake communities by imported species. *Global Change Biology*, 11, 2009-2015.

Huang, D., Haack, R.A. & Zhang, R. (2011). Does Global Warming Increase Establishment Rates of Invasive Alien Species? A Centurial Time Series Analysis. *PLOS ONE*, 6, e24733.

Hudson, L.N., Emerson, R., Jenkins, G.B., Layer, K., Ledger, M.E., Pichler, D.E. *et al.* (2013). Cheddar: analysis and visualisation of ecological communities in R. *Methods in Ecology and Evolution*, 4, 99-104.

Hui, C. & Richardson, D.M. (2019). How to invade an ecological network. *Trends in ecology & evolution*, 34, 121-131.

Iles, A.C. (2014). Towards predicting community level effects of climate: relative temperature scaling of metabolic and ingestion rates. *Ecology*, 95, 2657–2668.

Johnson, T.B. & Evans, D.O. (1990). Size-Dependent Winter Mortality of Young-of-the-Year White Perch: Climate Warming and Invasion of the Laurentian Great Lakes. *Transactions of the American Fisheries Society*, 119, 301-313.

Kent, A., Drezner, T.D. & Bello, R. (2018). Climate warming and the arrival of potentially invasive species into boreal forest and tundra in the Hudson Bay Lowlands, Canada. *Polar Biology*, 41, 2007-2022.

Lefcheck, J.S. (2016). piecewiseSEM: Piecewise structural equation modelling in r for ecology, evolution, and systematics. *Methods in Ecology and Evolution*, 7, 573-579.

Logan, J.A., Régnière, J. & Powell, J.A. (2003). Assessing the impacts of global warming on forest pest dynamics. *Frontiers in Ecology and the Environment*, 1, 130-137.

Lu, X., Siemann, E., He, M., Wei, H., Shao, X. & Ding, J. (2016). Warming benefits a native

species competing with an invasive congener in the presence of a biocontrol beetle. *New*

Phytologist, 211, 1371-1381.

Lu, X., Siemann, E., Shao, X., Wei, H. & Ding, J. (2013). Climate warming affects biological

invasions by shifting interactions of plants and herbivores. *Global Change Biology*, 19,

2339-2347.

Lurgi, M., Galiana, N., López, B.C., Joppa, L.N. & Montoya, J.M. (2014). Network complexity

and species traits mediate the effects of biological invasions on dynamic food webs.

Frontiers in Ecology and Evolution, 2, 36.

Lurgi, M., López, B.C. & Montoya, J.M. (2012a). Climate change impacts on body size and food

web structure on mountain ecosystems. *Philosophical Transactions of the Royal Society*

B: Biological Sciences, 367, 3050-3057.

Lurgi, M., López, B.C. & Montoya, J.M. (2012b). Novel communities from climate change.

Philosophical Transactions of the Royal Society B: Biological Sciences, 367, 2913-2922.

Mandrak, N.E. (1989). Potential invasion of the Great Lakes by fish species associated with

climatic warming. *Journal of Great Lakes Research*, 15, 306-316.

Mellin, C., Lurgi, M., Matthews, S., MacNeil, M.A., Caley, M., Bax, N. *et al.* (2016).

Forecasting marine invasions under climate change: Biotic interactions and demographic

processes matter. *Biological Conservation*, 204, 459-467.

Montoya, J.M. & Raffaelli, D. (2010). Climate change, biotic interactions and ecosystem

services. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365,

2013-2018.

Nelson, G.C. (2005). *Millennium ecosystem assessment: drivers of ecosystem change: summary*

chapter. World Resources Institute, Washington, DC.

Ockendon, N., Baker, D.J., Carr, J.A., White, E.C., Almond, R.E., Amano, T. *et al.* (2014).

Mechanisms underpinning climatic impacts on natural populations: altered species

interactions are more important than direct effects. *Global Change Biology*.

Petchey, O.L., McPhearson, P.T., Casey, T.M. & Morin, P.J. (1999). Environmental warming

alters food-web structure and ecosystem function. *Nature*, 402, 69-72.

R Development Core Team (2017). *R: a language and environment for statistical computing*, R

Foundation for Statistical Computing, Vienna, Austria.

Rall, B.C., Vucic-Pestic, O., Ehnes, R.B., Emmerson, M. & Brose, U. (2010). Temperature,

predator–prey interaction strength and population stability. *Global Change Biology*, 16,

2145-2157.

Reino, L., Moya-Laraño, J. & Cláudio Heitor, A. (2009). Using survival regression to study

patterns of expansion of invasive species: will the common waxbill expand with global

warming? *Ecography*, 32, 237-246.

Ricciardi, A., Blackburn, T.M., Carlton, J.T., Dick, J.T.A., Hulme, P.E., Iacarella, J.C. *et al.*

(2017). Invasion Science: A Horizon Scan of Emerging Challenges and Opportunities.

Trends in Ecology & Evolution, 32, 464-474.

Rip, J.M.K. & McCann, K.S. (2011). Cross-ecosystem differences in stability and the principle of

energy flux. *Ecology Letters*, 14, 733-740.

Romanuk, T.N., Zhou, Y., Brose, U., Berlow, E.L., Williams, R.J. & Martinez, N.D. (2009).

Predicting invasion success in complex ecological networks. *Philosophical Transactions*

of the Royal Society B: Biological Sciences, 364, 1743-1754.

Seifert, L., Weithoff, G., Gaedke, U. & Vos, M. (2015). Warming-induced changes in predation,

extinction and invasion in an ectotherm food web. *Oecologia*, 1-12.

Sentis, A., Binzer, A. & Boukal, D.S. (2017). Temperature-size responses alter food chain

persistence across environmental gradients. *Ecology Letters*, 20, 852-862.

Sentis, A., Haegeman, B. & Montoya, J.M. (2020). Stoichiometric constraints modulate the

effects of temperature and nutrients on biomass distribution and community stability.

bioRxiv, 589895, ver. 7 peer-reviewed and recommended by Peer Community in Ecology,

DOI: 10.1101/589895.

Sentis, A., Hemptinne, J.L. & Brodeur, J. (2014). Towards a mechanistic understanding of

temperature and enrichment effects on species interaction strength, omnivory and food-

web structure. *Ecology Letters*, 17, 785-793.

Simberloff, D., Martin, J.-L., Genovesi, P., Maris, V., Wardle, D.A., Aronson, J. *et al.* (2013).

Impacts of biological invasions: what's what and the way forward. *Trends in ecology &*

evolution, 28, 58-66.

Soetaert, K., Petzoldt, T. & Setzer, R.W. (2010). Solving differential equations in R: package

deSolve. *Journal of Statistical Software*, 33, 1-25.

Sofaer, H.R., Jarnevich, C.S. & Pearse, I.S. (2018). The relationship between invader abundance

and impact. *Ecosphere*, 9, e02415.

Sorte, C.J.B., Williams, S.L. & Zerebecki, R.A. (2010). Ocean warming increases threat of

invasive species in a marine fouling community. *Ecology*, 91, 2198-2204.

Stachowicz, J.J., Terwin, J.R., Whitlatch, R.B. & Osman, R.W. (2002). Linking climate change

and biological invasions: Ocean warming facilitates nonindigenous species invasions.

Proceedings of the National Academy of Sciences, 99, 15497-15500.

Thuiller, W., Richardson, D.M. & Midgley, G.F. (2008). Will climate change promote alien plant

invasions? In: *Biological invasions*. Springer, pp. 197-211.

Tylianakis, J.M., Didham, R.K., Bascompte, J. & Wardle, D.A. (2008). Global change and

species interactions in terrestrial ecosystems. *Ecology Letters*, 11, 1351-1363.

Tylianakis, J.M. & Morris, R.J. (2017). Ecological networks across environmental gradients.

Annual Review of Ecology, Evolution, and Systematics, 48, 25-48.

Urban, M.C. (2015). Accelerating extinction risk from climate change. *Science*, 348, 571-573.

Uszko, W., Diehl, S., Englund, G. & Amarasekare, P. (2017). Effects of warming on predator–

prey interactions – a resource-based approach and a theoretical synthesis. *Ecology Letters*,

20, 513-523.

Vasseur, D.A., DeLong, J.P., Gilbert, B., Greig, H.S., Harley, C.D., McCann, K.S. *et al.* (2014).

Increased temperature variation poses a greater risk to species than climate warming.

Proceedings of the Royal Society B: Biological Sciences, 281, 20132612.

Voigt, W., Perner, J., Davis, A.J., Eggers, T., Schumacher, J., Bährmann, R. *et al.* (2003).

Trophic levels are differentially sensitive to climate. *Ecology*, 84, 2444-2453.

Vucic-Pestic, O., Ehnes, R.B., Rall, B.C. & Brose, U. (2011). Warming up the system: higher

predator feeding rates but lower energetic efficiencies. *Global Change Biology*, 17, 1301-

1310.

Walther, G.-R., Roques, A., Hulme, P.E., Sykes, M.T., Pyšek, P., Kühn, I. *et al.* (2009). Alien

species in a warmer world: risks and opportunities. *Trends in ecology & evolution*, 24,

686-693.

Williams, R.J. & Martinez, N.D. (2000). Simple rules yield complex food webs. *Nature*, 404,

180-183.

Yodzis, P. & Innes, S. (1992). Body size and consumer-resource dynamics. *The American*

Naturalist, 139, 1151-1175.

Zhang, L., Takahashi, D., Hartvig, M. & Andersen Ken, H. (2017). Food-web dynamics under

climate change. *Proceedings of the Royal Society B: Biological Sciences*, 284, 20171772.

624   SUPPORTING INFORMATION

Additional Supporting Information may be downloaded via the online version of this article at

Wiley Online Library (www.ecologyletters.com).

Tables

Table 1. *Parameter values for mass and temperature dependencies*. Parameters extracted /

calculated from: growth (r in [1/s], (Savage et al., 2004)), metabolism (x in [1/s], (Ehnes et al.,

2011)), attack rate (α in [m^2 s^{-1}]) and handling time (h in [s]), both calculated from (Rall et al.,

2012). Metabolic rates were calculated using the conversion factor from Peters (1983).

	r_i	α_{ij}	h_{ij}	x_i
Intercept (I)	-15.68	-13.1	9.66	-16.54
Body mass scaling species i (b)	-0.25	0.25	-0.45	-0.31
Body mass scaling predator (c)		-0.8	0.47	
Activation energy (K)	-0.84	-0.38	0.26	-0.69

634   **Table 2.** *Network structural properties used in this study to quantify the structure of simulated*

635   *food webs.* The name of the properties, their abbreviations and a brief description are presented.

Property	*Abbrev.*	*Description*
Species richness	*S*	Number of species in the community.
Number of links	*L*	Total number of trophic interactions in the food web.
Links per species	*L/S*	Average number of links (i.e. interactions) per species.
Connectance	*C*	Fraction of realised links out of all the possible links in the food web. Calculated as: $C = L/S^2$
Standard deviation of generality	*GenSD*	Quantifies the variability of species' normalised number of prey (or their generality) $G_j = \frac{1}{L/S} \sum_{i=1}^{S} a_{ij}$, where a_{ij} are the elements of A, the adjacency matrix representing the food web and in which if $a_{ij} = 1$ there is a trophic interaction between prey i and predator j.
Standard deviation of vulnerability	*VulSD*	Quantifies the variability of species' normalised number of predators (or their vulnerability) $G_i = \frac{1}{L/S} \sum_{j=1}^{S} a_{ij}$
Mean food chain length	*MFCL*	Average length (i.e. number of links) of all the paths (food chains) running from each basal species to each top predator species in the food web.
Fraction of basal species	*B*	Fraction of basal species, out of the total number of

species, in the food web. Basal species are those with no prey (i.e. no incoming links).

Fraction of intermediate species	I	Fraction of species in the food web having both prey and predators (i.e. sitting in the 'middle' of the food web).
Fraction of top species	T	Fraction of top consumers in the food web (i.e. species that do not have predators).
Maximum similarity	$MxSim$	The average maximum trophic similarity across species in the network. Trophic similarity (s_{ij}) of a pair of species is the number of common prey and predator species divided by the pair's total number of predator and prey species. Thus, $MxSim = \frac{1}{S}\sum_{i=1}^{S} max_{\forall i \neq j}(s_{ij})$
Modularity	Q	A measure that quantifies the extent to which species within the same compartment share more interactions among themselves than with species in other compartments (see Appendix S1 in Supporting Information for details on modularity calculation).

637 **Figures**

638 **Figure 1.**

Figure 2.

 Figure legends

Figure 1. *Structural equation model (SEM) describing the direct and indirect effects of*

temperature on invasion success in complex food webs. Arrows indicate the direct effects of

predictor on response variables. Only predictors having a statistically significant effect (i.e. p-

value < 0.05) on invasion success are shown (see Table S1 for more details). Black and red

arrows represent positive and negative effects respectively. Solid, dashed, and dotted arrow styles

represent strongly (p-value < 0.001), intermediate (p-value < 0.01), and marginally (p-value <

0.05) statistically significant effects respectively. Model fit: Fisher's C = 22.67, p-value = 0.305,

degrees of freedom = 20.

Figure 2. *Effect sizes (mean ± s.d.) of successful invasions on complex food webs across*

temperature regimes on network (a) and community (b) properties. Effect sizes were quantified

as the ratio between the values of the network/community property after species introduction vs.

before introduction, minus unity. Negative values thus indicate negative effects of the invasion

on the community (i.e. the value after the invader's establishment is smaller than before the

introduction). Only effects on properties identified by SEMs as being significantly influenced by

temperature (Table S2) are shown. Only a subset of temperature regimes is shown to ease the

visualisation of the results.