1 **ALG-2 interacting protein-X (Alix) is required for activity-dependent bulk**
2 **endocytosis at brain synapses**
3

4 Marine H. Laporte [1,2,*], Kwang Il Chi[1*], Laura C. Caudal[3], Na Zhao[3], Marta Rolland[1], , José
5 Martinez-Hernandez[1,4], Magalie Martineau[5], Christine Chatellard[1,6] Eric Denarier[1], Vincent
6 Mercier[1,7], Florent Lemaître [1,8], Béatrice Blot[1], Eve Moutaux[1], Maxime Cazorla[1,9], David
7 Perrais[5], Fabien Lanté[1], Sandrine Fraboulet[1,10] , Fiona J. Hemming[1], Frank Kirchhoff[3], Rémy
8 Sadoul[1,6£]
9
10
1) Grenoble Institute Neurosciences, University Grenoble Alpes- INSERM U1216- CEA, Grenoble, France
2) Department of Cell Biology, University of Geneva, Geneva, Switzerland.
3) Molecular Physiology, Center for Integrative Physiology and Molecular Medicine (CIPMM), University of
Saarland, Homburg, Germany.
4) Department of Biochemistry and Molecular Biology, Faculty of Science and Technology, University of
the Basque Country (UPV/EHU), Leioa, Spain.
5) Interdisciplinary Institute for Neuroscience, University of Bordeaux-CNRS UMR 5297, Bordeaux, France
6) Institut de Biologie Structurale, University Grenoble Alpes- CNRS-CEA UMR 5075, Grenoble, France
7) Department of Biochemistry, University of Geneva, Geneva, Switzerland.
8) Département des Neurosciences, Centre de Recherche du centre hospitalier de Montréal, Canada
9) Institut de Neuroscience la Timone, CNRS UMR7289, Marseille, France
10) Institute for Advanced Biosciences, Univ. Grenoble Alpes - INSERM U1209 - CNRS UMR 5309, Grenoble,
France
24
25
26 *Equal contribution
27 £Corresponding author
28 email: remy.sadoul@univ-grenoble-alpes.fr

29 **Running title:** Alix and ADBE
30
31 Number of pages: 35
32 Items number: 7 figures, 2 movies.
33 Words count: abstract (176), introduction (660), discussion (1164).
34
35

Abstract

In chemical synapses undergoing high frequency stimulation, vesicle components can be retrieved from the plasma membrane via a clathrin-independent process called activity dependent bulk endocytosis (ADBE). Alix (ALG-2 interacting protein X)/ PDCD6IP) is an adaptor protein binding to ESCRT and endophilin-A proteins which is required for clathrin-independent endocytosis in fibroblasts. Alix is expressed in neurons and concentrates at synapses during epileptic seizures. Here, we used cultured neurons to show that Alix is recruited to presynapses where it interacts with, and concentrates endophilin-A during conditions triggering ADBE. Using Alix knockout (ko) neurons we showed that this recruitment, which requires interaction with the calcium-binding protein ALG-2, is necessary for ABDE. We also found that presynaptic compartments of Alix ko hippocampi display subtle morphological defects compatible with flawed synaptic activity and plasticity detected electrophysiologically. Furthermore, mice lacking Alix in the forebrain undergo less seizures during kainate-induced status epilepticus and reduced propagation of the epileptiform activity. These results thus show that impairment of ADBE due to the lack of neuronal Alix alters synaptic recovery during physiological or pathological repeated stimulations.

Keywords: PDCD6IP**,** Clathrin-independent-endocytosis, Alix knock-out, endophilin-A, ALG-2, bulk endocytosis, synaptic vesicle recycling, synaptic transmission, calcium, epileptic seizures.

Introduction

Neuronal communication in mammalian brain relies heavily on the activity-dependent release of chemical neurotransmitters from presynaptic boutons. Following fusion of synaptic vesicles (SV) with the presynaptic membrane, SV lipids and proteins are retrieved by endocytosis. Endocytosis avoids detrimental increase in the plasma membrane surface and allows recycling of the SV components to replenish the SV pool (Gan and Watanabe, 2018). At moderate levels of stimulation, retrieval of membrane involves clathrin-mediated (CME) and clathrin-independent endocytosis (CIE), in proportions which are still highly debated (Chanaday and Kavalali, 2018). Moreover, long-lasting high-frequency stimulations also lead to the clathrin-independent internalization of large stretches of presynaptic membranes. This calcium-dependent process, first discovered at the amphibian neuromuscular junction (Miller and Heuser, 1984) is referred to as activity-dependent bulk endocytosis (ADBE). It is meant to avoid abnormal increase of the synaptic bouton surface and to allow replenishment of SVs during sustained synaptic stimulations (Cheung and Cousin, 2012; Marxen et al., 1999). Thus, ADBE has been suggested to play key regulatory roles in physiological or pathological events like epilepsy, which are triggered and sustained by high frequency neuronal activity. However, decrypting the physiological role of ADBE has been hindered by the lack of identified molecules that are both specific and essential to this endocytosis mode.

We have recently demonstrated that ALG-2 interacting protein-X (Alix) is essential for clathrin-independent bulk endocytosis in fibroblasts (Mercier et al., 2016). Alix ko mice have normally organized but smaller brains (Campos et al., 2016; Laporte et al., 2017), a phenotype linked with an alteration of CIE in developing neurons (Laporte et al., 2017). In the adult brain, Alix is ubiquitously expressed but concentrates at hippocampal presynaptic terminals during epileptic seizures (Hemming et al., 2004). Alix is a cytosolic protein first identified through its calcium-dependent binding to the penta-EF-hand protein ALG-2 (apoptosis-linked gene 2) which helps opening and activating the protein (Chatellard-Causse et al., 2002; Missotten et al., 1999; Sadoul, 2006). Its activation leads to interaction with membranes as reported in the case of plasma membrane wounds, where ALG-2 binds to inflowing calcium and helps recruiting Alix to the membrane where the latter organizes repair (Scheffer et al., 2014). Alix also interacts with lipids (Bissig et al., 2013) and with several membrane modifying proteins among which endophilin-A proteins (A1, A2 and A3) (Chatellard-Causse et al., 2002). These

cytoplasmic proteins that contain an N-BAR (Bin/Amphiphysin/Rvs) domain capable of sensing

and generating membrane curvature (Kjaerulff et al., 2011), are major actors of CME at

synapses (Gad et al., 2000; Ringstad et al., 1999). They also drive CIE in fibroblasts (Boucrot et

al., 2014; Renard et al., 2014) and were shown to control the fast mode of CIE at ribbon

synapses (Llobet et al., 2011) as well as in hippocampal neurons (Kononenko et al., 2014;

Watanabe et al., 2018).

The role of Alix in bulk endocytosis in fibroblasts, its capacity to interact with endophilin-A

and to be recruited by calcium at membranes, together with its increased concentration at

hippocampal synapses during kainate-induced seizures, brought us to test its possible function

in ADBE.

Using cultured wildtype (wt) neurons, we now bring evidence that sustained synaptic

activity leads to calcium-dependent recruitment of ALG-2 at synapses. ALG-2 in turn interacts

with Alix, which binds and concentrates endophilin-A. This protein complex is necessary for

ADBE, which is selectively impaired in Alix ko neurons. We also report that synapse

morphology and function are both altered in Alix ko brains. This finding correlates with the

modifications in kinetics of synaptic recovery following prolonged stimulation that we

detected in hippocampal slices of mice in which Alix was selectively deleted in forebrain

neurons. In these mice, the number of seizures during status epilepticus induced by

intracortical kainate injections was reduced as well as the propagation of epileptiform activity

to the contralateral side of injection. Thus, our results show that some molecular mechanisms

involved in ADBE may also be involved in certain aspects of synaptic plasticity such as short-

term synaptic depression and recurrence of epileptic seizures.

Material and methods

<u>Plasmids</u>

Endophilin A2-mCherry was obtained by subcloning (In-Fusion Cloning kit, Clontech) endophilin A2 cDNA into a pmCherry-N1 vector (Clontech). GFP-ALG-2 was obtained by performing a reverse mutagenesis (Quick change II site directed mutagenesis kit, Stratagene) on a GFP-hALG-2 Y180A (a generous gift from Masatoshi Maki) to acquire GFP-hALG-2wt. hALG-2 E47A-E114A cDNA was kindly provided by Masatoshi Maki (Shibata et al., 2004) and was subcloned into a pEGFP-C1 vector (Clontech) to obtain GFP-hALG-2 E47A-E114A (GFP-ALG-2ΔCa).

All constructs containing Alix cDNA (wt or mutants) were obtained by subcloning the relevant cDNAs from pCI vectors harbouring Alix cDNA or its mutants. Alix I212D (AlixΔCHMP4B) and AlixΔPGY (AlixΔALG-2) cDNAs in pCI were generated by mutagenesis (Quick change II site directed mutagenesis kit, Stratagene) and Alix R757E (AlixΔendo) by in-fusion cloning, using the oligos given below.

mCherry-2Xflag-mAlix wt (mCherry-Alix) was obtained by subcloning 2xflag-mAlix wt cDNA into a pmCherry-C1 vector (Clontech). Alix-YFP was obtained by subcloning wild type Alix cDNA into a pEYFP-N1 vector (Clontech). GFP-flag-Alix (GFP-Alix) was described in (Mercier et al., 2016). GFP-Alix and its mutant forms (GFP-Alix R757E, GFP-AlixΔPGY) were obtained by subcloning the various cDNAs into a pEGFP-C1 vector (Clontech). DNA constructs used for the rescue experiments were prepared in two steps. First, IRES2-GFP cDNA was subcloned into pSIN lentiviral vector (kindly provided by F. Saudou) by using pIRES2-GFP (Clontech) as a template. Then the various cDNAs were subcloned into pSIN-IRES2-GFP.

Oligos used to generate mutants:

Alix I212D

sense: 5'-AAGATGAAAG ATGCCGACAT AGCTAAGCTG-3'

antisense: 5'-CAGCTTAGCT ATGTCGGCAT CTTTCATCTT -3'

Alix R757E

sense: 5'-CAGCCGAGCC TCCACCTCCT GTGCTTCCTG -3'

antisense: 5'-GAGGCTCGGC TGGAGGCTGG GGCTTAGCAG-3'

AlixΔPGY

sense: 5'-GCCACAGGCT CAGGGATGCC AAATGCCCAT GC-3'

145   antisense: 5'-GCATGGGCAT TTGGCATCCC TGAGCCTGTG GC -3'.

146 Antibodies

Antibody	Supplier (reference)	Species (type)	Dilution
Anti-Actin	Millipore (MAB1501R)	Mouse (monoclonal)	1/10000
Anti-Alix	Covalab (ab0204)	Rabbit (polyclonal)	1/10000
Anti GFAP	Abcam	Goat (polyclonal)	1/1000
Anti Iba1	Wako	Rabbit (polyclonal)	1/500
Anti-phospho-p44/42 MAPK (T202/Y204)	Cell signaling (9106)	Mouse (monoclonal)	1/1000
Anti-PSD95	NeuroMab (73-028)	Mouse (monoclonal)	1/2000
Anti-PSD95	Millipore (MAB1598)	Mouse (monoclonal)	1/500
Anti-Synapsin-1	Millipore (AB1543P)	Rabbit (polyclonal)	1/1000
Anti-Synaptophysin	Merck Millipore (MAB5258)	Mouse (monoclonal)	1/5000
Anti-Flag	Sigma-Aldrich (F3165)	Mouse (monoclonal)	1/1000
Anti-Flag	Sigma-Aldrich (F7425)	Rabbit (polyclonal)	1/1000
Anti-Mouse HRP	Jackson ImmunoResearch (115-035-166)	Goat (polyclonal)	1/5000
Anti-Rabbit HRP	Jackson ImmunoResearch (115-035-044)	Goat (polyclonal)	1/5000
Anti-Mouse Alexa Fluor 488	Invitrogen (A-11029)	Goat (polyclonal)	1/1000
Anti-Mouse Alexa Fluor 594	Invitrogen (A-11032)	Goat (polyclonal)	1/1000
Anti-Mouse Cy5	Jackson ImmunoResearch (115-175-146)	Goat (polyclonal)	1/500
Anti-Rabbit Alexa Fluor 488	Invitrogen (A-11034)	Goat (polyclonal)	1/1000
Anti-Rabbit Alexa Fluor 594	Invitrogen (A-11037)	Goat (polyclonal)	1/1000
Anti-Rabbit Cy5	Jackson ImmunoResearch (111-175-144)	Goat (polyclonal)	1/1000

Animals

Animals were handled and killed in conformity with European law and internal

regulations of INSERM. Pregnant Oncins France souche A (OFA; a substrain of Sprague Dawley)

rats (Charles River) were used for rat neuronal cultures. Alix ko C57BL/6 mouse pups (Laporte

et al., 2017; Mercier et al., 2016) and their control littermates referred to thereafter as Alix

wt, were also used for primary neuronal culture. Transgenic mice were held at the animal

facility of the Grenoble Institute for Neurosciences and fed *ad libitum*. All animals were held

at a twelve-hour light/dark cycle. One- to two-month-old Alix wt and Alix ko mice were used

for electrophysiological extracellular recordings, histochemistry and electron microscopy

studies.

Mice were anesthetized by intraperitoneal injection of 0.1 ml sodium pentobarbital (5.6% w/v;

CEVA Santé Animale) and treated as described in the corresponding sub-headings of the

material and methods section.

162   Emx1^{IREScre}/Alix$^{fl/fl}$ and Emx1^{IREScre} control mice, thereafter referred to control and Alix

cko, were used for the kainate injection epilepsy model and the electrophysiological. 8-11

weeks old mice were anesthetized under a mixture of 2 % isoflurane, 47.5 % O_2 and 47.5 %

N_2O. Transgenic mice were held at the animal facility of the CIPMM and fed *ad libitum*. All

animals were held at a twelve-hour light/dark cycle. This study was carried out in strict

accordance with the European and German guidelines for the welfare of experimental

animals. Animal experiments were approved by the Saarland state's "Landesamt für

Gesundheit und Verbraucherschutz" animal licence number 36/2016.

<u>Golgi staining</u>

172   2-month-old anesthetized mice were dislocated prior to brain dissection and 100 µm

thick coronal brain sections were cut on a vibratome in the hippocampal region. The dendritic

spines of hippocampal neurons from the CA1 *stratum radiatum* were visualized by the Golgi

impregnation technique. For this, we used the FD Rapid GolgiStain kit (FD NeuroTechnologies).

Brain sections were immersed in equal volumes of solutions A and B for 7 d and impregnated

with solution C for 48 h at 4°C. Then, the sections were washed twice in double-distilled water

and incubated for 10 min in a mixture solution D and solution E in double-distilled

water(1:1:2). Sections were washed twice, dehydrated with increasing concentrations of

ethanol, and mounted with epoxy resin (Fluka). Stacks of bright-field images with 0.3 µm

spacing were acquired with a Zeiss Axioskop 50 microscope with 63x oil objective (NA 1.4;

Plan-Apochromat) coupled to a CCD camera (CoolSnap ES; Roper Scientific) operated by

Metaview software (Molecular Devices). Images were analysed with ImageJ. The number of

dendritic spines (>1 μm protrusion) along portion of dentrite of 100 μm was counted with

ImageJ.

<u>Transmission electron microscopy of the CA1 hippocampus</u>

2-month-old mice mice were anesthetized and intracardially perfused with phosphate-

buffered 0.9% NaCl (PBS), followed by 0.1 M phosphate buffered 4% paraformaldehyde, pH

7.4, supplemented with 0.05% glutaraldehyde (Sigma). The brains were carefully removed,

postfixed for 4 h in the same fixative and 60 μm sections were cut with a vibratome. After

several washes in PBS, the sections were postfixed in 1% glutaraldehyde in the same buffer

for 10 min and processed for EM. This included treatment with osmium tetroxide (1% in 0.1

M PB), block staining with uranyl acetate, dehydration through a graded series of ethanol, and

flat embedding on glass slides in Durcupan (Fluka) resin. Regions of interest were cut at 70 nm

on an ultramicrotome (Reichert Ultracut E; Leica) and collected on one-slot copper grids.

Staining was performed on drops of 1% aqueous uranyl acetate, followed by Reynolds's lead

citrate. EM images were acquired in a JEOL-1200 electron microscope with a digital camera

(Veleta, SIS; Olympus) and analysed with ImageJ. Twenty images per animal from 3 animals

per genotype were used for quantification. The number of synapses per $μm^2$ was calculated.

A synapse was considered if it met 3 criteria: a presynaptic bouton filled with at least 10

synaptic vesicles (1) juxtaposed to the head of a dendritic spine with a clearly visible PSD (2)

and the presence of the neck in the section (3). Number of synaptic vesicles and areas of

presynaptic boutons were quantified in each synapse using the free-shape tool and the cell

counter plugins of ImageJ. We used the straight tool of ImageJ to measure the lengths of PSDs,

and head and neck diameters. Note that the head diameter was taken parallel to the PSD and

the neck diameter was perpendicular to the neck membranes.

<u>Electrophysiological recordings in CA1</u>

Field recordings

Alix wt and ko brain slices were prepared from 2-month-old C57BL/6 wt and Alix ko mice. The

brains were removed quickly and 350 μm-thick sagittal slices containing both cortex and

hippocampus were cut in ice-cold sucrose solution (2.5 mM KCl, 1.25 mM NaH_2PO_4, 10 mM

$MgSO_4$, 0.5 mM $CaCl_2$, 26 mM NaHCO3, 234 mM sucrose, 11 mM glucose, saturated with 95%

O_2 and 5% CO_2) with a Leica VT1200 blade microtome (Leica Microsystemes, Nanterre,

France). After cutting, hippocampi were extracted from the slice and transferred to

oxygenated Artificial Cerebro-Spinal Fluid (ACSF: 119 mM NaCl, 2.5 mM KCl, 1.25 mM

NaH_2PO_4, 1.3 mM $MgSO_4$, 2.5 mM $CaCl_2$, 26 mM $NaHCO_3$, 11 mM glucose) at 37± 1°C for 30

min and then kept at room temperature for at least 1 h before recordings. Each slice was

individually transferred to a submersion-type recording chamber and continuously superfused

(2 ml/min) with oxygenated ACSF. Extracellular recordings were obtained at 28°C from the

apical dendritic layers of the hippocampal CA1 area, using glass micropipettes filled with ACSF.

Field excitatory postsynaptic potentials (fEPSPs) were evoked by the electrical stimulation of

Schaffer collaterals afferent to CA1. The magnitude of the fEPSPs was determined by

measuring their slope. Signals were acquired using a double EPC 10 Amplifier (HEKA Elektronik

Dr. Schulze GmbH, Germany) and analysed with Patchmaster software (HEKA Elektronik Dr.

Schulze GmbH, Germany). For the induction of long-term potentiation, test stimuli were

delivered once every 15 s. Stimulus intensities were adjusted to produce 40-50% of the

maximal response. A stable baseline was recorded for at least 15 min. LTP was induced by high

frequency stimulation (4 trains delivered at 100 Hz with 5 min between each train). Average

value of fEPSP slope was expressed as a percentage of the baseline response.

Patch-clamp recordings

Alix$^{fl/fl}$ mice (Laporte et al., 2017) with heterozygous cre expression (Emx1^{IREScre}

(Emx$^{tm1(cre)Krj}$, MGI: 2684610)(Gorski et al., 2002) as well as control mice (Alix$^{fl/fl}$ x Emx^{wt}) were

decapitated under anaesthesia and brains were removed from the skull and immersed in an

ice-cold, oxygenated (5% CO_2/95% O_2, pH 7.4) slice preparation solution containing (in mM)

87 NaCl, 3 KCl, 25 $NaHCO_3$, 1.25 NaH_2PO_4, 3 $MgCl_2$, 0.5 $CaCl_2$, 75 sucrose and 25 glucose.

Coronal slices of 300 µm thickness were prepared with a vibratome (Leica VT 1200S, Nussloch,

Germany) and transferred to a nylon basket slice holder for incubation in ACSF containing (in

mM) 126 NaCl, 3 KCl, 25 $NaHCO_3$, 15 glucose, 1.2 NaH_2PO_4, 2 $CaCl_2$, and 2 $MgCl_2$ at 32 °C for

30min. Subsequently, slices were removed from the water bath and kept at room temperature

with continuous oxygenation prior to use.

Slices were transferred to the recording chamber continuously perfused with

oxygenated ACSF containing (in mM) 1 $MgCl_2$ and 2.5 $CaCl_2$ (2-5 mL/min). During recordings,

50 µM strychnine and 50 µM picrotoxin were added to block inhibitory synaptic transmission.

Pyramidal neurons were identified morphologically using the recording microscope (Axioskop

2 FS mot, Zeiss, Jena, Germany) with a 40x water immersion objective. Images were detected

with a QuantEM 512SC camera (Photometrics, Tucson, USA). Whole-cell membrane currents

were recorded by an EPC 10 USB amplifier (HEKA, Lambrecht, Germany), low pass filtered at

3 kHz and data acquisition was controlled by Patchmaster software (HEKA). The resistance of

patch pipettes (7–9 ΩM) were fabricated from borosilicate capillaries (OD: 1.5 mm; Sutter,

USA) using a Micropipette Puller (Model P-97, Sutter Instrument Co., CA). Patch pipettes were

filled with a solution containing (in mM) 125 cesium gluconate, 20 tetraethylammonium (TEA),

2 $MgCl_2$, 0.5 $CaCl_2$, 1 EGTA, 10 HEPES and 5 Na_2ATP (pH 7.2). The holding potential in voltage-

clamp mode was at -70 mV. Spontaneous excitatory postsynaptic currents (sEPSC) were

recorded under holding potential for 40 seconds in 5 minutes later after whole-cell

configuration. Here only <=-10 pA sEPSC was analyzed. Excitatory postsynaptic currents

(EPSCs) of pyramidal neurons in the hippocampal CA1 were evoked by stimulating Schaffer

collaterals of CA3 neurons with a concentric bipolar microelectrode (MicroProbes, USA).

Stimulus duration was 200 μs. To estimate short-term plasticity, 25 trains of stimuli were

applied to induce synaptic depression, and the depressing trains were repeated at 10 Hz with

an interval of 20 s to allow for recovery from synaptic depression. To investigate recovery from

depression, stimulation with gradually increased interval was applied. The stimulation

threshold was applied from 30 μA to 80 μA. All the experiments were conducted at room

temperature (RT) (22 – 24 ℃).

Data generated by Patch Master were loaded into Matlab (Mathworks, MA, USA) with

a module adapted from sigTOOL (Lidierth, 2009). EPSC traces from the same cells were

manually checked and pooled. The average of EPSC traces from each cell was used to analysis.

The readily releasable pool (RRP) size was estimated by using train-extrapolation method

(Neher, 2015; Thanawala and Regehr, 2016).The cumulative sum of the peak EPSC amplitudes

was plotted against the stimulation numbers. The last 10 data points were linearly fitted. The

RRP size was calculated by back-extrapolating the fitted line to the y-axis (stimulation ID = 0).

The release probability was calculated by p = Amplitude[1] / RRP_{train}. The fitted slop represents

the replenishment rate (Körber et al., 2015). The recovery was estimated with the decay

constant (τ) as $N_t = N_0 * exp\left(-\frac{t}{\tau}\right)$.

Data analysis was performed using routines that were custom written in Matlab.

EEG telemetry and unilateral intracortical kainate injection

We took advantage of the unilateral intracortical kainate injection model for human temporal

lobe epilepsy. Telemetric EEG transmitter implantation, kainate injection and data analysis

was adapted from Bedner and colleagues (Bedner et al., 2015).

Controls and alix cko were implanted with telemetric EEG transmitters (DSI PhysioTel®

ETA-F10, Harvard Biosciences, Inc. Holliston, Massachusetts, USA). The animals were placed in

a stereotaxic frame (Robot stereotaxic, Neurostar, Tübingen, Germany) for implantation of

depth electrodes at 3.4 mm posterior to bregma and bilaterally 1.6 mm from the sagittal

suture. After post-surgical care and recovery, mice were again placed in the stereotaxic frame

and injected with 70 nl of a 20 mM solution of kainate (Tocris, Wiesbaden-Nordenstadt,

Germany) in 0.9% NaCl, above the right dorsal hippocampus (1.9 mm posterior to bregma,

1.5 mm from sagittal suture and 1.3 mm from skull surface). Kainate was injected at a rate of

70 nl/ min with a 10 µl Nanofil syringe (34 GA blunt needle, World Precision Instruments,

Sarasota, FL, USA). The syringe was kept in place for 2 min after the injection was completed

to avoid liquid reflux.

Cages were placed on individual radio receiving plates (DSI PhysioTel® RPC-1, Data

Sciences International, St. Paul, USA), which record EEG signals and sent them, together with

the video recording (MediaRecorder Software, Noldus Information Technology, Wageningen,

Netherlands), to an input exchange matrix (DSI PhysioTel® Matrix 2.0 (MX2), Ponemah

software, DSI, Data Sciences International, St. Paul, USA). The animals were monitored for at

least 20 h post kainate injection. In our model the mortality rate associated to *status*

epilepticus is less than 5% in more than 50 mice with different genetic backgrounds over the

last 12 months.

EEG traces were analysed with the Neuroscore software (Version 3.3.1., Data Sciences

International, St. Paul, USA). Electrographic seizures were detected with the spike detection

protocol. Subsequently, an additional manual screen was employed to remove artifacts that

were eventually picked up. Seizures were characterized by high frequency spiking and ceased

with a postictal depression (flattening of EEG). Seizure detection was complemented by

synchronized video monitoring. Electrographic seizures were associated with behavioural

analogues of Racine stages II-V (Racine, 1972). The total duration of *status epilepticus* was

defined from the first electrographic seizure to the first seizure free period lasting 1 h.

For immunohistemistry analysis, 40 µm free-floating vibratome sections were
incubated for one hour in blocking buffer (5 % HS, 0.3 % Triton X in 1x PBS) at room
temperature. Sections were incubated with primary antibodies (see antibodies section)
diluted in blocking solution, overnight at 4°C. Secondary antibodies and DAPI (0.025 µg/ml,
Sigma) were diluted in blocking buffer and incubated for 2 h at room temperature.
315
Cell culture and transfection
Cortical and hippocampal neurons from rat E18 embryos and P0 Alix wt and ko mice
were prepared as previously described (Faure et al., 2006). Briefly, cortices and hippocampi
were dissected from E18 rat embryos or P0 mouse pups, treated with trypsin, and
mechanically dissociated. Dissociated cells were seeded at a density of $5 \times 10^4/cm^2$ in 100 mm
dishes for cortical neurons and $1.5 \times 10^4/cm^2$ onto acid-washed coverslips (either 14 mm
diameter or 25 mm; Marienfeld) in either 4-well plates or P35 dishes (ThermoScientific)
precoated for 4 h with 50 µg/ml poly-D-lysine (Sigma). Neurons were plated in DMEM
containing 10 unit/mL penicillin, 10 µg/mL streptomycin,0.5 mM L-glutamine and 10% of
inactivated horse serum (Invitrogen) Neurons were maintained in water-saturated 95% air/5%
CO_2 at 37°C. The seeding medium was replaced after 20 h with serum-free neuronal culture
medium (Neurobasal medium containing 2% B27 supplement, 10 unit/mL penicillin, 10 µg/mL
streptomycin, and 0.5 mM L-glutamine; Invitrogen). Neurons were transfected at 10 DIV as
previously described (Chassefeyre et al., 2015). Briefly, for each P35 dish, 2 µg plasmid DNA,
250 mM $CaCl_2$ were mixed with an equal volume of 2x BES-buffered saline and left to
precipitate for 20 min at room temperature. Neurons were placed in transfection medium
(Minimum Essential Medium containing 0.22% $NaHCO_3$, 20 mM D-glucose and 0.5 mM L-
glutamine) supplemented with 2% B27, before the DNA precipitate was added. They were
then incubated for 1.5 h at 37°C and 5% CO_2. Neurons were then washed by being placed in
transfection medium (pre-warmed to 37°C in 10% CO_2) for 20 min at 37°C and 5% CO_2. Finally,
they were transferred back into their conditioned medium.
Primary cultures of cerebellar granule neurons (CGN) were prepared from 6-day-old
C57BL/6 Alix wt and Alix ko pups as described previously (Trioulier et al., 2004), with some
modifications. The cerebella were removed, cleared of their meninges, and cut into 1-mm
pieces. They were then incubated at 37°C for 10 min in 0.25% trypsin-EDTA and DNAse (1500
U/mL). Trypsin was inactivated and cells were dissociated in culture medium (DMEM

containing 10% fetal bovine serum, 2 mM L-glutamine, 25 mM KCl, 10 mM HEPES and 10

unit/ml penicillin, 10 μg/ml streptomycin). After filtration on 70 μm cell strainers, neurons

were plated at 5.10^5 cell/cm^2 onto poly-D-lysine (10 μg/ml, Sigma) precoated coverslips.

Cytosine-β-D-arabinoside (10 μM, Sigma) was added after 1 day in vitro (DIV), to prevent the

growth of non-neuronal cells, until 8 DIV when neurons were used for activity dependent bulk

endocytosis experiment (see below).

<u>Multi-electrode array</u>

Dissociated hippocampal neurons were resuspended in neuronal medium and plated

at a 10^6 cells/cm^2 on poly-L-lysine-coated multi-electrode arrays comprising 59 extracellular

recording electrodes and one reference electrode (MEA-60 Multichannel Systems, MCS).

Electrodes were 30 μm in diameter and separated by a distance of 200 μm. MEA-60 plates

were connected to a 60-channel data acquisition system (USB-MEA64) and associated

amplifier (MEA1060-Inv-BC) powered by a PS40W power supply system (MCS, Germany). The

recording system was then placed in a humidified incubation chamber at 37°C and 5% CO_2.

Neurons were left for 5 min to equilibrate before recording. Basal spontaneous activity was

recorded for 3 min prior addition of drugs. Bicuculline (100 μM) and 4-aminopyridine (5 mM)

were then added and neuronal activity recorded for another 10 min. Signals were recorded

with a 1100 gain, sampled at 20 kHz and analyzed with MC Rack software (MCS, Germany).

Raw signals were first filtered with a Butterworth band pass filter (2nd order) between 200

and 800 Hz to remove electrical noise and low frequency signals. Spike detection analysis was

performed on filtered signals using a threshold of 6 standard deviations to mark out action

potentials. Burst events were identified as a minimum of 5 consecutive spikes with an

interspike interval lower than 100 ms. The minimum interval time between two bursts was

fixed at 1000 ms. Representative traces were exported using custom-made Matlab functions

(Matlab 2014b).

<u>Live imaging of synaptophysin-pHluorin upon electrical stimulation</u>

Hippocampal neurons from Alix wt and ko mice were transfected with synaptophysin-

pHluorin (Syp-pH) at 6 DIV. The recycling of synaptic vesicles was imaged in a buffer solution

containing 120 mM NaCl, 5 mM KCl, 2 mM $CaCl_2$, 2 mM $MgCl_2$, 5 mM glucose, 10 mM HEPES

adjusted to pH 7.4 and 270 mOsm/l. Experiments were carried out at 34°C. Neurons were

stimulated by electric field stimulation (platinum electrodes, 10 mm spacing, 1 ms pulses of

50 mA and alternating polarity at 5-40 Hz) applied by a constant current stimulus isolator (SIU-

102, Warner Instruments). The presence of 10 µM 6-cyano-7-nitroquinoxaline-2,3-dione

(CNQX) and 50 µM D,L-2-amino-5-phosphonovaleric acid (AP5) prevented recurrent activity.

Experiments were performed on an inverted microscope (IX83, Olympus) equipped with an

Apochromat N oil 100× objective (NA 1.49). Images were acquired with an electron multiplying

charge coupled device camera (QuantEM:512SC; Roper Scientific) controlled by MetaVue7.1

(Roper Scientific). Samples were illuminated by a 473-nm laser (Cobolt). Emitted fluorescence

was detected after passing a 525/50 nm filter (Chroma Technology Corp.). Time-lapse images

were acquired at 1 or 2 Hz with integration times from 50 to 100 ms.

Image analysis was performed with custom macros in Igor Pro (Wavemetrics) using an

automated detection algorithm as described previously (Martineau et al., 2017). The image

from the time series showing maximum response during stimulation was subjected to an "à

trous" wavelet transformation. All identified masks and calculated time courses were visually

inspected for correspondence to individual functional boutons. The intensity values were

normalized to the ten frames before stimulation.

<u>Live Fluorescence Imaging of Alix and endophilin recruitment to synapses</u>

For live imaging of protein recruitment, cultured hippocampal neurons were co-

transfected with Alix-mCherry or endophilin-A2-mCherry together with Syp-pH expression

vectors 2 to 4 days prior to imaging. All live imaging experiments were performed at 37°C and

images were acquired using a spinning disk confocal microscope (AxioObserver Z1) with a 63x

oil objective (NA 1.46, Zeiss). Transfected neurons were placed in basal medium for 10 min

and then mounted in an imaging chamber (POC-R2 Cell cultivation system, Zeiss). After 2 min

in basal medium, a 5x bicuculline/4AP solution was added (final concentration, 100 µM and 5

mM, respectively) for stimulation which lasted 5 min and the chamber was perfused with

basal medium at 3 ml/min for washing for 5min. ROI were drawn on 'presynapses' defined by

spots of synaptophysin-pHluorin that increased during stimulation. Fluorescence values were

measured in these ROI and then normalised to the initial fluorescence values (fluorescence

values prior to stimulation) using ImageJ.

<u>Immunofluorescence</u>

Cultured hippocampal neurons were fixed for 20 min at room temperature in
phosphate-buffered 4% paraformaldehyde supplemented with 4% sucrose. After three
washes in PBS, cells were permeabilized and blocked in PBS containing 0.3% Triton X-100 and
3% BSA for 15 min at room temperature. Coverslips were incubated for 1–2 h at room
temperature with primary antibodies (see 'antibodies' section) diluted in the blocking
solution. After washing in PBS, cells were incubated for 1 h with secondary antibodies
conjugated to Alexa Fluor 488, Alexa Fluor 594, or Cyanine 5 (Cy5), diluted in the blocking
solution. Coverslips were rinsed and mounted in Mowiol. Images were acquired on a Leica SPE
microscope using a 40x dry objective (NA 0.75, Leica) or a 100x oil immersion objective (NA
1.4, Leica) at 488 nm, 532 nm or 635 nm.

Synapse density: Cultured hippocampal neurons were fixed and stained with
antibodies at 14-16 DIV as described above. Presynaptic boutons (synapsin-1-positive) and
postsynaptic terminals (PSD95-positive) were selected using the Spot Detector plugin in Icy
software (de Chaumont et al., 2012) (wavelet detection with size filtering between 0.4 μm and
2 μm in diameter) on max image projections. Synapses were defined as spots of colocalization
between the detected presynaptic and postsynaptic terminals that were within 3 μm of each
other. Synapses were counted using the Colocalization Studio in ICY software (Lagache et al.,
2018)

Protein recruitment assays: Cultured hippocampal neurons were transfected with
expression vectors 24~48 h prior to stimulation. The transfected neurons were incubated in
basal medium (150 mM NaCl, 5 mM KCl, 1.3 mM $CaCl_2$, 10 mM HEPES and 33 mM D-glucose
at pH 7.4) for 10 min prior to stimulation. Coverslips were either treated with basal medium
alone or bicuculline/4AP solution (basal medium supplemented with 100 μM bicuculline and
5 mM 4-aminopyridine) for 5 min at 37°C and 5% CO_2. Coverslips were fixed and stained with
antibodies against synapsin-1 and PSD95 (see 'antibodies' section). Images were acquired on
a Leica SPE microscope using a 100x oil immersion objective (NA 1.4, Leica). The acquired
images were analysed using ImageJ. Regions of interest (ROI) were drawn on areas of axons
which colocalize with anti-synapsin-1 spots, but not with anti-PSD95 spots to ensure
presynaptic measurement. Fluorescence intensities were then measured in these ROI and
normalised to fluorescence intensities measured on other regions of axons to give the relative
fluorescence intensity at presynaptic boutons.

 <u>Calcium increase imaging upon neuronal stimulation</u>

Hippocampal neurons cultured on 24-well plates were placed in basal medium and

incubated with 1 µM Fluo-4-AM for 1 h at 37°C. After washing, neurons were left in 250 µl

basal medium for 5 min. Plates were then transferred to a Pherastar automatic plate reader

(BMG Labtech, Germany) set to 37 °C, to record fluorescence intensity at 0.1 Hz during 10min,

for each well, and with fluorescence measurement settings as described on

http://www.bmglabtech.com/media/35216/1043854.pdf. Neurons were stimulated with 50

µl 6x bicuculline/4AP solution (final concentration, 100 µM and 5 mM, respectively) after 5min

with automatic injection.

<u>Dextran uptake upon neuronal stimulation</u>

The protocol for dextran uptake was adapted from (Clayton and Cousin, 2009b). 15-17

DIV hippocampal neurons were stimulated with bicuculline/4AP solution for 5 min, at 37°C

and 5% CO_2, in the presence of 10 kDa tetramethylrhodamine-dextran (50 µM). Coverslips

were immediately washed several times in washing solution (basal medium supplemented

with 0.2% BSA and warmed to 37°C) to remove excess dextran. For ADBE inhibition, neurons

were stimulated in the presence of 2 µM GSK3 inhibitor (CT99021, Tocris) and placed in fresh

basal medium containing 2 µM GSK3 inhibitor for 10 min at 37°C and 5% CO_2. Neurons were

then fixed as previously described and imaged with a Leica SPE microscope using a 40x dry

immersion objective (NA 0.75, Leica). Analysis was performed on ImageJ. The number of

fluorescent spots was counted in a defined field of view (130µm x 130 µm) in thresholding

analysis with a diameter limit between 300 nm and 2 µm (resolution limit for the microscope

and maximum size of a nerve terminal).

For rescue experiments, cultured hippocampal neurons were transfected 2 to 4 d prior

to the day of experiment with the following constructs :Alix wt, Alix I212D (AlixΔCHMP4B),

AlixΔPGY (AlixΔALG-2), Alix R757E (AlixΔendo). Dextran uptake assay was performed as

described above. Images were acquired on a Leica SPE microscope using a 40x oil immersion

objective (NA 1.25, Leica) at 488 nm and 532 mm excitation. The analysis was performed on

ICY software. ROI were generated closely around axons by using 'thresholder'. Then dextran

spots within these ROI were counted by using 'spot detector' with size limit between 300 nm

and 2 µm. The length of axon per field of view was estimated by manually drawing 'Polyline

type ROI' over the axon images. The number of dextran spots per µm of axon was calculated

and expressed as ratio to control values.

<u>EM examination of ADBE in culture neurons</u>

Analysis of ADBE from cerebellar granule neurons was performed as described

previously with some modifications (Cheung et al., 2010). 8 DIV CGN were incubated in

hyperpolarizing medium (170 mM NaCl, 3.5 mM KCl, 0.4 mM KH_2PO_4, 20 mM TES, 5 mM

$NaHCO_3$, 5 mM D-glucose, 1.2 mM Na_2SO_4, 1.2 mM $MgCl_2$, 1.3 mM $CaCl_2$, pH7.4) for 10 min

prior to stimulation. Neurons were then incubated for 2 min with 10 mg/ml HRP (Sigma P8250)

in either the hyperpolarizing medium or a high-potassic solution containing 50 mM KCl (123.5

mM NaCl, 50 mM KCl, 0.4 mM KH_2PO_4, 20 mM TES, 5 mM $NaHCO_3$, 5 mM D-glucose, 1.2 mM

Na_2SO_4, 1.2 mM $MgCl_2$, 1.3 mM $CaCl_2$, pH7.4) before rapid washing in PBS and fixation in PBS-

glutaraldehyde 2% for 30 min. After three washes in Tris buffer 100 mM, endocytosed HRP

was revealed by incubation in Tris 100 mM containing 0.1% diaminobenzidine and 0.2% H_2O_2.

The cultures were then post-fixed in 1% osmium tetroxide, dehydrated and embedded in

Epon. Synapses were photographed with a JEOL-1200 electron microscope. Quantification

was performed as follows: HRP-positive structures were quantified per synapse and classified

as synaptic vesicles when the diameter was less than 100 nm and as bulk endosome when the

diameter was more than 100 nm.

For focused ion beam scanning electron microscopy (FIB-SEM), the block was mounted

on a pin, coated with gold, and inserted into the chamber of a HELIOS 660 Nanolab DualBeam

SEM/FIB microscope (FEI Co., Eindhoven, The Netherlands). Region of interest (ROI) was

prepared using focused ion beam (FIB) and ROI set to be approximatively 15 µm wide. During

the acquisition process, the image acquisition parameters of the electron beam were 2 kV and

0.4 nA and the thickness of the FIB slice between each image acquisition was 10 nm. The

segmentation of the synapse and ABDE was done with Amira software (Thermo Fisher

Scientist) and the movie obtained using Imaris (Oxford Instruments).

<u>Synaptosomal preparation from cortical neurons</u>

Synaptosome-enriched membranes from 15 DIV cortical neurons were prepared as described

previously with some modifications (Frandemiche et al., 2014). Briefly, cultured neurons were

stimulated for 15 min with a mixture of bicuculline/4AP (50 µM/2.5 mM). After a wash in

HBSS, neurons were homogenized by passing 15-20 times through 0.25G needle in cold buffer

containing 0.32 M sucrose, 10 mM HEPES, 15mM NaF, 15mM β-glycerophosphate and

protease inhibitors (Roche), pH 7.4. Samples were maintained at 4°C during all steps of the

experiment. Homogenates were cleared at 1000g for 10 min to remove nuclei and large

debris. The resulting supernatants were spun down at 12,000 g for 20 min to obtain a crude

membrane fraction and washed twice in HEPES buffer 4 mM containing 1 mM EDTA, 15 mM

NaF, 15 mM β-glycerophosphate and protease inhibitors (Roche), pH 7.4. The resulting pellet

was solubilized in 0.5% Triton X-100, 20 mM HEPES, 100 mM NaCl, 15mM NaF, 15 mM β-

glycerophosphate, pH 7.2, containing protease inhibitors (Roche) for 20 min at 4°C with mild

agitation and analysed by Western blot.

<u>Western blot</u>

Cells lysates were resuspended in Laemmli buffer and resolved by SDS-PAGE in 10%

polyacrylamide gels. Proteins were electro-transferred onto PVDF membranes that were then

blocked for 30 min in TBS containing 0.1% Tween 20 and 5% dry milk and incubated for 1 h to

overnight with primary antibodies diluted in the blocking solution. After washes in TBS–

Tween, the membranes were further incubated for 1 h with secondary antibodies coupled to

HRP, washed as before and incubated with luminescence-generating HRP substrate. Bound

antibodies were revealed by luminography on film.

<u>Statistical analysis</u>

The comparison of two groups was performed using two-sided Student's t-test or its non-

parametric correspondent, the Mann-Whitney test, if normality was rejected (Shapiro-Wilks

test). The comparisons of more than two groups were made using one or two way ANOVAs

followed by post-hoc tests (Holm Sidak's or Tukey's HSD) to identify all the significant group

differences. N indicates independent biological replicates. The graphs with error bars indicate

1 SEM (+/-) except for figure 5 where we used box plots showing distribution of medians

(whiskers = min and max values), figures 7 and supplementary figure 6 where median and

interquartile range (IQR) were represented. The significance level is denoted as usual

(*$p<0.05$, **$p<0.01$, ***$p<0.001$). All the statistical analyses were performed using Prism7

(Graphpad version 7.0a, April 2, 2016).

Results

Alix is recruited to synapses upon synaptic activation.

We used dissociated cortical neuron cultures to decipher the role of Alix at synapses. Western blot analysis during *in vitro* differentiation revealed that Alix expression strongly increases during synaptogenesis as indicated by the parallel rise in postsynaptic density protein 95 (PSD95) expression (**Figure 1A**). Moreover, synaptosome-enriched membranes prepared from mature cortical neurons contained Alix (**Figure 1B**). This synaptic pool increased when neuron cultures were incubated 15 min with the GABA$_A$ receptor antagonist bicuculline, together with a weak potassium channel blocker 4-aminopyridine (4AP)(**Figure 1B**), a treatment known to increase the frequency of action potential bursts (**Supplementary figure 3C**) and thereby induce sustained intracellular calcium elevation (**Supplementary figure 1A**) (Hardingham et al., 2001; Hardingham et al., 2002). Therefore, this observation suggests that Alix tends to concentrate at synapses undergoing prolonged stimulations. To further confirm this, we used live imaging to follow mCherry-Alix relocalization to synapses in mature hippocampal neurons. While the fluorescent signal was homogeneously distributed throughout the entire neuronal cytoplasm in resting conditions (**Figure 1C**, t= 0 min), it almost doubled in discrete spots within neuronal processes during synaptic activation (**Figure 1C**, t=2m20s arrowheads). These spots correspond to active presynaptic boutons as they were labelled with Synaptophysin-pHluorin (Syp-pH), which becomes fluorescent upon exocytosis at synapses (Granseth et al., 2006) (arrowheads **Figure 1C**; **Movie 1**). Alix recruitment to these sites is concomitant with the activation of glutamatergic synapses, being detectable a few seconds after addition of bicuculline/4AP to the culture medium and decreasing soon after washing the cells (**Figure 1 C, D**). Alix relocalization was also seen with Alix-YFP which concentrated during bicuculline/4AP treatment at sites confirmed to be at axonal boutons by synapsin-1 immunolabelling (**Figure 1E, F**). Anti-PSD95 immunostaining showed that Alix-YFP positive spots were juxtaposed to, but did not overlap with the postsynaptic density marker (**Figure 1G**). Accordingly, measuring Alix-YFP fluorescence intensities in dendritic spines before and during stimulation showed no accumulation of Alix at the post-synaptic level (**Supplementary figure 1B, C**), further demonstrating that synaptic activation leads to Alix recruitment only at presynaptic parts.

Recruitment of Alix partners, ALG-2 and endophilin-A, at activated synapses

In non-neuronal cells, Alix recruitment to plasma membrane depends on calcium binding

to ALG-2, a penta-EF-hand containing protein, which thereafter interacts and activates Alix

(Scheffer et al., 2014). In neurons, action potential depolarization induces massive and

transient calcium accumulation in the bouton, which triggers fusion of SV with the plasma

membrane. Adding the intra-cellular calcium chelators BAPTA-AM or EGTA-AM completely

abolished Alix-YFP recruitment (**Supplementary figure 2**). Interestingly, GFP-ALG-2

concentrated at presynaptic boutons upon synaptic activation, in contrast to an ALG-2 point-

mutant unable to bind calcium (ALG-2ΔCa) (Shibata et al., 2004) (**Figure 2A, B**). The use of

neurons prepared from Alix ko mice (Laporte et al., 2017; Mercier et al., 2016) showed that

ALG-2 recruitment does not depend on Alix, as it concentrates at stimulated synapses of Alix

ko neurons at the same level than in wt neurons (**Figure 2B, C** left histograms). On the

contrary, Alix recruitment to active synapses is tightly dependent on its capacity to interact

with ALG-2, as a mutated version of the protein unable to interact with ALG-2 (AlixΔALG-2)

(Suzuki et al., 2008; Trioulier et al., 2004) does not accumulate at presynaptic boutons upon

stimulation (**Figure 2C**, right histogram).

Among demonstrated interactors of Alix, endophilins-A are main regulators of endocytosis

at synapses and impact the number of SVs (Milosevic et al., 2011; Schuske et al., 2003). We

therefore tested if endophilins-A could be recruited to active synapses, similarly to Alix and

ALG-2. Endophilin-A2-mCherry was mainly detected at presynaptic boutons and its

fluorescence increased during bicuculline/4AP treatment (**Figure 2D, E; Movie 2**). Remarkably,

no such increase could be seen in Alix ko neurons, whereas Alix deleted of its endophilin

binding domain (AlixΔendo) (Chatellard-Causse et al., 2002), was still able to be recruited to

synapses upon activation (**Figure 2F**). This strongly suggests that the increase in endophilin

concentration during sustained synaptic activation requires Alix expression.

<u>Alix is required for ADBE</u>

Altogether, our observations indicate that calcium increase at synapses undergoing

sustained activity allows the transient recruitment of ALG-2, Alix and endophilin-A

sequentially to presynaptic parts. Because of the demonstrated roles of Alix in clathrin-

independent endocytosis, we next tested if the turnover of presynaptic vesicles might be

affected by the lack of the protein by quantifying Syp-pH fluorescence during electrical

stimulation. Syp-pH fluorescence increases during stimulation witnessing exocytosis. It decays

thereafter reflecting SV retrieval and vesicle re-acidification (Soykan et al., 2017). As shown in

figures 3A and B, no striking difference in the increase of fluorescence was detected between

Alix ko and wt neurons stimulated at 5 or 40 Hz. However, the time constant of the exponential

decay was slightly decreased in Alix ko neurons stimulated at 40 Hz, suggesting a higher rate

of Syp-pH endocytosis occurring at this frequency (**Figure 3B**).

We next tested if the proteins might intervene in activity-dependent bulk endocytosis

(ADBE) at synapses. Cultured cerebellar granular neurons (CGN) have been extensively used

to study this process (Cheung et al., 2010). Here, neurons were depolarized with 50 mM KCl,

in the presence of Horse Radish Peroxidase (HRP) that is endocytosed and fills vesicles and

endosomes as they form. Examination of wt synapses by electron microscopy showed that

depolarization dramatically increased the number of HRP-positive vacuoles. We verified using

FIB SEM and 3D reconstruction that these vacuoles had undergone fission from the plasma

membrane and could therefore be identified as bulk endosomes (**Figure 3C** yellow

arrowheads; **Figure 3D, E**). Other HRP-positive vesicles having the size of neurotransmitter

vesicles were also more numerous in depolarized synapses (**Figure 3C** blue arrowheads; **Figure**

3D, E). In Alix ko synapses, the number of depolarization-induced bulk endosomes was

strongly reduced (**Figure 3E, F**). Consistent with the increased endocytic rate detected by Syp-

pH in Alix ko neurons stimulated at 40 Hz (**Figure 3B**), the number of newly formed SV was

significantly increased, suggesting a mechanism compensating for ADBE deficiency (**Figure 3E,**

G).

Another way to assess for ADBE is by the use of fluorescent 10 kDa dextran, a fluid phase

cargo that accumulates inside bulk endosomes but fails to label vesicles formed by clathrin-

dependent mechanisms upon neuronal stimulation (Clayton and Cousin, 2009a). Hippocampal

neurons incorporated dextran when incubated in presence of bicuculline/4AP (**Figure 4A, B**).

A GSK3 inhibitor, known to block ADBE but no other modes of SV endocytosis (Clayton et al.,

2010), completely abolished the dextran labelling of wt hippocampal neurons

(**Supplementary figure 3A, B**). In contrast, Alix ko neurons failed to endocytose dextran

(**Figure 4A, B**) even though bicuculline/4AP stimulation increased calcium entry

(**Supplementary Figure 1A**) and neuronal activity (**Supplementary figure 3C-F**) in both Alix ko

and wt neurons. Noteworthy is that even though bicuculline/4AP increased activity in both

cases, the frequency of burst was significantly reduced in Alix ko neurons (**Supplementary**

figure 3C, D). As already shown (Morton et al., 2015), neurons also failed to endocytose

dextran when treated with calcium chelators BAPTA and EGTA (**Supplementary figure 3E**),

both of which blocked Alix recruitment to presynaptic boutons (**Supplementary figure 2**).

Finally, expression of GFP-Alix constructs in Alix ko neurons showed that the impairment in

ADBE observed in Alix ko cells is due solely to the absence of Alix, since Alix expression fully

restored the capacity of Alix ko neurons to endocytose 10 kDa dextran (**Figure 4 C, D**). Rescue

was also obtained with AlixΔ4B, a mutant unable to bind the ESCRT-III protein CHMP4B. This

was not the case with AlixΔALG-2 or AlixΔEndo which were not able to rescue dextran

endocytosis (**Figure 4D**). Noteworthy is that AlixΔALG-2 cannot be recruited to synapses but

both AlixΔendo and AlixΔ4B are. Thus, besides its requirement for ALG-2-binding to be

recruited at synapses, Alix needs to interact with endophilins but not with CHMP4B-ESCRT-III

in order to drive ADBE.

<u>Synaptic anomalies detected in Alix ko mice</u>

To study the effect of Alix deletion *in vivo*, we first compared brains of Alix ko (Laporte et

al., 2017) and wt mice and found that the density of synaptic contact as well as dendritic spines

in the CA1 hippocampal region of adult brains was not different (**Figure 5A, Supplementary**

Figure 4). This indicates that Alix is not required for synaptogenesis, a conclusion confirmed

in vitro since the number of synapses revealed by co-immunostaining with synapsin-1 and

PSD95 was similar between wt and Alix ko cultured hippocampal neurons (**Figure 5B**).

However, examination of the CA1 *stratum radiatum* of adult mice showed that Alix ko

synapses contained significantly fewer synaptic vesicles (SV) than wt (**Figure 5C, D**).

Furthermore, the size of postsynaptic densities, known to be strictly correlated with the

number of SV (Harris and Stevens, 1989), was similarly reduced in Alix ko synapses (**Figure 5F,**

G). Importantly, the surface of Alix ko synaptic boutons was also significantly increased

possibly suggesting plasma membrane accumulation in these synapses (**Figure 5C, E**). Finally,

at the postsynaptic level, we also noticed that the ratio between the diameter of the spine

head and that of the neck was changed in Alix ko animals, which can reflect defects in

maturation or plasticity of adult synapses (**Figure 5F, H**). Accordingly, field recordings in the

CA1 *stratum radiatum* of hippocampal slices revealed that long-term potentiation (LTP)

induced by tetanic stimulations was significantly impaired in Alix ko hippocampal slices

(**Supplementary Figure 5**).

<u>Voltage-clamp recordings uncovers anomalies at Alix ko synapses.</u>

In order to better characterize the effect of the lack of Alix on the physiology of synapses,

we next used Alix conditional-ko mice where Alix is deleted in neocortical and hippocampal

excitatory neurons (Gorski et al., 2002) (**Figure 6A**). We first recorded CA1 neurons of

hippocampal slices and found that the amplitude of spontaneous EPSCs was unaffected

whereas the frequency of events was significantly lower in Alix ko neurons (**Figure 6B, C**). This

was reminiscent of the reduced frequency of burst measured by MAE in dissociated

hippocampal cultures (**Supplementary figure 3C, D**). Stimulation of Schaffer collaterals

revealed that evoked EPSCs have a smaller amplitude in Alix ko brains than in wt animals

(**Figure 6D, E**). In order to estimate the amount of the readily releasable pool (RRP), we

performed repeated stimulations and integrated the EPSC amplitudes to give a cumulative

plot during 10 Hz trains. Linear regression fit to the last 10 data points was back extrapolated

to time 0 (dotted line) to estimate the cumulative EPSC before steady state depression. This

rough estimate of the RRP size (Schneggenburger et al., 1999) revealed that it is two times

smaller in Alix ko neurons (**Figure 6F**), in good agreement with the lower amounts of SVs

quantified in EM sections of synapses of the CA1 *stratum radiatum* (**Figure 5D**). Interestingly,

we could observe that the rate of recovery post-depression appears faster in Alix ko mice, a

possible mechanism of compensation for the reduced pool of synaptic vesicles (**Figure 6G, H,**

I).

Thus, these electrophysiological results show that synapses of Alix ko neurons which lack

ABDE have less SVs but increased rate of endocytosis, as already suggested in cultured

neurons by the increase of Syp-pH endocytosis and of newly formed SVs detected in Alix ko-

hippocampal and CB neurons, respectively.

<u>Alix conditional ko mice undergo less kainate induced acute seizures.</u>

We next examined the possible *in vivo* consequences of synaptic changes detected during

repeated stimulation of Alix ko neurons in hippocampal slices. Because of our earlier finding

of Alix increase at synapses of the rat hippocampus after kainaite injections (Hemming et al.,

2004) we tested how the lack of Alix might influence kainate induced epileptic seizures, a

mouse model for human temporal lobe epilepsy (Bedner et al., 2015). The unilateral

intracortical kainate injection induces the acute phase, *status epilepticus*, characterized by

high seizure frequency, during several hours. Seizure activity was determined by telemetric

EEG recording and synchronized video monitoring (**Figure 7A**). Here we used again the

conditional-ko mice where Alix is deleted in neocortical and hippocampal excitatory neurons

(**Figure 6A**). The total duration of *status epilepticus* (SE) was comparable in Alix ko and control

mice (**Figure 7B**). However, Alix cko mice experienced about 70 % less seizures than control

mice (**Figure 7C**). The average seizure duration was only minimally reduced in Alix ko

compared to control (**Figure 7D**). These results show that the absence of Alix selectively

reduces the number of high frequency events (single seizures), without affecting the total

duration of *status epilepticus*.

This kainate injection paradigm produces seizures originating from the hippocampus, which

can spread to other contralateral brain areas (Bedner et al., 2015). Microglial reactivity was

detected 24 h after kainate injection in ipsilateral and contralateral cortex and hippocampus

in both control and Alix cko animals (**Supplementary Figure 6A-E**). While the difference

between ipsi-and contralateral side was non-significant in control animals ($\approx$ 10%), Alix ko

animals displayed a significantly reduced contralateral microglial activation ($\approx$ 30%)

(**Supplementary Figure 6E**). No difference between Alix ko and control animals was detected

using GFAP immunolabelling, an expected result since astroglial activation peaks 4-7 days

after kainate injection (**Supplementary Figure 6F**). Thus, the reduced contralateral microglial

activation seen in Alix ko animals indicates a reduced propagation of the seizure activity and

corroborates the previous seizure quantification (**Figure 7C**).

In summary, Alix is specifically required for ADBE. Impairment in this process seems to

relate to impairments in synaptic function and plasticity in normal and pathological settings.

Discussion

To sustain neurotransmission and prevent expansion of the presynaptic plasma membrane, synaptic vesicle (SV) fusion is coupled to the endocytic recycling and regeneration of SV proteins and lipids. Vesicle components can be retrieved from the plasma membrane via clathrin scaffolds, or via clathrin-independent processes mediating fast and ultrafast endocytosis and, in the case of high frequency stimulation, bulk endocytosis (Gan and Watanabe, 2018). Using Alix ko cells, we recently discovered that Alix drives clathrin-independent-endocytosis (CIE) during ligand-induced endophilin-dependent endocytosis as well as bulk endocytosis (Laporte et al., 2017; Mercier et al., 2016). We had previously reported that Alix immunoreactivity in the rat hippocampus is strongly upregulated in synapses undergoing high frequency activation during kainate-induced epileptic seizures. This increase was presynaptic and only transient, as it was reversed soon after cessation of the seizures (Hemming et al., 2004). The aim of the present study was to decipher the role of Alix in endocytosis at synapses. We first used primary cultures of cortical and hippocampal neurons, which make networks undergoing spontaneous activities. Massive increases in spike frequency induced by 4AP and bicuculline resulted in elevated calcium entry (Hardingham et al., 2001) and triggered activity dependent bulk endocytosis (ADBE) together with a transient concentration of Alix at presynaptic parts. Quantifying bulk endocytosis using dextran uptake or electron microscopy observations showed that ADBE is strongly reduced in Alix ko neurons demonstrating that Alix is essential for ADBE. ADBE is triggered by high $[Ca^{2+}]$ in response to sustained activity (Paillart et al., 2003). Calcium and calmodulin-dependent phosphatase calcineurin were suggested to make a link between synaptic activity and formation of bulk endosomes through dynamin (Morton et al., 2015). Alix may represent another link as its recruitment and activity in ADBE seems dependent on calcium increase. Indeed, EGTA, which binds calcium at an approximately 100 times slower on-rate than BAPTA (Adler et al., 1991; Schneggenburger and Neher, 2005), and was already known to block ADBE without affecting synaptic release, was equally efficient as BAPTA in inhibiting Alix recruitment. Furthermore, Alix recruitment required its capacity to bind the calcium-binding protein ALG-2, which also concentrated at hyperactive synapses. ALG-2 is a cytosolic penta-EF hand-containing protein with two Ca^{2+} binding sites (Kd= 1.2 μM), and its interaction with Alix strictly depends on calcium (Missotten et al., 1999; Shibata et al., 2004; Trioulier et al., 2004). Conformational change and exposure of hydrophobic patches occur at μM concentrations of Ca^{2+}, suggesting

that ALG-2 functions as a calcium sensor (Maki et al., 2016). Indeed, in non-neuronal cells,

calcium entry provoked by membrane wounds leads to the sequential recruitment to the

membrane of ALG-2, Alix and ESCRT-III proteins necessary for membrane repair (Jimenez et

al., 2014; Scheffer et al., 2014). In neurons, the activity-dependent accumulation of ALG-2 at

synapses required Ca^{2+} binding as it was totally abolished by point-mutations within the two

Ca^{2+} binding sites. On the other hand, it did not require Alix, as ALG-2 recruitment also

occurred in Alix ko neuron. Thus, our observations highlight synaptic ALG-2 as an obvious

candidate for sensing calcium elevation and suggest the following scenario: sustained high

frequency depolarization leads to massive elevation of calcium in the synaptic bouton; Ca^{2+}

binding to ALG-2 causes its accumulation at the synaptic membrane and binding to cytosolic

Alix, whose recruitment to the membrane drives ADBE.

Little is known about the molecular mechanisms underlying the plasma membrane

deformation during ADBE. The proline rich domain of Alix interacts with the SH3 domains of

endophilins-A1-3 which contain N-BAR domains and have been shown to regulate clathrin-

dependent and -independent SV endocytosis at different synapses (Gad et al., 2000; Llobet et

al., 2011; Ringstad et al., 1999). BAR domains are dimerization domains able to induce,

stabilize and sense membrane curvature (Farsad et al., 2001; Kjaerulff et al., 2011).

Interestingly, a proteomic approach revealed the presence of endophilin-A1, Alix and ALG-2

in bulk endosomes (Kokotos et al., 2018). Furthermore, endophilin-A2, actin and dynamin

mediate a restricted type of CIE activated upon ligand binding to cargo receptors (Boucrot et

al., 2014; Renard et al., 2014). We previously showed that in fibroblasts, endophilin-A and Alix

act in the same CIE pathway, even if Alix is more promiscuous for ligands, as it is also involved

in fluid phase endocytosis. In this case interaction of Alix with endophilins favoured their

presence at the membrane (Mercier et al., 2016). We now found that endophilin-A2

decorated presynapses of resting neurons and that Alix was needed for its synaptic

enrichment during stimulation. An Alix point mutant unable to interact with endophilin was

recruited to synapses but could not rescue ABDE in Alix ko neurons, reminiscent of the

situation seen in fibroblasts (Mercier et al., 2016). Thus, a complex between endophilin-A and

Alix is required for ADBE. One function of Alix might be to enable endophilin-A binding onto

flat membranes of the peri-active zone to drive membrane bending in response to high

calcium concentrations (Boucrot et al., 2014; Kjaerulff et al., 2011). Interestingly, the

incapacity of Alix to interact with Chmp4B from ESCRT-III, known to alleviate its plasma

membrane repair ability (Jimenez et al., 2014; Scheffer et al., 2014), did not alter rescue of

ADBE. This result also discriminates Alix-driven ADBE from roles of the ESCRT system in the

degradation of SV proteins (Sadoul et al., 2017; Sheehan et al., 2016).

Electrophysiological recordings and EM observations in CA3-CA1 synapses both point to a

reduction in the number of synaptic vesicles due to the lack of Alix and possibly to the

associated defects in ADBE.   On the other hand, Syp-pH fluorescence and EM analysis

revealed that activity dependent synaptic endocytosis is increased in absence of Alix.

Accordingly, electrophysiological patch-clamp recordings show that Alix ko CA3-CA1 synapses

recover better from STD induced by repeated stimulations Thus, compensation by increased

endocytosis, is not sufficient to rescue the phenotype. This could be a first demonstration that

ADBE of the essential role of ADBE in regulating the number of synaptic vesicles and thereby

normal synaptic function.

Excitatory, glutamatergic neurotransmission plays a central role in the generation of seizure

activity (Albrecht and Zielińska, 2017; Barker-Haliski and White, 2015) and figures among the

primary anti-epileptic drug targets (Rogawski and Löscher, 2004). Sustained, intense and

synchronous neurotransmission as occurring during seizures requires replenishment of

synaptic vesicle pools mainly through activity-dependent bulk endocytosis (ADBE) (Cheung

and Cousin, 2012). Impairing ADBE *in vivo,* by deleting Alix in neocortical and hippocampal

excitatory neurons led to a significantly reduced number of seizures during SE, without

affecting the total duration of SE or the mean seizure duration. The present results suggest

that excitatory networks lacking Alix have a reduced capacity to initiate sequential seizures,

possibly due to an impaired replenishment of synaptic vesicles through ADBE. Seizure

termination and *status epilepticus* cessation however seem to be governed by distinct

mechanisms, probably involving inhibitory transmission to re-establish the balance between

excitation and inhibition in the brain. Our results highlight Alix ko mice as an invaluable tool

for exploring and understanding the exact role of ADBE at synapses undergoing normal and

pathological stimulations.

Acknowledgements

We thank J. Brocard for helping with data analysis, Y. Saoudi for helping with fluorescence microscopy, N. Liaudet from the Bioimaging core facility of the University of Geneva for his help with the segmentation and electron microscopy, F. Saudou for making the MEA experiments possible. We also thank K. Sadoul for critically reading the manuscript. This work was funded by France Alzheimer (R.S.), Ministère de l'Enseignement Supérieur et de la Recherche (K.I.C, M.R. and M.H.L.), Marie Sklodowska-Curie post-doctoral fellowship (M.M.); EC H2020 MSCA-ITN EU-GliapHD #722053 (L.C.C. and F.K.), Agence Nationale de la Recherche ANR (J.M.-H.) and Fondation pour la Recherche Medical FRM (E.M.).

Competing interests

On behalf of all authors, R.Sadoul declares no competing interests.

Author contributions

RS conceived the project. MHL participated to most of the in vitro experiments and analysed electron microscopy data under the supervision of SF. KIC performed and analysed most of the *in vitro* experiments concerning live imaging of Alix recruitment and dextran uptake. LCC performed and analysed the EEG telemetry and unilateral intracortical kainate injection. NZ performed and analysed patch clamp electrophysiological recordings. The experiments of LCC and NZ were done under the supervision of FK. MR and FLa performed field electrophysiological recordings. JM-H performed electron microscopy of mouse brains. MM and DP performed and analysed the live imaging of synaptophysin-pHluorin during electrical stimulation. CC performed mouse breeding, genotyping and participated in *in vitro* experiments. ED helped with data analysis of live imaging. VM performed FIB-SEM imaging. BB performed mouse genotyping and construction of plasmids. EM and MC performed and analysed multielectrode array experiments. FLe did some of the live imaging experiments of Alix recruitment *in vitro*. FJH performed electron microscopy of cultured neurons. RS and MHL wrote the manuscript with input from all other authors.

     Figure legends

Figure 1. Alix is recruited presynaptically during synaptic activation. A) Western blot analysis of
cortical neurons cultured for 3 to 15 DIV demonstrates the increase in Alix expression correlating with
synaptogenesis as illustrated by the increase in PSD95 expression. **B**) Western blot analysis of the
increase of Alix in synaptosome-enriched neuronal membranes upon neuronal stimulation by
bicuculline/4AP. Synaptophysin and PSD95 were used as pre- and postsynaptic markers, respectively.
The phosphorylated form of ERK (p-ERK) assessed the efficiency of the stimulation. **C**) Images from
time-lapse video microscopy of 15 DIV hippocampal neurons expressing both mCherry-Alix (mche-Alix)
and synaptophysin-pHluorin (Syp-pH) stimulated between 2 and 7 minutes with bicuculline/4AP.
White arrowheads indicate presynaptic boutons where Alix is recruited during stimulation. Scale bar:
10 µm. **D**) Profiles of mcherry-Alix and Syp-pH fluorescent increase at presynaptic boutons 1 and 2 of
panel **C**. Blue box indicates the bicuculline/4AP stimulation. **E**) 15 DIV hippocampal neurons expressing
Alix-YFP (green) stimulated for 5 min before fixation and stained with anti-synapsin-1 antibody (Syn,
magenta). Scale bar: 5 µm. **F**) Graph shows the presynaptic increase in Alix-YFP upon stimulation.
Presynaptic Alix-YFP corresponds to the ratio of YFP fluorescence between synapsin-positive and -
negative axonal regions. **G**) 15 DIV hippocampal neurons expressing Alix-YFP (green) stimulated for 5
min before fixation and stained with anti-synapsin-1 antibody (Syn, magenta) and anti-PSD95 (post-
synaptic, cyan) showing the selective recruitment of Alix to presynpase. Scale bar: 5 µm.
<u>Average +/- SEM, N, statistical analysis:</u> **B**) 2.07+/-0.34; 1.38+/-0.21; 1.00+/-0.01 for Alix,
Synaptophysin and PSD95 respectively. N=4 independent experiments, Alix versus PSD95, p=0.0187,
one-way ANOVA. **F**) 1.63+/-0.12; 3.27+/-0.44 for No Stim and Stim respectively. N=12 neurons per
condition from 4 independent experiments, p=0.0017, Unpaired t-test.

Figure 2. Interplay between Alix, ALG-2 and endophilin recruitments at activated synapses. All
experiments made use of 15 DIV hippocampal neurons expressing the indicated constructs. Neurons
were stimulated with bicuculline/4AP for 5 min before fixation and stained with anti-synapsin-1 (Syn,
magenta). **A**) GFP-ALG-2 (green) is recruited presynaptically upon stimulation. Scale bar: 5 µm. **B**)
Quantification of synaptic recruitments of GFP-ALG-2 and GFP-ALG-2ΔCa in Alix wt neurons shows that
ALG-2 presynaptic increase depends on its capacity to bind calcium. **C**) Alix recruitment depends on its
capacity to bind ALG-2 as GFP-AlixΔALG-2 is not recruited upon stimulation of Alix ko neurons.
However, ALG-2 does not require Alix to be recruited as GFP-ALG-2 is recruited presynaptically in Alix
ko neurons. **D**, **E**) Endophilin-mCherry (endo, green) concentrates at presynaptic parts of stimulated
neurons. Scale bar: 5 µm. **F**) Endophilin concentration following synaptic activation requires Alix as it
does not occur in Alix ko neurons. Alix recruitment does not depend on its binding to endophilins as
shown with GFP-AlixΔendo in Alix ko neurons.
<u>Average +/- SEM, N, statistical analysis:</u> **B)** 1.41+/-0.13; 2.24+/-0.11; 1.38+/-0.05; 1.18+/-0.15 for GFP-
ALG-2 no stim, GFP-ALG-2 stim, GFP-ALG-2ΔCa no stim, GFP-ALG-2ΔCa stim respectively. N=7 and n=10
neurons for GFP-ALG-2 not stimulated and stimulated, respectively and n=12 and n=6 neurons for GFP-
ALG-2ΔCa not stimulated and stimulated, respectively, from 3 experiments, p=0.0001, one-way
ANOVA. **C**) 1.37+/-0.07; 2.06+/-0.11; 1.54+/-0.05; 1.48+/-0.08 for GFP-ALG-2 no stim; GFP-ALG-2 stim,
GFP-AlixΔALG-2 no stim, GFP-AlixΔALG-2 stim respectively. N=17 neurons for both GFP-ALG-2 not
stimulated and stimulated, and n=18 and n=15 neurons for GFP-AlixΔALG-2 not stimulated and
stimulated, respectively, from 3 independent experiments, p=0.0001, one-way ANOVA. **D,E)** 1.92+/-
0.11; 3.48+/-0.33 for no stim and stim respectively. N=10 and n=12 neurons for not stimulated and
stimulated, respectively from 3 independent experiments, p=0.0001, Mann Whitney test. **F)** 2.37+/-
0.12; 2.66+/-0.08; 1.56+/-0.06; 2.43+/-0.32 for endophilin-mCherry no stim, endophilin-mCherry stim,
GFP-AlixΔendo no stim, GFP-AlixΔendo stim respectively. N=13 and n=15 neurons for endophilin-
mCherry not stimulated and stimulated, respectively, and n=10 and n=9 neurons for GFP-AlixΔendo
not stimulated and stimulated, respectively, from 3 independent experiments p=0.0031, one-way
ANOVA.

Figure 3. Alix is necessary for activity-dependent bulk endocytosis. A, **B**) Average traces of Syp-pH fluorescence in synaptic boutons of Alix wt and ko hippocampal neurons stimulated with 200 action potentials (AP) applied at 5 Hz (**A**) or 40 Hz (**B**). Insets show the exponential fit of fluorescence decay after stimulations in the fields imaged. **C**, **E**) Electron micrographs of cerebellar granule neurons stimulated in presence of free HRP to label newly-formed synaptic vesicle (blue arrowhead) and bulk endosomes (yellow arrowheads). **C**) Focused ion beam scanning electron microscopy (FIB-SEM); orthogonal views from different planes (xy, xz, yz) extracted from a stack used for the 3D reconstruction of wt presynaptic terminal shown in (**D**). Scale bar: 500nm. **D**) Two different views of the reconstructed synapse are shown where the membrane is represented in transparent gray and HRP-positive structures in red. Scale bar: 500nm. E) Transmission electron microscopy of HRP incubated CB Alix wt and Alix ko neurons. Scale bar: 100 nm. **F**, **G**) Quantification of the number of bulk endosomes (**F**) and synaptic vesicles (**G**) in Alix wt and ko neurons from images as shown in (E).

Average +/- SEM, N, statistical analysis: **A**, **B**) N= 16-46 fields of view per condition from 4 experiments for wt and 5 for ko mice, p=0.99 (**A**) and p=0.001 (**B**), one-way ANOVA. **F**, **G**) 0.52+/-0.17; 0.64+/-0.15; 3.28+/-0.19; 1.82+/-0.06 for Alix wt no stim, Alix ko no stim, Alix wt stim, Alix ko stim respectively (**F**). 4.43+/-1.02; 4.26+/-0.58; 7.46+/-0.49; 9.64+/-0.67 for Alix wt no stim, Alix ko no stim, Alix wt stim, Alix ko stim respectively (**G**). N= 44 and 51 neurons from 4 independent experiments, p<0.0001 (Alix wt no stim vs Alix wt stim, **F**), p=0.0015 (Alix ko no stim vs Alix ko stim, **F**), p=0.0001 (Alix wt stim vs Alix ko stim, **F**) and p=0.0376 (Alix wt no stim vs Alix wt stim, **G**), p=0.0014 (Alix ko no stim vs Alix ko stim, **G**), p=0.0393 (Alix wt stim vs Alix ko stim, **G**), one-way ANOVA.

Figure 4. Alix-driven bulk endocytosis requires its binding to ALG-2 and endophilin but not to CHMP4. A) Confocal images of Alix wt and Alix ko hippocampal neurons stimulated with Bicucculine/4AP in the presence of 10kDa dextran. Scale bar: 50 µm. **B**) Dextran uptake triggered by stimulation is strongly reduced in Alix ko neurons. **C**) Representative images of Dextran uptake by GFP-expressing Alix wt and Alix ko neurons (Alix wt, top and Alix ko,middle) or Alix ko neurons expressing both GFP and Alix (Alix ko+Alix, bottom). Scale bars: 10 µm. **D**) Dextran uptake is rescued in ko neurons expressing Alix wt and AlixΔChmp4B (AlixΔ4B), but not AlixΔALG-2 or AlixΔendo. % of dextran uptake corresponds to the number of dextran spots per µm expressed as percentages of the control (Alix wt) for each experiment.

Average +/- SEM, N, statistical analysis: **B**) 16.26+/-1.08; 11.86+/-2.05; 100+/-1.76; 15.10+/-1.76 for Alix wt no stim, Alix ko no stim, Alix wt stim, Alix ko stim respectively. N=4 experiments, p=0.0001, one-way ANOVA. **D**) 100+/-6.31; 33.27+/-3.32; 94.78+/-5.81; 45.89+/-9.36; 80.05+/-7.32; 44.59+/-1.85 for Alix wt, Alix ko, Alix ko + Alix, Alix ko + AlixΔChmp4B, Alix ko + AlixΔendo, Alix ko + AlixΔALG-2. N=17 neurons for Alix wt, n=13 neurons for Alix ko, n=11 for Alix ko + Alix, n=12 for Alix ko + AlixΔChmp4B, n=12 neurons for Alix ko + AlixΔendo, , and n=11 for Alix ko + AlixΔALG-2, all from 3-4 independent experiments, p=0.62 (Alix wt vs Alix AlixΔ4B) p=0.0001 for the other conditions, one-way ANOVA).

Figure 5. Morphological anomalies of Alix ko synapses. A) Representative electron micrographs of the CA1 stratum radiatum from Alix wt and ko brain sections. Presynaptic profiles are highlighted in blue, and dendritic spines are in purple. Scale bar: 2 µm. Graph shows no difference in synaptic density per µm^2 of Alix wt and ko brains. **B**) 15 DIV hippocampal neurons were stained with anti-PSD95 (magenta) and anti-synapsin-1 (green) antibodies. Immunolabelled objects were considered as synapses when both stainings were juxtaposed. Scale bar: 10 µm. Graph shows no difference in the number of synapses per µm^2 of Alix wt and ko neurons). **C-H**) Anomalies can be detected in synapses of Alix ko brains. **C**, **F**) Representative electron micrographs of CA1 from Alix wt and ko mice. Scale bar: 200 nm. Graphs represent the numbers of synaptic vesicles per µm^2 (**D**), presynaptic bouton surface area (**E**), PSD length (**F**), ratio between the diameter of the spine head and neck (**G**).

Average +/- SEM, N, statistical analysis: **A**) 0.40+/-0.02; 0.45+/-0.03 for Alix wt and Alix ko respectively. N=600 synapses from 3 animals per genotype, p=0.1395, Unpaired t-test. **B**) 0.13+/-0.01; 0.12+/-0.01 for Alix wt and ko respectively. N=7 independent experiments, p=0.5501, Unpaired t-test.

<u>Median (min to max), N, statistical analysis</u> **D)** 165.4 (69 to 278); 127.3 (31 to 336) for Alix wt and Alix
ko respectively. N=136 synapses from 3 animals, p=0.0009, Mann Whitney test). **E)** 0.13 (0.027 to
0.376); 0.16 (0.053 to 0.47) for Alix wt and Alix ko respectively. N=136 synapses from 3 animals,
p=0.0036, Mann Whitney test). **G)**: 0.76 (0.25 to 1.48) ; 0.54 (0.13 to 1.11) for Alix wt and Alix ko
respectively. N=136 synapses from 3 animals, p=0.0001, Mann Whitney test). **H)** 2.58 (0.85 to 5.60);
1.68 (0.58 to 4.78) for Alix wt and Alix ko respectively. N=136 synapses from 3 animals, p=0.0001, Mann
Whitney test.
945

Figure 6. Voltage-clamp recordings uncover anomalies at Alix ko synapses. **A)** Emx1 ^{IREScre} (Emx-Cre)
and Alix $^{fl/fl}$ mouse lines were crossbred to delete Alix in neocortical and hippocampal excitatory
neurons (Alix cko). **B)** No difference in spontaneous Excitatory Post Synaptic Current (sEPSC) amplitude
was detected between control and cko mice. **C)** The frequency of sEPSC in cko neurons is lower
compared to control. **D)** Representative traces showing short-term depression in response to 10 Hz
stimulation trains in control (black) and cko (grey) mice. **E)** Cumulative EPSC amplitudes in response to
10 Hz stimulation. Train-extrapolation is illustrated by the dashed line. **F)** The size of the readily
releasable pool, estimated by the train-extrapolation method, was significantly decreased in Alix cko
mice. **G**) Representative traces showing recovery from depression in response to 10 Hz stimulation
trains in control (black) and Alix cko (grey) mice. **H)** To evaluate the speed of recovery, all inward signals
were normalized to 1st inward current. Recovering signals with a single-exponential function revealed
the plateaus at which the capacities of recovery were saturated. **I)** Alix cko neurons slightly recover
faster than controls.
<u>Average +/- SEM, N, statistical analysis:</u> **B)** 9.79+/-0.53; 8.88+/-0.42 for controls and Alix cko
respectively. N=11 neurons from 5 control mice and N=17 neurons from 6 Alix cko mice, p=0.1875,
unpaired t test. **C)** 0.93+/-0.25; 0.44+/-0.10 for controls and Alix cko respectively. N=11 neurons from
5 control mice and N=17 neurons from 6 Alix cko mice, **p=0.0095, Mann Whitney test. **F)** 1.23+/-
0.15; 0.67+/-0:08 for controls and Alix cko respectively. N=12 neurons from 5 control mice, and N=10
neurons from 4 Alix cko mice, **p=0.0052, unpaired t test. **I)** 8.34+/-0.90; 5.56+/-0.69 for controls and
Alix cko respectively. N= 5 and 6 neurons from 3 mice from control and Alix cko respectively,
*p=0.0336, unpaired t test.
967

Figure 7. Alix cko mice develop less seizures during status epilepticus. A) Representative EEG traces
from control and Alix cko mice. **B)** Total duration of *status epilepticus* (SE) did not differ between Alix
cko and control mice. **C)** Alix cko mice experience about 66 % less seizures during SE than controls. **D)**
Mean seizure duration during SE was not affected in Alix cko mice compared to controls.
<u>Median + IQR, N, statistical analysis:</u> **A**, **B**) 1.6 h (IQR = 2.5); 1.7 h (IQR = 0.9) for control and Alix cko
mice respectively. N=11 mice for both genotype, p= 0.49, Mann-Whitney test. **C)** 18 seizures (IQR = 9);
8 seizures (IQR = 9) for control and Alix cko mice respectively. N=11 mice for both genotypes, *p= 0.03,
Mann-Whitney test. **D)** 32.9 s (IQR = 18.6); 25.7 h (IQR = 12.7) for control and Alix cko mice
respectively. N=11 mice for both genotype, p= 0.19, Mann-Whitney test.
977
978
979
References

Adler, E.M., Augustine, G.J., Duffy, S.N., and Charlton, M.P. (1991). Alien intracellular calcium
chelators attenuate neurotransmitter release at the squid giant synapse. J Neurosci *11*,
1496-1507.
Albrecht, J., and Zielińska, M. (2017). Mechanisms of Excessive Extracellular Glutamate
Accumulation in Temporal Lobe Epilepsy. Neurochem Res *42*, 1724-1734.
Barker-Haliski, M., and White, H.S. (2015). Glutamatergic Mechanisms Associated with
Seizures and Epilepsy. Cold Spring Harb Perspect Med *5*, a022863.

Bedner, P., Dupper, A., Huttmann, K., Muller, J., Herde, M.K., Dublin, P., Deshpande, T.,
Schramm, J., Haussler, U., Haas, C.A., *et al.* (2015). Astrocyte uncoupling as a cause of human
temporal lobe epilepsy. Brain *138*, 1208-1222.
Bissig, C., Lenoir, M., Velluz, M.C., Kufareva, I., Abagyan, R., Overduin, M., and Gruenberg, J.
(2013). Viral infection controlled by a calcium-dependent lipid-binding module in ALIX. Dev
Cell *25*, 364-373.
Boucrot, E., Ferreira, A.P., Almeida-Souza, L., Debard, S., Vallis, Y., Howard, G., Bertot, L.,
Sauvonnet, N., and McMahon, H.T. (2014). Endophilin marks and controls a clathrin-
independent endocytic pathway. Nature.
Campos, Y., Qiu, X., Gomero, E., Wakefield, R., Horner, L., Brutkowski, W., Han, Y.G., Solecki,
D., Frase, S., Bongiovanni, A., *et al.* (2016). Alix-mediated assembly of the actomyosin-tight
junction polarity complex preserves epithelial polarity and epithelial barrier. Nature
communications *7*, 11876.
Chanaday, N.L., and Kavalali, E.T. (2018). Time course and temperature dependence of
synaptic vesicle endocytosis. FEBS Lett *592*, 3606-3614.
Chassefeyre, R., Martinez-Hernandez, J., Bertaso, F., Bouquier, N., Blot, B., Laporte, M.,
Fraboulet, S., Coute, Y., Devoy, A., Isaacs, A.M., *et al.* (2015). Regulation of Postsynaptic
Function by the Dementia-Related ESCRT-III Subunit CHMP2B. J Neurosci *35*, 3155-3173.
Chatellard-Causse, C., Blot, B., Cristina, N., Torch, S., Missotten, M., and Sadoul, R. (2002).
Alix (ALG-2-interacting protein X), a protein involved in apoptosis, binds to endophilins and
induces cytoplasmic vacuolization. J Biol Chem *277*, 29108-29115.
Cheung, G., and Cousin, M.A. (2012). Adaptor protein complexes 1 and 3 are essential for
generation of synaptic vesicles from activity-dependent bulk endosomes. J Neurosci *32*,
6014-6023.
Cheung, G., Jupp, O.J., and Cousin, M.A. (2010). Activity-dependent bulk endocytosis and
clathrin-dependent endocytosis replenish specific synaptic vesicle pools in central nerve
terminals. J Neurosci *30*, 8151-8161.
Clayton, E.L., and Cousin, M.A. (2009a). The molecular physiology of activity-dependent bulk
endocytosis of synaptic vesicles. J Neurochem *111*, 901-914.
Clayton, E.L., and Cousin, M.A. (2009b). Quantitative monitoring of activity-dependent bulk
endocytosis of synaptic vesicle membrane by fluorescent dextran imaging. J Neurosci
Methods *185*, 76-81.
Clayton, E.L., Sue, N., Smillie, K.J., O'Leary, T., Bache, N., Cheung, G., Cole, A.R., Wyllie, D.J.,
Sutherland, C., Robinson, P.J., *et al.* (2010). Dynamin I phosphorylation by GSK3 controls
activity-dependent bulk endocytosis of synaptic vesicles. Nat Neurosci *13*, 845-851.
de Chaumont, F., Dallongeville, S., Chenouard, N., Herve, N., Pop, S., Provoost, T., Meas-
Yedid, V., Pankajakshan, P., Lecomte, T., Le Montagner, Y., *et al.* (2012). Icy: an open
bioimage informatics platform for extended reproducible research. Nature methods *9*, 690-
696.
Farsad, K., Ringstad, N., Takei, K., Floyd, S.R., Rose, K., and De Camilli, P. (2001). Generation
of high curvature membranes mediated by direct endophilin bilayer interactions. J Cell Biol
155, 193-200.
Faure, J., Lachenal, G., Court, M., Hirrlinger, J., Chatellard-Causse, C., Blot, B., Grange, J.,
Schoehn, G., Goldberg, Y., Boyer, V., *et al.* (2006). Exosomes are released by cultured cortical
neurones. Mol Cell Neurosci *31*, 642-648.

Frandemiche, M.L., De Seranno, S., Rush, T., Borel, E., Elie, A., Arnal, I., Lante, F., and
Buisson, A. (2014). Activity-dependent tau protein translocation to excitatory synapse is
disrupted by exposure to amyloid-beta oligomers. J Neurosci *34*, 6084-6097.
Gad, H., Ringstad, N., Low, P., Kjaerulff, O., Gustafsson, J., Wenk, M., Di Paolo, G., Nemoto,
Y., Crun, J., Ellisman, M.H.*, et al.* (2000). Fission and uncoating of synaptic clathrin-coated
vesicles are perturbed by disruption of interactions with the SH3 domain of endophilin.
Neuron *27*, 301-312.
Gan, Q., and Watanabe, S. (2018). Synaptic Vesicle Endocytosis in Different Model Systems.
Frontiers in cellular neuroscience *12*, 171.
Gorski, J.A., Talley, T., Qiu, M., Puelles, L., Rubenstein, J.L., and Jones, K.R. (2002). Cortical
excitatory neurons and glia, but not GABAergic neurons, are produced in the Emx1-
expressing lineage. J Neurosci *22*, 6309-6314.
Granseth, B., Odermatt, B., Royle, S.J., and Lagnado, L. (2006). Clathrin-mediated
endocytosis is the dominant mechanism of vesicle retrieval at hippocampal synapses.
Neuron *51*, 773-786.
Hardingham, G.E., Arnold, F.J., and Bading, H. (2001). Nuclear calcium signaling controls
CREB-mediated gene expression triggered by synaptic activity. Nat Neurosci *4*, 261-267.
Hardingham, G.E., Fukunaga, Y., and Bading, H. (2002). Extrasynaptic NMDARs oppose
synaptic NMDARs by triggering CREB shut-off and cell death pathways. Nat Neurosci *5*, 405-
414.
Harris, K.M., and Stevens, J.K. (1989). Dendritic Spines of Ca-1 Pyramidal Cells in the Rat
Hippocampus - Serial Electron-Microscopy with Reference to Their Biophysical
Characteristics. Journal of Neuroscience *9*, 2982-2997.
Hemming, F.J., Fraboulet, S., Blot, B., and Sadoul, R. (2004). Early increase of apoptosis-
linked gene-2 interacting protein X in areas of kainate-induced neurodegeneration.
Neuroscience *123*, 887-895.
Jimenez, A.J., Maiuri, P., Lafaurie-Janvore, J., Divoux, S., Piel, M., and Perez, F. (2014). ESCRT
machinery is required for plasma membrane repair. Science *343*, 1247136.
Kjaerulff, O., Brodin, L., and Jung, A. (2011). The structure and function of endophilin
proteins. Cell biochemistry and biophysics *60*, 137-154.
Kokotos, A.C., Peltier, J., Davenport, E.C., Trost, M., and Cousin, M.A. (2018). Activity-
dependent bulk endocytosis proteome reveals a key presynaptic role for the monomeric
GTPase Rab11. Proc Natl Acad Sci U S A *115*, E10177-E10186.
Kononenko, N.L., Puchkov, D., Classen, G.A., Walter, A.M., Pechstein, A., Sawade, L., Kaempf,
N., Trimbuch, T., Lorenz, D., Rosenmund, C.*, et al.* (2014). Clathrin/AP-2 mediate synaptic
vesicle reformation from endosome-like vacuoles but are not essential for membrane
retrieval at central synapses. Neuron *82*, 981-988.
Körber, C., Horstmann, H., Venkataramani, V., Herrmannsdörfer, F., Kremer, T., Kaiser, M.,
Schwenger, Darius B., Ahmed, S., Dean, C., Dresbach, T.*, et al.* (2015). Modulation of
Presynaptic Release Probability by the Vertebrate-Specific Protein Mover. Neuron *87*, 521-
533.
Lagache, T., Grassart, A., Dallongeville, S., Faklaris, O., Sauvonnet, N., Dufour, A., Danglot, L.,
and Olivo-Marin, J.C. (2018). Mapping molecular assemblies with fluorescence microscopy
and object-based spatial statistics. Nature communications *9*, 698.
Laporte, M.H., Chatellard, C., Vauchez, V., Hemming, F.J., Deloulme, J.C., Vossier, F., Blot, B.,
Fraboulet, S., and Sadoul, R. (2017). Alix is required during development for normal growth
of the mouse brain. Sci Rep *7*, 44767.

Lidierth, M. (2009). sigTOOL: A MATLAB-based environment for sharing laboratory-
developed software to analyze biological signals. J Neurosci Methods *178*, 188-196.
Llobet, A., Gallop, J.L., Burden, J.J., Camdere, G., Chandra, P., Vallis, Y., Hopkins, C.R.,
Lagnado, L., and McMahon, H.T. (2011). Endophilin drives the fast mode of vesicle retrieval
in a ribbon synapse. J Neurosci *31*, 8512-8519.
Maki, M., Takahara, T., and Shibata, H. (2016). Multifaceted Roles of ALG-2 in Ca(2+)-
Regulated Membrane Trafficking. Int J Mol Sci *17*.
Martineau, M., Somasundaram, A., Grimm, J.B., Gruber, T.D., Choquet, D., Taraska, J.W.,
Lavis, L.D., and Perrais, D. (2017). Semisynthetic fluorescent pH sensors for imaging
exocytosis and endocytosis. Nature communications *8*.
Marxen, M., Volknandt, W., and Zimmermann, H. (1999). Endocytic vacuoles formed
following a short pulse of K+ -stimulation contain a plethora of presynaptic membrane
proteins. Neuroscience *94*, 985-996.
Mercier, V., Laporte, M.H., Destaing, O., Blot, B., Blouin, C.M., Pernet-Gallay, K., Chatellard,
C., Saoudi, Y., Albiges-Rizo, C., Lamaze, C.*, et al.* (2016). ALG-2 interacting protein-X (Alix) is
essential for clathrin-independent endocytosis and signaling. Sci Rep *6*, 26986.
Miller, T.M., and Heuser, J.E. (1984). Endocytosis of synaptic vesicle membrane at the frog
neuromuscular junction. J Cell Biol *98*, 685-698.
Milosevic, I., Giovedi, S., Lou, X., Raimondi, A., Collesi, C., Shen, H., Paradise, S., O'Toole, E.,
Ferguson, S., Cremona, O.*, et al.* (2011). Recruitment of endophilin to clathrin-coated pit
necks is required for efficient vesicle uncoating after fission. Neuron *72*, 587-601.
Missotten, M., Nichols, A., Rieger, K., and Sadoul, R. (1999). Alix, a novel mouse protein
undergoing calcium-dependent interaction with the apoptosis-linked-gene 2 (ALG-2) protein.
Cell Death Differ *6*, 124-129.
Morton, A., Marland, J.R., and Cousin, M.A. (2015). Synaptic vesicle exocytosis and increased
cytosolic calcium are both necessary but not sufficient for activity-dependent bulk
endocytosis. J Neurochem *134*, 405-415.
Neher, E. (2015). Merits and Limitations of Vesicle Pool Models in View of Heterogeneous
Populations of Synaptic Vesicles. Neuron *87*, 1131-1142.
Paillart, C., Li, J., Matthews, G., and Sterling, P. (2003). Endocytosis and vesicle recycling at a
ribbon synapse. J Neurosci *23*, 4092-4099.
Racine, R.J. (1972). Modification of seizure activity by electrical stimulation. II. Motor
seizure. Electroencephalogr Clin Neurophysiol *32*, 281-294.
Renard, H., Simunovic, M., Lemiere, J., Boucrot, E., Garcia-Castillo, M.D., Arumugam, S.,
Chambon, V., Lamaze, C., Wunder, C., Kenworthy, A.K.*, et al.* (2014). Endophilin-A2 functions
in membrane scission in clathrin-independent endocytosis. Nature.
Ringstad, N., Gad, H., Low, P., Di Paolo, G., Brodin, L., Shupliakov, O., and De Camilli, P.
(1999). Endophilin/SH3p4 is required for the transition from early to late stages in clathrin-
mediated synaptic vesicle endocytosis. Neuron *24*, 143-154.
Rogawski, M.A., and Löscher, W. (2004). The neurobiology of antiepileptic drugs. Nat Rev
Neurosci *5*, 553-564.
Sadoul, R. (2006). Do Alix and ALG-2 really control endosomes for better or for worse. Bio
Cell *98*, 69-77.
Sadoul, R., Laporte, M.H., Chassefeyre, R., Chi, K.I., Goldberg, Y., Chatellard, C., Hemming,
F.J., and Fraboulet, S. (2017). The role of ESCRT during development and functioning of the
nervous system. Semin Cell Dev Biol.

Scheffer, L.L., Sreetama, S.C., Sharma, N., Medikayala, S., Brown, K.J., Defour, A., and Jaiswal,
J.K. (2014). Mechanism of Ca(2+)-triggered ESCRT assembly and regulation of cell membrane
repair. Nature communications *5*, 5646.
Schneggenburger, R., Meyer, A.C., and Neher, E. (1999). Released fraction and total size of a
pool of immediately available transmitter quanta at a calyx synapse. Neuron *23*, 399-409.
Schneggenburger, R., and Neher, E. (2005). Presynaptic calcium and control of vesicle fusion.
Curr Opin Neurobiol *15*, 266-274.
Schuske, K.R., Richmond, J.E., Matthies, D.S., Davis, W.S., Runz, S., Rube, D.A., van der Bliek,
A.M., and Jorgensen, E.M. (2003). Endophilin is required for synaptic vesicle endocytosis by
localizing synaptojanin. Neuron *40*, 749-762.
Sheehan, P., Zhu, M., Beskow, A., Vollmer, C., and Waites, C.L. (2016). Activity-Dependent
Degradation of Synaptic Vesicle Proteins Requires Rab35 and the ESCRT Pathway. J Neurosci
36, 8668-8686.
Shibata, H., Yamada, K., Mizuno, T., Yorikawa, C., Takahashi, H., Satoh, H., Kitaura, Y., and
Maki, M. (2004). The penta-EF-hand protein ALG-2 interacts with a region containing PxY
repeats in Alix/AIP1, which is required for the subcellular punctate distribution of the amino-
terminal truncation form of Alix/AIP1. J Biochem (Tokyo) *135*, 117-128.
Soykan, T., Kaempf, N., Sakaba, T., Vollweiter, D., Goerdeler, F., Puchkov, D., Kononenko,
N.L., and Haucke, V. (2017). Synaptic Vesicle Endocytosis Occurs on Multiple Timescales and
Is Mediated by Formin-Dependent Actin Assembly. Neuron *93*, 854-866 e854.
Suzuki, H., Kawasaki, M., Inuzuka, T., Okumura, M., Kakiuchi, T., Shibata, H., Wakatsuki, S.,
and Maki, M. (2008). Structural basis for Ca2+ -dependent formation of ALG-2/Alix peptide
complex: Ca2+/EF3-driven arginine switch mechanism. Structure *16*, 1562-1573.
Thanawala, M.S., and Regehr, W.G. (2016). Determining synaptic parameters using high-
frequency activation. J Neurosci Methods *264*, 136-152.
Trioulier, Y., Torch, S., Blot, B., Cristina, N., Chatellard-Causse, C., Verna, J.M., and Sadoul, R.
(2004). Alix, a protein regulating endosomal trafficking, is involved in neuronal death. J Biol
Chem *279*, 2046-2052.
Watanabe, S., Mamer, L.E., Raychaudhuri, S., Luvsanjav, D., Eisen, J., Trimbuch, T., Sohl-
Kielczynski, B., Fenske, P., Milosevic, I., Rosenmund, C.*, et al.* (2018). Synaptojanin and
Endophilin Mediate Neck Formation during Ultrafast Endocytosis. Neuron *98*, 1184-1197
e1186.
1158

FIGURE1

FIGURE2

FIGURE 3

FIGURE4

FIGURE5

FIGURE6

A

Control

Alix cko

1mV
2min

1mV
5sec

B

ns

SE duration (h)

Control   Alix cko

C

*

# of seizures in SE

Control   Alix cko

D

ns

Mean seizure duration (s)

Control   Alix cko

FIGURE7