Regulation of immune receptor kinases plasma membrane nanoscale landscape by a plant peptide hormone and its receptors

Julien Gronnier[1,2,7], Christina M. Franck[1], Martin Stegmann[2,8], Thomas A. DeFalco[1,2], Alicia Abarca[1], Michelle Von Arx[1], Kai Dünser[3], Wenwei Lin[4], Zhenbiao Yang[4], Jürgen Kleine-Vehn[5,6], Christoph Ringli[1], and Cyril Zipfel[1,2]

[1]Institute of Plant and Microbial Biology and Zurich-Basel Plant Science Center, University of Zurich, 8008 Zurich, Switzerland.

[2]The Sainsbury Laboratory, University of East Anglia, Norwich Research Park, NR4 7UH, Norwich, UK.

[3]Department of Applied Genetics and Cell Biology, University of Natural Resources and Life Sciences Vienna, 1190 Vienna, Austria.

[4]FAFU-UCR Joint Center for Horticultural Biology and Metabolomics Center, Haixia, Institute of Science and Technology, Fujian Agriculture and Forestry University, Fuzhou, Fujian, China.

[5]Faculty of Biology, Department of Molecular Plant Physiology (MoPP), University of Freiburg, 79104 Freiburg, Germany.

[6]Center for Integrative Biological Signalling Studies (CIBSS), University of Freiburg, 79104 Freiburg, Germany.

[7]Present address: University of Tübingen, Center for Plant Molecular Biology (ZMBP), 72076 Tübingen, Germany.

[8]Present address: Phytopathology, School of Life Sciences Weihenstephan, Technical University of Munich, Freising, Germany.

Correspondence: Cyril Zipfel, cyril.zipfel@botinst.uzh.ch

ABSTRACT

Spatial partitioning is a propensity of biological systems orchestrating cell activities in space and time. The dynamic regulation of plasma membrane nano-environments has recently emerged as a key fundamental aspect of plant signaling, but the molecular components governing it are still mostly unclear. The receptor kinase FERONIA (FER) controls complex formation of the immune receptor kinase FLAGELLIN SENSING 2 (FLS2) with its co-receptor BRASSINOSTEROID INSENSITIVE 1-ASSOCIATED KINASE 1 (BAK1), and this function is inhibited by the FER ligand RAPID ALKALANIZATION FACTOR 23 (RALF23). Here, we show that FER regulates the plasma membrane nanoscale organization of FLS2 and BAK1. Our study demonstrates that akin to FER, leucine-rich repeat (LRR) extensin (LRXs) proteins contribute to RALF23 responsiveness, regulate BAK1 nanoscale organization and immune signaling. Furthermore, RALF23 perception leads to rapid modulation of FLS2 and BAK1 nanoscale organization and its inhibitory activity on immune signaling relies on FER kinase activity. Our results suggest that perception of RALF peptides by FER and LRXs actively modulates the plasma membrane nanoscale landscape to regulate cell surface signaling by other receptor kinases.

INTRODUCTION

Multicellular organisms evolved sophisticated surveillance systems to monitor changes in their environment. In plants, receptor kinases (RKs) and receptor proteins (RPs) are the main ligand-binding cell-surface receptors perceiving self, non-self, and modified-self molecules (Hohmann, Lau and Hothorn, 2017). For example, recognition of pathogen-associated molecular patterns (PAMPs) by pattern recognition receptors (PRRs) initiates signaling events leading to PRR-triggered immunity (PTI) (Couto and Zipfel, 2016; Yu *et al.*, 2017). The *Arabidopsis thaliana* (hereafter Arabidopsis) leucine-rich repeat receptor kinases (LRR-RKs) FLAGELLIN SENSING 2 (FLS2) and EF-TU RECEPTOR (EFR) recognize the bacterial PAMPs flagellin (or its derived epitope flg22) and elongation factor-Tu (or its derived epitope elf18), respectively (Gómez-Gómez and Boller, 2000; Zipfel *et al.*, 2006). Both FLS2 and EFR form ligand-induced complexes with the co-receptor LRR-RK BRASSINOSTEROID INSENSITIVE 1-ASSOCIATED KINASE 1/SOMATIC EMBRYOGENESIS RECEPTOR KINASE 3 (BAK1/SERK3, hereafter BAK1) to initiate immune signaling, such as for instance, the production of apoplastic reactive oxygen species (ROS) (Chinchilla *et al.*, 2007; Schulze *et al.*, 2010; Schwessinger *et al.*, 2011; Sun *et al.*, 2013).

We previously showed that the *Catharanthus roseus* RECEPTOR-LIKE PROTEIN KINASE 1-LIKE (CrRLK1L) FERONIA (FER) and the LORELEI-LIKE-GPI ANCHORED PROTEIN 1 (LLG1) are required for flg22-induced FLS2-BAK1 complex formation (Stegmann *et al.*, 2017; Xiao *et al.*, 2019). Notably, the endogenous RAPID ALKALINIZATION FACTOR 23 (RALF23) is perceived by a LLG1-FER complex, which leads to inhibition of flg22-induced FLS2-BAK1 complex formation (Stegmann *et al.*, 2017; Xiao *et al.*, 2019). As such, although FER and LLG1 are positive regulator of PTI, RALF23 is a negative regulator. How these components regulate FLS2-BAK1 complex formation remains however unclear.

Members of the CrLKL1s family are involved in RALF perception (Ge *et al.*, 2017; Gonneau *et al.*, 2018; Liu *et al.*, 2021). Among them, FER plays a pivotal role in the perception of several Arabidopsis RALF peptides (Haruta *et al.*, 2014; Stegmann *et al.*, 2017; Gonneau *et al.*, 2018; Zhao *et al.*, 2018; Abarca, Franck and Zipfel, 2021; Liu *et al.*, 2021). In addition, cell-wall associated LEUCINE RICH REPEAT-EXTENSINs (LRXs) proteins are also involved in CrRLK1L-regulated pathways and were shown to be high-affinity RALF-binding proteins (Mecchia *et al.*, 2017; Zhao *et al.*, 2018; Dünser *et al.*, 2019; Moussu *et al.*, 2020). Structural and biochemical analyses have shown that RALF-binding by CrRLK1L/LLGs complexes and LRXs are mutually exclusive and mechanistically distinct from each other (Xiao *et al.*, 2019; Moussu *et al.*, 2020). While CrRLK1Ls and LRXs have emerged as

important RALF-regulated signaling modules, it is still unknown whether LRXs are also

involved in RALF23-mediated regulation of immune signaling.

Plasma membrane lipids and proteins dynamically organize into diverse membrane

domains giving rise to fluid molecular patchworks (Gronnier *et al.*, 2018; Jaillais and Ott,

2020). These domains are proposed to provide dedicated biochemical and biophysical

environments to ensure acute, specific and robust signaling events (Gronnier *et al.*, 2019;

Jacobson, Liu and Lagerholm, 2019). For instance, FLS2 localizes in discrete and static

structures proposed to specify immune signaling (Bücherl *et al.*, 2017). The cell wall is

thought to impose physical constraints on the plasma membrane, limiting the diffusion of its

constituents (Feraru *et al.*, 2011; Martinière *et al.*, 2012). Indeed, alteration of cell wall

integrity leads to aberrant protein motions at the plasma membrane (Martinière *et al.*, 2012;

McKenna *et al.*, 2019). Notably, perturbation of the cell wall affects FLS2 nanoscale

organization (McKenna *et al.*, 2019). Despite its imminent importance, it remains largely

unknown how the cell wall and its integrity modulate the organization of the plasma

membrane. Interestingly, both CrRLK1Ls and LRXs are proposed cell wall integrity sensors

and conserved modules regulating growth, reproduction and immunity (Franck, Westermann

and Boisson-Dernier, 2018; Herger *et al.*, 2019). However, their mode of action and potential

link to RALF perception, are still poorly understood.

Here, we show that FER regulates the plasma membrane nanoscale organization of

FLS2 and BAK1. Similarly, we show that LRXs contribute to RALF23 responsiveness,

regulate BAK1 nanoscale organization and immune signaling. Importantly, our work reveals

an unexpected uncoupling of FER and LRXs modes-of-action in growth and immunity. We

demonstrate that RALF23 perception leads to rapid modulation of FLS2 and BAK1

nanoscale organization and that its inhibitory activity on immune signaling requires FER

kinase activity. We propose that the regulation of the plasma membrane nanoscale

organization by RALF23 receptors underscores their role in the formation of protein

complexes and initiation of immune signaling.

RESULTS AND DISCUSSION

FER regulates membrane nanoscale organization of FLS2 and BAK1

We combined variable angle total internal reflection fluorescence microscopy (VA-TIRFM)

and single-particle tracking to analyze the lateral mobility of FLS2-GFP proteins in transgenic

Arabidopsis lines. Two lines expressing FLS2-GFP under the control of its native promoter

were crossed with two *FER* knock-out alleles, *fer-2* and *fer-4*. In line with previous reports

(Bücherl *et al.*, 2017; Tran *et al.*, 2020), we observed that FLS2-GFP localizes to static foci in

the wild-type (WT) (Movie S1). Consistently, FLS2-GFP single particle trajectories exhibit a

confined mobility behavior (Fig. S1, Movie S1). Comparative analysis of the diffusion

coefficient (D), which describes the diffusion properties of detected single particles, showed
that FLS2-GFP is more mobile in *fer* mutants than in WT (Fig. S1, S2 and Movie S1). To
analyze FLS2-GFP organization, we reconstructed images using a temporal averaging of
FLS2-GFP fluorescence observed across VA-TIRFM time series. Furthermore, individual
image sections were subjected to kymograph analysis. Using this approach, we found that
FLS2-GFP fluorescence is maintained into well-defined and static structures in WT, while it
appears more disperse and more labile in both *fer* mutants (Fig. 1A-B and S2). To
substantiate these observations, we used previously established spatial clustering index
(SCI) which describes protein lateral organization (Gronnier *et al.*, 2017; Tran *et al.*, 2020).
As expected, SCI of FLS2-GFP was lower in *fer-4* than in WT (Fig. 1C), indicating
disturbance in FLS2-GFP lateral organization.

In *Medicago truncatula* and in yeast, alteration of nanodomain localization has been linked to
impaired protein accumulation at the plasma membrane due to increased protein
endocytosis (Grossmann *et al.*, 2008; Liang *et al.*, 2018). To inquire for a potential defect in
FLS2 plasma membrane accumulation, we observed sub-cellular localization of FLS2-GFP
using confocal microscopy. The analysis revealed a decrease in FLS2-GFP accumulation in
fer mutants (Fig. S3). Altogether, these results show that *FER* is genetically required to
control FLS2-GFP nanoscale organization and accumulation at the plasma membrane.

To further characterize the impact of *FER* loss-of-function in RK organization, we analyzed
the behavior of BAK1-mCherry at the plasma membrane. Fluorescence recovery after
photobleaching experiments previously suggested that the vast majority of BAK1 molecules
are mobile (Hutten *et al.*, 2017). Consistent with this result, BAK1-mCherry was much more
mobile than FLS2-GFP in the WT background (Movie S2). Given that BAK1 is a common co-
receptor for multiple LRR-RK signaling pathways, we hypothesize that BAK1 might
dynamically associates with various pre-formed signaling platforms, such as FLS2
nanodomains (Fig. 1, Bücherl *et al.*, 2017). Under our experimental conditions, we were not
able to perform high-quality single particle tracking analysis for BAK1-mCherry (Movie S2;
see methods section). However, visual inspection of particles behavior suggests that BAK1-
mCherry is less mobile in *fer-4* than in WT (Movie S2). In good agreement, reconstructed
VA-TIRFM images and kymographs show that BAK1-mCherry fluorescence is more
structured and static in *fer-4* than in WT (Fig. 1F). Furthermore, we observed an increase of
BAK1-mCherry SCI in *fer-4* (Fig 1G). Confocal microscopy analysis did not reveal significant
differences in BAK1-mCherry plasma membrane accumulation between *fer-4* and WT
backgrounds (Fig. S4). Altogether, these data show that loss of *FER* perturbs FLS2 and
BAK1 nanoscale organization, albeit in an opposite manner (Fig. 1D and 1H). Previous
reports have similarly shown that altering the composition of the cell wall can lead to
opposed effects on the mobility of different proteins. For instance, inhibition of cellulose

synthesis increases the mobility of HYPERSENSITIVE INDUCED REACTION 1 (HIR1)

(Daněk *et al.*, 2020) but limits the mobility of LOW TEMPERATURE INDUCED PROTEIN 6B

(Lti6b) (Martinière *et al.*, 2012; Daněk *et al.*, 2020). Modification of pectin methyl esterification

status increases the mobility of FLS2 (McKenna *et al.*, 2019) but decreases the mobility of

FLOTILIN 2 (FLOT2) (Daněk *et al.*, 2020). Collectively, these observations suggest that

various membrane environments are differentially regulated by the cell wall and the proposed

cell wall integrity sensor FER.

LRX3, LRX4 and LRX5 regulate BAK1 nanoscale organization and PTI signaling

LRXs are dimeric, cell wall-localized, high-affinity RALF-binding proteins suggested to

monitor cell wall integrity in growth and reproduction (Baumberger, Ringli and Keller, 2001;

Mecchia *et al.*, 2017; Dünser *et al.*, 2019; Herger *et al.*, 2019, 2020; Moussu *et al.*, 2020).

Their extensin domain confers cell wall anchoring while their LRR domain mediates RALF

binding (Herger *et al.*, 2019; Moussu *et al.*, 2020). Among the Arabidopsis 11-member *LRX*

family, *LRX3*, *LRX4*, and *LRX5* are the most expressed in vegetative tissues, and the *lrx3*

lrx4 lrx5 triple mutant (hereafter *lrx3/4/5*) shows stunted growth and salt hypersensitivity

phenotypes reminiscent of the *fer-4* mutant (Zhao *et al.*, 2018; Dünser *et al.*, 2019).

Therefore, we hypothesized that LRXs also regulate immune signaling. Indeed, our co-

immunoprecipitation experiments showed that *lrx3/4/5* is defective in flg22-induced FLS2-

BAK1 complex formation (Fig. 2A). Consistently, flg22-induced ROS production was reduced

in *lrx3/4/5* similar to the levels observed in *fer-4* (Fig. 2B). In addition, we found that *lrx3/4/5*

was impaired in elf18-induced ROS production (Fig. 2C), suggesting that, as for FLS2-BAK1

complex formation, *LRX3/4/5* are required for complex formation between EFR and BAK1.

Thus, we conclude that LRX3/4/5 are positive regulators of PTI signaling.

We then asked whether, similar to FER, LRX3/4/5 regulate plasma membrane nanoscale

organization. We crossed lines expressing FLS2-GFP and BAK1-mCherry under the control

of their respective native promoter with *lrx3/4/5* background. However, despite several

attempts, we could not retrieve homozygous *lrx3/4/5* lines expressing FLS2-GFP.

Nonetheless, VA-TIRFM and confocal imaging showed that, like in *fer-4*, BAK1-mCherry is

more organized and more static in *lrx3/4/5* (Fig. 2D-E, Movie S3), and that BAK1-mCherry

plasma membrane localization is not affected by the loss of *LRX3/4/5* (Fig. S4). Thus, like in

fer mutants, perturbation in PTI signaling observed in *lrx3/4/5* correlates with alterations of

plasma membrane RK organization.

Formerly, LRX3, LRX4 and LRX5 have been proposed to sequester RALF peptides to

prevent internalization of FER and inhibition of its function (Zhao *et al.*, 2018). Following this

logic, defects in PTI observed in *lrx3/4/5* could be explained by a depletion of FER at the

plasma membrane. However, our confocal microscopy analysis and western blotting with

anti-FER antibodies indicated that FER accumulation and plasma membrane localization is

not affected in *lrx3/4/5* (Fig. S5). Furthermore, VA-TIRFM revealed that FER-GFP transiently

accumulates in dynamic foci, independently of *LRX3/4/5* (Fig. S6, Movie S4). Together,

these results suggest that LRX3/4/5 do not prevent RALF association with FER to modulate

PTI. Moreover, our results suggest that active monitoring by the proposed cell wall integrity

sensors FER and LRXs regulates plasma membrane nanoscale dynamics of RKs.

The ability of LRX3/4/5 to associate with RALF23 *in planta* (Zhao *et al.*, 2018) prompted us to

test if LRX3/4/5 are required for RALF23 responsiveness. Indeed, *LRX3, LRX4* and *LRX5*

were required for RALF23-induced inhibition of ROS production upon elf18 treatment (Fig.

S7A). Similarly, we observed a decrease in the inhibition of seedlings growth triggered by

RALF23 in *lrx3/4/5* compared to WT (Fig. S7B). Altogether, these data show that LRX3/4/5

contribute to RALF23 responsiveness (Fig. S7C), and that LRXs and FER have analogous

functions in regulating PTI.

We next asked if FER and LRX3/4/5 form a complex. For this, we made use of a deleted

version of LRX4 lacking its extensin domain (LRX4$^{\Delta E}$), previously used to assess protein

complex formation (Dünser *et al.*, 2019; Herger *et al.*, 2020). Consistent with previous reports

on transient expression assays, (Dünser *et al.*, 2019; Herger *et al.*, 2020), co-

immunoprecipitation experiments with stably transgenic Arabidopsis showed that FER

associates constitutively with LRX4$^{\Delta E}$-FLAG, and that RALF23 treatment does not modulate

this association (Fig. S8). This suggests that FER-LRX complex-mediated direct monitoring

of the cell wall (Dünser *et al.*, 2019; Herger *et al.*, 2019) is not regulated by RALF23. In

agreement with structural and biochemical analyses of RALF-binding by CrRLK1L/LLGs and

LRXs (Moussu *et al.*, 2020), FER-LLG1 and LRX3/4/5 may form distinct RALF23 receptor

complexes. As in the cases of pollen tube and root hair growth and integrity (Ge *et al.*, 2017;

Mecchia *et al.*, 2017; Moussu *et al.*, 2020; Dünser *et al.*, 2019; Herger *et al.*, 2020), future

investigations are thus needed to understand the exact molecular link between RALF-binding

LRXs and CrRLK1s.

214   Functional dichotomy of FER and LRXs in regulating growth and immunity

In line with previous reports our data shows that FER and LRXs can form a complex (Fig. S8;

Dünser *et al.*, 2019; Herger *et al.*, 2019). Moreover, they are known to associate with the cell

wall (Baumberger, Ringli and Keller, 2001; Feng *et al.*, 2018), and are proposed to relay co-

jointly its properties (Dünser *et al.*, 2019; Herger *et al.*, 2019). We thus asked if direct cell

wall sensing underlies FER and LRXs function in PTI. In the context of growth and cell

expansion, plants overexpressing LRX4$^{\Delta E}$, a truncated version of LRX4, are phenotypically

reminiscent of *lrx3/4/5* triple mutants and the *fer-4* single mutant (Dünser *et al.*, 2019). This

dominant negative effect is proposed to be caused by competition of the overexpressed

truncated LRX4$^{\Delta E}$ with endogenous LRXs and consequent loss of cell wall anchoring (Dünser

et al., 2019). Similarly, overexpression of LRX1$^{\Delta E}$ inhibits root hair elongation, phenocopying

loss of function of *LRX1* and *LRX2* (Herger *et al.*, 2020). By contrast, we observed that

LRX4$^{\Delta E}$ does not affect flg22-induced interaction between FLS2 and BAK1 (Fig. S9A). In

good agreement with this notion, overexpression of LRX4$^{\Delta E}$ did not affect flg22 nor elf18-

induced ROS production (Fig. S9B-C). To corroborate these results, we tested inhibition of

root growth triggered by flg22 treatment. Consistent with the positive role of FER and

LRX3/4/5 in PTI, we observed that *fer-4* and *lrx3/4/5* are hyposensitive to flg22 treatment

(Fig. S9D). By contrast, overexpression of LRX4$^{\Delta E}$ did not affect inhibition of root growth by

flg22 (Fig. S9D). In addition, we observed that LRX4$^{\Delta E}$ does not impact RALF23

responsiveness (Fig. S9E). Altogether, these data suggest that the function of LRX3/4/5 in

PTI is distinct from their role during growth.

The ectodomain of FER contains two malectin-like domains, malA and malB (Fig. 3A), which

share homology with malectin, a carbohydrate-binding protein from *Xenopus laevis* (Boisson-

Dernier, Kessler and Grossniklaus, 2011). Despite lacking the canonical carbohydrate-

binding site of malectin (Moussu *et al.*, 2018; Xiao *et al.*, 2019), malA and malB have been

proposed to bind pectin *in vitro* (Feng *et al.*, 2018) and FER-mediated cell wall-sensing

regulates morphogenesis (Duan *et al.*, 2010; Lin *et al.*, 2018). To investigate if direct cell

wall-sensing underlies FER`s function in regulating PTI, we used transgenic lines expressing

a FER truncated mutant lacking the malA-domain C-terminally fused to YFP (FER$^{\Delta malA}$-YFP)

in the *fer-4* mutant background (Fig. 3B). We observed that FER$^{\Delta malA}$-YFP did not

complement cell shape or root hair elongation defect of *fer-4* (Fig. 3C-D), emphasizing the

importance of malA in FER-regulated cell morphogenesis. In contrast, immuno-precipitation

assays showed that FER$^{\Delta malA}$-YFP fully complements flg22-induced complex formation

between endogenous FLS2 and BAK1 (Fig. 3E) as well as ROS production in response to

flg22 and elf18 (Fig. 3F-G). Altogether, these data suggest that malA-mediated cell wall

sensing underlies specific function(s) of FER in regulating growth and cell morphology, but is

dispensable for FER´s role in PTI. Interestingly, we observed that expression of FER$^{\Delta malA}$-

YFP restores inhibition of growth triggered by RALF23, suggesting that malB is sufficient for

RALF responsiveness (Fig. 3H), as suggested by its physical interaction with RALF23 (Xiao

et al., 2019). While we cannot formally exclude the implication of pectin-binding by malB in

regulating immunity, the contrasted context-dependent functionality of FER$^{\Delta malA}$-YFP

suggests that FER´s function in PTI is primarily mediated by RALF perception. Altogether,

our data indicate molecular and functional dichotomy of FER and LRXs in regulating growth

and immunity.

RALF23 alters FLS2 and BAK1 organization and function through active FER signaling.

We next asked if RALF23 activity is mediated by active FER signaling. We used a kinase-dead mutant, FER^{K565R} C-terminally fused to GFP, expressed in *fer* knock-out backgrounds and selected lines accumulating it to level similar to endogenous FER in WT background (Fig. S10 and S11; Chakravorty, Yu and Assmann, 2018)). Interestingly, we observed that FER^{K565R}-GFP complements *fer*'s defect in FLS2-BAK1 complex formation (Fig. S10A) and PAMP-induced ROS production (Fig. S10B-C). In contrast, we observed that inhibition of FLS2-BAK1 complex formation by RALF23 depends on FER kinase activity (Fig. S11B). Similarly, inhibition of elf18-induced ROS production and seedlings growth inhibition by RALF23 depended on FER kinase activity (Fig. S11C). Overall, these data show that inhibition by RALF23 is mediated by active FER signaling while FER's positive role in immune signaling is kinase activity-independent.

We next asked if inhibition of FLS2-BAK1 complex formation by RALF23 correlates with a modulation of FLS2 or BAK1 nanoscale organization. VA-TIRFM imaging showed an increase of FLS2-GFP mobility and an alteration of FLS2-GFP nanodomain organization within minutes of RALF23 treatment (Fig. S12 and S13, Movie S7, imaging performed 2 to 30 min post-treatment Fig. S14). Similarly, we observed that RALF23 treatment affects BAK1-mCherry nanoscale organization (Fig. 5, Fig. S14, Movie S2). These data suggest that RALF23 perception leads to rapid modification of FLS2 and BAK1 membrane organization and thereby potentially inhibits their association. It will be important in the future to identify the components mediating RALF23 signaling and modification of FLS2 and BAK1 nanoscale dynamics.

In sum, our study unravels the regulation of FLS2 and BAK1 nanoscale organization by the RALF receptors FER and LRX3/4/5. The function of RALF receptors in other processes might similarly rely on the regulation of RK nanoscale dynamics, and the identification of the corresponding regulated RKs is an exciting prospect for future investigation.

ACKNOWLEDGMENTS

We thank all present and past members of the Zipfel laboratory for fruitful discussions and comments on the manuscript. We thank the members of the Grossniklaus, Ringli, Sanchez-Rodriguez and Keller laboratories for sharing results and comments during our stimulating CCWI meetings. We thank Vera Gorelova, Yvon Jaillais and Birgirt Kemmerling for comments on the manuscript. This research was funded by the Gatsby Charitable Foundation (C.Z.), the University of Zürich (C.Z.), the European Research Council under the Grant Agreements 309858 and 773153 (grants PHOSPHinnATE and IMMUNO-PEPTALK to C.Z.) and 639678 (grant AuxinER to J.K-V), the Swiss National Science Foundation (grant

no. 31003A_182625 to C.Z. and 31003A_166577/1 to C.R.), and the Austrian science fund
(FWF; P 33044 to J.K-V). J.G., C.M.F. and T.A.D. were supported by Long-Term Fellowships
from the European Molecular Biology Organization (EMBO) (numbers 438-2018, 512-2019
and 100-2017, respectively), while M.S. was supported by a post-doctoral fellowship (STE
2448/1) from the Deutsche Forschungsgemeinschaft (DFG) and K.D. by a doctoral fellowship
from the Austrian Academy of Sciences (ÖAW). We thank Sarah Assman for providing
segregating lines of *fer-4*/pFER::FER^{K565R}-GFP, Sacco de Vries for kindly providing Col-
0/pBAK1::BAK1-mCherry line and Silke Robatzek for Col-0/pFLS2::FLS2-GFP lines, and
Nana Keinath for kindly offering *fer-2*/pFER::FER^{K565R}-GFP lines.

REFERENCES

Abarca, A., Franck, C. M. and Zipfel, C. (2021) 'Family-wide evaluation of RAPID
ALKALINIZATION FACTOR peptides', *Plant Physiology*. doi: 10.1093/PLPHYS/KIAB308.

Baumberger, N., Ringli, C. and Keller, B. (2001) 'The chimeric leucine-rich repeat/extensin
cell wall protein LRX1 is required for root hair morphogenesis in Arabidopsis thaliana', *Genes
and Development*, 15(9), pp. 1128–1139. doi: 10.1101/gad.200201.

Boisson-Dernier, A., Kessler, S. A. and Grossniklaus, U. (2011) 'The walls have ears: The
role of plant CrRLK1Ls in sensing and transducing extracellular signals', *Journal of
Experimental Botany*, pp. 1581–1591. doi: 10.1093/jxb/erq445.

Bücherl, C. A. *et al.* (2013) 'Visualization of BRI1 and BAK1(SERK3) Membrane Receptor
Heterooligomers during Brassinosteroid Signaling', *Plant Physiology*, 162(4), p. 1911. doi:
10.1104/PP.113.220152.

Bücherl, C. A. *et al.* (2017) 'Plant immune and growth receptors share common signalling
components but localise to distinct plasma membrane nanodomains', *eLife*, 6. doi:
10.7554/eLife.25114.

Chakravorty, D., Yu, Y. and Assmann, S. M. (2018) 'A kinase-dead version of FERONIA
receptor-like kinase has dose-dependent impacts on rosette morphology and RALF1-
mediated stomatal movements', *FEBS Letters*, 592(20), pp. 3429–3437. doi: 10.1002/1873-
3468.13249.

Chinchilla, D. *et al.* (2007) 'A flagellin-induced complex of the receptor FLS2 and BAK1
initiates plant defence', *Nature*, 448(7152), pp. 497–500. doi: 10.1038/nature05999.

Couto, D. and Zipfel, C. (2016) 'Regulation of pattern recognition receptor signalling in
plants', *Nature Reviews Immunology*, 16, pp. 537–552. doi: 10.1038/nri.2016.77.

Daněk, M. *et al.* (2020) 'Cell wall contributes to the stability of plasma membrane
nanodomain organization of *Arabidopsis thaliana* FLOTILLIN2 and HYPERSENSITIVE
INDUCED REACTION1 proteins', *The Plant Journal*, 101(3), pp. 619–636. doi:
10.1111/tpj.14566.

Duan, Q. *et al.* (2010) 'FERONIA receptor-like kinase regulates RHO GTPase signaling of
root hair development', *Proceedings of the National Academy of Sciences of the United
States of America*, 107(41), pp. 17821–17826. doi: 10.1073/pnas.1005366107.

Dünser, K. *et al.* (2019) 'Extracellular matrix sensing by FERONIA and
Leucine-Rich Repeat Extensins controls vacuolar expansion during cellular elongation in
Arabidopsis thaliana', *The EMBO Journal*, 38(7). doi: 10.15252/embj.2018100353.

Feng, W. *et al.* (2018) 'The FERONIA Receptor Kinase Maintains Cell-Wall Integrity during
Salt Stress through Ca 2+ Signaling', *Current Biology*, 28, pp. 666-675.e5. doi:
10.1016/j.cub.2018.01.023.

Feraru, E. *et al.* (2011) 'PIN polarity maintenance by the cell wall in Arabidopsis', *Current
Biology*, 21(4), pp. 338–343. doi: 10.1016/j.cub.2011.01.036.

Franck, C. M., Westermann, J. and Boisson-Dernier, A. (2018) 'Plant Malectin-Like Receptor
Kinases: From Cell Wall Integrity to Immunity and Beyond', *Annual Review of Plant Biology*,
69(1), pp. 301–328. doi: 10.1146/annurev-arplant-042817-040557.

Ge, Z. *et al.* (2017) 'Arabidopsis pollen tube integrity and sperm release are regulated by
RALF-mediated signaling', *Science*, 358(6370), pp. 1596–1600. doi:
10.1126/science.aao3642.

Göhre, V. *et al.* (2008) 'Plant Pattern-Recognition Receptor FLS2 Is Directed for Degradation
by the Bacterial Ubiquitin Ligase AvrPtoB', *Current Biology*, 18(23), pp. 1824–1832. doi:
10.1016/j.cub.2008.10.063.

Gómez-Gómez, L. and Boller, T. (2000) 'FLS2: An LRR receptor-like kinase involved in the
perception of the bacterial elicitor flagellin in Arabidopsis', *Molecular Cell*, 5(6), pp. 1003–
1011. doi: 10.1016/S1097-2765(00)80265-8.

Gonneau, M. *et al.* (2018) 'Receptor Kinase THESEUS1 Is a Rapid Alkalinization Factor 34
Receptor in Arabidopsis', *Current Biology*, 28(15), pp. 2452-2458.e4. doi:
10.1016/j.cub.2018.05.075.

Gronnier, J. *et al.* (2017) 'Structural basis for plant plasma membrane protein dynamics and
organization into functional nanodomains', *eLife*, 6, pp. 1–24. doi: 10.7554/eLife.26404.

Gronnier, J. *et al.* (2018) 'Divide and Rule: Plant Plasma Membrane Organization', *Trends in
Plant Science*, 23(10). doi: 10.1016/j.tplants.2018.07.007.

Gronnier, J. *et al.* (2019) 'Mechanisms governing subcompartmentalization of biological
membranes', *Current Opinion in Plant Biology*, 52. doi: 10.1016/j.pbi.2019.08.003.

Grossmann, G. *et al.* (2008) 'Plasma membrane microdomains regulate turnover of transport
proteins in yeast', *Journal of Cell Biology*, 183(6), pp. 1075–1088. doi:
10.1083/jcb.200806035.

Haruta, M. *et al.* (2014) 'A peptide hormone and its receptor protein kinase regulate plant cell
expansion', *Science*, 343(6169), pp. 408–411. doi: 10.1126/science.1244454.

Herger, A. *et al.* (2019) 'Leucine-Rich Repeat Extensin Proteins and Their Role in Cell Wall
Sensing', *Current Biology*. Cell Press, pp. R851–R858. doi: 10.1016/j.cub.2019.07.039.

Herger, A. *et al.* (2020) 'Overlapping functions and protein-protein interactions of LRR-
extensins in Arabidopsis', *PLOS Genetics*. Edited by G. K. Muday, 16(6), p. e1008847. doi:
10.1371/journal.pgen.1008847.

Hohmann, U., Lau, K. and Hothorn, M. (2017) 'The Structural Basis of Ligand Perception and
Signal Activation by Receptor Kinases', *Annu. Rev. Plant Biol*, 68, pp. 109–37. doi:
10.1146/annurev-arplant-042916.

Hutten, S. J. *et al.* (2017) 'Visualization of BRI1 and SERK3/BAK1 Nanoclusters in
Arabidopsis Roots', *Plos One*, 12(1), p. e0169905. doi: 10.1371/journal.pone.0169905.

Jacobson, K., Liu, P. and Lagerholm, B. C. (2019) 'The Lateral Organization and Mobility of
Plasma Membrane Components', *Cell*. Cell Press, pp. 806–819. doi:
10.1016/j.cell.2019.04.018.

Jaillais, Y. and Ott, T. (2020) 'The Nanoscale Organization of the Plasma Membrane and Its

Importance in Signaling: A Proteolipid Perspective', *Plant Physiology*, 182(4), pp. 1682–1696. doi: 10.1104/pp.19.01349.

Jaqaman, K. *et al.* (2008) 'Robust single-particle tracking in live-cell time-lapse sequences', *Nature Methods*, 5(8), pp. 695–702. doi: 10.1038/nmeth.1237.

Kadota, Y. *et al.* (2014) 'Direct Regulation of the NADPH Oxidase RBOHD by the PRR-Associated Kinase BIK1 during Plant Immunity', *Molecular Cell*, 54, pp. 43–55. doi: 10.1016/j.molcel.2014.02.021.

Liang, P. *et al.* (2018) 'Symbiotic root infections in Medicago truncatula require remorin-mediated receptor stabilization in membrane nanodomains', *Proceedings of the National Academy of Sciences of the United States of America*, 115(20), pp. 5289–5294. doi: 10.1073/pnas.1721868115.

Lin, W. *et al.* (2018) 'FERONIA's sensing of cell wall pectin activates ROP GTPase signaling in Arabidopsis', *bioRxiv*, 35(June), p. 269647. doi: 10.1101/269647.

Liu, C. *et al.* (2021) 'Pollen PCP-B peptides unlock a stigma peptide–receptor kinase gating mechanism for pollination', *Science*, 372(6538), pp. 171–175. doi: 10.1126/SCIENCE.ABC6107.

Martinière, A. *et al.* (2012) 'Cell wall constrains lateral diffusion of plant plasma-membrane proteins.', *Proceedings of the National Academy of Sciences of the United States of America*, 109(31), pp. 12805–10. doi: 10.1073/pnas.1202040109.

McKenna, J. F. *et al.* (2019) 'The cell wall regulates dynamics and size of plasma-membrane nanodomains in Arabidopsis', *Proceedings of the National Academy of Sciences of the United States of America*, 116(26), pp. 12857–12862. doi: 10.1073/pnas.1819077116.

Mecchia, M. A. *et al.* (2017) 'RALF4/19 peptides interact with LRX proteins to control pollen tube growth in Arabidopsis', *Science*, 358(6370), pp. 1600–1603. doi: 10.1126/science.aao5467.

Moussu, S. *et al.* (2018) 'Crystal structures of two tandem malectin-like receptor kinases involved in plant reproduction', *research papers Acta Cryst*, 74, pp. 671–680. doi: 10.1107/S205979831800774X.

Moussu, S. *et al.* (2020) 'Structural basis for recognition of RALF peptides by LRX proteins during pollen tube growth', *Proceedings of the National Academy of Sciences of the United States of America*, 117(13), pp. 7494–7503. doi: 10.1073/pnas.2000100117.

Perraki, A. *et al.* (2018) 'Phosphocode-dependent functional dichotomy of a common co-receptor in plant signalling', *Nature*, 561(7722), pp. 248–252. doi: 10.1038/s41586-018-0471-x.

Roux, M. *et al.* (2011) 'The Arabidopsis leucine-rich repeat receptor-like kinases BAK1/SERK3 and BKK1/SERK4 are required for innate immunity to hemibiotrophic and biotrophic pathogens', *Plant Cell*, 23(6), pp. 2440–2455. doi: 10.1105/tpc.111.084301.

Schindelin, J. *et al.* (2012) 'Fiji: an open-source platform for biological-image analysis', *Nature Methods 2012 9:7*, 9(7), pp. 676–682. doi: 10.1038/nmeth.2019.

Schulze, B. *et al.* (2010) 'Rapid heteromerization and phosphorylation of ligand-activated plant transmembrane receptors and their associated kinase BAK1', *Journal of Biological Chemistry*, 285(13), pp. 9444–9451. doi: 10.1074/jbc.M109.096842.

Schwessinger, B. *et al.* (2011) 'Phosphorylation-Dependent Differential Regulation of Plant Growth, Cell Death, and Innate Immunity by the Regulatory Receptor-Like Kinase BAK1', *PLoS Genetics*. Edited by D. S. Guttman, 7(4), p. e1002046. doi: 10.1371/journal.pgen.1002046.

Stegmann, M. *et al.* (2017) 'The receptor kinase FER is a RALF-regulated scaffold controlling
plant immuneTraduire I'll send you everything by tomorrow night that you could send to all
authors by the end of this week. 102/5000 Je vous enverrai tout demain soir que vous
pourriez envoy', 4(January), pp. 287–289. doi: 10.1126/science.aal2541.

Sun, Y. *et al.* (2013) 'Structural basis for flg22-induced activation of the Arabidopsis FLS2-
BAK1 immune complex', *Science*, 342(6158), pp. 624–628. doi: 10.1126/science.1243825.

Tinevez, J. Y. *et al.* (2017) 'TrackMate: An open and extensible platform for single-particle
tracking', *Methods*, 115, pp. 80–90. doi: 10.1016/j.ymeth.2016.09.016.

Tran, T. M. *et al.* (2020) 'The bacterial quorum sensing signal DSF hijacks Arabidopsis
thaliana sterol biosynthesis to suppress plant innate immunity', *Life Science Alliance*, 3(10).
doi: 10.26508/LSA.202000720.

Wagner, T. *et al.* (2017) 'Classification and Segmentation of Nanoparticle Diffusion
Trajectories in Cellular Micro Environments', *PLOS ONE*. Edited by B. Xu, 12(1), p.
e0170165. doi: 10.1371/journal.pone.0170165.

Xiao, Y. *et al.* (2019) 'Mechanisms of RALF peptide perception by a heterotypic receptor
complex', *Nature*, 572(7768), pp. 270–274. doi: 10.1038/s41586-019-1409-7.

Yu, X. *et al.* (2017) 'From Chaos to Harmony: Responses and Signaling upon Microbial
Pattern Recognition', *Annual Review of Phytopathology*, 55(1), pp. 109–137. doi:
10.1146/annurev-phyto-080516-035649.

Zhao, C. *et al.* (2018) 'Leucine-rich repeat extensin proteins regulate plant salt tolerance in
Arabidopsis', *Proceedings of the National Academy of Sciences of the United States of
America*, 115(51), pp. 13123–13128. doi: 10.1073/pnas.1816991115.

Zipfel, C. *et al.* (2006) 'Perception of the Bacterial PAMP EF-Tu by the Receptor EFR
Restricts Agrobacterium-Mediated Transformation', *Cell*, 125(4), pp. 749–760. doi:
10.1016/j.cell.2006.03.037.

FIGURE LEGENDS

Fig. 1 ┆ FER regulates the nanoscale organization of FLS2-GFP and BAK1-mCherry

A and E. FLS2-GFP and BAK1-mCherry nanodomain organization. Pictures are maximum projection images of FLS2-GFP (**A**) and BAK1-mCherry (**E**) in Col-0 and *fer-4* cotyledon epidermal cells.

B and F. Representative kymograph showing lateral organization of FLS2-GFP (**B**) and BAK1-mCherry (**F**) overtime in Col-0 and *fer-4*.

C and G. Quantification of FLS2-GFP (**C**) and BAK1-mCherry (**G**) spatial clustering index. Red crosses and red horizontal lines show mean and SEM, P values reports non-parametric Mann-Whitney test. Similar results were obtained in three independent experiments.

D and H. Proposed graphical summary of our observations for FLS2-GFP (**D**) and BAK1-mCherry (**H**) nanoscale dynamics.

Fig. 2 ┆ LRX3, LRX4 and LRX5 regulate PTI and BAK1-mCherry organization.

A. flg22-induced FLS2-BAK1 complex formation. Immunoprecipitation of FLS2 in Arabidopsis Col-0 and *lr3/4/5* seedlings either untreated or treated with 100 nM flg22 for 10 min. Blot stained with CBB is presented to show equal loading. Western blots were probed with α-FLS2, α-BAK1, or α-FER antibodies. Similar results were obtained in at least three independent experiments.

B-C. ROS production after elicitation with 100 nM elf18 (**B**), or 100 nM flg22 (**C**). Values are means of total photon counts over 40 min. Red crosses and red horizontal lines denote mean and SEM, *n* = 32. Conditions which do not share a letter are significantly different in Dunn's multiple comparison test (p< 0.0001).

D. BAK1-mCherry nanodomain organization. Pictures are maximum projection images (20 TIRFM images obtained at 2.5 frames per second) of BAK1-mCherry in Col-0, *fer-4* and *lrx3/4/5* cotyledon epidermal cells.

E. Representative kymograph showing lateral organization of BAK1-mCherry overtime in Col-0, *fer-4* and *lrx3/4/5*.

F. Quantification of BAK1-mCherry spatial clustering index, red crosses and red horizontal lines show mean and SEM. Conditions which do not share a letter are significantly different in Dunn's multiple comparison test (p< 0.001). Similar results were obtained in three independent experiments.

G. Proposed graphical summary of our observations for BAK1-mCherry nanoscale dynamics in *lrx3/4/5*.

Fig. 3 ┊ FER malectin A domain regulates cell morphogenesis not PTI

A. Proposed graphical representation of RALF23 perception by FER-LLG1 complex.

B. Morphology of 4-week-old Arabidopsis plants, scale bar indicates 5 cm.

C-D. Confocal microscopy pictures of 5-day-old seedlings cotyledon (**C**) and root (**D**) stained
with propidium iodide. Similar results were obtained in at least three independent
experiments.

E. flg22-induced FLS2-BAK1 complex formation. Immunoprecipitation of FLS2 in Arabidopsis
Col-0, *fer-4*, and *fer-4*/p35S::FER$^{\Delta MalA}$-YFP seedlings that were either untreated or treated
with 100 nM flg22 for 10 min. Blot stained with Coomassie brilliant blue (CBB) is presented to
show equal loading. Western blots were probed with α-FLS2, α-BAK1, or α-FER antibodies.
Similar results were obtained in at least three independent experiments.

F-G. ROS production after elicitation with 100 nM flg22 (**F**), or 100 nM elf18 (**G**). Values are
means of total photon counts over 40 min, $n = 8$. Red crosses and red horizontal lines denote
mean and SEM, respectively. Conditions which do not share a letter are significantly different
in Dunn's multiple comparison test ($p < 0.0001$).

H. Fresh weight of 12-day-old seedlings grown in the absence (mock) or presence of 1 µM of
RALF23 peptide. Fresh weight is expressed as relative to the control mock condition. Similar
results were obtained in at least three independent experiments. Conditions which do not
share a letter are significantly different in Dunn's multiple comparison test ($p < 0.001$).

Fig. 4 ┊ RALF23 perception regulates BAK1-mCherry organization.

A. BAK1-mCherry nanodomain organization (pBAK1::BAK1-mCherry). Pictures are
maximum projection images of BAK1-mCherry in Col-0 and *fer-4* cotyledon epidermal cells
with or without 1 µM RALF23 treatment (2 to 30 min).

B. Representative kymograph showing lateral organization of BAK1-mCherry overtime in
Col-0 and *fer-4* with or without 1 µM RALF23 treatment.

C. Quantification of BAK1-mCherry spatial clustering index, red crosses and red horizontal
lines show mean and SEM. Conditions which do not share a letter are significantly different
in Dunn's multiple comparison test ($p < 0.001$). Similar results were obtained in three
independent experiments.

D. Proposed graphical summary of our observations for BAK1-mCherry nanoscale dynamics
upon RALF23 treatment.

Fig. S1 ┊ Analysis of FLS2-GFP single particle dynamics in *fer-4*.

A. Representative images of FLS2-GFP single-particles tracked in Col-0 and *fer-4* cotyledon
epidermal cells of 5-day-old seedlings.

B. Quantification of FLS2-GFP diffusion coefficient (D). Similar results were obtained in three

independent experiments.

Fig. S2 ⦙ Analysis of FLS2-GFP organization and dynamics in *fer-2.*

A. Representative images of FLS2-GFP single-particles tracked in Col-0 and *fer-2* cotyledon

epidermal cells of 5-day-old seedlings.

B. Quantification of FLS2-GFP diffusion coefficient (D). Similar results were obtained in three

independent experiments.

C. Pictures are maximum projection images of FLS2-GFP in Col-0 and *fer-2* cotyledon

epidermal cells.

D. Quantification of FLS2-GFP spatial clustering index in Col-0 and *fer-2*. Red crosses and

red horizontal lines show mean and SEM, P values reports non-parametric Mann-Whitney

test. Similar results were obtained in three independent experiments.

Fig. S3 ⦙ FLS2-GFP accumulation at the PM is altered in *fer* mutants.

A-D. Representative confocal microscopy pictures of FLS2-GFP driven by its native promoter

in Col-0 and *fer-2* (**A**) and *fer-4* (**C**) cotyledon epidermal cells of 5-day-old seedlings. Scale

bar indicate 2 µm. Quantification of FLS2-GFP fluorescence intensity at the plasma

membrane in Col-0 and *fer-2* (**B**) and *fer-4* (**D**) cotyledon epidermal cells of 5-day-old

seedlings. Two-way student's t test. Similar results were obtained in at least three

independent experiments.

Fig. S4 ⦙ Subcellular localization of BAK1-mCherry in *fer-4* and *lrx3/4/5.*

A. Representative confocal microscopy pictures of BAK1-mCherry driven by its native

promoter in Col-0, *fer-4* and *lrx3/4/5* cotyledon epidermal cells of 5-day-old seedlings. Scale

bar indicates 20 µm.

B. Quantification of BAK1-mCherry fluorescence intensity at the plasma membrane. Letters

indicate a Dunn's multiple comparison statistical test.

Fig. S5 ⦙ *LRX3, LRX4* and *LRX5* are dispensable for FER plasma membrane

localization and accumulation.

A. Accumulation of endogenous FER detected by western blot. Total proteins were extracted

from 2-week-old seedlings. Blot stained with CBB is presented to show equal loading.

B-C. Representative confocal microscopy pictures of FER-GFP driven by its native promoter in *fer-4* and *lrx3/4/5* cotyledon epidermal cells (**B**) and root hair (**C**) of 5-day-old seedlings. Scale bar indicates 20 μm.

Fig. S6 ⫶ *LRX3, LRX4* and *LRX5* are dispensable for FER-GFP nanoscale organization.

A. FER-GFP nanodomain organization. Pictures are maximum projection images (20 TIRFM images obtained at 5 frames per second) of FER-GFP in *fer-4* and *lrx3/4/5* cotyledon epidermal cells.

B. Representative kymograph showing lateral organization of FER-GFP overtime in *fer-4* and *lrx3/4/5*.

C. Quantification of FER-GFP spatial clustering index. Red crosses and red horizontal lines show mean and SEM. Conditions which do not share a letter are significantly different in Dunn's multiple comparison test ($p < 0.001$). Similar results were obtained in three independent experiments.

D. Proposed graphical summary of our observations for FER-GFP nanoscale dynamics.

Fig. S7 ⫶ LRX3, LRX4 and LRX5 contribute to RALF23 responsiveness.

A. ROS production in Col-0 and *lr3/4/5* leaf discs treated with 100 nM elf18 and with or without 1 μM RALF23 co-treatment in 2 mM MES–KOH pH 5.8. Values are means of total photon counts over 40 min. Red crosses and red horizontal lines show mean and SEM, n = 12. Conditions which do not share a letter are significantly different in Dunn's multiple comparison test ($p < 0.0001$).

B. Fresh weight of 12-day-old seedlings grown in the absence (mock) or presence of 1 μM of RALF23 peptide. Fresh weight is expressed as relative to the control mock condition. Similar results were obtained in at least three independent experiments. Conditions which do not share a letter are significantly different in Dunn's multiple comparison test ($p < 0.001$).

C. Proposed graphical representation of LRX3/4/5 as potential receptors for RALF23.

Fig. S8 ⫶ RALF23 does not modulate constitutive association between FER and LRX4.

Immunoprecipitation of LRX4$^{\Delta E}$-FLAG in Arabidopsis seedlings untreated or treated with 1 μM RALF23 for 10 min. Western blots were probed with α-FLAG or α-FER antibodies. Blot stained with CBB is presented to show equal loading. Similar results were obtained in at least three independent experiments.

Fig. S9 ⦙ Overexpression of LRX4$^{\Delta E}$ does not affect PTI.

A. flg22-induced FLS2-BAK1 complex formation. Immunoprecipitation of FLS2 in Arabidopsis
seedlings either untreated or treated with 100 nM flg22 for 10 min. Blot stained with CBB is
presented to show equal loading. Western blots were probed with α-FLS2 and α-BAK1
antibodies.

B-C. ROS production after elicitation with 100 nM flg22 (B), or 100 nM elf18 (C). Values are
means of total photon counts over 40 min. Red crosses and red horizontal lines denote mean
and SEM. Conditions which do not share a letter are significantly different in Dunn's multiple
comparison test ($p < 0.0001$). Similar results were obtained in at least three independent
experiments.

D. Root length of 6-day-old seedlings incubated for 3 days in liquid MS medium with or
without indicated concentration of flg22. Red crosses and red horizontal lines denote mean
and SEM. Conditions which do not share a letter are significantly different in Brown-Forsythe
and Welch ANOVA multiple comparison test ($p < 0.001$), n= 9-12 seedlings per condition.
Similar results were obtained in at least three independent experiments.

E. Fresh weight of 12-day-old seedlings grown in the absence (mock) or presence of 1 μM of
RALF23 peptide. Fresh weight is expressed as relative to the control mock condition. Similar
results were obtained in at least three independent experiments. Conditions which do not
share a letter are significantly different in Dunn's multiple comparison test ($p < 0.001$).

Fig. S10 ⦙ FER kinase activity is dispensable to support PTI signaling.

A. flg22-induced FLS2-BAK1 complex formation. Immunoprecipitation of FLS2 in Arabidopsis
Col-0, *fer-4*, and *fer-4*/pFER::FER^{KD}-GFP seedlings that were either untreated or treated with
100 nM flg22 for 10 min. Blot stained with Coomassie brilliant blue (CBB) is presented to
show equal loading. Western blots were probed with α-FLS2, α-BAK1, α-BAK1-pS612 or α-
FER antibodies. Similar results were obtained in at least three independent experiments.

B-C. ROS production after elicitation with 100 nM flg22 (**A**), or 100 nM elf18 (**B**). Values are
means of total photon counts over 40 min, *n* = 16. Red crosses and red horizontal lines
denote mean and SEM, respectively. Conditions which do not share a letter are significantly
different in Dunn's multiple comparison test ($p < 0.0001$).

Fig. S11 ⦙ Inhibition of PTI signaling by RALF23 requires FER kinase activity.

A. Accumulation of endogenous FER and FER^{K565R}-GFP detected by western blot. Total
proteins were extracted from 2-week-old seedlings. Blot stained with CBB is presented to
show loading.

B. flg22-induced FLS2-BAK1 complex formation. Immunoprecipitation of FLS2 in Arabidopsis
fer-2, and *fer-2*/pFER::FER^{K565R}-GFP seedlings that were either untreated or treated with 100
nM of flg22 and 1 µM RALF23 for 10 min. Western blots were probed with α-FLS2, α-BAK1
antibodies.

C. ROS production after elicitation with 100 nM elf18, with and without 1 µM RALF23 co-
treatment. Values are means of total photon counts over 40 min, n = 10. Red crosses and
red horizontal lines denote mean and SEM, respectively. Conditions which do not share a
letter are significantly different in Dunn's multiple comparison test ($p < 0.05$). Similar results
were obtained in at least three independent experiments.

Fig. S12 ⦙ Analysis of FLS2-GFP single particle dynamics upon RALF23 treatment.

A. Representative super-resolved images of FLS2-GFP single-particles tracked in Col-0 and
fer-2 cotyledon epidermal cells of 5-day-old seedlings with or without 1 µM RALF23
treatment.

B. and quantification of FLS2-GFP diffusion coefficient (D) in Col-0 and *fer-2* with or without
1 µM RALF23 treatment. Data represent analysis of *ca* 1800 to 2900 single particles
observed across 14 to 18 cells. Individual data point represents the mean diffusion coefficient
for each cell. Red crosses and red horizontal lines show mean and SEM. Conditions which
do not share a letter are significantly different in Dunn's multiple comparison test ($p < 0.0001$).
Similar results were obtained in at least three independent experiments.

Fig. S13 ⦙ Analysis of FLS2-GFP organization upon RALF23 treatment.

A. Pictures are maximum projection images (20 TIRFM images obtained at 20 frames per
second) of FLS2-GFP in Col-0 and *fer-2* cotyledon epidermal cells with or without 1 µM
RALF23 treatment.
B. Representative kymograph showing lateral organization of FLS2-GFP overtime in Col-0
and *fer-2* cotyledon epidermal cells with or without 1 µM RALF23 treatment.
C. Quantification of FLS2-GFP spatial clustering index. Red crosses and red horizontal lines
show mean and SEM. Conditions which do not share a letter are significantly different in
Dunn's multiple comparison test ($p < 0.0001$). Similar results were obtained in at least three
independent experiments.
663
Fig. S14 ⦙ Time-resolved analysis of the spatial clustering index.

Data points correspond to the average of the spatial clustering index measured over 5 min
windows for FLS2-GFP (**A**) and BAK1-mCherry (**B**) with or without 1 µM RALF23 treatment.
The first data point (5 min), corresponds to the value obtained between 2- and 3-min post

treatment. Acquisition approximately started 2 min after treatments, the time required for
sample mounting.

Fig. S15 ¦ Linear regression analysis of the relationship between the spatial clustering
index and fluorescence intensity.

Spatial clustering index is calculated as the ratio of the mean of the 5 % highest values to the
mean of 5 % lowest values of fluorescence intensity obtain on a plot line. Here, spatial
clustering index values were plotted against the mean fluorescence intensity of the
corresponding plot line for FLS2-GFP (**A**) and BAK1-mCherry (**B-C**). Linear regression
analyses show poor to no correlation between the spatial clustering index and fluorescence
intensity (R^2 ranking from 0.14 and 0.015).

Movie S1 ¦ TIRFM imaging of FLS2-GFP in Col-0 and *fer-4*.

Representative stream images acquisition of FLS2-GFP particles observed at the surface of
5-day-old cotyledon epidermal cells by TIRF microscopy at 5 frame per second.

Movie S2 ¦ TIRFM imaging of BAK1-mCherry in Col-0 and *fer-4* with or without RALF23
treatment.

Representative stream images acquisition of BAK1-mCherry observed at the surface of 5-
day-old cotyledon epidermal cells by TIRF microscopy at 2.5 frame per second, in Col-0 (A
and B) and *fer-4* (C and D) with (B and D) or without (A and C) 1 μM RALF23 treatment.
Scale bar indicates 2 μm.

Movie S3 ¦ TIRFM imaging of BAK1-mCherry in Col-0 and *lrx3/4/5*.

Representative stream images acquisition of BAK1-mCherry observed at the surface of 5-
day-old cotyledon epidermal cells by TIRF microscopy at 2.5 frame per second, in Col-0 (A)
and *lrx3/4/5* (B). Scale bar indicates 2 μm.

Movie S4 ¦ TIRFM imaging of FER-GFP in *fer-4* and *fer-4*;*lrx3/4/5*.

Representative stream images acquisition of FER-GFP observed at the surface of 5-day-old
cotyledon epidermal cells by TIRF microscopy at 5 frame per second, in *fer-4* (A) and *fer-*
4;lrx3/4/5 (B). Scale bar indicates 2 μm.

Movie S5 ¦ TIRFM imaging of FLS2-GFP in Col-0 with or without RALF23 treatment.

Representative stream images acquisition of FLS2-GFP particles observed at the surface of
5-day-old cotyledon epidermal cells by TIRF microscopy at 5 frame per second. Scale bar
indicates 2 μm.

Movie S6 ¦ TIRFM imaging of FLS2-GFP in *fer-4* with or without RALF23 treatment.

Representative stream images acquisition of FLS2-GFP particles observed at the surface of
5-day-old cotyledon epidermal cells by TIRF microscopy at 5 frame per second. Scale bar
indicate 2 μm.

Movie S7 ⦂ TIRFM imaging of FLS2-GFP in Col-0 and *fer-2* with or without RALF23
treatment.

Representative stream images acquisition of FLS2-GFP particles observed at the surface of
5-day-old cotyledon epidermal cells by TIRF microscopy at 20 frame per second. Scale bar
indicates 2 μm

MATERIALS AND METHODS

Plant materials and growth

Arabidopsis thaliana ecotype Columbia (Col-0) was used as WT control. The *fer-4, fer-*
*4/*pFER::FER-GFP (Duan *et al.*, 2010), *fer-4/*pFER::FER^{KD}-GFP (Chakravorty, Yu and
Assmann, 2018), *fer-4/*p35S::FER$^{\Delta MalA}$-GFP (Lin *et al.*, 2018), Col- 0/pFLS2::FLS2-GFP
(Göhre *et al.*, 2008), *fer-2/*pFLS2::FLS2-GFP (Stegmann *et al.*, 2017), *lrx3/4/5*,
p35S::LRX4$^{\Delta E}$-Citrine and p35S::LRX4$^{\Delta E}$-FLAG (Dünser *et al.*, 2019) lines were previously
published. Col- 0/pFLS2::FLS2-GFP (Göhre *et al.*, 2008) was crossed with *fer-4* to obtain *fer-*
*4/*pFLS2::FLS2-GFP. Col-0/pBAK1::BAK1-mCherry (Bücherl *et al.*, 2013) was crossed with
fer-4 and *lrx3/4/*5 to obtain *fer-*4/pBAK1::BAK1-mCherry and *lrx3/4/5/*pBAK1::BAK1-mCherry.
*fer-4/*pFER::FER-GFP was crossed with *lrx3/4/5* to obtain *fer-4/lrx3/4/5;*pFER::FER-GFP. For
ROS burst assays, plants were grown in individual pots at 20-21 °C with a 10-h photoperiod
in environmentally controlled growth rooms. For seedling-based assays, seeds were surface-
sterilized using chlorine gas for 5 h and grown at 22 °C and a 16-h photoperiod on
Murashige and Skoog (MS) medium supplemented with vitamins, 1 % sucrose and 0.8 %
agar.

Synthetic peptides and chemicals

The flg22, elf18, and RALF23 peptides were synthesized by EZBiolab (United States) with a
purity of >95 %. All peptides were dissolved in sterile purified water.

ROS burst measurement

ROS burst measurements were performed as previously documented (Kadota *et al.*, 2014).
At least eight leaf discs (4 mm in diameter) per individual genotype were collected in 96-well
plates containing sterile water and incubated overnight. The next day, the water was
replaced by a solution containing 17 μg/mL luminol (Sigma Aldrich), 20 μg/mL horseradish

peroxidase (HRP, Sigma Aldrich) and the peptides in the appropriate concentration.
Luminescence was measured for the indicated time period using a charge-coupled device
camera (Photek Ltd., East Sussex UK). The effect of RALF23 on elf18-triggered ROS
production was performed as previously described (Stegmann *et al.*, 2017). Eight to ten leaf
discs per treatment and/or genotype were collected in 96-well plates containing water and
incubated overnight. The following day, the water was replaced by 75 µL of 2 mM MES-KOH
pH 5.8 to mimic the apoplastic pH. Leaf discs were incubated further for 4-5 h before adding
75 µL of a solution containing 40 µg/mL HRP, 1 µM L-O12 (Wako Chemicals, Germany) and
2X elicitor RALF peptide solution (final concentration 20 µg/mL HRP, 0.5 µM L-O12, 1x
elicitors). ROS production is displayed as the integration of total photon counts.

Root growth inhibition assay

Three-day-old Col-0, *fer-4*, *lrx3/4/5* and 35S::LRR4-Cit seedlings (n = 9-12) were transferred
for additional 3 days to 3 mL liquid ½ MS medium containing different concentrations (100
nM, 300 nM or 1 µM) of flg22 or the appropriate amount of solvent. The seedlings were then
placed on solid MS plates before scanning. Root length was measured using ImageJ.

Live cell imaging

For confocal microscopy and TIRF microscopy experiments, surface-sterilized seeds were
individually placed in line on square Petri dishes containing 1/2 MS 1 % sucrose, 0.8 %
phytoagar, stratified 2 d in the dark at 4 °C, then placed in a growth chamber at 22 °C and a
16-h photoperiod for 5 d. Seedlings were mounted between a glass slide and a coverslip in
liquid 1/2 MS, 1 % sucrose medium. To test the effect of RALF23 on FLS2-GFP dynamics
and nanodomain organization, seedlings were pre-incubated in 2 mM MES-KOH pH 5.8 for 3
to 4h prior treatment. Seedlings were image 2-30 min after treatment.

Confocal laser scanning microscopy (CLSM)

Confocal microscopy was performed using a Leica SP5 CLSM system (Leica, Wetzlar,
Germany) equipped with Argon, DPSS, He-Ne lasers, hybrid detectors and using a 63X 1.2
NA oil immersion objective. GFP was excited using 488 nm argon laser and emission
wavelengths were collected between 495 and 550 nm. mCherry was excited using 561 nm
He/Ne laser and emission wavelengths were collected between 570 and 640 nm. Propidium
iodide was imaged using 488 nm and 500-550 nm excitation and emission wavelengths,
respectively. In order to obtain quantitative data, experiments were performed using strictly
identical confocal acquisition parameters (*e.g.* laser power, gain, zoom factor, resolution, and
emission wavelengths reception), with detector settings optimized for low background and no
pixel saturation. Pseudo-color images were obtained using look-up-table (LUT) of Fiji
software (Schindelin *et al.*, 2012).

Total Internal Reflection Fluorescence (TIRF) microscopy

TIRF microscopy was performed using an inverted Leica GSD equipped with a 160x

objective (NA = 1.43, oil immersion), and an Andor iXon Ultra 897 EMCCD camera. Images

were acquired by illuminating samples with a 488 nm solid state diode laser set at 15 mW,

using a cube filter with an excitation filter 488/10 and an emission filter 535/50 for FLS2-GFP

and FER-GFP. optimum critical angle was determined as giving the best signal-to-noise.

Images time series were recorded at 20 frames per second (50 ms exposure time) for Figure

S11 and Figure S3; 5 frames per second for Figure 1 and Figure S6. To observe BAK1-

mCherry we could only use a 532 nm solid state diode laser (*c.a* 40% of maximum excitation

for mCherry), using a cube filter with an excitation filter 532/10 and an emission filter

600/100. To obtain a sufficient signal to noise ratio images time series were recorded at 2.5

frames per second (Figure 1, 2 and 5). Due to apparent high mobility of BAK1 and relatively

slow acquisition rate we couldn't asses with confidence the identity of fluorescent particles

from one time frame to another and therefore did not perform particle tracking analysis of

BAK1-mCherry.

Single particle tracking analysis

To analyse single particle tracking experiments, we used the plugin TrackMate 2.7.4

(Tinevez *et al.*, 2017) in Fiji (Schindelin *et al.*, 2012). Single particles were segmented frame-

by-frame by applying a LoG (Laplacian of Gaussian) filter and estimated particle size of 0.4

μm. Individual single particle were localized with sub-pixel resolution using a built-in

quadratic fitting scheme. Then, single particle trajectories were reconstructed using a simple

linear assignment problem (Jaqaman *et al.*, 2008) with a maximal linking distance of 0.4 μm

and without gap-closing. Only tracks with at least ten successive points (tracked for 500 ms)

were selected for further analysis. Diffusion coefficients of individual particles were

determined using TraJClassifier (Wagner *et al.*, 2017). For each particle, the slope of the first

four time points of their mean square displacement (MSD) plot was used to calculate their

diffusion coefficient according to the following equation: MSD= $(x-x_0)^2+(y-y_0)^2$ and D=MSD/4t,

where x0 and y0 are the initial coordinates, and x and y are the coordinates at any given

time, and t is the time frame.

Quantification of spatial clustering index

Genotype and/or treatment dependent variation in fluorescence intensity of FLS2-GFP and

fluorescence pattern of FLS2-GFP and BAK1-mCherry compromised the use of a unique set

of parameters to compute nanodomain size and density across the different experiments. To

uniformly quantify differences in membrane organization of both FLS2 and BAK1 across all

experiments we used the spatial clustering index which was shown to be largely insensitive

to variation in fluorescence intensity (Gronnier *et al.*, 2017). Quantifications were performed
as previously described (Gronnier *et al*., 2017). Briefly, fluorescence intensity was plotted
along an 8 µm long line on maximum projection TIRFM images, three plots were randomly
recorded per cell and, at least 8 cells per condition per experiment were analysed. For each
line plot, the spatial clustering index was calculated by dividing the mean of the 5 % highest
values by the mean of 5 % lowest values. Because the absence of correlation between
fluorescence intensity and spatial clustering index was assessed on confocal microscopy
images and for a single protein (Gronnier *et al.*, 2017), we decided to test if this was also the
case in our experimental conditions. Indeed, we consistently observed poor to no correlation
between variation in fluorescence intensity and values of spatial clustering index (Fig. S15).

Co-immunoprecipitation experiments

Twenty to thirty seedlings per plate were grown in wells of a 6-well plate for 2 weeks,
transferred to 2 mM MES-KOH, pH 5.8 and incubated overnight. The next day, flg22, (final
concentration 100 nM) and/or RALF23 (final concentration 1 µM) were added and incubated
for 10 min. Seedlings were then frozen in liquid N2 and subjected to protein extraction. To
analyse FLS2-BAK1 receptor complex formation, proteins were isolated in 50 mM Tris-HCl
pH 7.5, 150 mM NaCl, 10 % glycerol, 5 mM dithiothreitol, 1 % protease inhibitor cocktail
(Sigma Aldrich), 2 mM Na2MoO4, 2.5 mM NaF, 1.5 mM activated Na3VO4, 1 mM
phenylmethanesulfonyl fluoride and 0.5 % IGEPAL. For immunoprecipitations, α- rabbit
Trueblot agarose beads (eBioscience) coupled with α-FLS2 antibodies (Chinchilla *et al.,*
2007) or GFP-Trap agarose beads (ChromoTek) were used and incubated with the crude
extract for 3-4 h at 4 °C. Subsequently, beads were washed 3 times with wash buffer (50 mM
Tris-HCl pH 7.5, 150 mM NaCl, 1 mM phenylmethanesulfonyl fluoride, 0,1 % IGEPAL) before
adding Laemmli sample buffer and incubating for 10 min at 95 °C. Analysis was carried out
by SDS-PAGE and immunoblotting. To test the association between Flag-LRX4 and FER,
total protein from 60-90 seedlings per treatment per genotype was extracted as previously
described. For immunoprecipitations, M2 anti-Flag affinity gel (Sigma A2220-5ML) was used
and incubated with the crude extract for 2-3 h at 4 °C. Subsequently, beads were washed 3
times with wash buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1 mM phenylmethanesulfonyl
fluoride, 0,1 % IGEPAL) before adding Laemmli sample buffer and incubating for 10 min at
95 °C. Analysis was carried out by SDS-PAGE and immunoblotting.

Immunoblotting

Protein samples were separated in 10 % bisacrylamide gels at 150 V for approximately 2 h
and transferred into activated PVDF membranes at 100 V for 90 min. Immunoblotting was
performed with antibodies diluted in blocking solution (5 % fat-free milk in TBS with 0.1 %
(v/v) Tween-20). Antibodies used in this study: α-BAK1 (1:5 000; (Roux *et al.*, 2011); α-FLS2

(1:1000; (Chinchilla et al., 2007); α-FER (1:2000; (Xiao *et al.*, 2019), α-BAK1 pS612 (1:3000;

(Perraki *et al.*, 2018)), α-FLAG-HRP (Sigma Aldrich, A8592, dilution 1:4000); α -GFP (sc-

9996, Santa Cruz, used at 1:5000). Blots were developed with Pierce ECL/ ECL Femto

Western Blotting Substrate (Thermo Scientific). The following secondary antibodies were

used: anti-rabbit IgG-HRP Trueblot (Rockland, 18-8816-31, dilution 1:10000) for detection of

FLS2-BAK1 co-immunoprecipitation or anti-rabbit IgG (whole molecule)–HRP (A0545,

Sigma, dilution 1:10000) for all other western blots.

Statistical analysis

Statistical analyses were carried out using Prism 6.0 software (GraphPad). As mentioned in

the figure legend, statistical significances were assessed using non-parametric Kruskal-

Wallis bilateral tests combined with post-hoc Dunn's multiple pairwise comparisons, or using

a two-way non-parametric student's t test Mann-Whitney test.

Accession numbers

FER (AT3G51550), LRX3 (AT4G13340), LRX4 (AT3G24480), LRX5 (AT4G18670), RALF23

(AT3G16570), FLS2 (AT5G46330), BAK1 (AT4G33430).

Fig. 1 ¦ FER regulates the nanoscale organization of FLS2-GFP and BAK1-mCherry

Fig. 2 ¦ LRX3, LRX4 and LRX5 regulate PTI and BAK1-mCherry organization.

Fig. 3 ¦ FER malectin A domain regulates cell morphogenesis not PTI

Fig. 4 ¦ RALF23 perception regulates BAK1-mCherry organization.