Competition for electrons favors N$_2$O reduction in denitrifying *Bradyrhizobium*

isolates

Gao Y[1*], Mania D[1], Mousavi SA[2], Lycus P[1], Arntzen M[1], Woliy K[1†], Lindström K[2], Shapleigh JP[3],

Bakken LR[1] and Frostegård Å[1*]

[1] Faculty of Chemistry, Biotechnology and Food Sciences, Norwegian University of Life Sciences, Aas,

Norway

[2] Ecosystems and Environment Research programme, Faculty of Biological and Environmental Sciences,

and Helsinki Institute of Sustainability Science (HELSUS), University of Helsinki, Finland

[3] Department of Microbiology, Cornell University, Ithaca, New York, USA

[†]Present address: Department of Biology, Hawassa University, Hawassa, Ethiopia

[*]Corresponding author, E-mail asa.frostegard@nmbu.no; Tel. (+47) 67232473.

Summary

Bradyrhizobia are common members of soil microbiomes and known as N$_2$-fixing symbionts of

economically important legumes. Many are also denitrifiers, which can act as sinks or sources for

N$_2$O. Inoculation with compatible rhizobia is often needed for optimal N$_2$-fixation, but the choice

of inoculant may also have consequences for N$_2$O emission. Here, we analyzed the phylogeny

and denitrification capacity of *Bradyrhizobium* strains, most of them isolated from peanut-

nodules. All were dinitrifiers, but only ~1/3 could reduce N_2O while most others were net N_2O

producers. The N_2O-reducing isolates showed strong preference for N_2O- over NO_3^--reduction.

Such preference was also observed in a study of other bradyrhizobia and tentatively ascribed to

competition between the electron pathways to Nap (periplasmic NO_3^- reductase) and Nos (N_2O

reductase). Another possible explanation is lower abundance of Nap than Nos. Here, proteomics

revealed that Nap was instead more abundant than Nos, supporting the hypothesis that the

electron pathway to Nos outcompetes that to Nap. In contrast, *Paracoccus denitrificans,* which

has membrane-bond NO_3^- reductase (Nar), reduced N_2O and NO_3^- simultaneously. We propose

that the control at the metabolic level, favoring N_2O reduction over NO_3^- reduction, applies also

to other denitrifiers carrying Nos and Nap but lacking Nar.

Originality-Significance Statement

This study extends the current knowledge on denitrification in bradyrhizobia, which mostly

originates from studies of one model strain, by investigating the denitrification phenotypes of a

diverse collection of Bradyrhizobium isolates. Only 1/3 of them could reduce N_2O while the

others were net sources for this potent greenhouse gas. All N_2O-reducers showed strong

preference for N_2O over NO_3^-. We revealed by proteomics that this was not explained by

differences in the abundances of Nap (periplasmic nitrate reductase) and Nos (N_2O reductase),

which strengthens our hypothesis (Mania *et al*., 2020) of a metabolic control mechanism by

which Nos competes efficiently with Nap for electrons, making these organisms strong sinks for

N_2O. The findings highlight the potential importance of these organisms as N_2O sinks in natural

and agricultural ecosystems and pinpoint the need to take N_2O reduction into account, along

with N_2-fixation effectiveness, when searching for strains suitable for production of inoculants.

Introduction

Emissions of nitrous oxide (N_2O), the third most important anthropogenic greenhouse gas, have

been rising steadily during the past 150 years. Predictions for the rest of the century show either

continued N_2O emission increases or rates of decrease slower than the decreases occurring in

CO_2 emissions (Rogelj *et al.*, 2018; Thompson *et al.*, 2019). Agriculture is the main source of

anthropogenic N_2O emissions, with denitrification playing the major role in most soils (Thomson

et al., 2012). The primary cause of the emissions is excessive use of synthetic fertilizers (Reay *et*

al., 2012; Thompson *et al.*, 2019; Tian *et al.*, 2020), but cultivation of legumes also contributes

substantially with a global average of 1.29 (range 0.03-7.09) kg N_2O-N ha^{-1} $year^{-1}$ while the

corresponding estimate for fertilized agricultural fields was 3.22 (range 0.09-12.67) (Jensen *et al.*,

2012). In both cases, the ultimate driver of the emission is input of reactive nitrogen which

undergoes a series of redox reactions and eventually returns to the atmosphere as N_2O or N_2.

Our aim should be to minimize the N_2O/N_2 emission ratio of the system (Schlesinger, 2009). To

achieve this, novel mitigation options beyond "good agronomic practice" will be needed

(Davidson and Kanter, 2014). Several technical abatement options have been suggested,

including precision fertilization and use of nitrification inhibitors (Winiwarter *et al.*, 2018).

An alternative strategy is to enhance the populations of N_2O reducing microorganisms in the

soil. Selective *in situ* stimulation of N_2O reducing, indigenous soil organisms appears unrealistic,

while some options for the addition of N_2O reducing microbes may be feasible. One is the

suggestion to inoculate soybeans with rhizobia that are not only efficient N_2-fixers but also have

the capacity to reduce N_2O (Hénault and Revellin, 2011; Itakura *et al.*, 2013; Akiyama *et al.*, 2016).

At present, N_2O reduction is not taken into account when screening strains for their suitability as

inoculants and, in fact, several of the commercial inoculants currently used are only partial

denitrifiers (Zilli *et al.*, 2019). For example, only one of the five *Bradyrhizobium* strains most

commonly used as soybean inoculants in South America could reduce N_2O (Obando *et al.*, 2019).

The growing global need for plant-based proteins is expected to require new areas to be used for

legume cropping and will likely increase the demand for new legume varieties. Not all

combinations of legume cultivar and rhizobial strain result in an efficient symbiosis (Thrall *et al.*,

2011; Woliy *et al*, 2019). Importantly, indigenous soil populations of rhizobia may not be

compatible with a certain legume crop, especially if it has not been grown in that area before.

Therefore, the expansion of legume cropping is creating an increasing demand for a wider variety

of inoculants to ensure nodulation and optimal N_2 fixation efficacy (Santos *et al.*, 2019). This

offers a golden opportunity to mitigate N_2O emissions, providing that the inoculants are selected

among rhizobial strains that can reduce N_2O.

The purpose of the present study was to determine how denitrification phenotypes vary

across taxonomically diverse groups of bradyrhizobia, with particular focus on their capacity for

N_2O reduction, and to understand cellular mechanisms that make these organisms potential sinks

for N_2O. We determined the phylogenetic position of a set of *Bradyrhizobium* strains, obtained

from different culture collections, and analyzed their capacity for denitrification. Most of the

strains were isolated from nodules of peanut plants (*Arachis hypogaea*) growing in different

regions of China. Peanut is the fourth most important legume crop globally, with China being the

main producer (Arya *et al.*, 2016). We also included some bradyrhizobia from other parts of the

world, isolated from peanut and other legumes.

Several taxonomic groups within the genus *Bradyrhizobium* have been reported to denitrify

with many strains having truncated denitrification pathways lacking one or more of the steps

(Sameshima-Saito *et al.,* 2004; Zilli *et al.*, 2019; Mania *et al*., 2020). A complete denitrification

pathway in bradyrhizobia comprises the periplasmic reductase Nap for dissimilatory NO_3^-

reduction, encoded as a part of the *napEDABC* operon; the Cu-containing nitrite reductase NirK

encoded by *nirK* only or as a part of the *nirKV* operon, although cd_1-type NirS nitrite reductase

has also been reported (Sánchez and Minamisawa, 2018); the cytochrome *c* dependent nitric

oxide reductase cNor *(norCBQD* operon); and a clade I N_2O reductase Nos (*nosRZDYFLX* operon)

(Velasco *et al.*, 2001;Mesa *et al.*, 2002; Delgado *et al.*, 2003; Velasco *et al.*, 2004; Bedmar *et al.*,

2005). While the bradyrhizobia, along with several other denitrifiers, only have the periplasmic

Nap for dissimilatory NO_3^- reduction, other denitrifiers have the membrane-bound Nar, or both.

Nap is a periplasmic heterodimer (NapAB) consisting of the small electron transfer subunit NapB

and the catalytic NapA, which draws electrons from quinol via the membrane-bound *c*-type

tetrahaeme cytochrome NapC. Nar is a three-subunit enzyme (NarGHI) anchored in the

membrane by NarI, which is a transmembrane protein that draws electrons from quinol to the

catalytic NarG via NarH, both facing the cytoplasm (Richardson, 2000; Richardson *et al.*, 2001).

Nir, Nor and Nos receive electrons from the bc_1 complex via periplasmic small c-type cytochromes

(Berks *et al.*, 1995; Zumft, 1997). Clade I type Nos reductase may in addition draw electrons from

quinol via the membrane bound NosR (Zhang *et al.*, 2019).

In a recent study, Mania *et al.* (2020) screened strains within the genus *Bradyrhizobium* for

their denitrification phenotype. Most of them could perform two or more of the denitrification

steps, but only half of them could reduce N_2O. All the N_2O reducing strains displayed the same

strong preference for N_2O over NO_3^- reduction when incubated under anoxic conditions, which

was suggested to be due to competition between the electron pathways to Nap and Nos. The

quantification of denitrification gene transcripts showed 5-8 times lower transcript numbers of

nap compared to *nos,* however. Therefore, it could not be excluded that the phenomenon is due

to low abundance of Nap compared to Nos. Here we have investigated the denitrification geno-

and phenotypes for another collection of *Bradyrhizobium* isolates and deepened our

understanding of the competition for electrons between Nap and Nos by refined analyses of the

electron flow kinetics, in combination with the determination of the relative amounts of Nap and

Nos by proteomics. In addition, we compared the denitrification kinetics of the bradyrhizobia

with the denitrification model bacterium *Paracoccus denitrificans*, which holds a complete

denitrification pathway including both Nap and Nar (Zumft, 1997; Richardson, 2000). The results

showed that in all bradyrhizobia with complete denitrification, NO_3^- reduction was almost

completely hampered in the presence of N_2O. The proteomics analysis provided evidence that

this was not due to a low abundance of Nap, also showing that transcript abundance cannot be

directly translated into corresponding protein abundance. As expected, the NarG carrying *P.*

denitrificans showed a different denitrification phenotype with simultaneous reduction of NO_3^-

and N_2O.

Results

Phylogenetic distribution of denitrifying bradyrhizobia

The phylogenetic position of the 23 test strains was determined by MLSA analysis of four protein-

coding housekeeping genes (*atpD-glnII-recA-rpoB*) (Fig. 1). The analysis comprised 78

bradyrhizobial strains and included all hitherto described *Bradyrhizobium* reference species. All

but one of the 23 test strains listed in Table 1 were assigned to the *Bradyrhizobium japonicum*

supergroup (Fig. 1). The exception was *Bradyrhizobium lablabi* HAMBI 3052^T, which is a member

of the *Bradyrhizobium jicamae* supergroup according to the new genome-based taxonomy of

bradyrhizobia proposed by Avontuur *et al.* (2019). The test strains were placed in 11 lineages

(clades A-K) with high posterior probability (PP=1.00). Most test strains were clustered within the

clades of seven diverse bradyrhizobial species. Strains *Bradyrhizobium* sp. HAMBI 2134 and 2142

formed a monophyletic cluster (clade G) with high posterior probability (1.00). *Bradyrhizobium*

sp. strain HAMBI 2116 was accommodated in a lineage (clade A) distant from other bradyrhizobia.

The phylogenetic relationships revealed by the four-gene MLSA tree (Fig. 1) showed good

correspondence with the results from an analysis of all the HAMBI strains using the GTDB Tool

kit (Fig. S1). In both trees strains HAMBI 2299, 2153 and 2116 were most closely related to the

reference species *B. yuanmingense*; strains HAMBI 2125 (and 2126) and 2127 were closest to *B.*

ottawaense; strain HAMBI 2130 was closest to *B. shewense*; strains HAMBI 2115 was closest to

B. japonicum USDA6; strains HAMBI 2128 (as well as HAMBI 2129 and 2150, Table S1) were

closest to *B. arachidis*; strain HAMBI 3052 was closest to *B. lablabi.* Strains HAMBI 2149, 2133,

2136, 2137, 2151 and 2152 formed one cluster in the GTDB tree together with *B. japonicum* USDA

135. This cluster was close to *B. liaoningense*, as also found in the MLSA analysis (Clade F, Fig. 1),

which did not include *B. japonicum* USDA 135. Strains HAMBI 2134 (and 2135, Table S1) and 2142

formed a separate cluster in both trees.

We originally included 21 isolates from the HAMBI culture collection in our study, all

deposited with different strain names. Pairwise whole genome comparisons revealed that some

strains were very similar with ANI values >99.2% and in most cases >99.9 % (Table S1), which was

also supported by the clustering of similar strains in the GTDB tree (Fig. S1). Since the strains

within such clusters with ANI values >99% also showed practically identical denitrification

phenotypes, we present the result for only one of the strains in each such cluster in the main

paper, and the results for the others as supplementary information.

The denitrification end-product analysis showed that all test strains were denitrifiers, able to

reduce NO_3^- and/or NO_2^- to NO, N_2O or N_2. Seven of the 19 strains tested for denitrification in the

present study were complete denitrifers that reduced NO_3^- to N_2. The four Ethiopian strains

AC87j1, AC86d2, AC101b and AC70c were determined earlier to be complete denitrifiers (Mania

et al. 2020). All complete denitrifiers except *B. lablabi* (clade K) belonged to the *B. japonicum*

supergroup and were positioned in clades E, H, I and J. The clades E, H and J represented the

species *B. guangxiense*, *B. ottawaense* and *B. shewense*, respectively. Clade I, accommodating

strains *Bradyrhizobium* sp. HAMBI 2125 and AC70c, formed a separate, monophyletic group. The

other test strains had truncated denitrification pathways and were able to reduce NO_3^- and NO_2^-

to NO (5 strains) or N_2O (7 strains). The seven strains having N_2O as end-product were distributed

in clades A, C, D and G. Three strains grouped with *B. arachidis* in clade C*. Bradyrhizobium* sp.

strain HAMBI 2115 was placed in clade D along with *B. japonicum* USDA6^T. Three strains were

accommodated in clades A and G, neither of which included any of the hitherto described

Bradyrhizobium species. The strains with NO as denitrification end-product were found in clades

B, C and F, which also harbored the species *B. yuanmingense*, *B. arachidis* and *B. liaoningense*,

respectively. Strains HAMBI 2153 and 2149, as well as the strains highly similar to HAMBI 2149

(Table S1), were also examined using detailed gas kinetics to further examine whether they could

reduce NO (Fig. S2). Like in the end-product investigation they produced large amounts of NO

(14-15 µmol flask^{-1}) while no reduction of NO or N_2O was detected (Fig. S2).

Consistency between denitrification gene operons and end products

The genetic potential for denitrification, based on whole genome sequencing, and the

corresponding phenotype, determined by end-point analysis and gas kinetics, were compared

for 13 strains (Table S2), as well as for isolates with genomes with ANI values >99.2% to these

strains (Table S1). The five strains able to reduce NO_3^- to N_2 had complete operons for all four

reduction steps. For most of the other strains, with NO or N_2O as denitrification end -products,

the lack of a function was explained by complete lack of the corresponding operon. However,

two strains (HAMBI 2153 and HAMBI 2299) showed a discrepancy between denitrification

genotype and phenotype. They were apparently unable to reduce NO and accumulated µM

concentrations of NO, despite having the *nor* operon. Comparison of these strains with strains

having functional NO reduction revealed that strain HAMBI 2153 had a frame shift in *norB*, while

norQ was not in the correct reading frame. Strain HAMBI 2299 carried *nap, nir* and *nor* operons

with all expected genes present. The end-point analysis of this strain showed that after 10 days

of incubation with 1 mM NO_3^-, 0.5 mM NO_2^- and 1 ml N_2O in headspace, it had produced toxic

levels (>10 µM) of NO. Inspection of the amino acid sequences did not reveal any obvious

explanations for lack of Nor activity.

All N_2O-reducing Bradyrhizobium *strains preferred N_2O over NO_3^- as electron acceptor*

To determine if the complete denitrifiers showed a similar, strong preference for N_2O over NO_3^-

as reported by Mania *et al.* (2020) for other *Bradyrhizobium* strains, detailed gas kinetics was

analyzed. The results showed remarkably similar kinetics for all strains during and after the

transition from aerobic to anaerobic respiration in flasks to which N_2O had been added to the

headspace (Figs 2A, S3, S4, S5). Strain HAMBI 2125 was chosen as a representative for the

complete denitrifiers. The NO_2^- concentration in cultures of this strain was measured along with

the gas kinetics (Figs 2A, S3), allowing the estimation of NO_3^- concentrations by N-mass balance.

The first phase, marked in blue, was dominated by O_2 respiration. When the O_2 concentration

in the medium reached 4.6 µM, some NO_3^- reduction took place, measured as a slight

accumulation of NO_2^- and NO (0.7 and 1.7 µmol; not clearly visible in the figure due to scaling).

The small but significant decrease in N_2O during the aerobic phase was caused by sampling losses,

not by N_2O reduction or other reactions, as explained in Fig. S6. The same is seen for experiments

presented in Figs. 2B and S3-S5. Reduction of exogenous N_2O started when the O_2 concentration

in the liquid was approximately 3.0 µM and rapidly reached a rate of 10.4 ± 1.2 µmol N flask^{-1} h^{-1}

^{1}, matching the corresponding N_2 production rate of 9.7 ± 1.0 µmol N flask^{-1} h^{-1}. This phase of

rapid and stoichiometric reduction of N_2O to N_2, marked in pink, lasted until most of the

exogenous N_2O was reduced. No NO_3^--reduction took place during reduction of the exogenous

N_2O. When the exogenous N_2O was depleted, the rate of NO_3^- reduction increased and the rate

of N_2 production decreased instantaneously to 2.22 ± 0.18 µmol flask^{-1} h^{-1} (phase marked in

green). When NO_3^- reduction started, the NO_2^- concentration increased sharply and reached

approximately 200-350 µM (9.8-17.4 µmol flask^{-1}), accounting for 20-35% of the initial 50 µmol

NO_3^--N. The N_2 curve for strain HAMBI 2125 reached a stable plateau at 123 µmol N_2-N, which

accounts for the sum of N_2O- and NO_3-N present initially. The same experiment was conducted

with other strains, but without measuring NO_2^- (Figs S4, S5), and they all showed the same fast

stoichiometric conversion of N_2O to N_2 and a subsequent slower production of N_2 from NO_3^-,

lasting until the N_2O+NO_3^--N was completely recovered as N_2-N. Strain HAMBI 2125 controlled

NO during the entire incubation period, reaching a small peak of 11 nmol flask^{-1} (6.9 nM in the

liquid) around 41 hpi. For the other strains NO concentrations never exceeded 50 nM, and most

strains kept NO concentrations <20 nM during the entire incubation (Figs 2A, S3, S4, S5).

Nar carrying Paracoccus denitrificans *showed no preference for N_2O over NO_3^-*

In accordance with our hypothesis, the Nar carrying *P. denitrificans* reduced NO_3^- simultaneously

with the exogenously supplied N_2O, indicating no hampering of NO_3^- reduction by N_2O (Fig. 2B).

In contrast to all the examined *Bradyrhizobium* strains (Figs 2A, S3, S4, S5), *P. denitrificans* started

to reduce NO_3^- and headspace N_2O concomitantly when the O_2 concentration was close to

depletion and stoichiometrically accumulated NO_2^- from NO_3^- reduction. Nitrite reduction and

emergence of NO started when most of the NO_3^- had been reduced to NO_2^-.

Cell specific electron flow rates

The measured gas- and NO_2 kinetics were used to estimate the electron flow rates to the various

reductase (μmol e$^-$ flask^{-1} h^{-1}) for each time increment between gas samplings. To convert these

rates to cell specific electron flow rates (fmol cell^{-1} h^{-1}), we needed estimates of the cell density

throughout an incubation. For this, we used cumulated electron flow to the various electron

acceptors (calculated from the gas- and NO_2^--measurements), and the cell yield per mol electrons

to each electron acceptor, which was determined in a separate experiment (see Experimental

procedures and Fig. 5). In *Bradyrhizobium* strain HAMBI 2125 (Fig. 2C), the cell specific electron

flow was sustained at a more or less constant level (5-6 fmol e$^-$ cell^{-1} h^{-1}) through the transition

from respiring O_2 to respiring N_2O, until the external N_2O was depleted (phases marked blue and

pink in the corresponding gas kinetics Fig. 2A), but declined to ~3-4 fmol e$^-$ cell^{-1} h^{-1} when

depending on the activity of Nap. In *P. denitrificans* (Fig. 2D) there was instead a simultaneous

electron delivery to Nar and Nos during the phase when exogenous N_2O was being reduced

(marked pink in the corresponding gas kinetics, Fig. 2B). This was later followed by electron flow

to Nir, Nor and Nos, after the depletion of nitrate. The cell specific respiratory metabolism was

10-13 fmol e$^-$ cell^{-1} h^{-1} during the phase dominated by aerobic respiration, then increased to 15-

16 fmol e$^-$ cell^{-1} h^{-1} when respiring NO_3^- and N_2O, followed by a decrease to approximately 8 fmol

e$^-$ cell^{-1} h^{-1} during respiration of the accumulated NO_2^-.

Injection of N_2O to actively denitrifying cultures resulted in an immediate arrest of NO_3^- reduction

To investigate whether the preferred electron flow to N_2O in the foregoing experiments could be

explained by regulation at the transcriptional level (i.e. early expression of *nos*), N_2O was injected

into a culture that was actively reducing NO_3^- to N_2. A transcriptional regulation would not result

in an immediate switch to the exogenous N_2O as electron acceptor but would instead be seen as

a simultaneous reduction of all N-oxides present. This was tested on strain HAMBI 2125 (Fig. 3).

The cultures were incubated in flasks containing YMB with 2 mM initial NO_3^-, and with 1% O_2 but

without exogenous N_2O in the headspace. Denitrification (from NO_3^- reduction) commenced

when O_2 reached 4.3 μM and proceeded at a rate of 2.6 ± 0.4 μmol N flask^{-1} h^{-1}. At 32 hpi, when

the cells were actively reducing NO_3^- to N_2, approximately 300 μmol N_2O-N was injected. This led

to a complete suppression of NO_3^- reduction and rapid decline in NO accumulation. The measured

N_2 production rate was similar to the measured N_2O reduction rate (23.9 ±3.6 and 26.4 ±3.6 μmol

N flask^{-1} h^{-1}, respectively). The NO_3^- reduction recovered upon depletion of external N_2O,

reflected in an increase in NO and an N_2 production rate similar to that before the N_2O injection.

In an additional experiment, including all seven N_2O-reducing strains, exogenous N_2O was

present in the headspace from the start of the incubation (Fig. S5). As expected from the first

experiment (Fig. 2A), the exogenous N_2O was rapidly reduced, followed by NO_3^- reduction to N_2

at a slower rate. A second injection of N_2O into the actively denitrifying cultures resulted in an

immediate increase in the N_2 production rate, comparable with that during reduction of the initial

N_2O. These results supported the notion that N_2O was preferred as electron acceptor over NO_3^-.

The relative abundances of the reductases could not explain the preference for N_2O over NO_3^-

A proteomics analysis was done for strain HAMBI 2125. The abundance of NapA, NirK and NosZ

molecules, measured as intensities, were determined at different time points in cells sampled

during aerobic respiration (21% O_2), and during and after transition from 1% oxygen respiration

to denitrification (Fig. 4). The level of NorC molecules detected was two orders of magnitude

lower than those for the other reductases, suggesting that membrane-bound enzymes require a

different extraction protocol. Therefore, only values for the periplasmic reductases NapA, NirK

and NosZ were included here. The aerobic pre-incubation over several generations effectively

removed most denitrification reductases that may have lingered from earlier periods of anoxia,

seen from the signals from NapA and NirK being 293-585 times lower than those observed during

denitrification. Nos values were only 16-47 times lower, indicating low levels of constitutive

transcription of the *nos* gene and translation of the corresponding mRNA under aerobic

conditions. All three reductases increased in abundance during the O_2 depletion phase (blue) and

when O_2 approached depletion, the NapA/Nos and NapA/NirK ratios were 1.4±0.2 and 4.1±1.8

(sd, n=3). Interestingly, the NapA/Nos ratio was stable during the period of strong respiration of

exogenous N_2O, marked pink in Fig. 4, with values of 1.5± 0.3 and 1.2 ± 0.2, although practically

no NO_3^- reduction took place. The NapA/Nos ratio decreased to 0.7 during the phase of NO_3^-

reduction to N_2 (after depletion of exogenous N_2O). A similar pattern was found for the

NapA/NirK ratios, while the NosZ/NirK ratios stayed around 1.0 during the anaerobic phase.

Growth rate and yield per mol electrons for aerobic and anaerobic respiration

The experiment was designed to determine the growth rate and growth yield per mol electrons

for aerobic and anaerobic respiration (N_2O and NO_3^-), but also to inspect if cumulated electron

flow to the various electron acceptors could be used to predict cell density throughout a batch

incubation. The results summarized in Fig. 5 confirm that cumulated electron flow can be used

to predict increase in cell density (or dry weight, DW) throughout an incubation, provided that

the yield per electron to the various electron acceptors is known. The aerobic growth rate was

twice the anaerobic growth rate by N_2O respiration (0.063 and 0.031 h^{-1}). Growth from complete

denitrification (NO_3^- provided as e^- acceptor) was 0.024 h^{-1} (Fig. 5 A-C). The growth yield per mol

electrons was 7.17±0.38 g DW mol^{-1} e^- for aerobic respiration, 5.1±0.10 g DW mol^{-1} e^- from N_2O

respiration and 4.03±0.20 g DW mol^{-1} e^- for complete denitrification (NO_3^- to N_2) (Fig. 5D). In the

same experiment, we determined the DW per cell for cells grown by respiring O_2, N_2O and NO_3^-

(Fig. S7 B). Since the values for growth on the different electron acceptors were not significantly

different, they were regarded as replicates, giving an average cell DW of 615 ±54 fg (sd, n=12).

Discussion

The study comprised a set of taxonomically diverse *Bradyrhizobium* strains, as shown by the

phylogenetic tree (Figs 1, S1). This genus is commonly found among the most dominant

organisms in soil microbial communities (Delmont *et al*., 2012; Xu *at al*., 2014) and of large

environmental and economic value for their N_2-fixation ability in symbiosis with legumes. Based

on the MLSA analysis of four concatenated housekeeping genes (Fig. 1) the strains clustered in

eleven clades, covering various parts of the phylogenetic tree. Five strains in three clades did not

group with described *Bradyrhizobium* species, suggesting that three new species may be

delineated for these three clades. Our results from the denitrification analyses support other

studies showing that complete denitrification pathways are more common in the *B. japonicum*

supergroup than in other phylogenetic branches of *Bradyrhizobium* (Sameshima-Saito *et al.*, 2004**;**

Mania *et al*., 2020). The complete denitrifiers in the present study belonged to clusters related

to *B. shewense*, *B. ottawaense* and *B. guangxiense*, with the exception of one strain which

clustered with the distantly related *B. lablabi* (Fig. 1). Although this suggests that some taxonomic

groups of bradyrhizobia have higher frequency of complete denitrifiers than others, taxonomic

position cannot be used as an indicator of the denitrification end-product since even very closely

related strains may differ in this regard (Mania *et al.*, 2020).

Recent research has revealed several transcriptional and post-transcriptional mechanisms

that influence the amounts of intermediate products released from denitrifying organisms (Liu

et al., 2013; Torres *et al.*, 2016; Lycus *et al.*, 2018; Gaimster *et al.*,2019; Jiménez-Leiva *et al.*, 2019).

Although of value by itself, such information may be anecdotal. To understand the environmental

relevance of a regulatory mechanism, it is important to know how widespread it is. We found the

same preference for N_2O over NO_3^- as electron acceptor in all strains with complete

denitrification, as did Mania *et al.* (2020) for a different set of isolates. This indicates a

phenomenon common to bradyrhizobia. In theory, it could be explained by lower numbers of

NapA molecules compared to NosZ molecules. The results by Mania *et al.* (2020) for strain AC

87j1, which is closely related to strain HAMBI 2125, could not disprove this, since transcription

analysis showed 5-8 times lower copy numbers of transcripts coding for NapA compared to NosZ.

This prompted us to perform a proteomics analysis in the present study. The results (Fig. 4)

provided clear evidence that the strong N_2O reduction was not due to higher abundance of NosZ

vs NapA, thus supporting the hypothesis by Mania *et al.* (2020) that the electron pathway to NosZ

is inherently more competitive than that to NapA.

Electrons from the quinol pool are channeled to Nap via the membrane-bound NapC, and to

Nos, Nor and Nir via the membrane-bound bc_1 complex and periplasmic cytochromes (Shapleigh,

2013). Most likely, the electron pathway to NosZ via bc_1 is inherently more competitive than that

to NapAB via NapC**,** plausibly due to a combination of stronger affinity for quinol and a higher k_{cat}

of bc_1 versus NapC. This implies a general mechanism which may apply not only to bradyrhizobia,

but possibly to all denitrifiers that carry Nap and Nos but lack the membrane-bound Nar.

Verifying this will require testing of a wide range of organisms. As a first step, we included in this

study a comparison of the denitrification kinetics of the bradyrhizobia with *P. denitrificans*, which

carries Nar in addition to Nap and Nos. The results supported our hypothesis by clearly

demonstrating that, while the *Bradyrhizobium* strains left NO_3^- untouched as long as exogenously

supplied N_2O was present (Figs 2A, S3, S4, S5), *P. denitrificans* reduced these electron acceptors

simultaneously (Fig. 2B). This suggests that the electron pathways from the quinol pool to NarG

via NarI and NarH and to NosZ via the bc_1 complex and cytochrome c compete equally well for

electrons, at least in *P. denitrificans*.

The proteomics analysis of HAMBI strain 2125 showed that NosZ was synthesized under fully

aeriated conditions, while NapA and NirK were not. Before sampling for proteomics analysis, the

cultures had been incubated in flasks with vigorous stirring for at least 10 generations to secure

that denitrification reductases produced during earlier exposure to denitrifying conditions were

diluted to near extinction in the cells. This was the case for NapA and NirK in the sample from

21% O_2 (Fig. 4). NosZ molecules were, however, approximately 10 times more abundant than

NapA and NirK during this phase, which suggests constitutive, albeit low, synthesis of NosZ. The

transcription analysis of strain AC87j1 reported by Mania *et al.* (2020) provides no clear

explanation to this, though. In the presence of 5 μM O_2 in the liquid, there were < 0.3 *nosZ* and

nirK transcripts cell^{-1} with only slightly higher levels of *nosZ* than *nirK*. The *napA* transcription was

somewhat higher with 0.8 copies cell^{-1}. It is possible, still, that a low but steady production of Nos

takes place in these organisms irrespective of O_2 concentrations, in line with the higher β-

galactosidase activity observed for a *nos-lacZ* fusion of *B. japonicum* compared to a *nap-lacZ*

fusion (Bueno *et al.*, 2017; Torres *et al.*, 2017). This could be a strategy for cells to be prepared

for anaerobic conditions and avoid being entrapped in anoxia without enough energy to produce

the denitrification machinery (Lycus *et al.*, 2018). If so, it suggests that these bacteria may be

ready to reduce any N_2O produced by other organisms in their environment, thus acting as net

sinks for this greenhouse gas.

Seven strains lacked the last step of denitrification and thus had N_2O as end-product,

explained by the absence of the entire *nos* operon in all cases (Table S2). When exposed to

denitrifying conditions in the environment, such organisms will reduce all available N-oxides to

N_2O and thus act as net producers of this greenhouse gas. Three strains had NO as denitrification

end-product, but only HAMBI 2149 lacked the entire *nor* operon (Table S2). The other two strains,

HAMBI 2153 and 2299, contained *nor* operons but showed some sequence differences compared

to strains with functional NO reduction. Further investigations of the lacking NO reduction were,

however, beyond the scope of the present study. Organisms that produce NO but are incapable

of reducing it are apparently not uncommon (Casella *et al*., 2006; Falk *et al.*, 2010; Mania *et al.*,

2020). In the environment such organisms are most likely found in consortia with organisms that

express Nor or other NO-reducing enzymes.

The estimated growth yields were calculated based on the electron flow to the different

reductases (Fig. 5) and the proportions were expected to match that of the charge separations

per electron transport for the various respiratory pathways, as listed by (van Spanning *et al.*,

2007). The number of charge separations per 10 electrons ($q/10e^-$) are 50 for low affinity oxidase,

40 for high affinity oxidases, 30 for Nos alone, and 26 for the entire Nap, Nir, Nor, Nos

denitrification pathway (NO_3^- to N_2), which would a yield ratio of 100:60:52 for the three

pathways (O_2, N_2O, $NO_3^- \rightarrow N_2$) if we assume that low affinity oxidases are used, and 100:75:65 for

high affinity oxidases. Our measured yields were 12, 8.5 and 6.7 $*10^{12}$ cells per mol electrons for

the three pathways, i.e. a yield ratio of 100:71:56, which agrees reasonably with that predicted

for high affinity oxidases.

The emerging evidence that bradyrhizobia with complete denitrification deplete N_2O in their

surroundings before reducing available NO_3^- is promising from an environmental point of view

since they have the potential to act as net sinks for N_2O produced by other organisms in their

vicinity. In legume fields, nodules and other N-rich organic material will be degraded, releasing

NH_4^+, which will rapidly be metabolized by nitrifying bacteria and archaea, producing NO_3^- while

consuming O_2. This will create optimal conditions for denitrification. Legume cropped fields

account for a significant share of the global N_2O emissions (Jensen *et al.*, 2012), and it is likely

that coupled nitrification-denitrification is a main cause (Bakken and Frostegård, 2017). One way

of mitigating these emissions is to enhance the number of efficient N_2O reducing bacteria in the

soil microbial community. Rhizobial inoculants are used since long to increase the yield of

leguminous crops and at the same time avoid the need for synthetic fertilizers. Different

combinations of rhizobial strain and legume cultivar may differ substantially in nitrogen fixation

effectiveness, which is also affected by environmental conditions (Woliy *et al.*, 2019; Lindström

and Mousavi, 2020), and large efforts are put into the development of new inoculants that are

compatible with various types of legumes as well as soil types and climates (Martínez *et al.*, 2016).

The present study shows that *Bradyrhizobium* strains also differ with respect to N_2O reduction,

and suggests that new inoculants should be selected among rhizobia that express Nos.

Experimental Procedures

Bacterial strains

The rhizobial isolates were obtained from various culture collections. Table 1 shows a list of host

plant and geographic origin of the strains used in the phylogenetic analysis (all listed isolates;

details see (Van Rossum *et al.*, 1993; Van Rossum *et al.*, 1994; Zhang *et al.*, 1999; Chang *et al.*,

2011; Wang *et al.*,2013; Li *et al.*, 2015; Degefu *et al.*, 2017) and the strains analyzed for their

denitrification capacity (all except the four strains from Ethiopia earlier determined to have a

complete denitrification pathway (Mania *et al.,* 2020). Preparation of *Bradyrhizobium* cultures

and growth conditions followed Mania *et al.* (2020). All cultures were grown at 28 °C in Yeast

Mannitol Broth (YMB; 10 g l^{-1} D-mannitol, 0.5 g l^{-1} K$_2$HPO$_4$, 0.2 g l^{-1} MgSO$_4$·7H$_2$O, 0.1 g l^{-1} NaCl and

0.5 g l^{-1} yeast extract, pH 7.0; Schwartz, 1972).

Phylogenetic analyses

DNA was extracted using QIAamp DNA Mini Kit (Qiagen). The genome sequences were obtained

from another ongoing project: (JGI Gold study ID: Gs0134353), and their Genome accession

numbers are listed in Table S3. Genome sequences were analyzed using RAST (Rapid Annotations

using Subsystems Technology). Whole-genome average nucleotide identity (ANI) comparisons

were done using FastANI v.0.1.2. A phylogenetic tree that included all HAMBI and CCBAU strains

from this study (Table 1), as well as the Ethiopian strains AC70c, AC86d2, AC87j1 and AC101b and

the type strain of all species of the genus *Bradyrhizobium* (February 2020,

http://www.bacterio.net/bradyrhizobium.html), was constructed based on multilocus sequence

analysis (MLSA) of the four housekeeping genes *atpD* (ATP synthase F1, beta

subunit), *glnII* (glutamine synthetase type II), *recA* (recombinase A), *rpoB* (RNA polymerase, beta

subunit). The sequences from all HAMBI strains used in the MLSA analysis were obtained from

the present study. Amplicons of *rpoB* and *atpD* of the Ethiopian strains AC70c, AC86d2 and

AC101b and *recA* from AC70c were Sanger sequenced. For primers and PCR conditions, see Table

S4. All other sequences were obtained from GenBank (http://www.ncbi.nlm.nih.gov/genbank).

The GeneBank accession numbers for all housekeeping genes used are listed in Table S5. The

gene sequences were aligned by ClustalW (Larkin *et al.*, 2007) in Bioedit v7.0.5.3 (Hall, 1999), and

the alignments were edited manually. The best-fit models of nucleotide substitution were

selected based on Akaike information criterion (AIC) applying MEGA7 (Kumar *et al.*, 2016).

The Bayesian phylogenetic analyses of the concatenated loci (*atpD-recA-glnII-rpoB*) of 78

Bradyrhizobium strains were performed using the algorithm Metropolis coupled Markov chain

Monte Carlo (MCMCMC) twice for ten million generations using MrBayes v3.2.7a program

(Ronquist *et al.*, 2012). *Neorhizobium galegae* was used to root the tree. The general time

reversible plus gamma distribution plus invariable sites (GTR + G + I) was chosen as the best-fit

model of phylogenetic analyses of the four partition of the concatenated dataset (*atpD-glnII-*

recA-rpoB). The algorithms were analyzed using Tracer v1.7 (Rambaut *et al.*, 2018), and the

phylogenetic trees were visualized by Figtree v1.4.3 (https://beast.community/figtree). For

comparison with the MLSA tree based on four housekeeping genes, a phylogenetic tree of all the

HAMBI test strains listed in Table S2 was constructed using the Genome Taxonomy Database

Toolkit (GTDB-Tk) v0.3.2, which is based on a phylogeny inferred from the concatenation of 120

ubiquitous, single-copy proteins (Chaumeil *et al*., 2020).

Denitrification end products and gas kinetics analyses

Prior to these experiments the cultures were streaked for purity and single colonies were picked and checked for possible contaminating bacteria by sequencing of the 16S rRNA genes (for primers and PCR conditions see Table S4). Pure cultures were preserved as in glycerol (15%) at -80 °C. Cultures were raised from these frozen stocks in stirred batches of 50 ml YMB supplemented with filter sterilized KNO_3 and/or KNO_2 solutions. The flasks were sealed with sterilized, gas tight butyl rubber septa (Matrix AS, Norway). Air was removed by applying vacuum for 6 x 360 s, followed by 30 s He-filling. The desired amount of O_2 and/or N_2O was injected into the headspace, and the over-pressure released (Molstad *et al.*, 2007). Denitrification kinetics of *Bradyrhizobium* strains with complete denitrification pathways were compared to the model bacterium *Paracoccus denitrificans* PD1222, which was grown under the same conditions as the *Bradyrhizobium* strains, except that Sistrom´s medium supplemented with 2 mM KNO_3 was used (Sistrom, 1962; Bergaust *et al.*, 2010).

Denitrification end-products were determined for all strains as described previously (Lycus *et al.*, 2017). Detailed gas kinetics were measured following Mania *et al*. (2020). Pre-cultures were added to incubation flasks (n=3) supplied with 1 mM initial KNO_3 and, when indicated, also with 0.5 mM KNO_2. Unless stated otherwise, 0.7 mL O_2 (1 vol% in headspace) and 0.7 ml N_2O (approximately 50 µmol N flask^{-1}) were added to the headspace at the start of the experiment (0 hpi, hours post inoculation). For the determination of end-products, the amounts of O_2, NO, N_2O and N_2 in the headspace were measured at the start and after 10 days of incubation.

All strains with a complete denitrification pathway according to the end-point analysis, as

well as all HAMBI strains that had NO as end-product, were investigated further by analyzing gas

kinetics in response to oxygen depletion. To secure an inoculum in which any previously

synthesized denitrification enzymes had been diluted out by aerobic growth, all cultures were

precultured aerobically for at least 6-8 generations. This was done in uncapped flasks covered

with aluminum foil (to allow oxygen transport), with vigorous stirring (600 rpm) at 28 $^{\circ}$C. To

further ensure the absence of hypoxia, the cultures were transferred to new flasks before the

OD_{600} reached 0.3. Portions of the precultures were inoculated into 120 ml flasks containing 50

ml YMB supplemented with 1, 2 or 3 mM KNO_3 and with He, O_2 and N_2O in headspace as

described above for the end-product measurements. The flasks were placed in a robotized

incubation system that frequently measured O_2, NO, N_2O and N_2 in headspace (Molstad *et al.*,

2007; Molstad *et al.*, 2016). Leakage and sampling losses were taken into account when

calculating the rates of production/consumption of each gas, see Molstad *et al.* (2007) for details.

Earlier experiments determined that no gas production (NO, N_2O or N_2) took place from chemical

reactions between of NO_2^- and the medium under the conditions used (Mania *et al.*, 2020).

Nitrite measurements

Samples for NO_2^- measurements (0.1-1.0 ml) were taken manually from the liquid phase through

the rubber septum of the flasks using a sterile syringe. NO_2^- concentrations were determined

immediately after each sampling by drawing a 10 μL liquid sample which was injected into a

purging device containing 50% acetic acid with 1% (w/v) NaI (at room temperature), by which

NO_2^- is reduced instantaneously to NO (Cox, 1980; MacArthur *et al.*, 2007). The quantities of

resultant NO were measured by chemiluminescence using a NO analyser (Sievers NOA^{TM} 280i, GE

Analytical Instruments). The system was calibrated by injecting standards. Sampling for NO_2^-

measurements was done as frequently as the gas sampling.

Proteomics

Relative quantification of the denitrification reductases was done for the strain *Bradyrhizobium*

sp. HAMBI 2125, chosen as a representative for the complete denitrifiers (Fig. 1), which all

showed the same preference for N_2O over NO_3^- reduction (Figs 2A, S3, S4, S5). Cells were first

raised from frozen stock under strictly aerobic conditions as explained above. As the aerobic

preculture reached OD_{600} = 0.05 (2.9*10^7 cells ml^{-1}), it was used to inoculate 18 flasks (1 ml

inoculum flask^{-1}) containing YMB with 1 mM NO_3^- and with 1 ml N_2O (approximately 80 µmol

N_2O-N flask^{-1}) and 1 % O_2 in headspace. Samples for protein analysis were taken i) from the

aerobic inoculum; ii) just before transition to anaerobic respiration (22 hpi, when O_2 reached ~4.6

µM); iii) during the period of rapid reduction of exogenous N_2O (27 and 32 hpi); and iv) during

the period of NO_3^- reduction to N_2 (36 hpi). Three replicate flasks were harvested at each time

point. Gas kinetics, NO_2^- concentrations and sampling times for proteomics are shown in Fig. S8.

After measuring the OD_{600}, the samples were centrifuged (10 000 g, 4 °C, 10 min) and the cell

pellets were stored at -20 °C for protein extraction. In addition, we harvested and froze cells from

the aerobic preculture.

To extract the proteins, the cell pellets were thawed, resuspended in lysis buffer (20 mM Tris-

HCl pH8, 0.1 % v/v Triton X-100, 200 mM NaCl, 1 mM DTT) and treated with 3x45 s bead beating

with glass beads (particle size ≤ 106 µm Sigma) at maximum power and cooling on ice between

the cycles (MP Biomedicals™ FastPrep- 24™, Thermo Fischer Scientific Inc). Cell debris was

removed by centrifugation (10 000 g; 5 min) and the supernatant, containing water soluble

proteins, was used for proteomic analysis based on Orbitrap-MS as described before (Conthe *et*

al., 2019). To quantify the denitrification reductases we calculated the fraction of the individual

enzymes as percentages of the sum of all LFQ (label-free quantitation of proteins) abundances

for each time point.

Determinations of growth and yield

Aerobic and anaerobic growth was investigated in detail for *Bradyrhizobium* sp. HAMBI 2125 to

determine the growth yield per mol electrons transferred to O_2, N_2O, and to the entire

denitrification pathway. Four treatments were compared: 1) 7% O_2 in headspace; 2) 0% O_2, 4 mM

NO_3^-; 3) 0% O_2 and 10 ml N_2O; 4) 7% O_2, 4 mM NO_3^- and 10 ml N_2O. All treatments had an excess

of electron donor (>70 times more electrons than needed to reduce the electron acceptor). Pre-

cultures were started from single colonies and grown in YMB under fully oxic conditions (21% O_2)

for inoculation of the aerobic batch cultures (treatments 1 and 4), and under anoxic conditions

for inoculation of the anaerobic batches (treatments 2 and 3). The OD_{600} of the aerobic pre-

cultures was kept < 0.3 by regularly inoculating portions of the culture into flasks with fresh

medium. There were 6 replicate flasks for each treatment. Liquid samples were taken from three

of them every 6 h (0.85 ml) for growth measurements (OD_{600}). The three other replicates were

only monitored for gas kinetics and were then used to measure final cell dry weight, to assess

the growth yield per mol electrons to O_2 and N-oxides. For determinination of cell dry weight,

cultures were centrifuged, washed twice in distilled water and dried at 100 °C before weighing.

Cell numbers were determined by microscopic counts at selected sampling times, which gave a

conversion factor of $5.8*10^8$ cells ml^{-1} * OD$_{600}$$^{-1}$. The estimated yield from growth on O_2, N_2O and

NO_3^- was 12.0, 8.5 and 6.7 E12 cells mol^{-1} electrons, respectively (Figs 5, S7). The yields were used

to calculate cell density throughout the incubations, enabling the calculation of cell specific rates

of electron flows to the various electron acceptors, for details see Fig. S7.

Acknowledgements

This project was supported by Kingenta Ecological Engineering Co., LTD. Yuan Gao is grateful to

the China Scholarship Council (CSC) for financial support. We thank Dr. Xinghua Sui from China

Agricultural University for providing seven of the *Bradyrhziobium* strains.

Ethics declarations

558 The authors declare that they have no conflict of interest.

References

Akiyama, H., Hoshino, Y.T., Itakura, M., Shimomura, Y., Wang, Y., Yamamoto, A., *et al*. (2016) Mitigation

of soil N_2O emission by inoculation with a mixed culture of indigenous *Bradyrhizobium diazoefficiens*. *Sci*

Rep **6**: 32869.

Arya, S.S., Salve, A.R., Chauhan, S. (2016) Peanuts as functional food: a review. *J Food Sci Technol* **53**: 31–

41.

Avontuur, J.R., Palmer, M., Beukes, C.W., Chan, W.Y., Coetzee, M.P.A., Blom, J., *et al*. (2019) Genome-

informed *Bradyrhizobium* taxonomy: where to from here? *Syst Appl Microbiol* **42**: 427-439.

Bakken, L.R., Frostegård, Å. (2017) Sources and sinks for N_2O, can microbiologist help to mitigate N_2O

emissions? *Environ Microbiol* **19**: 4801-4805.

Bedmar, E.J., Robles, E.F., Delgado, M.J. (2005) The complete denitrification pathway of the symbiotic,

nitrogen-fixing bacterium *Bradyrhizobium japonicum*. *Biochem Soc Trans* **33**: 141–144.

Bergaust, L., Mao, Y., Bakken, L.R., Frostegård, Å. (2010) Denitrification response patterns during the

transition to anoxic respiration and posttranscriptional effects of suboptimal ph on nitrogen oxide

reductase in *Paracoccus denitrificans*. *Appl Environ Microbiol* **76**: 6387–6396.

Berks, B.C., Ferguson, S.J., Moir, J.W.B., Richardson, D.J. (1995) Enzymes and associated electron

transport systems that catalyse the respiratory reduction of nitrogen oxides and oxyanions. *Biochim*

Biophys Acta **1232**: 97–173.

Bueno, E., Robles, E.F., Torres, M.J., Krell, T., Bedmar, E.J., Delgado, M.J., *et al*. (2017) Disparate

response to microoxia and nitrogen oxides of the *Bradyrhizobium japonicum napEDABC*, *nirK* and

norCBQD denitrification genes. *Nitric Oxide* **68**: 137–149.

Casella, S., Shapleigh, J.P., Toffanin, A., Basaglia, M. (2006) Investigation into the role of the truncated

denitrification chain in *Rhizobium sullae* strain HCNT1. *Biochem Soc Trans* **34**: 130-132.

Chang, Y.L., Wang, E.T., Sui, X.H., Zhang, X.X., Chen, W.X. (2011) Molecular diversity and phylogeny of

rhizobia associated with *Lablab purpureus* (Linn.) grown in Southen China. *Syst Appl Microbiol* **34**: 276-

284.

Chaumeil, P.A., Mussig, A.J., Hugenholtz, P., Parks, D.H. (2020) GTDB-Tk: a toolkit to classify genomes

with the Genome Taxonomy Database. *Bioinformatics* **36**: 1925-1927.

Conthe, M., Lycus, P., Arntzen, M.Ø., Ramos da Silva, A., Frostegård, Å., Bakken, L.R., *et al*. (2019)

Denitrification as an N_2O sink. *Water Res* **151**: 381–387.

Cox, R.D. (1980) Determination of nitrate and nitrite at the parts per billion level by chemiluminescence.

Anal Chem **52**: 332–335.

Davidson, E.A., Kanter, D. (2014) Inventories and scenarios of nitrous oxide emissions. *Environ Res Lett*

9: 105012.

Degefu, T., Wolde-meskel, E., Woliy, K., Frostegård, Å. (2017) Phylogenetically diverse groups of

Bradyrhizobium isolated from nodules of tree and annual legume species growing in Ethiopia. *Syst Appl*

Microbiol **40**: 205-214.

Delgado, M.J., Bonnard, N., Tresierra-Ayala, A., Bedmar, E.J., Müller, P. (2003) The *Bradyrhizobium*

japonicum napEDABC genes encoding the periplasmic nitrate reductase are essential for nitrate

respiration. *Microbiology* **149**: 3395–3403.

Delmont, T.O., Prestat, E., Keegan, K.P., Faubladier, M., Robe, P., Clark, I.M., *et al*. (2012) Structure,

fluctuation and magnitude of a natural grassland soil metagenome. *ISME J* **6**: 1677-1687.

Falk, S., Liu, B., Braker, G. (2010) Isolation, genetic and functional characterization of novel soil *nirK*-type

denitrifiers. *Syst Appl Microbiol* **33**: 337–347.

Gaimster, H., Hews, C.L., Griffiths, R., Soriano-Laguna, M.J., Alston, M., Richardson, D.J., *et al*. (2019) A

central small RNA regulatory circuit controlling bacterial denitrification and N_2O emissions. *mBio* **10**:

e01165-19.

Hall, T.A. (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for

Windows 95/98/NT. *Nucleic Acids Symposium Series* **41***: 95–98.

Hénault, C., Revellin, C. (2011) Inoculants of leguminous crops for mitigating soil emissions of the

greenhouse gas nitrous oxide. *Plant Soil* **346**: 289-296.

Itakura, M., Uchida, Y., Akiyama, H., Hoshino, Y.T., Shimomura, Y., Morimoto, S., *et al*. (2013) Mitigation

of nitrous oxide emissions from soils by *Bradyrhizobium japonicum* inoculation. *Nat Clim Chang* **3**: 208–

212.

Jensen, E.S., Peoples, M.B., Boddey, R.M., Gresshoff, P.M., Hauggaard-Nielsen, H., Alves, J.R., *et al*.

(2012) Legumes for mitigation of climate change and the provision of feedstock for biofuels and

biorefineries. A review. *Agron Sustain Dev*. **32**: 329–364.

Jiménez-Leiva, A., Cabrera, J.J., Bueno, E., Torres, M.J., Salazar, S., Bedmar, E.J., *et al*. (2019) Expanding

the Regulon of the *Bradyrhizobium diazoefficiens* NnrR Transcription Factor: New Insights Into the

Denitrification Pathway. *Front Microbiol* **10**: 1926.

Kumar, S., Stecher, G., Tamura, K. (2016) MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0

for Bigger Datasets. *Mol Biol Evol* **33**: 1870–1874.

Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., Mcgettigan, P.A., McWilliam, H., *et al*. (2007)

Clustal W and Clustal X version 2.0. *Bioinformatics* **23**: 2947–2948.

Li, Y.H., Wang, R., Zhang, X.X., Young, P.W., Wang, E.T., Sui, X.H., *et al*. (2015) *Bradyrhizobium*

guangdongense sp. nov. and *Bradyrhizobium guangxiense* sp.nov., isolated from effectvie nodules of

peanut. *Int J Syst Evol Microbiol* **65**: 4655-4661.

Lindström, K., Mousavi, S.A. (2020) Effectiveness of nitrogen fixation in rhizobia. *Microb Biotechnol* **13**:

1314-1335.

Liu, B., Mao, Y., Bergaust, L., Bakken, L.R., Frostegård, Å. (2013) Strains in the genus *Thauera* exhibit

remarkably different denitrification regulatory phenotypes. *Environ Microbiol* **15**: 2816–2828.

Lycus, P., Bøthun, K.L., Bergaust, L., Shapleigh, J.P., Bakken, L.R., Frostegård, Å. (2017) Phenotypic and

genotypic richness of denitrifiers revealed by a novel isolation strategy. *ISME J* **11**: 2219–2232.

Lycus, P., Soriano-Laguna, M.J., Kjos, M., Richardson, D.J., Gates, A.J., Milligan, D.A., *et al*. (2018) A bet-

hedging strategy for denitrifying bacteria curtails their release of N_2O. *Proc Natl Acad Sci* **115**: 11820–

11825.

MacArthur, P.H., Shiva, S., Gladwin, M.T. (2007) Measurement of circulating nitrite and S-nitrosothiols

by reductive chemiluminescence. *J Chromatogr B* **851**: 93–105.

Mania, D., Woliy, K., Degefu, T., Frostegård, Å. (2020) A common mechanism for efficient N_2O reduction

in diverse isolates of nodule-forming bradyrhizobia. *Environ Microbiol* **22**: 17–31.

Martínez, J., Negrete-Yankelevich, S., Godinez, L.G., Reyes, J., Esposti, M.D., Martínez Romero, E. (2016)

Short-Term Evolution of Rhizobial Strains Toward Sustainability in Agriculture. In *Microbial Models: From*

Environmental to Industrial Sustainability. Springer: Singapore, pp. 277–292.

Mesa, S., Velasco, L., Manzanera, M.E., Delgado, M.J., Bedmar, E.J. (2002) Characterization of the

norCBQD genes, encoding nitric oxide reductase, in the nitrogen fixing bacterium *Bradyrhizobium*

japonicum. *Microbiology* **148**: 3553–3560.

Molstad, L., Dörsch, P., Bakken, L.R. (2016) Improved robotized incubation system for gas kinetics in

batch cultures. *ResearchGate*. https://doi.org/10.13140/RG.2.2.30688.07680.

Molstad, L., Dörsch, P., Bakken, L.R. (2007) Robotized incubation system for monitoring gases (O_2, NO,

N_2O N_2) in denitrifying cultures. *J Microbiol Methods* **71**: 202–211.

Obando, M., Correa-Galeote, D., Castellano-Hinojosa, A., Gualpa, J., Hidalgo, A., Alché, J.D., *et al*. (2019)

Analysis of the denitrification pathway and greenhouse gases emissions in *Bradyrhizobium* sp. strains

used as biofertilizers in South America. *J Appl Microbiol* **127**: 739–749.

Rambaut, A., Drummond, A.J., Xie, D., Baele, G., Suchard, M.A. (2018) Posterior summarization in

Bayesian phylogenetics using Tracer 1.7. *Syst Biol* **67**: 901–904.

Reay, D.S., Davidson, E.A., Smith, K.A., Smith, P., Melillo, J.M., Dentener, F., *et al*. (2012) Global

agriculture and nitrous oxide emissions. *Nat Clim Chang* **2**: 410–416.

Richardson, D.J. (2000) Bacterial respiration: a flexible process for a changing environment. *Microbiology*

146: 551–571.

Richardson, D.J., Berks, B.C., Russell, D.A., Spiro, S., Taylor, C.J. (2001) Functional, biochemical and

genetic diversity of prokaryotic nitrate reductases. *Cell Mol Life Sci* **58**: 165–178.

Rogelj, J., Shindell, D., Jiang, K., Fifita, S., Forster, P., Ginzburg, V., *et al*. (2018) Mitigation pathways

compatible with 1.5 °C in the context of sustainable development. In, Masson-Delmotte V, Zhai P,

Pörtner HO, Roberts D, Skea J, Shukla PR, et al. (eds). *Global Warming of 1.5 °C. An IPCC Special Report*

on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas

emission pathways, in the context of strengthening the global response to the threat ofclimate change,

sustainable development, and efforts to eradicate poverty;

https://www.ipcc.ch/site/assets/uploads/sites/2/2019/02/SR15_Chapter2_Low_Res.pdf.

Ronquist, F., Teslenko, M., Van Der Mark, P., Ayres, D.L., Darling, A., Höhna, S., *et al*. (2012) Mrbayes 3.2:

Efficient bayesian phylogenetic inference and model choice across a large model space. *Syst Biol* **61**:

539–542.

Sameshima-Saito, R., Chiba, K., Minamisawa, K. (2004) New method of denitrification analysis of

Bradyrhizobium field isolates by gas chromatographic determination of ^{15}N-labeled N_2. *Appl Environ*

Microbiol **70**: 2886–2891.

Sánchez, C., Minamisawa, K. (2018) Redundant roles of *Bradyrhizobium oligotrophicum* Cu-type (NirK)

and cd1-type (NirS) nitrite reductase genes under denitrifying conditions. *FEMS Microbiol Lett* **365**:

fny015.

Santos, M.S., Nogueira, M.A., Hungria, M. (2019) Microbial inoculants: reviewing the past, discussing the

present and previewing an outstanding future for the use of beneficial bacteria in agriculture. *AMB Expr*

9: 205.

Schlesinger, W.H. (2009) On the fate of anthropogenic nitrogen. *Proc Natl Acad Sci* **106**: 203–208.

Schwartz, W. (1972) Vincent, J.M, a manual for the practical study of the root-nodule bacteria (IBP

Handbuch no. 15 des international biology program, London). XI u. 164 S., 10 Abb., 17 tab., 7 Taf.

Oxford-Edinburgh 1970: Blackwell scientific Publ.,45 s. *Z Allg Mikrobiol* **12**: 1521–4028.

Shapleigh, J.P. (2013) The prokaryotes: Prokaryotic physiology and biochemistry. In Rosenberg E, DeLong

EF, Lory S, Stackebrandt E, Thompson F (eds). *The prokaryotes.* Springer: Berlin Heidelberg, pp 405–425.

Sistrom, W.R. (1962) The kinetics of the synthesis of photopigments in Rhodopseudomonas spheroides.

J Gen Microbiol **28**: 607–616.

Thompson, R.L., Lassaletta, L., Patra, P.K., Wilson, C., Wells, K.C., Gressent, A., *et al*. (2019) Acceleration

of global N$_2$O emissions seen from two decades of atmospheric inversion. *Nat Clim Chang* **9**: 993–998.

Thomson, A.J., Giannopoulos, G., Pretty, J., Baggs, E.M., Richardson, D.J. (2012) Biological sources and

sinks of nitrous oxide and strategies to mitigate emissions. *Philos Trans R Soc B.* **367**: 1157-1168.

Thrall, P.H., Laine, A-L., Broadhurst, L.M., Bagnall, D.J., Brockwell, J. (2011) Symbiotic effectiveness of

rhizobial mutualists varies in interactions with native Australian legume genera. *PLoS ONE* **6**: e23545.

Tian, H., Xu, R., Canadell, J.G., Thompson, R.L., Winiwarter, W., Suntharalingam, P., *et al*. (2020) A

comprehensive quantification of global nitrous oxide surces and sinks. *Nature* **586**: 248-256.

Torres, M.J., Bueno, E., Jiménez-Leiva, A., Cabrera, J.J., Bedmar, E.J., Mesa, S., *et al*. (2017) FixK$_2$ Is the

Main Transcriptional Activator of *Bradyrhizobium diazoefficiens nosRZDYFLX* Genes in Response to Low

Oxygen. *Front Microbiol* **8**: 1621.

Torres, M.J., Simon, J., Rowley, G., Bedmar, E.J., Richardson, D.J., Gates, A.J., *et al*. (2016) Nitrous oxide

metabolism in nitrate-reducing bacteria: physiology and regulatory mechanisms. In: Poole RK (eds).

Advances in microbial physiology, first edition*.* Elsevier, pp 353-432.

Van Rossum, D., Muyotcha, A., Van Verseveld, H.W., Stouthamer, A.H., Boogerd, F.C. (1993) Effects of

Bradyrhizobium strain and host genotype, nodule dry weight and leaf aera on groundnut *(Arachis*

hypogaea L. ssp. fastigiata) yield. *Plant and Soil* **154**: 279-288.

Van Rossum, D., Muyotcha, A., Van Verseveld, H.W., Stouthamer, A.H., Boogerd, F.C. (1994) Siderophore

production by *Bradyrhizobium* spp. Strains nodulating groundnut. *Plant and soil* **163**: 177-187.

Van Spanning, R., Richardson, D., Ferguson, S. (2007) Introduction to the biochemistry and molecular

biology of denitrification. In: Bothe H,Ferguson SJ, Newton WE (eds). *Biology of the nitrogen cylce*.

Elsevier: Amsterdam, The Netherlands, pp 3–21.

Velasco, L., Mesa, S., Delgado, M.J., Bedmar, E.J. (2001) Characterization of the *nirK* gene encoding the

respiratory, Cu-containing nitrite reductase of *Bradyrhizobium japonicum*. *Biochim Biophys Acta* **1521**:

130–134.

Velasco, L., Mesa, S., Xu, C., Delgado, M.J., Bedmar, E.J. (2004) Molecular characterization of

nosRZDFYLX genes coding for denitrifying nitrous oxide reductase of *Bradyrhizobium japonicum*. *Antonie*

Van Leeuwenhoek **85**: 229–235.

Wang, R., Chang, Y.L., Zheng, W.T., Zhang, D., Zhang, X.X., Sui, X.H., *et al*. (2013) *Bradyrhizobium*

arachidis sp. Nov., isolated from effective nodules of Arachis hypogaea grown in China. *Syst Appl*

Microbiol **36**: 101-105.

Winiwarter, W., Höglund-Isaksson, L., Klimont, Z., Schöpp, W., Amann, M. (2018) Technical opportunities

to reduce global anthropogenic emissions of nitrous oxide. *Environ Res Lett* **13**: 014011.

Woliy, K., Degefu, T., Frostegård, Å. (2019) Host Range and Symbiotic Effectiveness of N_2O Reducing

Bradyrhizobium Strains. *Front Microbiol* **10**: 2746.

Xu, Z.F., Hansen, M.A., Hansen, L.H., Jacquiod, S., Sørensen, S.J. (2014) Bioinformatic Approaches Reveal

Metagenomic Characterization of Soil Microbial Community. *PLoS ONE* **9**: e93445.

Zhang, L., Wüst, A., Prasser, B., Müller, C., Einsle, O. (2019) Functional assembly of nitrous oxide

reductase provides insights into copper site maturation. *Proc Natl Acad Sci* **116**: 12822–12827.

Zhang, X.P., Nick, G., Kaijalainen, S., Terefework, Z., Paulin, L., Tighe, S.W., *et al*. (1999) Phylogeny and

Diversity of *Bradyrhizobium Strains* isoalted from the Root Nodules of Peanut *(Arachis hypogaea)* in

Sichuan, China. *Syst Appl Microbiol* **22**: 378-386.

Zilli, .JÉ., Alves, B.J.R., Rouws, L.F.M., Simões-Araujo, J.L., de Barros Soares, L.H., Cassán, F., *et al*. (2019)

The importance of denitrification performed by nitrogen-fixing bacteria used as inoculants in South

America. *Plant Soil* **451**: 5-24.

Zumft, W.G. (1997) Cell biology and molecular basis of denitrification. *Microbiol Mol Biol Rev* **61**: 533–

616.

 Table and Figure legends

Table 1. Strains used in the current study. All listed strains were included in the MLSA-based phylogenetic

analysis (Fig. 1). The strains above the dashed line were also examined in the present study with regard

to their denitrification phenotypes. Strains AC 70c, AC 86d2, AC 86j1 and AC 101b were not examined for

denitrification since this was reported elsewhere (Mania *et al*., 2020).

Fig. 1. Phylogenetic tree of four concatenated housekeeping genes (*atpD- glnII- recA- rpoB*) from 78

Bradyrhizobium strains constructed based on a Bayesian inference analysis. Only the posterior

probabilities ≥0.95 are shown in the tree. The genus name *Bradyrhizobium* is abbreviated as *B*., and a "T"

at the end of each strain code shows the type strains. In cases where two or more isolates showed high

whole genome sequence similarity (>99 %) based on pairwise comparison of Average Nucleotide Identity

(ANI) (see Table S1), only one representative is shown in this tree. Colors indicate denitrification end-

products. A sequence analysis of the genes in the respective denitrification gene operons showed good

correspondence between genetic set-up and denitrification end products (Table S2). The only exception

was HAMBI 2299, which had a complete *nor* operon but did not reduce NO. Another non-NO reducing

strain, HAMBI 2153, lacked two genes in the *nor* operon.

Fig. 2. Competition for electrons between N_2O- and NO_3^- reductase; comparison of Nap- and Nar-carrying

organisms. Gas kinetics and corresponding electron flow to oxidases (aerobic respiration) and to the

different N-reductases (denitrification) in batch cultures of *Bradyrhizobium* strain HAMBI 2125 and

Paracoccus denitrificans. Both organisms perform complete denitrification but *Bradyrhizobium* (A, C)

carries only the periplasmic NapA for dissimilatory NO_3^- reduction while *P. denitrificans* (B, D) carries both

NapA and the membrane-bond NarG. The top panels (A, B) show the measured amounts of NO_2^-, NO, N_2O

and N_2 per flask, and the concentration of O_2 in the liquid. Note that the small but significant decrease in

N_2O during the aerobic phase was caused by sampling losses, not by N_2O reduction or other reactions, as

explained in Fig. S6. Background colors indicate the dominating respiratory activity. Blue: O_2 respiration;

Pink: N_2O respiration; Green: Complete denitrification (NO_3^- to N_2). The bottom panels (C, D) show the

calculated cell specific rates of electron flow (fmol cell^{-1} h^{-1}) to the terminal oxidases (V_{eO2}) and to the four

reductases ($V_{eNar/Nap}$, V_{eNir}, V_{eNor} and V_{eNos}). V_{eNir} and V_{eNor} were practically identical throughout (NO

remained very low), thus indistinguishable in the panels. Strain HAMBI 2125 was incubated in 4 replicate

flasks with YMB supplemented with 1 mM KNO_3, of which only one is shown here (gas- and NO_2^- kinetics

of replicate flasks are shown in Fig. S3). *P. denitrificans* was incubated in 50 ml Sistrom's medium with 2

mM KNO_3 (n=7, bars show standard errors). At the start of the incubation, all flasks contained ~1% O_2 in

the headspace (~10 μM in liquid) and ~ 1.2 ml N_2O (~90 μmol N_2O-N). $V_{eNar/Nap}$ and V_{eNir} represent rates

of electron transport to NO_3^- and NO_2^-, respectively (via NapA and NirK for *Bradyrhziobium* and via

NarG+Nap and NirS for *P. denitrificans*).

Fig. 3. Effect of injecting N_2O to an actively denitrifying culture. Gas kinetics for *Bradyrhizobium* strain

HAMBI 2125 during and after transition from aerobic respiration to N-oxide respiration. Triplicate cultures

were incubated in YMB medium supplemented with 2 mM KNO_3 and with 0.7 ml O_2 in the headspace

(around 10 μM O_2 in the medium). Approximately 4 ml N_2O (~300 μmol N_2O-N) were injected at 32 hpi,

indicated by the arrow, into actively denitrifying cultures respiring NO_3^-. Bars show standard errors (n=3).

Fig. 4. Abundance of denitrification reductases during and after transition from aerobic to anaerobic

respiration. The abundances of Nap, NirK and Nos were determined by proteomics and calculated as

percentages of the sum of all LFQ (label-free quantitation of proteins) abundances for each time point.

The first samples were taken from triplicate flasks of *Bradyrhizobium* strain HAMBI 2125 growing under

fully oxic conditions (21% O_2). Subsamples from these precultures were inoculated into flasks with 1% O_2,

and samples for proteomics were taken when the cultures approached O_2 depletion ([O_2] = 5.8±0.2 μM)

and during the anaerobic respiration. Concentrations of O_2 (grey shading) and N_2O (-Δ-) are schematically

indicated here to illustrate the time of their depletion; for actual values of [O_2] [N_2O], [N_2] and [NO_2^-], see

Fig. S8. Background colors indicate the dominating respiratory activity. Blue: O_2 respiration; Pink: N_2O

respiration; Green: Complete denitrification (NO_3^- to N_2). Bars show standard deviation (n=3).

Fig. 5. Growth and yield of *Bradyrhizobium* strain HAMBI 2125 when incubated in YMB at 28 °C and

provided with O_2, N_2O or NO_3^- as electron acceptor. All relevant gases were measured frequently using a

robotized system. Panels A-C show estimated g dry weight (DW) flask^{-1} throughout the incubations either

calculated from measured OD_{600} (□) using the factor 0.374 mg dry weight ml^{-1} OD_{600}^{-1} (Fig. S7) or calculated

from the measured cumulated electron flow (solid line) using measured yields per mol e$^-$ in panel D. The

specific growth rates (μ) were determined by regression of ln(OD_{600}) against time during the period of

exponential growth. Panel A. Growth using O_2 as sole electron acceptor (initial amount 180 μmol flask^{-1}).

<u>Panel B</u>. Growth using N_2O as sole electron acceptor (initial amount 800 µmol N flask^{-1}). <u>Panel C</u>. Growth

from complete denitrification (NO_3^- to N_2; initial amount of NO_3^- was 200 µmol flask^{-1}). <u>Panel D</u>. Yield per

mol e$^-$ obtained from growth using the different electron acceptors. The differences in yield were

statistically different (p<0.01). O_2 was depleted in the O_2 treatment after 55 h, while N_2O and NO_3^- were

not depleted during the incubations. Bars indicate standard deviation, n=3. Note: The initial value for DW,

estimated by cumulated electron flow, was set equal to that based on measured OD.

Fig. 1. Phylogenetic tree of four concatenated housekeeping genes (*atpD- glnII- recA- rpoB*) from 78 *Bradyrhizobium* strains constructed based on a Bayesian inference analysis. Only the posterior probabilities ≥0.95 are shown in the tree. The genus name *Bradyrhizobium* is abbreviated as *B*., and a "T" at the end of each strain code shows the type strains. In cases where two or more isolates showed high whole genome sequence similarity (>99 %) based on pairwise comparison of Average Nucleotide Identity (ANI) (see Table S1), only one representative is shown in this tree. Colors indicate denitrification end-products. A sequence analysis of the genes in the respective denitrification gene operons showed good correspondence between genetic set-up and denitrification end products (Table S2). The only exception was HAMBI 2299, which had a complete *nor* operon but did not reduce NO. Another non-NO reducing strain, HAMBI 2153, lacked two genes in the *nor* operon.

Fig. 2. Competition for electrons between N_2O- and NO_3^- reductase; comparison of Nap- and Nar-carrying organisms. Gas kinetics and corresponding electron flow to oxidases (aerobic respiration) and to the different N-reductases (denitrification) in batch cultures of *Bradyrhizobium* strain HAMBI 2125 and *Paracoccus denitrificans*. Both organisms perform complete denitrification but *Bradyrhizobium* carries only the periplasmic NapA for dissimilatory NO_3^- reduction while *P. denitrificans* carries both NapA and the membrane-bond NarG. The top panels (A, B) show the measured amounts of NO_2^-, NO, N_2O and N_2 per flask, and the concentration of O_2 in the liquid. Note that the small but significant decrease in N_2O during the aerobic phase was caused by sampling losses, not by N_2O reduction or other reactions, as explained in Fig. S6. Background colors indicate the dominating respiratory activity. Blue: O_2 respiration; Pink: N_2O respiration; Green: Complete denitrification (NO_3^- to N_2). The bottom panels (C, D) show the calculated cell specific rates of electron flow (fmol cell^{-1} h^{-1}) to the terminal oxidases (V_{eO2}) and to the four reductases ($V_{eNar/Nap}$, V_{eNir}, V_{eNor} and V_{eNos}). V_{eNir} and V_{eNor} were practically identical throughout (NO remained very low), thus indistinguishable in the panels. Strain HAMBI 2125 was incubated in 4 replicate flasks with YMB supplemented with 1 mM KNO_3, of which only one is shown here (gas- and NO_2^- kinetics of replicate flasks are shown in Fig. S3). *P. denitrificans* was incubated in 50 ml Sistrom's medium with 2 mM KNO_3 (n=7, bars show standard errors). At the start of the incubation, all flasks contained ~1% O_2 in the headspace (~10 µM in liquid) and ~ 1.2 ml N_2O

(~90 µmol N_2O-N). $V_{eNar/Nap}$ and V_{eNir} represent rates of electron transport to NO_3^- and NO_2^-, respectively (via NapA and NirK for *Bradyrhziobium* and via NarG+Nap and NirS for *P. denitrificans*).

Fig. 3. Effect of injecting N_2O to an actively denitrifying culture. Gas kinetics for *Bradyrhizobium* strain HAMBI 2125 during and after transition from aerobic respiration to N-oxide respiration. Triplicate cultures were incubated in YMB medium supplemented with 2 mM KNO_3 and with 0.7 ml O_2 in the headspace (around 10 μM O_2 in the medium). Approximately 4 ml N_2O (~300 μmol N_2O-N) were injected at 32 hpi, indicated by the arrow, into actively denitrifying cultures respiring NO_3^-. Bars show standard errors (n=3).

Fig. 4. Abundance of denitrification reductases during and after transition from aerobic to anaerobic respiration. The abundances of Nap, NirK and Nos were determined by proteomics and calculated as percentages of the sum of all LFQ (label-free quantitation of proteins) abundances for each time point. The first samples were taken from triplicate flasks of *Bradyrhizobium* strain HAMBI 2125 growing under fully oxic conditions (21% O_2). Subsamples from these precultures were inoculated into flasks with 1% O_2, and samples for proteomics were taken when the cultures approached O_2 depletion ([O_2] = 5.8±0.2 μM) and during the anaerobic respiration. Concentrations of O_2 (grey shading) and N_2O (-Δ-) are schematically indicated here to illustrate the time of their depletion; for actual values of [O_2] [N_2O], [N_2] and [NO_2^-], see Fig. S8. Background colors indicate the dominating respiratory activity. Blue: O_2 respiration; Pink: N_2O respiration; Green: Complete denitrification (NO_3^- to N_2). Bars show standard deviation (n=3).

Fig. 5. Growth and yield of *Bradyrhizobium* strain HAMBI 2125 when incubated in YMB at 28 °C and provided with O_2, N_2O or NO_3^- as electron acceptor. All relevant gases were measured frequently using a robotized system. Panels A-C show estimated g dry weight (DW) flask^{-1} throughout the incubations either calculated from measured OD_{600} ($\square$) using the factor 0.374 mg dry weight ml^{-1} OD_{600}^{-1} (Fig. S7) or calculated from the measured cumulated electron flow (solid line) using measured yields per mol e$^-$ in panel D. The specific growth rates (μ) were determined by regression of ln(OD_{600}) against time during the period of exponential growth. Panel A. Growth using O_2 as sole electron acceptor (initial amount 180 µmol flask^{-1}). Panel B. Growth using N_2O as sole electron acceptor (initial amount 800 µmol N flask^{-1}). Panel C. Growth from complete denitrification (NO_3^- to N_2; initial amount of NO_3^- was 200 µmol flask^{-1}). Panel D. Yield per mol e$^-$ obtained from growth using the different electron acceptors. The differences in yield were statistically different (p<0.01). O_2 was depleted in the O_2 treatment after 55 h, while N_2O and NO_3^- were not depleted during the incubations. Bars indicate standard deviation, n=3. Note: The initial value for DW, estimated by cumulated electron flow, was set equal to that based on measured OD.

Table 1. Strains used in the current study. All listed strains were included in the MLSA-based phylogenetic analysis (Fig. 1). The strains above the dashed line were also examined in the present study with regard to their denitrification phenotypes. Strains AC 70c, AC 86d2, AC 86j1 and AC 101b were not examined for denitrification since this was reported elsewhere (Mania *et al.*, 2020).

Strain*	Host plant	Geographic origin	Reference
HAMBI 2115	*Arachis hypogaea* L.	Zimbabwe	(Zhang *et al.*, 1999)
HAMBI 2116	*A. hypogaea* L.	Zimbabwe	(Zhang *et al.*, 1999)
HAMBI 2125	*A. hypogaea* L.	Sichuan, China	(Zhang *et al.*, 1999)
HAMBI 2127	*A. hypogaea* L.	Sichuan, China	(Zhang *et al.*, 1999)
HAMBI 2128	*Macrotyloma africanum*	Zimbabwe	(Van Rossum *et al.*, 1993)
HAMBI 2130	*Vigna unguiculata*	Zimbabwe	(Van Rossum *et al.*, 1994)
HAMBI 2134	*A. hypogaea* L.	Sichuan, China	(Zhang *et al.*, 1999)
HAMBI 2142	*A. hypogaea* L.	Sichuan, China	(Zhang *et al.*, 1999)
HAMBI 2149	*A. hypogaea* L.	Sichuan, China	(Zhang *et al.*, 1999)
HAMBI 2153	*A. hypogaea* L.	Israel	(Zhang *et al.*, 1999)
HAMBI 2299	*A. hypogaea* L.	Hawaii, USA	HAMBI**
HAMBI 3052^T	*Lablab purpureus*	Anhui, China	(Chang *et al.*, 2011)
CCBAU 051107^T	*A. hypogaea* L.	Hebei, China	(Wang *et al.*, 2013)
CCBAU 23155	*A. hypogaea* L.	Anhui, China	(Wang *et al.*, 2013)
CCBAU 33067	*A. hypogaea* L.	Jiangxi, China	(Wang *et al.*, 2013)
CCBAU 45332	*A. hypogaea* L.	Henan, China	(Wang *et al.*, 2013)
CCBAU 53344	*A. hypogaea* L.	Guangxi, China	(Li *et al.*, 2015)
CCBAU 53363^T	*A. hypogaea* L.	Guangxi, China	(Li *et al.*, 2015)
CCBAU 53429	*A. hypogaea* L.	Guangxi, China	(Li *et al.*, 2015)
AC70c	*Phaseolus vulgaris*	Akaki, Ethiopia	(Degefu et al., 2017)
AC86d2	*Cajanus cajan*	Awassa, Ethiopia	(Degefu et al., 2017)
AC87j	*Millettia ferruginea*	Awassa, Ethiopia	(Degefu et al., 2017)
AC101b	*Acacia saligna*	Leku, Ethiopia	(Degefu et al., 2017)

HAMBI: The culture collection at the Department of Microbiology, University of Helsinki.

CCBAU: Culture Collection of China Agricultural University, Beijing, China.

AC: The strains are kept in a local collection at Norwegian University of Life Sciences and available upon request (Å. Frostegård).

*) For information on donator and specimen history of HAMBI strains please consult https://kotka.luomus.fi/culture/bac

**) https://kotka.luomus.fi/view?uri=http://tun.fi/RAH.1747&type=document&page=1&spot=1