1 **Targeting of Mammalian Glycans Enhances Phage Predation in the**
2 **Gastrointestinal Tract**
3

Sabrina I. Green[1], Carmen Gu Liu[1], Xue Yu[1], Shelley Gibson[2], Wilhem Salmen[1],

Anubama Rajan[1], Hannah E. Carter[1], Justin R. Clark[1], Xuezheng Song[3], Robert F.

Ramig[1], Barbara W. Trautner[4,5], Heidi B. Kaplan[6], Anthony W. Maresso[1#]

Address correspondence to Anthony Maresso, maresso@bcm.edu.

[1]Department of Molecular Virology and Microbiology, Baylor College of Medicine, Houston, TX,

USA

[2]Department of Human Genetics, Baylor College of Medicine, Houston, TX, USA

[3]Department of Biochemistry, Emory Comprehensive Glycomics Core, Emory University School

of Medicine, Atlanta, GA, USA.

[4]Michael E. Debakey Veterans Affairs Medical Center, Houston, TX, USA.

[5]Department of Medicine, Baylor College of Medicine, Houston, TX, USA.

[6]Department of Microbiology and Molecular Genetics, McGovern Medical School, University of

Texas Health Science Center at Houston, Houston, Texas, USA.

Key Words

Phage therapy, *E. coli*, multidrug-resistance, bacteriophages, pathogen, pathobiont

Graphical Abstract

Model showing (1) mucins from the intestinal mucus layer inhibit phage infection, (2) phage
ES17 can bind to mucin and utilize other intestinal glycans as a receptor to infect and kill
mucus-coated bacteria, and (3) phages like ES17 can be utilized to coat the intestinal
epithelium by binding heparan sulfate glycans to protect from invasive pathogen infection.
33

Abstract

The human mucosal surface consists of a eukaryotic epithelium, a prokaryotic

microbiota, and a carbohydrate-rich interface that separates them. Bacteriophage

parasitize the prokaryotes but are not known to associate with eukaryotic cells. In the

gastrointestinal tract, the interaction of these two domains influences the health of the

host, especially colonization with invasive pathobionts. Antibiotics may be used but they

also kill protective commensals and lack the physio-chemical properties to be

specifically and optimally active in this complex milieu. Here, we report a novel phage

whose lytic cycle is enhanced in intestinal environments. The enhanced activity is

encoded in its tail fiber gene, whose protein product binds human heparan sulfated

proteoglycans and localizes the phage to the epithelial cell surface, thereby positioning

it near its bacterial host, a type of locational targeting mechanism. This finding offers the

prospect of developing epithelial-targeting phage to selectively remove invasive

pathobiont species from mucosal surfaces.

Introduction

The human mucosal surface is a diverse community comprised of bacteria, viruses,

fungi, and human epithelial, immune, and stem cells. The chemistry of the surface is

complex—commonly composed of proteins, lipids, nucleic acid, and small molecule

metabolites (Sansonetti, 2004; Nataro, 2005). The pH, ionic environment, and physio-

mechano properties also contribute to this dynamic surface. All of these factors

influence the biology, and, in turn, disease and treatment. Research over the last

decade has revealed the importance of mucosal homeostasis. Invasive pathobionts

may colonize, grow, and cause acute or chronic infections, some of which may become

systemic and life-threatening (Johnson JR, 2002; Russo and Johnson, 2003). The

bacterial composition of this intestinal community varies in association with human

illnesses such as cancer, diabetes, neurological illness, obesity, and cardiovascular

disease (De Martel *et al.*, 2012; Sampson and Mazmanian, 2015; Maruvada *et al.*,

2017; Vatanen *et al.*, 2018; Tang, Li and Hazen, 2019).

Antibiotics have revolutionized medicine. The meteoric rise of multidrug resistance

threatens a returns to pre-antibiotic days by 2050 (O'Neill, 2016). In addition, it is

increasingly recognized that antibiotics act as a commensal destroying napalm. The

antibiotics developed commercially have focused on inhibition of key processes

common to all bacterial species, for example, the inhibition of protein or cell wall

biosynthesis. Current antibiotics lack the specificity to selectively target a causative

pathobiont, which in the mucosal environment, might be the major driver of the disease

among the backdrop of hundreds of benign, even beneficial, symbionts. Moreover, the

core chemical backbones of antibiotics (e.g. β-lactam antibiotics) somewhat limit the

ability to develop new modified versions. Furthermore, these modifications improve

enzymatic activity (e.g. increase affinity or induced suicide inhibition) but has not

enhanced the drug's activity in the human mucosal environment. Mucosal-active,

targeted drugs that specifically control the bacterial composition of this complex

environment without perturbing ecological balance would promote the health of the

nasopharyngeal, lower respiratory, gastrointestinal, and urogenital epithelium.

The rise of multidrug resistance has precipitated the search for alternative antibacterial

approaches, including the use of bacteriophages (phages) which are bacterial viruses

used to treat infections. Although phage therapy has been used in Eastern Europe for

decades, only in the past few years has it been employed for compassionate use cases

in the U.S. and Europe and in clinical trials (Schooley *et al.*, 2017; Dedrick *et al.*, 2019;

Jault *et al.*, 2019). In theory, phage offer numerous advantages over antibiotics,

including that they can be specific towards a given species of bacteria resulting in

potential microbiome sparing, "generally regarded as safe" or GRAS attributes, allowing

them to be given in high doses, and diverse, as phage mixtures can be used to either

broaden the range or reduce the frequency of resistance or both (Loc-Carrillo and

Abedon, 2011; Grose and Casjens, 2014; Moye, Woolston and Sulakvelidze, 2018).

However, their greatest attribute has been speculated to be their abundance –

estimated to be 10^{31} on Planet Earth. The phage genosphere—all the collection of

genes and their unique biology—sometimes referred to as "viral dark matter", may

encode novel features that allow for enhanced lytic activity towards bacterial pathogens

(Youle, Haynes and Rohwer, 2012; Brum *et al.*, 2016; Terwilliger *et al.*, 2020).

Reasoning that this genosphere may include phage with unique phenotypes that

facilitate lytic activity against pathobionts at mucosal environments, we report here a

novel podovirus of the genus *Kuravirus* that infects pathogenic *E. coli* using positional

targeting to human heparan-sulfated proteoglycans.

Results

The gastrointestinal tract is prohibitive to phage therapy

Escherichia coli are gram-negative bacilli found in a variety of environments, including

the human intestinal microbiome, with > 90% of people colonized (Mitsuoka, Hayakawa

and Kimura, 1975). Although most strains are benign, *E. coli* is particularly troubling

because of its propensity to undergo horizontal transfer of genes encoding virulence

factors and antimicrobial resistance (Huang *et al.*, 2001; Antão, Wieler and Ewers,

2009; Price *et al.*, 2013; Mathers, Peirano and Pitout, 2015; Poole *et al.*, 2017). Thus,

several pathotypes have arisen that are associated with human disease. InPEC

(Intestinal pathogenic *E. coli*) are associated with diarrhea and gastroenteritis. ExPEC

(Extraintestinal pathogenic *E. coli*) are associated with systemic infections of the urinary

tract, brain, peritoneum, peripheral organs, blood, and in-dwelling devices, resulting in 9

million infections per year (Russo and Johnson, 2003; Colpan *et al.*, 2013; Poolman and

Wacker, 2016). The pandemic ExPEC sequence type 131 or ST131 possess a rare

combination of multidrug resistance and enhanced virulence (Pitout, 2012; Johnson *et*

al., 2013; Banerjee and Johnson, 2014). These strains readily colonize the human

intestine, which can become a reservoir, prior to extraintestinal infection (Nicolas-

Chanoine, Bertrand and Madec, 2014). Previous studies reported that phage HP3, a

lytic myovirus isolated from environmental reservoirs of ExPEC, could reduce ST131

bacteremia and disease severity in murine models of infection (Green *et al.*, 2017a; Ma

et al., 2018). Since the human gastrointestinal tract is the primary reservoir of ExPEC

ST131, we wondered if phage HP3 could act prophylactically to reduce or eliminate

ExPEC burden in the intestine. To test this, mice were orally gavaged with an ExPEC

ST131 clinical isolate then treated with phage or an antibiotic as illustrated in Figure 1A.

Untreated mice sustained stable bacterial colonization during the course of the

experiment (6 days) (Figure 1B). When phage HP3 were given to animals via water or a

daily gavage, the levels of ExPEC were indistinguishable from the untreated control at

the end of the experiment. However, no ExPEC was detected at any time point in the

antibiotic-treated group. Interestingly, phage HP3 was detected, and active, in the stool

of phage-treated mice, even on day 4, indicating the lack of ExPEC reduction was not

due to a lack of delivery to the intestinal environment or to inactivation of the phage

(Figure 1C). It should also be noted that despite having higher levels of phage upon

gavage during days 1-4, the phage was no more effective at removing ExPEC than

phage given in the water, suggesting that in this experiment there was no dose

dependent effect of phage. In addition, as much as 10^5 (PFU/g) phage were found in the

murine intestinal tissue (including cecum and colon) on day 6, the final day of the study,

yet there was little to no clearing of ExPEC compared to the untreated control in these

tissues (Figures 1D and 1E). Importantly, antibiotic treatment significantly reduced the

number of operational taxonomic units (OTUs) and diversity, as determined by the

Shannon diversity index, which takes into account species richness and distribution

(Figure 1Fi, ii, $p = 0.007$, $p = 0.009$). Also, a Principle component analysis (PCA) of beta

diversity demonstrated that mice in the antibiotic cohort clustered together and away

from the untreated and phage groups, suggesting antibiotics had a more profound effect

on the microbiome than did phage (Figure 1Fiii). Finally, we assessed phage killing in a

modified "cecal medium" or CM that is derived from the murine cecal contents and

designed to simulate the luminal complexity of the mammalian intestine (Figure 1G).

Whereas phage HP3 completely abolished ExPEC in LB (nearly a 9-log drop in levels

and no detectable live bacteria), and nearly abolished it in a slurry of fecal pellets taken

from the same murine host (~ 8-log drop in levels), there was little to no phage-based

killing in CM, despite recovering nearly 10^6-$10^{7.5}$ PFU/mL (Figures 1H and 1I) of phage.

The result was the same when the experiment was repeated anaerobically (data not

shown). These results mirrored those from the ExPEC colonization model and indicate

there is/are factor(s) present in the mammalian GI that inhibit this lytic phage.

Figure 1

Figure 1: The gastrointestinal tract is prohibitive to phage therapy
(A) Mouse colonization diagram. (B) Intestinal (fecal) colonization of ExPEC (C) and phage HP3
levels. (D) Day 6 intestinal tissue HP3 levels and (E) ExPEC levels. (Fi) OTU from 16S rRNA
analysis of fecal pellets day 6. (Fii) Shannon diversity values (Fiii) PCA of beta diversity.
Symbols (squares) indicate individual mice, except in (B) squares indicate mean. Exp.
performed 1X. N=8-10 where N indicates individual mice. **** p < 0.0001, NS not significant," =
none detected, # missing values due to histological analysis of 2 mice/group. Mean, ±SD
shown. (G) Cecal medium (CM) development. (H) ExPEC levels after 4.5 hours growth and (I)
phage HP3 levels. Experiment performed 1X. Squares (N=3) indicate individual biological
replicates from independent cultures. Mean, ±SD shown.

The inhibitory component is mucin

We wished to understand the reasons phage HP3 was ineffective in this intestinal

microenvironment. Previously, we had shown that phage HP3 utilizes calcium for

efficient infection, suggesting the effect may be nutritionally-based (Ma *et al.*, 2018).

However, addition of calcium did not increase phage HP3 killing in CM (data not

shown). We next tested whether the inhibition might be related to the presence of live

bacterial microbiota in CM. However, ExPEC killing with phage was not enhanced with

removal of the microbiota with a broad range of antibiotics, including inhibitors of protein

synthesis, cell wall, and DNA synthesis (Figure S1i-iii – note that the antibiotics

efficiently killed a commensal, antibiotic-sensitive, *E. coli* that was spiked into the CM,

iv). Furthermore, the inhibitory effect was maintained when a related, and equally as

effective phage EC1 was used in the same experiment, thereby indicating the inhibitory

effect was not solely due something specific for phage HP3 (data not shown). Arriving at

no resolution as to what the inhibitory factor may be, we decided to test more drastic

treatments for their ability to restore phage killing in CM. First, the CM was heat-treated

(Figure 2Ai, HT CM). Interestingly, heat treatment led to greater than a 6-log

improvement in ExPEC killing by phage HP3 (Figure 2Aii, $p < 0.0001$). Similarly, when

CM was filter-treated (Figure 2Ai, FT CM), there was no detectable levels of ExPEC in

the medium after treatment with phage HP3 (Figure 2Aii), and, this observation was

extended to another phage showing inhibition in CM, EC1 (Figure S2A, $p < 0.0001$).

Thus, the inhibitory component was large (retained on 0.22 micron filter) and sensitive

to boiling.

We reasoned intestinal mucins might fit this profile due to their highly associative and

sticky properties (captured on a filter) and as proteins they would be sensitive to heat.

Mucins are extensively glycosylated proteins found throughout the gastrointestinal

system, which form a layer between the intestinal epithelial cells (IECs) and the

commensal or pathogenic microbiota (Johansson and Hansson, 2016). Also, they can

function as receptors for microbes (McGuckin *et al.*, 2011). To test whether mucins

were inhibiting bacterial killing by phage, we devised another method whereby CM was

separated via high speed centrifugation into soluble (S CM) and insoluble (INS CM)

forms (Figure 2Bi). We hypothesized that INS CM would contain mucin, since large

intestinal mucins are normally present in this portion, and be inhibitory to phage killing,

whereas the S CM would not have these properties (Carlstedt *et al.*, 1993). Indeed,

phage killing in unprocessed CM or INS CM was inhibited to a greater extent than in S

CM (Figure 2Bii). To more directly test the hypothesis that mucin was the inhibitory

factor, the mucolytic drug N-acetyl cysteine (NAC) was added to INS CM and porcine

gastric mucin (1.5 % m/v) added to S CM (Figure 2Bi-ii). Indeed, a 5-log reduction in

phage killing of ExPEC was observed in INS CM upon addition of NAC, which also

improved killing to that seen in S CM ($p < 0.0001$). Perhaps more compelling, the

addition of mucin to S CM abrogated bacterial killing by phage to levels originally

observed in CM alone (not significant $p = 0.970$). These results indicate that the inhibitor

of phage killing in cecal medium is intestinal mucin.

Figure 2

Figure 2: The inhibitory component is mucin. (Ai) CM was heat-treated (HT, 100°C) or filter treated (FT) prior to assay. (Aii) ExPEC levels in HT CM and FT CM. (Bi) CM was separated into pellet (insoluble), homogenized and treated with N-acetyl cysteine (NAC), or into supernatant (soluble) and mucin added. Following assay modified CM was plated for (Bii) ExPEC levels. (Ci) A screen of phage killing in mucin was performed using the phage in Table 1. (Cii) ExPEC levels in LB + 1.5% mucin. (Ciii) LB alone. N=3-6, **** p < 0.0001, " = none detected. Exp. performed 1X (B), 2X (A and C). Squares indicate individual biological replicates from independent cultures. Mean, ±SD shown.

Along these lines, *E. coli* is known to use mucins as a source of carbon (Chang *et al.*,

2004). We reasoned that a murine host colonized with ExPEC may see a bloom upon

NAC treatment due to the drug liberating the mucins for bacterial consumption and thus

serve as a system to test if the reduction in aggregated mucin would promote phage

HP3's ability to kill ExPEC. Indeed, upon treatment with NAC for two weeks, ExPEC

levels in the small intestine were increased and phage HP3 reduced ExPEC levels, 2.5

logs, in the small intestine (Figure S2Bi-ii). A similar trend was observed in the large

intestine, but it was not substantial (<1 log reduction), though in both organ sections the

data were not significant (Figure S2Biii). The less pronounced effect in the large

intestine may be due to the thickness of mucus, and thus the lower likelihood for NAC to

be effective at breaking up this mucus. Also, NAC's known to be rapidly absorbed in the

small intestinal tissue, thereby losing its effect in the more distal large intestine (Aruoma

et al., 1989; Tsikas *et al.*, 1998).

Discovery of a mucin-enhanced phage

An estimated 10^{31} phages exist in nature. And the human gastrointestinal tract contains

a large diversity consisting of >10^{10} phages many of which are just beginning to be

discovered (Shkoporov and Hill, 2019; Sausset *et al.*, 2020). Reasoning that human

sewage or the feces of animals may contain phage that have evolved to target their host

in high mucin environments, such as the intestinal tract, we screened our phage library

(Gibson *et al.*, 2019) and other phages (Table 1) recently isolated from these

environments for enhanced activity in LB containing 1.5% mucin (Figure 2C). This is the

same concentration that prevented phage HP3 activity in soluble cecal medium and was

shown to provide strong inhibition in LB for up to 8 hours (Figure S3A). Surprisingly,

only a single phage, designated phage ES17, significantly killed bacteria in the LB

mucin medium (Figure 2Cii, $p < 0.0001$, about 1-log improvement). Surprisingly, phage

ES17 was active in mucin despite being much less effective (> 3-logs) than phages

HP3, J2W, Ult1, Shp1, or M1S, which completely killed ExPEC to undetectable levels in

LB medium alone (Figure 2Ciii, LB). Consistent with these data, when the amount of

mucin was varied from 0 to 2% and phages HP3 and ES17 were compared for lytic

activity against ExPEC, phage HP3 was highly effective as the concentration of mucin

was lowered to below 0.5%, but completely inhibited above that level (Figure S3B).

However, phage ES17 was most effective at concentrations in which HP3 was inactive

(0.5 – 1% mucin) and least active at concentrations below or above this range. Taken

together, these data suggests that phage ES17 harbors unique properties that facilitate

its ability to efficiently be lytic in the presence of mucin.

Phage ES17 is a rare C3 type phage whose activity is enhanced by mucin.

We sought to understand the molecular mechanism of phage ES17's enhanced ability

to find and lyse its bacterial host in mucin. Phage ES17 was determined to be a dsDNA

virus of the order *Caudovirales*, family *Podoviridae* and genus *Kuravirus*. The genus

Kuravirus was named for the River Kura in Tbilisi, Georgia where the original phage of

this genus, PhiEco32, which shows close genetic similarity to ES17, was first isolated

(Figure S4A) (Savalia *et al.*, 2008). Kuravirus phages have similar morphological

characteristics (Ren *et al.*, 2019), including an elongated C3 type capsid, which is

considered a rare morphotype for phages (only 13% of tailed phages have elongated

heads) and short tail fibers (Ren *et al.*, 2019). Electron microscopy confirmed ES17 to

have these same morphological characteristics (capsid >100 nm – Figure 3A). Similar to

other phages of this genus, ES17 has a small genome size consisting of 75,007 bps

with 123 predicted ORFs (Figure 3B).

In order to determine why this phage is distinct from other phages that lack activity in

mucin-rich environments, we examined the ability of phages ES17 and HP3 to adsorb to

their *E. coli* hosts. Previous data had shown that 98% of HP3 was adsorbed in 10

minutes, whereas only 32% of ES17 was adsorbed in that time (Gibson *et al.*, 2019).

We wondered whether the addition of mucin could improve phage ES17 adsorption and

inhibit the adsorption of phage HP3. A modified adsorption assay was utilized for this

experiment using ExPEC "coated" with mucin. Briefly, the bacteria were incubated in

different concentrations of mucin (0%-1.5%), pelleted, and washed to remove any

mucin that did not adhere to the bacterial surface (Figure 3Ci). Next, a standard

adsorption assay was conducted with phages ES17 and HP3. Interestingly, phage

ES17 showed no adsorption to ExPEC in 10 minutes in the absence of mucin; however,

if first incubated with 1.5% mucin, adsorption increased to 98% (Figure 3Cii). In

contrast, 98% of phage HP3 was adsorbed without mucin and the adsorption dropped

to 75% in its presence (Figure 3Ciii). Transmission electron microscopy (TEM) analysis

of these samples at 10 minutes demonstrated that only in the presence of mucin was

phage ES17 found bound to the bacterial surface (red arrows) (Fig 3Di-vi, 4 out of 11

images of ExPEC + mucin had visible evidence of ES17 compared to 0 out of 10 in the

absence of mucin). TEM imaging of phage HP3 adsorption showed the presence of

bound phage in images with and without mucin present (red arrows) (Figure S4Bi-ii). In

this case, 5/14 of the imaged fields showed phage HP3 bound with mucin-coated

ExPEC compared with 7/14 bound without mucin (Figure S4Biii). Finally, we adapted an

ELISA-like approach to determine if phage ES17 preferred binding to surfaces coated

with mucin, a hypothesis consistent with our data. Indeed, when mucin was bound to an

ELISA plate, phage ES17 bound to the mucin-surface at higher levels than phage HP3

(Figure 3E, $p = 0.041$ and $p = 0.002$). Taken together, using three different approaches,

these data obtained suggests that phage ES17 binds mucin, a property that may

enhance its ability to infect *E. coli* in mucin-rich environments.

Figure 3

A.

B. Phage ES17 genome

C. i. Mucin-coating

ii. ΦES17

iii. ΦHP3

D. ΦES17   No Mucin   Mucin-coated

vi.

E.

Figure 3: Phage ES17 is a rare C3 type phage with avidity for mucin. (A) TEM of ES17. (B) Genome of ES17. (Ci) ExPEC was incubated in 0%, 0.5% and 1.5% mucin, centrifuged, washed and phage added for an adsorption assay. (Cii) Phages ES17 and (Ciii) HP3 adsorption. Adsorption constant (K) and % adsorbed in 10 min indicated. Regression line generated from mean values of phage remaining for time points indicated. (Di-v) TEM of phage ES17 and ExPEC fixed after 10 min. adsorption. White bar is 500 nm. Red arrows indicate phage (Dvi) Attached phages per field view. (E) Mucin-ELISA results for phages HP3 and ES17. Squares indicate individual biological replicates from independent cultures. Exp. performed 1X (C) and (D), 2X (E). N=6, " = none detected. Mean, ±SD shown.

Phage ES17 may bind bacterial polysaccharides

As indicated above, phage ES17 is a close relative of the kuravirus phiEco32 and

annotated as the same phage type. ES17's putative tail fiber protein (ES17-TFP)

showed high homology (64% identical) to a tail fiber protein in another lytic podophage,

the T7-like bacteriophage LM33_P1, which also targets ST131 strains (Dufour *et al.*,

2016). This homology is in the C-terminal receptor binding region (Figure S5Ai). The N-

terminal region is similar to vB_EcoP_WFI101126 (47% identical), a phage with C3 like

morphology similar to phage ES17 (Figure S5Aii)(Korf *et al.*, 2019). The T7-like phage

tail fibers have recently been characterized and shown to have knob-like tips, which we

also observed in the EM images of phage ES17 (Figure 3A) (Garcia-Doval and van

Raaij, 2012). T7-like phage tail fibers have been shown to possess endosialidases that

target surface sugars, such as capsule-forming polysaccharides (Garcia-Doval and van

Raaij, 2012). A BLAST analysis revealed that ES17-TFP contains a putative

pectinesterase (E value 7.45e-03, 369 bp) with high homology to phage lyases (Figure

4A). Modeling of the predicted structure of ES17-TFP indicated it may contain a pectin

lyase fold (Figure S5B, IPR011050). These lyases cleave bacterial polysaccharides and

are found in biofilm-degrading phages (Latka *et al.*, 2017). Additionally, this enzyme

was not found in phage HP3. Thus, we formed the hypothesis that ES17 might degrade

and infect biofilms.

ExPEC is a primary cause of catheter-associated UTI (CAUTI), often associated with

the formation of biofilms on the catheter (Russo and Johnson, 2003). For this study, we

used a clinical isolate (DS515) from a CAUTI patient, which forms significant biofilms *in*

vitro (data not shown). We tested the effect of various *E. coli* phages on DS515 biofilms

using a colorimetric assay with a metabolic substrate, MTT, to determine the sensitivity

of the biofilm bacteria to the phage (Figure 4B). As determined in Figure 4C, of the 10

E. coli phages assessed, which were shown to be lytic towards the clinical isolate, only

two phages significantly educed *E. coli* viability after establishment of the biofilm, and

only one of these three, ES17, reduced the biofilm by more than half. Since phage HP3

showed moderate levels of killing of *E. coli* in biofilms, we compared its ability to phage

ES17 over a 6-log titration range. Strikingly, as little as 10^3 PFU/ml of phage ES17

statistically reduced biofilm viability, which was greater than that achieved with 10^9

PFU/ml of phage HP3 (Figures 4Di-ii). In addition, phage ES17 showed a dose

dependent response at each concentration with the greatest effect seen at 10^9 PFU/ml

(p < 0.0001) (Figure 4Dii). Also, we found that phage ES17 formed large plaques with

halos on DS515 and related clinical isolates (picture above Figure 4Dii). This is in

contrast to phage HP3 which showed no halo formation (picture above Figure 4Di). This

halo formation is indicative of EPS (extracellular polymetric substance)-degrading

activity (Knecht, Veljkovic and Fieseler, 2020).

We reasoned ES17-TFP may be responsible for the cleavage of the EPS in the biofilm,

thereby allowing phage ES17 to gain access to *E. coli* in the biofilm. Thus, we cloned

the DNA encoding ES17-TFP into an expression vector and purified it from *E. coli*. A

highly induced band of the expected molecular weight of ES17-TFP (107 KDa) was

observed upon SDS-PAGE separation of the eluted fractions (Figure S5C). This band

was not observed in the vector only control elutions (data not shown). When ES17-TFP

was added to the biofilms, no change in bacterial viability was observed, nor did it

enhance ES17 killing of *E. coli* in biofilms when added separately (Figure 4E). Likewise,

when incubated in CM, or examined for its ability to degrade mucus using a gel-shift

assay in an anti-mucin Western blot, it did not enhance ES17 activity or degrade mucus

(data not shown). Furthermore, phage ES17 by itself also did not demonstrate this

activity. These data suggests that phage ES17's unique killing abilities are not

dependent on degradation of polysaccharides (e.g. mucin or bacterial). This leaves

open the question of how this tail fiber protein may enhance killing in carbohydrate-rich

environments.

Figure 4: Phage ES17 may bind bacterial polysaccharides. (A) ES17 TFP domain compared
to vB_EcoP_WFI101126 lyase. (B) DS515 was grown o/n in TSB and inoculated on a plate for
48-hr biofilm growth. A phage library was screened with MTT used to quantitate biofilm forming
bacteria. (C) DS515 levels with phage library (N=3-8). (Di) DS515 levels with HP3 and (Dii)
ES17 (N=12). Plaques "P" from phages on pictured with or without halo "H" (E) ExPEC levels
with purified ES17-TFP in CM assay (N=2-8). Squares indicate individual biological replicates
from independent cultures. Exp. performed 2X. Mean, ±SD shown. Figure created with
BioRender.com.

Phage ES17 binds human heparan sulfated proteoglycans.

Phage ES17 harbors an enhanced ability relative to other *E. coli* phages to find its

bacterial host in environments in which carbohydrates are a prominent chemical

component (examples from above include cecal medium, mucin-rich broth, and

biofilms). A structural analysis of modeled ES17-TFP showed a high similarity to a

phage K5 lyase binding domain (E-value=4E-12). The predicted structure of ES17-TFP

(blue) and K5 lyase are pictured in Figure S6Ai (PDB:2X3H; red) with identical residues

colored yellow. Phage K5 binds K5 capsular polysaccharide and acts as a K5

polysaccharide lyase (Hanfling *et al.*, 1996). The K5 *E. coli* capsule is made of a

repeating disaccharide that is identical to the precursor of heparin and heparan sulfate

(HS), a linear polysaccharide, present in glycosaminoglycans (HSPGs, heparan sulfate

proteoglycans). (Figure S6Aii) These proteoglycans are found on mammalian cells and

in mucus (Monzon, Casalino-Matsuda and Forteza, 2006). Also, mucins with similar

structures to heparan sulfate/heparin (α-linked GlcNAC or N-Acetyl-D-glucosamine) are

present intestinally and found in PGM (Fujita *et al.*, 2011).

We reasoned that ES17's enhanced activity might be due to a novel ability to bind

mammalian polysaccharides found on glycoproteins. This would be a novel mechanism

by which a phage could localize to its host, thereby explaining its enhanced activity in a

mucin-rich environment and in EPS biofilms. To test this idea, we assessed the ability of

purified ES17-TFP to bind to a glycan array containing over ~ 860 unique glycan

structures from porcine gastric mucin (PGM), glycosaminoglycans (GAGs) and a variety

of synthetic and naturally sourced glycans generated by the Consortium for Functional

Glycomics (CFG) (see supplementary methods). No or very low relative fluorescent

units (RFU), a proxy for binding, was observed for a wide array of mammalian glycans,

including those purified from porcine gastric mucin (Figure S6Bi-ii). However,

surprisingly, there was an increase of several orders of magnitude in RFU (RFUs >

2000) observed for binding to the GAGs containing heparan sulfate (ID # 64-173), but

not the structurally similar GAGs hyaluronic acid #1-20 or chondroitin sulfate #21-63

(Figure S6Bii-iii). The finding that purified ES17-TFP binds human heparan sulfated

proteoglycans provides a possible mechanism to explain why phage ES17

demonstrates enhanced activity in intestinal environments.

ES17 binds to the surface of human intestinal enteroids (HEIMs)

Human intestinal enteroids (HIEs) are organotypic, higher-order cultures that have

become popular as surrogates to model the human intestine. They can be grown as 3-

dimensional structures complete with a lumen and crypt/villus axis or as 2-dimensional

monolayers that facilitate host-pathogen interactions (In *et al.*, 2016; Saxena *et al.*,

2016). These cultures are also useful because they express a variety of glycans found

in the human intestine, including mucins and proteoglycans (Saxena *et al.*, 2016).

Human intestinal enteroid monolayers (HIEMs) were derived from colonic stem cells

following differentiation for 5 days in high Wnt medium. Phage ES17 or HP3 were

added to confluent HIEMs for 1 hour, extensively washed, and visualized by

immunofluorescence microscopy using antibodies raised against each phage. Little to

no detectable phage HP3 was observed on the HIEMs intestinal epithelial cell (IEC)

surface, though antibodies generated robust signal and specificity towards the phage

when HP3 was fixed on slides alone (Figure 5A and S4C). Phage ES17 (green) bound

evenly to the IECs on the apical side, including areas where prominent Muc2 (red)

localization was observed, but also on areas where there was no muc staining (Figure

5Aii). In order to determine whether phage ES17 bound to the IECs of HIEMs via

heparan sulfate, we pre-treated the HIEMs with heparinase III to enzymatically remove

HSPGs and then assessed phage binding. Indeed, HIEMs treated with heparinase

significantly reduced the levels of bound ES17, both qualitatively and quantitatively

(Figures 5Bi-vii, $p < 0.0001$). Next, we tested whether this binding to the epithelial cell

surface also improved bacterial killing. Previous data had shown that ExPEC adheres

less to intestinal enteroids compared to the diarrhea-causing pathogen

Enteroaggregative *Escherichia coli* (EAEC) (data not shown). EAEC adheres to HIEMs

robustly in an aggregative, mesh-like pattern and similar to phage ES17 utilizes heparan

sulfate as a receptor to bind these human colonic cells (Rajan et al., 2020, paper *in*

review)(Poole, Rajan and Maresso, 2018; Rajan *et al.*, 2018). Thus, EAEC serves as an

ideal *E. coli* strain to test the hypothesis that phage targeting HSPGs can increase

killing. For this experiment we pre-coated HIEMs with phage ES17, washed, as detailed

above, then infected with EAEC strain 042. HIEMs were fixed and stained using a

Geimsa-Wright stain to visualize cells and bacteria, as previously described (Rajan *et*

al., 2018). Infected HIEMs showed robust bacterial adhesion to cells with an

aggregative phenotype (Figure 5Cii, red arrows). Phage coated HEIMs, however,

showed significantly reduced EAEC on the surface of the organoid (Figure 5Ciii-iv, **$p =$

0.007). Taken together, these data suggests phage ES17 may localize to both the

mucus layer and to the IEC surface by binding to HSPGs, which likely position the

phage to be in the exact location needed to find its bacterial target in the intestinal

microenvironment.

Figure 5

Figure 5: ES17 binds to the surface of human intestinal enteroids (HIEMs). HIEMs were incubated with phage for 1 hr, washed, then fixed and imaged for phage (Alexa Fluor 488, Green), intestinal cells (DAPI, Blue), and muc2 (Alexa Fluor 594; Red). (Ai) No phage added. (Aii) Phage ES17 or (Aiii) HP3 added. Monolayers pretreated with heparanase III then with phage. (Bi, ii) No phage. (Biii) Phage ES17 and (Biv) phage HP3 added. (Bv) Hep III treated HEIMs incubated with ES17 and (Bvi) HP3 (Bvii) Quantification of particles (phage) per well. (N=1-2). Images at 60X. Scale bars at 20 μm. HIEMs (Ci) uninfected or (Cii) Infected with EAEC 042 (Ciii) or pretreated with phage ES17 prior to EAEC. Images at 60X. Scale bars at 50 μm. (Civ) EAEC attached to HIEMs per field view. Mean, ±SD shown. Squares indicate individual biological replicates from independent cultures. Exp. performed 1X. Figure created with BioRender.com.

Phage ES17 kills ExPEC in the mammalian intestine

The finding that phage ES17 demonstrated enhanced lytic activity in the presence of

mucins, was the best of several screened phages in a mock luminal environment rich in

mucins, and binds the human organotypic culture IECs via HSPGs, which leads to

EAEC-killing on the bacterial surface, prompted an examination into whether this phage

could overcome the intestinal-induced inhibition of phage lytic activity towards colonized

ExPEC that was observed for phage HP3. We first tested if ES17 was effective in cecal

medium. Indeed, phage ES17 showed a 2.5-log improvement in ExPEC removal in this

environment compared to phage HP3 (Figure 6A, $p < 0.0001$). There was no effect on

the number of OTUs or the Shannon diversity index in this experiment, suggesting

phage ES17 was highly selective at removing only the target ExPEC strain (Figure 6Bi-

ii). We next tested the effect of phage ES17 on ExPEC in a murine intestine. Animals

were colonized with ExPEC as in Figure 1A and treated with either phage ES17 or HP3

(Figure 6C).The dose of phage was also increased from 10^9 PFU to 10^{10} to also

evaluate if giving more phage would improve HP3's ability to reduce ExPEC, especially

in more proximal segments as the phage slowly moves through the alimentary canal.

Examination of the small and large intestine on day 6 showed phage levels were high

across all groups (10^6-10^8 PFU/g intestinal tissue) (Figure 6D). Animals treated with

phage HP3 had no detectable CFU in small intestinal tissue (Figure 6E), but had

indistinguishable levels from that of the untreated control in the cecum. In contrast,

every animal treated with phage ES17, except one, had no detectable levels of ExPEC

in either the small or large intestine, suggesting that this lytic phage possess a unique

ability to target ExPEC in complex mucosal environments such as the large intestine.

Figure 6

Figure 6: ES17 kills in the mammalian intestine. (A) ExPEC levels in CM assay with ES17
(N=3). 16S rRNA analysis of CM including (Bi) OTU and (Bii) Shannon diversity index. (C) Mice
were gavaged with ExPEC then treated with no phage or gavaged daily doses of phage HP3 at
10^{10} PFU, or ΦES17 at 10^{10} PFU or both phages (2×10^{10} PFU total). (D) Intestinal (tissue) phage
levels (E) Intestinal (tissue) ExPEC levels. (N=10). Symbols (squares) indicate individual mice
or biological replicates derived from different cultures. Exp. performed 1X. NS not significant. " =
none detected. Mean, $\pm$SD shown.

Discussion

A limitation of all antibiotics is their broad killing activity and no inherent features to

function well in complex human environments, including human blood, urine, or at

mucosal surfaces throughout the body. In particular, the gastrointestinal tract (GIT) is a

highly complex system made up of diverse organs and tissues and within that a

diversity of cell types. Some of the cells and factors found at the mucosal surface may

influence intestinal capacity including: intestinal epithelial cells, enteroendocrine cells,

stem cells, mucus-secreting goblet cells, antimicrobial peptides (AMPs – from Paneth

cells) and secretory immunoglobulin A (Mammarappallil and Elsinghorst, 2000;

Johansson, Larsson and Hansson, 2011; Sato *et al.*, 2011; Gribble and Reimann, 2016;

Pickard *et al.*, 2017). To add to this complexity, the GIT is colonized by a large number

of microbes—archaea, fungi, protists and bacteria that have co-evolved with the host,

termed the human gut microbiota (Pickard *et al.*, 2017; Heintz-Buschart and Wilmes,

2018). A bacteriophage would have to navigate through this complex ecosystem to find

and infect its host. Since there are estimated to be >10^{10} PFU/g of phages already

present in the GIT, this suggests that phages can thrive in this environment (Hoyles *et*

al., 2014; Shkoporov *et al.*, 2018). Here, we report a novel anti-bacterial targeting

strategy that seems to have evolved to position a lytic bacteriophage in the exact niche

as its bacterial host. Our results indicate that; (i) phage HP3 that is lytic towards *E. coli*

ST131 *in vitro* and is very effective at eliminating bacteremia in a murine model of

sepsis, is ineffective when tested for this same property in the murine intestinal tract; (ii)

that the inhibition is due to intestinal mucins; (iii) a medium designed to mimic the

luminal environment can be used to identify phage with enhanced lytic activity in such

an environment; (iv) a rare C3 type phage, isolated from human wastewater, overcomes

this inhibitory activity due to an enhanced ability of the phage to bind to heparan-

sulfated proteoglycans present in mucus or immobilized on the surface of intestinal

epithelial cells, which likely drives the positional targeting of the phage to the exact

ecological niche as the host bacterium. In addition, our data reveal that this treatment

when compared to antibiotic treatment, did not alter the intestinal diversity of the

microbiota. Taken together these data suggest a new mechanism of predation by a

phage, one which may be highly useful for the killing of bacteria in intestinal

environments and biofilms, both of which would be desirable properties of a targeted,

specific antibacterial.

Potential narrow or limited ecological range of bacteriophages has not been as explored

as much as their narrow host range capabilities. Using an experimental process of

elimination, we determined that mucin can greatly inhibit phage infection. Mucins are

composed of tandem repeats of serine and threonine that act as an attachment sites for

o-linked glycans (N-acetlygalactosamine, N-acetylglucosamine, fucose, galactose and

sialic acid) (McGuckin *et al.*, 2011; Wang and Hasnain, 2017). Gel-forming mucins,

which make up the mucus intestinal layer are large polymers (up to 40 MDa) of mucins

attached via disulfide linkages (McGuckin *et al.*, 2011). When we incubated phages with

porcine gastric mucin, composed of the gel-forming mucin MUC 5AC (Sturmer *et al.*,

2018), we saw reduced or no bacterial killing. However, when we added N-acetyl

cysteine (NAC), which is known to exert a mucolytic effect by reducing the disulfide

bonds that keep these mucin polymers together, bacterial killing was restored (Aruoma

et al., 1989; Sadowska *et al.*, 2006). NAC is a drug has been tested and used in the

clinical setting to treat syndromes including cystic fibrosis (CF) and chronic obstructive

pulmonary disease (COPD). In both of these cases it loosens thick mucus in lungs, but

it can also serve as a treatment for acetaminophen overdose due to its antioxidant

action (Sadowska *et al.*, 2006; Saito, Zwingmann and Jaeschke, 2010; Bear, 2013).

Perhaps this drug could also be used as an adjuvant to phages to help treat patients

with bacterial infections in mucin-rich ecosystems such as the gut or the lungs of CF

patients. We tested oral NAC treatment in mice and found it to be more effective in

improving phage-mediated bacterial killing in the small intestine, but not in the large

intestine. NAC has been shown to be rapidly absorbed following oral dosing after only

60 minutes (Tsikas *et al.*, 1998). An oral dose of NAC has a short half-life of 2.5 hours

and 10% bioavailability in the gut (Tsikas *et al.*, 1998). It is likely that NAC did not exert

a mucolytic effect on the distal large intestine. Interestingly, we observed elevated

ExPEC levels in intestinal tissue with NAC treatment. It is possible that some pathogens

like ExPEC may thrive in environments with a reduced mucus layer.

Another possibility, distinct from adjuvating phage with NAC, would be to search for

phages that have evolved phenotypes that enhance their ability to encounter their

bacterial host in a specific ecosystem, such as the GIT. Phage ES17, originally isolated

from human sewage, is a member of the family *Podoviridae* and has a rare C3 type

elongated capsid morphology with paddle-like short tail fibers (Ackermann, 2001).

Previously, this phage was shown to be similar to PhiEco32, a rare Kuravirus which are

known for their prolate head (Gibson *et al.*, 2019). Phage ES17 showed a significant

killing effect in cecal medium (CM) and in a mouse model of intestinal colonization

compared to phage HP3. Phage ES17 bound mucin significantly better and coating

bacteria with mucin increased phage adsorption. There is precedence for this concept.

The bacteriophage adhering to mucus (BAM) model proposes that phages bind to

mucus via *hoc* proteins or Ig (immunoglobulin fold-like) domains present on the capsid

proteins of phages similar to T4 (Fraser *et al.*, 2006; Barr *et al.*, 2013, 2015). This

mechanism was suggested to explain how phages bind to the metazoan mucosal

surface and increase the probability of encountering a bacterial host, likely via a unique

type of controlled diffusion that increases spurious interactions with its bacterial target

(Barr *et al.*, 2013, 2015). These studies showed that T4 adsorption increased in the

presence of mucin from 65% to 80% absorbed phage within 10 minutes (Barr *et al.*,

2015). The adsorption changes in phage ES17 in the presence of mucin were all or

nothing from 0% to 98% within 10 minutes. This suggests that mucin may be a type of

"bridge receptor" for phage ES17 since this phage is ineffective at killing in the absence

of mucin, even at very high levels of phage. Such a model would make sense in the

context of *E. coli* embedded in a mucin matrix, perhaps breaking down mucin as a

source of carbon and thus binding it, thereby allowing the phage to adapt a capsular

polysaccharide binding mechanism to structurally related sugars that are prominent on

proteoglycans.

Bacteriophages have been shown to have a variety of carbohydrate-binding proteins

present on tail fibers in order to bind different sugars present on the bacterial cell

surface (Nobrega *et al.*, 2018). Some of the most diverse are capsular depolymerases,

which bind and break down capsular carbohydrates secreted by bacteria (Pires *et al.*,

2016). These capsular depolymerases are diverse because of the diversity of bacterial

capsule types. Capsule types have been shown to mimic some components of the

intestinal system, including sugars present in the mucus layer (Vimr and Lichtensteiger,

2002; Severi, Hood and Thomas, 2007). Bacteria can use these mechanisms to subvert

the immune system and to allow for the colonization of hosts (Vimr and Lichtensteiger,

2002; Kahya, Andrew and Yesilkaya, 2017). Phage ES17 was effective at killing

bacteria in biofilms and possessed a protein with putative capsular depolymerase

domains in a tail fiber protein. However, we could not detect any polysaccharide lyase

activity and instead detected strong and stable affinity for a human intestinal sugar, the

glycosaminoglycan sugar, heparan sulfate. This suggests a model whereby phage

ES17 adapted to "colonize" the mucosal environment by binding to mammalian sugars.

Heparan sulfate proteoglycans (HSPG) are expressed on the surface of many types of

cells and prominently on the surface of intestinal epithelial cells (Garcia *et al.*, 2016).

One of the more tantalizing discoveries here was that human enteroids derived from

colonic stem cells specifically bound phage ES17, but not other *E. coli* phages. HSPGs

consist of repeating units of sulfated polysaccharides—heparan sulfate (HS). HS is a

linear polysaccharide that begins as the precursor, heparan (disaccharide of α1,4-linked

N-acetylglucosamine (GlcNAc) and β1,4-linked glucuronic acid (GlcUA)) (Sugahara and

Kitagawa, 2002; Murphy *et al.*, 2004). Subsequent modifications due to sulfation and

epimerization leads to the mature form polysaccharide HS (Murphy *et al.*, 2004).

Bacteriophage K5 utilizes a tail spike lyase (*KflA*) to bind and degrade K5 capsule

present on *E. coli* strains (Hanfling *et al.*, 1996). Because the K5 capsule (heparoson;

β1,4-linked GlcUA and α1,4-linked GlcNAc) polysaccharide has been shown to be

structurally identical to heparan sulfate precursor, heparan, this enzyme can also act a

heparinase (Hanfling *et al.*, 1996; Roberts, 1996; Murphy *et al.*, 2004; O'Leary, Xu and

Liu, 2013). K5 heparan lyase cleaves the linkage of N-acetyl glucosamine and

glucuronic acid, but is inhibited by sulfated regions (Murphy *et al.*, 2004; O'Leary, Xu

and Liu, 2013). Further homology searches showed that the carbohydrate binding

domain found in ES17-TFP is structurally homologous to the binding domain present in

the tail spike in phage K5. Considering that heparanase III cleaves at the same regions

where K5 lyase would bind and degrades suggests ES17-TFP is utilizing the same

receptor. Indeed, the addition of heparanse III, from the soil bacteria *Flavobacterium*

heparinum, which specifically cleaves unmodified (unsulfated) NAc domains (GlcNAc-

GlcUA) and NA/NS domains (GlcNS (N-Sulfo-D-Glucosamine) and GlcNAc) of HS,

leads to abrogated phage ES17 binding to cells, meaning ES17 was highly specific for

this type of glycan (Murphy *et al.*, 2004; Park *et al.*, 2017). Structurally similar mucins to

heparan sulfate (Class III type, α-linked GlcNAc) suggest that this protein may also

target these intestinal mucins (Ota *et al.*, 1998; Fujita *et al.*, 2011). This interaction is

driven by phage ES17's tail fiber protein, which is not related to the *hoc* proteins or

contain an Ig-like fold as predicted from the BAM model. Thus, it would seem that

phage ES17 uses a novel mechanism to localize not only a mucin-rich surface, but also

directly to the mammalian epithelial surface (Figure 7). It is worth speculating that not

only would this be a way to intercept *E. coli* that would also colonize this surface, but it

could also be exploited therapeutically as a prophylactic "epithelium protecting" coating

that prevents the pathobiont from invading the mucosa. Several pathogens, including

bacteria and human pathogenic viruses have been shown to bind to HS and use them

as receptors (Alvarez-Dominguez *et al.*, 1997; Fleckenstein, Holland and Hasty, 2002;

Cagno *et al.*, 2019). It is tempting to speculate that this property might have been also

acquired by phage, as suggested here.

Figure 7: Model showing (1) mucins from the intestinal mucus layer inhibit phage infection, (2)
phage ES17 can bind to mucin and utilize other intestinal glycans as a receptor to infect and kill
mucus-coated bacteria, and (3) phages like ES17 can be utilized to coat the intestinal
epithelium by binding heparan sulfate glycans to protect from invasive pathogen infection.
Figure created with BioRender.com.

Materials and Methods

Bacterial strains and phages

ExPEC ST131 isolate JJ1901 was used in all ExPEC infections, except Figure S1Aiii

(JJ2528). Both isolates were previously obtained from Dr. Jim Johnson (University of

Minnesota)(Green *et al.*, 2017b). Commensal *E. coli* ECN (Figure S1iv) was isolated

from a human fecal sample. Deidentified strain DS515 is a clinical isolate obtained from

the clinical microbiology laboratory at the Houston Veterans Administration Hospital.

Prior to infections all strains were grown overnight at 37°C from a single colony streaked

on an LB agar plate. The intestinal pathogen EAEC 042 (serotype 44:H18), was

originally isolated from a child in Peru, and extensively characterized (Nataro *et al.*,

1985).

Phages HP3, ES12, ES17, ES19, ES21 and ES26 have been previously described and

characterized (Green *et al.*, 2017a; Gibson *et al.*, 2019). Phage 6914, 6915 and 6939

were recently isolated from sewage. All phages described in Table 1 were isolated by

single plaque isolation from environmental sources as described previously (Green *et*

al., 2017b; Gibson *et al.*, 2019).

Murine infections

Mixed ages (6-10 months) and sexes of BALB/c (Jackson laboratories, Bar Harbor, ME)

mice were used in mouse models of infection. Mice were kept in SPF environment at

Baylor College of Medicine CCM (Center for Comparative Medicine) Taub facility. All

methods performed on mice were approved in accordance with relevant guidelines and

regulations from "The Guide and Care and Use of Laboratory Animals" (National

Institute of Health) and approved by Baylor College of Medicine's Institutional Animal

Care and Use Committee (protocol AN-6372). For infections mice were kept in a

biohazard facility with sterile food and water. The mice were individually housed during

colonization experiments and bedding replaced with autoclaved techboard liners for

daily fecal collection. For colonization experiments, sample size was determined based

on previous colonization experiments in mice (Green *et al.*, 2017a). Mice received a 10^9

CFU dose of ExPEC strain JJ1901 via oral gavage. Rodent health was monitored daily

for indication of pain or disease. Bacterial colonization (fecal and intestinal) was

determined after homogenization and selective plating for the chloramphenicol resistant

strain JJ1901 on LB agar plates containing chloramphenicol and colony counting.

Purified phage in 3% (m/v) $NaHCO_3$ was administered either via gavage or in water 5%

(m/v) sucrose added *ad libitum*. All groups received sucrose and $NaHCO_3$ in water for

consistency. The antibiotic ampicillin (1g/500 ml) was administered in water. Phage

colonization was quantified after dilution of homogenates and serial plating on a double

agar overlay assay.

Ex vivo cecal model

A modified cecal assay was used for experiments (Theriot *et al.*, 2014). Briefly, cecal

contents from just euthanized mice were pooled and homogenized in sterile 0.09%

NaCl solution at a 1:5 dilution (mg:mls). The homogenate was centrifuged to remove

large particulates (2000 G for 30 sec.). The supernatant fluid was used for 4.5 hr phage

killing assays at an MOI of 10 at 37°C, shaking (255 RPM), as previously described (Ma

et al., 2018). All cecal and mucin experiments were performed using independent

bacterial cultures grown up from different colonies streaked on a plate. This was

considered a biological replicate. For FS CM, cecal supernatant was centrifuged (6000

G for 5 min.) and filtered through a 0.22 μm syringe filter. For HT CM, the supernatant

was heated at 100°F for 20 min in a hot water bath, then cooled to RT for infections.

Insoluble CM and soluble CM (supernatant) were isolated post high-speed

centrifugation (9000 G for 5 min) of CM. The insoluble pellet was resuspended in sterile

0.09% NaCl solution for infections (IN CM). For the mucin assays, porcine gastric mucin

type II (PGM; Sigma-Aldrich) was used at various concentrations diluted in phosphate

buffered saline (PBS). The mucolytic drug N-acetyl cysteine (NAC, Sigma-Aldrich, 5

mg/ml) was diluted in PBS for de-mucolytic assays.

Phage Sequencing and Annotation

The annotation figure was generated from previous data (Gibson *et al.*, 2019) using

Geneious Prime 2019.2.3 and adjusted in Inkscape.

Mucin-coating Adsorption and Imaging

For the adsorption curves, assays were performed at an MOI of 1 using mid-log phase

cultures independently grown from different colonies on a plate (this was considered a

biological replicate) and samples taken every 5 min for 10 min. Prior to adsorption, PGM

(0%, 0.5% and 1.5% m/v) was added to the bacterial cultures for 10 min. RT, shaking

(255 RPM). The cultures were centrifuged (6000 G for 5 min) and gently washed with

PBS. The adsorption rate constants (K) were determined from the natural log of the

slope of the adsorption curve versus the bacterial concentration. Some cultures were

fixed in glutaraldehyde solution after 10 min adsorption for TEM imaging. TEM imaging

was performed at the Texas Children's Hospital Center for Digestive Diseases. After

fixation the samples were dehydrated in ethanol and followed by evaporation of ethanol

and coating with gold for observation during TEM.

Mucin Binding ELISA Assay

Clear-walled Immulon 2 HB 96-well microtiter plate (Immunochemistry Technologies

#227) were used for the ELISA assays. PGM (200 μl of 1mg/ml) was added to a

microtiter plate and incubated at 4°C overnight. The next day the mucin was removed

and wells were washed twice with PBS. The phage was added to wells for 1 hr then

washed three times in PBST (PBS with 0.1% Tween-20). The wells were blocked with

BSA then incubated with antibodies for the phage overnight at 4°C. Following washing

steps, a HRP (horse radish peroxidase) conjugated antibody was added for 1 hr. To

assess phage binding, TMB solution was added until the wells turned light blue and

then a stop solution (2M H_2SO_4) was added. The absorbance was read at 450 nm. Each

well was considered a biological replicate for this experiment.

Biofilm Screening

Biofilm experiments were recently described but with some modifications (Mapes *et al.*,

2016). *E. coli* DS515 cultures were independently grown from different colonies on an

agar plate in tryptic soy broth (TSB) were diluted to an OD of 0.1 and seeded to

inoculate a 96 well plate for the biofilm formation assay. Each well was considered a

biological replicate. The plates were incubated statically at 37°C for 48 hrs. Following

incubation, planktonic cells were removed, and the wells were washed three times in

PBS. Phages diluted in TSB to 10^9 PFU/ml were added and incubated for 24 hrs at

37°C statically. The wells were washed three times in PBS then an MTT assay was

performed to determine the metabolic activity of the phage resistant biofilm-forming

bacteria. Protein quantification was determined using a Bradford assay (Biorad) using

BSA as a standard. Purified ES17-TFP was added at a concentration of 0.17 mg/ml to

biofilms with added phage.

HIEM Infection and Imaging

Human enteroid monolayers (HIEMs) were differentiated for 5 days (>90% confluent) as

described (Poole, Rajan and Maresso, 2018). For experiments each well containing

HIEMs was considered a biological replicate. HIEMs were incubated with phage at 10^8

PFU/ml in culture differentiation media for 1 hour at 37°C in the presence of 5% CO_2 in

a humidified incubator, then washed in PBS. The HIEMs were fixed in Clark's solution

for 10 min to preserve the mucus layer. The HIEMs were permeabilized and blocked

with 5% BSA in 0.1% Triton X-100 in PBS for 30 min at RT. Mucus was detected using

antibodies to MUC2 (1:200) (Abcam) and nuclei stained with 4′, 6′-diamidino-2-

phenylindole (DAPI) (300 nM) for 5 min at RT. Antibodies against phages HP3 and

ES17 were generated from whole virus (phage) injection into rabbits performed by

Pacific Immunology. A 13 week antibody production protocol consisted of 4

immunizations and anti-sera collection.

To selectively removed HSPG from GAG chains, enteroid cultures were pretreated with

heparinase III (Sigma, 2U/ml) for 2 hours, as described (Jiao *et al.*, 2007), followed by the

addition of phage. Images were captured using a Zeiss LSM 510 confocal microscope.

Represented images were adjusted equally for brightness and contrast using FIJI

software version 2.0.0. The images were adjusted equally for brightness and contrast.

Particle analysis was used to determine particles per well. The Cell counter program in

FIJI was used to quantitate bacteria (EAEC) per field of view.

EAEC infection and staining

For infection of HIEMs, the EAEC strain 042 was grown overnight and then sub-cultured

(1:100) for 2 hrs at 37°C. One microliter of the log-phase bacterial culture was added to

100 microliters of the cell culture media and added to chambered slides either with

phage-coated HIEMs (as described above) or untreated. Cultures were incubated for

3.5 hrs at 37°C in the presence of 5% CO_2 in a humidified incubator. Following growth,

bacteria were removed, by gently washing the cells in PBS then the remaining cells

were fixed (Hema 3 fixative) and stained using a Geimsa-Wright stain (Fisher scientific).

The infection and staining technique have been previously described (Rajan *et al.*,

2018).

Statistics

Statistical analysis was performed using PRISM 8 software. Microbiome 16S data was

analyzed using ATIMA (Agile Toolkit for Incisive Microbial Analyses) as described in

supplemental methods. For figures with log transformed data and groups >2

significance was determined using a one-way ANOVA analysis (Figures 1, 2, 3, 4, 5 and

6) or two way ANOVA when necessary (Figure 3E). For multiple comparison analysis,

the secondary test, Tukey, was used. For non-transformed data a normality test

(Shapiro-Wilk) was performed to determine normality before a one-way ANOVA

analysis was performed. Significance was determined to be a p value less than or equal

to 0.05. Unless otherwise stated all graphs show the mean and standard deviation.

Acknowledgements

This work is supported in part by grant from US Veterans Affairs (VA I01-RX002595),

Roderick D. MacDonald Research Fund at Baylor St. Luke's Medical Center, the Mike

Hogg Foundation, and Seed Funds from Baylor College of Medicine Seed Fund

We would like to thank Dr. Melinda Engevik for general expertise and protocol help

related to mucin work. We would also like to thank Hannah Johnson from the Integrated

Microscopy Core at Baylor College of Medicine for technical assistance related to

imaging.

Author Contributions

825   S.G., C.L., X.Y., W.S., A.R., H.C. and X.S. performed experiments and analyzed data.

826   J.C. performed bioinformatics studies. R.R., B.T., and H.K., contributed to design of

827   studies and edited the manuscript. S.G. designed studies and wrote the manuscript.

828   A.M. contributed to design and overall major goals of the study and edited the

829   manuscript.

References

Ackermann, H. W. (2001). Frequency of morphological phage descriptions in the year

2000. Arch. Virol. *5*, 843–857.

Alvarez-Dominguez, C. *et al.* (1997). Host cell heparan sulfate proteoglycans mediate

attachment and entry of Listeria monocytogenes, and the listerial surface protein ActA is

involved in heparan sulfate receptor recognition. Infect. Immun. *1*, 78–88.

Antão, E.M., Wieler, L. H. and Ewers, C. (2009). Adhesive threads of extraintestinal

pathogenic *Escherichia coli*. Gut. Pathog. *1*, 22.

Aruoma, O. I. *et al.* (1989). The antioxidant action of N-acetylcysteine: its reaction with

hydrogen peroxide, hydroxyl radical, superoxide, and hypochlorous acid. Free Radic.

Biol. Med. *6*, 593–597.

Banerjee, R. and Johnson, J. R. (2014). A new clone sweeps clean: The enigmatic

emergence of *Escherichia coli* sequence type 131. Antimicrob. Agents CH. *9*, 4997-

5004.

Barr, J. J. *et al.* (2013). Bacteriophage adhering to mucus provide a non-host-derived

immunity. Proc. Natl. Acad. Sci. *26*, 10771–10776.

Barr, J. J. *et al.* (2015). Subdiffusive motion of bacteriophage in mucosal surfaces

increases the frequency of bacterial encounters, Proc. Natl. Acad. Sci. *44*, 13675–

13680.

Bear, C. E. (2013). 50 years ago in the Journal of Pediatrics: the effect of N-

acetylcysteine on the viscosity of tracheobronchial secretions in cystic fibrosis of the

pancreas, J. Pediatr. *1*, 85.

Brum, J. R. *et al.* (2016). Illuminating structural proteins in viral "dark matter" with

metaproteomics. Proc. Natl. Acad. Sci. *9*, 2436-2441.

Cagno, V. *et al.* (2019). Heparan Sulfate Proteoglycans and Viral Attachment: True

Receptors or Adaptation Bias? Viruses. *7*. 596.

Carlstedt, I. *et al.* (1993). Characterization of two different glycosylated domains from

the insoluble mucin complex of rat small intestine. J. Biol. *25*, 18771-18781.

Chang, D. E. *et al.* (2004). Carbon nutrition of *Escherichia coli* in the mouse intestine.

Proc. Natl. Acad. Sci. *19*, 7427-7432.

Colpan, A. *et al.* (2013). *Escherichia coli* sequence type 131 (ST131) subclone h30 as

an emergent multidrug-resistant pathogen among US Veterans. Clin. Infect. Dis. *9*,

1256-1265.

Dedrick, R. M. *et al.* (2019). Engineered bacteriophages for treatment of a patient with a

disseminated drug-resistant Mycobacterium abscessus. Nat. Med. *5*, 730-733.

Dufour, N. *et al.* (2016). Bacteriophage LM33_P1, a fast-acting weapon against the

pandemic ST131-O25b: H4 *Escherichia coli* clonal complex. J. Antimicrob. *11*, 3072-

3080.

Fleckenstein, J. M., Holland, J. T. and Hasty, D. L. (2002). Interaction of an Outer

Membrane Protein of Enterotoxigenic *Escherichia coli* with Cell Surface Heparan

Sulfate Proteoglycans. Infect. Immun. *3*, 1530–1537.

Fraser, J. S. *et al.* (2006). Ig-like domains on bacteriophages: a tale of promiscuity and

deceit. J. Mol. Biol. *2*, 496–507.

Fujita, M. *et al.* (2011). Glycoside hydrolase family 89 α-N-acetylglucosaminidase from

clostridium perfringens specifically acts on GlcNAcα1,4Galβ1R at the non-reducing

terminus of O-glycans in fastric mucin. J .Biol. Chem. *8*, 6479-6489.

Garcia-Doval, C. and van Raaij, M. J. (2012). Structure of the receptor-binding carboxy-

terminal domain of bacteriophage T7 tail fibers. Proc. Natl. Acad. Sci .U S A. *24*, 9390–

9395.

Garcia, B. *et al.* (2016). Surface Proteoglycans as Mediators in Bacterial Pathogens

Infections. Front. Microbiol. *7*, 220.

Gibson, S. B. *et al.* (2019). Constructing and Characterizing Bacteriophage Libraries for

Phage Therapy of Human Infections. Front. Microbiol. *10,* 2537.

Green, S. I. *et al.* (2017a). Bacteriophages from ExPEC Reservoirs Kill Pandemic

Multidrug-Resistant Strains of Clonal Group ST131 in Animal Models of Bacteremia. Sci

Rep. *7*, 46151.

Gribble, F. M. and Reimann, F. (2016). Enteroendocrine Cells: Chemosensors in the

Intestinal Epithelium. Annu. Rev. Physiol. *78*, 277–299.

Grose, J. H. and Casjens, S. R. (2014). Understanding the enormous diversity of

bacteriophages: The tailed phages that infect the bacterial family Enterobacteriaceae,

Virology. *468*, 421–443.

Hanfling, P. *et al.* (1996). Analysis of the enzymatic cleavage (beta elimination) of the

capsular K5 polysaccharide of *Escherichia coli* by the K5-specific coliphage:

reexamination. J. Bacteriol. *15*, 4747–4750.

Heintz-Buschart, A. and Wilmes, P. (2018). Human Gut Microbiome: Function Matters.

Trends Microbiol. *7*, 563–574.

Hoyles, L. *et al.* (2014). Characterization of virus-like particles associated with the

human faecal and caecal microbiota. Res. Microbiol. *10*, 803–812.

Huang, S. H. *et al.* (2001). A novel genetic island of meningitic *Escherichia coli* K1

containing the ibeA invasion gene (GimA): Functional annotation and carbon-source-

regulated invasion of human brain microvascular endothelial cells. Funct. Integr.

Genomic. *5*, 312-322.

In, J. G. *et al.* (2016). Human mini-guts: New insights into intestinal physiology and

host-pathogen interactions. Nat. Rev. Gastroenterol. Hepatol. *11*, 633–642.

Jault, P. *et al.* (2019). Efficacy and tolerability of a cocktail of bacteriophages to treat

burn wounds infected by *Pseudomonas aeruginosa* (PhagoBurn): a randomised,

controlled, double-blind phase 1/2 trial. Lancet Infect. Dis. *1*, 35-45.

Jiao, X. *et al.* (2007). Heparan sulfate proteoglycans (HSPGs) modulate BMP2

osteogenic bioactivity in C2C12 cells. J. Biol. Chem. *2*, 1080–1086.

Johansson, M. E. V. and Hansson, G. C. (2016). Immunological aspects of intestinal

mucus and mucins. Nat. Rev. Immunol. *10*, 639–649.

Johansson, M. E. V., Larsson, J. M. H. and Hansson, G. C. (2011). The two mucus

layers of colon are organized by the MUC2 mucin, whereas the outer layer is a

legislator of host-microbial interactions. Proc. Natl. Acad. Sci. *108*, 4659–4665.

Johnson, J. R. *et al.* (2013). Abrupt emergence of a single dominant multidrug-resistant

strain of *Escherichia coli*. J. Infect. *6*, 919-928.

Johnson JR, R. T. (2002). Extraintestinal pathogenic *Escherichia coli:* "The other bad *E coli*". J. Lab. Clin. Med*. 3,* 155-162*.*

Kahya, H. F., Andrew, P. W. and Yesilkaya, H. (2017). Deacetylation of sialic acid by esterases potentiates pneumococcal neuraminidase activity for mucin utilization, colonization and virulence. PLOS Pathog. *3*, e1006263

Knecht, L. E., Veljkovic, M. and Fieseler, L. (2020). Diversity and Function of Phage Encoded Depolymerases. Front. Microbiol. *10*, 29449.

Korf, I. H. E. *et al.* (2019). Still Something to Discover: Novel Insights into *Escherichia coli* Phage Diversity and Taxonomy. Viruses. *5*, 454.

Latka, A. *et al.* (2017). Bacteriophage-encoded virion-associated enzymes to overcome the carbohydrate barriers during the infection process. Appl. Microbiol. *8*, 3103-3119.

Loc-Carrillo, C. and Abedon, S. T. (2011). Pros and cons of phage therapy. Bacteriophage. *2*, 111-114.

Ma, L. *et al.* (2018). Metals Enhance the Killing of Bacteria by Bacteriophage in Human Blood. Sci. Rep.*1,* 2326*.*

Mammarappallil, J. G. and Elsinghorst, E. A. (2000). Epithelial Cell Adherence Mediated

by the Enterotoxigenic *Escherichia coli* Tia Protein, *12*, 6595–6601.

Mapes, A. C. *et al.* (2016). Development of expanded host range phage active on

biofilms of multi-drug resistant *Pseudomonas aeruginosa*. Bacteriophage. 1, e1096995.

De Martel, C. *et al.* (2012). Global burden of cancers attributable to infections in 2008: A

review and synthetic analysis. Lancet Oncol. *6*, 607-615.

Maruvada, P. *et al.* (2017). The Human Microbiome and Obesity: Moving beyond

Associations. Cell Host Microbe. *5*, 589-599.

Mathers, A. J., Peirano, G. and Pitout, J. D. (2015). *Escherichia coli* ST131: The

quintessential example of an international multiresistant high-risk clone. Adv. Appl.

Microbiol. *90*, 109–154.

McGuckin, M. A. *et al.* (2011). Mucin dynamics and enteric pathogens. Nat. Rev

Microbiol. *4*, 265–278.

Mitsuoka, T., Hayakawa, K. and Kimura, N. (1975). The fecal flora of man, Zentralbl

Bakteriol. Orig. A. *4*, 499–511.

Monzon, M. E., Casalino-Matsuda, S. and Forteza, R. M. (2006). Identification of

glycosaminoglycans in human airway secretions. Am. J. Resp. Cell. Mol. *2*, 135–141.

1001

1002   Moye, Z. D., Woolston, J. and Sulakvelidze, A. (2018). Bacteriophage applications for

1003   food production and processing. Viruses. *2*, 135.

1004

1005   Murphy, K. J. *et al.* (2004). A new model for the domain structure of heparan sulfate

1006   based on the novel specificity of K5 lyase. J. Biol. Chem. *26*, 27239–27245.

1007

1008   Nataro, J. P. *et al.* (1985). Detection of an adherence factor of enteropathogenic

1009   *Escherichia coli* with a dna probe. J. Infect. *3*, 560-565.

1010

1011   Nataro, J. P. (2005) *Colonization of Mucosal Surfaces*. Edited by W. N. J. Nataro,

1012   James P. Cohen S. Paul, Mobley L.T. Harry. ASM. doi: 10.1128/9781555817619.

1013

1014   Nicolas-Chanoine, M. H., Bertrand, X. and Madec, J. Y. (2014). *Escherichia coli* st131,

1015   an intriguing clonal group. Clin. Microbiol. Rev. *3*, 543-574.

1016

1017   Nobrega, F. L. *et al.* (2018). Targeting mechanisms of tailed bacteriophages. Nat. Rev.

1018   Microbiol. *12*, 760–773.

1019

1020   O'Leary, T. R., Xu, Y. and Liu, J. (2013). Investigation of the substrate specificity of K5

1021   lyase A from K5A bacteriophage. J. Glycobiol.*1*, 132–141.

1022

1023 O'Neill, J. (2016). Antimicrobial Resistance : Tackling a crisis for the health and wealth

1024 of nations, Review on Antimicrobial Resistance.

1025

1026 Ota, H. *et al.* (1998). New monoclonal antibodies against gastric gland mucous cell-type

1027 mucins: A comparative immunohistochemical study. Histochem Cell Biol. 2, 113-119.

1028

1029 Park, H. *et al.* (2017). Heparan sulfate proteoglycans (HSPGs) and chondroitin sulfate

1030 proteoglycans (CSPGs) function as endocytic receptors for an internalizing anti-nucleic

1031 acid antibody. Sci .Rep.*1*, 14373.

1032

1033 Pickard, J. M. *et al.* (2017). Gut microbiota: Role in pathogen colonization, immune

1034 responses, and inflammatory disease. Immunol. Rev. *1*, 70–89.

1035

1036 Pires, D. P. *et al.* (2016). Bacteriophage-encoded depolymerases: their diversity and

1037 biotechnological applications. Appl. Microbiol .Biotechnol. *5*, 2141–2151.

1038

1039 Pitout, J. D. D. (2012). Extraintestinal pathogenic *Escherichia coli:* A combination of

1040 virulence with antibiotic resistance. Front. Microbiol. *3*, 9.

1041

1042 Poole, N. M. *et al.* (2017). Role for FimH in extraintestinal pathogenic *Escherichia coli*

1043 invasion and translocation through the intestinal epithelium. Infect. Immun. *11*. e00581-

1044 17.

1045

1046

1047 Poole, N. M., Rajan, A. and Maresso, A. W. (2018). Human Intestinal Enteroids for the

1048 Study of Bacterial Adherence, Invasion, and Translocation. Curr. Protoc. Microbiol. *1*,

1049 55.

1050

1051 Poolman, J. T. and Wacker, M. (2016). Extraintestinal pathogenic *Escherichia coli*, a

1052 common human pathogen: challenges for vaccine development and progress in the

1053 field. J. Infect. *1*, 6–13.

1054

1055 Price, L. B. *et al.* (2013). The epidemic of extended-spectrum-β-lactamase-producing

1056 *Escherichia coli* ST131 is driven by a single highly pathogenic subclone, H30-Rx. mBio.

1057 6, e00377-13.

1058

1059 Rajan, A. *et al.* (2018). Novel segment- and host-specific patterns of enteroaggregative

1060 *Escherichia coli* adherence to human intestinal enteroids. mBio. 1, e02419-17.

1061

1062

1063 Ren, H. *et al.* (2019). Genome sequence analysis of Vibrio parahaemolyticus lytic

1064 phage Vp_R1 with a C3 morphotype. Arch. Virol. *11*, 2865–2871.

1065

1066 Roberts, I. S. (1996). The biochemistry and genetics of capsular polysaccharide

1067 production in bacteria. Annu. Rev. Microbiol. *50*, 285–315.

1068

1069   Russo, T. A. and Johnson, J. R. (2003). Medical and economic impact of extraintestinal

1070   infections due to Escherichia coli: focus on an increasingly important endemic problem,

1071   Microbes. Infect. *5*, 449–456.

1072

1073   Sadowska, A. M. *et al.* (2006). Role of N-acetylcysteine in the management of COPD,

1074   Int. J. Chron. Obstruct. Pulmon. Dis. *4*, 425–434.

1075

1076   Saito, C., Zwingmann, C. and Jaeschke, H. (2010). Novel mechanisms of protection

1077   against acetaminophen hepatotoxicity in mice by glutathione and N-acetylcysteine.

1078   Hepatology.*1*, 246–254.

1079

1080   Sampson, T. R. and Mazmanian, S. K. (2015). Control of brain development, function,

1081   and behavior by the microbiome. Cell Host Microbe. *5*, 565–576.

1082

1083   Sansonetti, P. J. (2004). War and peace at mucosal surfaces. Nat. Rev. Immunol. *12*,

1084   953–964.

1085

1086   Sato, T. *et al.* (2011). Paneth cells constitute the niche for Lgr5 stem cells in intestinal

1087   crypts. Nature. *7330*, 415–418.

1088

1089   Sausset, R. *et al.* (2020). New insights into intestinal phages. Mucosal Immunol. 2, 205–

1090   215.

1091

1092 Savalia, D. *et al.* (2008). Genomic and Proteomic Analysis of phiEco32, a Novel

1093 *Escherichia coli* Bacteriophage. J. Mol. Biol. *3*, 774–789.

1094

1095 Saxena, K. *et al.* (2016). Human Intestinal Enteroids: a New Model To Study Human

1096 Rotavirus Infection, Host Restriction, and Pathophysiology. J. Virol.*1*, 43–56.

1097

1098 Schooley, R. T. *et al.* (2017). Development and use of personalized bacteriophage-

1099 based therapeutic cocktails to treat a patient with a disseminated resistant

1100 Acinetobacter baumannii infection. Antimicrob. Agents Chemother*. 10*, e00954-17.

1101

1102 Severi, E., Hood, D. W. and Thomas, G. H. (2007). Sialic acid utilization by bacterial

1103 pathogens. Microbiology. *9*, 2817–2822.

1104

1105 Shkoporov, A. N. *et al.* (2018). Reproducible protocols for metagenomic analysis of

1106 human faecal phageomes. Microbiome. *1*, 68.

1107

1108 Shkoporov, A. N. and Hill, C. (2019). Bacteriophages of the Human Gut: The "Known

1109 Unknown" of the Microbiome. Cell Host Microbe. *2*, 195–209.

1110

1111 Sturmer, R. *et al.* (2018). Commercial Porcine Gastric Mucin Preparations, also Used as

1112 Artificial Saliva, are a Rich Source for the Lectin TFF2: In Vitro Binding Studies.

1113 Chembiochem, *24*, 2598–2608.

1114

1115 Sugahara, K. and Kitagawa, H. (2002). Heparin and heparan sulfate biosynthesis,

1116 IUBMB Life. *4*, 163–175.

1117

1118 Tang, W. H. W., Li, D. Y. and Hazen, S. L. (2019). Dietary metabolism, the gut

1119 microbiome, and heart failure. Nat. Rev. Cardiol. *3*, 137–154.

1120

1121 Terwilliger, A. L. *et al.* (2020). Tailored Antibacterials and Innovative Laboratories for

1122 Phage (Φ) Research: Personalized Infectious Disease Medicine for the Most Vulnerable

1123 At-Risk Patients. PHAGE. 2, 66-74.

1124

1125 Theriot, C. M. *et al.* (2014). Antibiotic-induced shifts in the mouse gut microbiome and

1126 metabolome increase susceptibility to Clostridium difficile infection. Nat. Commun. *5*,

1127 3114.

1128

1129 Tsikas, D. *et al.* (1998). Analysis of cysteine and N-acetylcysteine in human plasma by

1130 high-performance liquid chromatography at the basal state and after oral administration

1131 of N-acetylcysteine. J. Chromatogr. B. Biomed. Sci. Appl. *1–2*, 55–60.

1132

1133 Vatanen, T. *et al.* (2018). The human gut microbiome in early-onset type 1 diabetes

1134 from the TEDDY study. Nature. *7728*, 589–594.

1135

1136 Vimr, E. and Lichtensteiger, C. (2002). To sialylate, or not to sialylate: that is the

1137   question. Trends Microbiol. *6*, 254–257.

1138

1139   Wang, R. and Hasnain, S. (2017). Analyzing the Properties of Murine Intestinal Mucins

1140   by Electrophoresis and Histology. BIO-PROTOCOL. doi: 10.21769/bioprotoc.2394.

1141

1142   Youle, M., Haynes, M. and Rohwer, F. (2012). Scratching the surface of biology's dark

1143   matter. Viruses. doi: 10.1007/978-94-007-4899-6_4.

1144

1145   **Tables**

Table 1. Properties of E. coli phages used in mucin screen.

Phages	ΦES12	ΦES17	ΦES19	ΦJ2W	ΦJ3M	ΦJ1M	ΦJ1S	ΦUlt1	ΦShp1	ΦM1M	ΦM1S	ΦT1S
Source	Human	Human	Human	Human	Human	Human	Human	Human	Sheep	Rabbit	Rabbit	Pigs
Source location	Sewage	Sewage	Sewage	Sewage	Sewage	Sewage	Sewage	Sewage	Animal facility	Animal facility	Animal facility	Animal facility
Isolation strain	JJ2050	JJ2547	DS104	JJ2528	JJ2528	JJ2528	JJ2528	JJ1901	JJ1901	JJ1901	JJ1901	JJ1901
Plaque size (mm)	0.5	0.5 – 1.0	0.5	0.5	0.5	0.5	0.5	0.5 – 1.0	0.5	0.5	0.5	0.5
Plaque morphology	Clear	Clear, halo	Clear, halo	Clear, halo	Clear	Clear	Clear	Clear	Clear	Clear	Clear	Clear
ST131 lysed (%)[a.]	9/13 (69)	9/13 (69)	11/13 (85)	-	-	-	-	12/13 (92)	12/13 (92)	11/13 (85)	12/13 (92)	10/13 (77)
Accession #	MN508614	MN508616	MN508617	-	-	-	-	-	-	-	-	-

[a.] ExPEC ST131 library of isolates described previously (Gibson et al., 2019).

1146

1147

1148

1149

1150

1151

1152

1153

1154

 Supplemental Figures, Methods and References

Figure S1

1156

1157 **Figure S1** (i) ExPEC levels in CM assay incubated with antibiotics (ii) chloramphenicol (Cm)
1158 ciprofloxacin (CIP) or (iii) ampicillin (AMP). (iv) Commensal bacterial levels incubated in
1159 antibiotics indicated. (N=3-5). Exp. performed 1X. Squares indicate individual biological
1160 replicates from independent cultures. NS not significant." = none detected. Mean, ±SD shown.
1161

Figure S2

1162

1163 **Figure S2** (A) ExPEC levels in CM (FS) with phage EC.1. (N=3) (Bi) Mice were gavaged daily
1164 with N-acetyl cysteine for 2 weeks. ExPEC was gavaged on day 0 following protocol from Figure
1165 1A. (Bii) Small intestinal ExPEC levels. (Biii) Large intestinal ExPEC levels. (N=9). Exp.
1166 performed 1X. Squares indicate individual biological replicates from independent cultures or
1167 individual mice. NS not significant." = none detected. Mean, ±SD shown.
1168

Figure S3

1169

1170

1171 **Figure S3** (A) ExPEC levels from 0 to 8 hours in LB plus mucin concentrations indicated and +/-
1172 phage HP3. (B) ExPEC levels with phages HP3 or ES17 (MOI of 10) in LB plus mucin
1173 concentrations indicated. (N=3). ****p < 0.0001. " = none detected. Exp. performed 1X. Squares
1174 indicate individual biological replicates from independent cultures. Mean, ±SD shown.
1175

1176

1177

Figure S4 (A) Graphical representation of BLAST analysis of phage ES17 (bottom axis;
MN508615) and phage phiECo32 (top axis; EU330206.1) using Kablammo web-based software
(Wintersinger and Wasmuth, 2015). Trapezoids drawn between the axis indicate individual
BLAST alignments between the two sequences. The stronger alignments are shaded darker.
(Bi-ii) TEM images of phage HP3 on ExPEC fixed 10 min post-incubation. White bar scale is 1
μm. Red arrows indicate phage. (Bii) Quantification of attached phages per field view. Mean,
±SD shown.
1185

1186

Figure S5 (Ai) BLAST analysis of ES17-TFP (top axis) and phage LM33_P1 tail fiber (bottom
axis, ref|YP_009324518.1|)) and (Aii) *E. coli* phage vB_EcoP_WFI101126 tail fiber (bottom axis,
gb|QBQ76440.1|) (B) Network pictured. An E-value of 1E-1 was used as a cut off for node
inclusion. Each node represents a single protein. Edges connect nodes with a BLAST E-value
of less than 1E-10. The yellow node is the ES17-TFP, red nodes are proteins that belong to the
pectin lysase fold/virulence factor family (IP011050), green nodes are proteins that are
predicted peptidases; blue nodes are phage tail fibers; and gray nodes are uncharacterized
proteins. (C) Protein SDS-PAGE gel showing purified ES-TFP pooled elutions and protein
ladder.

Figure S6

A. TFP predicted structure comparison to K5 lyase

i.

TFP   K5 Lyase (PDB: 2X3H)   Sugar-binding region   TFP   K5 Lyase

ii.

Heparosan/K5   Heparan-sulfate family

B. Glycan Microarray

i. PGM Array (Porcine Gastric Mucin)

ii. CF Array

iii. Glycosaminoglycans (GAGs)

#1-20 (Hyaluronic Acid)   #21-63 (Chondroitin sulfate)   #64-173 (Heparan Sulfate)

C. HP3 antibody specificity

ΦHP3   ΦES17
40X   40X

1196

Figure S6 (Ai) Structures show the predicted structure of tail fiber (blue), the structure of K5
lyase (PDB:2X3H; red). Identical residues between K5 lyase and ES17 tail fiber protein are
yellow. (Aii) Structures of heparosan or K5 capsule and heparan-sulfate derivative. (B) Glycan
microarray analysis in relative fluorescent units (RFUs) using purified ES17-TFP as the GBP
(glycan binding protein) and glycans derived from purified (Bi) porcine gastric mucin, (Bii) other
mammalian glycans (CF array, see methods) and (Biii) glycosaminoglycans. N=3. Experiment
was performed 3X. (C) Immunofluorescent staining of ES17 and HP3 phages fixed on a slide
using antibodies generated against HP3 and secondary fluorescently conjugated (Alexa Fluor
594; Red).
1206

1207 **Supplemental Methods**

1208

1209 Antibiotic treated CM

1210

1211 For antibiotic treated cecal medium, CM was incubated with chloramphenicol (10

1212 ug/mL, EMD Millipore) or ampicillin sodium salt (100 ug/mL) or ciprofloxacin (10 ug/mL)

1213 throughout the 4.5 hour infection using ExPEC JJ1901 (Figure S1ii) or JJ2528 (Figure

1214 S1iii) or commensal *E. coli* ECN (Figure S1iv).

1215

1216 NAC Mouse Model

1217

Mice (BALB/c) were housed individually and gavaged daily with NAC (N-acetyl cysteine)

at a concentration of 40 mg/mouse/day throughout the experiment as indicated (14

days). This was a tolerable concentration and shown to be effective in the mouse small

intestine (De Lisle, Roach and Jansson, 2007). Mice were infected as described in the

primary methods, then euthanized on day 6 and plated to determine bacterial

concentration in the intestinal contents.

1224

1225 Genome Comparison

1226 Genome comparison was performed using a BLAST analysis of phage ES17

1227 (MN508615) and phiECO32 (EU330206.1) and visualized using Kablammo software

1228 (Wintersinger and Wasmuth, 2015).

1229

1230 ES17-TFP Construct Generation

1231

The construct used to purify tail fiber protein (TFP) was prepared by Genewiz (South

Plainfield, NJ) using established protocols. The construct sequence for TFP was

synthesized from the phage ES17 sequence (MN508615.1) nucleotides 58,576-61,641,

ttaacgcagggcacgtctactattagcgaagcgcatagcctctgcctgagtagtgaactccgctacgaagtaatcagcga

ggcgtaggactactccgcctgtaccggctgtacttaggcgcaggttcactgaaccggaagcactagcctgagctacacca

aaggtactcacccactggccccccttagtaaatacttcacctacattgatatctccaccgatgcggaagtttgtaggggttcc

ctgtaccagttcagtgctaatgccccatacataccagttaccggaagttatctgaaatgcagcaccagtaagctcatactca

gcagcaccagggatttcgaagtacatgcagctctgagataactggccttgtggctgcaaactcccagtgctgcttgccgta

ccactaaattcagcagtacctccatcgaagggtttaactaccagcttattaccaggtacacgatagatgtatccttccggca

ggtatccacgcatactcatattaggtgtccagctttgcccaatcttgcctgtctgagccgctacagaacgtaccattacatcttt

gccggagctaccattcaggtacaattgatcagccacaaaactgctgttggcatctacagagatatccatgtggccatctga

attatctactcggcagccataagtaacaaggttggagttctgaagctctacgttattaaggtatgtgtactctgcaaagaagat

actggtatcaataagtttaaggacacgaggcttctgagatacaccatctacattaatagctggagtagtagcgatagcttcc

atccatacattggatacaccacatggggcagtagggcagtcaccatccttgttcttgaggtaaataccaccaccgccaga

ggcttccataatactatctttaatccaccagccgcctacaccacctccgaatgtaccaggcgcttctcctgttgtagcgttcaa

gtaaacacagtaagtgctaataccatcaaagtgcatgttctgaagggtgtcagagcctaagtgcatctcagaagcagact

gtgcccagtaaccccagtcacaaccttgcacgccaatgcctttccatgtattacctatattcccggaaggtttctgaatggcct

tgttaatgttatggatgtatacgtcactaaagttgtgacgaccagcatattggtcggtcgggtcaaaactaatacctacagcc

ccggttgtcccagaaccatctatagcaaaaccaccactatggcggaattgccaatgaggcgtagatgccttaacctgaat

agtgtctgtaaagtttgacggagctttcagtaccgtggctccaccaacaccctttaactgtgtactaccaatagtaatcgcttc

actgtattcaccggggggaaacaattaagtcgttttttggtaacgatgcatttaccgcagcttgtaaggtagggaaatctcctga

atgcttaatgtcgttgttacggacaagcaaacttctctcaccactgccaatgttttttagccccatctttactggaaagactctca

acagaattaccattttcaaaggtaataagggatgcaccttcacctgcattactacttttcagccttgccagcaaatctgcttca

cttgtatgcaactgtctcaaagcaaccgcgtcattatcatctacagcatcaggaattactggagattgtgtgaaagtaacctt

gttggtcatcgaggtgatatcagagttatcaccactcttagctgcaactagagctagtcgagcattctctgatgaagtagctcc

tgtaccacctttggtaataggtacaattgattcagtggaaaccaaacctaaaccaagtttagtacgtgcagcacttgcagta

gttgcccctgtgccaccctgagagatagtcacagggcgagcggaagacagctcttctcgtactgtaaatactcgggaacc

aactcctccagtccagcttacagagaaaactgagtagtagttattattgtttgtccctggcaccactcgtacctcacctcctcct

gtaactacgccaatgacttcaagatataacccataattggcaggataatttatctcagcaggggcatttgtaatattcgacatt

aatactgaatagttacctccattgacaaaattaaatgtttgccagtccatagccttcactttggttaaggtagggctaccaatcc

catgctcagcgagtccaatatttgtaatagctgcacttgtgtcggttatttcggataggttattagttacactaaaagctcccact

tgagcagctgtgtagtctccagtagcaggaactacagcacccttacgtccattgaatgtggagacagaatcaaaaaccac

tgaacctgtttgttcagcgccgccattgattgaaacagatcgaacagatggaatcaagttggcaatatctgacatgtcagct

cggtataaaccaccatcatctttatctacatagaaaccatagtcatcaacaacaggggtaccaacaggaacgcttccaat

atcaccgattgttccgtcagtaccatcacgccccggtttaccttctggaataccaaaagcaaagacacctgttgcattatcat

aagaccccgtagctaggcttcccggaggaagaagcgttgtttgaactgtcatgttagagataatgtcaagttgtgctagaac

gttttgagttgcctcttttgaaagctcagttgctgcattggtgttgtcaattgcagttcctgtggattgaatcaaacctaagagcac

ttgttcttgtgactcaaagtcactcttgagttgtttaatgtcttcttccagtagatgaccttgagcatacagacgttctacttcagcc

aaaatctcctctactgaggagtatttagtttgggacaacaatgcccaatatttagagtctgctgcatattcagctgctgatttata

agcaccaatagttccctcagtagctccaaattgtcctacattgttaacattcttaggtgcttggttgttataaacgatcat.

1272

1273   This sequence was ligated onto pET-21b (+) (ampicillin) vector using restriction

1274   enzymes 5' *BamHI* and 3' *XhoI* by recombination and delivered as a mini-scale DNA

1275   sample.

1276

1277   Protein Purification

1278

BL21 (DE3) cells were used for expression of His_6-TFP under the control of a T7

promoter expression. Cells were induced with Isopropyl ß-D-1-thiogalactopyranoside

(IPTG) 1mM for 2 hrs at 37°C. Then the culture was pelleted at 7000 RPM for 20 min

and resuspended in Binding buffer (50mM NaH_2PO_4, pH 8.0 0.5 M NaCl). Cells were

lysed using a French Press. Following lysing and centrifugation (15,000 RPM for 20

min.) the lysate was added to a Ni-NTA column for binding, washing 4 times in buffer

(Binding buffer with 20mM imidazole) and elution (binding buffer with 250 mM

imidazole). Buffer exchange was performed in PBS. An SDS-PAGE gel was run to

verify the size of the protein along with a ladder (Benchmark Protein Ladder).

1288

1289

1290   HIEM Generation

1291

Human Intestinal Enteroid (HIEs) were cultured from 3D enteroid cultures derived from

human colonic biopsy tissue using methods developed by Sato and Cleavers and 2-D

enteroid monolayers (HEIMs) developed using methods of Van Dussen et al. (Sato *et*

al., 2009, 2011; VanDussen *et al.*, 2015; Saxena *et al.*, 2016). Briefly, HIEs were

derived from human biopsy samples. The colonic crypts were isolated, embedded in

Matrigel (Corning) or collagen (Sigma) and incubated with complete growth media with

growth factors (CMGF+) for differentiation into 3D enteroid structures. Next, chambered

slides (Greinier Bio-One) were coated with Matrigel or collagen, dissociated, about 50

spheroids, (3D enteroids cultured CMGF$^+$ for 7 days) were added per well. HIEs were

dissociated using 0.05% trypsin/0.5mM EDTA and a gravity flow strainer. Cells were

allowed to grow and adhere 1-2 days with CMGF$^-$. Then the media was removed and

replaced with Differentiation media. Typically for experiment, HEIMs were used 4-7

days later when fully confluent (>90%) and differentiated. For our experiments this was

day 5. The components of media the CMGF+, CMGF- and Differentiation media are

described previously (Poole, Rajan and Maresso, 2018).

1307

1308 Glycan Microarray

1309 Glycan microarray analysis was performed by the Emory Comprehensive Glycomics

1310 Core using the methods described below.

1311

An oligosaccharide array composed of 560 defined glycans from the Consortium for

Functional Glycomics were obtained from MicroArray Core Facility of the Scripps

Research Institute (La Jolla, CA) (lists can be found at

http://www.functionalglycomics.org/static/consortium/resources/resourcecored2.shtml#

Glycolipid) as described(Heimburg-Molinaro *et al.*, 2011). The glycans were printed on

Nexterion 3-D hydrogel coating (H) slides using a contact printer in replicates of six. A

second microarray consisting of 138 glycans from purified porcine gastric mucin was

also included. And a third microarray consisted of 170 charged glycosaminoglycans

(GAG), including oligomers from hyaluronic acid, chondroitin sulfates and heparan,

obtained from the Emory Comprehensive Glycomics Core (Atlanta, GA). Glycans were

conjugated with the bifunctional linker (AEAB) and printed on Nexterion 3-D hydrogel

coated (H) slides using a contact printer Aushon 2470 (Quanterix, Billerica, MA) in

replicates of four. His-tagged TFP (up to 40 ug/mL) was diluted in PBS buffer with 1%

BSA and 0.05% Tween 20 and incubated on the microarray slides for 1 hr at RT. Slides

were washed with PBS buffer with 0.05% Tween 20 and PBS buffer. Alexa Flour 647

anti-His-tag mouse mAb (MBL International, Woburn, MA) was diluted to 5 µg/ml and

incubated on the slides for 1 hr. After washing and drying, the slides were scanned with

an InnoScan 1100 AL scanner and the data were processed using Mapix 8.2.5 software

(Innopsys, Chicago, IL). Mean relative fluorescence units (RFU) from replicate spots

were averaged after subtracting local median background, and the standard deviation

(STDEV) were calculated and plotted as error bars in the histogram plot of the glycan

array.

1334

1335 Bioinformatics modeling

1336

The phage genome was annotated using the RASTtk (Brettin *et al.*, 2015). The Enzyme

Function Initiative-Enzyme Similarity Tool (EFI-EST) was used to visualize BLAST

results of the tail fiber protein (TFP) (Gerlt *et al.*, 2015). The network visualization

showed that TFP was related to other tail fiber proteins and proteins that belong to the

pectin lyase fold/virulence factor family of proteins (IPR011050). HHpred was used to

identify other proteins with similar structures (Hildebrand *et al.*, 2009). The highest

scoring template was a Tailspike protein from *E. coli* bacteriophage HK620 (PDB: 2VJJ;

E-value = 9.4E-23). The other top five hits from Enterobacteria or *Escherichia* phages

were a tailspike protein from Enterobacteria phage SF6 (PDB: 2VBK; E-value=1.9E-14),

a putative endo-glycosidase tailspike protein (PDB: 6NW9; E-value=3.1E-14), a

colanidase tailspike protein from Enterobacteria phage phi92 (PDB:6E0V; E-

value=1.2E-13), a sugar-binding tailspike from *Salmonella* phage Det7 (PDB:6F7D; E-

value=2.1E-12), and a K5 lyase (E.C.4.2.2.7) tailspike protein from Enterobacteria

phage K5 (PDB: 2X3H; E-value=4E-12). All of these tailspike proteins bind and degrade

polysaccharides and are classified as members of the pectin lyase fold/virulence factor

family (IPR011050)(Freiberg *et al.*, 2003; Barbirz *et al.*, 2008; Schwarzer *et al.*, 2012;

Broeker *et al.*, 2019; Greenfield *et al.*, 2019).The K5 lyase has well-known heparanase

activity due to the similarities between K5 and heparan-sulfate (O'Leary, Xu and Liu,

2013). All six were used to produce a predicted structure of the ES17 tail fiber protein in

Modeler (Zimmermann *et al.*, 2018). Structures were aligned using UCSF Chimera.

Chimera's matchmaker function was used to structurally align the predicted tail fiber 2

protein with the HK620 tailspike and the K5 lyase. Solvent exposed identical residues

between ES17 tail fiber protein 2 and K5 lyase were determined using Chimera's

match-align to determine what residues were both identical and within 5 angstroms of

each other in the structural alignment.

1362

1363 16S rRNA Gene Sequencing

1364

1365 The Center for Metagenomics and Microbiome Research (CMMR) at Baylor College of

1366 Medicine performed 16S rRNA sequencing and analysis.

1367

The 16S rRNA gene sequencing methods were adapted from the methods developed for

the Earth Microbiome Project (Caporaso *et al.*, 2012) and NIH-Human Microbiome Project

(Caporaso *et al.*, 2012; Huttenhower *et al.*, 2012). Briefly, bacterial genomic DNA was

extracted using the Qiagen MagAttract PowerSoil DNA Kit (formerly sold by MO BIO as

PowerMag Soil DNA Isolation Kit). The 16S rDNA V4 region was amplified by PCR using

primers 515F and 806R containing Illumina adapters and a single-index barcodes

(Caporaso *et al.*, 2012). The amplicons were visualized with gel electrophoresis and

purified to remove primer-dimers and non-specific amplicons. The samples were

quantified using the Qubit® fluorometer and pooled at a DNA mass of 100 ng per sample.

The amplicon pool was sequenced on the Illumina MiSeq platform using the reagent kit

v2 (2 × 250 bp) and the paired-end protocol yielding paired-end reads that overlaped

almost completely.

1380

The raw data files in Binary Base Call (BCL) format created by the MiSeq run were first

converted into the FASTQ format and demultiplexed based on the single-index barcodes

using the Illumina BCL2FASTQ software. The demultiplexed FASTQ read pairs were then

merged using USEARCH v7.0.1090 (Edgar, 2010) 'fastq_mergepairs' function, requiring

read pairs to overlap by at least 50 bp, a merged length of at least 252 bp, a truncation

quality above 5, and zero differences in the overlapping region. The merged files were

then filtered further, using the USEARCH70 'fastq_filter' program, and only allowing for a

maximum expected error of 0.05. Merged reads were then combined into a single FASTQ

file, which was filtered for PhiX using Bowtie2 v.2.3.4.3 (Langmead and Salzberg, 2012)

and the 'very-sensitive' parameter setting. After removing PhiX reads, the FASTQ file was

transformed into a FASTA format file. The FASTA file was run through USEARCH70

'derep_fulllength' program and the reads were sorted by size using usearch70 'sortbysize'

program. The reads were clustered into operational taxonomic units (OTUs) at a similarity

cutoff value of 97% using the UPARSE algorithm (Edgar, 2013). This was accomplished

in an iterative stepwise manner in increments of 0.4% using USEARCH70 'cluster_otus'

function. The output from the previous increment was fed into the next iteration, until a

maximum of 3.2% was reached. All the intermediate files of this step were filtered for any

chimeras and after the last run through the above loop, the final output was run through

USEARCH70 'uchime_ref' program against the GOLD database (Kyrpides, 1999;

Mukherjee *et al.*, 2019), using only the plus strand and allowing for no chimeras, in order

to create a clustered OTU file with no chimeras. The OTU file was mapped against an

optimized version of the latest current SILVA Database (Quast *et al.*, 2013) containing

only sequences from the V4 region of the 16S rRNA gene to determine taxonomies. This

step was performed using USEARCH70 'usearch_global' function, specifying the identity

threshold to 96.8%. Abundances were recovered by mapping the demultiplexed reads to

the OUT file and all files created in the loop were then run through a program developed

in-house that resolved the iterative UPARSE steps, creating an OTU table in BIOM format

and removing the chimera and singleton reads. The BIOM file was summarized, recording

the number of reads per sample, and merged with a file that was generated for the overall

read statistics, to produce a final summary file with read statistics and taxonomy

information. ATIMA (Agile Toolkit for Incisive Microbial Analyses) was used to generate

beta diversity graphs.

Supplemental References

Barbirz, S. *et al.* (2008). Crystal structure of *Escherichia coli* phage HK620 tailspike:

podoviral tailspike endoglycosidase modules are evolutionarily related. Mol. Microbiol.

2, 303–316.

Brettin, T. *et al.* (2015). RASTtk: a modular and extensible implementation of the RAST

algorithm for building custom annotation pipelines and annotating batches of genomes.

Sci. Rep.*5*, 8365.

Broeker, N. K. *et al.* (2019). Time-resolved DNA release from an O-antigen-specific. J.

Biol. Chem. *31*, 11751–11761.

Caporaso, J. G. *et al.* (2012). Ultra-high-throughput microbial community analysis on the

Illumina HiSeq and MiSeq platforms, ISME J. doi: 10.1038/ismej.2012.8.

1429

1430   Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST',

1431   Bioinformatics. doi: 10.1093/bioinformatics/btq461.

1432

1433   Edgar, R. C. (2013). UPARSE: Highly accurate OTU sequences from microbial

1434   amplicon reads. Nat. Methods. doi: 10.1038/nmeth.2604.

1435

1436   Freiberg, A. *et al.* (2003). The tailspike protein of Shigella phage Sf6. A structural

1437   homolog of Salmonella phage P22 tailspike protein without sequence similarity in the

1438   beta-helix domain. J. Biol. Chem. *3*, 1542–1548.

1439

1440   Gerlt, J. A. *et al.* (2015). Enzyme Function Initiative-Enzyme Similarity Tool (EFI-EST):

1441   A web tool for generating protein sequence similarity networks. Biochim. Biophys. Acta.

1442   *8*,1019–1037.

1443

1444   Greenfield, J. *et al.* (2019). Structure and tailspike glycosidase machinery of ORF212

1445   from E. coli O157:H7 phage CBA120 (TSP3). Sci. Rep.*1*, 7349.

1446

1447   Heimburg-Molinaro, J. *et al.* (2011). Preparation and analysis of glycan microarrays.

1448   Curr. Protoc. Protein. Sci. doi: 10.1002/0471140864.ps1210s64.

1449

1450   Hildebrand, A. *et al.* (2009). Fast and accurate automatic structure prediction with

1451   HHpred. Proteins. *9*, 128–132.

1452

1453   Huttenhower, C. *et al.* (2012). Structure, function and diversity of the healthy human

1454   microbiome. Nature. doi: 10.1038/nature11234.

1455

1456   Kyrpides, N. C. (1999). Genomes Online Database (GOLD 1.0): A monitor of complete

1457   and ongoing genome projects world-wide. Bioinformatics. doi:

1458   10.1093/bioinformatics/15.9.773.

1459

1460   Langmead, B. and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2',

1461   Nat. Methods. doi: 10.1038/nmeth.1923.

1462

1463   De Lisle, R. C., Roach, E. and Jansson, K. (2007). Effects of laxative and N-

1464   acetylcysteine on mucus accumulation, bacterial load, transit, and inflammation in the

1465   cystic fibrosis mouse small intestine. Am. J. Physiol. Gastrointest. Liver Physiol. doi:

1466   10.1152/ajpgi.00195.2007.

1467

1468   Mukherjee, S. *et al.* (2019). Genomes OnLine database (GOLD) v.7: Updates and new

1469   features. Nucleic. Acids Res. Publ. doi: 10.1093/nar/gky977.

1470   O'Leary, T. R., Xu, Y. and Liu, J. (2013) 'Investigation of the substrate specificity of K5

1471   lyase A from K5A bacteriophage. Glycobiology. *1*, 132–141.

1472

1473   Poole, N. M., Rajan, A. and Maresso, A. W. (2018). Human Intestinal Enteroids for the

1474   Study of Bacterial Adherence, Invasion, and Translocation. Curr. Protoc. Microbiol. *1*,

1475 e55.

1476

1477 Quast, C. *et al.* (2013). The SILVA ribosomal RNA gene database project: Improved

1478 data processing and web-based tools. Nucleic Acids Res. Spec. Publ. doi:

1479 10.1093/nar/gks1219.

1480

1481 Sato, T. *et al.* (2009). Single Lgr5 stem cells build crypt-villus structures in vitro without

1482 a mesenchymal niche. Nature. doi: 10.1038/nature07935.

1483

1484 Sato, T. *et al.* (2011). Long-term expansion of epithelial organoids from human colon,

1485 adenoma, adenocarcinoma, and Barrett's epithelium. Gastroenterology. doi:

1486 10.1053/j.gastro.2011.07.050.

1487

1488 Saxena, K. *et al.* (2016). Human Intestinal Enteroids: a New Model To Study Human

1489 Rotavirus Infection, Host Restriction, and Pathophysiology. J. Virol.*1*,43–56.

1490

1491 Schwarzer, D. *et al.* (2012). A multivalent adsorption apparatus explains the broad host

1492 range of phage phi92: a comprehensive genomic and structural analysis. J. Virol.*19*,

1493 10384–10398.

1494

1495 VanDussen, K. L. *et al.* (2015). Development of an enhanced human gastrointestinal

1496 epithelial culture system to facilitate patient-based assays. Gut. doi: 10.1136/gutjnl-

1497 2013-306651.

1498

1499   Wintersinger, J. A. and Wasmuth, J. D. (2015). Kablammo: An interactive, web-based

1500   BLAST results visualizer. Bioinformatics. doi: 10.1093/bioinformatics/btu808.

1501

1502   Zimmermann, L. *et al.* (2018). A Completely Reimplemented MPI Bioinformatics Toolkit

1503   with a New HHpred Server at its Core. J. Mol. Biol. *15*, 2237–2243.

1504

1505

1506

1507

1508
1509

1510