Antifeedant activities of L-arabinose to caterpillars of the cotton bollworm

***Helicoverpa armigera* (Hübner)**

Running title: Effects of L-arabinose on caterpillars

Longlong Sun[a#], Zhenzhen Hong[a#], Ying Ma[b*], Wenhua Hou[a], Long Liu[a], Xincheng Zhao[a], Fengming

Yan[a], Xinming Yin[a], Chenzhu Wang[c], Qingbo Tang[a*]

[a] Department of Entomology, College of Plant Protection, Henan Agricultural University, Zhengzhou

450002, China;

[b] College of Agronomy, Henan Agricultural University, Zhengzhou 450002, China;

[c] State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology,

Chinese Academy of Sciences, Beijing 100101, China;

[#] These authors contribute equally to this work.

*****Correspondence authors**:

Qingbo Tang, Department of Entomology,College of Plant Protection, Henan Agricultural University,

Zhengzhou, Henan 450002, China. Email: qbtang@henau.edu.cn, qingbotang@126.com

Ying Ma, College of Agronomy, Henan Agricultural University, Zhengzhou, Henan 450002, China. Email:

maying999@126.con

ABSTRACT

Exploring botanical biopesticide is one of the eco-friendly approaches for pest control in current crop protection. L-arabinose, a plant-specific and less absorptive pentose, is known for its selective inhibitory effect on the sucrose digestion in mammals. In this study, we investigated the effects of L-arabinose on the feeding preference, the peripheral gustatory perception, the larval development, as well as the activity of intestinal sucrase of an insect pest species, the cotton bollworm *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae), in an attempt to explore the antifeedant activities of this pentose. The results showed that L-arabinose deterred the feeding preferences of *H. armigera* caterpillars for its two host plants and two phagostimulants, the tobacco, the pepper, the sucrose and the fructose. Gustatory receptor neurons (GRNs) sensitive to L-arabinose were not found in the peripheral maxillary sensilla of *H. armigera* caterpillars, but the sensitivities of GRNs sensitive to sucrose, fructose and tobacco saps were suppressed by the additions of L-arabinose. Exposure of *H. armigera* caterpillars to dietary L-arabinose resulted in a prolonged larval developmental duration, a suppressed activity of intestinal sucrase and a reduced glucose level in midgut. *In vitro*, L-arabinose inhibited activities of the intestinal sucrase of *H. armigera* caterpillars in an uncompetitive manner. Taken together, these findings demonstrate that L-arabinose is a behavioral, gustatory and physiological inhibitor to caterpillars of the extremely generalist pest species *H. armigera*, suggesting the great potential of L-arabinose to be an effective antifeedant.

Keywords: L-arabinose, inhibitor, *Helicoverpa armigera*, feeding preference, taste sensitivity, sucrase activity

Introduction

It is estimated that the insect pests are responsible for damaging between 20 and 40% of global crop production annually (FAO 2019). Though the strategy of the integrated pest management (IPM) has been extensively accepted in the past decades, the use of conventional synthetic insecticides is still the main approach for pest control in current crop protection. In comparison with conventional insecticides, many attentions have been paid to the behavioral manipulation of insect pests for their management in recent years (Dancewicz et al. 2020; Karkanis and Athanassiou 2021; Koul 2005; Walia et al. 2017; Wroblewska-Kurdyk et al. 2020). Of particular interests are nontoxic or less-toxic substances of natural origin that show selective activities toward insect pests but not toward non-target beneficial organisms and the environment (Akhtar et al. 2010; Junhirun et al. 2018; Koul et al. 2003; Ponsankar et al. 2020).

Most natural substances with behavioral manipulating activities could be perceived by the chemosensory system of insect pests, resulting in appetitive feeding behaviors or aversive feeding behaviors(Jiang et al. 2020; Mitri et al. 2009; Sollai et al. 2017; Sollai et al. 2015; Wada-Katsumata et al. 2013). Examples of manipulating activities of insect pests include repellent, antifeedant, deterrent, as well as growth inhibition and negative effects on both fecundity and longevity of the target species (Dancewicz et al. 2020; Hasheminia et al. 2013; Karkanis and Athanassiou 2021; Koul et al. 2003; Mao et al. 2019). Lots of substances originated from plants such as azadirachtin, rotenone, matrine, nicotine, pyrethrins have been well developed and commercially registered in some countries (Cheng et al. 2020; Isman 2020; Pavela 2016; Siegwart et al. 2015; Walia et al. 2017). But there should still be more botanical substances deserved to be explored for biological control of insect pests.

L-arabinose (abbreviated as LA), a naturally-occurring and widely present pentose, is a

plant-specific sugar accounting for 5-10% of cell wall saccharides in varieties of plants

including corn, rice, wheat, barley, rye, and oats (Bengtsson 1990; Garleb et al. 1989;

Saulnier et al. 1995). LA is also known as a less absorptive sugar with a sweet taste and

has been recognized to have a selective inhibitory effect on the intestinal sucrase activity in

mammals, thereby resulting in a delay of sucrose digestion (Halschoujensen et al. 2015;

Seri et al. 1996; Shibanuma and Houda 2011; Yamamoto 2013). In recent years, LA is

widely used as additives in human food products with the intention to combat obesity and

diabetes (DuBois and Prakash 2012; van Opstal et al. 2019).

In regard of inhibiting activity, we wonder whether LA has inhibitory effects on the

digestion of sucrose in insect pests, and thereby LA may have the potential to be a

phytochemical antifeedant. A pioneering report in 2010 found that the exposure of the

sweet potato whitefly *Bemisia tabaci* to dietary arabinose resulted in low survivals of the

nymphs and reduced mean day survivals of adults, suggesting a possible antifeedant

activity of arabinose on this piercing insect species (Hu et al. 2010), but the proofs of how

LA modulate the chemosensory perception and the feeding behaviors of insects were

absent.

The cotton bollworm *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae) is an

extreme generalist agricultural pest in the world, feeding on at least 161 host plant species

in 49 plant families (Fitt 1989; Zalucki et al. 1986). In our preliminary studies, we observed

that the feedings of *H. armgiera* caterpillars could be deterred by LA. Then we postulated

that LA might be an antifeedant to this herbivorous chewing mouthpart pest species. In this

study, we investigated the effects of LA on the feeding behavioral, the peripheral gustatory

perception, the larval development, as well as the activities of intestinal sucrase both in *vitro*

and in *vivo*, with the aim of exploring the antifeedant effects and the possible mechanism of

this pentose to the lepidopteran pest species.

Materials and Methods

Insects

The colony of *H. armigera* was maintained in the laboratory under conditions of photoperiod

(L16:D8), temperature (27 ± 1°C) and 75% relative humidity. Larvae of were reared on an

standard artificial diet (Jiang et al. 2010) or standard artificial diet containing different

concentrations of LA (Sigma-Aldrich) described as below. Adults were supplied with a 10%

v/v solution of sucrose in distilled water with vitamins.

Feeding choice assay

Six behavioral experiments were set to investigate the effects of LA on the feeding

preferences of *H. armigera* caterpillars for plant leaves treated by different concentrations of

LA (Table 1). Leaves from two plant species, the pepper *Capsicum frutescens* L. "Yu-Yi"

and the tobacco *Nicotiana tabacum* "K326", were used to be the feeding mediums

respectively, in an attempt to investigate whether the preferences may change if different

medium leaves were used. Of the 6 experimental settings, experiment II, IV and VI were

used to investigate whether LA could counteract the stimulation effects of sucrose and

fructose after LA was mixed to sucrose or fructose. While experiment III was used to

investigate whether caterpillars more prefer sucrose if the control leaves were treated by

sucrose.

A dual-choice bioassay (Wang et al. 2017) was performed to test feeding preferences

of the 5[th] instar *H. armigera* caterpillars for leaf discs. In general, leaf discs (10 mm

diameter) were punched from fresh plant leaves using a conventional hand-held punch,

followed by immersed in a treatment solution or in a control solution (as shown in Table 1)

for 5 min. For a test of a single caterpillar, a moist medium-speed qualitative filter paper

(Φ11cm, Jiaojie, China) was put on the bottom of a Petri dish (12 cm diameter).Then four

leaf discs treated with a treatment solution (A) and four discs with the control solution (B)

were put on the moist filter paperin an ABABABAB fashion around the circumference. At

test initiation, a single caterpillar was placed in the middle of the dish and the leaf disc

consumption was observed at regular intervals until at least two of the four disks of either

treatment (A) or control (B) had been consumed. Remaining areas of all discs consumed

were measured using transparency film (PP2910, 3M Corp.). The feeding preference index

was calculated as follows.

Preference index= (area of treatment-disc consumed- area of control-disc consumed) /

(area consumed of all discs)

For each setting of choice scenario, the mean consumption of all caterpillars was

calculated.

Electrophysiological recordings

Four electrophysiological experiments were set to investigate the effects of LA on the

sensitivity of gustatory receptor neurons (GRNs) in the styloconic sensilla located on the

galea of 5[th] instar *H. armigera* larvae by using the single sensillum recording (Roessingh et

al. 1999; van Loon 1990). As shown in Table 2, experiment I and II investigated whether

sensitive GNRs to LA were in the lateral sensillum and the medial sensillum. Experiment III

investigated whether LA have effects on the sensitivities of the sucrose-best GRN in the

lateral sensillum to sucrose. Experiment IV investigated whether LA have effects on the

sensitivities of the GRNs sensitive to tobacco leaf saps in the medial sensillum. The

sucrose best GRN in the lateral sensillum and the GRNs sensitive to tobacco-leaf-sap best

in the medial sensillum have been identified in our previous work (Ma et al. 2016; Tang et al.

2014). The leaf saps were extracted as per our previous work (Tang et al. 2014).

Before stimulation, an excised 5[th] instar caterpillar head was mounted on a silver wire

electrode, which was connected to the input of a pre-amplifier (Syntech Taste Probe DTP-1,

Hilversum, The Netherlands). Glass micropipette was filled with stimulating solution and

touched the tip of a sensillum under the control of a micromanipulator. The stimuli and the

stimulating sequence in a single caterpillar were shown in Table 2. Amplified signals were

digitized by an A/D-interface (Syntech IDAC-4) and sampled into a personal computer.

The analysis of responding spikes to different stimuli were performed with the aid of

Autospike version 3.7 software (Syntech, The Netherlands). In general, impulse waveforms

were classified into different groups according to the amplitude, waveform (shape), and

phasic temporal pattern, as well as by visual inspection, and then were counted. In most

recordings, three or four impulse spikes were identified and labeled as small (S),

intermediate (M1 or/and M2), and large (L), which best responded to water,

phagostimulant/deterrent, and salt, respectively (Sollai et al. 2014; Sun et al. 2021).The

mean impulse frequency ($spk.s^{-1}$) of a GRN to one stimulus in the first 1000 ms was

calculated.

Effects of LA on Larval development

Six groups of *H. armigera* neonatal caterpillars were reared by a standard artificial diet

(Jiang et al. 2010) containing different concentration of LA from 0, 6.66 mM, 66.6 mM,

199.8 mM, 333 mM, to 500 mM until the last instar, respectively. The larval developmental

duration and the mortality of caterpillars were recorded. Caterpillars were maintained

through pupation to adult emergence, and then the pupal developmental duration as well as

the pupal weights was recorded.

Effects of LA on enzymatic activity

Six groups of neonate caterpillars of *H. armigera* were reared by standard artificial diets

containing different concentrations of LA from 0, 6.66 mM, 66.6 mM, 199.8 mM, 333 mM, to

500 mM until the second day of the 5th instar. Then the midguts of caterpillars were

extracted and weighted. Midguts were hand homogenized in PBS (pH 7.4) at a ratio of 1.0

g midgut to 6.0 ml PBS, and then were centrifuged at 2500 g for 10 min at 4°C.

Supernatants were used to analyze sucrase activity, sucrose concentration, and glucose

concentration by manufacturer protocols (No. A082-2 for sucrase activity, No.A099 for

sucrose, and No.F006 for glucose; Nanjing Jiancheng Bioengineering Institute, Nanjing,

China).

Kinetic analysis of sucrase inhibition by LA

Midguts of the fifth instar caterpillars of *H. armigera* were dissected and then were rinsed

with ice-cold saline (10%) followed by weighing and hand-homogenization. Homogenized

tissues were centrifuged at 2500 rpm for 10 min at 4°C. Supernatants were transferred to a

fresh tube and stored at -20°C until use.

For the kinetic studies of the inhibitory effects of LA on sucrase activity in caterpillars,

midgut supernatants prepared as above were incubated with increasing concentrations of

sucrose (0, 10 mM, 20 mM, 30 mM, 40 mM, 50 mM, 60 mM) in the absence or presence of

LA (6.66 mM or 9.99 mM) or acarbose (7.35 µM or 11.79 µM; Sigma-Aldrich). Selected

concentrations of these sugars were determined based on preliminary experiments.

The volume of standard reaction assay was 90 µL. After the reaction mixtures incubated

at 37°C for 5 min, reactions were terminated by incubating in boiling water for 5 min. After

cooldown, the D-glucose concentration was measured using a Glucose Oxidase Method

(No. F006; Nanjing Jiangcheng Bioengineering Company, Nanjing, China). Specific activity

was calculated as micromoles of substrate hydrolyzed per milligram protein per min. Each

assay was repeated at least three times and results were analyzed by Lineweaver-Burk

plots.

Data analysis and statistics

All data are presented as mean ± SEM. A one-sample t-test was used to compare the

means of the preference indices of caterpillars between for control leaves and for treatment

leaves ($P < 0.05$). The levels of significance were as follows: NS (non-significant): $P \geq 0.05$,

*: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$. For statistical analysis of neuronal responding

frequency to stimuli, the original responding frequencies were square-root transformed if

the variances were not equal. A one-way ANOVA followed by a Scheffé test was used to

compare the mean responding frequency of the same GRN to stimuli with different

concentrations ($P < 0.05$).

The data of the developmental parameters and the midgut enzyme activities were log-

transformed for uneven sample sizes. Means were compared by one-way ANOVA followed

by a Scheffé Test ($P < 0.05$). All statistical analyses were performed using SPSS version

10.0 (SPSS Inc., Chicago, USA).

Results

Inhibition of LA on feeding behaviors

The results of the feeding preference of the 5[th] instar caterpillars of *H. armigera* for LA

treated tobacco leaves and for pure distilled water treated tobacco leaves were shown in

figure 1. It showed that low concentrations of LA from 0.0001 to1.0 mM had no significant

effects on the preferences of caterpillars (Fig. 1A: 0.0001mM, $t(107) = 0.409$, $P = 0.686$;

0.001mM, $t(86) = -0.935$, $P = 0.352$; 0.01mM, $t(160) = -1.157$, $P = 0.249$; 0. 1mM, $t(117) = -$

1.319, $P = 0.190$; 1.0 mM, $t(224) = -1.679$, $P = 0.095$). But high concentrations of LA ≥ 10

mM significantly deterred the feeding of caterpillars (Fig. 1A: 10mM, $t(272) = -2.77$, $P =$

0.006; 66.6mM, t(243) = -3.491, P = 0.001; 133mM, t(190) = -5.88, P < 0.0001; 200mM,

t(311) = -6.359, P < 0.0001; and 333mM, t(100) = -5.743, P < 0.0001).

We also found LA could counteract the phagostimulant effects of sucrose on *H.*

armigera caterpillars. Caterpillars significantly preferred 200 mM sucrose-treated leaves

over control leaves (t (81) = 5.117, P < 0.0001), whereas caterpillars preferred similarly to

control leaves and leaves treated by mixtures of 200 mM sucrose with 100, 300, and 400

mM LA, respectively (200mM sucrose + 100mM LA, t(154) = -0.048, P = 0.962; 200mM

sucrose +300mM LA, t(55) = -0.156, P = 0.877; 200mM sucrose +400mM LA, t(56) = 0.138,

P = 0.891) (Fig. 1B). With the concentration of LA increasing in mixtures, caterpillars

exhibited aversive feedings to mixture-treated leaves (200mM sucrose + 500 mM LA, t(83)

= -4.224, P = 0.0001; 200 mM sucrose + 600 mM LA, t(81) = -3.118, P = 0.0025; 200 mM

sucrose + 800 mM LA, t(23) = -2.876, P = 0.0085) (Fig. 1B).

Next, we examined whether *H. armigera* caterpillars could distinguish between tobacco

leaves treated by 10 mM sucrose and leaves treated by mixtures of 10 mM sucrose with

different concentrations of LA. It showed that low concentrations of LA from 1.0 mM to 100

mM in mixtures had no significant effects on the preferences (10 mM sucrose + 1mM LA,

t(110) = 1.363, P = 0.176; 10 mM sucrose + 10mM LA, t (105) = -0.301, P = 0.764; 10 mM

sucrose + 100mM LA, t(70) = -1.121, P = 0.266), but leaves treated by mixtures containing

LA ≥ 200 mM drove significant aversive feedings of caterpillars (10 mM sucrose + 200 mM

LA, t(69) = -3.362, P = 0.001; 10 mM sucrose + 300 mM LA, t(46) = -4.396, P < 0.0001; 10

mM sucrose + 400 mM LA, t(77) = -13.09, P < 0.0001) (Fig. 1C).

We also found that LA could counteract the stimulating effect of another

phagostimulant, the fructose. Single 200 mM fructose significantly elicited appetitive

feedings of *H. armigera* caterpillars (t(101) = 6.722, P < 0.0001), but mixtures containing

200 mM fructose with 300mM or 400mM LA induced a similar preferences as controls (200

mM fructose + 300 mM LA, t(102) = -1.911, P = 0.059; 200 mM fructose + 400 mM LA,

t(109) = -0.677, P = 0.500) (Fig. 1D). When the concentration of LA in mixtures increased to

500 mM and 600 mM, mixture-treated leaves significantly drove aversive feedings of *H.*

armgiera caterpillars (200 mM fructose+ 500 mM LA, t(113) = -4.115, P = 0.0001; 200 mM

fructose + 600 mM LA, t(100) = -0.3.572, P = 0.0005) (Fig. 1D).

When the pepper leaves were used as the plant leaf medium, LA still showed inhibiting

effects on feedings of *H. armigera* caterpillars. Caterpillars showed more and more aversive

feedings for LA-treated pepper leaves with the concentration of LA increasing (Fig. 2A:

0.01mM LA, t(62) = 0.124, P = 0.902; 0.1mM LA, t(252) = -1.315, P = 0.190; 1.0 mM LA,

t(214) = -1.493, P = 0.137; 10mM LA, t(325) = -2.251, P = 0.025; 100mM LA, t(155) = -

2.028, P = 0.044; 200mM LA, t(115) = -2.244, P = 0.027; 300mM LA, t(95) = -5.211, P <

0.0001; 400mM LA, t(107) = -4.360, P < 0.0001).

Similarly, LA also could counteract the phagostimulant effects of sucrose on *H.*

armigera caterpillars when the pepper leaves were used as the medium leaves. Mixtures

containing 200 mM sucrose with LA from 100 mM to 500 mM still induced the preference of

caterpillars for sucrose (Fig. 2B: 200 mM sucrose + 0 LA, t(62) = 5.718, P < 0.0001; 200

mM sucrose + 100 mM LA, t(33) = 4.630, P = 0.0001; 200 mM sucrose + 300 mM LA, t(41)

= 2.899, P = 0.0060; 200 mM sucrose + 500 mM LA, t(38) = 4.521, P = 0.0001). Whereas

LA in mixtures ≥ 600 mM, the preferences for sucrose were significantly counteracted (Fig.

2B: 200 mM sucrose + 600 mM LA, t(76) =-0.564, p = 0.574; 200 mM sucrose + 800 mM

LA, t(75) = -0.851, p = 0.398). When LA in mixtures increased to 1000 mM, caterpillars

significantly preferred control leaves over mixture-treated leaves (Fig. 2B: 200 mM sucrose

+ 1000 mM LA, t(71) = -2.110, p = 0.038).

Inhibition of LA on gustatory sensitivity

Gustatory perception in *H. armigera* caterpillars are mainly dominated by two pairs of

gustatory styloconic sensilla, the lateral sensilla and the medial sensilla (Sun et al. 2021;

Tang et al. 2014; Zhou et al. 2010), located on the maxillary galea of mouthparts (Figs. 3A

and 3B). But our current results showed LA from 0.0001 mM to 10 mM stimulated very low

responses, which is similar to those induced by 2 mM KCl in the response to the two

styloconic sensilla (Fig. 3C), suggesting non-specific GRN housed in both the two sensilla

were sensitive to LA.

We then investigated whether LA affect the sensitivities of GRNs sensitive to other plant

substances, such as the phagostimulant sucrose and the leaf saps from the host plant

tobacco, by testing the responses of sensillum to mixtures containing LA and these

substances. Firstly we tested the effects of LA on the sensitivity of the sucrose-best GRN in

the lateral sensillum of the 5th instar *H. armigera* caterpillars. The results showed that both

2.0 mM KCl and 10 mM LA only induced weak responses of a water-like L-GRN and a salt-

like S-GRN, while 1.0 mM sucrose induced activities of three GRNs in the lateral sensillum

of caterpillars, including the weak responses of L-GRN and S-GRN, and the strong

response of the sucrose-best M1-GRN (Fig. 4A). But the sensitivity of the sucrose-best M1-

GRN decreased with the concentration of LA increased in mixtures including 1.0 mM

sucrose (ANOVA, $P < 0.05$). Even the sensitivities of the L-GRN and the M1-GRN were

almost completely suppressed when the concentration of LA in mixtures reached to 200mM

and 300mM (Fig. 4A, Fig. 4B). Finally, when the lateral sensillum was repeated stimulated

by 1.0 mM sucrose in the absence of LA, the activity of the sucrose-best M1-GRN

recovered completely (Figs. 4A, 4B).

We also found that LA suppressed the sensitivities of GNRs in the medial sensillum of *H.*

armigera caterpillars sensitive to tobacco leaf saps. As shown in figure 5, pure tobacco leaf

saps induced the activities of 4 GRNs in the medial sensillum including the strong response

of the M1-GRN, the relatively strong response of the M2-GNR, the weak responses of L-

GNR and S-GNR, but the activities of the M1-GRN and the M2-GRN decreased with the

concentration of LA in mixtures increasing. Even the activities of the M1-GRN, M2-GRN

and L-GRN almost were completely inhibited when LA reached to 300 mM and 400 mM in

mixtures (Figure 5A and 5B). Finally, when the medial sensillum was repeated stimulated

by tobacco leaf saps in the absence of LA, the activity of the GRNs sensitive to tobacco

leaves recovered completely (Figs. 5A, 5B).

Larval development prolonged after exposed to LA

The larval development duration, the larval mortality rate, the puapl development duration,

and the pupal weight of *H. armgiera* originated form neonates reared on artificial diets

containing serial concentrations of LA (6.66 mM, 66.6 mM, 199.8 mM, 333 mM, and 500

mM) were also examined. The results show that dietary LA has significant effects on the

larval developmental duration, the pupal developmental duration and the pupal weights of *H.*

armigera (one-way ANOVA: larval developmental duration, $F_{(5, 473)} = 48.928$, $P < 0.0001$;

pupal developmental duration, $F_{(5, 379)} = 5.384$, $P = 0.02$; pupal weights, $F_{(5, 466)} =$

11.125, $P < 0.0001$) (Fig. 6). At the concentrations of 333 mM and 500 mM of LA, the larval

developmental duration was significantly prolonged compared to the control caterpillars

(Scheffé Test: $P < 0.001$; Fig. 6A). The larval mortalities were not significantly affected by

dietary LA (one-way ANOVA: $F_{(5, 18)} = 0.283$, $P = 0.917$, Fig. 6B). But the pupal duration

and the pupal weight were significantly decreased under 500 mM LA (Scheffé Test followed

ANOVA: pupation duration duration, $P = 0.005$; pupal weights, $P < 0.001$; Fig. 6C and 6D).

The developmental assays demonstrate that the consumption of LA by *H. armigera*

caterpillars could result in the delaying of larval development.

Suppressed activity of digestive enzymes after exposure to LA

After exposures of *H. armigera* neonates to dietary LA until the 5[th] instar stage, the sucrase activity in midguts of the 5[th] instar *H. armgiera* caterpillars were suppressed at a dose-dependent manner (6.66 mM – 500 mM; One-way ANOVA: $F_{(5, 134)} = 26.755$, $P < 0.0001$) (Fig. 7A). The sucrase activities in midguts of caterpillars exposed to 6.66 mM and 66.6 mM dietary LA were similar as those of the control caterpillars (One-way ANOVA, $P = 0.692$), while the sucrase activities of caterpillars exposed to 199.8 mM, 333 mM and 500 mM dietary LA were significantly lower than those of caterpillars exposed to 6.66 mM and 66.6 mM dietary LA and the control caterpillars (one-way ANOVA: $P < 0.01$, all results; Fig. 7A).

As expected, the sucrose concentration in midgut of caterpillars exposed to dietary LA increased at a dose-dependent manner (One-way ANOVA: $F_{(5, 113)} = 39.750$, $P < 0.0001$) (Fig. 7B). The sucrose concentration of caterpillars exposed to 199.8 mM LA was significantly higher than that of the control caterpillars (Scheffé's Tests: $P = 0.023$; Fig. 7B), but was similar to that of caterpillars exposed to 6.66 mM and 66.6 mM LA (Scheffé's Tests: $P = 0.926$; Fig. 7B). With the dietary LA increasing to 333 mM and 500 mM, the sucrose concentration in midguts of caterpillars were significantly higher than that of caterpillars exposed to low concentrations of dietary LA and the control (Scheffé's Tests: $P < 0.0001$, all results; Fig. 7B).

Correspondingly, the glucose levels in midgut of LA-fed caterpillars decreased in a dose-dependent manner from 6.66 mM to 500 mM dietary LA (One-way ANOVA, $F_{(5, 230)} = 17.590$, $P < 0.0001$). Caterpillars exposed to 0, 6.66 mM and 66.6 mM dietary LA were similar at their midgut glucose levels (Scheffé's Tests followed ANOVA: $P = 0.08$; Fig. 7C). But the glucose levels in caterpillars exposed to 199.8 mM dietary LA were significantly lower than that in control caterpillars and caterpillars exposed to 6.66 mM LA (Scheffé's Tests: $P < 0.01$; Fig. 7C). With the concentration of LA in diets increasing to 333 mM and

500 mM, the glucose levels were even lower than that of caterpillars exposed to 66.6 mM

dietary LA (Scheffé's Tests: all $P < 0.05$; Fig. 7C).

Kinetic analysis of sucraseinhibition by LA

We also found that LA inhibited the activity of intestinal sucrase of *H. armigera* caterpillars

in an uncompetitive manner *in vitro* based on the Lineweaver-Burk analyses, displaying

parallel lines of $1/V$ for the different LA concentrations (Fig. 8A). The maximum reaction

velocity (V_{max}) decreased from 7.31 to 5.85 mmol/min/mg protein and the K_m decreased

from 7.24 to 5.15 mM with increasing of the LA concentration from 0 to 9.99 mM. At

concentrations of 6.66 and 9.99 mM, LA reduced sucrase activity 12.24% and 26.81%,

respectively, at V_{max} with the K_i of 30.66 mM/L.

As the positive control, the other alpha-glucosidase inhibitor, acarbose, inhibited sucrase

activities by a fully competitive manner (Fig. 8B). The V_{max} was similar from 8.94 to 8.89

mmol/min/mg protein and the K_m increased from 2.14 to 4.71 mM with increasing of the

acarbose concentration from 0 to 9.99 μM (Fig. 8B). The apparent K_i was calculated as

54.08 μM/L.

Discussion

The present study is the first report that LA could inhibit the feeding behaviors of

herbivorous insects. Tobacco is one of the high value hosts of *H. armigera* (Fitt 1989), but

our result showed tobacco leaves exposed to ≥ 10 mM LA significantly deterred the

feedings of *H. armigera* caterpillars. Under both field and lab conditions, pepper leaves

were not the preferred choice of *H. armgiera* caterpillars (Liu et al. 2009; Ruan and Wu

2001; Tang et al. 2006), whereas our data showed *H. armgiera* preferred control pepper

leaves over leaves exposed to ≥ 10 mM LA. Sucrose and fructose are two phagostimulants

of lots of lepidopteran species (Bernays and Chapman 2000; 2001; Simmonds et al. 1992;

Sun et al. 2021; Vickerman and de Boer 2002), additions of sucrose or fructose on tobacco

or pepper leaves could significantly induce preferences of *H. armigera* caterpillars, but the

preferences could be counteracted after LA mixed to sucrose or fructose in our present

study. These comprehensive behavioral results show that LA is an antifeedant compound

or deterrent substance to caterpillars of the generalist herbivorous *H. armigera*.

The definition of an antifeedant could be: "a peripherally mediated behavior-modifying

substance resulting in feeding deterrence" (Isman 1994; Koul 2005). In detail, antifeedants

could be of pre-ingestive feeding inhibitors that act through gustatory receptors eliciting

rejection of the plant material by changing the sensory code from "acceptable" to

"unacceptable" (Dancewicz et al. 2020; Frazier and Chyb 1995). Our results demonstrated

that the additions of LA to sucrose and tobacco leaf saps resulted in the suppressing of the

sensitivities of peripheral GRNs to these substances at dose-dependent patterns, which is

consistent with the tendency models of LA effects on the feeding behaviors for these

substances, suggesting that the suppressions may contribute to the deterrent effects of LA

on *H. armigera* caterpillars.

LA sensitive GRNs were not found in the peripheral maxillary sensillum of *H. armigra*

caterpillars, indicating the indirect detection of LA in this *Helicoverpa* species. Although

indirect detection of sweeteners have not yet been reported in herbivorous insects, indirect

detection of deterrents by suppressing the activity of sugar GRNs have been found in

Drosophila (Meunier et al. 2003) and the honeybee *Apis mellifera* (de Brito Sanchez et al.

2014). Our study showed the activities of GRN sensitive to phagostimulants and host leaf

saps were suppressed by LA, but the mechanism of indirect detection of LA and the

suppressing coding retains to be identified.

The impacts of LA on behaviors and taste could be further reflected by the prolonged

larval development and the suppressed intestinal sucrase activity of *H. armgiera* caterpillars

after LA ingestion. The depressed sucrase activity in midgut resulted in the accumulations

of sucrose, the decreasing of glucose level and thereby the prolonged development, which

is similar to the effects of prolonged duration in other insect caterpillars after digestions of

toxic substances or deterrents (Glendinning 2002; Sollai et al. 2017; Yadav et al. 2010).

Consequently, our results of behaviors, gustation, development and intestinal physiology

provide a comprehensive understanding of the effects of LA, demonstrating that LA is an

effective antifeedent to the polyphagous herbivorous insect *H. armigera*.

However, our results of dietary LA suppressing the sucrase activity is not consistent with

the effects of LA on the piercing-sucking mouthpart whitefly *B. tabaci* that dietary LA

exposures didn't change the activity of α-glucosidase although LA exposure resulted in a

significant drop in the mean day survival of this species (Hu et al. 2010). We postulate two

possibilities may contribute to the inconsistence of α-glucosidase activity between the two

insect species. Firstly, in our present experiment, we investigated the effects of LA on the

intestinal sucrase activity but not the activities of all α-glucosidases because of the latter

including sucrase, maltase and glucoamylase. In mammals, dietary LA showed no inhibitory

effect on the activities of intestinal maltase and glucoamylase (Halschou-Jensen et al. 2015;

Seri et al. 1996). But the activity of the whole α-glucosidases were investigated in the work

of *B. tabaci* (Hu et al. 2010), then we suppose that the activity of total α-glucosidase in *B.*

tabaci may cover the different changes of activity of different α-glucosidases, such as the

sucrase and the maltase. Secondary, dietary LA may indeed have no obvious effect on the

sucrase activity in *B. tabaci* because of great differences of digestive system between

chewing mouthpart caterpillars and the hemipteran *B. tabaci*. In the work of *B. tabaci*,

dietary LA resulted in a high mortality whereas our work showed the mortality of *H.*

armigera caterpillars was not significantly changed by LA exposure.

LA is one of the components of plant biopolymers such as cereal fiber and pectin in a

variety of crops. But no reports show that crops containing natural LA deter feedings of

herbivorous insects, which may due to the low concentration of LA in plants (Kotake et al.

2016) and the tolerance of low concentration of deterrents in herbivorous insects (Ahn et al.

2011; Sun et al. 2019; Wink and Theile 2002). As one of plant specific primary metabolites,

the metabolic pathway of LA may be of importance to plant pollen development (Kotake et

al. 2016). Then its physiological functions in plants may be quite different from most

reported antifeedants, such as azadirachtin, rotenone, matrine, nicotine, pyrethrins,

servicing as defense compounds (Cheng et al. 2020; Isman 2020; Pavela 2016; Siegwart et

al. 2015; Walia et al. 2017).

As a non-caloric sugar, the physiological and ecological roles of LA in plants is not quite

clear (Kotake et al. 2016), but it is obvious that LA doesn't service as nutrients to animals.

The negative effects of LA on a variety of insects and thereby the phenotypic profiles in

these insects may be different. In *Drosophila*, LA services as a peripheral sweetener, but

LA exposures could be remembered as a negative experience and leaded to a so-called

caloric frustration memory, resulting in a reduced response to LA, which may be involved in

the post-ingestion process of this compound (Musso et al. 2017). Indeed, during the feeding

processes of our work, some *H. armigera* caterpillars vomited after consumption of the LA-

treated plant leaves, indicating LA evoked an emesis response of digestive disruption,

which suggests the post-ingestive feedback process may also mediate the evaluation of

this pentose. We are not clear whether the emesis response of *H. armgiera* caterpillars is

related to the suppressed activity of intestinal sucrase in midgut, but it seems this non-

caloric sugar had serious negative effects on the digestive system of caterpillars.

Together with our findings, our study demonstrated that LA could be an effective feeding

antifeedant towards *H. armgiera*, depending on its inhibiting activities on caterpillars from

behaviors to physiologies. We also investigated whether the antifedant effects of LA could

be observed in other insect species. Indeed, our preliminary experiments suggested LA

also could inhibit the feedings in another *Helicoverpa* species, the tobacco budworm *H.*

assulta, and the eusocialinsect species, the termite *Reticulitermes chinensis* (Isoptera:

Rhinotermitidae), but the antifeedant effect was not obvious in another extreme pest

species, the fall armyworm *Spodoptera frugiperda* (J. E. Smith) (unpublished data). As a

matter of fact, exploring and application of potential antifeedants would be very helpful for

pest control in the current and future IPM strategy. Our work provides substantial evidence

of an unusual botanical sugar which could service as a potential biological antifeedant for

pest control.

Acknowledgments

We thank Dr. Wenge Chen, Ms. Juan Qu, Dr. Huijuan Yang, and Mr. Zhongwei Sun for

kindly helping in the kinetic analysis of enzyme reaction, insect rearing, supplying the

tobacco seedlings, and supplying the pepper seedlings, respectively. The work was

supported by the National Natural Science Foundation of China (Grant No. 31672367 &

31861133019), the Science and Technology Program of Henan Province of China (Grant

No. 202102110072) and the Key Science and Technology Research Project of Henan

Province of China (Grant No. 201300111500).

Author Contributions

QT, YM, XZ, and FY designed the research. LS and ZH performed most of the experiments.

YM, WH and LL performed some of the behavioral and electrophysiological experiments.

YM, LS, ZH analyzed the data. QT, YM, XY and CW wrote the manuscript. All authors read

and approved the final manuscript.

Conflict of Interest Declaration

The authors have declared that no competing interest exists.

References

Ahn SJ, Badenes-Perez FR,Heckel DG (2011) A host-plant specialist, *Helicoverpa assulta*, is more tolerant to capsaicin from *Capsicum annuum* than other noctuid species. J Insect Physiol 57: 1212-1219. https://doi.org/10.1016/j.jinsphys.2011.05.015.

Akhtar Y, Yu Y, Isman MB,Plettner E (2010) Dialkoxybenzene and dialkoxyallylbenzene feeding and oviposition deterrents against the cabbage looper, *Trichoplusia ni*: potential insect behavior control agents. J Agric Food Chem 58: 4983-4991. https://doi.org/10.1021/jf9045123.

Bengtsson S (1990) Isolation and chemical characterization of water soluble arabinoxylans in rye grain. Carbohydr Polym 12: 267-277. https://doi.org/10.1016/0144-8617(90)90068-4.

Bernays EA,Chapman RF (2000) A neurophysiological study of sensitivity to a feeding deterrent in two sister species of *Heliothis* with different diet breadths. J Insect Physiol 46: 905-912. https://doi.org/10.1016/S0022-1910(99)00197-3.

Bernays EA,Chapman RF (2001) Taste cell responses in the polyphagous arctiid, *Grammia geneura*: towards a general pattern for caterpillars. J Insect Physiol 47: 1029-1043. https://doi.org/10.1016/s0022-1910(01)00079-8.

Cheng X, He H, Wang WX, Dong F, Zhang H, Ye J, Tan C, Wu Y, Lv X, Jiang X,Qin X (2020) Semi-synthesis and characterization of some new matrine derivatives as insecticidal agents. Pest Manag Sci 76: 2711-2719. https://doi.org/10.1002/ps.5817.

Dancewicz K, Szumny A, Wawrzenczyk C,Gabrys B (2020) Repellent and antifeedant activities of citral-derived lactones against the peach potato aphid. Int J Mol Sci 21: 8029. https://doi.org/10.3390/ijms21218029.

de Brito Sanchez MG, Lorenzo E, Su S, Liu F,Giurfa M (2014) The tarsal taste of honey bees: behavioral and electrophysiological analyses. Front Behav Neurosci 8. https://doi.org/10.3389/fnbeh.2014.00025.

DuBois GE,Prakash I (2012) Non-caloric sweeteners, sweetness modulators, and sweetener enhancers. Annu Rev Food Sci Technol 3: 353-380. https://doi.org/10.1146/annurev-food-022811-101236.

FAO (2019) New standards to curb the global spread of plant pests and diseases. Web page of the Food and Agriculture Organization of the United Nations. http://www.fao.org/news/story/en/item/1187738/icode/. Accessed 16 November, 2020.

Fitt GP (1989) The ecology of *Heliothis* species in relation to agroecosystems. Annu Rev Entomol 34: 17-52. https://doi.org/10.1146/annurev.en.34.010189.000313.

Frazier JL,Chyb S (1995) Use of feeding inhibitors in insect control. In: Chapman RF,de Boer G (ed) Regulatory Mechanisms in Insect Feeding, Chapman & Hall, New York, NY, USA, pp. 364-381.

Garleb KA, Bourquin LD,Fahey GC (1989) Neutral monosaccharide composition of various fibrous substrates: a comparison of hydrolytic procedures and use of anion-exchange high-performance liquid-chromatography with pulsed amperometric detection of monosaccharides. J Agric Food Chem 37: 1287-1293. https://doi.org/10.1021/jf00089a018.

Glendinning JI (2002) How do herbivorous insects cope with noxious secondary plant compounds in their diet? Entomol Exp Appl 104: 15-25. https://doi.org/10.1046/j.1570-7458.2002.00986.x.

Halschou-Jensen K, Bach Knudsen KE, Nielsen S, Bukhave K,Andersen JR (2015) A mixed diet supplemented with L-arabinose does not alter glycaemic or insulinaemic responses in healthy human subjects. Br J Nutr 113: 82-88. https://doi.org/10.1017/S0007114514003407.

Halschoujensen K, Knudsen KE, Nielsen S, Bukhave K,Andersen JR (2015) A mixed diet supplemented with L-arabinose does not alter glycaemic or insulinaemic responses in healthy human subjects. Br J Nutr 113: 82-88. https://doi.org/10.1017/S0007114514003407.

Hasheminia SM, Sendi JJ, Jahromi KT,Moharramipour S (2013) Effect of milk thistle, *Silybium marianum*, extract on toxicity, development, nutrition, and enzyme activities of the small white butterfly, *Pieris rapae*. J Insect Sci 13: 146. https://doi.org/10.1673/031.013.14601.

Hu JS, Gelman DB, Salvucci ME, Chen YP,Blackburn MB (2010) Insecticidal activity of some
reducing sugars against the sweet potato whitefly, *Bemisia tabaci,* Biotype B. J Insect Sci 10:
203. https://doi.org/10.1673/031.010.20301.
Isman MB (1994) Botanical insecticides and antifeedants: new sources and perspectives. Pestic Res
J 6: 11-19.
Isman MB (2020) Commercial development of plant essential oils and their constituents as active
ingredients in bioinsecticides. Phytochem Rev 19: 235-241. https://doi.org/10.1007/s11101-
019-09653-9.
Jiang J, Ding S, Zhang Y,Wang Y (2010) Effects of artificial diet on the growth & development and
fecundity of *H. armigera*. J Henan Agr Univ 44: 78-82. https://doi.org/10.16445/j.cnki.1000-
2340.2010.01.007.
Jiang X, Xu H, Zheng N, Yin X,Zhang L (2020) A Chemosensory Protein Detects Antifeedant in
Locust (*Locusta migratoria*). Insects 12: 1. https://doi.org/10.3390/insects12010001.
Junhirun P, Pluempanupat W, Yooboon T, Ruttanaphan T, Koul O,Bullangpoti V (2018) The study of
isolated alkane compounds and crude extracts from *Sphagneticola trilobata* (Asterales:
Asteraceae) as a candidate botanical insecticide for lepidopteran larvae. J Econ Entomol 111:
2699-2705. https://doi.org/10.1093/jee/toy246.
Karkanis AC,Athanassiou CG (2021) Natural insecticides from native plants of the Mediterranean
basin and their activity for the control of major insect pests in vegetable crops: shifting from
the past to the future. J Pest Sci 94: 187-202. https://doi.org/10.1007/s10340-020-01275-x.
Kotake T, Yamanashi Y, Imaizumi C,Tsumuraya Y (2016) Metabolism of L-arabinose in plants. J
Plant Res 129: 1-12. https://doi.org/10.1007/s10265-016-0834-z.
Koul O (2005) Insect Antifeedants. CRC Press, Boca Raton, USA.
Koul O, Daniewski WM, Multani JS, Gumulka M,Singh G (2003) Antifeedant effects of the limonoids
from *Entandrophragma candolei* (Meliaceae) on the gram pod borer, *Helicoverpa armigera*
(Lepidoptera: Noctuidae). J Agric Food Chem 51: 7271-7275.
https://doi.org/10.1021/jf0304223.
Liu Z, Gong P, Heckel DG, Wei W, Sun J,Li D (2009) Effects of larval host plants on over-wintering
physiological dynamics and survival of the cotton bollworm, *Helicoverpa armigera* (Hubner)
(Lepidoptera: Noctuidae). J Insect Physiol 55: 1-9.
https://doi.org/10.1016/j.jinsphys.2008.07.017.
Ma Y, Li JJ, Tang QB, Zhang XN, Zhao XC, Yan FM,van Loon JJA (2016) Trans-generational
desensitization and within-generational resensitization of a sucrose-best neuron in the
polyphagous herbivore *Helicoverpa armigera* (Lepidoptera: Noctuidae). Sci Rep 6: 39358.
https://doi.org/10.1038/srep39358.
Mao G, Tian Y, Sun Z, Ou J,Xu H (2019) Bruceine D isolated from *Brucea Javanica* (L.) Merr. as a
systemic feeding deterrent for three major lepidopteran pests. J Agric Food Chem 67: 4232-
4239. https://doi.org/10.1021/acs.jafc.8b06511.
Meunier N, Marion-Poll F, Rospars JP,Tanimura T (2003) Peripheral coding of bitter taste in
Drosophila. J Neurobiol 56: 139-152. https://doi.org/10.1002/neu.10235.
Mitri C, Soustelle L, Framery B, Bockaert J, Parmentier ML,Grau Y (2009) Plant insecticide L-
canavanine repels *Drosophila* via the insect orphan GPCR DmX. PLoS Biol 7: e1000147.
https://doi.org/10.1371/journal.pbio.1000147.
Musso P, Lampin-Saint-Amaux A, Tchenio P,Preat T (2017) Ingestion of artificial sweeteners leads
to caloric frustration memory in *Drosophila*. Nat Commun 8: 1803.
https://doi.org/10.1038/s41467-017-01989-0.
Pavela R (2016) History, presence and perspective of using plant extracts as commercial botanical
insecticides and farm products for protection against insects - a review. Plant Prot Sci 52:
229-241. https://doi.org/10.17221/31/2016-PPS.
Ponsankar A, Sahayaraj K, Senthil-Nathan S, Vasantha-Srinivasan P, Karthi S, Thanigaivel A,
Petchidurai G, Madasamy M,Hunter WB (2020) Toxicity and developmental effect of
cucurbitacin E from *Citrullus colocynthis* L. (Cucurbitales: Cucurbitaceae) against *Spodoptera*

litura Fab. and a non-target earthworm *Eisenia fetida* Savigny. Environ Sci Pollut Res Int 27:
23390-23401. https://doi.org/10.1007/s11356-019-04438-1.

Roessingh P, Hora KH, van Loon JJA,Menken SBJ (1999) Evolution of gustatory sensitivity in
Yponomeuta caterpillars: sensitivity to the stereo-isomers dulcitol and sorbitol is localised in
a single sensory cell. J Comp Physiol A 184: 119-126.
https://doi.org/10.1007/s003590050311.

Ruan YM,Wu KJ (2001) Performances of the cotton bollworm, *Helicoverpa armigera* on different
food plants. Acta Entomol Sin 44: 205-212. https://doi.org/10.16380/j.kcxb.2001.02.013.

Saulnier L, Vigouroux J,Thibault JF (1995) Isolation and partial characterization of feruloylated
oligosaccharides from maize bran. Carbohydr Res 272: 241-253.
https://doi.org/10.1016/0008-6215(95)00053-V.

Seri K, Sanai K, Matsuo N, Kawakubo K, Xue C,Inoue S (1996) L-arabinose selectively inhibits
intestinal sucrase in an uncompetitive manner and suppresses glycemic response after
sucrose ingestion in animals. Metabolism 45: 1368-1374. https://doi.org/10.1016/S0026-
0495(96)90117-1.

Shibanuma K,Houda K (2011) Determination of the transient period of the EIS complex and
investigation of the suppression of blood glucose levels by L-arabinose in healthy adults. Eur
J Nutr 50: 447-453. https://doi.org/10.1007/s00394-010-0154-3.

Siegwart M, Graillot B, Lopez CB, Besse S, Bardin M, Nicot PC,Lopez-Ferber M (2015) Resistance to
bio-insecticides or how to enhance their sustainability: a review. Front Plant Sci 6.
https://doi.org/10.3389/Fpls.2015.00381.

Simmonds MS, Simpson SJ,Blaney WM (1992) Dietary selection Behaviour in *Spodoptera Littoralis*:
the effects of conditioning diet and conditioning period on neural responsiveness and
selection behaviour. J Exp Biol 162: 73-90.

Sollai G, Biolchini M, Solari P,Crnjar R (2017) Chemosensory basis of larval performance of *Papilio*
hospiton on different host plants. J Insect Physiol 99: 47-57.
https://doi.org/10.1016/j.jinsphys.2017.02.007.

Sollai G, Tomassini Barbarossa I, Masala C, Solari P,Crnjar R (2014) Gustatory sensitivity and food
acceptance in two phylogenetically closely related papilionid species: *Papilio hospiton* and
Papilio machaon. PLoS One 9: e100675. https://doi.org/10.1371/journal.pone.0100675.

Sollai G, Tomassini Barbarossa I, Solari P,Crnjar R (2015) Taste discriminating capability to different
bitter compounds by the larval styloconic sensilla in the insect herbivore *Papilio hospiton*
(Gene). J Insect Physiol 74: 45-55. https://doi.org/10.1016/j.jinsphys.2015.02.004.

Sun L, Hou W, Zhang J, Dang Y, Yang Q, Ma Y,Tang Q (2021) Plant metabolites drive different
responses in caterpillars of two closely related *Helicoverpa* species. Front Physiol.
https://doi.org/10.3389/fphys.2021.662978.

Sun Z, Shi Q, Li Q, Wang R, Xu C, Wang H, Ran C, Song Y,Zeng R (2019) Identification of a
cytochrome P450 *CYP6AB60* gene associated with tolerance to multi-plant allelochemicals
from a polyphagous caterpillar tobacco cutworm (*Spodoptera litura*). Pestic Biochem Physiol
154: 60-66. https://doi.org/10.1016/j.pestbp.2018.12.006.

Tang Q, Huang L, Wang C, Zhan H,van Loon JJA (2014) Inheritance of electrophysiological
responses to leaf saps of host- and nonhost plants in two *Helicoverpa* species and their
hybrids. Arch Insect Biochem Physiol 86: 19-32. https://doi.org/10.1002/arch.21154.

Tang Q, Jiang J, Yan Y, van Loon JJA,Wang C (2006) Genetic analysis of larval host-plant
preference in two sibling species of *Helicoverpa*. Entomol Exp Appl 118: 221-228.
https://doi.org/10.1111/j.1570-7458.2006.00387.x.

van Loon JJA (1990) Chemoreception of phenolic acids and flavonoids in larvae of two species of
Pieris. J Comp Physiol A 166: 889-899. https://doi.org/10.1007/bf00187336.

van Opstal AM, Kaal I, van den Berg-Huysmans AA, Hoeksma M, Blonk C, Pijl H, Rombouts S,van
der Grond J (2019) Dietary sugars and non-caloric sweeteners elicit different homeostatic
and hedonic responses in the brain. Nutrition 60: 80-86.
https://doi.org/10.1016/j.nut.2018.09.004.

Vickerman DB,de Boer G (2002) Maintenance of narrow diet breadth in the monarch butterfly
614   caterpillar: response to various plant species and chemicals. Entomol Exp Appl 104: 255-269.
615   https://doi.org/DOI 10.1046/j.1570-7458.2002.01012.x.
616   Wada-Katsumata A, Silverman J,Schal C (2013) Changes in taste neurons support the emergence of
617   an adaptive behavior in cockroaches. Science 340: 972-975.
618   https://doi.org/10.1126/science.1234854.
619   Walia S, Saha S, Tripathi V,Sharma KK (2017) Phytochemical biopesticides: some recent
620   developments. Phytochem Rev 16: 989-1007. https://doi.org/10.1007/s11101-017-9512-6.
621   Wang Y, Ma Y, Zhou D, Gao S, Zhao X, Tang Q, Wang C,van Loon JJA (2017) Higher plasticity in
622   feeding preference of a generalist than a specialist: experiments with two closely related
623   *Helicoverpa* species. Sci Rep 7: 17876. https://doi.org/10.1038/s41598-017-18244-7.
624   Wink M,Theile V (2002) Alkaloid tolerance in *Manduca sexta* and phylogenetically related sphingids
625   (Lepidoptera : Sphingidae). Chemoecology 12: 29-46. https://doi.org/10.1007/s00049-002-
626   8324-2.
627   Wroblewska-Kurdyk A, Dancewicz K, Gliszczynska A,Gabrys B (2020) New insight into the behaviour
628   modifying activity of two natural sesquiterpenoids farnesol and nerolidol towards Myzus
629   persicae (Sulzer) (Homoptera: Aphididae). Bull Entomol Res 110: 249-258.
630   https://doi.org/10.1017/S0007485319000609.
631   Yadav J, Tan CW,Hwang SY (2010) Spatial variation in foliar chemicals within radish (*Raphanus*
632   *sativus*) plants and their effects on performance of *Spodoptera litura*. Environ. Entomol. 39:
633   1990-1996. https://doi.org/10.1603/EN10118.
634   Yamamoto H (2013) Physiological functions and food application of L-arabinose. Oleoscience 13:
635   429-434. https://doi.org/10.5650/oleoscience.13.429.
636   Zalucki MP, Daglish G, Firempong S,Twine PH (1986) The biology and ecology of *Heliothis armigera*
637   (Hübner) and *H. punctigera* Wallengren (Lepidoptera:Noctuidae) in Australia - What do we
638   know? Aust J Zool 34: 779-814. https://doi.org/10.1071/ZO9860779.
639   Zhou D, van Loon JJ,Wang CZ (2010) Experience-based behavioral and chemosensory changes in
640   the generalist insect herbivore Helicoverpa armigera exposed to two deterrent plant
641   chemicals. J Comp Physiol A Neuroethol Sens Neural Behav Physiol 196: 791-799.
642   https://doi.org/10.1007/s00359-010-0558-9.

 Table and Figure legends

Table 1 Settings of the dual-choice feeding assay of *H. armigera* caterpillars for LA-

treated plant leaves.

Note: The tobacco leaves and the pepper leaves were used as the medium leaf discs, respectively.

LA: L-arabinose, Su.: sucrose; Fru.: fructose. Note in experiment II, III, IV and VI, leaves were

treated by mixtures of LA and other sugar.

Table 2 The substances with different concentrations used in the

electrophysiological recordings.

Note: LA: L-arabinose, Su.: sucrose; Tob.: tobacco leaf saps. 2 mM KCl was the adequate

electrolyte solvent.

Figure 1 Effects of LA on the feeding preferences of *H. armigera* caterpillars for

tobacco leaves. Columns represent the mean preference indices ± SEM of the 5th instar

caterpillars for tobacco leaves treated with LA/mixtures or water (control) in dual-choice

assays. **A:** Preferences between leaves treated by LA with different concentrations and

water-treated leaves (control). **B:** Preferences between water-treated leaves (control) and

leaves treated by mixtures of 200 mM sucrose (Su.) plus LA with different concentrations. **C:**

Preferences between 10 mM sucrose-treated leaves and leaves treated by mixtures of 10

mM sucrose plus LA with different concentrations. **D:** Preferences between water-treated

leaves (control) and leaves treated by mixtures of 200 mM fructose (Fru.) plus LA with

different concentrations. A one-sample t-test was used to compare the means of the

preference indices for treatment leaves and for control leaves in each experimental setting

($P < 0.05$).

Figure 2 Effects of LA on the feeding preference of *H. armigera* caterpillars for pepper leaves. Columns represent the mean preference indices ± SEM of the 5th instar caterpillars for pepper leaves treated with LA/mixtures or water (control) in dual-choice assays. **A:** Preference between for leaves treated by LA with different concentrations and for water-treated leaves (control). **B:** Preference between for water-treated pepper leaves (control) and for pepper leaves treated by mixtures of 200 mM sucrose (Su.) plus LA with different concentrations. A one-sample t-test was used to compare the means of the preference indices for treatment leaves and for control leave in each experimental setting ($P < 0.05$).

Figure 3 Morphology and electrophysiological response of gustatory sensilla of *H. armigera* caterpillars. A: Ventral view of the head showing the mouthparts of fifth instar caterpillars. **B:** Close-up view of the medial and the lateral sensillum styloconicum located on a galea. Images were taken by the digital microscope (VHX-600, Keyence, Japan). **C:** Responding curves of the lateral sensillum and the medial sensillum to series concentrations of LA in 2.0 mM KCl.

Figure 4 Representative responding traces and effects of LA on the sensitivity of the sucrose best GRN in the lateral sensillum. A: Representative traces (200 ms) and spikes identified from traces. Different traces were obtained from a single lateral sensillum in the same caterpillar to different stimuli with the stimulating order from up to down. L, M1, and S represent the identified spikes of GRNs sensitive to water, sucrose and 2mM KCl, respectively. **B:** Curves of responding frequency ± SEM of identified GRNs to 1.0 mM sucrose (Su.) and mixtures containing 1.0 mM sucrose and different concentrations of LA.

The Scheffé's post hoc test for one-way ANOVA was used to compare the mean response

frequency of M1 to different stimuli ($P < 0.05$). Columns having no lowercase letters in

common differ significantly ($P < 0.05$).

Figure 5 Representative responding traces and effects of LA on the sensitivities of

GRNs sensitive to tobacco leaf saps in the medial sensillum. A: Representative traces

(200 ms) and spikes identified from traces. Different traces were obtained from a single

medial sensillum in the same caterpillar to different stimuli with the stimulating order from

up to down. L, M1, M2, and S represent the identified spikes of GRNs sensitive to water,

sugars, deterrents and 2mM KCl, respectively. **B:** Curves of responding frequency ± SEM

of the identified GRNs to mixtures containing tobacco saps and different concentrations of

LA. The Scheffé's post hoc test for one-way ANOVA was used to compare the mean

response frequency of M1 and M2 to different stimuli ($P < 0.05$). Columns having no capital

letters or lowercase letters in common differ significantly *($P < 0.05$)*.

Figure 6 Effects of LA on the development of *H. armigera*. Comparison of the larval

developmental duration (**A**) and the larval mortality rate (**B**) after feeding larvae on artificial

diets containing different concentrations of LA from neonate until pupation. Comparison of

pupal developmental duration (**C**) and pupal weight (**D**) of pupae originated from neonatal

caterpillars rearing on artificial diets containing different concentrations of LA until pupation.

The Scheffé's post hoc test for one-way ANOVA was used to compare the means of

developmental parameters of caterpillars reared on different dietary LA ($P < 0.05$). Columns

having no lowercase letters in common differ significantly ($P < 0.05$). C: concentration of LA;

N: the number of caterpillars tested. Groups: repeated numbers of the larval mortality were

investigated.

Figure 7 Effects of LA on the intestinal physiology of *H. armigera* caterpillars.

Comparison of the sucrase activity (**A**), sucrose concentration (**B**), and the glucose

concentrations (**C**) in midgut of the 5th instar caterpillars exposed to artificial diets containing

different concentrations of LA from neonate until pupation. The Scheffé's post hoc test for

one-way ANOVA was used to compare the means of physiological parameters of

caterpillars exposing to different concentrations of LA ($P < 0.05$). Columns having no

lowercase letters in common differ significantly ($P < 0.05$). C: concentration of LA; N: the

number of caterpillars tested.

Figure 8 Kinetic analysis of sucrase inhibition by LA (A) and acarbose (B). Results

represent the Lineweaver-Burk analysis. Supernatants extracted from midgut homogenates

of the 5th instar caterpillar were incubated with increasing concentrations of sucrose in the

presence or absence of inhibitor: (**A**) 6.66 mM LA (□), 9.99 mM LA (□), and the control in

the absence of LA (○); (**B**) 7.53 μM acarbose (□), 11.79 μM acarbose (□), and the control

in the absence of acarbose (○). R^2 was the coefficient of determination which predicts the

goodness of fit for the linear regression model. *V* was the velocity of the reaction.

739   **Table 1**

Medium discs	Tobacco leaf				Pepper leaf	
Experiments	**I**	**II**	**III**	**IV**	**V**	**VI**
Treatments	LA (10^{-4}, 10^{-3}, 10^{-2}, 10^{-1}, 1.0, 10, 67, 133, 200, 333 mM)	200 mM Su. + (0, 100, 300, 400, 500, 600, 800 mM) LA	10 mM Su. + (1.0, 10, 100, 200, 300, 400 mM) LA	200 mM Fru. + (0, 300, 400, 500, 600 mM) LA	LA (10^{-2}, 10^{-1}, 1.0, 10, 100, 200, 300, 400 mM)	200 mM Su. + (0, 100, 300, 500, 600, 800, 1000 mM) LA
Control	pure water	pure water	10 mM sucrose	pure water	pure water	pure water

744   **Table 2**

Experiments	I	II	III	IV
Sensillum	Lateral sensillum	Medial sensillum	Lateral sensillum	Medial sensillum
Stimuli and stimulating sequence	LA (0, 10^{-4}, 10^{-3}, 10^{-2}, 10^{-1}, 1.0, 10 mM)	LA (10^{-4}, 10^{-3}, 10^{-2}, 10^{-1}, 1.0, 10 mM)	2.0 mM KCl, 10 mM LA, 1.0 mM Su., 1.0 mM Su. + (0.1, 1.0, 10, 100, 200, 300 mM) LA, 1.0 mM Su.	Tob., Tob. + (10, 100, 200, 300, 400 mM) LA, Tob.

751   Figures

752   **Figure 1**

773   **Figure 2**

Figure 3

Figure 4

Figure 5

Figure 6

891   **Figure 7**

917 **Figure 8**