TITLE PAGE

Title: PHYLOGENETIC ANALYSIS OF SARS-COV-2 IN THE FIRST MONTHS SINCE ITS

EMERGENCE

Authors: Matías J. PERESON[a,b], Laura MOJSIEJCZUK[a,b], Alfredo P. MARTÍNEZ[c], Diego M.

FLICHMAN[b,d], Gabriel H. GARCIA[a], Federico A. DI LELLO[a,b]

Affiliations:

[a]Universidad de Buenos Aires. Facultad de Farmacia y Bioquímica. Instituto de

Investigaciones en Bacteriología y Virología Molecular (IBaViM). Buenos Aires, Argentina.

[b]Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Ciudad Autónoma

de Buenos Aires, Argentina.

[c]Virology Section, Centro de Educación Médica e Investigaciones Clínicas Norberto Quirno

"CEMIC". Buenos Aires, Argentina.

[d]Instituto de Investigaciones Biomédicas en Retrovirus y Síndrome de Inmunodeficiencia

Adquirida (INBIRS) – Consejo Nacional de Investigaciones Científicas y Técnicas

(CONICET), Universidad de Buenos Aires, Buenos Aires, Argentina.

Corresponding author:

Dr. Federico Alejandro Di Lello, Facultad de Farmacia y Bioquímica, Universidad de Buenos

Aires, Instituto de Investigaciones en Bacteriología y Virología Molecular (IBaViM). Junín

956, 4º piso, (1113), Ciudad Autónoma de Buenos Aires, Argentina.

Phone: +54 11 5287 4472, Fax: +54 11 5287 4662, E-mail: fadilello@ffyb.uba.ar

Running Title: Phylogenetic analysis and evolution of SARS-CoV-2

ABSTRACT

During the first months of SARS-CoV-2 evolution in a new host, contrasting hypotheses

have been proposed about the way the virus has evolved and diversified worldwide. The aim

of this study was to perform a comprehensive evolutionary analysis to describe the human

outbreak and the evolutionary rate of different genomic regions of SARS-CoV-2.

The molecular evolution in nine genomic regions of SARS-CoV-2 was analyzed using three

different approaches: phylogenetic signal assessment, emergence of amino acid

substitutions, and Bayesian evolutionary rate estimation in eight successive fortnights since

the virus emergence.

All observed phylogenetic signals were very low and trees topologies were in agreement

with those signals. However, after four months of evolution, it was possible to identify

regions revealing an incipient viral lineages formation despite the low phylogenetic signal,

since fortnight 3. Finally, the SARS-CoV-2 evolutionary rate for regions nsp3 and S, the ones

presenting greater variability, was estimated to values of 1.37×10^{-3} and 2.19×10^{-3}

substitution/site/year, respectively.

In conclusion, results obtained in this work about the variable diversity of crucial viral regions

and the determination of the evolutionary rate are consequently decisive to understand

essential feature of viral emergence. In turn, findings may allow characterizing for the first

time, the evolutionary rate of S protein that is crucial for vaccines development.

KEYWORDS: SARS-CoV-2, Phylogeny, Evolution, Evolutionary Rate

Introduction

Coronaviruses belong to *Coronaviridae* family and have a single strand of positive-sense RNA genome of 26 to 32 kb in length [1]. They have been identified in different avian hosts as well as in various mammals including bats, mice, dogs, etc. [2,3]. Periodically, new mammalian coronaviruses are identified. In late December 2019, Chinese health authorities identified groups of patients with pneumonia of unknown cause in Wuhan, Hubei Province, China [4]. The pathogen, a new coronavirus called SARS-CoV-2 [5], was identified by local hospitals using a surveillance mechanism for "pneumonia of unknown etiology" [4,6,7]. The pandemic has spread rapidly and, to date, more than 22 million confirmed cases and nearly 750,000 deaths have been reported in just over a six months period [8]. This rapid viral spread raises interesting questions about the way its evolution is driven during the pandemic. From the SARS-CoV-2 genome, 16 non-structural proteins (nsp1-16), 4 structural proteins [spike (S), envelope (E), membrane (M) and nucleoprotein (N)], and other proteins essential to complete the replication cycle are translated [9,10]. The large amount of information currently available allows knowing, as never before, the real-time evolution history of a virus since its interspecies jump [11]. Most studies published to date have characterized the viral genome and evolution by analyzing complete genomes sequences [12,13,14,15]. Despite this, until now, the viral genomic region providing the most accurate information to characterize SARS-CoV-2, could not be established. This lack of information prevent from investigating its molecular evolution and monitoring biological features affecting the development of antiviral and vaccines. Therefore, the aim of this study was to perform a comprehensive viral evolutionary analysis in order to describe the human outbreak and the molecular evolution rate of different genomic regions of SARS-CoV-2.

Materials and Methods

Datasets

In order to generate a dataset representing different geographic regions and time evolution

of the SARS-CoV-2 pandemic from December 2019 to April 2020, data of all the complete

genome sequences available at GISAID (https: //www.gisaid.org /) on April 18, 2020 were

collected. Data inclusion criteria were: a.- complete genomes, b.- high coverage level, and

c.- human hosts only (no other animals, cell culture, or environmental samples). Complete

genomes were aligned using MAFFT against the Wuhan-Hu-1 reference genome

(NC_045512.2, EPI_ISL_402125). The resulting multiple sequence alignment (dataset 1)

was split in nine datasets corresponding to nine coding regions: a.- four structural proteins

[envelope (E), nucleocapsid (N), spike (S), Orf3a], b.- four nonstructural proteins (nsp1,

nsp3, Orf6, and nsp14), and c.- an unknown function protein (Orf8).

More than six thousand SARS-CoV-2 publicly available nucleotide sequences were

downloaded. After data selection according to the inclusion criteria, 1616 SARS-CoV-2

complete genomes were included in dataset 1. Sequences of this dataset 1 came from 55

countries belonging to the five continents as follow: Africa: 39 sequences, Americas: 383

sequences, Asia: 387 sequences, Europe: 686 sequences and Oceania: 121 sequences.

After elimination of sequences with indeterminate or ambiguous positions, the number of

analyzed sequences for each region were: nsp1, 1608; nsp3, 1511; nsp14, 1550; S, 1488;

Orf3a, 1600; E, 1615; Orf6, 1616; Orf8, 1612; and N, 1610. Finally, nucleotide sequences

were grouped by fortnight (FN) according to their collection date. Table 1 summarizes the

number of sequences per fortnight since the beginning of the pandemic up to FN 8. On the

other hand, Dataset 2 was created using only variable sequences of each region analyzed in

Dataset 1. Thus, Dataset 1 was used for the analysis of amino acid substitutions and

Dataset 2 was used for the phylogenetic signal analysis and the Bayesian coalescent trees

construction.

Phylogenetic signal

To determine the phylogenetic signal of each of the nine generated alignments, Likelihood

Mapping analyzes were carried out [16], using the Tree Puzzle v5.3 program [17] and the

Quartet puzzling algorithm. This algorithm allowed analyzing the tree topologies that can be

completely solved from all possible quartets of the n alignment sequences using maximum

likelihood. An alignment with defined tree values greater than 70-80% presents strong

support from the statistical point of view [17]. Identical sequences were also removed with

ElimDupes   (Available   at

https://www.hiv.lanl.gov/content/sequence/elimdupesv2/elimdupes.html) as they increase

computation time and provide no additional information about data phylogeny. The best-fit

evolutionary model to each dataset was selected based on the Bayesian Information

Criterion obtained with the JModelTest v2.1.10 software [18].

Analysis of amino acid substitutions

Entropy-One   (Available   at

https://www.hiv.lanl.gov/content/sequence/ENTROPY/entropy_one.html) was used to

determining in dataset 1 the frequency of amino acids at each position for the nine genomic

regions analyzed and evaluating their permanence in the eight investigated fortnights.

Bayesian coalescence and phylogenetic analysis

To study the relationship between SARS-CoV-2 sequences, nine regions of the virus

genome were investigated by Bayesian analyses. Phylogenetic trees were constructed using

Bayesian inference with MrBayes v3.2.7a [19]. Each gene was analyzed independently with

the same dataset used for the phylogenetic signal analysis so that non-identical sequences

were included in the analysis. Analyses were run for five million generations and sampled

every 5000 generations. Convergence of parameters [effective sample size (ESS) ≥ 200,

with a 10% burn-in] was verified with Tracer v1.7.1 [20]. Phylogenetic trees were visualized

with FigTree v1.4.4.

Evolutionary rate

The estimation of the nucleotide evolutionary rate was made with the Beast v1.10.4 program

package [21]. Analyses were run at the CIPRES Science Gateway server [22]. Three hundred

and twelve sequences without indeterminations corresponding to the nsp3 (5835nt) and S

(3822nt) genes were randomly selected from dataset 1. The sequences represent all the

fortnights and most of the geographical locations sampled until April 17. Temporal calibration

was performed by date of sampling. The appropriate evolutionary model was selected as

described above for phylogenetic signal analysis. The TIM model of nucleotide substitution

was used for nsp3 and, the HKY model of nucleotide substitution for S. The analysis was

carried out under a relaxed (uncorrelated lognormal) molecular clock model suggest by

Duchene & col. [23] and with an exponential demographic, proper for early viral samples from

an outbreak [24]. Independent runs were performed for each dataset and a Markov chain

Monte Carlo (MCMC) with a length of 1.3×10^9 steps, sampling every 1.3×10^6 steps, was set.

The convergence of the "meanRate" parameters [effective sample size (ESS) ≥ 200, burn-in

10%] was verified with Tracer v1.7.1 [20]. Additionally, in order to verify the obtained results,

15 independent replicates of the analysis were performed with the time calibration

information (date of sampling) randomized as described by Rieux & Khatchikian, 2017 [25].

Finally, the obtained parameters for real data and the randomized replicates were compared.

Results

Phylogenetic signal

Using bioinformatics tools, a phylogenetic signal study was carried out in order to identify the

most informative SARS-CoV-2 genomic regions. The likelihood mapping analysis showed

that most genes has very poor phylogenetic signal with high values in central region which

represents the area of unresolved quartets (Figure 1). Accordingly, genes could be

separated into three groups. A group with little or no phylogenetic signal (E, Orf6, Orf8, nsp1,

and nsp14), a second group with low phylogenetic signal (Orf3a and N), and a last group

with relatively more phylogenetic signal (S and nsp3) but still low to be considered a robust

one (unresolved quartets >40%).

Analysis of amino acid substitutions

The analysis of amino acids substitutions by fortnights was useful to study the viral

evolutionary dynamics in the context of the beginning of the pandemic. By analyzing different

time periods amino acid sequences, changes were observed in 5 out of 9 genomic regions

and only in 14 out of the 4975 (0.28%) evaluated residues. In most of the regions, except

nsp1, nsp14, E, and Orf6, 2 to 6 amino acids were selected since FN3 and remain

unchanged until the end of the follow up period (Table 2). Particularly, in Orf8 region, early

selection of two amino acid substitutions (V62L and L84S) was observed from FN2. On the

other hand, in the S region, the D614G substitution started with less than 2% in FN3 and

FN4 and reached 88% in the last fortnight. In a similar way, the Q57H (Orf3a) substitution

went from 6% to 34% while L84S (Orf8) start to be selected in FN2 and reached 6% by FN8.

The R203K and G204R substitutions of the N region was selected in FN4 and increased

their population proportion with values greater than 20% towards the end of the follow up

period. Moreover, selection of a great number of sporadic substitutions remaining in the

population for a short period (1-3 fortnights) was observed in the nine analyzed regions.

Indeed, 333 (6.83%) of the analyzed positions presented at least one substitution throughout

the eight fortnights. Table 3 summarizes the number of variable positions, number of

mutations, and number of sequences with mutations by region.

Bayesian coalescence analysis

In this study, trees were performed by Bayesian analysis instead of by distance, likelihood,

or parsimony methods. Consistently with the phylogenetic signal analysis, trees for nsp1, E,

and Orf6 showed a star-like topology. Nevertheless, different proportions of clades formation

could be observed in trees of Orf8, nsp14, Orf3a, N, S, and nsp3 regions (Figure 2). Finally,

from mentioned regions, nsp3 and S showed a better clade constitution. This analysis

allowed to differentiate regions presenting a diversification process (nsp3, nsp14, Orf3a, S,

Orf8, and N) from those that even after four months showed an incipient one (nsp1, E, and

Orf6). Furthermore, this nucleotide analysis is complemented by the previous study of amino

acid variations in each region. However, it is important to note that due to the low

phylogenetic signal observed for each region, results can only be considered as preliminary.

Evolutionary rate

Nsp3 and S sequences were selected to perform the evolutionary rate analysis since both

regions provided the best phylogenetic information among studied regions. The observed

evolutionary rate for nsp3 protein of SARS-CoV-2 was estimated to be 1.37×10^{-3} (ESS 782)

nucleotide substitutions per site per year (s/s/y) (95% HPD interval 9.16×10^{-4} to 1.91×10^{-3}).

On the other hand, the corresponding figures for S were estimated to be 2.19×10^{-3} (ESS

383) nucleotide s/s/y (95% HPD interval 3.19×10^{-3} to 1.29×10^{-3}). In both genomic regions,

date-randomization analyses showed no overlapping between the 95% HPD substitution-

rate intervals obtained from real data and from date-randomized datasets. This fact suggests

that the original dataset has enough temporal signal to perform analyses with temporal

calibration based on tip-dates (Figure 3).

Discussion

The phylogenetic characterization of an emerging virus is crucial to understand the way the

virus and the pandemic will evolve. Thereby, a detailed study of the SARS CoV-2 genome

allows, on the one hand, to contribute to the knowledge of viral diversity in order to detect

the most suitable regions to be used as antivirals or vaccines targets. On the other hand, the

large amount of information that is continuously generated, is allowing studying the SARS

CoV-2 genome and describing a new viral real time evolution like never before.

In the present study, the molecular evolution and viral lineages of SARS-CoV-2 in nine

genomic regions, during eight successive fortnights, was analyzed using three different

approaches: phylogenetic signal assessment, emergence of amino acid substitutions, and

Bayesian evolutionary rate estimation. In this context, the observed phylogenetic signals of

nine coding regions were very low and the obtained trees were consistent with this finding,

showing star-like topologies in some viral regions (nsp1, E, and Orf6). However, after a four

months evolution period, it was possible to identify regions (nsp3, S, Orf3a, Orf8, and N)

revealing an incipient formation of viral lineages, despite the phylogenetic signal, both at the

nucleotide and amino acid levels from FN3. Based on these findings, the SARS-CoV-2

evolutionary rate was estimated, for the first time, for the two regions showing higher

variability (S and nsp3).

As regards the phylogenetic signal, several simulation studies has proven that for a set of

sequences to be considered robust, the central and lateral areas representing the

unresolved quartets, must not be greater than 40% [16]. In this regard, none of the nine

analyzed regions met this requirement. Three regions (E, nsp1, and Orf6) presented values

of 100% unresolved quartets. Most regions (nsp14, Orf3a, Orf8, and N) reached values

higher than 85%. Only in regions nsp3 and S the number of unresolved quartets dropped to

~ 60%. Thus, despite being a virus with an RNA genome, the short time elapsed since its

emergence, and possibly genetic restrictions have led to a constrained evolution of SARS-

CoV-2 in these months. For this reason, it is expected that trees generated from SARS-CoV-

2 partial sequences in the first months of the pandemic are unreliable for defining clades.

Therefore, they should be analyzed with great caution.

Since Bayesian analysis allows to infer phylogenetic patterns from tree distributions, it

represents a more reliable tool to compare different evolutionary behaviors. Bayesian

analysis helps to obtain a tree topology that is closer to reality in the current conditions of

SARS-CoV-2 pandemic [26]. The phylogenetic analysis for nsp1, E, and Orf6 regions

confirmed the star-like topologies in accordance to a lower diversification of these regions

using the sequences available up to FN8 (Figure 2). Trees generated from nsp14 and Orf8

are at an intermediate point, where the formation of small clusters can be observed. In fact,

a mutation at position 28,144 (Orf8: L84S) has been proposed as a possible marker for viral

classification [27,28]. Finally, trees obtained from regions Orf3a, N, nsp3, and S showed the

best clade formation. Indeed, in the most variable regions nsp3 and S, it can be clearly seen

that sequences are separated into two large groups. Despite the aforementioned for the

nsp3 and S regions, even clusters with very high support values should be taken with

precaution and longer periods should be considered to obtain more accurate phylogeny

data. However, even when data are not the most accurate to study the spread or clade

formation [29, 30], they provide a good representation of the way the virus is evolving.

The analysis of amino acids frequencies allowed identifying different degree of region

conservation throughout the viral genome as a consequence of positive and negative

pressures. In particular, nsp3, S, Orf8, and N showed some substitutions in high

frequencies. This would indicate, as other authors previously report, the frequent circulation

of polymorphisms due to significant positive pressure [13,27,31]. Additionally, since S and N are

among candidates to be used in the formulation of vaccines and antibody treatment, it will be

important to monitor these substitutions in different geographic regions in order to improve

treatment and vaccination efficacy [32,33,34]. In particular, the appearance of the D614G variant

in the third week and its rapid increase until reaching a prevalence of 88% in the eighth week

could reflect an improvement in viral fitness, as several studies reported [35].

Contrarily, in regions nsp1, nsp14, E, and Orf6 no substitutions were selected and lasted

during the first 4 months of the pandemic. This would suggest that these are regions with

constraints to change due to the great negative selection pressure, as it has been recently

reported [13].

In the present study, the evolutionary rate for SARS-CoV-2 genes was estimated by

analyzing a large number of sequences, which were carefully curated and had a good

temporal and spatial structure. Additionally, the most phylogenetically informative regions of

the genome (nsp3 and S) were used for analysis, reinforcing the results confidence.

Previous studies on SARS-CoV-2 have reported similar data ranging from 1.79×10^{-3} to

6.58×10^{-3} s/s/y for the complete genome [6,36]. However, in both articles, small datasets of

complete genomes were used (N=32 and 54, respectively). As studies were performed early

in the outbreak and due to datasets temporal structure, analysis could have led to less

precise estimates of the evolutionary rate [23]. On the other hand, another study from van

Dorp et al. (2020), analyzing 7,666 sequences has obtained different results with a

remarkably low evolutionary rate (6×10^{-4} nucleotide/genome/year) [15]. However, it is

important to consider that van Dorp et al. (2020) estimate the evolutionary rate using the

complete genome, including several highly conserved genomic regions, while in our work,

the estimation was performed with the most variable regions of the genome. Additionally,

tests randomizing the dates of nsp3 and S datasets were carried out; they showed that these

partial genomic regions have enough temporal signal. In this context, our results (1.37×10^{-3}

s/s/y for NSp3 and 2.19×10^{-3} s/s/y for S) are in close agreement with those published for

SARS-CoV genome, which have been estimated between 0.80 to 3.01 x 10^{-3} s/s/y [37-39](The

Chinese SARS Molecular Epidemiology Consortium, 2004, Vega et al. 2004, Zhao et al.

2004). Moreover, our values are in the same order magnitude as other RNA viruses [40].

Even though we should be cautious with these results interpretation, the date-randomization

analysis indicated a robust temporal signal.

In addition, the importance of separately studying the evolutionary rate in S region arises

from the fact that it represents the main target for antiviral agents and vaccines since it

includes the SARS-CoV-2 binding receptor domain (RBD), a crucial structure for the virus to

enter host cells and binding site for neutralizing antibodies [41].

Despite limitations of the evolutionary study of an emerging virus, where the selection

pressures are still low and therefore its variability is also low, this work has a great strength:

it lies on the extremely careful selection of a big sequence dataset to be analyze. First, it was

considered selected sequences to have a good temporal signal and spatial (geographic)

structure. Secondly, much attention was paid to the elimination of sequences with low

coverage and indeterminacies that could generate a noise for the phylogenetic analysis of a

virus that is beginning to evolve in a new host.

The appearance of a new virus means an adaptation challenge. The SARS-CoV-2 overcome

the spill stage and shows a significantly higher spread than SARS-CoV and MERS-CoV,

thus becoming itself the most important pandemic of the century. In this context, the results

obtained in this work about the variable diversity of nine crucial viral regions and the

determination of the evolutionary rate, are consequently decisive to understanding essential

feature of viral emergence. Nevertheless, monitoring SARS-CoV-2 population will be

required to determine the evolutionary course of new mutations as well as to understand the

way they affect viral fitness in human hosts.

Competing interest: On behalf of all authors, the corresponding author states that there is
no conflict of interest.

Funding: None

Declaration of Author Contributions

MJP: Data curation, acquisition of data, analysis and interpretation of data, drafting the
article, final approval of the version to be submitted.

LM: Data curation, acquisition of data, analysis and interpretation of data, revising the article
critically for important intellectual content, final approval of the version to be submitted.

APM: Data curation, Validation, revising the article critically for important intellectual content,
final approval of the version to be submitted.

DMF: Data curation, Validation, drafting the article, final approval of the version to be
submitted.

GG: Data curation, acquisition of data, analysis and interpretation of data, drafting the article,
final approval of the version to be submitted.

FAD: Conception and design of the study, acquisition of data, analysis and interpretation of
data, drafting the article, final approval of the version to be submitted.

Acknowledgements

MJP, LM, DMF, and FAD are members of the National Research Council (CONICET). We
would like to thank to the researchers who generated and shared the sequencing data from
GISAID (https://www.gisaid.org/) and Mrs. Silvina Heisecke from CEMIC-CONICET for
providing language assistance.

REFERENCES

[1] Su S, Wong G, Shi W, et al. Epidemiology, genetic recombination, and pathogenesis of

coronaviruses. *Trends in Microbiol*ogy 2016; 24, 490-502.

https://doi.org/10.1016/j.tim.2016.03.003

[2] Cavanagh D. Coronavirus avian infectious bronchitis virus. *Veterinary Research* 2007*;*

38, 281-297. https://doi.org/10.1051/vetres:2006055

[3] Ismail MM, Tang AY & Saif YM. Pathogenicity of turkey coronavirus in turkeys and

chickens. *Avian Diseases* 2003; 47, 515-522. https://doi.org/10.1637/5917

[4] Zhu N, Zhang D, Wang W, et al. A Novel Coronavirus from Patients with Pneumonia in

China, 2019. *The New England Journal of Medicine* 2020; 382, 727-733.

https://doi.org/10.1056/NEJMoa2001017

[5] Coronaviridae Study Group of the International Committee on Taxonomy of Viruses. The

species *Severe acute respiratory syndrome-related coronavirus*: classifying 2019-nCoV and

naming it SARS-CoV-2. *Nature Microbiology 2020;* 5, 536-544.

https://doi.org/10.1038/s41564-020-0695-z

[6] Li X, Wang W, Zhao X, et al. Transmission dynamics and evolutionary history of 2019-

nCoV. *Journal of Medical Virology* 2020a; 92, 501-511. https://doi.org/10.1002/jmv.25701

[7] Li Q, Guan X, Wu P, et al. Early Transmission Dynamics in Wuhan, China, of Novel

Coronavirus-Infected Pneumonia. *The New England Journal of Medicine* 2020b; 382, 1199-

1207. https://doi.org/10.1056/NEJMoa2001316

[8] World Healt Organization, 2020. Coronavirus disease (COVID-19) Situation Report –

118. Retrieved from: https://www.who.int/docs/default-source/coronaviruse/situation-

reports/20200517-covid-19-sitrep-118.pdf?sfvrsn=21c0dafe_6 (15 August 2020, date last

accessed).

[9] Cui J, Li F & Shi ZL. Origin and evolution of pathogenic coronaviruses. *Nature Reviews*

Microbiology 2019; 17, 181-192. https://doi.org/10.1038/s41579-018-0118-9

[10] Luk HKH, Li X, Fung J, et al. Molecular epidemiology, evolution and phylogeny of SARS

coronavirus. *Infection Genetics and Evolution* 2019; 71, 21-30.

https://doi.org/10.1016/j.meegid.2019.03.001

[11] Zhou P, Yang XL, Wang XG, et al. A pneumonia outbreak associated with a new

coronavirus of probable bat origin. *Nature* 2020; 579, 270-273.

https://doi.org/10.1038/s41586-020-2012-7

[12] Benvenuto D, Giovanetti M, Salemi M, et al. The global spread of 2019-nCoV: a

molecular evolutionary analysis. *Pathogens and Global Health 2020;* 114, 64-67.

https://doi.org/10.1080/20477724.2020.1725339

[13] Cagliani R, Forni D, Clerici M, et al. Computational inference of selection underlying the

evolution of the novel coronavirus, SARS-CoV-2. *Journal of Virology* 2020;

https://doi.org/10.1128/JVI.00411-20

[14] Phan T. Genetic diversity and evolution of SARS-CoV-2. *Infection Genetics and*

Evolution 2020; 81, 104260. https://doi.org/10.1016/j.meegid.2020.104260

[15] van Dorp L, Acman M, Richard D, et al. Emergence of genomic diversity and recurrent

mutations in SARS-CoV-2. *Infection Genetics and Evolution 2020;* 5, 104351.

https://doi.org/10.1016/j.meegid.2020.104351

[16] Strimmer K & von Haeseler A. Likelihood-mapping: A simple method to visualize

phylogenetic content of a sequence alignment. *Proceedings of the National Academy of*

Sciences of the USA 1997; 94, 6815-6819. https://doi.org/10.1073/pnas.94.13.6815

[17] Schmidt HA, Strimmer K, Vingron M, et al. TREE-PUZZLE: Maximum likelihood

phylogenetic analysis using quartets and parallel computing. *Bioinformatics* 2002; 18, 502-

504. https://doi.org/10.1093/bioinformatics/18.3.502

[18] Darriba D, Taboada GL, Doallo R, et al. jModelTest 2: more models, new heuristics and

parallelcomputing. *Nature Methods* 2012; 9, 772. https://doi.org/10.1038/nmeth.2109

[19] Ronquist F, Teslenko M, van der Mark P, et al. MrBayes 3.2: efficient Bayesian

phylogenetic inference and model choice across a large model space. *Systematic Biology*

2012; 61, 539-542. https://doi.org/10.1093/sysbio/sys029

[20] Rambaut A, Drummond AJ, Xie D, et al. Posterior summarization in Bayesian

phylogenetics using Tracer 1.7. *Systematic Biology* 2018; 67, 901-904.

https://doi.org/10.1093/sysbio/syy032

[21] Suchard MA, Lemey P, Baele G, et al. Bayesian phylogenetic and phylodynamic data

integration using BEAST 1.10. *Virus Evolution* 2018; 4, vey016.

https://doi.org/10.1093/ve/vey016

[22] Miller MA, Pfeiffer X, & Schwartz T. Creating the CIPRES Science Gateway for

inference of large phylogenetic trees. *Gateway Computing Environments Workshop* 2010; 1-

8. https://doi.org/10.1109/GCE.2010.5676129

[23] Duchene S, Featherstone L, Haritopoulou-Sinanidou M, et al. Temporal signal and the

phylodynamic threshold of SARS-CoV-2. *bioRxiv* 2020; [Preprint].

https://doi.org/10.1101/2020.05.04.077735

[24] Grassly NC & Fraser C. Mathematical models of infectious disease transmission. *Nat

Rev Microbiol 2008;* 6, 477-487. https://doi.org/10.1038/nrmicro1845

[25] Rieux A & Khatchikian CE. tipdatingbeast: an r package to assist the implementation of

phylogenetic tip-dating tests using beast. *Molecular Ecology Resources* 2017; 17, 608-613.

https://doi.org/10.1111/1755-0998.12603

[26] Drummond AJ, Ho SY, Phillips MJ, et al. Relaxed phylogenetics and dating with

confidence. *PLoS Biology* 2006; 4, e88. https://doi.org/10.1371/journal.pbio.0040088

[27] Tang X, Wu C, Li X, et al. On the origin and continuing evolution of SARS-CoV-2.

National Science Review 2020; 0, 1-12. https://doi.org/10.1093/nsr/nwaa036

[28] Yin C. Genotyping coronavirus SARS-CoV-2: methods and implications. *Genomics*

2020; 30318-30319. https://doi.org/10.1016/j.ygeno.2020.04.016

[29] Mavian C, Marini S, Prosperi M, et al. A snapshot of SARS-CoV-2 genome availability

up to 30th March, 2020 and its implications. JMIR Public Health Surveill 2020; 6, e19170

https://doi.org/10.2196/19170

[30] Sánchez-Pacheco SJ, Kong S, Pulido-Santacruz P, et al. Median-joining network

analysis of SARS-CoV-2 genomes is neither phylogenetic nor evolutionary. *Proceedings of*

the National Academy of Sciences of the USA 2020; 117, 9241–9243.

https://doi.org/10.1073/pnas.2007062117

[31] Issa E, Merhi G, Panossian B, et al. S.SARS-CoV-2 and ORF3a: Nonsynonymous

Mutations, Functional Domains, and Viral Pathogenesis. *mSystems* 2020 [Preprint].

https://doi.org/10.1128/mSystems.00266-20

[32] Ahmed SF, Quadeer AA & McKay MR. Preliminary Identification of Potential Vaccine

Targets for the COVID-19 Coronavirus (SARS-CoV-2) Based on SARS-CoV Immunological

Studies. *Viruses 2020;* 12, 254. https://doi.org/10.3390/v12030254

[33] Callaway E. The race for coronavirus vaccines: a graphical guide. *Nature* 2020*;* 580,

576-577. https://doi.org/10.1038/d41586-020-01221-y

[34] Koyama T, Weeraratne D, Snowdon JL, et al. Emergence of Drift Variants That May

Affect COVID-19 Vaccine Development and Antibody Treatment. *Pathogens* 2020; 9, 324.

https://doi.org/10.20944/preprints202004.0024.v1

[35] Li Q, Wu J, Nie J, et al. The Impact of Mutations in SARS-CoV-2 Spike on Viral

Infectivity and Antigenicity. *Cell* 2020; S0092-8674(20)30877-1. Advance online publication.

https://doi.org/10.1016/j.cell.2020.07.012

[36] Giovanetti M, Benvenuto D, Angeletti S, et al. The first two cases of 2019-nCoV in Italy:

Where they come from? *Journal of Medical Virology* 2020; 92, 518-521.

https://doi.org/10.1002/jmv.25699

[37] The Chinese SARS Molecular Epidemiology Consortium. Molecular Evolution of the

SARS Coronavirus During the Course of the SARS Epidemic in China. *Science* 2004; 303,

1666-1669. https://doi.org/10.1126/science.1092002

[38] Vega VB, Ruan Y, Liu J, et al. Mutational dynamics of the SARS coronavirus in cell

culture and human populations isolated in 2003. *BMC Infectious Diseases* 2004, 4, 32.

https://doi.org/10.1186/1471-2334-4-32

[39] Zhao Z, Li H, Wu X, et al. Moderate mutation rate in the SARS coronavirus genome and

its implications. *BMC Evolutionary Biology* 2004; 4, 21. https://doi.org/10.1186/1471-2148-4-

21

[40] Sanjuán R. From molecular genetics to phylodynamics: evolutionary relevance of

mutation rates across viruses. *PLoS Pathogens* 2012; 8, e1002685. https://doi.org/

10.1371/journal.ppat.1002685

[41] Ju B, Zhang Q, Ge J, et al. Human neutralizing antibodies elicited by SARS-CoV-2

infection. *Nature* 2020; 115–119. https://doi.org/10.1038/s41586-020-2380-z

Table 1. Number of SARS-CoV-2 sequences by fortnight (Temporal structure)

Fortnight	Date	Median of analyzed sequences (Q1-Q3)
FN1	12/24/2019 to 12/31/2019	15
FN2	01/01/2020 to 01/15/2020	19
FN3	01/16/2020 to 01/31/2020	145 (136-145.5)
FN4	02/01/2020 to 02/15/2020	119 (113-120)
FN5	02/16/2020 to 03/02/2020	258 (247-259)
FN6	03/03/2020 to 03/17/2020	403 (390-406)
FN7	03/18/2020 to 04/01/2020	447 (416-450)
FN8	04/02/2020 to 04/17/2020	199 (197-201)
TOTAL		**1488 to 1616**

FN: Fortnight; Q1=quartile 1, Q3=quartile 3. The total number of sequences is variable depending on the analyzed region (nsp1, 1608; nsp3, 1511; nsp14, 1550; S, 1488; Orf3a, 1600; E, 1615; Orf6, 1616; Orf8, 1612; and N, 1610)

450   **Table 2.** Amino acids selected by region and fortnight. The number indicates the amino

451   acids location in its protein.

Region	Amino acid substitution	Amino acid percentage by FN							
		FN1	FN2	FN3	FN4	FN5	FN6	FN7	FN8
nsp3	A58T	0	0	0	1.0	6.0	3.0	3.0	2.5
	P135L	0	0	0.8	0	0	1.5	0.5	2.5
S	D614G	0	0	1.5	1.8	37.0	64.0	75.0	88.0
Orf3a	Q75H	0	0	0	0	6.0	22.0	23.0	34.0
	G196V	0	0	0	0	0.8	4.0	0.9	0.5
	G251V	0	0	8.0	24.0	8.0	9.0	10.0	3.0
Orf8	V62L	0	5.0	1.0	3.3	0.0	1.5	1.3	3.0
	L84S	0	42.0	37.0	21.0	21.0	18.0	7.0	6.0
N	P13L	0	0	0	0	1.0	1.0	2.5	0.5
	S197L	0	0	0	0	1.1	5.0	0.9	0.5
	S202N	0	0	3.5	4.2	0	0.5	2.2	2.5
	R203K	0	0	0	0	17.0	19.0	24.0	23.0
	G204R	0	0	0	0	17.0	19.0	24.0	23.0
	I292T	0	0	0	0	2.0	0.2	0.2	0.5

452   Only regions where amino acid change was selected and remained until the last analyzed fortnight
453   are shown. FN: Fortnight; aa: amino acid

456   **Table 3.** Number of variable positions, number of mutations, and number of sequences with

457   mutation by region

Region	Nº of variable aa positions (%)	Nº of aa substitutions	Nº of sequences with aa substitutions (%)
nsp1 (180aa)	3 (1.7)	37	37 (2.4)
nsp3 (1945aa)	158 (8.1)	322	294 (19.3)
nsp14 (527aa)	6 (1.4)	83	83 (5.5)
S (1273aa)	76 (5.9)	1013	904 (59.4)
Orf3a (275aa)	11 (4)	491	468 (30.7)
E (75aa)	5 (6.7)	6	6 (0.4)
Orf6 (60aa)	7 (11.6)	9	9 (0.6)
Orf8 (121aa)	14 (11.6)	312	288 (18.9)
N (419aa)	53 (12.6)	760	470 (30.9)
Total (4875aa)	333 (6.8)	3033	-

458   aa: amino acid

FIGURE LEGENDS

Figure 1

Phylogenetic signal for SARS-CoV-2 datasets. Presence of phylogenetic signal was

evaluated by likelihood mapping, unresolved quartets (center) and partly resolved quartets

(edges) for genomes available on April 17 for the nine analyzed regions: nsp1 (29

sequences), nsp3 (225 sequences), nsp14 (65 sequences), S (183 sequences), Orf3a (74

sequences), E (11 sequences), Orf6 (12 sequences), Orf8 (23 sequences), and N (113

sequences). Presence of strong phylogenetic signal (<40% unresolved quartets) was not

reached for any region.

Figure 2

Bayesian trees of 29 sequences of nsp1 (540nt), 225 sequences of nsp3 (5835nt), 65

sequences of nsp14 (1581nt), 183 sequences of S (3822nt), 74 sequences of Orf3a (828nt),

11 sequences of E (228nt), 12 sequences of Orf6 (186nt), 23 sequences of Orf8 (366nt),

and113 sequences of N (1260nt). Scale bar represents substitutions per site.

Figure 3

Comparison of the evolutionary rates estimated using BEAST for the original dataset and the

date-randomized datasets (312 sequences). This analysis was performed for regions nsp3

(5835nt) and S (3822nt). s.s.y = substitutions/site/year.

Figure 1

Figure 2

Figure 3