Crosstalk between chloroplast protein import and the SUMO system revealed through genetic and molecular investigation

Samuel James Watson[1†], Na Li[1], Yiting Ye[2], Feijie Wu[2,3], Qihua Ling[1,2,3‡], R. Paul Jarvis[1,3]*

[1]Department of Plant Sciences, University of Oxford, Oxford OX1 3RB, UK.

[2]National Key Laboratory of Plant Molecular Genetics, CAS Centre for Excellence in Molecular Plant Sciences, Institute of Plant Physiology and Ecology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai, China.

[3]Department of Biology, University of Leicester, Leicester LE1 7RH, UK.

[†]Present address: Cambridge Institute for Therapeutic Immunology and Infectious Disease, Jeffrey Cheah Biomedical Centre, University of Cambridge, Cambridge CB2 0AW, UK.

*For correspondence: paul.jarvis@plants.ox.ac.uk

Abstract

The chloroplast proteome contains thousands of different proteins that are encoded by the nuclear genome. These proteins are imported into the chloroplast via the action of the TOC translocase and associated downstream systems. Our recent work has revealed that the stability of the TOC complex is dynamically regulated by the ubiquitin-dependent chloroplast-associated protein degradation (CHLORAD) pathway. Here, we demonstrate that the stability of the TOC complex is also regulated by the SUMO system. *Arabidopsis* mutants representing almost the entire SUMO conjugation pathway can partially suppress the phenotype of *ppi1*, a pale yellow mutant lacking the Toc33 protein. This suppression is linked to the increased stability of TOC proteins and improvements in chloroplast development. In addition, we demonstrate using molecular and biochemical experiments that the SUMO system directly targets TOC proteins. Thus, we have identified a regulatory link between the SUMO system and chloroplast protein import.

Introduction

The chloroplast is a membrane-bound organelle that houses photosynthesis in all green plants
(Jarvis and Lopez-Juez, 2013). Chloroplasts have an unusual evolutionary history – they are the
integrated descendants of a cyanobacterial ancestor that entered the eukaryotic lineage via
endosymbiosis. Although chloroplasts retain small genomes, almost all of the proteins required for
chloroplast development and function are now encoded by the central, nuclear genome (Jarvis,
2008). These proteins must be imported into the organelle after synthesis in the cytosol, and this
import is mediated by the coordinate action of the TOC and TIC complexes (the translocons at the
outer and inner envelope membranes of chloroplasts) (Jarvis, 2008).

The TOC complex contains three major components: the Omp85 (outer membrane protein, 85 kDa)-
related protein, Toc75, which serves as a membrane channel (Schnell et al., 1994; Tranel et al.,
1995), and two GTPase-domain receptor proteins, Toc33 and Toc159 (Hirsch et al., 1994; Kessler et
al., 1994; Perry and Keegstra, 1994; Jarvis et al., 1998; Jarvis, 2008). Toc33 and Toc159 project into
the cytosol and bind incoming preproteins.

The key components of the TOC complex were identified more than two decades ago (Hirsch et al.,
1994; Kessler et al., 1994; Schnell et al., 1994; Tranel et al., 1995; Jarvis, 2008). However, the
regulation of the activity and stability of the complex was, until recently, poorly understood. Major
insights came from a forward genetic screen for suppressors of the pale yellow Toc33 mutant, *ppi1*
(Ling et al., 2012). In this screen, SP1 (SUPPRESSOR OF PPI1 LOCUS 1), a novel RING-type E3 ubiquitin
ligase, was identified. A series of *sp1* mutations were shown to partially suppress the phenotypic
defects of *ppi1* with respect to chlorosis, chloroplast development, and chloroplast protein import.
In addition, SP1 function was shown to promote plastid interconversion events (for example, the
development of the chloroplast from its precursor organelle, the etioplast). Later work
demonstrated that SP1 function is also important for abiotic stress tolerance, by enabling
optimisation of the organellar proteome via protein import regulation (Ling and Jarvis, 2015). Thus,
through SP1, the ubiquitin-proteasome system promotes TOC complex degradation and
reconfiguration in response to developmental and/or environmental stimuli.

Ubiquitinated TOC proteins are extracted from the chloroplast outer envelope membrane and
degraded in the cytosol. Recent work identified two proteins that physically associate with SP1 and
promote the membrane extraction of TOC proteins (Ling et al., 2019). These are SP2, an Omp85-type
β-barrel channel protein that was identified in the same genetic screen as SP1, and Cdc48, a well-
characterised cytosolic AAA+ chaperone ATPase that provides the motive force for the extraction of
proteins from the chloroplast outer envelope. The three proteins – SP1, SP2 and Cdc48 – together
define a new pathway for the ubiquitination, membrane extraction, and degradation of chloroplast
outer envelope proteins, which has been named chloroplast-associated protein degradation, or
CHLORAD. In addition to CHLORAD, there exist cytosolic ubiquitin-dependent systems that also
contribute to chloroplast biogenesis, by regulating the levels of unimported preproteins (Lee et al.,
2009; Grimmer et al., 2020), and by controlling the stability of the Toc159 receptor prior to its
integration into the outer envelope membrane (Shanmugabalaji et al., 2018).

The discovery of SP1 and the CHLORAD pathway demonstrated that the TOC complex is not static
but, instead, can be rapidly ubiquitinated and degraded in response to developmental and
environmental stimuli. To complement this work, we decided to explore whether the TOC complex is
also regulated by the SUMO system. This work was motivated by the results of a high-throughput
screen for SUMO substrates in *Arabidopsis* (Elrouby and Coupland, 2010). This screen suggested that
Toc159, a key component of the TOC complex, is a SUMO substrate. SUMOylation is intricately

involved in plant development and stress adaptation, and so we were interested to determine
whether the TOC complex is targeted by the SUMO system, and whether any such SUMOylation is
functionally important. As crosstalk between the SUMO system and the ubiquitin-proteasome
system is common, we reasoned that answering these questions might provide insights into the
regulation of SP1 and the CHLORAD pathway.

To explore the relationship between chloroplast protein import and the SUMO system, we carried
out a comprehensive series of genetic, molecular and biochemical experiments. Mutants
representing most components of the *Arabidopsis* SUMO pathway were found to partially suppress
the phenotype of the chlorotic Toc33 null mutant, *ppi1*, with respect to leaf chlorophyll
accumulation, chloroplast development, and TOC protein abundance. Conversely, overexpression of
either *SUMO1* or *SUMO3* enhanced the severity of the *ppi1* phenotype. Moreover, the E2 SUMO
conjugating enzyme, SCE1, was found to physically interact with the TOC complex in bimolecular
fluorescence complementation experiments; and TOC proteins were seen to physically associate
with SUMO proteins in immunoprecipitation assays. In combination, our data conclusively
demonstrate significant crosstalk between the SUMO system and chloroplast protein import, and
emphasise the complexity of the regulation of the TOC translocase.

Results

89 The E2 SUMO conjugating enzyme mutant *sce1-4*, and the E3 SUMO ligase mutants *siz1-4* and *siz1-*
90 *5*, partially suppress the phenotype of Toc33 mutant, *ppi1*

Two key components of the CHLORAD pathway, SP1 and SP2, were identified in a forward genetic
screen for suppressors of *ppi1,* an *Arabidopsis* Toc33 null mutant (Ling et al., 2012; Ling et al., 2019).
Both *sp1* and *sp2* mutants can partially suppress the *ppi1* phenotype with respect to chlorophyll
accumulation, chloroplast development, and TOC protein abundance. To investigate whether the
TOC complex is targeted by the SUMO system, we obtained several *Arabidopsis* SUMO system
mutants, crossed them with *ppi1*, and carefully examined the phenotypes of the resulting double
mutants. This reverse genetic approach was possible because the basic architecture of the
Arabidopsis SUMO system is remarkably simple. In the SUMO pathway, the ubiquitin-like SUMO
modifier protein is conjugated to substates by the coordinated action of E1 activating enzymes, E2
conjugating enzymes, and E3 SUMO ligases. Although thousands of proteins are SUMOylated in
Arabidopsis, there is just one known E1 SUMO activating enzyme, one known E2 SUMO conjugating
enzyme, and only two known E3 SUMO ligases of canonical function (Saracco et al., 2007; Ishida et
al., 2009).

First, we analysed *sce1-4*, a weak mutant allele of the sole E2 SUMO conjugating enzyme gene in
Arabidopsis, which is an essential gene (Saracco et al., 2007). The *sce1-4* mutant shows a moderate
reduction in the expression of SCE1 and in global levels of SUMOylation, but it displays no obvious
visible phenotypic defects under steady-state conditions (Saracco et al., 2007). The *ppi1 sce1-4*
double mutant was phenotypically characterised, and, intriguingly, it appeared greener than the
ppi1 single mutant (Figure 1A; Figure 1 – Supplement 1A). This was linked to a moderate increase in
leaf chlorophyll concentration (Figure 1B; Figure 1 – Supplement 1B). Next, we asked whether the
phenotypic suppression observed in *ppi1 sce1-4* was linked to changes in the development of
chloroplasts. The chloroplasts of *ppi1 sce1-4* were visualised via transmission electron microscopy.
Interestingly, the chloroplasts of the *ppi1 sce1-4* double mutant appeared larger and better
developed than those of the *ppi1* control (Figure 1C). The transmission electron micrographs were
quantitatively analysed, and the *ppi1 sce1-4* chloroplasts were indeed found to be significantly larger

than those of *ppi1* (Figure 1D), with larger, more interconnected thylakoidal granal stacks (Figures 1E
and 1F).

SUMO conjugation is usually dependent on the action of E3 SUMO ligases. In *Arabidopsis*, the best
characterised E3 SUMO ligase is SIZ1 (Kurepa et al., 2003; Miura et al., 2005; Saracco et al., 2007).
SIZ1 is not essential, but null mutants display severely dwarfed phenotypes. In order to include SIZ1
in our genetic analysis, we obtained two new T-DNA insertion alleles, and named them *siz1-4* and
siz1-5. While both mutants were visibly similar to the published mutants (Miura et al., 2005; Liu et
al., 2019), *siz1-4* showed a milder phenotype with only moderate growth retardation when grown to
maturity. We mapped the integration sites of the T-DNA insertions in these two mutants (Figure 1 –
Supplement 2A), and showed that both display a strong reduction in *SIZ1* transcript by RT-PCR
analysis (Figure 1 – Supplement 2B). In addition, both mutants displayed defects in global
SUMOylation in response to heat shock, similar to the published alleles (Figure 1 – Supplement 3).
The two new *siz1* mutants were crossed with *ppi1* and the resulting double mutants were
phenotypically characterised. Both the *ppi1 siz1-4* and the *ppi1 siz1-5* double mutants appeared
greener than the *ppi1* control (Figure 1G; Figure 1 – Supplement 1C). In addition, the double mutants
showed dramatic increases in leaf chlorophyll concentration relative to *ppi1* (Figure 1H; Figure 1 –
Supplement 1D). Next, we asked whether the phenotypic suppression observed in *ppi1 siz1-4* and
ppi1 siz1-5 was linked to changes in the abundance of TOC proteins. To this end, protein samples
were taken from the two double mutants and relevant control plants and resolved via
immunoblotting. Both double mutants displayed clear increases in the abundance of Toc159 and
Toc75, two core components of the TOC complex, relative to *ppi1* (Figures 1I, 1J, and 1K).

The suppression effects mediated by the SUMO system mutants are specific

As discussed in the previous section, the SUMO system is encoded by a remarkably small number of
genes in *Arabidopsis*. As a consequence, SUMO system mutants have highly pleiotropic molecular
and physiological phenotypes. We therefore asked whether the partial suppression of *ppi1* by SUMO
system mutants was specific to the *ppi1* background. We crossed *sce1-4* with *tic40-4* and *hsp93-V-1*,
two TIC-complex-associated mutants. These mutants are chlorotic, due to defects in protein import
across the chloroplast inner membrane, and in this respect are highly similar to *ppi1* (Kovacheva et
al., 2005). Significantly, the resulting double mutants, *tic40-4 sce1-4* and *hsp93-V-I sce1-4*, were
indistinguishable from *tic40-4* and *hsp93-V-1*, their respective single mutant controls (Figures 2A and
2C). Moreover, the double mutants did not display changes in leaf chlorophyll accumulation relative
to the single mutant controls (Figures 2B and 2D). We therefore concluded that the suppression
effects observed in *ppi1 sce1* plants were background-specific and associated with the TOC complex.

Next, we asked whether the *sce1-4* and *siz1-4* single mutants display an increase in chlorophyll
concentration even in the wild-type background. However, neither mutant appeared greener than
wild-type plants (Figures 2E and 2G; Figure 1 – Supplement 1A and 1C) or displayed an increase in
leaf chlorophyll concentration (Figures 2F and 2H; Figure 1 – Supplement 1B and 1D). We therefore
concluded that the suppression effects mediated by the SUMO mutants were synthetic phenotypes
specific to the *ppi1* background.

BiFC analysis reveals that SCE1 physically interacts with TOC proteins

Our reverse genetic experiments revealed a genetic interaction between the E2 SUMO conjugating
enzyme, SCE1, and protein import across the chloroplast outer membrane. To determine whether
SCE1 directly interacts with the TOC complex, we carried out bimolecular fluorescence
complementation (BiFC) experiments in *Arabidopsis* protoplasts. The *SCE1* coding sequence was

cloned into a vector that C-terminally appends the C-terminal half of YFP (cYFP) to its insert. This
construct was co-expressed with various other constructs encoding TOC proteins bearing the
complementary, N-terminal moiety of the YFP protein (nYFP), appended N-terminally; or with a
negative control construct encoding ΔOEP7 bearing the nYFP moiety appended C-terminally. In this
system, protein-protein interactions are inferred via the detection of a YFP signal, caused by the
nYFP and cYFP fragments coming together to reconstitute a functional YFP protein.

Strikingly, SCE1-nYFP was found to physically associate with all tested TOC proteins – nYFP-Toc159,
nYFP-Toc132, nYFP-Toc34 and nYFP-Toc33 (Figure 3). Moreover, these interactions were
concentrated at the periphery of the chloroplasts, placing them in an appropriate subcellular context
for the *in situ* regulation of the chloroplast protein import machinery. Conversely, SCE1-cYFP was not
found to physically associate with the negative control protein ΔOEP7-nYFP. The ΔOEP7 protein
comprises the transmembrane domain of plastid protein OEP7, which is sufficient to efficiently
target the full-length YFP protein to the chloroplast outer membrane (Lee et al., 2001); thus cYFP-
ΔOEP7 serves as a location-specific negative control.

174   Manipulating the expression of three SUMO isoforms alters the phenotypic severity of *ppi1*

To further explore the genetic interaction between chloroplast protein import and the SUMO
system, we crossed *ppi1* with several SUMO protein mutants. There are three major SUMO isoforms
in *Arabidopsis* – SUMO1, SUMO2, and SUMO3. The *SUMO1* and *SUMO2* genes are expressed at a
relatively high level throughout the plant and are largely functionally redundant (Saracco et al.,
2007; van den Burg et al., 2010). In addition, they are highly similar to each other in terms of amino
acid sequence (Saracco et al., 2007). In contrast, at steady state, *SUMO3* is expressed at a relatively
low level throughout the plant, while the SUMO3 amino acid sequence is significantly divergent with
respect to the other two SUMO isoforms (van den Burg et al., 2010).

First, we analysed *SUMO1* and *SUMO2*. We obtained *sum1-1* and *sum2-1*, two previously
characterised null mutants (Saracco et al., 2007), and crossed them with *ppi1*. To account for the
functional redundancy between these two genes, we also sought a *ppi1 sum1-1 sum2-1* triple
mutant. However, as *SUMO1* and *SUMO2* are collectively essential, *ppi1 sum1-1 sum2-1* plants that
were homozygous with respect to *ppi1* and *sum2-1*, but heterozygous with respect to the *sum1-1*
mutation, were selected from a segregating population. The double and triple mutants were
phenotypically characterised, and all three appeared larger and greener than the *ppi1* control plants
(Figure 4A). Moreover, the double and triple mutants showed corresponding increases in leaf
chlorophyll concentration, with the triple mutant showing a larger increase than the double mutants
(Figure 4B). These were synthetic effects, as the *sum1-1*, *sum2-1*, and *sum1-1 sum2-1* single and
double mutants did not appear greener than wild-type plants, or show increases in chlorophyll
accumulation (Figure 4 – Supplement 1). We therefore concluded that the *sum1-1* and *sum2-1*
mutants can additively suppress the phenotype of *ppi1*.

To complement the preceding experiment, we generated transgenic plants overexpressing *SUMO1*
in the *ppi1* background. The *SUMO1* coding sequence was cloned into a vector carrying a strong,
constitutive promotor (cauliflower mosaic virus 35S) upstream of the cloning site. The resulting
construct was stably introduced into the *ppi1* background via *Agrobacterium*-mediated
transformation. Two lines carrying a single, homozygous transgene insert were identified and taken
forward for analysis. The overexpression of *SUMO1* was confirmed in both lines by RT-PCR (Figure 4
– Supplement 2A). Significantly, both lines displayed an accentuation of the *ppi1* phenotype: the
plants were significantly smaller and paler than the *ppi1* control plants (Figure 4C), and showed
decreases in leaf chlorophyll concentration (Figure 4D).

Next, we turned our attention to *SUMO3*. We obtained *sum3-1*, a previously characterised null
mutant (van den Burg et al., 2010), and crossed it with *ppi1*. The resulting double mutant was
phenotypically characterised, but it did not appear obviously different from the *ppi1* control (Figure
4E). Correspondingly, it did not display any clear increase in leaf chlorophyll concentration relative to
ppi1 (Figure 4F). To complement this experiment, we generated transgenic plants overexpressing
SUMO3 in the *ppi1* background, using the approach described above, and a line carrying a single,
homozygous insert was identified and taken forward for analysis. The overexpression of *SUMO3* was
confirmed by RT-PCR (Figure 4 – Supplement 2B). Interestingly, the transgenic plants showed a
striking increase in the severity of the *ppi1* phenotype: the plants were severely dwarfed and paler
than the *ppi1* control (Figure 4G), and displayed a significant decrease in leaf chlorophyll
accumulation (Figure 4H). These findings are particularly noteworthy when considered alongside a
previous report which explored the consequences of overexpressing *SUMO3* in wild-type plants (van
den Burg et al., 2010). In that study, *SUMO3* overexpression was not found to alter the appearance
of the transgenic plants, which implies a degree of specificity in the phenotypic accentuation
observed here.

Biochemical analysis reveals SUMOylation of TOC proteins *in vivo*

The genetic and molecular experiments described thus far strongly suggested that TOC proteins are
SUMOylated. However, to our knowledge, conclusive evidence that chloroplast-resident proteins are
SUMOylated is currently lacking. To investigate whether chloroplast proteins may be SUMOylated,
we isolated chloroplasts from seedlings by cell fractionation and analysed them by anti-SUMO
immunoblotting. For this analysis, we employed a proven commercial antibody against SUMO1,
which is one of the most abundant SUMO isoforms in *Arabidopsis* making it more tractable for
analysis, and which furthermore is known to accumulate in response to heat and other stresses
(Kurepa et al., 2003; van den Burg et al., 2010). To enhance detection of SUMOylated proteins in our
samples, we subjected some of the seedlings to heat shock before chloroplast isolation and/or
treatment with 10 mM N-ethylmaleimide (NEM) during chloroplast isolation. NEM is a potent
inhibitor of SUMO-specific proteases (Hilgarth and Sarge, 2005). Importantly, we detected protein
SUMOylation in the isolated chloroplast samples, and this SUMOylation was increased by NEM
treatment (Figure 5 – Supplement 1).

Next, we carried out a series of biochemical experiments to determine whether TOC proteins are
SUMOylated. In the first of these, the *SCE1* coding sequence was cloned into a vector that appends a
C-terminal YFP tag (Karimi et al., 2002). We confirmed that the resulting SCE1-YFP construct delivers
good expression and the expected nucleocytoplasmic fluorescence pattern when transiently
expressed in protoplasts (Figure 5 – Supplement 2A). Then, we transfected a large volume of
protoplasts with the SCE1-YFP construct (or with a YFP-HA negative control construct), and
performed immunoprecipitation (IP) using YFP-Trap magnetic beads. The samples were analysed by
immunoblotting. The YFP-HA and SCE1-YFP fusion proteins both showed robust expression and
strong recovery in the IP elutions (Figure 5A). Remarkbly, the SCE1-YFP fusion protein was found to
be associated with native Toc159 and Toc132, but not with the negative control proteins Tic110 or
Tic40 (Figure 5A). Conversely, YFP-HA did not associate with any of the tested proteins. Given that
SCE1 is a promiscuous enzyme that associates with thousands of proteins (Elrouby and Coupland,
2010), and that these interactions are likely to be transient, it is remarkable that TOC co-elution was
detectable in this experiment.

In the second experiment, we cloned the *SUMO1*, *SUMO2* and *SUMO3* coding sequences into a
vector that appends an N-terminal YFP tag to its insert (Karimi et al., 2002), a modification which
previous studies have shown to be tolerated (Ayaydin and Dasso, 2004). All three constructs

expressed well and showed the expected nucleocytoplasmic fluorescence pattern when transiently
expressed in protoplasts (Figure 5 – Supplement 2B). The three constructs were expressed in parallel
in protoplasts alongside the YFP-HA negative control construct. As in the previous experiment, the
protoplasts were subjected to YFP-Trap immunoprecipitation, and the samples were subsequently
analysed by immunoblotting. Remarkably, all three YFP-SUMO proteins were found to physically
associate with Toc159, although YFP-SUMO3 clearly bound Toc159 with the greatest affinity (Figure
5B; Figure 5 – Supplement 3). Moreover, inspection of an extended exposure of the anti-YFP blot
revealed a number of higher molecular weight bands that we interpret to be SUMO adducts and
indicative of the functionality of the fusions (Figure 5 – Supplement 4). In contrast with the SUMO
fusions, the YFP-HA negative control did not associate with Toc159, and none of the four YFP fusion
proteins physically associated with Tic40, a negative control protein (Figure 5B).

The immunoprecipitation experiment described above identified SUMO3 as having the highest
affinity for Toc159. To extend our analysis of SUMO3 to include another TOC proteins, and to
provide more direct evidence for TOC protein SUMOylation, the experiment was repeated with
modifications, as follows. Protoplasts were co-transfected with YFP-SUMO3 and Toc33-HA, or YFP-
HA and Toc33-HA; in each case, Toc33 was transiently overexpressed to aid detection of this
component and its adducts. Upon co-expression of these construct pairs, the protoplast samples
were subjected to YFP-Trap immunoprecipitation analysis, as described earlier. In accordance with
the Toc159 result (Figure 5B), YFP-SUMO3, but not YFP-HA, was found to physically associate with
Toc33-HA (Figure 5C). Moreover, bands of the exact expected molecular weight for Toc33-HA
bearing one or two YFP-SUMO3 moieties (75 and 114 kDa) were also detected. These bands were
accompanied by a high molecular weight smear at the top of the immunoblot, which is indicative of
complex, multisite or chain SUMOylation.

Discussion

This work has revealed a genetic and molecular link between the SUMO system and chloroplast
protein import. The genetic experiments demonstrated that SUMO system mutations can suppress
the phenotype of the Toc33 mutant, *ppi1*, while the molecular and biochemical experiments
indicated that TOC proteins associate with key SUMO system proteins and are SUMOylated. Visible
suppression effects observed in the *ppi1* / SUMO system double mutants were linked to
improvements in chloroplast development and enhanced accumulation of key TOC proteins. Thus,
our results suggest that SUMOylation acts to destabilise the TOC complex, and that when such
SUMOylation is perturbed the TOC proteins are stabilised. We interpret that TOC complexes
containing Toc34, Toc75 and Toc159 accumulate at higher levels in *ppi1* / SUMO system double
mutants, and that this synthetically improves the double mutant phenotypes relative to the *ppi1*
control. Importantly, each core TOC protein, including all of those analysed in this study, was
predicted with high probability to have one or more SUMOylation sites (Table 1) (Zhao et al., 2014;
Beauclair et al., 2015).

The *ppi1* suppression effects described here are remarkably similar to those mediated by the *sp1*
and *sp2* mutations (Ling et al., 2012; Ling et al., 2019). Like *sp1* and *sp2*, SUMO system mutations can
partially suppress *ppi1* with respect to chlorophyll concentration, TOC protein accumulation, and
chloroplast development. This similarity suggests that SUMOylation may regulate the activity of the
CHLORAD pathway. This is an attractive hypothesis, as both SUMOylation and the CHLORAD
pathway are activated by various forms of environmental stress (Kurepa et al., 2003; Ling and Jarvis,
2015; Ling et al., 2019). One possibility is that the SUMOylation of TOC proteins promotes their
CHLORAD-mediated degradation. Indeed, as already noted, the ability to carry out SUMOylation is

negatively correlated with the stability of TOC proteins in the context of the developed plants
studied here. However, it should be kept in mind that SUMOylation can both promote and
antagonise the effects of ubiquitination, in different situations (Desterro et al., 1998; Ahner et al.,
2013; Liebelt and Vertegaal, 2016); and so our results do not preclude the possibility that
SUMOylation may have different consequences for chloroplast biogenesis in other contexts. The
Toc159 receptor is regulated by SP1 when integrated into the outer envelope membrane (Ling et al.,
2012; Ling et al., 2019), but by a different E3 ligase when it exists as a cytosolic precursor during the
earliest stages of development before germination (Shanmugabalaji et al., 2018). Thus, regulation by
SUMOylation might be similarly different in these two distinct developmental contexts.

The precise mechanisms underpinning the observed negative regulation of the TOC apparatus by
SUMOylation are currently unknown. One possibility is that the SUMOylation of TOC proteins
promotes their association with SP1. SUMOylation can modify protein-protein interactions, and
some RING-type E3 ubiquitin ligases specifically recognise SUMOylated substrates (Sriramachandran
and Dohmen, 2014). However, these SUMO-targeted ubiquitin ligases (STUbLs) typically contain
SUMO-interacting motifs (SIMs) which guide the ligases to SUMO proteins conjugated to their
substrates, and these are not apparent in SP1 (data not shown) (Zhao et al., 2014). However, SP1
forms a complex with SP2 and very likely additional cofactors, and these could hypothetically
provide a SUMO binding interface. Another possibility is that SUMOylation could be involved in the
recruitment of Cdc48 from the cytosol. Two important Cdc48 cofactors are Ufd1 and Npl4, and the
former contains a SUMO-interacting motif which can guide Cdc48 to SUMOylated proteins (Nie et
al., 2012; Baek et al., 2013). Moreover, the SUMO-mediated recruitment of Cdc48 has important
roles in the maintenance of genome stability in yeast (Bergink et al., 2013).

The biochemical experiments described in this manuscript indicate that, of the three SUMO isoforms
tested, SUMO3 binds TOC proteins with the highest affinity. However, there is an apparent
incongruence between the results of these experiments and the results of the genetic experiments.
While the *sum1-1* and *sum2-1* mutants were found to additively suppress *ppi1*, the *sum3-1* mutant
did not suppress *ppi1*. At face value, this seems puzzling; however, it can be explained by the relative
abundance of the three SUMO proteins *in planta*. SUMO1 and SUMO2 are highly abundant relative
to SUMO3, which is, at steady state, very weakly abundant (van den Burg et al., 2010). The
immunoprecipitation data shown in Figure 5B indicated that SUMO1 and SUMO2 can weakly
interact with Toc159, and so it is likely that these two isoforms can compensate for the loss of
SUMO3 in the *sum3-1* mutant. Although SUMO3 associates with TOC proteins with the highest
affinity, the higher abundance of the other two SUMO proteins may facilitate such compensation. It
is also noteworthy that, when overexpressed, SUMO3 accentuates the *ppi1* phenotype to a far
greater extent than does SUMO1.

It is now well established that the regulation of chloroplast protein import has critical roles in plant
development and stress acclimation (Sowden et al., 2018; Watson et al., 2018). Here, we
demonstrate regulatory crosstalk between the SUMO system and chloroplast protein import, and
present results which are consistent with a model in which SUMOylation modulates the activity or
effects of the CHLORAD pathway. The precise nature of the links between these two critically
important control systems will be the subject of future investigation.

Materials and methods

Plant material and growth conditions

All *Arabidopsis thaliana* plants used in this work were of the Columbia-0 (Col-0) ecotype. The
mutants used in most of the analyses (*ppi1*, *sce1-4*, *sum1-1*, *sum2-1*, *sum3-1*, *hsp93-V-I*, *tic40-4*)
have been described previously (Jarvis et al., 1998; Kovacheva et al., 2005; Saracco et al., 2007; van
den Burg et al., 2010). The *siz1-4* (SAIL_805_A10) and *siz1-5* (SALK_111280) mutants were obtained
from the Salk Institute Genomic Analysis Laboratory (SIGnAL) (Alonso et al., 2003), via the
Nottingham *Arabidopsis* Stock Centre (NASC). Each line was verified via PCR genotyping (see Table 2
for primer sequences) and phenotypic analysis (including the double and triple mutants). The
positions of the T-DNA insertions were mapped via PCR. The following primer pairs were used to
generate diagnostic amplicons from gDNA: LB1 and Siz1-Seq-1R (for mapping *siz1-4*), and LBb1 and
Siz1-Seq-3R (for mapping *siz1-5*) (see Table 2D and 2E). The amplicons were sequenced and the
positions of the T-DNA insertions inferred from the sequence data.

In most experiments, plants were grown on soil (80% (v/v) compost (Levington M2), 20% (v/v)
vermiculite (Sinclair Pro, medium particle size)). However, where plants were grown for selection of
transformants or for chloroplast isolation, seeds were surface sterilised and sown on petri plates
containing Murashige-Skoog (MS) agar medium. The plates were stored at 4°C for 48 hours before
being transferred to a growth chamber. Both soil-grown and plate-grown plants were kept in a
growth chamber (Percival Scientific) under long-day conditions (16 hours light, 8 hours dark). The
light intensity was approximately 120 µE m^{-2} s^{-1}, the temperature was held constant at 20°C, and the
humidity was held constant at approximately 70% (relative humidity).

Chlorophyll measurements

359 Chlorophyll measurements were taken from mature rosette leaves in each instance. A handheld
360 Konica-Minolta SPAD-502 meter was used to take each measurement, and the raw values were
361 converted into chlorophyll concentration values (nmol/mg tissue) via published calibration
362 equations (Ling et al., 2011).

Chloroplast isolation and protein extraction

Chloroplasts were isolated from 14-day-old, plate-grown seedlings as described previously (Flores-
Pérez and Jarvis, 2017). Some of the seedlings were heat-shocked immediately prior to chloroplast
isolation. To do this, the plates containing the seedlings were wrapped in clingfilm and placed into a
water bath (42°C for one hour, followed by a one hour recovery period at 22°C). Protein samples
were prepared from the isolated chloroplasts by extraction using SDS-PAGE sample buffer, as well as
from whole 14-day-old seedlings as previously described (Kovacheva et al., 2005). In some cases, the
samples were treated with 10 mM N-ethylmaleimide (Hilgarth and Sarge, 2005); this was added
directly to the protein extraction buffer (whole seedling samples), or to the chloroplast isolation
buffer following polytron homogenization and all subsequent buffers (chloroplast samples).

Plasmid constructs

The constructs used in the BiFC experiments were generated as follows. The coding sequences of
SCE1, *SIZ1*, *TOC159*, *TOC132*, *TOC34* and *TOC33* were PCR amplified from wild-type cDNA (see Table
2 for primer sequences). In the case of *ΔOEP7*, the first 105 base pairs of the *OEP7* coding sequence
were amplified; this encodes a truncated sequence which is sufficient to efficiently target the full-
length YFP protein to the chloroplast outer envelope membrane (Lee et al., 2001). The inserts were
cloned into one of the following complementary vectors: pSAT4(A)-cEYFP-N1 (*SCE1*), pSAT4-nEYFP-
C1 (*TOC159*, *TOC132*, *TOC34*, *TOC33*), or pSAT4(A)-nEYFP-N1 (*ΔOEP7*), which were described
previously (Tzfira et al., 2005; Citovsky et al., 2006).

The constructs used in the immunoprecipitation experiments were generated as follows. The coding
sequences of *SCE1*, *SUMO1*, *SUMO2* and *SUMO3* were PCR amplified from wild-type cDNA using
primers bearing 5' attB1 and attB2 adaptor sequences (see Table 2 for primer sequences). The
amplicons were then cloned into pDONR221 (Invitrogen), a Gateway entry vector. The inserts from
the resulting entry clones were then transferred to one of two destination vectors: p2GWY7 (*SCE1*)
or p2YGW7 (*SUMO1*, *SUMO2*, *SUMO3*); the former appends a C-terminal YFP tag to its insert, and
the latter appends an N-terminal YFP tag to its insert (Karimi et al., 2002; Karimi et al., 2005). The
Toc33-HA and YFP-HA constructs have been described previously (Ling et al., 2019).

The constructs used to generate transgenic plants were generated as follows. The coding sequences
of *SUMO1* and *SUMO3* were PCR amplified from wild-type cDNA using primers bearing 5' attB1 and
attB2 adaptor sequences (see Table 2 for primer sequences). The inserts were then cloned into
pDONR201 (Invitrogen), a Gateway entry vector. The inserts from the resulting entry clones were
then transferred to the pH2GW7 binary destination vector (Karimi et al., 2002; Karimi et al., 2005).

Transient expression assays

Protoplasts were isolated from mature rosette leaves of wild-type *Arabidopsis* plants and
transfected in accordance with an established method (Wu et al., 2009; Ling et al., 2012). In the BiFC
experiments, 100 µL protoplast suspension (containing approximately 10^5 protoplasts) was
transfected with 5 µg plasmid DNA; in the immunoprecipitation experiments, 600 µL protoplast
suspension (containing approximately 6×10^5 protoplasts) was transfected with 30 µg plasmid DNA.
In both cases, the samples were analysed after 15-18 hours.

Stable plant transformation

Transgenic lines carrying the *SUMO1-OX* or *SUMO3-OX* constructs were generated via
Agrobacterium-mediated floral dip transformation (Clough and Bent, 1998). Transformed plants (T_1
generation) were selected on MS medium containing phosphinothricin. Multiple T_2 families were
analysed in each case, and lines bearing a single T-DNA insertion were taken forward for further
analysis. Transgene expression was analysed by semi-quantitative RT-PCR as described previously
(Kasmati et al., 2011) (see Table 2 for primer sequences).

Transmission electron microscopy

Transmission electron micrographs were recorded using mature rosette leaves as previously
described (Huang et al., 2011). Images were taken from three biological replicates (different leaves
from different individual plants), and at least 10 images were taken per replicate. The images were
analysed using ImageJ (Schneider et al., 2012). The freehand tool was used to measure the plan area
of the chloroplasts. For this, between 9 and 28 chloroplasts were analysed for each biological
replicate (i.e., for each plant), and then an average value for each replicate was calculated and used
for statistical comparisons. The analysis of chloroplast ultrastructure was performed as in previous
work (Huang et al., 2011). For this, between 3 and 8 chloroplasts were analysed per biological
replicate, and the data were processed as above.

BiFC experiments

The BiFC experiments were carried out as described previously (Ling et al., 2019). Protoplasts were
co-transfected with two constructs encoding fusion proteins bearing complementary fragments of
the YFP protein (nYFP and cYFP) (Citovsky et al., 2006). After transfection and overnight incubation,
the protoplasts were imaged using a Leica TCS SP5 laser scanning confocal microscope equipped
with a Leica HC Plan Apochromat CS2 63.0x UV water immersion lens with a numerical aperture

(N.A.) of 1.2. YFP was excited with an Argon-ion laser at 514 nm, selected using an acousto-optic
tuneable filter (AOTF), and was detected using a 525-600 nm bandpass filter and a photomultiplier.
Chlorophyll fluorescence was simultaneously excited with 514 nm excitation and detected with a
680-700 nm bandpass filter using a photomultiplier. Images were collected in 8-bit resolution with
the pinhole set at 111.5 μm (1 Airy Unit), using 16-line averaging and a scan speed of 400 Hz. The
image size was 512 x 512 pixels, with an (x,y) pixel size of 0.239 μm. Images were processed in the
Leica Application Suite (LAS) software.

Immunoblotting and immunoprecipitation

Protein extraction and immunoblotting were performed as described previously (Kovacheva et al.,
2005). Total protein samples were extracted from 50 mg of intact, pooled seedlings after two weeks
of growth. To detect proteins, we used an anti-SUMO1 antibody (Ab5316, Abcam), an anti-Toc75-III
antibody (Kasmati et al., 2011), an anti-Toc159 antibody (Bauer et al., 2000), an anti-Toc132
antibody (Ling et al., 2012), an anti-Toc33 antibody (Kasmati et al., 2011), an anti-Tic110 antibody
(Aronsson et al., 2010), an anti-Tic40 antibody (Kasmati et al., 2011), and an anti-green fluorescent
protein antibody (Sigma, SAB4301138). In most cases, the secondary antibody used was goat anti-
rabbit immunoglobulin G (IgG) conjugated with horseradish peroxidase (Sigma, 12-348); and protein
bands were visualised via chemiluminescence using an ECL Plus Western blotting detection kit (GE
Healthcare) and an LAS-4000 imager (Fujifilm). However, in the case of Figure 5 – Supplement 1, the
secondary antibody was goat anti-rabbit IgG conjugated with alkaline phosphatase (Sigma, A3687),
and the membrane was incubated with BCIP/NBT chromogenic substrate (Sigma, B3679).

The immunoprecipitation (IP) experiments were carried out as described previously (Ling et al.,
2019). Constructs encoding YFP-conjugated fusion proteins (YFP-HA, SCE1-YFP, YFP-SUMO1, YFP-
SUMO2, YFP-SUMO3) were transiently expressed in protoplasts. In some cases, the constructs were
co-expressed with a construct encoding Toc33-HA. The protoplasts were solubilised using IP buffer
containing 1% Triton X-100, and the resulting lysates were incubated with GFP-Trap beads
(Chromotek). After four washes in IP buffer, the protein samples were eluted by boiling in SDS-PAGE
loading buffer, and then analysed by immunoblotting.

Statistical analysis

453   The data from each experiment were analysed in R. In most cases, two-tailed T-tests were
454   performed. However, in one case, a one-way ANOVA was performed in conjunction with a Tukey
455   HSD test (as indicated in the figure legend). The figures are annotated to indicate the level of
456   significance, as follows: ns, not significant; *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; ****, $p<0.0001$;
457   *****, $p<0.00001$.

SUMO site prediction

The amino acid sequences of Toc159, Toc132, Toc120, Toc90, Toc75, Toc33 and Toc34 were
retrieved from The *Arabidopsis* Information Resource (TAIR) website (Berardini et al., 2015). The
GPS-SUMO algorithm was applied to all seven sequences (http://sumosp.biocuckoo.org/online.php)
(Zhao et al., 2014). The 'high stringency' setting was applied. The *p*-values were generated by the
GPS-SUMO algorithm, and hits that were accompanied by *p*-values exceeding $p>0.05$ were manually
removed. The JASSA algorithm was also applied to all seven amino acid sequences
(http://www.jassa.fr/index.php) (Beauclair et al., 2015). In this case, the 'high cut-off' setting was
applied. The GPS-SUMO and JASSA algorithms use fundamentally different methodologies (Chang et
al., 2018).

References and notes

Ahner, A., Gong, X., and Frizzell, R.A. (2013). Cystic fibrosis transmembrane conductance regulator degradation: cross-talk between the ubiquitylation and SUMOylation pathways. FEBS Journal **280,** 4430-4438.

Alonso, J.M., Stepanova, A.N., Leisse, T.J., Kim, C.J., Chen, H., Shinn, P., Stevenson, D.K., Zimmerman, J., Barajas, P., Cheuk, R., Gadrinab, C., Heller, C., Jeske, A., Koesema, E., Meyers, C.C., Parker, H., Prednis, L., Ansari, Y., Choy, N., Deen, H., Geralt, M., Hazari, N., Hom, E., Karnes, M., Mulholland, C., Ndubaku, R., Schmidt, I., Guzman, P., Aguilar-Henonin, L., Schmid, M., Weigel, D., Carter, D.E., Marchand, T., Risseeuw, E., Brogden, D., Zeko, A., Crosby, W.L., Berry, C.C., and Ecker, J.R. (2003). Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. Science **301,** 653-657.

Aronsson, H., Combe, J., Patel, R., Agne, B., Martin, M., Kessler, F., Jarvis, P. (2010). Nucleotide binding and dimerization at the chloroplast pre-protein import receptor, atToc33, are not essential in vivo but do increase import efficiency. Plant Journal **63,** 297-311.

Ayaydin, F., and Dasso, M. (2004). Distinct in vivo dynamics of vertebrate SUMO paralogues. Molecular Biology of the Cell **15,** 5208-5218.

Baek, G.H., Cheng, H., Choe, V., Bao, X., Shao, J., Luo, S., and Rao, H. (2013). Cdc48: a swiss army knife of cell biology. Journal of Amino Acids **2013,** 183421.

Bauer, J., Chen, K., Hiltbunner, A., Wehrli, E., Eugster, M., Schnell, D., and Kessler, F. (2000). The major protein import receptor of plastids is essential for chloroplast biogenesis. Nature **403,** 203-207.

Beauclair, G., Bridier-Nahmias, A., Zagury, J.F., Saib, A., and Zamborlini, A. (2015). JASSA: a comprehensive tool for prediction of SUMOylation sites and SIMs. Bioinformatics **31,** 3483-3491.

Berardini, T.Z., Reiser, L., Li, D., Mezheritsky, Y., Muller, R., Strait, E., and Huala, E. (2015). The Arabidopsis information resource: Making and mining the "gold standard" annotated reference plant genome. Genesis **53,** 474-485.

Bergink, S., Ammon, T., Kern, M., Schermelleh, L., Leonhardt, H., and Jentsch, S. (2013). Role of Cdc48/p97 as a SUMO-targeted segregase curbing Rad51-Rad52 interaction. Nature Cell Biology **15,** 526-532.

Chang, C.C., Tung, C.H., Chen, C.W., Tu, C.H., and Chu, Y.W. (2018). SUMOgo: Prediction of sumoylation sites on lysines by motif screening models and the effects of various post-translational modifications. Scientific Reports **8,** 15512.

Citovsky, V., Lee, L.Y., Vyas, S., Glick, E., Chen, M.H., Vainstein, A., Gafni, Y., Gelvin, S.B., and Tzfira, T. (2006). Subcellular localization of interacting proteins by bimolecular fluorescence complementation *in planta*. Journal of Molecular Biology **362,** 1120-1131.

Clough, S.J., and Bent, A.F. (1998). Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. Plant Journal **16,** 735-743.

Desterro, J.M., Rodriguez, M.S., and Hay, R.T. (1998). SUMO-1 modification of IkappaBalpha inhibits NF-kappaB activation. Molecular Cell **2,** 233-239.

Elrouby, N., and Coupland, G. (2010). Proteome-wide screens for small ubiquitin-like modifier (SUMO) substrates identify Arabidopsis proteins implicated in diverse biological processes. Proceedings of the National Academy of Sciences of the United States of America **107,** 17415-17420.

Flores-Pérez, Ú., and Jarvis, P. (2017). Isolation and suborganellar fractionation of Arabidopsis chloroplasts. Methods in Molecular Biology **1511,** 45-60.

Grimmer, J., Helm, S., Dobritzsch, D., Hause, G., Shema, G., Zahedi, R.P., and Baginsky, S. (2020). Mild proteasomal stress improves photosynthetic performance in Arabidopsis chloroplasts. Nature Communications **11,** 1662.

Hilgarth, R.S., and Sarge, K.D. (2005). Detection of sumoylated proteins. Methods in Molecular
Biology **301,** 329-338.
Hirsch, S., Muckel, E., Heemeyer, F., von Heijne, G., and Soll, J. (1994). A receptor component of the
chloroplast protein translocation machinery. Science **266,** 1989-1992.
Huang, W.H., Ling, Q.H., Bedard, J., Lilley, K., and Jarvis, P. (2011). In vivo analyses of the roles of
essential Omp85-related proteins in the chloroplast outer envelope membrane. Plant
Physiology **157,** 147-159.
Inaba, T., Alvarez-Huerta, M., Li, M., Bauer, J., Ewers, C., Kessler, F., and Schnell, D.J. (2005).
Arabidopsis Tic110 is essential for the assembly and function of the protein import
machinery of plastids. Plant Cell **17,** 1482-1496.
Ishida, T., Fujiwara, S., Miura, K., Stacey, N., Yoshimura, M., Schneider, K., Adachi, S., Minamisawa,
K., Umeda, M., and Sugimoto, K. (2009). SUMO E3 ligase HIGH PLOIDY2 regulates endocycle
onset and meristem maintenance in Arabidopsis. Plant Cell **21,** 2284–2297.
Jarvis, P. (2008). Targeting of nucleus-encoded proteins to chloroplasts in plants. New Phytologist
179, 257-285.
Jarvis, P., and Lopez-Juez, E. (2013). Biogenesis and homeostasis of chloroplasts and other plastids.
Nature Reviews Molecular Cell Biology **14,** 787-802.
Jarvis, P., Chen, L.J., Li, H.M., Pete, C.A., Fankhauser, C., and Chory, J. (1998). An *Arabidopsis*
mutant defective in the plastid general protein import apparatus. Science **282,** 100-103.
Karimi, M., Inze, D., and Depicker, A. (2002). GATEWAY vectors for *Agrobacterium*-mediated plant
transformation. Trends in Plant Science **7,** 193-195.
Karimi, M., De Meyer, B., and Hilson, P. (2005). Modular cloning in plant cells. Trends Plant Science
10, 103-105.
Kasmati, A.R., Töpel, M., Patel, R., Murtaza, G., and Jarvis, P. (2011). Molecular and genetic
analyses of Tic20 homologues in *Arabidopsis thaliana* chloroplasts. Plant Journal **66,** 877-
889.
Kessler, F., Blobel, G., Patel, H.A., and Schnell, D.J. (1994). Identification of two GTP-binding
proteins in the chloroplast protein import machinery. Science **266,** 1035-1039.
Kovacheva, S., Bedard, J., Patel, R., Dudley, P., Twell, D., Rios, G., Koncz, C., and Jarvis, P. (2005). *In*
vivo studies on the roles of Tic110, Tic40 and Hsp93 during chloroplast protein import. Plant
Journal **41,** 412-428.
Kurepa, J., Walker, J.M., Smalle, J., Gosink, M.M., Davis, S.J., Durham, T.L., Sung, D.Y., and Vierstra,
R.D. (2003). The small ubiquitin-like modifier (SUMO) protein modification system in
Arabidopsis. Accumulation of SUMO1 and -2 conjugates is increased by stress. Journal of
Biological Chemistry **278,** 6862-6872.
Lee, S., Lee, D.W., Lee, Y., Mayer, U., Stierhof, Y.D., Jurgens, G., and Hwang, I. (2009). Heat shock
protein cognate 70-4 and an E3 ubiquitin ligase, CHIP, mediate plastid-destined precursor
degradation through the ubiquitin-26S proteasome system in *Arabidopsis*. Plant Cell **21,**
3984-4001.
Lee, Y.J., Kim, D.H., Kim, Y.W., and Hwang, I. (2001). Identification of a signal that distinguishes
between the chloroplast outer envelope membrane and the endomembrane system in vivo.
Plant Cell **13,** 2175-2190.
Liebelt, F., and Vertegaal, A.C. (2016). Ubiquitin-dependent and independent roles of SUMO in
proteostasis. American Journal of Physiology Cell Physiology **311,** C284-C296.
Ling, Q., and Jarvis, P. (2015). Regulation of chloroplast protein import by the ubiquitin E3 ligase SP1
is important for stress tolerance in plants. Current Biology **25,** 2527-2534.
Ling, Q., Huang, W., and Jarvis, P. (2011). Use of a SPAD-502 meter to measure leaf chlorophyll
concentration in Arabidopsis thaliana. Photosynthesis Research **107,** 209-214.
Ling, Q., Huang, W., Baldwin, A., and Jarvis, P. (2012). Chloroplast biogenesis is regulated by direct
action of the ubiquitin-proteasome system. Science **338,** 655-659.

Ling, Q., Broad, W., Trösch, R., Töpel, M., Demiral Sert, T., Lymperopoulos, P., Baldwin, A., and Jarvis, R.P. (2019). Ubiquitin-dependent chloroplast-associated protein degradation in plants. Science **363**.

Liu C., Yu H., Li L. (2019). SUMO modification of LBD30 by SIZ1 regulates secondary cell wall formation in *Arabidopsis thaliana*. PLOS Genetics **15**, e1007928.

Miura, K., Rus, A., Sharkhuu, A., Yokoi, S., Karthikeyan, A.S., Raghothama, K.G., Back, D., Koo, Y.D., Jin, J.B., Bressan, R.A., Yun, D.J., and Hasegawa, P.M. (2005). The Arabidopsis SUMO E3 ligase SIZ1 controls phosphate deficiency responses. Proceedings of the National Academy of Sciences of the United States of America **102,** 9734-9734.

Nie, M., Aslanian, A., Prudden, J., Heideker, J., Vashisht, A.A., Wohlschlegel, J.A., Yates, J.R., and Boddy, M.N. (2012). Dual recruitment of Cdc48 (p97)-Ufd1-Npl4 ubiquitin-selective segregase by small ubiquitin-like modifier protein (SUMO) and ubiquitin in SUMO-targeted ubiquitin ligase-mediated genome stability functions. Journal of Biological Chemistry **287,** 29610-29619.

Perry, S.E., and Keegstra, K. (1994). Envelope membrane proteins that interact with chloroplastic precursor proteins. Plant Cell **6,** 93-105.

Saracco, S.A., Miller, M.J., Kurepa, J., and Vierstra, R.D. (2007). Genetic analysis of SUMOylation in Arabidopsis: conjugation of SUMO1 and SUMO2 to nuclear proteins is essential. Plant Physiology **145,** 119-134.

Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of image analysis. Nature Methods **9,** 671-675.

Schnell, D.J., Kessler, F., and Blobel, G. (1994). Isolation of components of the chloroplast protein import machinery. Science **266,** 1007-1012.

Shanmugabalaji, V., Chahtane, H., Accossato, S., Rahire, M., Gouzerh, G., Lopez-Molina, L., and Kessler, F. (2018). Chloroplast biogenesis controlled by DELLA-TOC159 interaction in early plant development. Current Biology **28,** 2616-2623 e2615.

Sowden, R.G., Watson, S.J., and Jarvis, P. (2018). The role of chloroplasts in plant pathology. Essays in Biochemistry **62,** 21-39.

Sriramachandran, A.M., and Dohmen, R.J. (2014). SUMO-targeted ubiquitin ligases. Biochimica et Biophysica Acta **1843,** 75-85.

Tranel, P.J., Froehlich, J., Goyal, A., and Keegstra, K. (1995). A component of the chloroplastic protein import apparatus is targeted to the outer envelope membrane via a novel pathway. The EMBO Journal **14,** 2436-2446.

Tzfira, T., Tian, G.W., Lacroix, B., Vyas, S., Li, J., Leitner-Dagan, Y., Krichevsky, A., Taylor, T., Vainstein, A., and Citovsky, V. (2005). pSAT vectors: a modular series of plasmids for autofluorescent protein tagging and expression of multiple genes in plants. Plant Molecular Biology **57,** 503.

van den Burg, H.A., Kini, R.K., Schuurink, R.C., and Takken, F.L.W. (2010). *Arabidopsis* small ubiquitin-like modifier paralogs have distinct functions in development and defense. Plant Cell **22,** 1998-2016.

Watson, S.J., Sowden, R.G., and Jarvis, P. (2018). Abiotic stress-induced chloroplast proteome remodelling: a mechanistic overview. Journal of Experimental Botany **69,** 2773-2781.

Wu, F.H., Shen, S.C., Lee, L.Y., Lee, S.H., Chan, M.T., and Lin, C.S. (2009). Tape-Arabidopsis Sandwich - a simpler Arabidopsis protoplast isolation method. Plant Methods **5,** 16.

Zhao, Q., Xie, Y., Zheng, Y., Jiang, S., Liu, W., Mu, W., Liu, Z., Zhao, Y., Xue, Y., and Ren, J. (2014). GPS-SUMO: a tool for the prediction of sumoylation sites and SUMO-interaction motifs. Nucleic Acids Research **42**, W325-W330.

Acknowledgements

We are very grateful to Alistair Haslam for initiating the genetic experiments. We also thank Dr Errin Johnson and Raman Dhaliwal (Dunn School of Pathology, University of Oxford) for assistance with the electron microscopy, and Professor Jane Langdale for advice during the course of the work. We are grateful to Pedro Bota and Rita Ross for technical support. This work was supported by the Oxford Interdisciplinary Bioscience Doctoral Training Partnership (DTP), and by grants from the Biotechnology and Biological Sciences Research Council (BBSRC) (grant numbers BB/K018442/1, BB/N006372/1, BB/R016984/1 and BB/R009333/1) to RPJ.

Competing interests

The application of CHLORAD as a technology for crop improvement is covered by a patent application (no. WO2019/171091 A).

(5 weeks)   (3 weeks)

Figure 1

The E2 SUMO conjugating enzyme mutation, *sce1-4*, and the E3 SUMO ligase mutations, *siz1-4* and
***siz1-5*, suppress the phenotype of plastid protein import mutant, *ppi1*.**

(A) The *ppi1 sce1-4* double mutant appeared greener than *ppi1* after approximately five weeks of
growth on soil. **(B)** The *ppi1 sce1-4* double mutant showed enhanced accumulation of chlorophyll
relative to *ppi1* after approximately five weeks of growth on soil. Measurements were taken from
the plants shown in (A) on the day of photography, as well as additional similar plants. There were
significant differences between the *ppi1* and *ppi1 sce1-4* plants (Two-tailed t-test, unpaired samples,
T = 6.15, *p* = 0.000049). **(C)** Transmission electron microscopy revealed improved chloroplast
development in mature rosette leaf mesophyll tissue of *ppi1 sce1-4* plants relative to *ppi1*. Plants

that had been grown on soil for approximately four weeks were analysed, and representative images
are shown. Scale bar = 2 μm. **(D)** Chloroplast plan area was elevated in *ppi1 sce1-4* relative to *ppi1*.
The transmission electron microscopy dataset was quantified. There were significant differences
between the *ppi1* and *ppi1 sce1-4* plants (Two-tailed t-test, unpaired samples, T = 4.65, *p* =
0.009674). **(E, F)** Thylakoid membrane development was increased in *ppi1 sce1-4* relative to *ppi1*.
The number of stacked thylakoidal lamellae per granum (E), and the number of stromal thylakoidal
lamellae emanating from each granum (granal interconnections) (F), was analysed using the
transmission electron microscopy dataset. There were significant differences between the *ppi1* and
ppi1 sce1-4 plants (Two-tailed t-test, unpaired samples: T = 5.53, *p* = 0.005221 [E]; T = 3.38, *p* =
0.0277 [F]). **(G)** The *ppi1 siz1-4* and *ppi1 siz1-5* double mutants appeared greener than *ppi1* after
different periods of growth on soil. The plants were photographed after three weeks of growth (right
panel) and then again after five weeks of growth (left panel). **(H)** The *ppi1 siz1-4* double mutant
showed enhanced accumulation of chlorophyll relative to *ppi1* after approximately five weeks of
growth on soil. Measurements were taken from the plants shown in (G) on the day of photography,
as well as additional similar plants. There were significant differences between the *ppi1* and *ppi1*
siz1-4 plants (Two-tailed t-test, unpaired samples, T = 11.01, *p* < 0.00001). **(I)** TOC protein
accumulation was improved in *ppi1 siz1-4* and *ppi1 siz1-5* relative to *ppi1*. Analysis of the levels of
Toc75 and Toc159 in *ppi1 siz1-4*, *ppi1 siz1-5*, and relevant control plants was conducted by
immunoblotting. Protein samples were taken from whole seedlings that had been grown on soil for
approximately two weeks (the plants shown in Figure 1 – Supplement 1C on the day of
photography). Tic110, a TIC-associated protein, was included as a compartment-specific loading
control. Migration positions of standards are displayed to the left of the gel images, and sizes are
indicated in kDa. Unprocessed membrane images are displayed in Source data 1. **(J, K)** Toc159 and
Toc75 protein accumulation was improved in *ppi1 siz1-4* and *ppi1 siz1-5* relative to *ppi1*. Specific
bands in (I) and in Source data 1 were quantified. Error bars indicate standard deviation from the
mean. The numbers above the graph indicate the number of biological replicates per sample. There
were significant differences between the *ppi1* and *ppi1 siz1-4* samples (One-tailed t-test, unpaired
samples: T = 2.26, *p* = 0.032316 [J, Toc159]; T = 2.93, *p* = 0.021334 [K, Toc75]) and between the *ppi1*
and *ppi1 siz1-5* samples (One-tailed t-test, unpaired samples: T = 2.76, *p* = 0.01639 [J, Toc159]; T =
3.36, *p* = 0.014118 [K, Toc75]). In all bar charts, error bars indicate standard deviation from the
mean, and open red circles indicate individual data points. The numbers above the graphs indicate
the number of biological replicates per sample. Statistical significance is indicated as follows: *,
p<0.05; **, *p*<0.01; ****, *p*<0.0001; *****, *p*<0.00001.

Figure 1 – Supplement 1

The E2 SUMO conjugating enzyme mutation, *sce1-4*, and the E3 SUMO ligase mutations, *siz1-4* and
siz1-5*, suppress the phenotype of plastid protein import mutant, *ppi1.

(A) The *ppi1 sce1-4* double mutant appeared greener than *ppi1* after approximately three weeks of
growth on soil, while the single *sce1-4* mutant appeared no greener than the wild-type control. **(B)**
The *ppi1 sce1-4* double mutant showed enhanced accumulation of chlorophyll relative to *ppi1* after
approximately three weeks of growth on soil (T = 8.19, *p* < 0.0001), while the single *sce1-4* mutant
displayed no enhanced accumulation of chlorophyll relative to the wild-type control. Measurements
were taken from the plants shown in (A) on the day of photography, as well as additional similar
plants. **(C)** The *ppi1 siz1-4* and *ppi1 siz1-5* double mutants appeared greener than *ppi1* after
approximately two weeks of growth on soil, while the single *siz1-4* and *siz1-5* mutants appeared no
greener than the wild-type control. **(D)** The *ppi1 siz1-4* and *ppi1 siz1-5* double mutants showed
enhanced accumulation of chlorophyll relative to *ppi1* after approximately two weeks of growth on
soil (T = 7.16, *p* < 0.0001 and T = 8.26, *p* < 0.0001 respectively), while the single *siz1-4* and *siz1-5*
mutants displayed no enhanced accumulation of chlorophyll relative to the wild-type control.
Measurements were taken from the plants shown in (C) on the day of photography, as well as
additional similar plants. In all bar charts, error bars indicate standard deviation from the mean, and
open red circles indicate individual data points. The numbers above the graphs indicate the number
of biological replicates per sample. Statistical significance is indicated as follows: *****, *p*<0.00001.

Figure 1 – Supplement 2

Molecular analysis of the *siz1-4* and *siz1-5* mutants.

(A) Schematic representation of the *SIZ1* (At5g60410) gene, annotated with the positions of the *siz1* T-DNA insertions. The *siz1-4* and *siz1-5* insertion sites were verified in this study by DNA sequencing. The black boxes indicate exons, while the interconnecting black lines represent introns. The white boxes indicate untranslated regions. **(B)** Analysis of *SIZ1* mRNA expression in *siz1-4* and *siz1-5*. RNA samples were taken from two-week-old seedlings and used as a template for cDNA synthesis. Primers specific to *SIZ1* and the reference gene *eIF4E1* were used in two parallel RT-PCR experiments. gDNA, genomic DNA. Migration positions of standards are displayed to the right of the gel images, and sizes are indicated in base pairs.

Figure 1 – Supplement 3

The *siz1-4* and *siz1-5* mutants display reduced global SUMOylation in response to heat shock.

14-day-old seedlings of the indicated genotypes were subjected to heat shock (42°C for 1 hour,
followed by a 1-hour recovery period at 22°C). Protein samples were taken from whole seedlings and
analysed by immunoblotting. The *siz1-4* and *siz1-5* mutants displayed a moderate reduction of global
SUMOylation, and this reduction was similar in magnitude to the reduction observed in the
previously characterised *siz1-2* and *siz1-3* mutants. The unprocessed membrane images are
displayed in Source data 1.

Figure 2

Genetic analysis reveals specificity of the suppression mediated by the *sce1-4* and *siz1-4*
mutations.

(A) The *tic40-4 sce1-4* double mutant did not appear greener than *tic40-4* after approximately four
weeks of growth on soil. **(B)** The *tic40-4 sce1-4* double mutant did not show an enhanced
accumulation of chlorophyll relative to *tic40-4* after approximately four weeks of growth on soil.
Measurements were taken from the plants shown in (A) on the day of photography, as well as
additional similar plants. There were no significant differences between the *tic40-4* and *tic40-4 sce1-*
4 plants (Two-tailed t-test, unpaired samples, T = 1.25, *p* = 0.280106). **(C)** The *hsp93-V-1 sce1-4*
double mutant did not appear greener than *hsp93-V-1* after approximately four weeks of growth on
soil. **(D)** The *hsp93-V-1 sce1-4* double mutant did not show an enhanced accumulation of chlorophyll
relative to *hsp93-V-1* after approximately four weeks of growth on soil. Measurements were taken
from the plants shown in (C) on the day of photography, as well as additional similar plants. There
were no significant differences between the *hsp93-V-1* and *hsp93-V-1 sce1-4* plants (Two-tailed t-
test, unpaired samples, T = 0.18, *p* = 0.860702). **(E)** The *sce1-4* single mutant did not appear greener
than wild-type plants after approximately four weeks of growth on soil. **(F)** The *sce1-4* single mutant
did not show an enhanced accumulation of chlorophyll relative to wild-type plants after

approximately four weeks of growth on soil. Measurements were taken from the plants shown in (E)
on the day of photography, as well as additional similar plants. There were no significant differences
between the *sce1-4* and wild type plants (Two-tailed t-test, unpaired samples, T = 0.38, *p* =
0.708484). **(G)** The *siz1-4* single mutant did not appear greener than wild-type plants after
approximately five weeks of growth on soil. **(H)** The *siz1-4* single mutant did not show an enhanced
accumulation of chlorophyll relative to wild-type plants after approximately five weeks of growth on
soil. Measurements were taken from the plants shown in (G) on the day of photography, as well as
from additional similar plants. There were no significant differences between the *siz1-4* and wild
type plants (T = 0.96, *p* = 0.370055). In all bar charts, error bars indicate standard deviation from the
mean, and open red circles indicate individual data points. The numbers above the graphs indicate
the number of biological replicates per sample. Statistical significance is indicated as follows: ns, not
significant.

Figure 3

SCE1 physically interacts with TOC proteins *in vivo.*

Bimolecular fluorescence complementation (BiFC) analysis of SCE1 protein-protein interactions was performed by imaging *Arabidopsis* protoplasts co-expressing proteins fused to complementary N-terminal (nYFP) and C-terminal (cYFP) fragments of the YFP protein, as indicated. Chlorophyll autofluorescence images were used to orientate the YFP signals in relation to the chloroplasts. SCE1 physically associated with all tested TOC proteins. The images shown are representative confocal micrographs indicating associations between SCE1 and Toc159, Toc132, Toc34 and Toc33. In contrast, SCE1 did not physically associate with ΔOEP7 (comprising the transmembrane domain of OEP7, which is sufficient to direct targeting to the chloroplast outer envelope), which served as a negative control. Scale bars = 10 μm.

Figure 4

Genetic interactions between *ppi1* and the genes encoding three SUMO isoforms.

(A) The *ppi1 sum1-1*, *ppi1 sum2-1*, and *ppi1 sum1-1† sum2-1* double and triple mutants appeared

greener than *ppi1* after approximately four weeks of growth on soil. The dagger symbol indicates

that the triple mutant was heterozygous with respect to the *sum1-1* mutation. **(B)** The *ppi1 sum1-1*,

ppi1 sum2-1, and *ppi1 sum1-1† sum2-1* double and triple mutants showed enhanced accumulation

of chlorophyll relative to *ppi1* after approximately four weeks of growth on soil. Measurements were
taken from the plants shown in (A) on the day of photography, as well as additional similar plants.
There were significant differences between the samples, as measured via a one-way ANOVA (F =
26.21, p = 6.65 × 10-14). A post-hoc Tukey HSD test indicated that there was a significant difference
between the *ppi1* and *ppi1 sum1-1* samples (p < 0.00001). There were also significant differences
between the *ppi1* and *ppi1 sum2-1* samples (p < 0.00001), and between the *ppi1* and *ppi1 sum1-1†*
sum2-1 samples (p < 0.00001). **(C)** The *ppi1* SUMO1 overexpression (OX) lines appeared smaller and
paler than *ppi1* after approximately four weeks of growth on soil. **(D)** The *ppi1* SUMO1 OX lines
showed reduced accumulation of chlorophyll relative to *ppi1* after approximately four weeks of
growth on soil. Measurements were taken from the plants shown in (C) on the day of photography,
as well as additional similar plants. There were significant differences between the *ppi1* and *ppi1*
SUMO1 OX #1 plants (Two-tailed t-test, unpaired samples, T = 2.27, p = 0.026832), and between the
ppi1 and *ppi1* SUMO1 OX #2 plants (Two-tailed t-test, unpaired samples, T = 2.49, p = 0.01634). **(E)**
The *ppi1 sum3-1* double mutant did not appear greener than *ppi1* after approximately four weeks of
growth on soil. **(F)** The *ppi1 sum3-1* double mutant did not show an enhanced accumulation of
chlorophyll relative to *ppi1* after approximately four weeks of growth on soil. Measurements were
taken from the plants shown in (E) on the day of photography, as well as additional similar plants.
There were no significant differences between the *ppi1* and *ppi1 sum3-1* plants (Two-tailed t-test,
unpaired samples, T = 0.54, p = 0.59407). **(G)** The *ppi1* SUMO3 overexpression line appeared smaller
and paler than *ppi1* after approximately three weeks of growth on soil. **(H)** The *ppi1* SUMO3
overexpression line showed reduced accumulation of chlorophyll relative to *ppi1* after
approximately three weeks of growth on soil. Measurements were taken from the plants shown in
(G) on the day of photography, as well as additional similar plants. There were significant differences
between the *ppi1* SUMO3 OX plants and *ppi1* (Two-tailed t-test, unpaired samples, T = 2.99, p =
0.008688). In all bar charts, error bars indicate standard deviation from the mean, and open red
circles indicate individual data points. The numbers above the graphs indicate the number of
biological replicates per sample. Statistical significance is indicated as follows: ns, not significant; *,
p<0.05; **, *p*<0.01; ****, *p*<0.0001.

A

B

Figure 4 – Supplement 1

The *sum1-1*, *sum2-1*, and *sum1-1† sum2-1* single and double mutants do not show enhanced
chlorophyll accumulation relative to wild-type plants.

(A) The *sum1-1*, *sum2-1*, and *sum1-1† sum2-1* mutants did not appear greener than wild-type plants
after approximately four weeks of growth on soil. **(B)** The *sum1-1*, *sum2-1*, and *sum1-1† sum2-1*
mutants did not show enhanced chlorophyll accumulation relative to wild-type plants after
approximately four weeks of growth on soil. The dagger symbol indicates that the double mutant
was heterozygous with respect to the *sum1-1* mutation. Error bars indicate standard deviation from
the mean, and open red circles indicate individual data points. The numbers above the graph
indicate the number of biological replicates per sample.

Figure 4 – Supplement 2

Analysis of the expression of the 35S:*SUMO1* and 35S:*SUMO3* transgenes in the selected transformants by RT-PCR.

Total RNA was extracted from the rosette tissue of plants that had been grown on soil for approximately four weeks. The expression of *SUMO1* **(A)** or *SUMO3* **(B)**, and of the control gene, *eIF4E1*, was analysed by semi-quantitative RT-PCR. A limited number of amplification cycles (*n* = 26) were employed to prevent saturation. Migration positions of standards are displayed to the left of the gel images, and sizes are indicated in base pairs.

Figure 5

Immunoprecipitation analysis reveals that TOC proteins are SUMO targets.

(A) SCE1 physically associated with native TOC proteins. *Arabidopsis* protoplasts expressing YFP-HA
(left panel) or SCE1-YFP (right panel) were solubilised and subjected to immunoprecipitation (IP)
analysis. The YFP-HA construct served as a negative control. In both cases, three samples were
analysed: The 'Total lysate' sample (total protein extract from solubilised protoplasts); the 'Flow
through' sample (the total protein sample after incubation with anti-YFP beads); and the 'IP' sample
(the eluted fraction of the total protein sample that bound to the anti-YFP beads). The samples were
analysed by immunoblotting, revealing that SCE1-YFP, but not the YFP-HA control, was associated

with native Toc159 and Toc132 (indicated by the two arrows). Neither SCE1-YFP nor YFP-HA was
associated with native Tic110 or Tic40, which were included as negative control proteins. **(B)** All
three SUMO isoforms physically associated with native Toc159. Protoplasts expressing YFP-HA, YFP-
SUMO1, YFP-SUMO2 or YFP-SUMO3 were solubilised and subjected to IP analysis as in (A). In all four
cases, two samples (the 'Total lysate' and the 'IP' samples) were analysed by immunoblotting.
Toc159 was resolved on an 8% acrylamide gel for four hours to maximise the resolution of high
molecular weight bands. All three YFP-SUMO proteins were found to associate with native Toc159;
however, YFP-SUMO3 immunoprecipitated Toc159 with the greatest efficiency. None of the four YFP
fusion proteins associated with native Tic40, which served as a negative control protein. **(C)** YFP-
SUMO3 physically associated with Toc33-HA and related high molecular weight species. Protoplasts
co-expressing YFP-SUMO3 or YFP-HA together with Toc33-HA were solubilised and subjected to IP
analysis as in (A). In both cases, two samples (the 'Total lysate' and 'IP' samples) were analysed by
immunoblotting. The results showed that YFP-SUMO3, but not YFP-HA, was associated with Toc33-
HA (indicated by the arrow). Bands corresponding to the molecular weight of Toc33-HA bearing one,
two or several YFP-SUMO3 motifs were detected on the membrane (indicated by the upper arrows).
The predicted molecular weight of YFP-SUMO3 is approximately 38.9 kDa. Neither YFP-SUMO3 nor
YFP-HA was associated with Tic40, which served as a negative control protein. The asterisk indicates
a nonspecific band. Short (SE) and long (LE) exposures are shown. Migration positions of standards
are displayed to the left of the gel images, and sizes are indicated in kDa. The unprocessed
membrane images are displayed in Source data 1.
854

Figure 5 – Supplement 1

Chloroplast resident proteins are SUMOylated.

Anti-SUMO1 immunoblot of protein samples taken from whole seedlings (left hand side) and

isolated chloroplasts (right hand side). Where indicated, seedlings were exposed to heat shock (42°C

for one hour) and/or incubated with 10 mM N-ethylmaleimide (NEM) to aid the detection of

SUMOylated proteins. The anti-SUMO1 antibody shows significant cross-reactivity with SUMO2.

Migration positions of standards are displayed to the left of the gel images, and sizes are indicated in

kDa.

Figure 5 – Supplement 2

Analysis of the expression of the YFP-tagged constructs used in the immunoprecipitation experiments by confocal microscopy.

The expression of the SCE1-YFP construct **(A)**, and of the three YFP-SUMO constructs **(B)**, was analysed and confirmed by imaging transfected *Arabidopsis* protoplasts. Chlorophyll autofluorescence images were employed to orientate the YFP signals in relation to the chloroplasts. Representative confocal micrographs show a typical protoplast in each case. Scale bars = 10 µm.

Figure 5 – Supplement 3

All three SUMO isoforms physically associated with native Toc159, with SUMO3 showing the
strongest association.

(A) The immunoprecipitation experiment shown in Figure 5B was repeated. Tic110 was included as
an additional negative control. The samples were run on a 10% acrylamide gel for 1 hour, as
opposed to an 8% acrylamide gel for four hours (as in Figure 5B). Migration positions of standards
are displayed to the left of the gel images, and sizes are indicated in kDa. The unprocessed
membrane images are displayed in Source data 1. **(B)** The Toc159 bands in the eluted fractions in the
two experiments, in (A) and in Figure 5B, were quantified. Error bars indicate standard deviation
from the mean, and open red circles indicate individual data points.

Figure 5 – Supplement 4

All three YFP-SUMO probes are conjugation-competent.

The membrane shown is the same as the one presented in Figure 5B (YFP panel). In this case, the
membrane was visualised using a long exposure to aid the detection of weakly abundant protein
bands. As well as the YFP-SUMO monomers, additional high molecular bands and smears were seen
in the IP samples, indicating the capture of large numbers of SUMOylated proteins. This shows that
all three YFP-SUMO probes could be successfully conjugated, as expected (Ayaydin and Dasso,
2004). The positions of the YFP-SUMO monomers (lower pair of arrows), and of the various SUMO
adducts (vertical bar) potentially including di-SUMO (upper pair of arrows), are indicated. Note that
YFP-SUMO3 migrated more slowly than the other two fusions, as expected. The asterisks indicate
non-specific bands. Migration positions of standards are displayed to the left of the gel images, and
sizes are indicated in kDa.

Table 1

Bioinformatic analysis predicts that the core TOC proteins in *Arabidopsis* contain SUMOylation
sites and SUMO interaction motifs.

The GPS-SUMO algorithm was applied to the amino acid sequences of Toc159, Toc132, Toc120,
Toc90, Toc75, Toc33 and Toc34 using the 'high stringency' setting, and the results generated are
shown in columns 2, 3 and 4. 'Consensus' sites fall within canonical SUMO site motifs: ψ-K-X-E
(where ψ indicates a hydrophobic amino acid, and X indicates any amino acid residue). 'Non-
consensus' sites do not fall within canonical SUMO site motifs; analysis shows that ~40% of
SUMOylation may occur at non-consensus sites (Zhao et al., 2014). 'SUMO Interaction' sites are
predicted to mediate the non-covalent interaction between proteins and SUMO peptides. The JASSA
algorithm was also applied to the amino acid sequences using the 'high cut-off' setting (see column
5). aa denotes amino acids.

Protein name	Position (aa)	*p*-value	Type	Also predicted by JASSA (high stringency)?
atToc159 (At4g02510) 1503 aa	95	0.021	Consensus	No
	106	0.034	Consensus	Yes
	126	0.032	Consensus	Yes
	144	0.02	Consensus	Yes
	151	0.02	Consensus	No
	246-250	0.005	SUMO Interaction	Yes
	408-412	0.009	SUMO Interaction	Yes
	486-490	0.009	SUMO Interaction	Yes
	498	0.022	Consensus	No
	502	0.049	Non-consensus	No
	539	0.026	Consensus	No
	1300	0.049	Non-consensus	No
	1370	0.01	Consensus	Yes
atToc132 (At2g16640) 1206 aa	30	0.006	Consensus	Yes
	66	0.036	Consensus	Yes
	352	0.002	Consensus	No
	895	0.005	Consensus	No
	1077	0.014	Consensus	No
atToc120 (At3g16620) 1084 aa	52	0.008	Consensus	No
	57	0.031	Consensus	No
	209	0.05	Non-consensus	No
	777	0.006	Consensus	No
	959	0.013	Consensus	No
atToc90 (At5g20300) 793 aa	191	0.027	Consensus	Yes
	481	0.017	Consensus	No
	711	0.02	Consensus	No
	786	0.042	Non-consensus	Yes

atToc75 (At3g46740) 818 aa	434	0.049	Non-consensus	No
	513	0.029	Consensus	No
atToc33 (At1g02280) 297 aa	291	0.044	Non-consensus	No
atToc34 (At5g05000) 313 aa	290	0.026	Consensus	No
	298	0.043	Non-consensus	Yes

919 **Table 2**

920 **Primers used during the course of this study.**

921 (A) Primers used in restriction cloning procedures.

Primer name	Sequence*	Used to generate...
SCE1 F (HindIII)	AA<u>AAGCTT</u>ATGGCTAGTGGAATCGCTC	pSAT4(A)-nEYFP-N1 SCE1
SCE1 R (EcoRI)	AA<u>GAATTC</u>GACAAGAGCAGGATACTGCTTG	pSAT4(A)-nEYFP-N1 SCE1
Toc159-5 F (EcoRI)	AA<u>GAATTC</u>AATGGACTCAAAGTCGGTT	pSAT4-cEYFP-C1-B Toc159
Toc159-3 R (SalI)	AA<u>GTCGAC</u>TTAGTACATGCTGTACTT	pSAT4-cEYFP-C1-B Toc159
Toc132-5 F (XhoI)	AA<u>CTCGAGC</u>TATG GGAGATGGGACTGAG	pSAT4-cEYFP-C1-B Toc132
Toc132-3 R (SmaI)	AA<u>CCCGGG</u>TCATTGTCCATATTGCGT	pSAT4-cEYFP-C1-B Toc132
Toc33-5 F (HindIII)	AG<u>AAGCTT</u>CGATGGGGTCTCTCGTTCGT	pSAT4-cEYFP-C1-B Toc33
Toc33-3 R (XbaI)	AA<u>TCTAGA</u>TTAAAGTGGCTTTCCACT	pSAT4-cEYFP-C1-B Toc33
Toc34-5 F (HindIII)	AG<u>AAGCTT</u>CGATGGCAGCTTTGCAAACG	pSAT4-cEYFP-C1-B Toc34
Toc34-3 R (XbaI)	AA<u>TCTAGA</u>TCAAGACCTTCGACTTGC	pSAT4-cEYFP-C1-B Toc34
OEP7 F (XhoI)	<u>CTCGAG</u>ATGGGAAAAACTTCGGGA	pSAT4(A)-cEYFP-N1 ΔOEP7
OEP7-35 R (KpnI)	<u>GGTACCG</u>GAATTTATCGAGGAAAGG	pSAT4(A)-cEYFP-N1 ΔOEP7

924   (B) Primers used in Gateway cloning procedures.

Primer name	Sequence*	Used to generate…
SCE1 Gateway F	<u>GGGGACAAGTTTGTACAAAAAAGCAGGCTCCA</u>TGGCTAGTGGAATCGCTC	p2GWY7 SCE1
SCE1 Gateway R	<u>GGGGACCACTTTGTACAAGAAAGCTGGGTTGA</u>CAAGAGCAGGATACTGC	p2GWY7 SCE1
SUMO1 Gateway F	<u>GGGGACAAGTTTGTACAAAAAAGCAGGCTCCT</u>CTGCAAACCAGGAGGAAGACAAG	p2YGW7 SUMO1
SUMO1 Gateway R	<u>GGGGACCACTTTGTACAAGAAAGCTGGGTTTC</u>AGGCCGTAGCACCACC	p2YGWY SUMO1
SUMO2 Gateway F	<u>GGGGACAAGTTTGTACAAAAAAGCAGGCTCCT</u>CTGCTACTCCGGAAGAAGAC	p2YGW7 SUMO2
SUMO2 Gateway R	<u>GGGGACCACTTTGTACAAGAAAGCTGGGTT</u>CTAAAAGCAGAAGAGCTTCAGGCC	p2YGW7 SUMO2
SUMO3 Gateway F	<u>GGGGACAAGTTTGTACAAAAAAGCAGGCTCCT</u>CTAACCCTCAAGATGACAAGCCC	p2YGW7 SUMO3
SUMO3 Gateway R	<u>GGGGACCACTTTGTACAAGAAAGCTGGGTT</u>TTAAAGCCCATTATGATCGAAAAGC	p2YGW7 SUMO3

926   (C) Primers used in RT-PCR experiments.

Primer name	Sequence*	Used to amplify…
SUMO1 F(2)	AAAAAGCAGGCTCCACAAAAGCCACGGCCAATTAG	*SUMO1*
SUMO1 R(2)	AGAAAGCTGGGTTCCATTCATATCACACACAAGCCC	*SUMO1*
SUMO3 F	ACAGACTGGAGTTTTTGTTTC	*SUMO3*
SUMO3 R	CTCATGAGTCATTTACACACACG	*SUMO3*
eIF4E1 F	AAGATTTGAGAGGTTTCAAGCGGTGTAAG	*eIF4E1*
eIF4E1 R	AAACAATGGCGGTAGAAGACACTC	*eIF4E1*
Siz1-HindIII-F	AA<u>AAGCTT</u>ATGGATTTGGAAGCTAATTGTAAG	*SIZ1* (for *siz1-4*)
Siz1-seq-1R	TCTGCATTGTGCTTGCAC	*SIZ1* (for *siz1-4*)
Siz1-F	GGATTATCTTCCAGTAATAGGCAAG	*SIZ1* (for *siz1-5*)

| Siz1-R | CCCGACTGAGCTGAAGCATC | *SIZ1* (for *siz1-5*) |

928   (D) Primers used to genotype mutants.

Primer name	Sequence	Used to genotype…
SUMO1 F(2)	AAAAAGCAGGCTCCACAAAAGCCACGGCCAATTAG	*sum1-1*
SUMO1 R(2)	AGAAAGCTGGGTTCCATTCATATCACACACAAGCCC	*sum1-1*
SUMO2 F	CGTTGTTGGTACTTGGTTGG	*sum2-1*
SUMO2 R	CAAAACTCTAAACTGGTCGG	*sum2-1*
SUMO3 F	ACAGACTGGAGTTTTTGTTTC	*sum3-1*
SUMO3 R	CTCATGAGTCATTTACACACACG	*sum3-1*
SCE1 F	CGCCGCGAAATCTGGACC	*sce1-4*
SCE1 R	TTCCTCTCTTCAGCTAAACG	*sce1-4*
SIZ1 F(2)	GCAAACAGGGAAAGAAGCAGG	*siz1-4*
SIZ1 R(2)	CATTGAGTCTGTTTCTAGCG	*siz1-4*
LBb1	GCGTGGACCGCTTGCTGCAACT	SALK lines (left border)
LB1	GCCTTTTCAGAAATGGATAAATAGCCTTGCTTCC	SAIL lines (left border)

930   (E) Primers used to map the *siz1-4* and *siz1-5* T-DNA insertions.

Primer name	Sequence	Used to map…
Siz1-Seq-1R	TCTGCATTGTGCTTGCAC	*siz1-4* T-DNA insertion
Siz1-Seq-3R	TGACAACCACTGTATGCAGG	*siz1-5* T-DNA insertion

* Underlining indicates restriction sites (A and C), or attB1 and attB2 recombination sequences (B).

Additional files

Source data 1

Immunoblot source data.

Uncropped immunoblot images for Figures 1 and 5, and their supplements.

Transparent reporting form