Molecular analyses reveal consistent food web structure with elevation in rainforest *Drosophila* - parasitoid communities

Authors:

[1]Christopher T. Jeffs, (Joint first)
[1,2]J. Christopher D. Terry (Joint first) 0000-0002-0626-9938
[3]Megan Higgie 0000-0002-2397-0240
[4]Anna Jandová
[4]Hana Konvičková
[4,5]Joel J. Brown 0000-0002-3608-6745
[4]Chia Hua Lue
[6]Michele Schiffer 0000-0002-5203-2480
[7]Eleanor K. O'Brien 0000-0001-5145-7340
[7,8]Jon Bridle
[4]Jan Hrcek (co-last) 0000-0003-0711-6447
[1]Owen T. Lewis (co-last, corresponding author) 0000-0001-7935-6111

Affiliations:

1, Department of Zoology, University of Oxford, Oxford, UK,
2, School of Biological and Chemical Sciences, Queen Mary University of London, Mile End Road, London E1 4NS, UK
3, College of Science and Engineering, James Cook University, Townsville QLD 4811, Australia
4, Biology Centre of the Czech Academy of Sciences, Institute of Entomology, Ceske Budejovice, Czech Republic
5, University of South Bohemia, Faculty of Science, Ceske Budejovice, Czech Republic
6, Daintree Rainforest Observatory, James Cook University, Cape Tribulation, Australia.
7, School of Biological Sciences, University of Bristol, Bristol, BS8 1UD UK
8, Department of Genetics, Environment and Evolution, University College London, London, UK,

Abstract

The analysis of interaction networks across spatial environmental gradients is a powerful approach to investigate the responses of communities to global change. Using a combination of DNA metabarcoding and traditional molecular methods we built bipartite *Drosophila*-parasitoid food webs from six Australian rainforest sites across gradients spanning 850 m in elevation and 5° Celsius in mean temperature. Our cost-effective hierarchical approach to network reconstruction separated the determination of host frequencies from the detection and quantification of interactions. The food webs comprised 5-9 host and 5-11 parasitoid species at each site, and showed a lower incidence of parasitism at high elevation. Despite considerable turnover in the relative abundance of host *Drosophila* species, and contrary to some previous results, we did not detect significant changes to fundamental metrics of network structure including nestedness and specialisation with elevation. Advances in community ecology depend on data from a combination of methodological approaches. It is therefore especially valuable to develop model study systems for sets of closely-interacting species that are diverse enough to be representative, yet still amenable to field and laboratory experiments.

Keywords

DNA metabarcoding, elevational gradients, Drosophila, host-parasitoid, quantitative network

Introduction

The geographic distribution of a species is determined by its intrinsic abiotic limits and by its interactions with other species. Individual populations of species do not respond to environmental changes in isolation from the species with which they interact (Davis et al. 1998, Gilman et al. 2010, Urban et al. 2016). Furthermore, interspecific interactions themselves respond to environmental change in diverse ways (Tylianakis et al. 2008, Thierry et al. 2019, Guiden et al. 2019). To understand how species' distributions will respond to global change, it is therefore important to consider this wider community context.

One valuable approach for understanding how ecological communities will respond to environmental change is to study how food webs and other networks of interspecific interactions vary along elevational gradients (Tylianakis and Morris 2017, Pellissier et al. 2018, Baiser et al. 2019, Gravel et al. 2019). Temperature changes rapidly with elevation, so transects spanning elevations on mountain slopes can approximate aspects of climate change, while avoiding many of the confounding factors such as dispersal barriers or differences in seasonality that complicate analyses of latitudinal gradients (Hodkinson 2005, Körner 2007, Malhi et al. 2010, Morris et al. 2015). There is an expectation that predation and parasitism risk increase with temperature (Roslin et al. 2017, Libra et al. 2019), but it is unclear how such changes are expressed at the community level. Quantitative food webs should be particularly valuable in exploring changes in species interactions with changes in temperature, because they incorporate the relative abundances and interaction frequencies of component species (Memmott and Godfray 1994). The structure of communities can be summarised with network-level metrics that can reveal shifts that are not apparent in simple connectance food webs or in qualitative descriptors of diversity such as species richness (Tylianakis et al. 2007, Delmas et al. 2019).

To date, there have been relatively few network analyses along elevational transects (Pellissier et al. 2018), in part because of the substantial research effort required to document them through direct observation (for example, by observing herbivores feeding on plants, or parasites associated with hosts). These traditional approaches to describing networks are highly labour-intensive, even for a single network. Rapid, accurate and cost-effective methods to characterise interaction networks are therefore needed to assess the community-wide consequences of environmental change.

Recent developments in molecular methods offer the potential to document interaction networks with lower 'per-interaction' effort, while also addressing biases and deficiencies such as those arising from insufficient sampling. Traditional methods of food web construction are also often prone to sampling biases (Gibson et al. 2011) and can miss cryptic species and associated interactions (Derocles et al. 2015). By contrast, DNA-based approaches can be faster, more efficient and taxonomically more comprehensive, allowing the simultaneous resolution of interactions and identification of morphologically-cryptic species (Hrček and Godfray 2015). Molecular approaches to the construction of ecological networks have become increasingly common over the last decade (Clare 2014, Evans et al. 2016, Derocles et al. 2018, Roslin et al. 2019, Evans and Kitson 2020). Although high throughput, 'next generation sequencing' (NGS) methods are now routinely used to characterise bulk samples, their use for characterising trophic interactions in food webs has proved more challenging. Recently-developed nested DNA metabarcoding allows large numbers of pooled samples to be processed in a single NGS run without losing resolution to individual samples (Kitson

et al. 2019). This method has been used successfully to identify interactions involving a single species
and its consumers (Kitson et al. 2019). However, it is unclear whether such success is scalable to
more complex interaction networks.

Food webs involving insect parasitoids, whose larvae feed exclusively on, and kill, a single arthropod
host (Godfray 1994), have considerable potential for studying the processes that structure
communities as they are both experimentally tractable while retaining meaningful levels of
complexity (Hrček and Godfray 2015). Host-parasitoid interactions are amenable to molecular
identification (Hrček et al. 2011, Wirta et al. 2014, Derocles et al. 2015, Kitson et al. 2019), with the
caveat that the method describes parasitoid attack rate rather than the rate of successful parasitism
per se.

Host-parasitoid interactions are important drivers of insect population dynamics (Hassell 2000) and
climate change is expected to cause distribution shifts in parasitoids (Burrows et al. 2014), alter host-
location and utilisation, and change the balance between parasitoid virulence and the immune
response of their hosts (Hance et al. 2007). The consequences for host-parasitoid communities and
the functions and services that they mediate are likely to be pronounced (Jeffs and Lewis 2013,
Derocles et al. 2018, Thierry et al. 2019). Previous work has found increased specialisation within
host-parasitoid networks with increasing elevation (Maunsell et al. 2015). However, other sources of
information about the effect of temperature, such as latitudinal trends, do not show a consistent
trend in specialisation or other community-level properties (Morris et al. 2014, Moles and Ollerton
2016). Meta-analyses have found scale dependence of patterns (Galiana et al. 2019), but there is a
need for additional studies to assess general trends in network structure.

Here we explore how food webs might respond to predicted climate change by describing the
community of wild frugivorous vinegar flies (Diptera: Drosophilidae) and their parasitoids across two
elevational transects in Australian rainforest. Biotic interactions (O'Brien et al. 2017, 2020) and
thermal tolerance (Kellermann et al. 2012) have been shown to determine *Drosophila* species
ranges. In this study, we resolve and quantify food web composition and structure using molecular
methods. We demonstrate how nested DNA metabarcoding can be used to generate multiple
quantitative networks across landscapes, opening up opportunities to use this and similar systems as
models in community ecology research.

Methods

Study sites

The Australian Wet Tropics World Heritage Area is an area of rainforest with exceptional levels of
biodiversity (Stork et al. 2011) which lies close to Queensland's northeast coast between Cooktown
and Townsville (15-19'S, 145-146.30'E). We established study sites along two transects through the
rainforest along elevation gradients at Paluma Range Road in Paluma Range National Park and
Kirrama Range Road within Girramay National Park (Figure 1). Our study sites spanned elevations
from 59 – 916 m above sea level and mean temperatures ranged from 21°C at the higher elevation
sites to 26 °C at lower sites (SI 1). Variability in temperature was not strongly related to elevation
over this period, although the lower elevations show higher variation annually. The temperature
differences across our elevation transects are broadly consistent with temperature shifts expected
over the next 50-100 years under models of climate change (IPCC 2014). We selected seven forested
sites along each transect spanning the three 'ecologically significant climatic zones' identified by

Webb and Tracey (1981) on the basis of changes in forest structure with elevation.

Study system

We focused on the food web of interactions involving native, forest-dwelling, fruit-feeding
Drosophila species and their associated parasitoids (Hymenoptera). Well-established laboratory
practices and fundamental knowledge of *Drosophila* and their parasitoids (Kraaijeveld and Godfray
1997, Markow and O'Grady 2008, Prevost 2009) make these interacting species a valuable model
system in community ecology. In our study area, *Drosophila* assemblages associated with rotting
fruit are well characterised (Hangartner et al. 2015), although less is known about their parasitoids.

Figure 1. Location of transects in north-eastern Australia. Location of sites are coloured by the mean
temperature recorded by dataloggers at each site during the research period, and labelled by the elevation
(metres above sea level). White crosses show sites for which we constructed interaction networks. Map tiles
by Stamen Design, under CC BY 3.0. Data by OpenStreetMap, under ODbL.

Sampling Method

At each site, *Drosophila*-parasitoid communities were sampled using bottle traps baited with
fermented banana. Bottle traps consisted of 1.5 L plastic water bottles with two 8cm x 10cm
windows cut in the side to allow flies and wasps to enter, following Markow and O'Grady (2006). A
rain-shield was placed over the lid of the bottle and the line was coated with Tanglefoot® insect
barrier (The Tanglefoot Company, Grand Rapids, USA) to prevent ants compromising the traps (see
image in SI 1). Bottle traps were hung 1.5 m above the ground and separated by a minimum of 5 m.
We set the traps in shaded locations at least 5 m from a roadside edge or canopy gap.

Previous work has established that banana bait samples the native generalist *Drosophila* spp. in our
focal communities effectively (Schiffer *pers. obs.*, Carson and Okada (1983). Traps were baited with
50 g of banana bait (mashed 'Cavendish' bananas supplemented with bakers' yeast), and strips of
cardboard were added to facilitate *Drosophila* pupation. Traps were exposed during the period 11[th]
March – 12[th] April 2016, corresponding to mid-late wet season when insect activity is high.
Parasitoids of *Drosophila* attack either larvae or pupae, but all emerge from pupae. Thus, by

sampling *Drosophila* pupae we were able to document the network of host-parasitoid trophic
interactions because pupae contain information about both the host and any parasitoid infecting it.

To capture the variation in colonisation speed and pupation time of different *Drosophila* species and
across the elevation gradient, traps were open for either long (24 days), medium (14-15 days), or
short (11-12 days) exposure periods at all sites. We did not find evidence of exhaustion of baits,
indicating that competition for food was unlikely to consistently exclude components of the
community. The relative frequency of the host species likely changes through the stages of the
lifecycle through the differential impact of competition and lifespan. Our surplus-food design
therefore most closely represents the relative frequency of the natural population in the early larval
stage, before larval competition acts.

For each of the sites we sorted fly pupae from 15 randomly selected trap samples, five from each of
the three exposure-time categories. We excluded traps if they had been compromised by ants or
mammals, were exceptionally dry, or had been flooded by rain. Traps were either sorted on the day
of collection or frozen at -15°C and sorted later. Pupae were stored individually in separate wells of
96-well plates filled with 96% ethanol and stored at -15°C. A random subset of pupae from the high
and low elevation sites were reared in sealed, ventilated 96-well plates in a 22°C incubator to yield
adult fly and parasitoid material. These were identified morphologically (*Drosophila* to species level
by MS and parasitoid to genus level by C-HL) and used to build a DNA reference library (Table S2.1).

Hierarchical Network Construction Approach

Quantitative interaction networks were built for three sites from both transects representing the
highest, lowest and most central elevations. Pupa samples were randomly drawn from across the
three exposure times in proportion to the number caught in each. Quantitative networks were
generated by a hierarchical sampling strategy. First, a 'core' network was built from a random
selection of 182 pupae per site, where both host and parasitoid (if present) were identified by
molecular methods (detailed below). In this stage of network construction, 722 hosts were
successfully identified in 86-138 samples per site. Of these, a total of 179 were found to be
parasitoid positive using PCR, and in 147 cases the parasitoids were successfully identified.

Next, to optimise the number of interactions sampled from each network, all available additional
samples (total 2659, 339-564 per site) were screened for parasitoid DNA using PCR. Those found to
be parasitized were then sequenced to identify the host and parasitoid. This resulted in an additional
289 identified interactions (15-119 per network). These two sets of data were combined into an
overall 'enriched' quantitative network (Figure S2.2). This used interaction frequencies derived
from both sets of data, scaled by relative host frequencies and overall host-specific parasitism rates
from the 'core' network data. In 5 cases an observed interaction in the enriching set could not be
associated with a parasitism rate as that host was not observed to be parasitised in the core dataset
at that site. These interactions were assigned a relative frequency of 0 in the enriched network.
However, these interactions were included in the specification of the site-specific binary interaction
matrices used for the calculation of site dissimilarity and qualitative network metrics.

Molecular methods to describe network structure

We used a combination of DNA metabarcoding and classic molecular methods. We first performed
DNA metabarcoding on a selection of individually tagged pupal samples ("core" samples). By

comparing the sequences to a custom reference library, we were able to identify hosts and estimate
overall host diversity. Contrary to expectations based on previous work (Kitson et al. 2019),
metabarcoding did not allow parasitoid identification and therefore we could not use this approach
to document and quantify host-parasitoid interactions. We therefore developed a number of classic
molecular tools to obtain this missing information and allow cost-effective identification of
thousands of samples. The classic tools were: i) multiplex PCR to identify the host species, followed
by ii) Sanger sequencing of samples not identified by the multiplex, and iii) parasitoid detection and
identification using PCR and Sanger sequencing. Full details are given in Supplementary Information
2.

Patterns in *Drosophila* occurrence and parasitism rate

To measure rates of parasitism, we supplemented the data described above with additional
screening results from 350 samples from each of the remaining four sampling sites along each
transect, generating a grand total of 6780 parasitism assays. To explore trends in parasitism we fit
minimal generalised linear models (GLMs) with a quasi-binomial error distribution, using elevation
(metres a.s.l.) and transect (Kirrama or Paluma) as predictors. The significance of individual terms
was assessed using F-tests. Since host-species turnover may mask or drive underlying elevational
effects if species have different susceptibilities, we also fitted the model including *Drosophila* species
as an additional categorical predictor. This analysis was restricted to three abundant *Drosophila*
species observed at all six core sites (*D. pallidifrons, D. pseudoananassae* and *D. rubida*).

Network metrics across the elevation gradient

We calculated five standard network statistics for each site, using the *bipartite* R package (Dormann
et al. 2008): *H2'*, which measures overall level of specialisation in the network (Blüthgen et al. 2006);
weighted NODF, a metric of nestedness that corrects for network size and connectance (Almeida-
Neto et al. 2008); *quantitative modularity*, which assesses the extent to which the network is divided
into discrete compartments (Beckett 2016); *network vulnerability*, the mean number of parasitoid
species attacking each host; and *network generality*, the mean number of host species that each
parasitoid attacks. We tested for the impact of elevation on each metric with a linear model,
including the matrix size (number of hosts × number of parasitoids, following Morris et al. (2014)
and transect (Kirrama or Paluma) as additional predictors.

To assess the precision with which we can infer these metrics given our sample sizes, we used
bootstrapping of our available observations. For the enriched networks, we resampled each
sampling stage separately. Bootstrapping network in this manner provides an indication of the
degree of uncertainty in a metric value. However, because the networks can only become smaller,
the results are not unbiased and should be interpreted with caution.

Drivers of interaction turnover

Interaction β-diversity represents differences in the set of observed interactions between sites
(Poisot et al. 2012, Graham and Weinstein 2018). Whole-network dissimilarity (β_{WN}) between two
sites can be partitioned into that attributable to species not being present at both sites (β_{ST}) and
that attributable to interactions only occurring at one site despite both species being present at both
(β_{OS}). Because methods partitioning quantitative similarity measures are not yet well established,
we used the qualitative Jaccard similarity metric (Novotny 2009) and a customised version of the

betalinkr function from the *bipartite* R package that takes into account host species observed at a
site that were not observed interacting with parasitoids. We enforced $\beta_{WN} = \beta_{ST} + \beta_{OS}$ by dividing
β_{ST} and β_{OS} through by $\beta_{WN}/(\beta_{ST} + \beta_{OS})$ following Poisot et al. (2012). For additional comparison,
we calculated the Jaccard similarity between the host species found at each site (β_{Host}). For each of
these metrics, we calculated pairwise dissimilarities between each our six core sites. We used
multiple regression on distance matrices (Legendre et al. 1994) via the *ecodist* R package (Goslee
and Urban 2007) to investigate whether these dissimilarities are driven by elevational or between-
transect differences.

Results

Parasitism Rate

Overall parasitism rate declined significantly with elevation across our 14 sites. For each 100 m rise
in elevation, the probability of being parasitized decreased by 14% (coefficient estimate $\beta_{elevation}$= -
1.49 km^{-1}, se = 0.49, Pr(>F$_{1,11}$) = 0.00865). Parasitism was significantly higher overall at Paluma than
at Kirrama ($\beta_{Paluma} = 1.06$, Pr(>F$_{1, 11}$)= 0.0023). The elevational trend was largely driven by data
from the Paluma transect (Figure 2a), although the interaction term between elevation and transect
was not statistically significant (Pr(>F$_{1,10}$) = 0.135). However, the high over-dispersion in the data (the
dispersion parameter was estimated to be 14.8) makes it challenging to identify interaction terms
with confidence.

Parasitism rates of widespread *Drosophila* species at the 6 focal sites showed similar patterns (Figure
2b). Model selection did not identify any host-specific transect (Pr(>F$_{2,8}$) = 0.90) or elevation
(Pr(>F$_{2,10}$) =0.21) effects but did find a significantly greater decline in parasitism rate with elevation
at Paluma (Pr(>F$_{1,11}$) = 0.035). At Kirrama, there was only a negligible effect of elevation ($\beta_{elevation_K}$=
-0.16 km^{-1}, se: 0.85). Full statistical results and model selection procedures are given in SI 3.

Figure 2. a) The fraction of pupae from traps at each site observed to be parasitized in relation to elevation. b)
Parasitism rates at the six core sites for the three most abundant species in relation to elevation. In both
subfigures, lines show the fitted minimal models.

Network Structure

Across the six focal sites we found 12 *Drosophila* species, 15 parasitoid species (principally Figitidae,
Diapriidae and Braconidae, Table S2.3), together making up 55 different interaction pairs. The host

species found at Kirrama were a subset of those observed at Paluma. The six enriched quantitative
networks are shown in Figure 3. There was no significant relationship between any of the tested
network metrics and elevation (p>0.05 in all cases, Figure 4 & SI 3). Including network size as an
additional co-predictor did not change this. Support intervals from resampling are shown in Figure 4.

Figure 3. Quantitative interaction networks from each site, calculated using the enriching method described in
the main text. The width of the boxes corresponds to the relative frequency that each species or interaction
was observed. Networks are arranged in elevation order, with Kirrama on the left and Paluma on the right and
labelled by the elevation of the site. Networks were drawn using the *bipartiteD3* R package. Parasitoid wasps
are labelled by taxon: A= *Asobara*, B = Braconidae, C = Chalcidoidea, D= Diapriidae, G= *Ganapsis*, LM =
Leptolamina, LP = *Leptopilina*, P = Pteromalidae, U = Unidentified.

Figure 4. Consistency in network-level metrics across the elevational transect, with no statistically significant effect of elevation in all cases. Error bars show bootstrapped 95% support intervals from resampling the observations at each site. Shaded areas show 95% confidence intervals based on the elevation-only model.

Drivers of network dissimilarity

Species turnover (β_{ST}) was a more important driver of qualitative dissimilarity between networks (β_{WN}) than changes in observed interactions (β_{OS}, Figure 5a). Although dissimilarity tended to be higher for pairs of networks from locations differing greatly in elevation (Figure 5a), neither transect differences nor elevational differences significantly explained any of the components of network dissimilarity (all unadjusted p > 0.1). Overall host community composition disimilarity (β_{Host}) was not significantly related to elevational differences (p = 0.45) or transect differences (p = 0.78).

Figure 5. a) Effect of elevational difference on between-site community β-diversity measures. Whole-network interaction dissimilarity (β_{WN}) was large, and predominantly composed of species turnover (β_{ST}), although there was also a notable contribution derived from changing interactions between present species (β_{OS}).

Overall Jaccard dissimilarity in host species composition (β_{Host}) was also notable. Although all the dissimilarity
measures show an apparent increase with elevational difference, these trends were not significant. Lines show
a direct linear model through each dissimilarity metric, ignoring transect differences. **b)** Relative observation
frequency of the host species observed at each site. Host species are arranged in order of the centre of mass
of their relative abundances. The dark grey square shows where *D. pandora* was observed in the enriching
dataset, but not the core dataset, making its relative abundance indeterminable.

Discussion

Our study demonstrates the capacity of molecular methods to sample multiple interaction networks
of a model system efficiently and to shed light on the underlying ecological processes. We found
evidence for a reduction in parasitism frequency at higher elevations in line with previous results
from comparable systems (Kimura and Suwito 2015), although this was largely driven by one of our
two transects. Despite this, we found no evidence for major changes to food web network structure
along an elevation gradient that spanned a temperature range comparable to the ecological range of
many of the focal species. This contrasts with the results from previous studies of host parasitoid
food webs by Maunsell et al. (2015) and Morris et al. (2015) and suggests that any shifts in network
structure with elevation are unlikely to be consistent and that, by extension, the response of host-
parasitoid networks to climate change may not be easily predicted (Hance et al. 2007).

The lack of significant trends in network structure with elevation are consistent with the absence of
marked trends in these metrics across latitudes documented by Morris et al. (2014) and the
apparent constancy across landscapes in more focussed studies of antagonistic interaction networks
(Kemp et al. 2017, Redmond et al. 2019).

Sampling networks across gradients involves unavoidable trade-offs when
distributing sampling effort between sites, sampling periods and transects, in
order to best capture any spatio-temporal changes in network structure. The
sample size for each of our networks is still rather small, with a number of
interactions represented by only a single observation. This makes it likely that
many interactions remain unobserved (Jordano 2016). Nonetheless, the
quantitative network metrics we used are frequently found to be amongst the
most robust to moderate levels of sampling (Banašek-Richter et al. 2004,
Fründ et al. 2016, Vizentin-Bugoni et al. 2016, Henriksen et al. 2019).
With six sites, the power of our tests to detect small to moderate impacts of
elevation is inevitably limited. Nevertheless, we did not detect a trend in any of
the suite of five network structure metrics we tested. Any impact of elevation
along our transect on network structure appears to be minor compared to that
which we observed in the parasitism rate and species turnover. Identifying drivers
of community change

We found a high degree of network dissimilarity between sites at both Kirrama and Paluma, which
derived largely from species turnover (both host and parasitoid) rather than changes in the observed
interactions. Although the transects have similar temperature profiles, the vegetation structure on

the Kirrama transect had been more heavily disrupted by a cyclone five years before the sampling
period. However, we did not identify any significant differences between the transects. We did not
identify any significant effect of elevational differences on community differences, as measured by
binary Jaccard dissimilarity. This pattern of higher-level consistency mirrors that found in a number
of systems (Kaartinen and Roslin 2012, Kemp et al. 2017, Gravel et al. 2019, Redmond et al. 2019)
and suggests that there are mechanisms channelling communities towards particular structures,
despite external changes.

Although there are known shifts in relative abundance of host species in our study system with
elevation (for example, *D. birchii* has higher abundance at cool, high-elevation sites than in warmer,
low-elevation sites (O'Brien et al. 2017)), many of the host species occur across the full range of
elevations sampled, leaving only a minority of high or low elevation specialists to drive any
qualitative differences (Figure 5b). In terms of the interactions themselves, we found relatively high
levels of generality – the mean level of specificity (as measured by *H2'*) in our communities (0.42)
was notably lower than the mean of 0.65 found in a cross-latitude survey of host-parasitoid
networks (Morris et al. 2014). Although the majority of species were found across the elevation
range, their relative abundances varied considerably with elevation (Figure 5b). Differences in
network composition may be difficult to detect with the binary network dissimilarity methods we
used, and emphasise changes in comparatively rare species, which are more likely to be influenced
by sampling effects (Chao et al. 2005). This highlights the importance of recording abundance rather
than spatial range limits in determining species' responses to temperature.

To address this challenge, Staniczenko et al. (2017) proposed that changes in relative interaction
frequency could be used to link quantitative changes in interaction frequencies to environmental
changes. We attempted to use such an approach within a Bayesian framework to relate elevation to
per-capita interaction frequencies (*not shown*). However, we could not confidently identify any
strong evidence for individual interactions being modified across the gradient, even when restricting
analysis to the most widely observed interactions. Without independent external estimates of the
relative abundance of parasitoid species, analysis of changing interaction rate is severely restricted.
More fundamentally, variation in the identity and strength of links within networks along
environmental gradients may be driven by any combination of turnover in species composition,
changes in species abundances and by abiotic influences on species interactions (Pellissier et al.
2018). Moving from 'how' to 'why' is a critical challenge in network ecology, but feedbacks between
the relative abundance of each species and the interactions between them generate fundamental
circularities (Dormann et al. 2017, Moreira et al. 2018). An essential part of future work to identify
the drivers of community composition will be controlled manipulative experiments (Derocles et al.
2018). This and other *Drosophila*-parasitoid systems offer exceptional potential for field
manipulations (O'Brien et al. 2017, 2020), identification of physiological traits (Kellermann et al.
2012) and laboratory experiments (Davis et al. 1998).

Sampling strategy and molecular methods

Despite the per-sample cost of molecular methods falling significantly in recent years, identification
effort is still a finite and valuable resource. Our hierarchical sequencing approach was helpful in
increasing the number of identified interactions in our dataset for a given amount of laboratory
effort, an approach that can help to maximise scarce resources and particularly relevant in species-
rich interaction networks. One drawback is that the additional data wrangling required to construct

quantitative networks from the resulting data can complicate subsequent analyses, and in particular
to propagate estimates of uncertainty in network structure.

Our sampling strategy was only a single snapshot of the interaction network at each of our sites, and
therefore does not resolve seasonal or annual turnover in interactions (Kemp et al. 2017, Lue et al.
2018). Our approach of gathering pupae samples after a variety of exposure times enables us to
characterise a community where development time varies among species and with temperature.
However, a number of challenges remain. For example, shorter exposure periods are likely to
underestimate the abundance of host species with longer development times. Species with rapid
development times may therefore be over-represented in our sampling, although we did not detect
consistent differences in the host composition between our exposure periods. Furthermore, rather
than parasitism rate, our interactions are more accurately described as attack-rates, since it is not
necessarily the case that a parasitoid egg or larva within a host would develop successfully, kill the
host and emerge as an adult (Condon et al. 2014). Rearing to emergence is the only way to establish
the result of a parasitism event (Hrček and Godfray 2015), but this is much more labour intensive,
introduces other biases (such as survival in laboratory conditions), and would negate the advantages
of the molecular approach.

DNA metabarcoding alone was unable to describe our host-parasitoid networks without
supplementary use of conventional molecular methods. Although Kitson et al. (2019) demonstrated
the DNA metabarcoding method on a single host and parasitoid species, we were unable to obtain
network data with this method alone in our multispecies system. This was despite pure parasitoid
DNA from our reference samples amplifying well with the primers we used. The reason is probably
amplification bias to host DNA and low relative parasitoid/host DNA concentration. We estimate
that increasing sequencing coverage per sample by several orders of magnitude would be necessary
to characterise parasitoids and their links to hosts reliably. This would not currently be cost-
effective, but might become more realistic in the near future as sequencing costs fall. We
recommend using a single-step PCR with larger number of primer tags in future studies, since the
nested metabarcoding approach involving two PCRs may be more prone to mistagging. We used a
strategy for detecting mistagging developed by Kitson et al (2019) and found it to be a crucial aspect
for producing reliable and verifiable DNA metabarcoding data. We therefore recommend
incorporating mistagging controls to DNA metabarcoding runs in addition to negative and positive
controls.

Conclusion

We documented pronounced turnover of *Drosophila* species relative abundance across an elevation
and temperature gradient. A strong reduction in parasitism with elevation was not associated with a
detectable change in overall network structure. DNA metabarcoding proved useful to characterise
these previously unstudied ecological networks, but it did not prove possible to use this approach
"out of the box" to obtain network data. Rather, metabarcoding data had to be supplemented with
data from classic molecular methods and a reliable DNA reference library. Ecological communities
are highly complex and communities will have to respond to multiple changes under global change.
To understand the patterns in these networks and to distinguish the principal drivers behind them,
manipulative experiments in the field and laboratory will be needed. A focus on tractable model
systems will allow an in-depth investigation into fundamental ecological dynamics of natural
systems.

Author Contributions:

CTJ: Led fieldwork, drafting manuscript, sampling design, performed molecular work. JCDT: Led statistical analysis and writing of manuscript. MH: Contributed to funding application, supervised sampling and fieldwork. AJ: Contributed to development of molecular methods, performed molecular work, HK: Contributed to development of molecular methods, performed molecular work. JJB: Performed molecular work. CHL: Parasitoid taxonomy, built parasitoid reference library, MS: *Drosophila* taxonomy. EO-B: Contributed to funding application, sampling design and fieldwork. JB: Conceived project, secured funding, contributed to sampling design. JH: Secured funding, led molecular work, writing. OTL: Conceived project, secured funding, study design, writing. All authors contributed to editing of the final manuscript.

CTJ and JCDT are joint-first authors. JH and OTL are co-last authors.

Funding:

NERC grant: NE/N010221/1 to OTL and JB

Czech Science Foundation grant 17-27184Y to JH

Acknowledgments:

We thank Adam Bajgar, Michaela Borovanská, Eva Chocholová for help in the molecular lab; Melanie Thierry and Nicholas Pardikes for help with live flies and parasitoids for the barcoding library, Ary Hoffman for COI barcodes; Adriana Uzqueda, Jaimie Hopkins and Leah Carr for help with field collection and sample sorting; and Lexie Edwards for help with logistics. Fieldwork was conducted under permit WITK16977516 to CTJ from Queensland's Department of Environment and Heritage Protection.

Code and Data availability:

Code to replicate all analyses is available on Github at https://github.com/jcdterry/AusDrosTransect_CodeandData and if accepted will be archived on Zenodo. Raw data is also archived in a long-term data repository (EIDC https://doi.org/10.5285/85657c4c-54c9-4d02-a262-455ea1c38d95).

References

Almeida-Neto, M. et al. 2008. A consistent metric for nestedness analysis in ecological systems: reconciling concept and measurement. - Oikos 117: 1227–1239.

Baiser, B. et al. 2019. Ecogeographical rules and the macroecology of food webs. - Global Ecology and Biogeography 28: 1204–1218.

Banašek-Richter, C. et al. 2004. Sampling effects and the robustness of quantitative and qualitative food-web descriptors. - Journal of Theoretical Biology 226: 23–32.

Beckett, S. J. 2016. Improved community detection in weighted bipartite networks. - Royal Society Open Science 3: 140536.

Blüthgen, N. et al. 2006. Measuring specialization in species interaction networks. - BMC Ecology 6: 9.

Burrows, M. T. et al. 2014. Geographical limits to species-range shifts are suggested by climate velocity. - Nature 507: 492–495.

Carson, H. L. and Okada, T. 1983. Drosophilidae from banana traps over an altitudinal transect in Papua New Guinea, II. Frequency of species at eight collecting sites. - International Journal of Entomology 25: 142–151.

Chao, A. et al. 2005. A new statistical approach for assessing similarity of species composition with
incidence and abundance data. - Ecology Letters 8: 148–159.

Clare, E. L. 2014. Molecular detection of trophic interactions: Emerging trends, distinct advantages,
significant considerations and conservation applications. - Evolutionary Applications 7: 1144–1157.

Condon, M. A. et al. 2014. Lethal Interactions Between Parasites and Prey Increase Niche Diversity in a
Tropical Community. - Science 343: 1240–1244.

Davis, A. J. et al. 1998. Individualistic species responses invalidate simple physiological models of
community dynamics under global environmental change. - Journal of Animal Ecology 67: 600–612.

Delmas, E. et al. 2019. Analysing ecological networks of species interactions. - Biological Reviews 94: 16–36.

Derocles, S. A. P. et al. 2015. Determining plant - Leaf miner - parasitoid interactions: A DNA barcoding
approach. - PLoS ONE 10: 1–18.

Derocles, S. A. P. et al. 2018. Climate warming alters the structure of farmland tritrophic ecological
networks and reduces crop yield. - Molecular Ecology 27: 4931–4946.

Dormann, C. F. et al. 2008. Introducing the bipartite Package: Analysing Ecological Networks. - R News 8: 8–
11.

Dormann, C. F. et al. 2017. Identifying Causes of Patterns in Ecological Networks: Opportunities and
Limitations. - Annual Review of Ecology, Evolution, and Systematics 48: 559–584.

Evans, D. M. and Kitson, J. J. 2020. Molecular ecology as a tool for understanding pollination and other
plant–insect interactions. - Current Opinion in Insect Science 38: 26–33.

Evans, D. M. et al. 2016. Merging DNA metabarcoding and ecological network analysis to understand and
build resilient terrestrial ecosystems. - Functional Ecology 30: 1904–1916.

Fründ, J. et al. 2016. Sampling bias is a challenge for quantifying specialization and network structure:
Lessons from a quantitative niche model. - Oikos 125: 502–513.

Galiana, N. et al. 2019. The geographical variation of network structure is scale dependent: understanding
the biotic specialization of host–parasitoid networks. - Ecography 42: 1175–1187.

Gibson, R. H. et al. 2011. Sampling method influences the structure of plant-pollinator networks. - Oikos
120: 822–831.

Gilman, S. E. et al. 2010. A framework for community interactions under climate change. - Trends in
Ecology & Evolution 25: 325–331.

Godfray, HCJ. 1994. Parasitoids Behavioral and Evolutionary Ecology. - Princeton University Press.

Goslee, S. C. and Urban, D. L. 2007. The ecodist package for dissimilarity-based analysis of ecological data. -
Journal of Statistical Software 22: 1–19.

Graham, C. H. and Weinstein, B. G. 2018. Towards a predictive model of species interaction beta diversity. -
Ecology Letters 21: 1299–1310.

Gravel, D. et al. 2019. Bringing Elton and Grinnell together: a quantitative framework to represent the
biogeography of ecological interaction networks. - Ecography 42: 401–415.

Guiden, P. W. et al. 2019. Predator–Prey Interactions in the Anthropocene: Reconciling Multiple Aspects of
Novelty. - Trends in Ecology & Evolution 34: 616–627.

Hance, T. et al. 2007. Impact of Extreme Temperatures on Parasitoids in a Climate Change Perspective. -
Annual Review of Entomology 52: 107–126.

Hangartner, S. B. et al. 2015. A collection of Australian Drosophila datasets on climate adaptation and
species distributions. - Scientific Data 2: 150067.

Hassell, M. P. 2000. The Spatial and Temporal Dynamics of Host-Parasitoid Interactions. - Oxford University
Press.

Henriksen, M. v. et al. 2019. The effect of network size and sampling completeness in depauperate
networks (E Thébault, Ed.). - Journal of Animal Ecology 88: 211–222.

Hodkinson, I. D. 2005. Terrestrial insects along elevation gradients: Species and community responses to
altitude. - Biological Reviews of the Cambridge Philosophical Society 80: 489–513.

Hrček, J. and Godfray, H. C. J. 2015. What do molecular methods bring to host–parasitoid food webs? -
Trends in Parasitology 31: 30–35.

Hrček, J. et al. 2011. Molecular detection of trophic links in a complex insect host-parasitoid food web. -
Molecular Ecology Resources 11: 786–794.

IPCC 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth
Assessment Report of the Intergovernmental Panel on Climate Change.

Jeffs, C. T. and Lewis, O. T. 2013. Effects of climate warming on host-parasitoid interactions. - Ecological
Entomology 38: 209–218.

Jordano, P. 2016. Sampling networks of ecological interactions. - Functional Ecology 30: 1883–1893.

Kaartinen, R. and Roslin, T. 2012. High temporal consistency in quantitative food web structure in the face
of extreme species turnover. - Oikos 121: 1771–1782.

Kellermann, V. et al. 2012. Upper thermal limits of Drosophila are linked to species distributions and
strongly constrained phylogenetically. - Proceedings of the National Academy of Sciences 109: 16228–
16233.

Kemp, J. E. et al. 2017. Invariant antagonistic network structure despite high spatial and temporal turnover
of interactions. - Ecography 40: 1315–1324.

Kimura, M. T. and Suwito, A. 2015. Altitudinal patterns of abundances and parasitism in frugivorous
drosophilids in west Java, Indonesia. - Journal of Natural History 49: 1627–1639.

Kitson, J. J. N. et al. 2019. Detecting host–parasitoid interactions in an invasive Lepidopteran using nested
tagging DNA metabarcoding. - Molecular Ecology 28: 471–483.

Körner, C. 2007. The use of "altitude" in ecological research. - Trends in Ecology and Evolution 22: 569–574.

Kraaijeveld, A. R. and Godfray, H. C. J. 1997. Trade-off between parasitoid resistance and larval competitive
ability in Drosophila melanogaster. - Nature 389: 259–261.

Legendre, P. et al. 1994. Modeling Brain Evolution from Behavior: A Permutational Regression Approach. -
Evolution 48: 1487.

Libra, M. et al. 2019. Elevational contrast in predation and parasitism risk to caterpillars in a tropical
rainforest. - Entomologia Experimentalis et Applicata: eea.12851.

Lue, C. H. et al. 2018. Geographic and seasonal variation in species diversity and community composition of
frugivorous drosophila (Diptera: Drosophilidae) and their Leptopilina (Hymenoptera: Figitidae)
Parasitoids. - Environmental Entomology 47: 1096–1106.

Malhi, Y. et al. 2010. Introduction: Elevation gradients in the tropics: Laboratories for ecosystem ecology
and global change research. - Global Change Biology 16: 3171–3175.

Markow, T. A. and O'Grady, P. 2006. Drosophila: a Guide to Species Identification and Use. - Academic
Press (Elsevier).

Markow, T. A. and O'Grady, P. 2008. Reproductive ecology of Drosophila. - Functional Ecology 22: 747–759.

Maunsell, S. C. et al. 2015. Changes in host-parasitoid food web structure with elevation (T Webb, Ed.). -
Journal of Animal Ecology 84: 353–363.

Memmott, J. and Godfray, H. C. J. 1994. The use and construction of parasitoid webs. - In: Hawkins, B. A.
and Sheehan, W. (eds), Parasitoid Community Ecology. Oxford University Press, pp. 300– 318.

Moles, A. T. and Ollerton, J. 2016. Is the notion that species interactions are stronger and more specialized
in the tropics a zombie idea? - Biotropica 48: 141–145.

Moreira, X. et al. 2018. Elevational gradients in plant defences and insect herbivory: recent advances in the
field and prospects for future research. - Ecography 41: 1485–1496.

Morris, R. J. et al. 2014. Antagonistic interaction networks are structured independently of latitude and
host guild. - Ecology Letters 17: 340–349.

Morris, R. J. et al. 2015. Food web structure changes with elevation but not rainforest stratum. - Ecography
38: 792–802.

Novotny, V. 2009. Beta diversity of plant-insect food webs in tropical forests: A conceptual framework. -
Insect Conservation and Diversity 2: 5–9.

O'Brien, E. K. et al. 2017. Testing for local adaptation and evolutionary potential along altitudinal gradients
in rainforest Drosophila: beyond laboratory estimates. - Global Change Biology 23: 1847–1860.

O'Brien, E. K. O. et al. 2020. Fitness effects of competition within and between species change across
species' ranges, and reveal limited local adaptation in rainforest Drosophila. - bioRxiv doi:
10.1101/395624

Pellissier, L. et al. 2018. Comparing species interaction networks along environmental gradients. - Biological
Reviews 93: 785–800.

Poisot, T. et al. 2012. The dissimilarity of species interaction networks (F Jordan, Ed.). - Ecology Letters 15:
1353–1361.

Prevost, G. 2009. Parasitoids of Drosophila. - Academic Press.

Redmond, C. M. et al. 2019. High specialization and limited structural change in plant-herbivore networks
along a successional chronosequence in tropical montane forest. - Ecography 42: 162–172.

Roslin, T. et al. 2017. Higher predation risk for insect prey at low latitudes and elevations. - Science 356:
742–744.

Roslin, T. et al. 2019. Introduction: Special issue on species interactions, ecological networks and
community dynamics – Untangling the entangled bank using molecular techniques. - Molecular
Ecology 28: 157–164.

Staniczenko, P. P. A. et al. 2017. Predicting the effect of habitat modification on networks of interacting
species. - Nature Communications 8: 792.

Stork, N. E. et al. 2011. Status and Threats in the Dynamic Landscapes of Northern Australia's Tropical
Rainforest Biodiversity Hotspot: The Wet Tropics. - In: Zachos, F. E. and Habel, J. C. (eds), Biodiversity
Hotspots. Springer Berlin Heidelberg, pp. 311–332.

Thierry, M. et al. 2019. Mechanisms structuring host–parasitoid networks in a global warming context: a
review. - Ecological Entomology: een.12750.

Tylianakis, J. M. and Morris, R. J. 2017. Ecological Networks Across Environmental Gradients. - Annual
Review of Ecology, Evolution, and Systematics 48: 25–48.

Tylianakis, J. M. et al. 2007. Habitat modification alters the structure of tropical host-parasitoid food webs.
- Nature 445: 202–5.

Tylianakis, J. M. et al. 2008. Global change and species interactions in terrestrial ecosystems. - Ecology
Letters 11: 1351–1363.

Urban, M. C. et al. 2016. Improving the forecast for biodiversity under climate change. - Science in press.

Vizentin-Bugoni, J. et al. 2016. Influences of sampling effort on detected patterns and structuring processes
of a Neotropical plant-hummingbird network. - Journal of Animal Ecology 85: 262–272.

Webb, L. J. and Tracey, J. G. 1981. Australian rainforests: patterns and change. - In: Keast, A. (ed), Ecological
biogeography of Australia. W Junk, pp. 605–694.

Wirta, H. K. et al. 2014. Complementary molecular information changes our perception of food web
structure. - Proceedings of the National Academy of Sciences 111: 1885–1890.

595