Cellular events of acute, resolving or progressive COVID-19 in SARS-CoV-2 infected non-human primates

Fahlberg MD[1], Blair RV[1,2], Doyle-Meyers LA[1], Midkiff CC[1], Zenere G[1], Russell-Lodrigue KE[1], Monjure CJ[1,2], Haupt EH[1], Penney TP[1], Lehmicke G[1], Threeton BM[1], Golden N[1], Datta PK[1,2], Roy CJ[1,2], Bohm RP[1,2], Maness NJ[1,2], Fischer T[1,2], Rappaport J[1,2], and Vaccari M[1,2]*.

[1]Tulane National Primate Research Center, Covington, LA
[2]Tulane University School of Medicine, New Orleans, LA

*To whom correspondence should be addressed:

Monica Vaccari, Ph.D.

Associate Professor, Tulane National Primate Research Center,

18703 Three Rivers Road, Covington, LA 70433

Email: mvaccari@ tulane.edu

Abstract

We investigated the immune events following SARS-CoV-2 infection, from the acute inflammatory state up to four weeks post infection, in non-human primates (NHP) with heterogeneous pulmonary pathology. The acute phase was characterized by a robust and rapid migration of monocytes expressing CD16 from the blood and concomitant increase in CD16+ macrophages in the lungs. We identified two subsets of interstitial macrophages (HLA-DR+ CD206–), a transitional CD11c+ CD16+ cell population that was directly associated with IL-6 levels in plasma, and one long lasting CD11b+ CD16+ cell population. Strikingly, levels of monocytes were a correlate of viral replication in bronchial brushes and we discovered TARC (CCL17) as a new potential mediator of myeloid recruitment to the lungs. Worse disease outcomes were associated with high levels of cell infiltration in lungs including CD11b+ CD16hi macrophages and CD11b+ neutrophils. Accumulation of macrophages was long-lasting and detectable even in animals with mild or no signs of disease. Interestingly, animals with anti-inflammatory responses including high IL-10:IL-6 and kynurenine to tryptophan ratios had less signs of disease. Our results unravel cellular mechanisms of COVID-19 and suggest that NHP may be appropriate models to test immune therapies.

The COVID-19 pandemic is a major public health crisis, causing a global medical emergency. There are currently no effective vaccines or drugs approved to completely treat or prevent COVID-19, and much remains unknown about the pathogenesis of the disease. COVID-19 presents with highly variable outcomes, and patients that develop symptoms exhibit a wide range of disease, spanning from mild (fever, cough, shortness of breath) to severe (dyspnea,

pneumonia, rapidly progressing radiographic changes)[1]. It is currently estimated that up to 15%

of COVID-19 patients progress to acute respiratory distress syndrome (ARDS)[2], which is the

major cause of death among fatal SARS-CoV-2 cases. Studies elucidating molecular and

immunological details of SARS-CoV-2 infection are underway, and many of the reported

clinical observations point to the disease being at least partly the result of an excessive host

response aimed to clear the virus but instead contributing to disease development[3]. This

hypothesis is supported by studies on the closely related coronavirus SARS-CoV-1[4-6], and by

clinical reports showing elevated levels of the proinflammatory cytokine IL-6 in SARS-CoV-2

infected patients, and profound dysregulation of the myeloid cell compartment, particularly

among those with severe symptoms[3,7-9].

Interestingly, the increase in IL-6 coincides with the upregulation of chemokines responsible for

myelopoiesis and monocyte recruitment to the lungs[3,10] suggesting that peripheral inflammatory

monocytes and tissue macrophages may play a role in the cytokine storm seen in severe COVID-

19 patients[11]. The underlying inflammatory state and increased myelopoiesis seen in older

individuals may be responsible for the high mortality rate seen in this group[12]. A better

understanding of immunopathology of COVID-19 will aid the development of targeted

immunotherapies and effective vaccines.

Non-human primates (NHP) have been used as animal models to study acute respiratory diseases

such as SARS and MERS[13]. In these studies, African green monkeys (AGM) have been shown to

be more susceptible to severe disease than rhesus macaques (RM)[14]. Regardless of species

differences, and given the overlap between NHP and with human immune systems, both NHP

species have been useful models to study host responses to coronaviruses[5]. We and others have

shown that both AGM and RM can be successfully infected with SARS-CoV-2 via the intranasal

and the intratracheal route, or by aerosol[15-18]. Following exposure, AGM and RM have high

levels of virus replication in the respiratory tract for up to two weeks, and both species develop

pulmonary pathology characteristic of COVID-19, that varies from very mild to severe[19]. Other

groups have described the very early immune events in SARS-CoV-2 infected rhesus macaques,

with perturbation of monocytes populations, increased macrophages in lungs, and cellular

activation in blood all happening a few days following the infection[16].

Here, we extend on these findings by describing the kinetics of immune events following

infection, spanning from the acute phase up to four weeks following infection, in the blood and

the lungs of four African green monkeys (AGM) and four rhesus macaques (RM) with a

heterogeneous spectrum of COVID-19. Overall, we observed a multiphasic mobilization of

innate and adaptive cells from the blood to the lung that is strikingly common to both species.

The acute inflammatory phase was characterized by an increase in the frequency of monocytes

that was positively associated with bronchial viral replication. CD11b+ macrophages and

neutrophils were detected in the lungs of all infected animals, however, with greater numbers in

severe disease. Consistent with the resolution phase of inflammation in the lung[20], the second

phase was characterized by a switch toward type 2 responses in both species. We identified two

populations of interstitial macrophages (HLA-DR+ CD206–) in BAL; 1) CD11c+ CD16^{hi}

population increasing during the acute phase of infection that was associated with IL-6 levels in

the plasma, and 2) a long lasting CD11b+ and CD16^{hi} population accumulating in lungs of both

AGM and RM, particularly in those with increased disease severity. Animals with more

pronounced clinical signs had a higher level of infiltrates in lungs when compared with animals

with no or mild signs, lower IL-10:IL-6 and kynurenine /tryptophan ratios (anti-inflammatory:

pro-inflammatory) cytokines in plasma[21]. By investigating temporal and spatial changes in

immune cells our study links together findings in both NHP and human patients and unravels

new cellular events of COVID-19[22,23]. Further, we show that NHP are highly relevant models to

study COVID-19 immunopathology and suggest that they may be appropriate models to test

immune therapies and vaccines for SARS-CoV-2.

Results
104
Increased levels of monocytes and chemokines after infection is associated with viral load

and disease severity. Four adult RM and four adult AGM were exposed to SARS-CoV-2 by

either aerosol or mucosal challenge including buccal, intranasal, intratracheal, and conjunctival

(multi-route) exposure (**Supplementary Table 1**). The study concluded at 4 weeks post

exposure (**Fig. 1A**)[18]. All eight animals had detectable viral load by RT-PCR on nasal and

pharyngeal swabs and bronchial brushes by week 1 (**Fig. 1B**, **Supplementary Table 2**). Viral

RNA was also detected in pharyngeal, nasal, buccal, rectal and vaginal swabs. No differences in

viral load (VL) were observed between the different routes of exposure or between the two

species at week 1 (**Supplementary Table 2 and Fig. 1B**). Of note, by week 2 AGM had higher

VL than RM in the bronchial brushes and by week 3 VL in RM was mostly undetectable while it

remained above detection levels in AGM until study completion (**Fig. 1B** and **Supplementary**

Table 2). Similar to humans, the course of disease varied widely across animals, as detailed in

Table 1. Of the four AGM enrolled, two animals developed severe respiratory signs and were

euthanized on days 8 (NC34) and 22 (NC33) post-infection, following established endpoint

criteria for NHP study protocols, and as we previously reported[18]. Of the two remaining AGM,

NC38 had multifocal mild to moderate interstitial pneumonia scattered throughout all lung lobes

and NC40 was mostly asymptomatic and had scant inflammation in all lung lobes at necropsy.

Three of the four RM (GH99, HD09, and FR04) developed pneumonia characterized by a

granulomatous to pyogranulomatous inflammatory response. This was severe in GH99, mild in

HD09, and minimal in FR04. Histopathology of the fourth RM (HB37) revealed a lymphocytic

vasculitis and proliferative vasculopathy in the right middle lung lobe.

Clinical signs observed during the course of the study included mild tachypnea (day 11) and

cough (day 15) in GH99 (**Supplementary Table 3**). Intermittent mild to moderate pyrexia was

observed in HB37 (day 1-21 post infection). Radiographic changes included a nodule in the right

caudal lung lobe (day 11) in GH99, and HD09 also developed a mild, focal increased opacity in

the ventral lung field at day 11 (data not shown).Viral loads in the bronchial alveolar brushes at

all time points, temperature, pulse oxygen levels (SpO$_2$) and ranking did not differ between

female (n = 3) and males (n = 5) or between animals exposed to different routes of infection

(multi route n = 4 and aerosol n = 4), possibly due to the small number of animals per group

(**Supplementary Table 2 - 4**).

Viral load was not significantly associated with pulmonary pathology, however we observed an

inverse association between viral load at week 1 and disease severity ranking (1 to 8 from most

severe to least severe) based on the above observations and histopathology findings (AGM: R =

0.8 ; RM: R = –0.4, all animals: P = 0.057, R = – 0.71, by the Spearman correlation test) (**Fig.**

1C, **Supplementary Fig. 1A** and **1B** (**Supplementary Table 5**).

Interestingly, virus-driven immunological changes were remarkably similar between species at

this time (**Fig. 1** and **Supplementary Fig. 1**). To investigate overall changes driven by viral

replication in both groups, we combined their immunological analysis, however at the same time

we maintained different specifications for each species (teal = RM and magenta = AGM). Data

and statistics are also shown separately per group in **Supplementary Figure 1**. At 1 week after

infection both AGM and RM had an increase in the absolute number and frequency of

monocytes in the blood (P = 0.065, and P = 0.02 respectively, by the Tukey multiple comparison

test, n = 8) compared to baseline levels (**Fig. 1D** and **1E**). This change was significant in AGM

alone (absolute number: P = 0.0006, n = 4), and was also observed in all RM, particularly GH99

(**Supplementary Fig. 1C - 1F**). Of note, the increase in absolute number of monocytes remain

significant when GH99 was removed from the statistical analysis (P = 0.00002, **Supplementary**

Fig. 1G). The absolute number of neutrophils in blood did not change over time

(**Supplementary Fig. 1H**). Interestingly, the absolute number of monocytes at week 1 was

strongly positively associated with the viral load levels in the bronchial brush samples (P =

0.002, R = 0.93 by the Spearman test) when all the animals were combined (**Fig. 1F**) (AGM: R =

0.8; R: R = 1, not significant in separated groups, **Supplementary Fig. 1I** and **1J**), and remained

significant when GH99 was removed from the statistical analysis (**Supplementary Fig. 1K**).

We then used the CD14 and CD16 markers to discriminate between three circulating monocytes

subtypes: classical (CD14hi CD16low), intermediate (CD14hi CD16hi) and non-classical

(CD14low CD16hi) within the total (CD45, live , HLA-DR+ CD16hi) monocyte population by

flow cytometry (**Fig. 1G - 1I** and **Supplementary Fig. 2**). Subsets were gated as shown in

Supplementary Figure 2A, using a strategy that excludes CD16+ NK cells that are negative for

HLA-DR[24]. At week 1, the frequency of classical monocytes was significantly increased in

blood, as seen in humans[25], and the non-classical population was profoundly depleted (increased

levels baseline versus week 1: classical P = 0.007 and decreased non-classical P = 0.005 in all

animals combined, n = 4) (**Fig. 1H** and **1I**, and **Supplementary Fig. 2B – 2E**). As we observed

for the frequency, the absolute counts of classical and non-classical monocytes at week 1 were

also significantly increased and decreased, respectively, compared to the baseline values

(**Supplementary Figure 2F - G**). Interestingly, pre infection levels were never completely

restored even after 3 weeks post-infection (week 1 vs. week 3: classical P = 0.03 and non-

classical P = 0.03), as the number of circulating CD16hi monocytes increased overtime (non-

classical week 1 versus week 3: P = 0.0019). These results suggest emerging myelopoiesis and

rapid and robust recruitment of patrolling monocytes in tissues following infection in both

species. We did not observe changes in the frequency or number of intermediate monocytes

(**Supplementary Fig. 2H** and **2I**). In line with what we observed for the total monocyte count, at

week 1 the counts for the subsets of monocytes were positively associated with viral replication

detected at the bronchial alveolar sites (classical: R = 0.98; P = 0.06, non-classical: R = 0.96; P =

0.03 and intermediate R = 0.94; P = 0.06, by the Pearson test), further suggesting that the robust

myelopoiesis is therefore driven by the virus in the lung. Within the total monocytes, the

frequency of non-classical monocytes was significantly associated with lower viral loads

(classical: R = -0.9867; P = 0.0133, by the Pearson Test, adjusted for multiple comparison P =

0.07). It is important to note that none of these analyses were significant when the values were

adjusted for multiple comparisons (**Supplementary Table 6**).

Recent studies in humans have described an increase in the frequency of CD14 positive cells

expressing low levels of HLA-DR in COVID-19 patients. These cells phenotypically resemble

suppressive subsets of myeloid derived suppressor cells (MDSCs)[8] thus they may decrease

disease driven inflammation. We and others have previously identified MDSCs in blood of

monkeys by flow cytometry as HLA-DR[low] CD14[hi] cells within the CD45/live population[26,27].

Some of the animals in our study showed increased MDSCs frequency at week 3 compared to

baseline levels, however these changes were not significant (**Supplementary Fig. 2L**). We

measured Arginase 1 activity as a functional assay to assess the function of immunosuppressive

myeloid cells at the same time points (baseline and week 3) and we did not observe significant

changes in plasma overtime (**Supplementary Fig. 2M**)[28].

We analyzed a group of eight chemokines known to be involved in the mobilization of innate

cells, including monocytes, from the blood to the lung: IP-10 (CXCL-10), MCP-1 (CCL2),

MCP-4 (CCL13), Eotaxin, TARC (CCL17), MDC (CCL22), MIP-1α and MIP-1β

(**Supplementary Fig 3A - 3H**). Some of the chemokines measured at baseline in plasma differed

between the two species (MCP-4, Eotaxin, MIP-1b and MDC) as shown by the principle

component analysis in **Supplementary Fig. 3I** (**Supplementary Tables 7 - 9**). Plasma levels of

TARC (CCL17), a ligand for CCR4 that is involved in Th2 cell recruitment[29], were strongly

associated with the frequency of monocytes in blood (P = 0.003, R = 0.95, Spearman test,

adjusted for multiple comparison: P = 0.02) (**Fig. 1J**). Similarly, TARC/CCL17 was also

associated with viral replication in BAL at week 1 (**Fig. 1K** and **Supplementary Fig. 3J** and

3K) (P = 0.02, R = 0.857, adjusted for multiple comparisons: P = 0.2, and R = 0.927 and P=

0.016 when GH99 is excluded from the analysis), as was IP-10 (P = 0.05, R = 0.78, adjusted for

multiple comparisons) when all animals were grouped together (**Fig. 1K** and **Supplementary**

Fig. 3L), a ligand for CXCR3 that is associated with disease severity and predicts progression of

COVID-19 in humans[30]. To gain a better understanding of a common milieu possibly driving

monocyte migration observed in both groups we analyzed changes in the levels of these markers

with respect to the baseline (**Fig. 1L** and **Supplementary Table 8**). Overall this analysis

revealed that animals with the most severe disease outcomes had higher levels of chemokines at

week 1, including IP-10, TARC (CCL17).

214
Myelomonocytic cells infiltrate BAL and are associated with IL-6. Influx of macrophages has

been observed in the lung of severe cases of COVID-19 in humans[31]. To monitor cell trafficking

into the lungs, we collected bronchoalveolar lavage (BAL) at baseline and after each week

following infection, up to week 3. As expected, the majority of the cells found in BAL at

baseline were alveolar macrophages (AM)[32]. These cells are readily identified by the mannose

receptor C type 1 (CD206) marker, express high levels of HLA-DR, and are positive for the

scavenger receptor CD163 (**Fig. 2, Supplementary Fig. 4A** and **4B**), we observed a decrease in

the proportion of AM at week 1 post infection in animals of both species, corresponding with an

influx of myeloid HLA-DR+ CD206– CD163– cells (**Fig. 2A** and **2B**, **Supplementary Fig. 4C**).

Levels of AM similar to pre-infection were restored at week 2 in all but one animal (GH99), and

a concomitant increase of CD86 mean fluorescent intensity (MFI) in this population was

observed suggesting an M1 activation state[33] (**Supplementary Fig. 4D**). An increase was also

seen in the frequency of CD206+ CD163–that have been previously described as endothelial

cells which facilitate lymphocyte trafficking and they are detectable in BAL during inflammatory

events (**Fig. 2C**)[34].

T-distributed stochastic neighbor embedding (tSNE) analysis revealed changes in the phenotype

of CD16, CD11b, and CD11c positive populations most prominently at 1-2 weeks post infection

(**Fig. 2D - 2F**). In contrast to what we observed in the blood, there was an increase of CD16+

HLA-DR+ cells in BAL over time (**Fig. 2G**, and **Supplementary Fig. 4E** and **4F**). We observed

a transient increase in the level of CD11c+ CD16+ myelomonocytic cells, which function has

been described as patrolling, and they participate in rapid tissue invasion and monocyte survival

(baseline vs. week 1: P = 0.015) (**Fig. 2H**)[35,36]. The frequency of CD11c+ macrophages in BAL

at week 3 was strongly associated with the levels of the proinflammatory IL-6 cytokine in the

plasma at the same time point (week 3: P = 0.003, R = 0.96, Spearman test) (**Supplementary**

Fig. 5). The CD11b marker is required for monocyte migration to sites of inflammation[37], and is

highly expressed on myeloid cells recruited from the blood and differentiating into

macrophages[38]. We also observed an increase in the percentage of CD11b+ CD16+ HLA-DR+

CD206– myelomonocytic cells in BAL at week 2 and 3 following infection (**Fig. 2I**), a

population that is involved in regulating lung inflammatory response[20,33,39,40]. The gating strategy

for the two myelomonocytic cells populations is shown in **Supplementary Figure 4G**.

Interstitial macrophages accumulate in lungs of infected animals. Lungs were collected for

all animals at the end of the study, corresponding to 4 weeks post infection (1 week following the

last BAL collection shown in Figure 2), with the exception of NC33 and NC34. These two

animals had higher levels of IL-6 at necropsy compared to the other six monkeys, and NC34 had

symptoms similar to ARDS[18] (**Fig. 3**). Overall, infiltration of macrophages was observed in the

lungs of SARS-CoV-2 infected AGM and RM at necropsy and was more robust in animals with

severe disease (**Fig. 3A**). We further characterized macrophages population in the lung by

staining with the AM marker CD206 (in blue) and with CD11b integrin (in green) and CD16

(red) (**Fig. 3B**). Low numbers of macrophages were seen in the lungs of a non-infected control

RM, scattered throughout the interstitium and alveolar spaces, as shown by the arrows. In the

control this population was bi-morphic and composed of CD11b negative CD16+ CD206–

(arrow) in the alveolar septa or CD16+ CD206+ cells within the alveolar spaces (arrowhead)

(**Fig. 3B**). In all animals that were infected we could detect CD11b+ CD16+ macrophages. FR04

(RM ranking = 7) lungs had macrophages scattered throughout the interstitium and alveolar

spaces. The phenotypic population was heterogeneous and contained low numbers of CD11b+

CD16+ CD206– macrophages. The most severe RM, GH99 (ranking = 3), had large numbers of

macrophages multifocally surrounding small airways denoted by the asterisks. Of note this

animal had the higher absolute number of monocytes in the blood at week 1 (**Supplementary**

Fig. 1D). Low numbers of CD11b+ CD16+ CD206– cells were scattered throughout the

predominant population of CD11b+ CD16+ CD206+ macrophages. Macrophages were also

scattered throughout the interstitium and alveolar spaces. NC34 and NC33 (AGM ranking = 1

and 2 respectively) had multifocal to coalescing areas of hemorrhage (NC33 indicated by the

arrows), and pulmonary parenchyma infiltrated by multifocal aggregates of macrophages (NC34)

with infiltration by low to moderate numbers of macrophages. The phenotypic population was

heterogenous and composed of moderate numbers of CD11b+ CD16+ CD206– macrophages.

Aggregates were determined by staining animals FR04, GH99, and NC34 with CD68/CD163

(red) and CD3: (blue). Animal FR04 (RM ranking = 7) had isolated CD3+ cells scattered

throughout the lung. GH99 (RM ranking = 3) had cell aggregate with mixed populations of

CD3+ and CD68/CD163+ cells in isolated area of inflammation. NC34 (AGM ranking = 1) had

rare, small aggregates of CD3+ cells (**Supplementary Figure 6**). Altogether these findings

present a picture of long-lasting lung inflammation that, albeit more pronounced in animals with

severe signs of disease, is also present in animals with no detectable viral load and mild or no

signs of disease after 4 weeks from the start of infection.

Neutrophils accumulate in animals with severe disease. In this study we did not observe

changes in the absolute number of neutrophils in blood during the course of the infection

(**Supplementary Figure 1H**). Interestingly, studies in monkeys and humans showed increased

levels of neutrophils and suggested deregulation of their maturation, with CD11b being one of

the markers defining pre-neutrophils[8,9,41-43]. We stained lungs from animals NC33, NC34, GH99,

FR04 and NC40 with anti-myeloperoxidase (MPO) antibody for neutrophils and with anti-

CD11b antibody, and we included two controls, one RM and one AGM (**Figure 4A**). MPO+

cells[44] were elevated in animals with severe disease (NC33, NC34). GH99 had areas with

neutrophil accumulation and the remaining animals were low. Controls had very few neutrophils

in the lungs (**Figure 4B**). CD11b+ MPO+ neutrophils were only present in infected animals and

were also high in NC33 (**Figure 4C**). Accordingly, neutrophils and lymphocytes could be seen in

animals with severe disease but not in animals with moderate disease (**Supplementary Fig. 7**).

NC34 (necropsy day 8 post infection; rank = 1, severe) had lymphocytic aggregates that were

perivascular and intermixed with neutrophils and histiocytes. In NC38 (necropsy day 28, rank 5,

moderate) lymphocytic aggregates are scattered within the alveolar septa and predominately

mixed with histiocytes and rare neutrophils.

NKG2a+ cells and PD-1+ T cells decrease in blood and increase in BAL overtime.

Infiltration of lymphocytes was observed in the lungs of SARS-CoV-2 infected AGM and RM at

necropsy and was more robust in animals with severe disease (**Fig. 5A**). We first analyzed the

kinetics of innate NK cells subsets, that we defined as HLA-DR– CD3– cells expressing the

NKG2a receptors, which are known to traffic to the lung[45] (**Fig. 5B - D**, and **Supplementary**

Fig. 8). Interestingly, the frequencies of NKG2a+ cells significantly decreased in peripheral

blood of both AGM and RM over the course of infection (baseline vs. week 3, P = 0.03) (**Fig.**

5B). Consistent with the pattern described for the myeloid population, we also observed a

concomitant increase in the frequencies of NKG2a+ cells in BAL (baseline vs. week 3, P = 0.05)

(**Fig. 5C** and **5D**, **Supplementary Fig. 8A - 8C**). Four out of four AGM had decreased NKG2a+

cells in peripheral blood, coinciding with an increase of the same cells in BAL. While the pattern

of NKG2a+ cells in peripheral blood of RM was not statistically significant, nonetheless in this

species there was a marked increase of those cells in the BAL. These results suggest that the loss

of NKG2a+ cells in the PBMCs and the gain of NKG2a+ cells in the BAL may coincide with an

egress of these cells from peripheral blood to tissues. Accordingly, the frequency of NKG2a+

cells expressing CXCR3, a marker associated with NK cell migration to the lung[46] also

significantly decreased in blood over time (baseline vs. week 3, P = 0.03, **Fig. 5E**). Despite this

change in lymphocyte phenotype, the absolute number of lymphocytes did not change over the

first 2 weeks of infection (**Fig. 5F**) however it is important to note that all 3 remaining AGM had

an increase at week 3 (**Supplementary Fig. 8D – 8E**).

tSNE analysis revealed an increase in the level of PD-1 at weeks 2 and 3 post-infection,

overlapping with spatial regions corresponding to both CD4+ T and CD8+ T cells (**Fig. 5G and**

Supplementary Fig. 9A). In fact, the frequency of PD-1+ CD4+ T cells was elevated in the

blood of both RM and AGM (**Fig. 5H**, and **5I**, respectively). During the first week following the

infection we observed an increase in IP-10 and IFN-γ (**Supplementary Fig. 9B**), however we

also observed a switch toward Th2 type responses characterized by the increase of Th2 cytokines

IL-5 and IL-13 over time (**Supplementary Fig. 9B**). Accordingly, a gradual increase of Th2 type

cells (defined as CXCR3– CCR6 – CD4+ T cells)[24] was seen in the blood within the PD-1

positive population (**Fig. 5J**). The increase of PD-1+ Th2 cells was significantly and positively

associated with IL-4 cytokine levels (P = 0.016, R = 0.94, Spearman test) (**Supplementary Fig.**

9C). Moreover, we observed an increased level of PD-1+ CD8+ T cells in blood and BAL (BAL:

baseline vs. week 2, P = 0.015; baseline vs. week 3, P = 0.03) (data not shown and **Fig. 5K**).

Animals divided by species are shown in **Supplementary Fig. 9D** and **9E**. Associations between

viral replication and the frequency of PD1+ CD4 T cells in blood and PD1+ CD8 T cells in BAL

at week 3 were found (**Supplementary Fig. 9G- I),** however, they were most likely driven by

the AGM but not RM which had no detectable viral replication in bronchial brushes at that

specific time point.

The ratio between the anti-inflammatory and proinflammatory responses is associated with

disease progression. The pro-inflammatory cytokine IL-6 has been described as a strong

predictor to disease progression in humans[3,7]. We measured IL-6 and IL-10 in the plasma of all

the animals (**Fig. 6A**). While IL-6 levels alone did not associate with disease severity in NHPs,

the ratio of IL-10 to IL-6 levels measured in plasma associated with both the ranking of disease

severity and with the pathological score that was obtained by analyzing the extension of the

inflammation and edema of the lungs (R = 0.76 P = 0.037; and R = –0.76, P = 0.03, by the

Spearman correlation test, all animals) (**Fig. 6B** and **C**). To investigate further, we also measured

the levels of tryptophan and kynurenine in plasma at baseline, week 1, week 3 and at necropsy

(**Supplementary Figure 10**). Kynurenine was elevated and significantly increased at week 3

compared to baseline levels (P= 0.013 adj.). The kynurenine (Kyn)/ tryptophan (Tryp) ratio has

been used as a measurement of the indoleamine 2,3-dioxygenase (IDO) activity and is often

associated with T regulatory cell function [21]. In line with the observed association for the IL-

10:IL-6 ratio, the Kyn /Tryp ratio was also negatively correlated with the pathology score (1 =

mild – 18 = severe) suggesting that animals with higher levels of immune suppression had less

inflammation in the lung and better disease outcome (**Figure 6D**).

These findings show a complex series of immune events in the lung of that are most likely

initiated as a host response to the virus and then evolving overtime toward less antiviral

responses, as summarized in **Figure 6E.**

DISCUSSION
356
There is an urgent need to develop safe and effective strategies to prevent and treat COVID-19.

The development of such strategies requires a clear understanding of the immune cells involved

in COVID-19 and their unique role in pathogenesis. Clinical observations in patients and few

studies on non-human primates have started to shed light on the immune aspects involved in

COVID-19[8,9,11,22,47,48]. Here, we further characterized immune events following SARS-CoV-2

infection in two NHPs models recapitulating crucial aspects of COVID-19 lung disease[16,18].

Altogether our findings provide the first evidence in both AGM and RM of a multiphasic

immune response characterized by 1) an initial inflammatory phase with monocyte recruitment

to the lung 2) a gradual switch from a type 1 to type 2 response, and 3) a "make it or break it"

phase with either an increase in anti-inflammatory cytokines IL-10 and regulatory cell subsets

with suppressive activity, or IL-6, a pro-inflammatory cytokine associated with disease

progression in humans, as summarized in **Figure 6E**.

By examining the temporal and spatial distribution of monocytes, myelomonocytic cells and

macrophages in blood and BAL and lungs, we confirmed and extended on studies in humans and

NHPs suggesting increased myelopoiesis with recruitment of myeloid cells from the blood to

lungs during the infection that is initially driven by the virus replicating in the lung[3,16,22]. Similar

to what has been described by Wilk *et al.* and Thevarajan *et al.* in humans, we observed an

increase of classical CD14hi monocytes, and a corresponding drop in the frequency of

monocytes expressing CD16 in the peripheral blood[22,23]. Importantly, we extend on these

findings by showing a simultaneous increase in the frequency patrolling CD16hi HLA-DR+

monocytes in BAL occurring in both species and by describing the kinetics of two CD16+ HLA-

DR+ CD206– macrophages populations that preferentially populate the alveolar interstitium and

exhibit features of antigen-presenting cells[49]. CD11c+ CD16+ cells arising from intravascular

CD16hi patrolling monocytes in the acute phase and associating with IL-6 levels, and a CD11b+

macrophage population known to be involved in resolution of inflammation in the lung[20,33,39,40].

Higher frequencies of CD11b+ monocytes with increased disease severity has been observed in

the blood of humans[8]. The origin and the exact function of these interstitial macrophages remains

to be determined.

Differently than what was observed in humans and previously reported data in RM, we did not

detect an increase in the number of neutrophils, as these responses may have come up earlier

than 1 week after infection as shown by others[16]. At necropsy, CD11b+ MPO+ neutrophils were

present in the lungs of animals with severe disease as observed in humans. More studies are

needed to determine if SARS-CoV-2 results in the accumulation of immature neutrophils as seen

in humans [8,9].

392
Common to both species was the observation that animals with higher levels of myeloid and

lymphoid cell infiltration in the lung had worse disease outcomes. Several studies in both SARS-

CoV-1 and CoV-2 have reported depleted peripheral NK cell counts in severe patients[11,22,25,47].

Our data support the notion of NKG2a+ cell trafficking to the lung during SARS-CoV-2

infection, most likely mediated by IP-10 (CXCL-10), which promotes the recruitment of

CXCR3+ Th1 thymocytes and NKs[50,51]. Expression of NKG2a has been linked to both activation

and inhibition of inflammation[52]; the function in SARS-CoV-2 remains to be investigated. In the

periphery, we also observed increased levels of PD-1 on T cells. This is consistent with findings

in humans describing of an exhaustive T cell phenotype during SARS-CoV-1 and CoV-2

infection[11,53].

For the first time, we identified TARC (CCL17) as a possible mediator of myeloid recruitment in

the lung during SARS-CoV-2. Both dendritic cells and CD11b+ macrophages in the lung have

been identified as a source of CCL17 in the airways in other pulmonary inflammatory

diseases[54,55].These cells promote the recruitment of T helper type 2 (Th2) lymphocytes into the

airways[54]. Remarkably, we observed a gradual increase of the recruitment of Th2 lymphocytes

into the airway and of Th2 type cytokines such as IL-5 and IL-13. These findings show a

complex series of immune events in the lung of that are most likely initiated as a host response to

the virus, and that they may evolve into other less effective antiviral responses as either an

attempt to tissue repair/resolution of inflammation, as summarized in **Figure 6E,** or as the result

of virus direct immunomodulatory effects on host responses[6].

Intrinsic species-specific susceptibility to disease progression has been described for NHP in

SARS-CoV-1[6] with AGM being more susceptible than rhesus to severe disease. There were

unexpected similarities in the kinetics and the quality of the immune responses analyzed in RM

and AGM, particularly during the acute phase of infection, however AGM did not clear the

infection as well as RM and two AGM had most severe symptoms[56]. While the number of

animals enrolled in this study was not sufficient to look at species-specific differences and their

effect on disease progression, we found that a higher ratio of IL-10: IL-6 in plasma was

associated with disease severity. Undoubtedly, more studies are needed to prove or disprove the

single contribution of IL-10 and IL-6 in the outcome of COVID-19 disease and to identify the

source of these two potential key players[57]. The notion that dynamically balanced pro and anti-

immune responses to SASR-CoV-2 play an important role in disease progression and outcome is

supported by the association of the pathological score and the Kyn /Tryp suggesting a role of T

regulatory cells in resolution of inflammation of the lungs in COVID-19.

428
Overall, this study uncovers critical steps in the immune events of the host response to SARS-

CoV-2 and describes new correlates of viral replication and disease progression. Importantly, we

show viral driven inflammation/cell recruitment to the lung occurs rapidly, and lasts up to four

weeks post infection, which may explain in part the observed slow recovery from COVID-19

seen humans. Finally, these findings have the potential to aid the interpretation of the results of

the human trials that are currently underway for several vaccine candidates and may be

instrumental in generating new targeted therapeutic strategies aimed at resolving the severe

immune deregulation of the lung of COVID-19 patients.

Methods
439
Ethical statement on animal use and SARS-CoV-2 handling. Animals and infection. The

Institutional Animal Care and Use Committee of Tulane University reviewed and approved all

the procedures for this study. The Tulane National Primate Research Center is fully accredited

by the AAALAC. All animals were cared for in accordance with the ILAR Guide for the Care

and Use of Laboratory Animals 8th Edition. The Tulane University Institutional Biosafety

Committee approved the procedures for sample handling, inactivation, and transfer from BSL3

containment.

Animals and infection. Four adult-aged African green monkeys (Caribbean origin) and four

rhesus macaques (Indian ancestry) were exposed to SARS-CoV-2; 2019-nCoV/USA-WA1/2020

409 (MN985325.1). The virus stock was prepared in Vero E6 cells and the sequence confirmed

by PCR and/or Sanger sequencing. Plaque assays were performed in Vero E6 cells. The rhesus

macaques were from the Tulane National Primate Research Center breeding colony pathogen

free for simian type D retrovirus (SRV), macacine herpesvirus 1 (B virus), simian

immunodeficiency virus (SIV), simian T cell lymphotropic/leukemia virus (STLV), measles

virus (MV) and tuberculosis (TB). The African green monkeys were wild caught and at the

TNPRC for over a year before being assigned to this study. To mimic different possible routes of

infection in humans, animals were exposed to the virus either by aerosol (inhaled dose of 2.0 x

10^3 and 2.5 x 10^3 $TCID_{50}$) or by inoculating a cumulative dose of 3.61 x 10^6 PFU through

multiple routes (oral, nasal, intratracheal, conjunctival) (**Supplementary Table 1**).

The multi route exposure was given to four animals, one adult RM male (GH99, 14 years old),

one adult RM female (HD09, 13 years old) and two AGM, one aged male (NC40, 16 years old,

approximately) and one aged female (NC33, 16 years old, approximately). An additional two

adult males RM (FR04 and HB37, 15 and 13 years old, respectively) and one aged male and

female AGM: NC34 and NC38 (16 years old, approximately) were exposed by aerosol

(**Supplementary Table 1**). Rhesus macaques tend to live approximately 20 years in captivity,

while African green monkeys live approximately 15 years[58]. The age of the macaques is

generally multiplied by three when compared to humans, and GH99 had signs of aged lungs that

were not associated with COVID-19.

Cell count and CO₂ and SpO₂ pressure measurements. Total white blood cell counts, white

blood cell differentials, red blood cell counts, platelet counts, hematocrit values, total

hemoglobin concentrations, mean cell volumes, mean corpuscular volumes, and mean

corpuscular hemoglobin concentrations were analyzed from blood collected in tubes containing

EDTA using a SYSMEX XN-1000v Hematology Analyzer in the Level 2 Lab and a Sysmex XS-

1000 in the Level 3 Lab. CO_2 measurements were taken using the Piccolo Express Chemistry

Analyzer in the Level 3 lab and is part of the Metabolic Comprehensive Panel. SpO_2 values were

obtained using the Masimo Rad-57 Handheld Pulse Oximeter (**Supplementary Table 3**).

Sample collection and processing. Blood EDTA was collected by venipuncture and bronchial

brush and BAL by bronchoscopy. For the latter, the bronchoscope is introduced into the trachea

and directed into one of the bronchi. The bronchoscope is advanced to the point where the

diameter of the bronchoscope approximates the dimensions of the bronchus. The bronchial brush

is passed through the channel of the bronchoscope. The brush is gently advanced until the lumen

of the bronchus or bronchioles approximates the dimensions of the brush. The brush is then be

gently advanced back and forth against the mucosa to gather cells. The brush is removed from

the channel of the bronchoscope and the brush is placed into 200 μL of DNA/RNA Shield X1

(Cat.# R1200, Zymo Research, Irvine, CA). The bronchoscope remains in place while two

aliquots of 20 ml of saline each are instilled into the bronchus and subsequently aspirated. This

procedure is then repeated in the opposite lung.

Blood was layered over Ficoll-Plaque Plus (GE Healthcare17144003) and centrifuged for 30

minutes in cap-locked adaptors. Differential migration of cells during centrifugation results in

the formation of layers containing different cell types and allows collection of peripheral blood

mononuclear cells (PBMCs) from other populations. PBMCs were collected, counted and

aliquoted in tubes or resuspended in freezing media containing Fetal Bovine Serum and DMSO.

For bronchoalveolar lavage (BAL), cells and supernatant were separated by centrifugation (15

minutes at 2,000 x g at RT) in cap-locked adaptors and cells were lysed in Ammonium Chloride

Potassium (ACK) lysing buffer at room temperature for 5 minutes (Gibco# A1049201) washed

in 2% FBS, and resuspended in PBS and counted under laminar flux hood in BCL3. Fresh cells

were stained for flow cytometry.

BAL and PBMCs staining. Freshly obtained PBMCs and cells obtained from BAL were stained

with appropriate antibody cocktails and incubated for 30 minutes at room temperature.

Antibodies are listed in **Supplementary Table 10.** Cells were fixed with 4% or 2% PFA-BSA

overnight. Samples were run on a FACSAria Fusion flow cytometer (Becton Dickinson) within a

Biosafety Cabinet, and each sample was run for at least five minutes. For all panels, the

LIVE/DEAD Fixable Aqua Dead cell Stain kit was used (ThermoFisher #L34957). Monocytes

subsets were gated as described in **Supplementary Figure 2A** and as previously described[24].

Alveolar macrophages were gated based on the size and the expression of HLA-DR, CD206 and

CD163 markers. Myeloid infiltrates were gated as cells negative for CD206 and CD163 and

positive for HLA-DR, CD16 and CD11b. The gating strategy is shown in **Supplementary**

Figure 4G. NKG2a+ cells were gated on the CD3 negative parent population as shown (BAL

FR04 week 2) in **Supplementary Figure 8A.** Th1, Th2, and Th17 cells were gated as previously

described[24]. tSNE analyses were performed in FlowJo using the opt-SNE algorithm and plots

were created in R using the ggplot2 package[59].

Histopathology and scoring. Tissue samples were fixed in Z-fix (Anatech), embedded in

paraffin and 5 μm thick sections were cut, adhered to charged glass slides and stained with

hematoxylin and eosin. All slides were scanned on a Zeiss Axio Scan.Z1 digital slide scanner.

Images were acquired using HALO software (Indica Labs). Quantification of pathologic lesions

was performed with HALO software. Machine learning (Random Forest) algorithms were

trained by a board -certified pathologist to recognize fluidic and cellular inflammation in the

lungs of the RM and AGM. Scores were assigned to fluidic and cellular inflammation,

separately, based on the percentage of lung affected. Fluidic and cellular inflammation scores

were summated for all lobes and animals were ranked from 1 (highest sum) to 8 (lowest sum).

526
Plasma cytokines and chemokines. Plasma was collected by spinning and was thawed before

use. Cytokines were measured using 497 Mesoscale Discovery. IFN-γ, IL-4, IL-6, IL-8, IL-10

and IL-13 were part of the V-Plex Proinflammatory Panel 1, 10-Plex. V-plex. The chemokine

panel (#K15049D, Mesoscale Discovery, Rockville, Maryland) was also used, following the

instructions of the kit. The plate was read on a MESO Quickplex 500 SQ120 machine.]

Heatmaps were generated using the 'pheatmap' package in R[60,61]. Data were normalized by

dividing raw values at week 1 by baseline values for each animal, followed by the application of

$\log_2$. Values below the limit of detection were replaced with the lowest limit of detection value

based on the standard curve for each run, or with the lowest value detected during the run,

whichever was smaller. Bubble plots were generated using the 'ggplot2' package in R, using the

same normalized data shown in the heatmap[59]. Scatter plots were drawn using raw data points

and display Pearson's correlation coefficients and a 95% confidence interval.

RNA isolation. Swab and bronchial brush samples were collected in 200 μL of

DNA/RNA Shield X1 (Cat.# R1200, Zymo Research, Irvine, CA) and extracted for Viral RNA

(vRNA) using the Quick-RNAViral kit (Cat.# R1034/5, Zymo Research). The Viral RNA

Buffer dispensed directly to the swab in the DNA/RNA Shield. The swab was directly inserted

into the spin column. The vRNA was eluted (45μL) and 5 μL was added in a 0.1 mL fast 96-well

optical microtiter plate (Cat.# 4346906, ThermoFisher, CA).

Viral RNA detection. RT-qPCR reaction TaqPath1-Step MultiplexMaster Mix was used (Cat.#

A28527, ThermoFisher) along with the 2019-nCoV RUO Kit (Cat.# 10006713, IDTDNA,

Coralville, IA) targeting the N1 amplicon of N gene of SARS2-nCoV19 (accession MN908947).

The master mix was added to the microtiter plates covered with optical film (cat. #4311971;

ThermoFisher) and then was vortexed and pulse centrifuged. The RT-qPCR program consisted

of incubation at 25°C for 2 minutes, RT incubation at 50°C for 15 minutes, and an enzyme

activation at 95°C for 2 minutes followed by 40 cycles of denaturing step at 95°C for 3 seconds

and annealing at 60°C for 30 seconds. Fluorescence signals were detected with an Applied

Biosystems QuantStudio6 Sequence Detector. Data were captured and analyzed with Sequence

Detector Software v1.3 (Applied Biosystems, Foster City, CA). Viral copy numbers were

calculated by plotting Cq Values obtained from unknown (i.e. test) samples against a standard

curve representing known viral copy numbers. The limit of detection of the assay was 10 copies

per reaction volume. A 2019-nCoVpositive control (Cat.#10006625, IDTDNA) were analyzed in

parallel with every set of test samples. A non-template control (NTC) was also included.

Immunohistochemistry. Five μm sections of Formalin-fixed, paraffin-embedded lung were

mounted on charged glass slides, baked overnight at 56°C and passed through Xylene, graded

ethanol, and double distilled water to remove paraffin and rehydrate tissue sections. A

microwave was used for heat induced epitope retrieval, which was performed sequentially with

high and low pH solutions from Vector Labs (H-3301 and H-3300). CD16+ and MPO+ cells

were detected using MACH3 AP kits (Biocare Medical M3M532 or M3R533) and permanent

red substrate (Dako K0640). MPO+ cells CD3+ cells CD11b+ cells CD68+/CD163+ cells, and

CD206+ cells were detected using antibodies described in **Supplementary Table 10**. Slides

were incubated with a blocking buffer comprised of either 1% normal donkey serum or 10%

normal goat serum for 40 minutes. All primary antibodies were incubated for 60 minutes and all

secondary antibodies for 40 minutes at room temperature, with thorough washing in between.

DAPI (4',6-diamidino-2-phenylindole) was used to label the nuclei of each section. Slides were

mounted using a homemade anti-quenching mounting media containing Mowiol (Calbiochem

#475904) and DABCO (Sigma #D2522) and imaged with a Zeiss Axio Slide Scanner.

Arginase Activity. Arginase activity was analyzed on Plasma using the Arginase Activity Assay

Kit (MAK112, Sigma-Aldrich, St. Louis, MO) following the manufacturer instructions. Briefly,

samples were thawed on ice and, in order to deplete the urea, 40 μl of plasma were loaded in an

Amicon®Ultra 10K centrifugal filter (UFC501096 EMD Millipore), diluted with pure water to

500 μl, and centrifuged at 13,000 × g for 30 min at 4°C. Following centrifugation, the eluted

solution was discarded. Filtered samples were then diluted with pure water to 500 μl, and

centrifuged at 13,000 × g for 30 min at 4°C. At the end of centrifugation, the remaining volume

of each sample was measured, and ultra-pure water was added to reach a final volume of 40 μl.

Each sample was loaded into 2 wells of a 96-well plate (20 μl/well), representing the sample well

and the sample blank well, and 20 μl/well of ultra-pure water were added to each well. Together

with samples, the plate was loaded with urea standard and water as positive and negative

controls, respectively. Samples were loaded in singlicate, whereas controls were loaded in

duplicate. Ten microliters of 5X substrate buffer, composed of Arginine Buffer and Mn Solution,

were added to the wells except for sample blank wells, and they were incubated for 120 min at

37°C. Following the incubation, 200 μl of Urea Reagent, composed of Reagents A and B, was

added to each well to stop the reaction. Finally, 10 μl of 5X Substrate Buffer was added to the

sample blank wells to have the same reagents proportion of reagents as the Sample Wells. After

mixing, the plate was incubated for 60 min at RT, and finally acquired with microplate

spectrophotometer Power Wave XS2 (BioTek Instruments, Winooski, VT) to measure the

absorbance at 430 nm (A_{430}) of each well. The arginase activity was determined per the

following equation: [(sample well) - (blank)/(urea- blank)] X [(1mM x50x 103)/40ul x

120min)], according to the manufacture instructions.

Kynurenine and Tryptophan Plasma Levels. Tryptophan and Kynurenine plasma

concentrations were measured by using the Tryptophan ELISA (Rocky Mountain Diagnostics,

Colorado Springs, CO, USA, Catalog #BA E-2700) and Kynurenine ELISA commercial kits

(Rocky Mountain Diagnostics, Colorado Springs, CO, USA, Catalog #BA E-2200). For

tryptophan measurement, 20 μl of plasma were precipitated, the recovered supernatants were

derivatized, and the product was used to perform the ELISA according to manufacturer

instructions. For kynurenine assay, 10 μl of plasma were acylated and used to perform the

ELISA according to manufacturer instructions. The data are presented as the ratio between

kynurenine and tryptophan (Krn/Try) levels.

Statistical analysis. Correlation analyses were performed using the non-parametric Spearman

rank correlation method (two-tailed, 95% confidence) using exact permutation P values when

ranks were used, or Peason's correlation test when both variables were continuous. Analysis of

variance (ANOVA) was used for comparing statistical differences between multiple groups,

followed by Dunn's multiple comparison tests. Kruskal-Wallis ANOVA or Mann-Whitney tests

were used. All statistical analyses were performed using GraphPad Prism (version 8.4.2

GraphPad Software, La Jolla California USA) and R software (URL http://www.R-project.org/).

Correlation plots were created using the 'Performance Analytics' and 'corrr' plots, respectively,

in R[61-63].

Acknowledgments

We would like to thank Dr. Joseph (JC) Mudd and Dr. Cariappa Annaiah for critical reading of
the manuscript and Dr. Amitinder (Miti) Kaur, Megan Varnado, Kaitlin Didier from the TNPRC
Flow Cytometry Core for their help. We thank Angela Birnbaum for reviewing and optimizing
all technical SOPs and overseeing the safety of this study.
We would also like to thank Fast Grant Funding for COVID-19 Science for partially funding this
work and the NIH for supporting this work through the TNPRC base grant (P51 OD011104 59).
628

629   **Competing Interests statement**
630   **All data will be made available upon acceptance.**

631   These authors declare no competing interests.

632
Author Contributions
MDF analyzed data composed figures and wrote the manuscript, BRV was the lead pathologist and
LADM was the project veterinarian, contributed to study design, and writing of the IACUC and
manuscript. CCM designed IHC panels and performed the IHC staining, GZ wrote the manuscript, KERL
was a project veterinarian, made clinical assessments and collected samples, CJM processed and analyzed
samples for RT-qPCR, contributed; EHH provided the mesoscale data, TPP stained the samples with GL
and BMT. GL contributed to study design, provided administrative support, and aided with sample
processing and archiving. NG contributed to study design, study coordination, sample processing, and
SOP development. PKD processed and analyzed viral load data. CJR conceived and performed aerosol

experiments. RPB contributed to study design, analysis of clinical and imaging results, and writing the manuscripts. NJM performed staining of BAL and contributed to writing the manuscript. TF contributed to the animal study design and planning and contributed to writing the manuscript. JR designed and supported the animal study and helped with the writing of the manuscript. MV conceived and supported the study, run experiments, analyzed data and wrote the manuscript.

REFERENCES

1. Guan, W.J.*, et al.* Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* **382**, 1708-1720 (2020).
2. Sun, P.*, et al.* Clinical characteristics of hospitalized patients with SARS-CoV-2 infection: A single arm meta-analysis. *J Med Virol* **92**, 612-617 (2020).
3. Blanco-Melo, D.*, et al.* Imbalanced Host Response to SARS-CoV-2 Drives Development of COVID-19. *Cell* **181**, 1036-1045.e1039 (2020).
4. Weiss, S.R. Forty years with coronaviruses. *J Exp Med* **217**(2020).
5. Clay, C.*, et al.* Primary severe acute respiratory syndrome coronavirus infection limits replication but not lung inflammation upon homologous rechallenge. *J Virol* **86**, 4234-4244 (2012).
6. Graham, R.L., Donaldson, E.F. & Baric, R.S. A decade after SARS: strategies for controlling emerging coronaviruses. *Nat Rev Microbiol* **11**, 836-848 (2013).
7. Ruan, Q., Yang, K., Wang, W., Jiang, L. & Song, J. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive Care Med* **46**, 846-848 (2020).
8. Silvin, A.*, et al.* Elevated Calprotectin and Abnormal Myeloid Cell Subsets Discriminate Severe from Mild COVID-19. *Cell* (2020).
9. Schulte-Schrepping, J.*, et al.* Severe COVID-19 Is Marked by a Dysregulated Myeloid Cell Compartment. *Cell* (2020).
10. Xiong, Y.*, et al.* Transcriptomic characteristics of bronchoalveolar lavage fluid and peripheral blood mononuclear cells in COVID-19 patients. *Emerg Microbes Infect* **9**, 761-770 (2020).
11. Zhou, Y.*, et al.* Pathogenic T cells and inflammatory monocytes incite inflammatory storm in severe COVID-19 patients. *National Science Review* (2020).
12. Crooke, S.N., Ovsyannikova, I.G., Poland, G.A. & Kennedy, R.B. Immunosenescence and human vaccine immune responses. *Immun Ageing* **16**, 25 (2019).
13. Gretebeck, L.M. & Subbarao, K. Animal models for SARS and MERS coronaviruses. *Curr Opin Virol* **13**, 123-129 (2015).
14. McAuliffe, J.*, et al.* Replication of SARS coronavirus administered into the respiratory tract of African Green, rhesus and cynomolgus monkeys. *Virology* **330**, 8-15 (2004).
15. Chandrashekar, A.*, et al.* SARS-CoV-2 infection protects against rechallenge in rhesus macaques. *Science* (2020).
16. Singh, D.K.*, et al.* SARS-CoV-2 infection leads to acute infection with dynamic cellular and inflammatory flux in the lung that varies across nonhuman primate species. *BioRxiv* (2020).
17. Yu, J.*, et al.* DNA vaccine protection against SARS-CoV-2 in rhesus macaques. *Science* (2020).

18.   Blair, R.V*., et al.* ARDS and Cytokine Storm in SARS-CoV-2 Infected Caribbean
Vervets. *BioRxiv* (2020).
19.   Munster, V.J*., et al.* Respiratory disease in rhesus macaques inoculated with SARS-CoV-
2. *Nature* (2020).
20.   Aggarwal, N.R., King, L.S. & D'Alessio, F.R. Diverse macrophage populations mediate
acute lung inflammation and resolution. *Am J Physiol Lung Cell Mol Physiol* **306**, L709-
725 (2014).
21.   Fallarino, F*., et al.* The combined effects of tryptophan starvation and tryptophan
catabolites down-regulate T cell receptor zeta-chain and induce a regulatory phenotype in
naive T cells. *J Immunol* **176**, 6752-6761 (2006).
22.   Wilk, A.J*., et al.* A single-cell atlas of the peripheral immune response in patients with
severe COVID-19. *Nat Med* (2020).
23.   Thevarajan, I*., et al.* Breadth of concomitant immune responses prior to patient recovery:
a case report of non-severe COVID-19. *Nat Med* **26**, 453-455 (2020).
24.   Vaccari, M*., et al.* HIV vaccine candidate activation of hypoxia and the inflammasome in
CD14. *Nat Med* **24**, 847-856 (2018).
25.   Wen, W*., et al.* Immune cell profiling of COVID-19 patients in the recovery stage by
single-cell sequencing. *Cell Discov* **6**, 31 (2020).
26.   Vaccari, M*., et al.* Myeloid Cell Crosstalk Regulates the Efficacy of the
DNA/ALVAC/gp120 HIV Vaccine Candidate. *Front Immunol* **10**, 1072 (2019).
27.   Lin, A*., et al.* Rhesus Macaque Myeloid-Derived Suppressor Cells Demonstrate T Cell
Inhibitory Functions and Are Transiently Increased after Vaccination. *J Immunol* **200**,
286-294 (2018).
28.   Bronte, V., Serafini, P., Mazzoni, A., Segal, D.M. & Zanovello, P. L-arginine metabolism
in myeloid cells controls T-lymphocyte functions. *Trends Immunol* **24**, 302-306 (2003).
29.   Anderson, C.A*., et al.* A degradatory fate for CCR4 suggests a primary role in Th2
inflammation. *J Leukoc Biol* **107**, 455-466 (2020).
30.   Yang, Y*., et al.* Plasma IP-10 and MCP-3 levels are highly associated with disease
severity and predict the progression of COVID-19. *J Allergy Clin Immunol* (2020).
31.   Pence, B.D. Severe COVID-19 and aging: are monocytes the key? *Geroscience* (2020).
32.   Cai, Y*., et al.* In vivo characterization of alveolar and interstitial lung macrophages in
rhesus macaques: implications for understanding lung disease in humans. *J Immunol* **192**,
2821-2829 (2014).
33.   Johnston, L.K., Rims, C.R., Gill, S.E., McGuire, J.K. & Manicone, A.M. Pulmonary
macrophage subpopulations in the induction and resolution of acute lung injury. *Am J*
Respir Cell Mol Biol **47**, 417-426 (2012).
34.   Irjala, H*., et al.* Mannose receptor is a novel ligand for L-selectin and mediates
lymphocyte binding to lymphatic endothelium. *J Exp Med* **194**, 1033-1042 (2001).
35.   Auffray, C*., et al.* Monitoring of blood vessels and tissues by a population of monocytes
with patrolling behavior. *Science* **317**, 666-670 (2007).
36.   Landsman, L*., et al.* CX3CR1 is required for monocyte homeostasis and atherogenesis by
promoting cell survival. *Blood* **113**, 963-972 (2009).
37.   Prieto, J., Eklund, A. & Patarroyo, M. Regulated expression of integrins and other
adhesion molecules during differentiation of monocytes into macrophages. *Cell Immunol*
156, 191-211 (1994).

38.   Tam, J.W., Kullas, A.L., Mena, P., Bliska, J.B. & van der Velden, A.W. CD11b+ Ly6Chi
Ly6G- immature myeloid cells recruited in response to Salmonella enterica serovar
Typhimurium infection exhibit protective and immunosuppressive properties. *Infect*
Immun **82**, 2606-2614 (2014).
39.   Liao, M.*, et al.* Single-cell landscape of bronchoalveolar immune cells in patients with
COVID-19. *Nat Med* **26**, 842-844 (2020).
40.   Steinmüller, C., Franke-Ullmann, G., Lohmann-Matthes, M.L. & Emmendörffer, A.
Local activation of nonspecific defense against a respiratory model infection by
application of interferon-gamma: comparison between rat alveolar and interstitial lung
macrophages. *Am J Respir Cell Mol Biol* **22**, 481-490 (2000).
41.   Scapini, P., Marini, O., Tecchio, C. & Cassatella, M.A. Human neutrophils in the saga of
cellular heterogeneity: insights and open questions. *Immunol Rev* **273**, 48-60 (2016).
42.   Kwok, I.*, et al.* Combinatorial Single-Cell Analyses of Granulocyte-Monocyte Progenitor
Heterogeneity Reveals an Early Uni-potent Neutrophil Progenitor. *Immunity* **53**, 303-
318.e305 (2020).
43.   Ng, L.G., Ostuni, R. & Hidalgo, A. Heterogeneity of neutrophils. *Nat Rev Immunol* **19**,
255-265 (2019).
44.   Reumaux, D., de Boer, M., Meijer, A.B., Duthilleul, P. & Roos, D. Expression of
myeloperoxidase (MPO) by neutrophils is necessary for their activation by anti-
neutrophil cytoplasm autoantibodies (ANCA) against MPO. *J Leukoc Biol* **73**, 841-849
(2003).
45.   Cong, J. & Wei, H. Natural Killer Cells in the Lungs. *Front Immunol* **10**, 1416 (2019).
46.   Carlin, L.E., Hemann, E.A., Zacharias, Z.R., Heusel, J.W. & Legge, K.L. Natural Killer
Cell Recruitment to the Lung During Influenza A Virus Infection Is Dependent on
CXCR3, CCR5, and Virus Exposure Dose. *Front Immunol* **9**, 781 (2018).
47.   National Research Project for SARS, B.G. The involvement of natural killer cells in the
pathogenesis of severe acute respiratory syndrome. *Am J Clin Pathol* **121**, 507-511
(2004).
48.   Kuri-Cervantes, L.*, et al.* Immunologic perturbations in severe COVID-19/SARS-CoV-2
infection. *bioRxiv* (2020).
49.   Schyns, J.*, et al.* Non-classical tissue monocytes and two functionally distinct populations
of interstitial macrophages populate the mouse lung. *Nat Commun* **10**, 3964 (2019).
50.   Jiang, D.*, et al.* Regulation of pulmonary fibrosis by chemokine receptor CXCR3. *J Clin*
Invest **114**, 291-299 (2004).
51.   Agostini, C.*, et al.* Cxcr3 and its ligand CXCL10 are expressed by inflammatory cells
infiltrating lung allografts and mediate chemotaxis of T cells at sites of rejection. *Am J*
Pathol **158**, 1703-1711 (2001).
52.   Creelan, B.C. & Antonia, S.J. The NKG2A immune checkpoint - a new direction in
cancer immunotherapy. *Nat Rev Clin Oncol* **16**, 277-278 (2019).
53.   Li, C.K.*, et al.* T cell responses to whole SARS coronavirus in humans. *J Immunol* **181**,
5490-5500 (2008).
54.   Medoff, B.D.*, et al.* CD11b+ myeloid cells are the key mediators of Th2 cell homing into
the airway in allergic inflammation. *J Immunol* **182**, 623-635 (2009).
55.   Alferink, J.*, et al.* Compartmentalized production of CCL17 in vivo: strong inducibility in
peripheral dendritic cells contrasts selective absence from the spleen. *J Exp Med* **197**,
585-599 (2003).

56.   Woolsey, C*., et al.* Establishment of an African green monkey model for COVID-19.
780   *bioRxiv* (2020).
781   57.   Han, H*., et al.* Profiling serum cytokines in COVID-19 patients reveals IL-6 and IL-10
782   are disease severity predictors. *Emerg Microbes Infect* **9**, 1123-1130 (2020).
783   58.   Simmons, H.A. Age-Associated Pathology in Rhesus Macaques (Macaca mulatta). *Vet
784   Pathol* **53**, 399-416 (2016).
785   59.   Wickham, H. ggplot2: Elegant Graphics for Data Analysis. (Springer-
786   Verlag, New   York, 2016).
787   60.   Kolde, R. pheatmap: Pretty Heatmaps (2018).
788   61.   Team, R.C. R: A language and environment for statistical computing. . (Foundation for
789   Statistical Computing, Vienna, Austria, 2019).
790   62.   Peterson, B.G. & Carl, P. PerformanceAnalytics: Econometric Tools for Performance
791   and Risk Analysis.   . (2020).
792   63.   Kuhn, M., Jackson, S. & Cimentada, J. corrr: Correlations in R. (2020).

793
794
795   **Figure Legend**
796
Figure 1. Monocytes are a correlate of bronchoalveolar viral replication. (**A**) Schematic of

study design. Arrows represent time of inoculation with SARS-CoV-2 (day 0) or sacrifice (up to

28 days post infection). Unless stated otherwise, rhesus macaque data are shown in teal, and

African green monkey data are shown in magenta. (**B**) SARS-CoV-2 RNA levels in bronchial

brushes over time. (**C**) Association between SARS-CoV-2 load in bronchial brushes and ranking

of disease severity in all eight animals (most severe case = 1; mildest case = 8). (**D**) Changes in

monocyte count in blood before and after infection. (**E**) Proportion of total monocytes (HLA-

DR+ CD16+ and/or CD14+ out of CD45+ leukocytes) in PBMCs over time (baseline and week

1= 8 animals, week 2,3 = 7 animals). (**F**) Correlation between SARS-CoV-2 viral load in

bronchial brushes and absolute count of monocytes in blood at one-week post-infection (1wk pi)

(n= 8). (**G**) Representative flow cytometry changes in non-classical monocytes after the

infection in blood (1wk pi). (**H**) Percent of classical (CD14hi CD16low) and (**I**) non classical

monocytes (CD14low CD16hi) out of total monocytes in PBMCs (baseline = 8 animals week 1 =

4 animals, week 2, 3= 7 animals). (**J**) Correlation between monocytes absolute numbers in blood

and TARC (CCL17) in plasma (1wk pi). (**K**) Relationship between fold change of chemokines

and SARS-CoV-2 viral load in bronchoalveolar brushes (1wk pi). (**L**) Heatmap displaying log2

fold-change of 8 chemokines involved with traffic of monocytes to the lung. (1wk pi). Animals

are ordered by ranking of disease severity.

Figure 2. Myelomonocytic cells increase in BAL is associated with IL-6. (**A**) Decrease of

alveolar macrophages (HLA-DR+ CD206+ CD163+ cells) in the BAL after infection. (**B**)

Increase in the frequency of myelomonocytic infiltrates (HLA-DR+ CD206– CD163–) and (**C**)

of endothelial cells (HLA-DR+ CD206– CD163+) at infection in BAL (1wk pi). (**D**) tSNE plots

displaying kinetics of expression of CD16, CD11b and CD11c markers on live/ HLA-DR+ cells

populations in BAL before and after infection. (**E**) Gating of CD11c+ and CD11b+ populations

on the tSNE map based on antigen expression and discrete density clustering. (**F**)

Characterization of populations shown in (**E**). The heatmap depicts fold-change values of each

antigen compared to channel values of negative-expressing cell populations. (**G**) Increase in the

percentage of CD16+ HLA-DR+ cells and (**H**) in the percentage of interstitial macrophages

(CD16+ CD206– HLA-DR+) expressing of CD11c and (**I**) CD11b in BAL overtime.

Figure 3. Interstitial macrophages are increased in the lung of SARS-CoV-2 infected

animals. (**A**) Macrophage infiltration varied from minimal to severe in both rhesus and AGM

infected with SARS-CoV-2. (**B**) Fluorescent immunohistochemistry in the lung of AGM and

RM showing infiltration of CD16+ CD11b+ CD206– macrophages (arrows), with lesser

numbers of CD16+ CD206+ macrophages (arrowheads). Inset showing a higher magnification

of CD16+ CD11b+ CD206–macrophages (arrows). Asterix shows macrophages multifocally

surrounding small airways in GH99. White: DAPI (nuclei); Green: CD11b; Red: CD16; Blue:

CD206.

Figure 4. CD11b+ neutrophils accumulate in lung of infected animals with severe disease.

(**A**) Fluorescent immunohistochemistry in the lung of AGM and RM showing infiltration

of MPO+ neutrophils and MPO+ CD11b+ neutrophils. MPO and CD11b staining. White: DAPI

(nuclei); Green: CD11b; Red: MPO. The number of MPO positive cells (red, arrows) increases

in proportion to the severity of pulmonary disease from rare in the control (HK11), to mild in

FR04, moderate in GH99, and severe in NC34. Dual positive MPO+ CD11b+ positive cells

(arrowheads) are present in the most severely affected animals (NC34). (**B**) Mean number of

MPO+ cells, the standard deviation represents variability in the number of positive cells/area and

(**C**) and MPO+ CD11b+ per mm2 in lung necropsy.

Figure 5. Lymphocytes infiltration in the lungs is associated with disease severity. (**A**)

Histopathologic findings showing infiltration of lymphocytes in the lung of AGM (NC40 and

NC34) and RM (FR04 and GH99) with mild (NC40 and FR04), moderate (GH99), and severe

disease (NC34). (**B**) Frequency of NKG2a+ cells in the blood or (**C**) BAL of 4 AGM and 4 RM

at baseline, 2 weeks and 3 weeks post SARS-CoV-2 infection by flow cytometry. (**D**)

Representative flow cytometry plot showing decreased frequency of NGK2a+ cells in PBMCs

and increased frequency in BAL to baseline in at week 3 post SARS-CoV-2 infection respect to

baseline. One animal for each species is shown. (**E**) Decrease in the frequency of CXCR3+

NKG2a+ cells in the blood at 2 weeks, and 3 weeks post SARS-CoV-2 infection compared to the

baseline. (**F**) Absolute number of lymphocytes before and after infection. (**G**) tSNE plots

displaying kinetics of PD-1 expression (in red) on lymphocytic cell populations over time. (**H**)

Frequency of PD-1+ Th2 (CXCR3– CCR6–) CD4+ T cells is increased in RM and (**I**) in AGM

overtime following the infection. (**J**) Changes in the percentage of Th2 (purple), Th1 (blue) and

Th17 (grey) cells within the total PD-1+ CD4+ T cell population. (**K**) Significant increase in the

percentage of PD-1+ CD8+ T cells in BAL.

Figure 5. Increased IL-6 in the absence of IL-10 is associated with disease progression.

Levels of IL-6 and IL-10 in plasma overtime (pg/ml) **(A)** and **(B)** correlation between the ratio of

IL-10 to IL-6 levels measured in plasma and the ranking of disease severity or **(C)** with

pathology score**. (D)** Association between Kyn /Tryp ratio in plasma (pg/ml) and pathology

score. (**E**) Summary of immune events in lungs and blood of SARS-CoV-2 infected animals,

from the 1-acute phase 2- switch to Th2 type responses and 3- resolution phase.

Table 1: Heterogeneous clinical and pathological outcomes following SARS-CoV-2 infection

Animal ID	Clinical Findings	Clinical Score	Pathology Findings	Pathology Score	Ranking
NC34	Severe reduction in SpO2, moderate tachypnea, and severe hypothermia just prior to euthanasia for severe respiratory distress at 8 days post infection. Less than 24 hours earlier, the only abnormality was a moderate reduction in SpO2. Radiography appeared unremarkable until euthanasia when moderate to severe alveolar pattern with lobar sign and air bronchograms in right lung fields was observed.	9	Bronchointerstitial lung pattern in all lobes that was most severe in the right caudal lung lobe. Microscopically all lung lobes exhibited similar changes with the right caudal and right accessory lung lobe being the most severely affected. The lung lesions were comprised of diffuse alveolar damage with hyaline membrane formation, type II pneumocyte hyperplasia, and rare multinucleated giant syncytia. Some regions showed early evidence of organizing pneumonia and fibrosis.	19	1
NC33	Severe reduction in SpO2, severe tachypnea, and severe hypothermia just prior to euthanasia for severe respiratory distress at 22 days post infection. Less than 24 hours earlier, the only abnormality was a mild reduction in SpO2. Radiography appeared unremarkable until euthanasia when moderate to severe alveolar pattern in right middle/caudal lung fields and mild alveolar pattern left caudal lung fields were observed.	10	Severe diffuse alveolar damage of the right lower lung lobe. Other lobes exhibited suppurative alveolar infiltrate with lesser fibrin and edema. The right lung lobe contains peracute to acute lesions of necrosis, fibrin, and hemorrhage with minimal cellular infiltrate.	14	2
GH99	Transient cough beginning on 15 days post infection. Radiography showed significant changes at 11 days post infection which improved through 27 days post infection.	1	Severe bronchopneumonia of the right lower lung with granulomatous to pyogranulomatous inflammatory response.	10	3
NC38	Minor intermittent reductions in SpO2	0	Mild interstitial pneumonia with low numbers of multinucleated giant cells and rare atypical pneumocyte hyperplasia.	8	4
HD09	Mild focal increase opacity ventral lung field at 11 days post infection.	1	Minimal to mild inflammation scattered throughout the lungs.	9	5
HB37	Intermittent mild to moderate pyrexia from 1-21 days post infection with mild dehydration from 14-21 days post infection.	1	Proliferative lymphocytic vasculitis and proliferative vasculopathy in the right middle lung lobe. Scattered interstitial inflammation throughout the remaining lung lobes.	5	6
FR04	Unremarkable	0	Bronchioles exhibited multifocal, mild mineralization.	6	7
NC40	Minor intermittent reductions in SpO2	1	Minimal changes within lungs. Mild multifocal thickening of the alveolar septa in the right lower lobe associated with a mild inflammatory infiltrate.	4	8

870
871 Clinical findings are summarized for the duration of infection. Clinical scores, pathology findings, and
872 pathology scores reflect measurements at time of necropsy. Rankings were determined based on a

873 combination of the clinical findings and scores throughout the duration of infection and the pathology
874 findings and scores at time of necropsy (1 = severe; 8 = mild).
875 SpO_2 = oxygen saturation.
876

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6