1 **Cellular events of acute, resolving or progressive COVID-19 in SARS-CoV-2**
2 **infected non-human primates**
3
4

5 Fahlberg MD[1], Blair RV[1,2], Doyle-Meyers LA[1], Midkiff CC[1], Zenere G[1], Russell-
6 Lodrigue KE[1], Monjure CJ[1,2], Haupt EH[1], Penney TP[1], Lehmicke G[1], Threeton
7 BM[1], Golden N[1], Datta PK[1,2], Roy CJ[1,2], Bohm RP[1,2], Maness NJ[1,2], Fischer T[1,2],
8 Rappaport J[1,2], and Vaccari M[1,2]*.
9
10 [1]Tulane National Primate Research Center, Covington, LA
11 [2]Tulane University School of Medicine, New Orleans, LA
12

14 *To whom correspondence should be addressed:

16 Monica Vaccari, Ph.D.

17 Associate Professor, Tulane National Primate Research Center 21 18703

18 Three Rivers Road, Covington, LA 70433

19 Email: mvaccari@ tulane.edu

Abstract

We investigated the immune events following SARS-COV-2 infection, from the acute inflammatory state up to four weeks post infection, in non-human primates (NHP) with heterogeneous pulmonary pathology. The acute phase was characterized by a rapid migration of CD16+ monocytes from the blood and concomitant increase in CD16+ macrophages in the lungs. We identified two subsets of interstitial macrophages (HLA-DR+ CD206–), a transitional CD11c+ CD16+ population that was directly associated with IL-6 levels in plasma, and one long lasting CD11b+ CD16+ population. Strikingly, monocytes were a correlate of viral replication in bronchial brushes and levels of TARC (CCL17), and worse disease outcomes were associated with high levels of cell infiltration in lungs and CD11b+ CD16+ macrophages accumulation. Importantly, this accumulation was long-lasting and detectable even in animals with mild or no signs of disease. Interestingly, animals with less signs of disease had a high IL-10:IL-6 ratio. Our results unravel cellular mechanisms of COVID-19 and validate NHP as models to test immune therapies.

The COVID-19 pandemic is a major public health crisis, causing a global medical emergency. There are currently no effective vaccines or drugs approved to completely treat or prevent COVID-19, and much remains unknown about the pathogenesis of the disease. COVID-19 presents with highly variable outcomes, and patients that develop symptoms exhibit a wide range of disease, spanning from mild (fever, cough, shortness of breath) to severe (dyspnea, pneumonia, rapidly progressing radiographic changes)[1]. It is currently estimated that up to 15% of COVID-19 patients progress to acute respiratory distress syndrome (ARDS)[2], which is the major cause of death among fatal SARS-CoV-2 cases.

Studies elucidating molecular and immunological details of SARS-CoV-2 infection are

underway, and many of the reported clinical observations point to the disease being at least

partly the result of an excessive host response aimed to clear the virus but instead contributing to

disease development[3]. This hypothesis is supported by studies on the closely related coronavirus

SARS-CoV-1[4-6], and by clinical reports showing elevated levels of the proinflammatory cytokine

IL-6 in SARS-CoV-2 infected patients, particularly among those with severe symptoms[3,7].

Interestingly, the increase in IL-6 coincides with the upregulation of chemokines responsible for

myelopoiesis and monocyte recruitment to the lungs[3,8] suggesting that peripheral inflammatory

monocytes and tissue macrophages may play a role in the cytokine storm seen in severe COVID-

19 patients[9]. The underlying inflammatory state and increased myelopoiesis seen in older

individuals may be responsible for the high mortality rate seen in this group[10]. A better

understanding of immunopathology of COVID-19 will aid the development of targeted

immunotherapies and effective vaccines.

Non-human primates (NHP) have been used as animal models to study acute respiratory diseases

such as SARS and MERS[11]. In these studies, African green monkeys (AGM) have been shown to

be more susceptible to severe disease than rhesus macaques (RM)[12]. Regardless of species

differences, and given the overlap between NHP and with human immune systems, both NHP

species have been useful models to study host responses to coronaviruses[55]. We and others have

shown that both AGM and RM can be successfully infected with SARS-CoV-2 via the intranasal

and the intratracheal route, or by aerosol[13-16]. Following exposure, AGM and RM have high

levels of virus replication in the respiratory tract for up to two weeks, and both species develop

pulmonary pathology characteristic of COVID-19, that varies from very mild to severe[17]. Other

groups have described the very early immune events in SARS-CoV-2 infected rhesus macaques,

with perturbation of monocytes populations, increased macrophages in lungs, and cellular

activation in blood all happening a few days following the infection[18].

Here, we extend on these findings by describing the kinetics of immune events following

infection, spanning from the acute phase up to four weeks following infection, in the blood and

the lungs of four African green monkeys (AGM) and four rhesus macaques (RM) with a

heterogeneous spectrum of COVID-19. Overall, we observed a multiphasic mobilization of

innate and adaptive cells from the blood to the lung that is strikingly common to both species.

The acute inflammatory phase was characterized by an increase in the frequency of monocytes

that was positively associated with bronchial viral replication. Consistent with the resolution

phase of inflammation in the lung[19] , the second phase was characterized by a switch toward type

2 responses in both species. We identified two populations of interstitial macrophages (HLA-

DR+ CD206–) in BAL; 1) CD11c+ CD16+ population increasing during the acute phase of

infection that was associated with IL-6 levels in the plasma, and 2) a long lasting CD11b+ and

CD16+ population accumulating in lungs of both AGM and RM, particularly in those with

increased disease severity. Animals with more pronounced clinical signs had a higher level of

infiltrates in lungs when compared with animals with no or mild signs, and higher ratio of IL-

6:IL-10 (pro-inflammatory: anti-inflammatory) cytokines in plasma.

By investigating temporal and spatial changes in immune cells our study links together findings

in both NHP and human patients and unravels new cellular events of COVID-19[20,21]. Further, we

validate the use of NHP as a model to study immunopathology and to test immune therapies and

vaccines for SARS-CoV-2.

Results
99
Increased levels of monocytes and chemokines after infection is associated with viral load

and disease severity. Four adult RM and four adult AGM were exposed to SARS-CoV-2 by

either aerosol or mucosal challenge including buccal, intranasal, intratracheal, and conjunctival

(multi-route) exposure (**Supplementary Table 1**). The study concluded at 4 weeks post

exposure (**Fig. 1A**)[16]. All eight animals had detectable viral load by RT-PCR on nasal and

pharyngeal swabs and bronchial brushes by week 1 (**Fig. 1B**, **Supplementary Table 2**). Viral

RNA was also detected in pharyngeal, nasal, buccal, rectal and vaginal swabs. No differences in

viral load (VL) were observed between the different routes of exposure or between the two

species at week 1. Of note, by week 2 AGM had higher VL than RM in the bronchial brushes

and by week 3 VL in RM was mostly undetectable while it remained above detection levels in

AGM until study completion (**Fig. 1B** and **Supplementary Table 2**). Similar to humans, the

course of disease varied widely across animals, as detailed in **Table 1**. Of the four AGM

enrolled, two animals developed severe respiratory signs and were euthanized on days 8 (NC34)

and 22 (NC33) post-infection, following established endpoint criteria for NHP study protocols,

and as we previously reported[16]. Of the two remaining AGM, NC38 had multifocal mild to

moderate interstitial pneumonia scattered throughout all lung lobes and NC40 was mostly

asymptomatic and had scant inflammation in all lung lobes at necropsy. Three of the four RM

(GH99, HD09, and FR04) developed pneumonia characterized by a granulomatous to

pyogranulomatous inflammatory response. This was severe in GH99, mild in HD09, and

minimal in FR04. Histopathology of the fourth RM (HB37) revealed a lymphocytic vasculitis

and proliferative vasculopathy in the right middle lung lobe. Viral load was not significantly

associated with pulmonary pathology, however we observed an inverse association between viral

load at week 1 and disease severity ranking (1 to 8 from most severe to least severe) based on the

above observations and histopathology findings (AGM: $R = 0.8$; RM: $R = -0.4$, all animals: P

$= 0.057$, $R = - 0.71$, by the Spearman correlation test) (**Fig. 1C** and **SF 1A** and **1B**). Clinical

signs observed during the course of the study included mild tachypnea (day 11) and cough (day

15) in GH99. Intermittent mild to moderate pyrexia was observed in HB37 (day 1-21 post

infection). Radiographic changes included a nodule in the right caudal lung lobe (day 11) in

GH99, and HD09 also developed a mild, focal increased opacity in the ventral lung field at day

11 (**data not shown**).

Interestingly, virus-driven immunological changes were remarkably similar between species at

this time (**Fig. 1** and **SF 1**). To investigate overall changes driven by viral replication in both

groups, we combined their immunological analysis, however at the same time we maintained

different specifications for each species (teal = RM and magenta = AGM). Data and statistics are

also shown separately per group in **Supplementary Figure 1**. At 1 week after infection both

AGM and RM had an increase in the absolute number and frequency of monocytes in the blood

($P = 0.065$, and $P = 0.02$ respectively, by the Tukey multiple comparison test, $n = 8$) compared to

baseline levels (**Fig. 1D** and **1E**). This change was significant in AGM alone (absolute number: P

$= 0.0006$, $n = 4$), and was also observed in all RM, particularly GH99 (**SF 1C-1F**). The absolute

number of neutrophils did not change over time (**SF 1G**). Interestingly, the absolute number of

monocytes at week 1 was strongly positively associated with the viral load levels in the bronchial

brush samples ($P = 0.002$, $R = 0.93$ by the Spearman test) when all the animals were combined

(**Fig. 1F**) (AGM: R = 0.8; RM R = 1, not significant in separated group, **SF 1H** and **1I**). We then

used the CD14 and CD16 markers to discriminate between three circulating monocytes subtypes:

classical (CD14+ CD16–), intermediate (CD14+ CD16+) and non-classical (CD14– CD16+)

within the total (HLA-DR+ CD16+ and/or CD14+) monocyte population by flow cytometry

(**Fig. 1G - 1I** and **SF 1J- 1N**). At week 1, the frequency of classical monocytes was significantly

increased in blood, as seen in humans[22], and the non-classical population was profoundly

depleted (increased levels baseline versus week 1: classical P = 0.007 and decreased non-

classical P = 0.005 in all animals combined, n = 4) (**Fig. 1H** and **1I**, and **SF 1J- 1K**).

Pre-infection levels were restored at 3 weeks post-infection (week 1 vs. week 3: classical P =

0.03 and non-classical P = 0.03). These results suggest a rapid and robust recruitment of

patrolling monocytes with antiviral activity in tissues following infection in both species.

We analyzed a group of eight chemokines known to be involved in the mobilization of innate

cells, including monocytes, from the blood to the lung: IP-10 (CXCL-10), MCP-1 (CCL2),

MCP-4 (CCL13), Eotaxin, TARC (CCL17), MDC (CCL22), MIP-1a and MIP-1b (**SF 2A-**

2H). Some of the cytokines measured at baseline in plasma differed between the two species

(MCP-4, Eotaxin, MIP-1b and MDC, **Supplementary Table 3** and **SF 2I**). Plasma levels of

TARC (CCL17), a ligand for CCR4 that is involved in Th2 cell recruitment[23], were strongly

associated with the frequency of monocytes in blood (P = 0.003, R = 0.95, Spearman test,

adjusted for multiple comparison: P = 0.02) (**Fig. 1J**). Similarly, TARC was also associated with

viral replication in BAL at week 1 (**Fig. 1K** and **SF 2J**) (P = 0.02, R = 0.857, adjusted for

multiple comparisons: P = 0.2), as was IP-10 (P = 0.05, R = 0.78, adjusted for multiple

comparisons) when all animals were grouped together (**Fig. 1K** and **SF 2K**), a ligand for CXCR3

that is associated with disease severity and predicts progression of COVID-19 in humans[24]. To

gain a better understanding of a common milieu possibly driving monocyte migration observed

in both groups we analyzed changes in the levels of these markers with respect to the baseline.

Overall this analysis revealed that animals with the most severe disease outcomes had higher

levels of chemokines at week 1, including IP-10, TARC (CCL17) (**Fig. 1L**).

169
Myelomonocytic cells infiltrate BAL and are associated with IL-6. Influx of macrophages has

been observed in the lung of severe cases of COVID-19 in humans[25]. To monitor cell trafficking

into the lungs, we collected bronchoalveolar lavage (BAL) at baseline and after each week

following infection, up to week 3. As expected, the majority of the cells found in BAL at

baseline were alveolar macrophages (AM)[26]. These cells are readily identified by the mannose

receptor C type 1 (CD206) marker, express high levels of HLA-DR, and are positive for the

scavenger receptor CD163 (**Fig. 2, SF 3A** and **3B**), we observed a decrease in the proportion of

AM at week 1 post infection in animals of both species, corresponding with an influx of myeloid

HLA-DR+ CD206– CD163– cells (**Fig. 2A** and **2B**, **SF 3C**). Levels of AM similar to pre-

infection were restored at week 2 in all but one animal (GH99), and a concomitant increase of

CD86 mean fluorescent intensity (MFI) in this population was observed suggesting an M1

activation state[27] (**SF 3D**). An increase was also seen in the frequency of CD206+ CD163– non-

macrophages that have been previously described as endothelial cells which facilitate

lymphocyte trafficking and they are detectable in BAL during inflammatory events (**Fig. 2C**)[28].

T-distributed stochastic neighbor embedding (tSNE) analysis revealed changes in the phenotype

of CD16, CD11b, and CD11c positive populations most prominently at 1-2 weeks post infection

(**Fig. 2D - 2F**). In contrast to what we observed in the blood, there was an increase of CD16+

HLA-DR+ cells in BAL over time (**Fig. 2G**, and **SF 3E** and **3F**). We observed a transient

increase in the level of CD11c+ CD16+ macrophages, which function has been described as

patrolling, and they participate in rapid tissue invasion and monocyte survival (baseline vs. week

1: P = 0.015) (**Fig. 2H**)[29,30]. The frequency of CD11c+ macrophages in BAL at week 3 was

strongly associated with the levels of the proinflammatory IL-6 cytokine in the plasma at the

same time point (week 3: P = 0.003, R = 0.96, Spearman test) (**SF 4**). The CD11b marker is

required for monocyte migration to sites of inflammation[31], and is highly expressed on myeloid

cells recruited from the blood and differentiating into macrophages[32]. We also observed an

increase in the percentage of CD11b+ CD16+ HLA-DR+ CD206– macrophages in BAL at week

2 and 3 following infection (**Fig. 2I**), a population that is involved in regulating lung

inflammatory response[19,27,33,34]. The gating strategy for the two macrophages populations is

shown in **Supplementary Figure 3G**.

Interstitial macrophages accumulate in lungs of infected animals. Lungs were collected for

all animals at the end of the study, corresponding to 4 weeks post infection (1 week following the

last BAL collection shown in Figure 2), with the exception of NC33 and NC34. These two

animals had higher levels of IL-6 at necropsy compared to the other six monkeys, and NC34 had

symptoms similar to ARDS[16] (**Fig. 3**). Overall, infiltration of macrophages was observed in the

lungs of SARS-CoV-2 infected AGM and RM at necropsy and was more robust in animals with

severe disease (**Fig. 3A**). We further characterized macrophages population in the lung by

staining with the AM marker CD206 (in blue) and with CD11b integrin (in green) and CD16

(red) (**Fig. 3B**). Low numbers of macrophages were seen in the lungs of a non-infected control

RM, scattered throughout the interstitium and alveolar spaces, as shown by the arrows. In the

control this population was bi-morphic and composed of CD11b negative CD16+ CD206–

(arrow) in the alveolar septa or CD16+ CD206+ cells within the alveolar spaces (arrowhead)

(**Fig. 3B**). In all animals that were infected we could detect CD11b+ CD16+ macrophages. FR04

(RM ranking = 7) lungs had macrophages scattered throughout the interstitium and alveolar

spaces. The phenotypic population was heterogeneous and contained low numbers of CD11b+

CD16+ CD206– macrophages. The most severe RM, GH99 (ranking = 3), had large numbers of

macrophages multifocally surrounding small airways denoted by the asterisks. Of note this

animal had the higher absolute number of monocytes in the blood at week 1 (**SF 1D**). Low

numbers of CD11b+ CD16+ CD206– cells were scattered throughout the predominant

population of CD11b+ CD16+ CD206+ macrophages. Macrophages were also scattered

throughout the interstitium and alveolar spaces. NC34 and NC33 (AGM ranking = 1 and 2

respectively) had multifocal to coalescing areas of hemorrhage (NC33 indicated by the

arrows), and pulmonary parenchyma infiltrated by multifocal aggregates of macrophages (NC34)

with infiltration by low to moderate numbers of macrophages. The phenotypic population was

heterogenous and composed of moderate numbers of CD11b+ CD16+ CD206– macrophages.

Altogether these findings present a picture of long-lasting lung inflammation that, albeit more

pronounced in animals with severe signs of disease, it is also present in animals with no

detectable viral load and mild or no signs of disease after 4 weeks from the infection.

NKG2a+ cells and PD-1+ T cells decrease in blood and increase in BAL overtime.

Infiltration of lymphocytes was observed in the lungs of SARS-CoV-2 infected AGM and RM

at necropsy and was more robust in animals with severe disease (**Fig. 4A**). We first analyzed the

kinetics of innate NK cells subsets, that we defined as HLA-DR– CD3– cells expressing the

NKG2a receptors, which are known to traffic to the lung[35] (**Fig. 4B-D**, and **SF 5A** and **5B**)

Interestingly, the frequencies of NKG2a+ cells significantly decreased in peripheral blood of

both AGM and RM over the course of infection (baseline vs. week 3, P = 0.03) (**Fig. 4B**).

Consistent with the pattern described for the myeloid population, we also observed a

concomitant increase in the frequencies of NKG2a+ cells in BAL (baseline vs. week 3, P = 0.05)

(**Fig. 4C** and **4D**, **SF 5A** and **5B**). Four out of four AGM had decreased NKG2a+ cells in

peripheral blood, coinciding with an increase of the same cells in BAL. While the pattern of

NKG2a+ cells in peripheral blood of RM was not statistically significant, nonetheless in this

species there was a marked increase of those cells in the BAL. These results suggest that the loss

of NKG2a+ cells in the PBMCs and the gain of NKG2a+ cells in the BAL may coincide with an

egress of these cells from peripheral blood to tissues. Accordingly, the frequency of NKG2a+

cells expressing CXCR3, a marker associated with NK cell migration to the lung[36] also

significantly decreased in blood over time (baseline vs. week 3, P = 0.03, **Fig. 4E**). Despite this

change in lymphocyte phenotype, the absolute number of lymphocytes did not change over the

first 2 weeks of infection (**Fig. 4F**) though it is important to note that all 3 remaining AGM had

an increase at week 3 (**SF 5C**).

tSNE analysis revealed an increase in the level of PD-1 at weeks 2 and 3 post-infection,

overlapping with spatial regions corresponding to both CD4+ T and CD8+ T cells (**Fig. 4G**). In

fact, the frequency of PD-1+ CD4+ T cells was elevated in the blood of both RM and AGM (**Fig.**

4H, and **4I**, respectively). During the first week following the infection we observed an increase

in IP-10 and IFN-γ (**SF 5E**), however we also observed a switch toward Th2 type responses

characterized by the increase of Th2 cytokines IL-5 and IL-13 over time (**SF 3E**). Accordingly, a

gradual increase of Th2 type cells (defined as CXCR3– CCR6 – CD4+ T cells)[37] was seen in the

blood within the PD-1 positive population (**Fig. 4J**). The increase of PD-1+ Th2 cells was

significantly and positively associated with IL-4 cytokine levels (P = 0.016, R = 0.94, Spearman

test) (**SF 5F**). Moreover, we observed an increased level of PD-1+ CD8+ T cells in blood and

BAL (BAL: baseline vs. week 2, P = 0.015; baseline vs. week 3, P = 0.03) (data not shown and

Fig. 4K). Animals divided by species are shown in **SF 5G** and **5H**.

The ratio between the anti-inflammatory cytokine IL-10 and the pro inflammatory

cytokine IL-6 is associated with disease progression. The pro-inflammatory cytokine IL-6 has

been described as a strong predictor to disease progression in humans[3,7]. We measured IL-6 and

IL-10 in the plasma of all the animals (**Fig. 5A**). While IL-6 levels alone did not associate with

disease severity in NHPs, the ratio of IL-10 to IL-6 levels measured in plasma associated with

both the ranking of disease severity and with the pathological score that was obtained by

analyzing the extension of the inflammation and edema of the lungs (R = 0.76 P = 0.037; and R

= –0.76, P = 0.03, by the Spearman correlation test, all animals) (**Fig. 5B** and **C**). These findings

show a complex series of immune events in the lung of that are most likely initiated as a host

response to the virus and then evolving overtime toward less antiviral responses, as summarized

in **Figure 5D**.

DISCUSSION

There is an urgent need to develop safe and effective strategies to prevent and treat COVID-19.

The development of such strategies requires a clear understanding of the immune cells involved

in COVID-19 and their unique role in pathogenesis. Clinical observations in patients and few

studies on non-human primates have started to shed light on the immune aspects involved in

COVID-19[9,20,38,39]. Here, we further characterized immune events following SARS-CoV-2

infection in two NHPs models recapitulating crucial aspects of COVID-19 lung disease[14,16]

(**Table 1**). Altogether our findings provide the first evidence in both AGM and RM of a

multiphasic immune response characterized by 1) an initial inflammatory phase with monocyte

recruitment to the lung 2) a gradual switch from a type 1 to type 2 response, and 3) a "make it or

break it" phase with either an increase in anti-inflammatory cytokines IL-10, or IL-6 a pro-

inflammatory cytokines associated with disease progression in humans, as summarized in **Figure**

5D.

By examining the temporal and spatial distribution of monocytes and macrophage in blood and

BAL and lungs, we confirmed and extended on studies in humans and NHPs suggesting

increased myelopoiesis with recruitment of myeloid cells from the blood to lungs during the

infection that is initially driven by the virus replicating in the lung[3,14,20]. Similar to what has been

described by Wilk *et al.* and Thevarajan *et al.* in humans, we observed an increase of CD14+

monocytes, and a corresponding drop in the frequency of CD16+ monocytes in the peripheral

blood[20,21]**.** Importantly, we extend on these findings by showing a simultaneous increase in the

frequency patrolling CD16+ HLA-DR+ monocytes in BAL occurring in both species and by

describing the kinetics of two CD16+ HLA-DR+ CD206– macrophages populations that

preferentially populate the alveolar interstitium and exhibit features of antigen-presenting

cells[40]: CD11c+ CD16+ cells arising from intravascular CD16+ patrolling monocytes in the

acute phase and associating with IL-6 levels, and a CD11b+ macrophage population known to be

involved in resolution of inflammation in the lung[19,27,33,34]. The origin and the exact function of

these interstitial macrophages remains to be determined.

Differently than was observed in humans and previous reported data in RM, we did not detect an

increase in the number of neutrophils as these responses may have come up earlier than 1 week

after infection as shown by others[18].

307
Common to both species was the observation that animals with higher levels of myeloid and

lymphoid cell infiltration in the lung had worse disease outcomes. Several studies in both SARS-

CoV-1 and CoV-2 have reported depleted peripheral NK cell counts in severe patients[9,20,22,38].

Our data support the notion of NKG2a+ cell trafficking to the lung during SARS-CoV-

2 infection, most likely mediated by IP-10 (CXCL-10), which promotes the recruitment of

CXCR3+ Th1 thymocytes and NKs[41,42]. Expression of NKG2a has been linked to both activation

and inhibition of inflammation[43]; the function in SARS-CoV-2 remains to be investigated. In the

periphery, we also observed increased levels of PD-1 on T cells. This is consistent with findings

in humans describing of an exhaustive T cell phenotype during SARS-Cov-1 and Cov-2

infection[9,44].

For the first time, we identified TARC (CCL17) as central to myeloid recruitment in the lung

during SARS-CoV-2. Both dendritic cells and CD11b macrophages in the lung have been

identified as a source of CCL17 in the airways in other pulmonary inflammatory diseases[45,46].

These cells promote the recruitment of T helper type 2 (Th2) lymphocytes into the airway[45].

Remarkably, we observed a gradual increase of the recruitment of Th2 lymphocytes into the

airway Th2 cell frequency and of Th2 type cytokines such as IL-5, IL-13. These findings show a

complex series of immune events in the lung of that are most likely initiated as a host response to

the virus, and that they may evolve into other less effective antiviral responses as either an

attempt to tissue repair/resolution of inflammation, as summarized in **Figure 5D** or as the result

of virus direct immunomodulatory effects on host responses[6].

Intrinsic species-specific susceptibility to disease progression have been described for NHP in

SARS-CoV-1[6] with AGM being more susceptible than rhesus to severe disease. There were

unexpected similarities in the kinetics and the quality of the immune responses analyzed in RM

and AGM, particularly during the acute phase of infection, however AGM did not clear the

infection as well as RM and two AGM had most severe symptoms[47]. While the number of

animals enrolled in this study was not sufficient to look at species-specific differences and their

effect on disease progression, we found that a higher ratio of IL-6:IL-10 cytokine in plasma was

associated with disease severity. Undoubtedly, more studies are needed to prove or disprove the

single contribution of IL-10 and IL-6 in the outcome of COVID-19 disease and to identify the

source of these two potential key players[48].

Overall, this study uncovers critical steps in the immune events of the host response to SARS-

CoV-2 and describes new correlates of viral replication and disease progression. Importantly, we

show viral driven inflammation/cell recruitment to the lung occurs rapidly, and lasts up to four

weeks post infection, which may explain in part the observed slow recovery from COVID-19

seen humans. Finally, these findings have the potential to aid the interpretation of the results of

the human trials that are currently underway for several vaccine candidates and may be

instrumental in generating new targeted therapeutic strategies aiming at resolving the severe

immune deregulation of the lung of COVID-19 patients.

Methods

Ethical statement on animal use and SARS-CoV-2 handling. Animals and infection. The

Institutional Animal Care and Use Committee of Tulane University reviewed and approved all

the procedures for this study. The Tulane National Primate Research Center is fully accredited

by the AAALAC. All animals were cared for in accordance with the ILAR Guide for the Care

and Use of Laboratory Animals 8th Edition. The Tulane University Institutional Biosafety

Committee approved the procedures for sample handling, inactivation, and transfer from BSL3

containment.

Animals and infection. Four adult-aged African green monkeys (Caribbean origin) and four

rhesus macaques (Indian ancestry) were exposed to SARS-CoV-2; 2019-nCoV/USA-WA1/2020

409 (MN985325.1). The virus stock was prepared in Vero E6 cells and the sequence confirmed

by PCR and/or Sanger sequencing. Plaque assays were performed in Vero E6 cells. The rhesus

macaques were from the Tulane National Primate Research Center breeding colony pathogen

free for simian type D retrovirus (SRV), macacine herpesvirus 1 (B virus), simian

immunodeficiency virus (SIV), simian T cell lymphotropic/leukemia virus (STLV), measles

virus (MV) and tuberculosis (TB). The African green monkeys were wild caught and at the

TNPRC for over a year before being assigned to this study. To mimic different possible routes of

infection in humans, animals were exposed to the virus either by aerosol (inhaled dose of 2.0 x

10^3 and 2.5 x 10^3 TCID$_{50}$) or by inoculating a cumulative dose of 3.61 x 10^6 PFU through

multiple routes (oral, nasal, intratracheal, conjunctival) (**Supplementary Table 1**).

The multi route exposure was given to four animals, one adult RM male (GH99, 14 years old),

one adult RM female (HD09, 13 years old) and two AGM, one aged male (NC40, 16 years old,

approximately) and one aged female (NC33, 16 years old, approximately). An additional two

adult males RM (FR04 and HB37, 15 and 13 years old, respectively) and one aged male and

female AGM: NC34 and NC38 (16 years old, approximately) were exposed by aerosol

(**Supplementary Table 1**).

379
Cell count and CO_2 and SpO_2 pressure measurements. Total white blood cell counts, white

blood cell differentials, red blood cell counts, platelet counts, hematocrit values, total

hemoglobin concentrations, mean cell volumes, mean corpuscular volumes, and mean

corpuscular hemoglobin concentrations were analyzed from blood collected in tubes containing

EDTA using a SYSMEX XN-1000v Hematology Analyzer in the Level 2 Lab and a Sysmex XS-

1000 in the Level 3 Lab. CO_2 measurements were taken using the Piccolo Express Chemistry

Analyzer in the Level 3 lab and is part of the Metabolic Comprehensive Panel. SpO_2 values were

obtained using the Masimo Rad-57 Handheld Pulse Oximeter.

388
Sample collection and processing. Blood EDTA was collected by venipuncture and bronchial

brush and BAL by bronchoscopy. For the latter, the bronchoscope is introduced into the trachea

and directed into one of the bronchi. The bronchoscope is advanced to the point where the

diameter of the bronchoscope approximates the dimensions of the bronchus. The bronchial brush

is passed through the channel of the bronchoscope. The brush is gently advanced until the lumen

of the bronchus or bronchioles approximates the dimensions of the brush. The brush is then be

gently advanced back and forth against the mucosa to gather cells. The brush is removed from

the channel of the bronchoscope and the brush is placed into 200 μL of DNA/RNA Shield X1

(Cat.# R1200, Zymo Research, Irvine, CA). The bronchoscope remains in place while two

aliquots of 20 ml of saline each are instilled into the bronchus and subsequently aspirated. This

procedure is then repeated in the opposite lung.

Blood was layered over Ficoll-Plaque Plus (GE Healthcare17144003) and centrifuged for 30

minutes in cap-locked adaptors. Differential migration of cells during centrifugation results in

the formation of layers containing different cell types and allows collection of peripheral blood

mononuclear cells (PBMCs) from other populations. PBMCs were collected, counted and

aliquoted in tubes or resuspended in freezing media containing Fetal Bovine Serum and DMSO.

For bronchoalveolar lavage (BAL), cells and supernatant were separated by centrifugation (15

minutes at 2,000 x g at RT) in cap-locked adaptors and cells were lysed in Ammonium Chloride

Potassium (ACK) lysing buffer at room temperature for 5 minutes (Gibco# A1049201) washed

in 2% FBS, and resuspended in PBS and counted under laminar flux hood in BCL3. Fresh cells

were stained for flow cytometry.

BAL and PBMCs staining. Freshly obtained PBMCs and cells obtained from BAL were stained

with appropriate antibody cocktails and incubated for 30 minutes at room temperature.

Antibodies are listed in **Supplementary Table 4.** Cells were fixed with 4% or 2% PFA-BSA

overnight. Samples were run on a FACSAria Fusion flow cytometer (Becton Dickinson) within a

Biosafety Cabinet, and each sample was run for at least five minutes. For all panels, the

LIVE/DEAD Fixable Aqua Dead cell Stain kit was used (ThermoFisher #L34957). Alveolar

macrophages were gated based on the size and the expression of HLA-DR, CD206 and CD163

markers. Myeloid infiltrates were gated as cells negative for CD206 and CD163 and positive for

HLA-DR, CD16 and CD11b. The gating strategy is shown in **Supplementary Figure 3G**.

NKG2a+ cells were gated on the CD3 negative parent population. Th1, Th2, and Th17 cells were

gated as previously described[37]. tSNE analyses were performed in FlowJo using the opt-SNE

algorithm and designed in R using the ggplot2 package[49].

Histopathology and scoring. Tissue samples were fixed in Z-fix (Anatech), embedded in

paraffin and 5 μm thick sections were cut, adhered to charged glass slides and stained with

hematoxylin and eosin. All slides were scanned on a Zeiss Axio Scan.Z1 digital slide scanner.

Images were acquired using HALO software (Indica Labs). Quantification of pathologic lesions

was performed with HALO software. Machine learning (Random Forest) algorithms were

trained by a board -certified pathologist to recognize fluidic and cellular inflammation in the

lungs of the RM and AGM. Scores were assigned to fluidic and cellular inflammation,

separately, based on the percentage of lung affected. Fluidic and cellular inflammation scores

were summated for all lobes and animals were ranked from 1 (highest sum) to 8 (lowest sum).

Plasma cytokines and chemokines. Plasma was collected by spinning and was thawed before

use. Cytokines were measured using 497 Mesoscale Discovery. IFN-γ, IL-4, IL-6, IL-8, IL-10

and IL-13 were part of the V-Plex Proinflammatory Panel 1, 10-Plex. V-plex. The chemokine

panel (#K15049D, Mesoscale Discovery, Rockville, Maryland) was also used, following the

instructions of the kit. The plate was read on a MESO Quickplex 500 SQ120 machine.

Heatmaps were generated using the 'pheatmap' package in R[50,51]. Data were normalized by

dividing raw values at week 1 by baseline values for each animal, followed by the application of

$\log_2$. Values below the limit of detection were replaced with the lowest limit of detection value

based on the standard curve for each run, or with the lowest value detected during the run,

whichever was smaller. Bubble plots were generated using the 'ggplot2' package in R, using the

same normalized data shown in the heatmap[49]. Scatter plots were drawn using raw data points

and display Pearson's correlation coefficients and a 95% confidence interval.

RNA isolation. Swab and bronchial brush samples were collected in 200 μL of

DNA/RNA Shield X1 (Cat.# R1200, Zymo Research, Irvine, CA) and extracted for

Viral RNA (vRNA) using the Quick-RNAViral kit (Cat.# R1034/5, Zymo Research). The Viral

RNA Buffer dispensed directly to the swab in the DNA/RNA Shield. The swab was directly

inserted into the spin column. The vRNA was eluted (45μL) and 5 μL was added in a 0.1 mL fast

96-well optical microtiter plate (Cat.# 4346906, ThermoFisher, CA).

454
Viral RNA detection. RT-qPCR reaction TaqPath1-Step MultiplexMaster Mix was used (Cat.#

A28527, ThermoFisher) along with the 2019-nCoV RUO Kit (Cat.# 10006713, IDTDNA,

Coralville, IA) targeting the N1 amplicon of N gene of SARS2-nCoV19 (accession

MN908947). The master mix was added to the microtiter plates covered with optical film (cat.

#4311971; ThermoFisher) and then was vortexed and pulse centrifuged. The RT-qPCR

program consisted of incubation at 25°C for 2 minutes, RT

incubation at 50°C for 15 minutes, and an enzyme activation at 95°C for 2 minutes followed

by 40 cycles of denaturing step at 95°C for 3 seconds and annealing at 60°C for 30 seconds.

Fluorescence signals were detected with an Applied Biosystems QuantStudio6

Sequence Detector. Data were captured and analyzed with Sequence Detector Software v1.3

(Applied Biosystems, Foster City, CA). Viral copy numbers were calculated by plotting Cq

Values obtained from unknown (i.e. test) samples against a standard curve representing known

viral copy numbers. The limit of detection of the assay was 10 copies per reaction volume. A

2019-nCoVpositive control (Cat.#10006625, IDTDNA) were analyzed in parallel with every set

of test samples. A non-template control (NTC) was also included.

Immunohistochemistry. 5 μm sections of Formalin-fixed, paraffin-embedded lung were

mounted on charged glass slides, baked overnight at 56_{o}C and passed through Xylene, graded

ethanol, and double distilled water to remove paraffin and rehydrate tissue sections. A

microwave was used for heat induced epitope retrieval. Slides were heated in a high pH solution

(Vector Labs H-3301), rinsed in hot water and transferred to a heated low pH solution (Vector

Labs H-3300) where they were allowed to cool to room temperature. Once cool, slides were

incubated with Background Punisher (Biocare Medical BP974H) for 10 minutes, washed and

incubated with serum free protein block (Dako X0909) for 20 minutes. Immediately following

blocking, a mouse anti-CD16 primary antibody was diluted in the serum free protein block and

added to the slides for 60 minutes. Slides were washed and a MACH3 AP kit (Biocare Medical

M3M532) was used to label the primary antibody. Both the MACH3 mouse probe and polymer

were incubated for 20 minutes with washes in between. Signal was developed with Permanent

Red substrate (Dako K0640) for 13 minutes to allow for visualization of the CD16+ cells. Slides

were thoroughly washed and incubated with a blocking buffer comprised of 1% normal donkey

serum (NDS) for 40 minutes. A rabbit anti-CD11b primary antibody was diluted in NDS and

added to the slides for 60 minutes. After washing, slides were incubated for 40 minutes with a

fluorescently tagged secondary antibody. The staining process was then repeated, on all slides,

for the labeling of CD206. Following washes, DAPI (4',6-diamidino-2-phenylindole) was used

to label the nuclei of each section. Slides were mounted using a homemade anti-quenching

mounting media containing Mowiol (Calbiochem #475904) and DABCO (Sigma #D2522) and

imaged with a Zeiss Axio Slide Scanner. Antibodies are shown in **Supplementary Table 4.**

Statistical analysis. Correlation analyses were performed using the non-parametric Spearman

rank correlation method (two-tailed, 95% confidence) using exact permutation P values when

ranks were used, or Peason's correlation test when both variables were continuous. Analysis of

variance (ANOVA) was used for comparing statistical differences between multiple groups,

followed by Dunn's multiple comparison tests. Kruskal-Wallis ANOVA or Mann-Whitney tests

were used. All statistical analyses were performed using GraphPad Prism (version 8.4.2

GraphPad Software, La Jolla California USA) and R software (URL http://www.R-

project.org/). Correlation plots were created using the 'Performance Analytics' and 'corrr' plots,

respectively, in R[51-53].

Acknowledgments

We would like to thank Dr. Joseph (JC) Mudd and Dr. Cariappa Annaiah for critical reading of
the manuscript and Amitinder (Miti) Kaur, Megan Varnado, Kaitlin Didier from the TNPRC
Flow Cytometry Core for their help. We thank Angela Birnbaum for reviewing and optimizing
all technical SOPs and overseeing the safety of this study.
We would also like to thank Fast Grant Funding for COVID-19 Science for partially funding this
work and the NIH for supporting this work through the TNPRC base grant (P51 OD011104 59).
509

510 **Competing Interests statement**
511

512 These authors declare no competing interests.

513
Author Contributions
MDF analyzed data composed figures and wrote the manuscript,
BRV was the lead pathologist and LADM was the project veterinarian, contributed to study design, and
writing of the IACUC and manuscript. CCM designed IHC panels and performed the IHC staining, GZ
wrote the manuscript, KERL was a project veterinarian, made clinical assessments and collected samples,
CJM processed and analyzed samples for RT-qPCR, contributed; EHH provided the mesoscale data, TPP
stained the samples with GL and BMT. GL contributed to study design, provided administrative support,
and aided with sample processing and archiving. NG contributed to study design, study coordination,
sample processing, and SOP development. PKD processed and analyzed viral load data. CJR conceived
and performed aerosol experiments. RPB contributed to study design, analysis of clinical and imaging
results, and writing the manuscripts. NJM performed staining of BAL and contributed to writing the
manuscript. TF contributed to the animal study design and planning and contributed to writing the
manuscript. JR designed and supported the animal study and helped with the writing of the manuscript.
MV conceived and supported the study, run experiments, analyzed data and wrote the manuscript.

REFERENCES

1.   Guan, W.J.*, et al.* Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* **382**, 1708-1720 (2020).

2.   Sun, P.*, et al.* Clinical characteristics of hospitalized patients with SARS-CoV-2 infection: A single arm meta-analysis. *J Med Virol* **92**, 612-617 (2020).

3.   Blanco-Melo, D.*, et al.* Imbalanced Host Response to SARS-CoV-2 Drives Development of COVID-19. *Cell* **181**, 1036-1045.e1039 (2020).

4.   Weiss, S.R. Forty years with coronaviruses. *J Exp Med* **217**(2020).

5.   Clay, C.*, et al.* Primary severe acute respiratory syndrome coronavirus infection limits replication but not lung inflammation upon homologous rechallenge. *J Virol* **86**, 4234-4244 (2012).

6.   Graham, R.L., Donaldson, E.F. & Baric, R.S. A decade after SARS: strategies for controlling emerging coronaviruses. *Nat Rev Microbiol* **11**, 836-848 (2013).

7.   Ruan, Q., Yang, K., Wang, W., Jiang, L. & Song, J. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive Care Med* **46**, 846-848 (2020).

8.   Xiong, Y.*, et al.* Transcriptomic characteristics of bronchoalveolar lavage fluid and peripheral blood mononuclear cells in COVID-19 patients. *Emerg Microbes Infect* **9**, 761-770 (2020).

9.   Zhou, Y.*, et al.* Pathogenic T cells and inflammatory monocytes incite inflammatory storm in severe COVID-19 patients. *National Science Review* (2020).

10.   Crooke, S.N., Ovsyannikova, I.G., Poland, G.A. & Kennedy, R.B. Immunosenescence and human vaccine immune responses. *Immun Ageing* **16**, 25 (2019).

11.   Gretebeck, L.M. & Subbarao, K. Animal models for SARS and MERS coronaviruses. *Curr Opin Virol* **13**, 123-129 (2015).

12.   McAuliffe, J.*, et al.* Replication of SARS coronavirus administered into the respiratory tract of African Green, rhesus and cynomolgus monkeys. *Virology* **330**, 8-15 (2004).

13.   Chandrashekar, A.*, et al.* SARS-CoV-2 infection protects against rechallenge in rhesus macaques. *Science* (2020).

14.   Singh, D.K.*, et al.*

15.   Yu, J.*, et al.* DNA vaccine protection against SARS-CoV-2 in rhesus macaques. *Science* (2020).

16.   Blair, R.V.*, et al.* ARDS and Cytokine Storm in SARS-CoV-2 Infected Caribbean Vervets. *BioRxiv* (2020).

17.   Munster, V.J.*, et al.* Respiratory disease in rhesus macaques inoculated with SARS-CoV-2. *Nature* (2020).

18.   Singh, D.K.*, et al.* SARS-CoV-2 infection leads to acute infection with dynamic cellular and inflammatory flux in the lung that varies across nonhuman primate species. *BioRxiv* (2020).

19.   Aggarwal, N.R., King, L.S. & D'Alessio, F.R. Diverse macrophage populations mediate acute lung inflammation and resolution. *Am J Physiol Lung Cell Mol Physiol* **306**, L709-725 (2014).

573   20.   Wilk, A.J.*, et al.* A single-cell atlas of the peripheral immune response in patients with
574   severe COVID-19. *Nat Med* (2020).
575   21.   Thevarajan, I.*, et al.* Breadth of concomitant immune responses prior to patient recovery:
576   a case report of non-severe COVID-19. *Nat Med* **26**, 453-455 (2020).
577   22.   Wen, W.*, et al.* Immune cell profiling of COVID-19 patients in the recovery stage by
578   single-cell sequencing. *Cell Discov* **6**, 31 (2020).
579   23.   Anderson, C.A.*, et al.* A degradatory fate for CCR4 suggests a primary role in Th2
580   inflammation. *J Leukoc Biol* **107**, 455-466 (2020).
581   24.   Yang, Y.*, et al.* Plasma IP-10 and MCP-3 levels are highly associated with disease
582   severity and predict the progression of COVID-19. *J Allergy Clin Immunol* (2020).
583   25.   Pence, B.D. Severe COVID-19 and aging: are monocytes the key? *Geroscience* (2020).
584   26.   Cai, Y.*, et al.* In vivo characterization of alveolar and interstitial lung macrophages in
585   rhesus macaques: implications for understanding lung disease in humans. *J Immunol* **192**,
586   2821-2829 (2014).
587   27.   Johnston, L.K., Rims, C.R., Gill, S.E., McGuire, J.K. & Manicone, A.M. Pulmonary
588   macrophage subpopulations in the induction and resolution of acute lung injury. *Am J
589   Respir Cell Mol Biol* **47**, 417-426 (2012).
590   28.   Irjala, H.*, et al.* Mannose receptor is a novel ligand for L-selectin and mediates
591   lymphocyte binding to lymphatic endothelium. *J Exp Med* **194**, 1033-1042 (2001).
592   29.   Auffray, C.*, et al.* Monitoring of blood vessels and tissues by a population of monocytes
593   with patrolling behavior. *Science* **317**, 666-670 (2007).
594   30.   Landsman, L.*, et al.* CX3CR1 is required for monocyte homeostasis and atherogenesis by
595   promoting cell survival. *Blood* **113**, 963-972 (2009).
596   31.   Prieto, J., Eklund, A. & Patarroyo, M. Regulated expression of integrins and other
597   adhesion molecules during differentiation of monocytes into macrophages. *Cell Immunol*
598   **156**, 191-211 (1994).
599   32.   Tam, J.W., Kullas, A.L., Mena, P., Bliska, J.B. & van der Velden, A.W. CD11b+ Ly6Chi
600   Ly6G- immature myeloid cells recruited in response to Salmonella enterica serovar
601   Typhimurium infection exhibit protective and immunosuppressive properties. *Infect
602   Immun* **82**, 2606-2614 (2014).
603   33.   Liao, M.*, et al.* Single-cell landscape of bronchoalveolar immune cells in patients with
604   COVID-19. *Nat Med* **26**, 842-844 (2020).
605   34.   Steinmüller, C., Franke-Ullmann, G., Lohmann-Matthes, M.L. & Emmendörffer, A.
606   Local activation of nonspecific defense against a respiratory model infection by
607   application of interferon-gamma: comparison between rat alveolar and interstitial lung
608   macrophages. *Am J Respir Cell Mol Biol* **22**, 481-490 (2000).
609   35.   Cong, J. & Wei, H. Natural Killer Cells in the Lungs. *Front Immunol* **10**, 1416 (2019).
610   36.   Carlin, L.E., Hemann, E.A., Zacharias, Z.R., Heusel, J.W. & Legge, K.L. Natural Killer
611   Cell Recruitment to the Lung During Influenza A Virus Infection Is Dependent on
612   CXCR3, CCR5, and Virus Exposure Dose. *Front Immunol* **9**, 781 (2018).
613   37.   Vaccari, M.*, et al.* HIV vaccine candidate activation of hypoxia and the inflammasome in
614   CD14. *Nat Med* **24**, 847-856 (2018).
615   38.   National Research Project for SARS, B.G. The involvement of natural killer cells in the
616   pathogenesis of severe acute respiratory syndrome. *Am J Clin Pathol* **121**, 507-511
617   (2004).

39. Kuri-Cervantes, L.*, et al.* Immunologic perturbations in severe COVID-19/SARS-CoV-2
infection. *bioRxiv* (2020).
40. Schyns, J.*, et al.* Non-classical tissue monocytes and two functionally distinct populations
of interstitial macrophages populate the mouse lung. *Nat Commun* **10**, 3964 (2019).
41. Jiang, D.*, et al.* Regulation of pulmonary fibrosis by chemokine receptor CXCR3. *J Clin
Invest* **114**, 291-299 (2004).
42. Agostini, C.*, et al.* Cxcr3 and its ligand CXCL10 are expressed by inflammatory cells
infiltrating lung allografts and mediate chemotaxis of T cells at sites of rejection. *Am J
Pathol* **158**, 1703-1711 (2001).
43. Creelan, B.C. & Antonia, S.J. The NKG2A immune checkpoint - a new direction in
cancer immunotherapy. *Nat Rev Clin Oncol* **16**, 277-278 (2019).
44. Li, C.K.*, et al.* T cell responses to whole SARS coronavirus in humans. *J Immunol* **181**,
5490-5500 (2008).
45. Medoff, B.D.*, et al.* CD11b+ myeloid cells are the key mediators of Th2 cell homing into
the airway in allergic inflammation. *J Immunol* **182**, 623-635 (2009).
46. Alferink, J.*, et al.* Compartmentalized production of CCL17 in vivo: strong inducibility in
peripheral dendritic cells contrasts selective absence from the spleen. *J Exp Med* **197**,
585-599 (2003).
47. Woolsey, C.*, et al.* Establishment of an African green monkey model for COVID-19.
bioRxiv (2020).
48. Han, H.*, et al.* Profiling serum cytokines in COVID-19 patients reveals IL-6 and IL-10
are disease severity predictors. *Emerg Microbes Infect* **9**, 1123-1130 (2020).
49. Wickham, H. ggplot2: Elegant Graphics for Data Analysis. (Springer-
Verlag, New   York, 2016).
50. Kolde, R. pheatmap: Pretty Heatmaps (2018).
51. Team, R.C. R: A language and environment for statistical computing. . (Foundation for
Statistical Computing, Vienna, Austria, 2019).
52. Peterson, B.G. & Carl, P. PerformanceAnalytics: Econometric Tools for Performance
and Risk Analysis.   . (2020).
53. Kuhn, M., Jackson, S. & Cimentada, J. corrr: Correlations in R. (2020).
648
649

Figure Legend

Figure 1. Monocytes are a correlate of bronchoalveolar viral replication. (**A**) Schematic of

study design. Arrows represent time of inoculation with SARS-CoV-2 (day 0) or sacrifice (up to

28 days post infection). Unless stated otherwise, rhesus macaque data are shown in teal, and

African green monkey data are shown in magenta. (**B**) SARS-CoV-2 RNA levels in bronchial

brushes over time. (**C**) Association between SARS-CoV-2 load in bronchial brushes and ranking

of disease severity in all eight animals (most severe case = 1; mildest case = 8). (**D**) Changes in

monocyte count in blood before and after infection. (**E**) Proportion of total monocytes (HLA-

DR+ CD16+ and/or CD14+ out of CD45+ leukocytes) in PBMCs over time. (**F**) Correlation

between SARS-CoV-2 viral load in bronchial brushes and absolute count of monocytes in blood

at one-week post-infection (1wk pi). (**G**) Representative flow cytometry changes in non-

classical monocytes after the infection in blood (1wk pi). (**H**) Percent of classical (CD14+

CD16–) and (**I**) non classical monocytes (CD14– CD16+) out of total monocytes in PBMCs. (**J**)

Correlation between monocytes absolute numbers in blood and TARC (CCL17) in plasma (1wk

pi). (**K**) Relationship between fold change of chemokines and SARS-CoV-2 viral load in

bronchoalveolar brushes (1wk pi). (**L**) Heatmap displaying log2 fold-change of 8 chemokines

involved with traffic of monocytes to the lung. (1wk pi). Animals are ordered by ranking of

disease severity.

Figure 2. Myelomonocytic cells increase in BAL is associated with IL-6. (**A**) Decrease of

alveolar macrophages (HLA-DR+ CD206+ CD163+ cells) in the BAL after infection. (**B**)

Increase in the frequency of myelomonocytic infiltrates (HLA-DR+ CD206– CD163–) and (**C**)

of endothelial cells (HLA-DR+ CD206– CD163+) at infection in BAL (1wk pi). (**D**) tSNE plots

displaying kinetics of expression of CD16, CD11b and CD11c markers on live/ HLA-DR+ cells

populations in BAL before and after infection. (**E**) Gating of CD11c+ and CD11b+ populations

on the tSNE map based on antigen expression and discrete density clustering. (**F**)

Characterization of populations shown in (**E**). The heatmap depicts fold-change values of each

antigen compared to channel values of negative-expressing cell populations. (**G**) Increase in the

percentage of CD16+ HLA-DR+ cells and (**H**) in the percentage of interstitial macrophages

(CD16+ CD206– HLA-DR+) expressing of CD11c and (**I**) CD11b in BAL overtime.

Figure 3. Interstitial macrophages are increased in the lung of SARS-CoV-2 infected animals. (**A**) Macrophage infiltration varied from minimal to severe in both rhesus and AGM infected with SARS-CoV-2. (**B**) Fluorescent immunohistochemistry in the lung of AGM and RM showing infiltration of CD16+ CD11b+ CD206– macrophages (arrows), with lesser numbers of CD16+ CD206+ macrophages (arrowheads). Inset showing a higher magnification of CD16+ CD11b+ CD206–macrophages (arrows). Asterix shows macrophages multifocally surrounding small airways in GH99. White: DAPI (nuclei); Green: CD11b; Red: CD16; Blue: CD206.

Figure 4. Lymphocytes infiltration in the lungs is associated with disease severity. (**A**) Histopathologic findings showing infiltration of lymphocytes in the lung of AGM (NC40 and NC34) and RM (FR04 and GH99) with mild (NC40 and FR04), moderate (GH99), and severe disease (NC34). (**B**) Frequency of NKG2a+ cells in the blood or (**C**) BAL of 4 AGM and 4 RM at baseline, 2 weeks and 3 weeks post SARS-CoV-2 infection by flow cytometry. (**D**) Representative flow cytometry plot showing decreased frequency of NGK2a+ cells in PBMCs and increased frequency in BAL to baseline in at week 3 post SARS-CoV-2 infection respect to baseline. One animal for each species is shown. (**E**) Decrease in the frequency of CXCR3+ NKG2a+ cells in the blood at 2 weeks, and 3 weeks post SARS-CoV-2 infection compared to the baseline. (**F**) Absolute number of lymphocytes before and after infection. (**G**) tSNE plots displaying kinetics of PD-1 expression (in red) on lymphocytic cell populations over time. (**H**) Frequency of PD-1+ Th2 (CXCR3– CCR6–) CD4+ T cells is increased in RM and (**I**) in AGM overtime following the infection. (**J**) Changes in the percentage of Th2 (purple), Th1 (blue) and

Th17 (grey) cells within the total PD-1+ CD4+ T cell population. (**K**) Significant increase in the

percentage of PD-1+ CD8+ T cells in BAL.

Figure 5. Increased IL-6 in the absence of IL-10 is associated with disease progression.

Levels of IL-6 and IL-10 in plasma overtime (pg/ml) **(A)** and **(B)** correlation between the ratio of

IL-10 to IL-6 levels measured in plasma and the ranking of disease severity. (**C**) Levels of IL-6

and IL-10 after infection in plasma (pg/ml) in individual animals. (**D**) Summary of immune

events in lungs and blood of SARS-CoV-2 infected animals, from the 1-acute phase 2- switch to

Th2 type responses and 3- resolution phase.

715
716
717
718 **Table 1: Heterogeneous clinical and pathological outcomes following SARS-CoV-2 infection**

Animal ID	Clinical Findings	Clinical Score	Pathology Findings	Pathology Score	Ranking
NC34	Severe reduction in SpO$_2$, moderate tachypnea, and severe hypothermia just prior to euthanasia for severe respiratory distress at 8 days post infection. Less than 24 hours earlier, the only abnormality was a moderate reduction in SpO$_2$. Radiography appeared unremarkable until euthanasia when moderate to severe alveolar pattern with lobar sign and air bronchograms in right lung fields was observed.	9	Bronchointerstitial lung pattern in all lobes that was most severe in the right caudal lung lobe. Microscopically all lung lobes exhibited similar changes with the right caudal and right accessory lung lobe being the most severely affected. The lung lesions were comprised of diffuse alveolar damage with hyaline membrane formation, type II pneumocyte hyperplasia, and rare multinucleated giant syncytia. Some regions showed early evidence of organizing pneumonia and fibrosis.	19	1
NC33	Severe reduction in SpO$_2$, severe tachypnea, and severe hypothermia just prior to euthanasia for severe respiratory distress at 22 days post infection. Less than 24 hours earlier, the only abnormality was a mild reduction in SpO$_2$. Radiography appeared unremarkable until euthanasia when moderate to severe alveolar pattern in right middle/caudal lung fields and mild alveolar pattern left caudal lung fields were observed.	10	Severe diffuse alveolar damage of the right lower lung lobe. Other lobes exhibited suppurative alveolar infiltrate with lesser fibrin and edema. The right lung lobe contains peracute to acute lesions of necrosis, fibrin, and hemorrhage with minimal cellular infiltrate.	14	2
GH99	Transient cough beginning on 15 days post infection. Radiography showed significant changes at 11 days post infection which improved through 27 days post infection.	1	Severe bronchopneumonia of the right lower lung with granulomatous to pyogranulomatous inflammatory response.	10	3
NC38	Minor intermittent reductions in SpO$_2$	0	Mild interstitial pneumonia with low numbers of multinucleated giant cells and rare atypical pneumocyte hyperplasia.	8	4
HD09	Mild focal increase opacity ventral lung field at 11 days post infection.	1	Minimal to mild inflammation scattered throughout the lungs.	9	5
HB37	Intermittent mild to moderate pyrexia from 1-21 days post infection with mild dehydration from 14-21 days post infection.	1	Proliferative lymphocytic vasculitis and proliferative vasculopathy in the right middle lung lobe. Scattered interstitial inflammation throughout the remaining lung lobes.	5	6
FR04	Unremarkable	0	Bronchioles exhibited multifocal, mild mineralization.	6	7
NC40	Minor intermittent reductions in SpO$_2$	1	Minimal changes within lungs. Mild multifocal thickening of the alveolar septa in the right lower lobe associated with a mild inflammatory infiltrate.	4	8

719
720

721 Clinical findings are summarized for the duration of infection. Clinical scores, pathology findings,
722 and pathology scores reflect measurements at time of necropsy. Rankings were determined based
723 on a combination of the clinical findings and scores throughout the duration of infection and the
724 pathology findings and scores at time of necropsy (1 = severe; 8 = mild).
725 SpO_2 = oxygen saturation.
726
727

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Supplementary Table 1: Breakdown of routes of exposure and viral inoculation

Animal ID	Exposure (dose)	Age	Sex	Species	Base Weight (kg)
GH99	Cumulative dose of 3.61×10^6 PFU including oral, intratracheal, intranasal, and conjunctival (both eyes)	13.9	Male	*Macaca mulatta*, Indian ancestry	15.5
HD09		12.87	Female	*Macaca mulatta*, Indian ancestry	7.7
NC40		16 (est)	Male	*Chlorocebus aethiops*	7.0
NC33		16 (est)	Female	*Chlorocebus aethiops*	3.7
HB37	Aerosol (approx. 2.0×10^3 TCID$_{50}$)	12.96	Male	*Macaca mulatta*, Indian ancestry	11.9
FR04		14.93	Male	*Macaca mulatta*, Indian ancestry	12.3
NC34		16 (est)	Female	*Chlorocebus aethiops*	4.3
NC38		16 (est)	Male	*Chlorocebus aethiops*	7.4

Ages of monkeys NC40, NC33, NC34, and NC38 are approximate.

Supplementary Table 2: Virus loads in clinical specimens

Week 1

Specimen	HB37	FR04	NC38	NC34	NC33	NC40	GH99	HD09
Bronchial Brush	4.99E+05	1.64E+04	5.81E+06	3.22E+07	4.65E+06	1.42E+06	1.26E+07	1.23E+04
Buccal Swab	3.67E+03	4.74E+04	Not Detected	1.66E+06	Not Detected	1.07E+04	Inconclusive	2.55E+03
Nasal Swab	1.41E+06	1.52E+08	1.94E+04	6.45E+07	1.29E+05	1.05E+07	4.42E+04	1.42E+09
Pharyngeal Swab	5.03E+04	2.12E+06	2.47E+11	2.36E+09	7.00E+06	1.15E+08	Inconclusive	6.54E+06
Rectal Swab	3.81E+03	9.90E+05	4.95E+03	4.32E+03	1.73E+07	6.22E+08	8.22E+03	Not Detected
Vaginal Swab	Not Tested	Not Tested	Not Tested	3.03E+05	5.16E+04	Not Tested	Not Tested	Not Detected

Week 2

Specimen	HB37	FR04	NC38	NC34	NC33	NC40	GH99	HD09
Bronchial Brush	Inconclusive	2.46E+03	2.16E+04	Not Tested	2.92E+06	6.87E+03	1.95E+03	Inconclusive
Buccal Swab	Not Tested	1.71E+04	Inconclusive	Not Tested	6.40E+03	6.95E+03	3.78E+04	Not Detected
Nasal Swab	Not Tested	5.78E+06	2.07E+04	Not Tested	3.27E+06	9.54E+04	1.03E+05	2.89E+03
Pharyngeal Swab	3.77E+04	3.11E+03	5.61E+04	Not Tested	1.45E+03	9.21E+04	4.65E+03	Inconclusive
Rectal Swab	2.63E+04	1.55E+08	4.69E+05	Not Tested	3.93E+09	1.11E+05	2.71E+04	Not Detected
Vaginal Swab	Not Tested	Not Tested	Not Tested	Not Tested	8.92E+04	Not Tested	Not Tested	Not Detected

Week 3

Specimen	HB37	FR04	NC38	NC34	NC33	NC40	GH99	HD09
Bronchial Brush	Not Detected	Inconclusive	1.12E+05	Not Tested	3.04E+07	1.41E+06	Not Detected	Not Detected
Buccal Swab	Not Detected	Inconclusive	Inconclusive	Not Tested	Not Detected	Not Detected	Not Detected	Not Detected
Nasal Swab	Not Detected	7.53E+04	2.67E+04	Not Tested	2.77E+05	1.42E+07	2.03E+04	4.13E+04
Pharyngeal Swab	Not Tested	Not Detected	Not Detected	Not Tested	1.05E+04	8.95E+04	Inconclusive	4.28E+04
Rectal Swab	Not Detected	1.31E+04	3.78E+06	Not Tested	1.45E+04	3.45E+04	Not Detected	Not Detected
Vaginal Swab	Not Tested	Not Tested	Not Tested	Not Tested	Inconclusive	Not Tested	Not Tested	Not Detected

Samples which were not tested were due to lack of sample availability.

Supplementary Table 3: Rotation table derived from PCA

	PC1	PC2	PC3	PC4	PC5	PC6	PC7	PC8
Eotaxin	0.51349854	0.144318924	0.04547296	0.10908504	-0.2471479	0.22538446	-0.68436694	-0.3482608
IP-10	-0.25227747	0.559991260	-0.07652314	0.02243312	0.4106839	-0.16189141	0.05487606	-0.6469353
MCP-1	0.11727127	0.519454673	0.49526810	0.25731776	0.2000296	0.44221279	0.14981145	0.3832805
MCP-4	0.51986265	-0.002353372	0.07196805	-0.22069430	-0.2665758	0.11724746	0.67825244	-0.3619612
MDC	0.45844363	-0.050897556	0.21350047	0.37354660	0.2448526	-0.72839142	0.02537330	0.1047317
MIP-1a	0.15406887	0.251746497	-0.77979454	0.47174206	-0.1062296	0.09007799	0.16325863	0.1903004
MIP-1b	0.39295357	-0.047373837	-0.29263668	-0.51006598	0.6544815	0.14743768	-0.10172114	0.1926359
TARC	0.01858282	0.572304116	-0.04804510	-0.50043369	-0.3994979	-0.39152001	-0.09228374	0.3130139

All elements used for PCA shown.
PC = Principal Component

Supplementary Table 4: Antibodies used in flow cytometry and immunofluorescent microscopy panels

Antibodies used for flow cytometry

Marker	Fluorochrome	Clone	Ref.	Company	Known Reactivity
PD-1	APC	EH12.2H7	329908	Biolegend	RM & AGM
CD56	Alexa 700	B159	557919	BD	RM _ AGM not reported
CD183 (CXCR3)	APC/Cyanine7	G025H7	353722	Biolegend	RM & AGM
CD196 (CCR6)	BV605	G034E3	353420	Biolegend	RM & AGM
CD4	BV797	L200	563914	BD	RM & AGM
CD3	PE	SP34-2	552127	BD	RM & AGM
CD3	BV650	SP34-2	563918	BD	RM & AGM
CD20	BV650	2H7	563780	BD	RM & AGM
CD95	PE/Cyanine5	DX2	559773	BD	RM & AGM
CD8a	Texas Red	RPA-T8	MHCD0817	Life technology	RM & AGM
CD159a (NKG2a)	PE/Cyanine7	Z199	B10246	Beck Coulter	RM & AGM
CD16	BV711	3G8	563127	BD	RM & AGM
CD16	BUV496	3G8	564653	BD	RM & AGM
CD86	FITC	FUN-1	557343	BD	RM _ AGM not reported
CD11c	APC	3.9	17-0116-42	eBioscence	RM & AGM
HLA-DR	APC/Cyanine7	L243	307618	Biolegend	RM & AGM
CD14	V450	M5E2	561390	BD	RM & AGM
CD163	BV711	GHI/61	563889	BD	RM _ AGM not reported
CD184 (CXCR4)	PECF594	12G5	562389	BD	RM & AGM
CD45	BV786	D058-1283	563861	BD	RM & AGM
CX3CR1	PE	K0124E1	355704	Biolegend	RM _ AGM not reported
CD11b	PE/Cyanine5	ICRF44	301308	Biolegend	RM & AGM
CD206	PE/Cyanine7	19.2	25-2069-42	BD	RM & AGM

Antibodies used for immunofluorescent microscopy

Primary Antibody	Species	Company	Ref.	Dilution	Secondary Antibody
CD206	Goat	R&D	AF2535	1:50	Donkey anti-goat 647 (far red)
CD11b	Rabbit	Abcam	Ab133357	1:20	Donkey anti-rabbit 488 (green)
CD16	Mouse IgG2a	Novocastra	NCL-CD16	1:40	Permanent Red
DAPI	Dye - N/A	Invitrogen	D1306	1:20,000	N/A

Supplementary Figure 1. Kinetics of monocytes in Rhesus Macaques and African Green Monkeys. Association between SARS-CoV-2 load in bronchial brushes and ranking of disease severity in AGM (**A**) and RM (**B**) (most severe case = 1; mildest case = 8). Changes in monocyte count in blood before and after infection in AGM (**C**) and RM (**D**). Proportion of total monocytes (HLA-DR+ CD16+ and/or CD14+ out of CD45+ leukocytes) in PBMCs from AGM (**E**) and RM (**F**) over time. (**G**) Changes in the absolute number of neutrophils in blood over time in all eight animals. Correlation between SARS-CoV-2 viral load in bronchial brushes and absolute count of monocytes in blood at one-week post-infection (1wk pi) in AGM (**H**) and RM (**I**). Percent of classical (CD14+ CD16–) monocytes (CD14– CD16+) out of total monocytes in PBMCs from AGM (**J**) or RM (**K**). Percent of non classical monocytes (CD14– CD16+) out of total monocytes in PBMCs in AGM (**L**) or RM (**M**). (**N**) Percent of intermediate monocytes (CD14+ CD16+) out of total monocytes in PBMCs from all eight animals over time.

Supplementary Figure 2. Chemokines promote cell migration of monocytes from the blood to the lung following infection. Levels of chemokines in the peripheral blood over time, including (**A**) Eotaxin, (**B**) IP-10, (**C**) MCP-1 (CCL2), (**D**) MCP-4 (CCL13), (**E**) MDC (CCL22), (**F**) MIP-1a, (**G**), MIP-1b, and (**H**) TARC (CCL17). (**I**) PCA plot comparing baseline levels of chemokines between rhesus macaques and African Green monkeys. (**J**) TARC (CCL17) levels and (**K**) IP-10 levels in blood and SARS-CoV-2 viral load in bronchial brushes one week post-infection.

Supplementary Figure 3. Changes in the frequency of alveolar macrophages and infiltrates in BAL during infection by flow cytometric analysis. (**A**) **Left panel.** Representative light scatter (forward scatter and side scatter) analysis of lymphocytes, macrophages, and polymorphonuclear cells (PMNC) in BAL. Cell populations are located according to their relative size (forward scatter) and granularity (side scatter). **Right panels.** Overlaid histograms showing fluorescent intensities of CD206 and HLA-DR of lymphocytes, macrophages, and PMNC as shown on the left panels. (**B**) Representative pseudocolor dot plot displaying alveolar macrophages, endothelial cells, monocytes, and/or dendritic cells distinguished by differential expression of CD163 and CD206. (**C**) Range of phenotypes of CD11b+ and CD11c+ myeloid cells in BAL and their change in proportion out of total HLA-DR+ cells over time. (**D**) tSNE plot displaying kinetics of CD86 expression on total HLA-DR+ cells after infection. Percent of CD16+ HLA-DR+ CD206- cells in BAL over time in AGM (**E**) and RM (**F**). (**G**) Representative gating strategy showing classification of CD45+HLA-DR+CD16+ myeloid cells which express either CD11c or CD11b. Top panel showing a sample with CD11c+ cells, and bottom panel showing a sample without CD11c+ cells used to help determine optimal gate position.

Supplementary Figure 4. Association of myeloid subsets in the BAL with IL-6 plasma levels.
Correlation plot between myeloid subsets in BAL and levels of IL-6 (pg/mL) in the blood 3 weeks post infection using Pearson's method. Numbers inside boxes on the diagonal right half of the correlation plot represent the coefficient of correlation (*r*) and asterisks denote significance. One red asterisk corresponds with p <. 05, two red asterisks correspond with p < .01, and three red asterisks correspond with p < .001. Correlation plots in the bottom left display the actual data points.

Supplementary Figure 5. Lymphocyte response to SARS-CoV-2 in Rhesus Macaques and African Green Monkeys. Percent of NKG2a+ on CD3- cells in BAL in AGM (**A**) and RM (**B**). Changes in lymphocyte count in blood before and after infection in AGM (**C**) and RM (**D**). (**E**) Median levels of IP-10, IFN-γ, IL-13, and IL-5 in plasma (pg/mL) from all eight animals over time. (**F**) Correlation between IL-4 levels in the plasma (pg/mL) and percent of PD-1+ Th2 cells in the blood at week 3 post infection. Proportion of PD-1+ CD8+ T cells in BAL from AGM (**G**) and RM (**H**) after infection.