Title: Arc1 modulates microbiota-responsive developmental and metabolic traits in

Drosophila

Running Title: Arc1 and *Acetobacter* promote fly growth

Keywords: *Drosophila*, microbiota, *Acetobacter*, Arc1, insulin signaling, metabolism

Scott A. Keith*, Cassandra Bishop*[†], Samantha Fallacaro*[†], Brooke M. McCartney*

* Department of Biological Sciences, Carnegie Mellon University, Pittsburgh, PA 15213

[†] Equal contribution

Corresponding author:

Brooke M. McCartney

600D Mellon Institute

Department of Biological Sciences

Carnegie Mellon University

4400 Fifth Avenue

Pittsburgh, PA 15213

Phone: (412) 268-5195

Email: bmccartney@cmu.edu

SUMMARY

Drosophila Arc1 exhibits microbiota-dependent, tissue-specific differential expression, mitigates the impacts of germ-free rearing on insulin signaling and growth rate, but is dispensable for metabolic homeostasis in *Acetobacter*-colonized flies.

ABSTRACT

Perturbations to animal-associated microbial communities (the microbiota) have deleterious effects on various aspects of host fitness, including dysregulated energy metabolism. However, the molecular processes underlying these microbial impacts on the host are poorly understood. In this study, we identify a novel connection between the microbiota and the neuronal factor Arc1 that affects metabolism and development in *Drosophila*. We find that *Arc1* exhibits tissue-specific microbiota-dependent expression changes, and that flies bearing a null mutation of *Arc1* complete larval development at a dramatically slowed rate compared to wild-type animals. In contrast, monoassociation with a single *Acetobacter sp.* isolate was sufficient to enable *Arc1* mutants to develop at a wild-type rate. These developmental phenotypes are highly sensitive to composition of the larval diet, suggesting the growth rate defects of GF flies lacking Arc1 reflect metabolic dysregulation. Additionally, we show that pre-conditioning the larval diet with *Acetobacter sp.* partially accelerates *Arc1* mutant development, but live bacteria are required for the full growth rate promoting effect. Finally, GF *Arc1* mutants display multiple traits consistent with reduced insulin signaling activity that are reverted by association with *Acetobacter sp.*, suggesting a potential mechanism underlying the microbe-dependent developmental phenotypes. Our results reveal a novel role for Arc1 in modulating insulin

signaling, metabolic homeostasis, and growth rate that is specific to the host's microbial

and nutritional environment.

INTRODUCTION

The physiology and life history traits of animals are shaped in remarkable ways by

interactions with commensal and beneficial microorganisms (the microbiota). For many

metazoans, microbial symbionts play integral roles in post-embryonic development and

physiology to yield fit and fertile adults (McFall-Ngai et al., 2013; Robertson et al., 2019).

Thus, perturbations to the microbiota can have profoundly deleterious consequences for

its animal host. For example, germ free (GF) or antibiotic-treated mice exhibit decreased

body fat (Smith et al., 2007), abnormal intestinal epithelial architecture (Hayes et al.,

2018), fewer differentiated immune cells (Ekmekciu et al., 2017), and

neurodevelopmental defects (Sampson and Mazmanian, 2015). In humans, dysbiosis of

the gut microbiota has been implicated in the pathogenesis of a wide range of disorders,

such as Type 2 diabetes (Larsen et al., 2010), obesity (Shen et al., 2013), and autism

(Gilbert et al., 2013). However, the molecular factors that actuate microbial influence on

host physiology and development are not comprehensively understood. *Drosophila*

melanogaster and its gut microbiota are an ideal model to discover such factors given

Drosophila's extensive genetic resources, the low-diversity and readily-cultured bacterial

communities associated with laboratory fly cultures, and the consequent technical ease

of generating GF and gnotobiotic fly populations (Broderick and Lemaitre, 2012; Douglas,

2018; Martino et al., 2017).

From a screen to discover microbiota-responsive neuronal genes, we identified

Drosophila Activity-regulated cytoskeleton associated protein 1 (*Arc1*) as being

differentially expressed in GF flies. *Arc1* is a *Drosophila* homolog of mammalian

Arc/Arg3.1, an immediate early gene and master regulator of synaptic plasticity in the

brain (Carmichael and Henley, 2018; Shepherd and Bear, 2011). Arc transcription is

highly upregulated by synaptic activity and while the brain is encoding novel information

into neural circuits (Chen et al., 2020; Guzowski et al., 1999). Accordingly, reduced Arc

expression impairs memory formation and learning ability in rodents (Guzowski et al.,

2000; Shandilya and Gautam, 2020), and defects in human Arc function have been linked

to a variety of neurological and neurodevelopmental disorders, including Alzheimer's

disease (Bi et al., 2018), autism spectrum disorders (Alhowikan, 2016), and schizophrenia

(Fromer et al., 2014). Arc partially regulates synaptic plasticity by interacting with

endocytic machinery to regulate synaptic surface presentation of AMPA-type glutamate

receptors (Chowdhury et al., 2006; DaSilva et al., 2016; Wall and Corrêa, 2018). Both

mammalian *Arc* and fly *Arc1* encode retroviral group-specific antigen-like amino acid

sequences, and are predicted to have independently derived from ancient Ty3/Gypsy

retrotransposons (Ashley et al., 2018; Campillos et al., 2006; Cottee et al., 2019;

Pastuzyn et al., 2018). Recently, it was shown that Arc and Arc1 proteins can self-

assemble into capsid-like structures that package and transport mRNAs (including

Arc/Arc1 mRNA) into cultured neuronal cell lines and across synapses *in vivo*, constituting

a novel mechanism of cell-cell communication (Ashley et al., 2018; Erlendsson et al.,

2019; Pastuzyn et al., 2018).

As with mammalian *Arc*, *Drosophila Arc1* expression is strongly upregulated by

neuronal activation (Guan et al., 2005; Mattaliano et al., 2007; Montana and Littleton,

2006; Mosher et al., 2015). At the larval neuromuscular junction (NMJ), Arc1 protein-

mediated transfer of *Arc1* mRNA from the boutons of motoneurons to post-synaptic

myocytes appears to be required for appropriate synapse maturation and synaptic

plasticity (Ashley et al., 2018). Further, *Arc1* loss-of-function mutants have elevated fat

levels, altered metabolomic profiles, are starvation resistant, and differentially express a

repertoire of enzymes involved in central carbon metabolism (Mattaliano et al., 2007;

Mosher et al., 2015). These phenotypes may be at least partially attributable to Arc1

activity in the brain, where its expression in adults is concentrated in a subset of large

neurons in the pars intercerebralis (Mattaliano et al., 2007). However, the precise cellular

pathways and mechanisms through which it affects metabolism are not known.

In *Drosophila,* metabolic functions directly impact developmental timing and whole-

organism growth that occurs exclusively during the larval stages (Edgar, 2006). Fly larvae

progress through three developmental instars, increasing in size approximately 200-fold

prior to pupariation and metamorphosis (Robertson, 1963). Both the timing of

developmental progression and the magnitude of larval growth are genetically regulated

by multiple intersecting nutrient-responsive inter-organ signaling pathways, including the

insulin/insulin-like growth factor (IIS) pathway (Brogiolo et al., 2001; Rulifson et al., 2002),

the 20-hydroxyecdysone biosynthetic pathway (Buhler et al., 2018; McBrayer et al.,

2007), and target-of-rapamycin (TOR) signaling (Colombani et al., 2003; Layalle et al.,

2008). IIS and TOR signaling are highly functionally conserved between *Drosophila* and

vertebrates (Edgar, 2006; Gilbert, 2008). The microbiota also has a significant impact on

fly metabolism; GF larvae exhibit prolonged larval development and stunted growth

compared to conventionally-reared or gnotobiotic flies (Newell and Douglas, 2014; Shin

et al., 2011; Storelli et al., 2011; Storelli et al., 2017; Wong et al., 2014). Commensal

bacteria appear to impact Drosophila growth through numerous mechanisms, including

nutritional provisioning and activation of host signaling pathways via secreted metabolites

(Chaston et al., 2014; Consuegra et al., 2020; Kamareddine et al., 2018; Keebaugh et al.,

2018; Matos et al., 2017; Sannino et al., 2018; Shin et al., 2011). However, many gaps

remain in our understanding of the molecular mechanisms linking the bacterial microbiota

to larval metabolic function and growth dynamics.

Here we reveal *Arc1* as a novel host factor that modulates the microbiota's impact

on larval growth. *Arc1* is transcriptionally altered in tissue-specific patterns in GF flies,

and loss of *Arc1* dramatically exacerbates the developmental growth delay of GF larvae.

We further show that a single *Acetobacter* isolate is sufficient to restore normal larval

development in *Arc1* mutants, and that pre-conditioning the larval diet with this

Acetobacter sp. partly restores a normal growth rate to GF *Arc1* mutants. We further show

that loss of both *Arc1* and the microbiota results in multiple traits consistent with reduced

insulin/insulin-like growth factor (IIS) pathway activity, and that monoassociation with

Acetobacter sp. reduces the severity of these phenotypes. Together our data reveal an

experimental system wherein a single microbiota member supports the health of a

metabolically destabilized host genotype. Further, this work demonstrates a previously

unrecognized role for *Arc1* in altering insulin signaling in flies following microbiota

removal.

RESULTS

The microbiota promote larval development of *Arc1*-deficient *Drosophila* hosts

To identify host neuronal molecules and pathways that may be impacted by the

microbiota, we conducted a transcriptomic screen to identify *Drosophila* genes that are

differentially expressed in adult fly heads upon elimination of the bacterial microbiota.

From this work, we identified *Arc1* among the genes that were most responsive to host

microbial condition (Keith et al., 2019, bioRxiv). Specifically, we found that *Arc1*

transcripts are elevated in the heads of adult wild-type *Drosophila* grown under germ-free

conditions (GF; microbiologically sterile) compared to flies grown in gnotobiotic (GNO)

polyassociation with a four-species bacterial community consisting of two *Acetobacter*

(*Acetobacter sp.*, *A. pasteurianus*) and two *Lactobacillus* (*L. brevis*, *L. plantarum*) isolates

(Figure 1A). Notably, in published RNA-seq datasets comparing the gut transcriptomes

of microbiota-associated *Drosophila* to GF guts, *Arc1* is among the significantly

differentially expressed genes (Bost et al., 2017; Dobson et al., 2016; Guo et al., 2014;

Petkau et al., 2017); consistent with these studies, we found that *Arc1* is transcriptionally

decreased in the adult GF gut (Figure 1A). These microbiota-sensitive transcript level

changes occur in multiple wild-type fly lines, but both changes were not observed in every

wild type line (Figure 1A). Together our data demonstrate that *Arc1* is a microbiota-

responsive gene, and the transcript response exhibits tissue specificity.

Arc1 regulates lipid homeostasis and central carbon metabolite levels in

Drosophila larvae (Mosher et al., 2015), and an *Arc1* loss of function mutant exhibits

enhanced starvation resistance in adult flies (Mattaliano et al., 2007). Intriguingly, recent

studies have shown that these and other metabolic traits are also impacted by the

microbiota, depending on dietary conditions and host genetic background (reviewed in

Douglas, 2018). One of the major organism-level consequences of GF-induced metabolic

dysregulation in *Drosophila* is a prolonged larval growth period (Strigini and Leulier,

2016). Thus, to test for a physiologically relevant interaction between the microbiota and

Arc1 function, we raised wild-type flies (w^{1118}) and a fly line bearing a CRISPR-mediated

deletion of *Arc1* (*Arc1^{E8}*; Ashley et al., 2018) from embryo to adulthood either GF or GNO,

and monitored their larval growth rate. Consistent with many previous reports (Chaston

et al., 2014; Consuegra et al., 2020; Erkosar et al., 2015; Kamareddine et al., 2018;

Keebaugh et al., 2018; Newell and Douglas, 2014; Sannino et al., 2018; Shin et al., 2011;

Storelli et al., 2011; Storelli et al., 2017; Wong et al., 2014), we found that GNO wild-type

animals completed larval development faster than their GF siblings (Figure 1B). On our

laboratory's routine diet (see Material and Methods) at 23ºC, wild-type GNO cultures take

~7 days to complete larval development, and this period is extended by ~1.5 days under

GF conditions (Figure 1B, Table S1). Strikingly, the larval growth delay induced by

microbiota elimination was dramatically extended in larvae lacking *Arc1*. While *Arc1^{E8}*

mutant larvae grown under GNO conditions developed at a rate indistinguishable from

wild-type GNO larvae, GF *Arc1^{E8}* animals took on average ~12 days to complete larval

development (Figure 1B). These differences in larval growth rate were reflected in the

rate of adult emergence, with wild-type and *Arc1^{E8}* GNO animals eclosing ~12.5 days

after embryo deposition, wild-type GF eclosing after ~14 days, and *Arc1^{E8}* GF adults

emerging asynchronously between 16-20 days (Figure S1A). We observed a similar,

though less protracted, developmental delay for two independently-generated *Arc1* loss-

of-function alleles (*Arc1^{esm18}* and *Arc1^{esm113}*; Mattaliano et al., 2007) under GF conditions

(Figures S1B, C), and in GF animals transheterozygous for *Arc1^{E8}* and *Arc1^{esm113}* (Figure

1C). Further, ablation of *Arc1*-expressing cells using *Arc1-GAL4* (Mattaliano et al., 2007)

to drive expression of the pro-apoptotic *reaper* also extended the developmental period

of GF larvae (Figure 1D). All larvae bearing loss of function mutations in *Arc1* or ablation

of *Arc1* expressing cells developed at the wild type rate when grown with the GNO

bacterial community. *Arc2* is another *Drosophila* Arc homolog in a genomic locus adjacent

to *Arc1*; the two proteins likely represent an ancestral duplication event (Pastuzyn et al.,

2018). In contrast to *Arc1* depletion, a P-element insertion in the *Arc2* 3' UTR which

decreased *Arc2* expression (Figure S1D) had no effect on the developmental rate of GF

larvae (Figure S1E). Together these results suggest a reciprocal host gene-microbe

interaction: the bacterial microbiota facilitates the appropriate larval growth rate of *Arc1*-

deficient larvae, and Arc1 is a novel host protein that modulates the physiological

response to microbiota loss during larval development.

Monoassociation with *Acetobacter sp.* is sufficient to promote larval development

of *Arc1* mutant larvae

We next asked whether the ability of our polymicrobial GNO community to promote

larval growth rate was attributable to the effects of individual bacterial taxa or collective

effects of the community. To this end, we generated wild-type and *Arc1* null fly cultures

in monoassociation with each of the four bacteria and measured time to complete larval

development. Wild-type larvae monoassociated with each of the four bacteria developed

significantly faster than their GF siblings, with the two *Acetobacter* isolates promoting

slightly faster development than the two *Lactobacillus* isolates (Figure 2A). In contrast,

only $Arc1^{E8}$ mutant larvae monoassociated with *Acetobacter sp.* developed at an identical

rate to polyassociated GNO larvae; *Arc1* mutants associated with *A. pasteurianus*, *L.*

brevis, and *L. plantarum* all developed at the same rate, ~2 days faster than GF, but ~2.5-

3 days slower than GNO and *Acetobacter sp.* monoassociated larvae (Figure 2B). While

we did find significant differences in bacterial load among the four isolates, these

differences did not correlate with the impact of each bacteria on host developmental rate

for either genotype (Figure S2).

Because *Acetobacter sp.* alone was sufficient to enable *Arc1^{E8}* larvae to achieve

a wild-type growth rate, we asked whether it would also be necessary in the four-species

GNO community. Surprisingly, while mutant larvae associated with *A. pasteurianus*, *L.*

brevis, and *L. plantarum* alone developed substantially slower than those associated with

the four-species group or *Acetobacter sp.*, the three together in a GNO community lacking

Acetobacter sp. were sufficient to promote normal development. However, for

experimental tractability we focused all subsequent investigation on wild type and *Arc1*

mutants monoassociated with *Acetobacter sp.*.

While the delayed pupariation rate of wild-type GF larvae reflects a moderately

extended duration of all three larval instars (Figure 2C; Storelli et al., 2011), *Arc1^{E8}* GF

larvae undergo substantially prolonged L1 and L2 phases (Figure 2C). Further, GF

animals of both genotypes increase in size at a more gradual rate than those with

Acetobacter sp., but this effect is highly exaggerated in *Arc1* mutants, suggesting a longer

time to attain the critical weights that trigger each molt and metamorphosis (Figure 2D;

Mirth et al., 2005). The extended larval period could reflect reduced food consumption,

but we did not observe feeding differences in *Arc1*-deficient larvae under either microbial

condition (Figure S3). Along with a lengthened time to pupariation, both wild-type and

Arc1^{E8} GF larvae develop into smaller pupae, but, as with the developmental delay, this

size reduction is greater for animals lacking *Arc1* (Figure 2E). These data suggest that

loss of *Arc1* exacerbates the effects of GF rearing on both larval growth rate and growth

capacity.

Host diet can modulate the *Arc1*-dependent GF larval developmental delay

The developmental rate of wild type GF larvae is particularly sensitive to the concentration of dietary yeast, the primary source of ingested amino acids: the developmental lag of GF compared to conventional (CV) or GNO animals increases on diets with low yeast content, while feeding high-yeast diets enables GF larvae to develop at the same rate as microbe-associated larvae (Shin et al., 2011; Storelli et al., 2011; Wong et al., 2014). To determine how the loss of Arc1 impacts the dietary sensitivity of GF larvae, we reared *Acetobacter sp.*-associated and GF wild-type and *Arc1^{E8}* larvae on nine simplified diets containing systematically varied concentrations of dextrose and yeast, and monitored time to pupariation (Figure 3).

As expected, in wild-type animals "high-yeast" (10%) diets eliminated the developmental gap between *Acetobacter sp.*-associated and GF larvae, regardless of glucose concentration, while on "low-yeast" (3%) diets wild-type GF larvae were consistently delayed. The exception to this was the 3% yeast-10% glucose diet, which substantially slowed the developmental rate of all conditions, a known effect of high-sugar, low-protein diets (Musselman et al., 2011; Wong et al., 2014). Interestingly, on 5% yeast diets, increasing glucose concentration moderately accelerated the development of wild-type GF larvae, with a statistically significant delay observed only at the lowest glucose level. Overall, these data are congruent with published observations that GF rearing predominantly sensitizes wild-type flies to reductions in dietary protein content.

GF larvae lacking *Arc1* generally showed enhanced developmental sensitivity to dietary composition; GF *Arc1^{E8}* animals developed more slowly than GF wild-type on five out of the nine tested diets (Figure 3). Notably, GF *Arc1^{E8}* larvae were even delayed on

two "high-yeast" diets (10% yeast-3% glucose and 10%yeast-10% glucose), where the

wild type GF larvae developed at same rate as wild type *Acetobacter sp.* animals*.* On

most diets, *Arc1* mutants associated with *Acetobacter sp.* developed at the same rate as

wild-type *Acetobacter sp.*-associated larvae. One notable exception was the 3% yeast-

5% glucose diet, where *Acetobacter sp.* failed to have any growth rate promoting activity

for *Arc1^{E8}* larvae.

Importantly, while *Arc1*-deficient larvae grown GF still displayed a lengthened

duration of larval development on multiple diets, none of the tested formulations yielded

a magnitude of delay comparable to that observed on our laboratory's routine diet (Figure

1B). To compare the nutritional contents of the yeast-glucose diets to our cornmeal-yeast-

molasses diet (see Materials and Methods), we utilized the *Drosophila* Dietary

Composition Calculator to determine the protein:carbohydrate (P:C) ratio of each recipe

(Figure 3; Lesperance and Broderick, 2020). Relative to most of the tested yeast-glucose

diets, our laboratory diet has a low P:C ratio of 0.06. Interestingly, the yeast-glucose diet

with the P:C ratio closest to that of our diet (3% yeast-10% glucose; P:C 0.09) did not

recapitulate the microbe-dependent developmental rate trends observed on our diet;

specifically, *Acetobacter sp.*-associated wild-type and *Arc1* mutant larvae developed

substantially slower on this yeast-glucose diet than on our diet. This may indicate that the

full growth rate promoting effects of this *Acetobacter sp.* are specific to the more complex

nutritional substrate of the cornmeal-yeast-molasses diet and less dependent on the

relative proportions of these two macronutrients. For example, cornmeal provides sugars

in the form of complex polysaccharides that require a multi-step breakdown process

involving amylases and maltases, which have previously been shown to be microbiota

responsive in *Drosophila* larvae (Erkosar et al., 2017). Additionally, differences in

Acetobacter sp. developmental support could be due to different levels of bacterial

population growth on the various diets.

Together, these results indicate that the importance of *Arc1* function during GF

larval development is highly dependent on the host's nutritional environment, and suggest

that *Arc1* may be particularly important for GF animals' metabolic responses to dietary

amino acid availability.

Live *Acetobacter sp.* populations are required for optimal *Arc1* mutant

developmental rate

We next sought to investigate the mechanisms by which *Acetobacter sp.*

association enables *Arc1*-deficient animals to complete larval development at the same

rate as wild-type monoassociated animals. Given that increasing the concentration of

dietary yeast can substantially diminish the prolonged larval stage of GF *Arc1* mutant

larvae, we hypothesized that bacteria consumed with the diet may serve as a

supplemental food source supporting *Arc1^{E8}* development, a mechanism for which there

is precedent in wild-type *Drosophila* (Bing et al., 2018; Keebaugh et al., 2018; Storelli et

al., 2017). To test this prediction, we inoculated GF wild-type and *Arc1^{E8}* cultures with

heat-killed *Acetobacter sp.* cells daily throughout the larval developmental period until the

entire population pupariated. Administration of heat-killed *Acetobacter* resulted in a slight,

but not statistically significant, acceleration of developmental rate for GF *Arc1* mutants

(Figure 4A). Feeding with dead *Acetobacter* cells had no effect on wild-type GF larval

developmental rate (Figure 4A). These data suggest *Acetobacter sp.* does not solely

serve as a nutritional supplement to promote *Arc1* mutant development.

The *Acetobacteraceae* are distinguished by the ability to generate acetic acid

through partial oxidation of ethanol and other substrates (Saichana et al., 2015). Two

studies showed that acetic acid/acetate consumption enables GF larvae or larvae

associated with an *Acetobacter* mutant deficient for acetic acid production to develop at

a rate comparable to those associated with live bacteria (Kamareddine et al., 2018; Shin

et al., 2011); in contrast, acetic acid has also been reported to yield no effect on growth

in a GF context, or, at higher concentrations, further extends the GF larval growth delay

(Kim et al., 2017; Shin et al., 2011). Consistent with the latter results, we found that

providing GF wild type or *Arc1^{E8}* larvae with dietary acetic acid at different concentrations

either had no impact on growth rate, or increased the duration of larval development

(Figure 4C, C'). This suggests that any acetic acid generation by our *Acetobacter sp.*

isolate has minimal impact on larval growth under our dietary and husbandry conditions.

We also found that daily inoculation with filtered supernatant from planktonic *Acetobacter*

sp. cultures had no effect on either wild-type or *Arc1^{E8}* GF larvae, providing no evidence

to support a role for metabolite(s) secreted in *Acetobacter sp.* planktonic culture in

promoting larval growth rate (Figure 4B).

The association of larval and adult *Drosophila* with their commensals involves

bacterial proliferation on the flies' diet and consequent continual ingestion of live and dead

bacterial cells associated with the food bolus (Ludington and Ja, 2020). The host and its

microbial partners therefore share a dietary niche, with the bacteria utilizing the flies' food

as a carbon source (Blum et al., 2013; Martino et al., 2018; Storelli et al., 2017). Prior

work suggests that certain microbe-affected metabolic traits in *Drosophila*, including

growth rate, result from bacterial utilization of dietary nutrients (Consuegra et al., 2020;

Huang and Douglas, 2015; Storelli et al., 2017). These microbiota-dependent host traits

therefore arise from bacterial modification of the flies' diet, resulting in an altered

nutritional intake. We hypothesized that bacterial dietary modification might underlie

Acetobacter sp. support of *Arc1* mutant development. To test this hypothesis, we

inoculated GF food vials with *Acetobacter sp.* planktonic culture, allowed colonization of

the diet for five days, and then heat-treated the vials to kill all bacteria, resulting in an

Acetobacter sp.-conditioned but microbiologically sterile larval food substrate.

When fed the conditioned diet, wild-type GF larvae also developed slightly, though

not significantly, faster than wild-type GF larvae on regular sterile food, and on un-

inoculated sterile food subjected to the same heat treatment used to sterilize conditioned

media (Figure 4D). In contrast, *Acetobacter sp.*-conditioned food substantially

accelerated GF *Arc1^{E8}* larval development compared to GF larvae on untreated and

heated control diets, though still ~1.5 days slower than *Arc1* mutants associated with live

Acetobacter sp. (Figure 4D'). These data suggest that *Acetobacter sp.* modification of the

larval diet may be an important, though not exclusive, mechanism by which this bacterial

isolate promotes larval growth rate.

Germ free rearing alters metabolic defects of *Arc1* mutant animals and reduces

insulin signaling

The rate of larval growth and development in *Drosophila* is an organism-level

readout of systemic energy metabolism. Specifically, growth is promoted by the activation

of multiple circulating-hormone-regulated signaling pathways, including the

insulin/insulin-like signaling (IIS) pathway, which are responsive to and downstream of

nutrient intake (Edgar, 2006). Following food consumption, insulin-like peptides (Ilps) are

synthesized by neuroendocrine cells in the brain and released into the hemolymph

(Géminard et al., 2009). Ilps then circulate to principal metabolic tissues, including the fat

body, and activate the insulin receptor (InR; Brogiolo et al., 2001; DiAngelo and Birnbaum,

2009). This results in the nuclear exclusion of the transcription factor Foxo, the

suppression of starvation responses, and growth promotion (Jünger et al., 2003; Kramer

et al., 2003). Decreased IIS activity results in a larval growth delay, or in the most severe

cases a complete growth arrest (reviewed in Baker and Thummel, 2007). Furthermore, in

wild-type animals fed nutrient restrictive diets, individual commensal bacteria, including

some *Acetobacter*, may promote larval growth and development in part via activation of

host IIS (Kamareddine et al., 2018; Shin et al., 2011; Storelli et al., 2011). *Arc1* has been

previously implicated in metabolic homeostasis (Mosher et al., 2015), but not IIS signaling

(Mattaliano et al., 2007). Further, we have shown that the growth rate of GF *Arc1* mutants

is delayed and sensitive to dietary composition (Figure 3), consistent with metabolic

dysregulation. Thus, we hypothesized that the developmental delay of *Arc1^{E8}* GF larvae

is a consequence of reduced IIS activity, and that *Acetobacter sp.* supports an appropriate

Arc1^{E8} growth rate by restoring proper IIS signaling.

In addition to a reduction in larval growth rate, loss of IIS signaling results in other

growth defects. This includes a reduction in wing size, due to a reduction in cell number

and cell size (Figure 5A; Rulifson et al., 2002). While wild-type GF larvae have a

decreased larval growth rate (Figure 1B), and final larval size (Figure 2E), we did not

observe any alteration to wing growth properties: cell size (inferred by cell density), cell

number, and wing size were indistinguishable from that of *Acetobacter sp.*-associated

wild type animals (Fig. 5B-D). This may suggest that larval growth rate is particularly

sensitive to metabolic perturbations under our culture conditions.

Arc1 mutants grown under GF conditions exhibited a significant reduction in cell

size, consistent with reduced IIS activity, whereas *Acetobacter sp.*-associated mutant

wings had wild-type sized cells (Figure 5A,B). Surprisingly, though, loss of Arc1 increased

cell number, regardless of microbial condition (Figure 5A,C). Since cell size was

unaffected in *Arc1^{E8}-Actobacter sp.* flies, this resulted in larger wings for these animals,

whereas GF *Arc1^{E8}* wings were the same size as wild-type wings, due to their reduced

cell size (Figure 5A,D). While increased IIS activity is associated with larger wings, this is

accompanied by an increase in cell size (Brogiolo et al., 2001). This suggests that Arc1

may also contribute to wing growth through unknown additional mechanisms.

In addition to growth changes, loss of IIS pathway activity also results in resistance

to adult starvation conditions (Broughton et al., 2005). *Arc1^{esm18}* mutants are likewise

starvation resistant (Mattaliano et al., 2007). We observed the same phenotype, with

Arc1^{E8} Acetobacter sp. females taking longer to succumb to starvation than wild type

Acetobacter sp. and wild type GF adults. This starvation resistance was strongly

enhanced in GF *Arc1^{E8}* animals, which survived full nutrient deprivation ~two days longer

than wild type animals (Figure 5E). Reduced mobilization of energy stores under

starvation conditions is another consequence of reduced IIS activity (Broughton et al.,

2005; Luong et al., 2006). Thus, the strongly enhanced starvation survival of GF *Arc1*

mutants, are consistent with deficient IIS function in the absence of *Arc1* and the

microbiota.

Activation of the insulin receptor (InR) results in the activation of PI3K increasing

the membrane phospholipid PIP_3 (Britton et al., 2002). This produces a membrane-

associated binding site for proteins containing the Plextrin homology (PH) domain,

including the IIS effector Akt (Verdu et al., 1999). tGPH is a GFP-tagged PH domain that

has been widely used as an *in vivo* reporter of PI3K activity and IIS pathway activation

(Britton et al., 2002). In *Acetobacter sp.*-associated wild-type and *Arc1^{E8}* larvae and in

wild-type GF larvae, we observed a strong membrane localization of tGPH in the fat body

that was indistinguishable between those genotypes and conditions (Fig. 5F), suggesting

that IIS signaling is not significantly perturbed in these larvae. In contrast, we found that

the membrane localization of tGPH was strongly reduced in the larval fat body of GF

Arc1^{E8} animals (Fig. 5F), indicative of reduced IIS pathway activation. Together these

data demonstrate that aside from the larval growth delay, wild type GF larvae do not have

phenotypes associated with reduced IIS signaling. However, when *Arc1* is also

eliminated, these animals exhibit many hallmarks of IIS deficiency. Likewise, *Arc1* mutant

Acetobacter sp. animals generally do not display evidence of IIS signaling defects. But

the starvation resistance of *Arc1^{E8}* animals suggests that *Arc1* mutants may be sensitized

to reduced IIS signaling, which is exacerbated by GF rearing. Thus, *Acetobacter sp.* is

largely sufficient to promote IIS activity in *Arc1* mutants, suggesting a potential

mechanism by which *Acetobacter sp.* may promote organismal homeostasis in *Arc1*-

deficient hosts.

DISCUSSION

Here we reveal a novel and unexpected connection between the host gene *Arc1*

and the bacterial microbiota in *Drosophila*: we demonstrate that *Arc1* transcript levels are

responsive to the microbiota, and find that Arc1 functions to suppress metabolic and

fitness deficiencies induced by microbiota removal. *Arc1* exhibits microbiota-dependent

expression, as its transcript levels are elevated or repressed in a tissue-specific manner

in GF animals. Under our standard dietary conditions, GF rearing of wild type animals or

Arc1 mutation in GNO animals independently induced no defects or only moderate

defects in host metabolism-related traits. However, when both the microbiota and *Arc1*

were lost, strong IIS-dependent metabolic phenotypes emerged. In animals lacking *Arc1*,

association with one specific *Acetobacter sp.* alone was sufficient to restore normal IIS

signaling and developmental rate. Our data suggest that *Acetobacter sp.* functions in this

capacity through multiple mechanisms, one of which involves dietary modification. Thus,

in GF *Arc1* mutants, we propose that host genotype and microbial condition converge on

the IIS pathway, leading to growth defects.

<u>Role of Arc1 in insulin signaling in GF flies</u>

We found that GF *Drosophila* lacking Arc1 exhibit phenotypes consistent with

reduced IIS activity. Connections between Arc1/Arc and IIS/insulin activity have been

considered previously. Arc expression can be strongly induced in cultured human

neuroblasts by treatment with exogenous insulin (Kremerskothen et al., 2002). Arc1 is

strongly expressed in one small cluster of large neurons in each lobe of both the larval

brain and the pars cerebralis region of the adult brain that are proximal to and partially

overlap with the insulin producing cells (IPCs; Mattaliano et al., 2007; Mosher et al., 2015).

Further, Mattaliano et al. (2007) found that selectively restoring Arc1 expression in the

IPCs was sufficient to revert the starvation resistance of *Arc1* mutants. Yet beyond

starvation resistance and the increased larval fat levels reported by Mosher et al. (2015),

loss of Arc1 resulted in no additional traits suggestive of reduced IIS signaling (Mattaliano

et al., 2007). However, our data now reveal a microbiota-dependent component of the

Arc1-IIS connection, as we primarily observed IIS-related phenotypes in *Arc1* mutants

grown under GF conditions. Further, Arc1 metabolic phenotypes are strongly influenced

by nutritional composition, as we found that the developmental rate of GF *Arc1^{E8}* animals

varied considerably on various dietary formulations.

The IIS pathway coordinates nutritional intake, metabolism, and growth in a

mechanistically conserved manner between *Drosophila* and vertebrates. The *Drosophila*

IIS pathway is functionally analogous to the combined roles of the mammalian insulin

pathway, which primarily regulates blood glucose levels, and the Insulin-like growth factor

(IGF) pathway, which coordinates organismal growth (Edgar, 2006). Our data suggest

that Arc1 is a regulator of IIS signaling, whose role is revealed by GF rearing. In addition,

Arc1 likely impacts metabolism- and growth-regulating cellular processes beyond IIS,

which may or may not contribute to the growth delay of GF *Arc1^{E8}* larvae. This is

evidenced by the increased wing size and enhanced starvation resistance of *Arc1*

mutants even in the presence of *Acetobacter sp.* (Figure 5D,E). Possible additional

pathways influenced by Arc1 include Hippo signaling, which controls organ growth (Zhao

et al., 2011), and ecdysone biosynthesis, which is nutrient responsive and regulates

developmental rate (Edgar, 2006).

How might Arc1 promote IIS-pathway activity? Mammalian Arc modulates synaptic

plasticity in part by enhancing the endocytosis, and consequently reducing the membrane

availability, of AMPA receptors (Chowdhury et al., 2006; DaSilva et al., 2016; Wall and

Corrêa, 2018). Although it has not been shown for Arc1, this molecular function suggests

that Arc proteins may have the ability to regulate the recycling of other receptors, like the

insulin receptor. An alternative mechanism is suggested by recent reports that *Drosophila*

Arc1 and mammalian Arc encode polypeptides that can self-assemble into mRNA-

containing capsid-like structures, which are released from mammalian and *Drosophila*

cell lines, and *Drosophila* neurons at the larval NMJ (neuromuscular junction) in

extracellular vesicles (EVs; Ashley et al., 2018; Erlendsson et al., 2019; Pastuzyn et al.,

2018). These Arc/Arc1 capsid-containing EVs are taken up by recipient cells in both

cultured mouse hippocampal neurons and at the NMJ, constituting a novel mechanism of

intercellular mRNA transfer (Ashley et al., 2018; Pastuzyn et al., 2018). The functional

significance of this mechanism in cell types and tissues outside the NMJ that may be

relevant here (e.g. IPC-adjacent neurons) has not been studied, but it is tempting to

speculate that Arc1-mediated mRNA transfer may play a role in Arc1's metabolic function.

Interestingly, Arc1 is broadly expressed; in addition to its expression in the brain

(Mattaliano et al., 2007; Mosher et al., 2015), Arc1 is also expressed in the prothoracic

gland (Mattaliano et al., 2007) and is transcriptionally enriched in the gut (FlyAtlas; Leader

et al., 2018), where we and others have observed microbiota-dependent transcript level

changes (Figure 1A; Bost et al., 2017; Dobson et al., 2016; Guo et al., 2014; Petkau et

al., 2017). Additionally, knockdown of the transcription factor Seven-up, which reduces

IIS activity, leads to strongly increased *Arc1* expression in the larval fat body (Musselman

et al., 2018). This implicates Arc1 in multiple organs that control energy homeostasis. We

predict that Arc1 capsid-dependent intercellular mRNA transport and/or endocytic based

receptor cycling might act at multiple nodes in the systemic inter-organ signaling

dynamics of IIS.

Acetobacter sp. support of *Arc1* mutant larval growth

While loss of *Arc1* significantly exacerbated the developmental delay of GF larvae,

we found that monoassociation with *Acetobacter sp.* was sufficient for *Arc1* mutants to

develop at the same rate as wild-type *Acetobacter sp.*-associated larvae (Figure 2B). The

reduced growth rate of GF *Drosophila* is one of the most consistent and reproducible

microbiota-dependent host traits documented in the literature. As such, the mechanistic

basis underlying microbial growth promotion has been investigated (reviewed in Strigini

and Leulier, 2016). Our study revealed both similarities and differences between

published mechanisms of bacterial impact on larval growth, and the ways *Acetobacter*

sp. promotes development in an *Arc1*-deficient host.

In laboratory *Drosophila* cultures, bacterial populations predominantly grow on the

diet substrate, and are ingested along with the food (Ludington and Ja, 2020). We found

that pre-conditioning the larval diet with *Acetobacter sp.* substantially accelerated GF

Arc1 mutant development, suggesting that an interaction between *Acetobacter sp.* and

the larval diet is a key feature of this bacterial isolate's growth promoting activity (Figure

4D'). There is precedent for bacterial modification of the diet altering host metabolism and

growth in *Drosophila*. On nutritionally rich diets, *Acetobacter tropicalis* can prevent

excessive accumulation of triglycerides in adult flies by metabolizing glucose in the food

and reducing its availability to the host (Huang and Douglas, 2015). On nutritionally poor

diets, *L. plantarum* depletes the levels of sugars and branched-chain amino acids, and

increase the levels of glycolysis and fermentation products to promote larval growth

(Storelli et al., 2017). Because acetic acid bacteria frequently predominate the *Drosophila*

microbiota, the impacts of acetic acid on fly physiology and development have been

investigated (Kamareddine et al., 2018; Kim et al., 2017; Shin et al., 2011). As in Kim et

al. (2017), we found that acetic acid either increased or had no effect on the

developmental delay of wild-type or Arc1 mutant larvae (Figure 4C,C'), suggesting acetic

acid production by our *Acetobacter sp.* isolate likely has a minimal impact on *Arc1* mutant

development under our dietary conditions.

Importantly, our data reveal that association with a live population of *Acetobacter*

sp. is required for optimal growth rate promotion of *Arc1* mutants. This suggests that

additional mechanisms beyond dietary modification support larval development in the

absence of *Arc1*. In wild-type flies, commensal bacteria have been shown to promote

growth through mechanisms involving interactions between bacterial cell wall

components and gut cells. For example, D-alanylated teichoic acids in the *L. plantarum*

cell wall induce expression of intestinal peptidases, facilitating larval growth on protein-

poor diets (Consuegra et al., 2020; Matos et al., 2017). Further, the microbiota-responsive

immune deficiency (IMD) pathway, which is activated in enterocytes by DAP-type

peptidoglycan present in gram-negative and *Lactobacilli* cell walls, has been shown to

promote metabolic homeostasis and larval development (Davoodi et al., 2018;

Kamareddine et al., 2018).

Taken together, our data suggest that the full impact of live *Acetobacter sp.* on

Arc1 mutant metabolism and development involves a combinatorial effect of dietary

modification and direct bacterial-host cell interactions. These two distinct modes of

microbiota activity are analogous to the modes by which the microbiota influence host

physiology in mammals. Bacterial breakdown of certain macro-nutrients (e.g. complex

polysaccharides) in the human gut has been linked to health and disease states

(Cockburn and Koropatkin, 2016). In contrast, other functions, such as immune cell

maturation and maintenance of gut epithelial architecture, appear to require bacterial cell-

derived antigens (Sekirov et al., 2010).

<u>Proposed evolutionary conservation of Arc-microbiota-insulin connections</u>

Our data raise the question of whether microbiota- and diet-responsive metabolism

regulation is a conserved function for Arc proteins. It has been shown that chronically

feeding mice a high fat diet (HFD), which induces insulin resistance and diabetic-like

phenotypes as well as cognitive impairments, leads to suppressed Arc expression in the

cerebral cortex and hippocampus (Chen et al., 2020; Mateos et al., 2009). Comparably,

flies reared on a HFD exhibited reduced memory formation and reduced Arc1 expression

in the head (p<0.05, 1.35 fold-change reduction; Rivera et al., 2019). In rodent models, it

is well established that HFD perturbs the composition of the gut microbiota in a manner

that can mediate diabetic and other disease phenotypes (Kim et al., 2012; Murphy et al.,

2015). Our data thus raise the intriguing and previously unconsidered possibility that Arc

function in the brain plays a role linking diet-induced gut dysbiosis to cognitive and

metabolic impairments.

MATERIALS AND METHODS

***Drosophila* stocks and general rearing**

The following fly stocks were used in this study: w^{1118}, Canton-S, and Oregon-R are long-

term lab stocks originally from the Bloomington Drosophila Stock Center (BDSC), Top

Banana (kind gift from Dr. Michael Dickinson), *y[1] w[1]* (BDSC #1495), *w^{1118}; Arc1^{E8}* (kind

gift from Dr. Vivian Budnik and Dr. Travis Thomson), *w[*]; Arc1^{esm113}* (BDSC #37531),

w[*]; Arc1^esm18 (BDSC #37530), *w[*];; P{w[+mC]=Arc1-GAL4.M}3* (BDSC #37533), *w^1118;*

P{UAS-rpr.C}14   (BDSC   #5824),   *y[1]w[67c23];*

P{w[+mC]y[+mDint2]=EPgy2}Arc2[EY21260] (BDSC #22466), *w^118;; P{tGPH}4* (BDSC

#8164). Non-experimental fly stocks were maintained at 21-22°C. Our laboratory utilizes

a yeast-cornmeal-molasses diet of the following recipe (percentages are given as wt/vol

or vol/vol throughout Methods): 8.5% molasses (Domino Foods), 7% cornmeal (Prairie

Mills Products), 1.1% active dry yeast (Genesee Scientific), 0.86% gelidium agar

(MoorAgar). The diet is boiled for ~30-45 minutes, cooled to 60-65°C, supplemented with

0.27% propionic acid (Sigma) and 0.27% methylparaben (Sigma) and dispensed to

polypropylene vials. All experiments in this study utilized this diet formulation except those

presented in Figure 2, which were conducted on yeast-glucose diets. Yeast-glucose diets

consisted of the indicated proportions of active dry yeast (Genesee Scientific), dextrose

(Fisher Scientific), and gelidium agar (MoorAgar), and were prepared as described in

(Koyle et al., 2016). Diets were mixed, autoclaved, cooled to 60-65°C, supplemented with

propionic acid and methylparaben, and dispensed to autoclaved vials.

Bacterial stocks

The *Acetobacter sp.*, *Acetobacter pasteurianus*, *Lactobacillus plantarum*, and

Lactobacillus brevis stocks utilized in this study were all isolated from conventionally

reared Top Banana *Drosophila* cultures in our laboratory. Adult flies were surface

sterilized in 10% sodium hypochlorite and 70% ethanol, rinsed three times and

homogenized in phosphate buffered saline (PBS). Serial dilutions of fly homogenates

were plated on de Man, Rogosa, and Sharpe (MRS; Weber Scientific) and acetic acid-

ethanol (AE; 0.8% yeast extract, 1.5% peptone, 1% dextrose, 0.5% ethanol, 0.3% acetic

acid; Blum et al., 2013) agar plates. Colonies with distinct morphology were streaked for

isolation. Bacterial taxonomies were assigned by PCR amplification and sequencing of

the 16S rRNA gene using universal bacterial primers 8F (5'-

AGAGTTTGATCTGGCTCAG-3') and 1492R (5'-GGMTACCTTGTTACGACTT-3'; Eden

et al., 1991). Sequences were searched against the NCBI nr/nt database via blastn

(Altschul et al., 1990; Camacho et al., 2009; Morgulis et al., 2008) and taxonomies were

assigned based on >97% sequence homology. The 16S rRNA sequence for isolate A22

(*Acetobacter sp.*) bore >97% similarity with >five different *Acetobacter* species so we do

not assign a species-level taxonomic classification in this report.

Generation of germ free and gnotobiotic fly cultures

Germ free and gnotobiotic *Drosophila* cultures were generated according to established

methods (Koyle et al., 2016). Synchronous populations of embryos were collected on

apple juice agar plates. In a sterile biosafety cabinet, embryos were treated with 50%

sodium hypochlorite solution for two minutes to eliminate exogenous microbes and

remove the chorion. Embryos were then rinsed twice in 70% ethanol, twice in sterilized

milliQ water, and once in sterilized embryo wash (2% Triton X-100, 7% NaCl). Sterilized

embryos were then pipetted into autoclaved food vials to generate germ free cultures. To

generate gnotobiotic flies, overnight cultures of bacterial isolates were grown in MRS

broth (30°C with shaking for *Acetobacter* isolates and 37°C static for *Lactobacilli*). Sterile

food vials were inoculated with 40μL of overnight cultures (OD~1) immediately prior to

the addition of sterilized fly embryos. For polyassociated (GNO) flies, vials were

inoculated with 40μL of a 1:1:1:1 mixture of overnight cultures of the four indicated

bacteria.

All experimental *Drosophila* cultures were maintained in an insect incubator at 23°C, 70%

humidity, on a 12:12 light-dark cycle.

Larval and adult animals were confirmed as germ free or mono-/polyassociated by

homogenization in sterile PBS and plating on MRS and AE agar plates.

We did not maintain GF, GNO, or monoassociated flies over multiple generations; all

experiments utilized independently-derived germ free or gnotobiotic animals.

Developmental timing measurements

Synchronous populations of embryos were collected in a six-hour time window and

treated as described above to generate vials of defined microbial conditions. For

pupariation and eclosion rate analysis, the number of pupae formed or empty pupal

cases, respectively, were counted daily until 100% of the population had pupariated or

eclosed. The duration of larval development is strongly affected by crowding conditions

in the food (Klepsatel et al., 2018). Also, variable and unpredictable numbers of embryos

do not survive the bleach and ethanol washes employed to generate germ free and

gnotobiotic cultures (Koyle et al., 2016; Troha and Buchon, 2019; unpublished

observations). Therefore, vials containing fewer than ten and greater than forty animals

were omitted from analyses as either under- or over-crowded, respectively.

Larval instars were determined via mouth hook and/or posterior spiracle morphologies

(Oldroyd, 1951).

Larval length and pupal volume were measured from images using Fiji (Schindelin et al.,

2012). For pupal volume, length (l) and width (w) of each pupa were measured and

volume calculated as previously described (Layalle et al., 2008; Redhai et al., 2020):

$V=4/3\pi(l/2)(w/2)^2$.

 Larval feeding assays

Larval feeding was assessed via dye consumption (Buhler et al., 2018; Libert et al., 2007;

Mosher et al., 2015; Shin et al., 2011). Stage-matched pre-wandering third instar larvae

were transferred to autoclaved fly food containing 1.8% FD&C Red #40 dye (Ward's

Science,) and allowed to feed for 3 hr at 23°C. Guts were then dissected from 20 larvae

in PBS and homogenized via bead beating in 1mL PBS. Homogenates were centrifuged

at 13,000 rpm for 1 minute to pellet debris and absorbance of the supernatant was

measured at 490nm with a Tecan Spark microplate reader. Weighed amounts of dyed

food were similarly homogenized in PBS and A_{490} readings collected to generate a

standard curve, which was used to calculate μg food consumed per larva.

Mouth hook contraction rates were assayed as described (Bhatt and Neckameyer, 2013).

Pre-wandering third instar larvae were transferred to apple juice plates coated with yeast

paste, given 30 seconds to acclimate, and contractions were counted manually for 30

seconds.

Quantifying bacterial loads for monoassociated larvae

Pre-wandering third instar larvae (8-10 animals per replicate) were removed from the food

and surface sterilized in 10% sodium hypochlorite for 1 minute. Larvae were then rinsed

three times in PBS and homogenized in 125μL PBS via bead beating for 30 seconds.

Homogenates were serially diluted in PBS and dilutions were plated on AE (for

Acetobacter) or MRS (for *Lactobacilli*) agar plates. Plates were incubated for 2-3 days at

either 30°C (AE) or 37°C (MRS), and resultant colonies were counted manually from

dilution plates bearing ~50-400 colonies. Bacterial loads were calculated as colony-

forming units (CFUs) per larva, as previously described (Koyle et al., 2016), and log

transformed for analysis.

Dietary treatments

Acetobacter sp.-conditioned diet

To generate *Acetobacter sp.*-conditioned fly food (Figure 4), autoclaved food vials were

inoculated with *Acetobacter sp.* overnight culture, as described above. Inoculated vials

were incubated for five days at 23°C. Vials were then incubated at 65°C for 3 hr to kill

bacteria. Sterility of the conditioned diet was confirmed by plating food and larval

homogenates on AE plates, which consistently yielded no bacterial growth. Heated diet

controls consisted of un-inoculated, autoclaved GF vials incubated at 65°C for 3 hr.

Heat-killed bacterial feeding

For experiments feeding GF larvae dead bacterial cells (Figure 4A), overnight cultures of

Acetobacter sp. were heat-killed at 65°C for 1 hr and autoclaved food vials were

inoculated with 40μL of heat-killed suspension prior to the addition of GF embryos. Vials

were further inoculated with 40μL heat-killed bacterial suspension daily until 100% of the

population had pupariated. Successful heat killing was confirmed for each daily inoculum

by plating undiluted heat-treated bacterial suspension on AE plates, and by plating larval

homogenates, which consistently yielded no bacterial growth.

Cell-free supernatant feeding

For *Acetobacter sp.* supernatant treatments (Figure 4B), overnight cultures of

Acetobacter sp. (3mL in MRS, grown as described above) were filtered twice through

0.22μm PVDF sterile membrane filters (Genesee Scientific). Autoclaved food vials were

inoculated with 40μL of filtered media immediately prior to addition of GF embryos, and

vials were further inoculated with 40μL of filtered media daily until 100% of the population

pupariated. Absence of live bacterial cells was confirmed by plating daily filtered media

on AE plates, and plating larval homogenates.

<u>Acetic acid-supplemented diets</u>

For acetic acid supplementation of GF larval fly cultures (Figure 4C, C'), fly food was

autoclaved and allowed to cool to ~50-60°C. Glacial acetic acid (Fisher Scientific) was

then added to defined volumes of food to the indicated final concentrations (vol/vol 0.1%

and 0.2%). For sodium acetate supplementation, solid sodium acetate (Sigma) was

added to autoclaved, cooled food to a final concentration of 50mM. Supplemented diets

were mixed thoroughly and dispensed to pre-autoclaved empty vials.

Wing analysis

Wings from adult females were dissected, mounted in Aqua-mount (Thermo Scientific),

and imaged with a QICAM-IR Fast 1394 camera (Q-Imaging) on a Zeiss Axioskop2 Plus

microscope. All measurements were collected using Fiji. Wing area was measured along

wing margin, excluding the hinge. For cell size analysis, trichome density was measured

with FijiWings2.3 software using the 150px trichome density feature (Dobens and

Dobens, 2013). Cell count per wing was determined by multiplying the cell density

measurement by the total area of the wing.

Starvation resistance

Five days post-eclosion, adult mated female flies were transferred to 1% agar-water vials,

8-10 flies per vial. Flies were transferred to fresh agar-water vials daily. Survival was

monitored daily until 100% of the population succumbed. The experiment was conducted

three times, ~50-120 animals per condition per replicate, and data were combined for

analysis.

RT-qPCR

Heads and guts (proventriculus to hindgut, excluding crop and malpighian tubules) from

5-7 days-post-eclosion adult male flies (8-10 animals per replicate) were dissected in ice

cold PBS and homogenized immediately in Trizol reagent (Thermo Fisher). RNA was

extracted using the Direct-zol RNA Miniprep kit (Zymo Research) exactly following the

manufacturer's protocol. High quality RNA ($A_{260nm/280nm}$ ~2; 500ng) was used as template

for cDNA synthesis using the qScript cDNA synthesis kit (QuantaBio). Product from cDNA

synthesis reactions was used for qPCR with the PerfeCTa SYBR Green Supermix

(QuantaBio) in an Applied Biosystems 7300 Real Time PCR System instrument. Data

were normalized to *Rpl32* and expression fold changes were calculated using the $2^{-\Delta\Delta Ct}$

method. Primers used in this study: *Rpl32* (5'- ATGCTAAGCTGTCGCACAAATG-3' and

5'- GTTCGATCCGTAACCGATGT-3'; Ponton et al., 2011), *Arc1* (5'-

CATCATCGAGCACAACAACC-3' and 5'-CTACTCCTCGTGCTGCTCCT-3'; Mosher et

al., 2015), *Arc2* (5'-CGTGGAGACGTATAAAGAGGTGG-3' and 5'-

GACCAGGTCTTGGCATCCC-3'; FlyPrimerBank (Hu et al., 2013)).

Fluorescence imaging

Fat bodies from pre-wandering third instar larvae were dissected in ice cold PBS and

fixed in 4% paraformaldehyde-PBS (Electron Microscopy Sciences) for 30 min at room

temperature. Fat bodies were rinsed three times in PBS and mounted in aqua-poly/mount

(Polysciences). Images were captured on a spinning disc microscope with a Celesta 1W

light engine (Lumencor), an X-Light V2 scan head (Crest Optics), and a Prime95B CMOS

camera (Photometrics) on a Zeiss Axiovert 200M using Metamorph software (Molecular

Devices).

Statistical analysis

Statistical tests were conducted and figures generated using R version 3.5.1 (R Core

Team, 2019). For development data, the average time to pupariation was calculated for

each vial from the number of individuals pupariating on each day until the entire

population completed larval development. These per-vial values from at minimum three

replicates were used for statistical analyses; full sample sizes and statistical test output

for all development experiments in this study are reported in Table S1. Throughout,

within-genotype comparisons among different treatments were conducted via one-way

analysis-of-variance (ANOVA), while comparisons among different genotypes and

treatments were conducted via two-way ANOVA, as indicated in the figure legends. Post-

hoc analysis among significantly different factors were conducted via Tukey test using the

"lsmeans" package (Lenth, 2016). RT-qPCR data were analyzed via Student's t-test.

Starvation survival data were compared via Cox proportional-hazards model analysis

using the "survival" package (Therneau, 2012). The threshold of statistical significance

was considered $p<0.05$.

ACKNOWLEDGMENTS

We thank members of the McCartney, Hiller, and Mitchell labs for helpful discussions

during the performance of the study. We thank Rory Eutsey (Hiller lab, Carnegie Mellon

University) for technical assistance with qPCR experiments. We would like to thank Dr.

John Woolford for providing feedback on the manuscript. The Top Banana fly stock was

a generous gift from Dr. Michael Dickinson's lab (CalTech). The *Arc1^{E8}* fly stock was a

generous gift from Dr. Vivian Budnik and Dr. Travis Thomson (University of

Massachusetts Medical School). We thank Bloomington Drosophila Stock Center for

providing other fly stocks. We would like to thank the Woolford, Mitchell, and Hinman labs

and the Molecular Biosensor and Imaging Center at Carnegie Mellon University for

reagents and equipment.

Funding for this work was provided by a Charles E. Kaufman Foundation New Initiative

Grant and a Carnegie Mellon University ProSEED/BrainHub seed grant to B.M.M.

S.A.K. was supported by NSF Graduate Research Fellowship DGE 1252522 and DGE

1745016.

No competing interests declared.

REFERENCES

Alhowikan, A. M. (2016). Activity-Regulated Cytoskeleton-Associated Protein
Dysfunction May Contribute to Memory Disorder and Earlier Detection of Autism
Spectrum Disorders. *Med. Princ. Pract.* **25**, 350-354.
Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J. (1990). Basic
local alignment search tool. *J. Mol. Biol.* **215**, 403–10.
Ashley, J., Cordy, B., Lucia, D., Fradkin, L. G., Budnik, V. and Thomson, T. (2018).
Retrovirus-like Gag Protein Arc1 Binds RNA and Traffics across Synaptic Boutons.
Cell **172**, 262-274.e11.
Baker, K. D. and Thummel, C. S. (2007). Diabetic Larvae and Obese Flies-Emerging
Studies of Metabolism in Drosophila. *Cell Metab.* **6**, 257–266.
Bhatt, P. K. and Neckameyer, W. S. (2013). Functional analysis of the larval feeding
circuit in Drosophila. *J. Vis. Exp.* **81**, e51062.
**Bi, R., Kong, L. L., Xu, M., Li, G. D., Zhang, D. F., Li, T., Fang, Y., Zhang, C., Zhang,
B. and Yao, Y. G.** (2018). The Arc Gene Confers Genetic Susceptibility to
Alzheimer's Disease in Han Chinese. *Mol. Neurobiol.* **55** 1217-1226.
Bing, X. L., Gerlach, J., Loeb, G. and Buchon, N. (2018). Nutrient-dependent impact
of microbes on Drosophila suzukii development. *MBio* **9**, e02199-17.
Blum, J. E., Fischer, C. N., Miles, J. and Handelsman, J. (2013). Frequent
Replenishment Sustains the Beneficial Microbiome of Drosophila melanogaster.
MBio **4**, e00860-13.
**Bost, A., Franzenburg, S., Adair, K. L., Martinson, V. G., Loeb, G. and Douglas, A.
E.** (2017). How gut transcriptional function of *Drosophila melanogaster* varies with

the presence and composition of the gut microbiota. *Mol. Ecol.* **27**, 1848-1859.

Britton, J. S., Lockwood, W. K., Li, L., Cohen, S. M. and Edgar, B. A. (2002).
Drosophila ' s Insulin / PI3-Kinase Pathway Coordinates Cellular Metabolism with
Nutritional Conditions. *Dev Cell* **2**, 239–249.

Broderick, N. A. and Lemaitre, B. (2012). Gut-associated microbes of *Drosophila*
melanogaster. *Gut Microbes* **3**, 307–321.

Brogiolo, W., Stocker, H., Ikeya, T., Rintelen, F., Fernandez, R. and Hafen, E.
(2001). An evolutionarily conserved function of the Drosophila insulin receptor and
insulin-like peptides in growth control. *Curr Biol* **11**, 213–221.

Broughton, S. J., Piper, M. D. W., Ikeya, T., Bass, T. M., Jacobson, J., Driege, Y.,
Martinez, P., Hafen, E., Withers, D. J., Leevers, S. J., et al. (2005). Longer
lifespan , altered metabolism , and stress resistance in Drosophila from ablation of
cells making insulin-like ligands. *Proc Natl Acad Sci U S A* **102**, 3105–3110.

Buhler, K., Clements, J., Winant, M., Bolckmans, L., Vulsteke, V. and Callaerts, P.
(2018). Growth control through regulation of insulin signalling by nutrition-activated
steroid hormone in drosophila. *Dev.* **145**, dev165654.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K. and
Madden, T. L. (2009). BLAST+: Architecture and applications. *BMC Bioinformatics*
10, 421.

Campillos, M., Doerks, T., Shah, P. K. and Bork, P. (2006). Computational
characterization of multiple Gag-like human proteins. *Trends Genet.* **22**, 585-589.

Carmichael, R. E. and Henley, J. M. (2018). Transcriptional and post-translational
regulation of Arc in synaptic plasticity. *Semin. Cell Dev. Biol.* **77**, 3-9.

Chaston, J., Newell, P. and Douglas, A. (2014). Metagenome-wide association of
microbial determinants of host phenotype in Drosophila melanogaster. *MBio* **5**,
e01631-14.

Chen, T. J., Chen, S. S., Wang, D. C. and Hung, H. S. (2020). High-fat diet reduces
novelty-induced expression of activity-regulated cytoskeleton-associated protein. *J.*
Cell. Physiol. **235**, 1065–1075.

Chowdhury, S., Shepherd, J. D., Okuno, H., Lyford, G., Petralia, R. S., Plath, N.,
Kuhl, D., Huganir, R. L. and Worley, P. F. (2006). Arc/Arg3.1 Interacts with the
Endocytic Machinery to Regulate AMPA Receptor Trafficking. *Neuron*. **52**, 445-459.

Cockburn, D. W. and Koropatkin, N. M. (2016). Polysaccharide Degradation by the
Intestinal Microbiota and Its Influence on Human Health and Disease. *J. Mol. Biol.*
428, 3230–3252.

Colombani, J., Raisin, S., Pantalacci, S., Radimerski, T., Montagne, J. and
Léopold, P. (2003). A nutrient sensor mechanism controls Drosophila growth. *Cell*.
114, 739-749.

Consuegra, J., Grenier, T., Baa-Puyoulet, P., Rahioui, I., Akherraz, H., Gervais, H.,
Parisot, N., Da Silva, P., Charles, H., Calevro, F., et al. (2020). Drosophila-
associated bacteria differentially shape the nutritional requirements of their host
during juvenile growth. *PLoS Biol.* **18**, e3000681.

Cottee, M. A., Letham, S. C., Young, G. R., Stoye, J. P. and Taylor, I. A. (2019).
Structure of D. melanogaster ARC1 reveals a repurposed molecule with
characteristics of retroviral Gag. *Sci Adv* **6**, eaay6354.

DaSilva, L. L. P., Wall, M. J., de Almeida, L. P., Wauters, S. C., Januário, Y. C.,

Müller, J. and Corrêa, S. A. L. (2016). Activity-regulated cytoskeleton-associated
protein controls AMPAR endocytosis through a direct interaction with clathrin-
adaptor protein 2. *eNeuro*. **3**, ENEURO.0144-15.2016.
Davoodi, S., Galenza, A., Panteluk, A., Deshpande, R., Ferguson, M., Grewal, S.
and Foley, E. (2018). The Immune Deficiency Pathway Regulates Metabolic
Homeostasis in Drosophila. *J Immunol.* **202**, 2747-2759.
DiAngelo, J. R. and Birnbaum, M. J. (2009). Regulation of Fat Cell Mass by Insulin in
Drosophila melanogaster. *Mol. Cell. Biol.* **29**, 6341-6352.
Dobens, A. C. and Dobens, L. L. (2013). Fijiwings: An open source toolkit for
semiautomated morphometric analysis of insect wings. *G3 Genes, Genomes,*
Genet. **3**, 1443-1449.
Dobson, A. J., Chaston, J. M. and Douglas, A. E. (2016). The Drosophila
transcriptional network is structured by microbiota. *BMC Genomics* **17**, 975.
Douglas, A. E. (2018). The Drosophila model for microbiome research. *Lab Anim. (NY).*
47, 157–164.
Eden, P. A. E., Schmidt, T. M., Blakemore, R. P. and Pace, N. R. (1991).
Phylogenetic analysis of *Aquaspirillum magnetotacticum* using polymerase chain
reaction-amplified 16S rRNA-specific DNA. *Int. J. Syst. Bacteriol.* **41**, 324–325.
Edgar, B. A. (2006). How flies get their size: genetics meets physiology. *Nat rev Genet.*
7, 907–916.
Ekmekciu, I., von Klitzing, E., Fiebiger, U., Escher, U., Neumann, C., Bacher, P.,
Scheffold, A., Kühl, A. A., Bereswill, S. and Heimesaat, M. M. (2017). Immune
responses to broad-spectrum antibiotic treatment and fecal microbiota
transplantation in mice. *Front. Immunol.* **8**, 397.
Erkosar, B., Storelli, G., Mitchell, M., Bozonnet, L., Bozonnet, N. and Leulier, F.
(2015). Pathogen Virulence Impedes Mutualist-Mediated Enhancement of Host
Juvenile Growth via Inhibition of Protein Digestion. *Cell Host Microbe* **18**, 445–455.
Erkosar, B., Kolly, S., van der Meer, J. R. and Kawecki, T. J. (2017). Adaptation to
Chronic Nutritional Stress Leads to Reduced Dependence on Microbiota in
Drosophila melanogaster. *MBio* **8**, e01496-17.
Erlendsson, S., Morado, D. R., Shepherd, J. D. and Briggs, J. A. G. (2019).
Structures of virus-like capsids formed by the drosophila neuronal Arc proteins. *Nat*
Neurosci. **23**, 172-175.
Fromer, M., Pocklington, A. J., Kavanagh, D. H., Williams, H. J., Dwyer, S.,
Gormley, P., Georgieva, L., Rees, E., Palta, P., Ruderfer, D. M., et al. (2014). De
novo mutations in schizophrenia implicate synaptic networks. *Nature.* **506**, 179-
184.
Géminard, C., Rulifson, E. J. and Léopold, P. (2009). Remote Control of Insulin
Secretion by Fat Cells in Drosophila. *Cell Metab.* **10**, 199–207.
Gilbert, L. I. (2008). Drosophila is an inclusive model for human diseases, growth and
development. *Mol. Cell. Endocrinol.* **293**, 25-31.
Gilbert, J. A., Krajmalnik-Brown, R., Porazinska, D. L., Weiss, S. J. and Knight, R.
(2013). Toward Effective Probiotics for Autism and Other Neurodevelopmental
Disorders. *Cell* **155**, 1446–1448.
Guan, Z., Saraswati, S., Adolfsen, B. and Littleton, J. T. (2005). Genome-wide
transcriptional changes associated with enhanced activity in the Drosophila

nervous system. *Neuron*. **48**, 91-107.

Guo, L., Karpac, J., Tran, S. L. and Jasper, H. (2014). PGRP-SC2 promotes gut
immune homeostasis to limit commensal dysb, iosis and extend lifespan. *Cell* **156**,
109–122.

Guzowski, J. F., McNaughton, B. L., Barnes, C. A. and Worley, P. F. (1999).
Environment-specific expression of the immediate-early gene Arc in hippocampal
neuronal ensembles. *Nat. Neurosci.* **2**, 1120-1124.

Guzowski, J. F., Lyford, G. L., Stevenson, G. D., Houston, F. P., McGaugh, J. L.,
Worley, P. F. and Barnes, C. A. (2000). Inhibition of activity-dependent arc protein
expression in the rat hippocampus impairs the maintenance of long-term
potentiation and the consolidation of long-term memory. *J. Neurosci.* **20**, 3993-
4001.

Hayes, C. L., Dong, J., Galipeau, H. J., Jury, J., McCarville, J., Huang, X., Wang, X.
Y., Naidoo, A., Anbazhagan, A. N., Libertucci, J., et al. (2018). Commensal
microbiota induces colonic barrier structure and functions that contribute to
homeostasis. *Sci. Rep.* **8**, 14184.

Hu, Y., Sopko, R., Foos, M., Kelley, C., Flockhart, I., Ammeux, N., Wang, X.,
Perkins, L., Perrimon, N. and Mohr, S. E. (2013). FlyPrimerBank: an online
database for Drosophila melanogaster gene expression analysis and knockdown
evaluation of RNAi reagents. *G3 (Bethesda).* **3**, 1607-1616.

Huang, J.-H. and Douglas, A. E. (2015). Consumption of dietary sugar by gut bacteria
determines *Drosophila* lipid content. *Biol. Lett.* **11**, 20150469.

Jünger, M. A., Rintelen, F., Stocker, H., Wasserman, J. D., Végh, M., Radimerski,
T., Greenberg, M. E. and Hafen, E. (2003). The Drosophila Forkhead transcription
factor FOXO mediates the reduction in cell number associated with reduced insulin
signaling. *J. Biol.* **2**, 20.

Kamareddine, L., Robins, W. P., Berkey, C. D., Mekalanos, J. J. and Watnick, P. I.
(2018). The Drosophila Immune Deficiency Pathway Modulates Enteroendocrine
Function and Host Metabolism. *Cell Metab.* **28**, 449-462.e5.

Keebaugh, E. S., Yamada, R., Obadia, B., Ludington, W. B. and Ja, W. W. (2018).
Microbial Quantity Impacts Drosophila Nutrition, Development, and Lifespan.
iScience. **4**, 247-259.

Keith, S. A., Eutsey, R., Lee, H., Solomon, B., Oliver, S., Kingsford, C., Hiller, N. L.
and McCartney, B. M. (2019). Identification of Microbiota-Induced Gene
Expression Changes in the Drosophila melanogaster Head. *bioRxiv*.

Kim, K. A., Gu, W., Lee, I. A., Joh, E. H. and Kim, D. H. (2012). High Fat Diet-Induced
Gut Microbiota Exacerbates Inflammation and Obesity in Mice via the TLR4
Signaling Pathway. *PLoS One*. **7**, e47713.

Kim, G., Huang, J. H., McMullen, J. G., Newell, P. D. and Douglas, A. E. (2017).
Physiological responses of insects to microbial fermentation products: insights from
the interactions between Drosophila and acetic acid. *J. Insect Physiol.* **106**, 13-19.

Klepsatel, P., Procházka, E. and Gáliková, M. (2018). Crowding of Drosophila larvae
affects lifespan and other life-history traits via reduced availability of dietary yeast.
Exp. Gerontol. **110**, 298-308.

Koyle, M. L., Veloz, M., Judd, A. M., Wong, A. C.-N., Newell, P. D., Douglas, A. E.
and Chaston, J. M. (2016). Rearing the Fruit Fly Drosophila melanogaster Under

Axenic and Gnotobiotic Conditions. *J. Vis. Exp.* **113**, 54219.

Kramer, J. M., Davidge, J. T., Lockyer, J. M. and Staveley, B. E. (2003). Expression
of Drosophila FOXO regulates growth and can phenocopy starvation. *BMC Dev.*
Biol. **3**, 5.

Kremerskothen, J., Wendholt, D., Teber, I. and Barnekow, A. (2002). Insulin-induced
expression of the activity-regulated cytoskeleton-associated gene (ARC) in human
neuroblastoma cells requires p21ras, mitogen-activated protein kinase/extracellular
regulated kinase and src tyrosine kinases but is protein kinase C-independ.
Neurosci. Lett. **321**, 153–156.

Larsen, N., Vogensen, F. K., Van Den Berg, F. W. J., Nielsen, D. S., Andreasen, A.
S., Pedersen, B. K., Al-Soud, W. A., Sørensen, S. J., Hansen, L. H. and
Jakobsen, M. (2010). Gut microbiota in human adults with type 2 diabetes differs
from non-diabetic adults. *PLoS One* **5**, e9085.

Layalle, S., Arquier, N. and Léopold, P. (2008). The TOR Pathway Couples Nutrition
and Developmental Timing in Drosophila. *Dev. Cell* **15**, 568–577.

Leader, D. P., Krause, S. A., Pandit, A., Davies, S. A. and Dow, J. A. T. (2018).
FlyAtlas 2: A new version of the Drosophila melanogaster expression atlas with
RNA-Seq, miRNA-Seq and sex-specific data. *Nucleic Acids Res.* **46**, D809–D815.

Lenth, R. V. (2016). Least-squares means: The R package lsmeans. *J. Stat. Softw.*

Lesperance, D. N. A. and Broderick, N. A. (2020). Meta-analysis of Diets Used in
Drosophila Microbiome Research and Introduction of the Drosophila Dietary
Composition Calculator (DDCC). *G3; Genes|Genomes|Genetics* g3.401235.2020.

Libert, S., Zwiener, J., Chu, X., VanVoorhies, W., Roman, G. and Pletcher, S. D.
(2007). Regulation of Drosophila life span by olfaction and food-derived odors.
Science. **315**, 1133-1137.

Ludington, W. B. and Ja, W. W. (2020). Drosophila as a model for the gut microbiome.
PLoS Pathog. **16**, e1008398.

Luong, N., Davies, C. R., Wessells, R. J., Graham, S. M., King, M. T., Veech, R.,
Bodmer, R. and Oldham, S. M. (2006). Activated FOXO-mediated insulin
resistance is blocked by reduction of TOR activity. *Cell Metab.* **4**, 133-142.

Martino, M., Ma, D. and Leulier, F. (2017). Microbial influence on Drosophila biology.
Curr. Opin. Microbiol. **38**, 165–170.

Martino, M. E., Joncour, P., Leenay, R., Gervais, H., Shah, M., Hughes, S., Gillet,
B., Beisel, C. and Leulier, F. (2018). Bacterial Adaptation to the Host's Diet Is a
Key Evolutionary Force Shaping Drosophila-Lactobacillus Symbiosis. *Cell Host*
Microbe. **24**, 109–119.

Mateos, L., Akterin, S., Gil-Bea, F. J., Spulber, S., Rahman, A., Björkhem, I.,
Schultzberg, M., Flores-Morales, A. and Cedazo-Mínguez, A. (2009). Activity-
regulated cytoskeleton-associated protein in rodent brain is down-regulated by high
fat diet in vivo and by 27-hydroxycholesterol in vitro. *Brain Pathol.* **19**, 69-80.

Matos, R. C., Schwarzer, M., Gervais, H., Courtin, P., Joncour, P., Gillet, B., Ma, D.,
Bulteau, A.-L., Martino, M. E., Hughes, S., et al. (2017). D-Alanylation of teichoic
acids contributes to Lactobacillus plantarum-mediated Drosophila growth during
chronic undernutrition. *Nat. Microbiol.* **2**, 1635-1647.

Mattaliano, M. D., Montana, E. S., Parisky, K. M., Littleton, J. T. and Griffith, L. C.
(2007). The Drosophila ARC homolog regulates behavioral responses to starvation.

Mol. Cell. Neurosci. **36**, 211–221.

McBrayer, Z., Ono, H., Shimell, M. J., Parvy, J. P., Beckstead, R. B., Warren, J. T., Thummel, C. S., Dauphin-Villemant, C., Gilbert, L. I. and O'Connor, M. B. (2007). Prothoracicotropic Hormone Regulates Developmental Timing and Body Size in Drosophila. *Dev. Cell.* **13**, 857-871.

McFall-Ngai, M., Hadfield, M. G., Bosch, T. C. G., Carey, H. V, Domazet-Lošo, T., Douglas, A. E., Dubilier, N., Eberl, G., Fukami, T., Gilbert, S. F., et al. (2013). Animals in a bacterial world, a new imperative for the life sciences. *Proc. Natl. Acad. Sci.* **110**, 3229–3236.

Mirth, C., Truman, J. W. and Riddiford, L. M. (2005). The role of the prothoracic gland in determining critical weight for metamorphosis in Drosophila melanogaster. *Curr. Biol.* **15**, 1769-1807.

Montana, E. S. and Littleton, J. T. (2006). Expression profiling of a hypercontraction-induced myopathy in Drosophila suggests a compensatory cytoskeletal remodeling response. *J. Biol. Chem.* **10**, 421.

Morgulis, A., Coulouris, G., Raytselis, Y., Madden, T. L., Agarwala, R. and Schäffer, A. A. (2008). Database indexing for production MegaBLAST searches. *Bioinformatics*, **24**, 1757–1764.

Mosher, J., Zhang, W., Blumhagen, R. Z., D'Alessandro, A., Nemkov, T., Hansen, K. C., Hesselberth, J. R. and Reis, T. (2015). Coordination between Drosophila Arc1 and a specific population of brain neurons regulates organismal fat. *Dev. Biol.* **405**, 280–290.

Murphy, E. A., Velazquez, K. T. and Herbert, K. M. (2015). Influence of high-fat diet on gut microbiota: A driving force for chronic disease risk. *Curr. Opin. Clin. Nutr. Metab. Care*. **18**, 515-520.

Musselman, L. P., Fink, J. L., Narzinski, K., Ramachandran, P. V., Hathiramani, S. S., Cagan, R. L. and Baranski, T. J. (2011). A high-sugar diet produces obesity and insulin resistance in wild-type Drosophila. *Dis. Model. Mech.* **4**, 842-849.

Musselman, L. P., Fink, J. L., Maier, E. J., Gatto, J. A., Brent, M. R. and Baranski, T. J. (2018). Seven-up is a novel regulator of insulin signaling. *Genetics*. **208**, 1643-1656.

Newell, P. D. and Douglas, A. E. (2014). Interspecies Interactions Determine the Impact of the Gut Microbiota on Nutrient Allocation in Drosophila melanogaster. *Appl. Environ. Microbiol.* **80**, 788–796.

Oldroyd, H. (1951). Biology of Drosophila. *Nature*.

Pastuzyn, E. D., Day, C. E., Kearns, R. B., Kyrke-Smith, M., Taibi, A. V., McCormick, J., Yoder, N., Belnap, D. M., Erlendsson, S., Morado, D. R., et al. (2018). The Neuronal Gene Arc Encodes a Repurposed Retrotransposon Gag Protein that Mediates Intercellular RNA Transfer. *Cell*. **172**, 275-288.e18.

Petkau, K., Ferguson, M., Guntermann, S. and Foley, E. (2017). Constitutive Immune Activity Promotes Tumorigenesis in Drosophila Intestinal Progenitor Cells. *Cell Rep.* **20**, 1784–1793.

Ponton, F., Chapuis, M. P., Pernice, M., Sword, G. A. and Simpson, S. J. (2011). Evaluation of potential reference genes for reverse transcription-qPCR studies of physiological responses in Drosophila melanogaster. *J. Insect Physiol.* **57**, 840–850.

Redhai, S., Pilgrim, C., Gaspar, P., Giesen, L. van, Lopes, T., Riabinina, O.,
Grenier, T., Milona, A., Chanana, B., Swadling, J. B., et al. (2020). An intestinal
zinc sensor regulates food intake and developmental growth. *Nature*. **580**, 263-268.
Rivera, O., McHan, L., Konadu, B., Patel, S., Sint Jago, S. and Talbert, M. E. (2019).
A high-fat diet impacts memory and gene expression of the head in mated female
Drosophila melanogaster. *J. Comp. Physiol. B.* **189**, 179-198.
Robertson, F. W. (1963). The ecological genetics of growth in Drosophila: 6. The
genetic correlation between the duration of the larval period and body size in
relation to larval diet. *Genet. Res.* **4**, 74-92.
Robertson, R. C., Manges, A. R., Finlay, B. B. and Prendergast, A. J. (2019). The
Human Microbiome and Child Growth – First 1000 Days and Beyond. *Trends*
Microbiol. **27**, 131-147.
Rulifson, E. J., Kim, S. K. and Nusse, R. (2002). Ablation of insulin-producing neurons
in files: Growth and diabetic phenotypes. *Science.* **296**, 1118-1120.
Saichana, N., Matsushita, K., Adachi, O., Frébort, I. and Frebortova, J. (2015).
Acetic acid bacteria: A group of bacteria with versatile biotechnological
applications. *Biotechnol. Adv.* **33**, 1260-1271.
Sampson, T. R. and Mazmanian, S. K. (2015). Control of brain development, function,
and behavior by the microbiome. *Cell Host Microbe* **17**, 565–576.
Sannino, D. R., Dobson, A. J., Edwards, K., Angert, E. R. and Buchon, N. (2018).
The Drosophila melanogaster Gut Microbiota Provisions Thiamine to Its Host. *MBio*
9, e00155-18.
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: An open-
source platform for biological-image analysis. *Nat. Methods*. **9**, 676-682.
Sekirov, I., Russell, S. L., Antunes, L. C. M. and Finlay, B. B. (2010). Gut Microbiota
in Health and Disease. *Physiol. Rev.* **90**, 859–904.
Shandilya, M. C. V. and Gautam, A. (2020). Hippocampal Arc Induces Decay of
Object Recognition Memory in Male Mice. *Neuroscience*. **431**, 193-204.
Shen, J., Obin, M. S. and Zhao, L. (2013). The gut microbiota, obesity and insulin
resistance. *Mol. Aspects Med.* **34**, 39–58.
Shepherd, J. D. and Bear, M. F. (2011). New views of Arc, a master regulator of
synaptic plasticity. *Nat. Neurosci.* **14**, 279–284.
Shin, S. C., Kim, S.-H., You, H., Kim, B., Kim, A. C., Lee, K.-A., Yoon, J.-H., Ryu, J.-
H. and Lee, W.-J. (2011). Drosophila Microbiome Modulates Host Developmental
and Metabolic Homeostasis via Insulin Signaling. *Science.* **334**, 670–674.
Smith, K., McCoy, K. D. and Macpherson, A. J. (2007). Use of axenic animals in
studying the adaptation of mammals to their commensal intestinal microbiota.
Semin. Immunol. **19**, 59-69.
Storelli, G., Defaye, A., Erkosar, B., Hols, P., Royet, J. and Leulier, F. (2011).
Lactobacillus plantarum promotes drosophila systemic growth by modulating
hormonal signals through TOR-dependent nutrient sensing. *Cell Metab.* **14**, 403–
414.
Storelli, G., Strigini, M., Grenier, T., Bozonnet, L., Schwarzer, M., Daniel, C., Matos,
R. and Leulier, F. (2017). Drosophila Perpetuates Nutritional Mutualism by
Promoting the Fitness of Its Intestinal Symbiont Lactobacillus plantarum. *Cell*

Metab. **27**, 362-377.
Strigini, M. and Leulier, F. (2016). The role of the microbial environment in Drosophila
post-embryonic development. *Dev. Comp. Immunol.* **64**, 39–52.
Team, R. C. (2019). R: A Language and Environment for Statistical Computing. *Vienna,*
Austria.
Therneau, T. (2012). A Package for Survival Analysis in S. R package version. *Survival*
(Lond).
Troha, K. and Buchon, N. (2019). Methods for the study of innate immunity in
Drosophila melanogaster. *Wiley Interdiscip. Rev. Dev. Biol.* **8**, e344.
Verdu, J., Buratovicht, M. A., Wilder, E. L. and Birnbaum, M. J. (1999). Cell-
autonomous regulation of cell and organ growth in Drosophila by Akt/PKB. *Nat. Cell*
Biol. **1**, 500-506.
Wall, M. J. and Corrêa, S. A. L. (2018). The mechanistic link between Arc/Arg3.1
expression and AMPA receptor endocytosis. *Semin. Cell Dev. Biol.* **77**, 17-24.
Wong, A. C., Dobson, A. J. and Douglas, A. E. (2014). Gut microbiota dictates the
metabolic response of Drosophila to diet. *J Exp Biol* **217**, 1894–1901.
Zhao, B., Tumaneng, K. and Guan, K. L. (2011). The Hippo pathway in organ size
control, tissue regeneration and stem cell self-renewal. *Nat. Cell Biol.* **13**, 877-883.

FIGURES

Figure 1. *Arc1* exhibits tissue-specific microbiota-dependent expression changes,

and loss of *Arc1* exacerbates the developmental delay of GF larvae. A. RT-qPCR

analysis of *Arc1* transcripts in heads and guts of 5-day-old adult male wild-type flies.

Individual points represent normalized values for each replicate: n=3-4 per condition, 10

animals per replicate. Error bars represent standard error. * $p<0.05$, ** $p<0.01$, ***

$p<0.001$, n.s. = not significant Student's t-test. **B,C.** Developmental time courses of

GNO and GF wild-type/control *vs. Arc1* mutant animals, and **D.** larvae with *Arc1*-

expressing cells ablated. *Arc1>*: *Arc1-Gal4* crossed to w^{1118} background control,

Arc1>rpr: *Arc1-GAL4* crossed to *UAS-reaper*. Error bars represent standard error.

Conditions sharing letters are not statistically different from one another, two-way

ANOVA with Tukey's post-hoc test. For all developmental rate experiments, see Table

S1 for full sample sizes and statistical results.

Figure 2. Monoassociation with *Acetobacter sp.* promotes development of *Arc1*

mutant larvae. A, B. Time to pupariation for wild-type and *Arc1* mutant animals

monoassociated with each of the four bacterial isolates comprising the GNO condition.

GNO–3spp.: gnotobiotic community consisting of *A. pasteurianus*, *L. brevis*, and *L.*

plantarum. Conditions sharing letters are not statistically different from one another,

one-way ANOVA with Tukey's post-hoc test. **C.** Percentage of larvae in the indicated

instar stage daily until 100% of the population reaches the pre-wandering or wandering

third instar stage. Data represent pooled percentages from three biological replicates for

each day, ~30-60 animals per day. **D.** Larval length for both genotypes and microbial

conditions over time. Each data point represents an individual larva, three biological

replicates, ~10-20 larvae per replicate. Error bars represent standard error. **E.** Pupal

volume for each genotype and microbial condition. Each data point represents an

individual pupa. Conditions that share a letter are not statistically different from one

another, two-way ANOVA with Tukey's post-hoc test.

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157

Figure 3. Host diet impacts microbial effects on wild-type and *Arc1* mutant developmental rate. Time to pupariation for wild-type and *Arc1^E8 Acetobacter sp.*-associated and GF larvae reared on diets consisting of the indicated concentrations (weight/volume) of yeast and dextrose. Red dashed lines indicate the time to ~50% pupariation for *Arc1^E8 Acetobacter sp.* and *Arc1^E8* GF animals on the 3% glucose-5% yeast diet as an arbitrary reference to facilitate visual comparisons among diets. Values in blue represent the protein:carbohydrate ratio for each diet as calculated with the Drosophila Diet Composition Calculator (Lesperance and Broderick, 2020). Conditions that share a letter are not statistically different from one another, two-way ANOVA with Tukey's post-hoc test.

Figure 4. *Acetobacter sp.*-conditioned diet accelerates the development of *Arc1*

mutants. A. Daily administration of heat-killed *Acetobacter sp.* planktonic culture has

minimal effect on developmental rate of GF wild-type or *Arc1^{E8}* larvae. Two-way ANOVA

with Tukey's post-hoc test. **B.** Daily administration of filtered supernatant from

Acetobacter sp. planktonic culture has no effect on developmental rate of GF wild-type

or *Arc1^{E8}* larvae. Two-way ANOVA with Tukey's post-hoc test. **C, C'.** Rearing GF wild-

type and *Arc1* mutant larvae on diets containing acetic acid either further extends or has

no effect on the larval developmental delay. Each panel analyzed by one-way ANOVA

with Tukey's post-hoc test. **D, D'.** Rearing GF larvae on sterile diet that has been pre-

conditioned with *Acetobacter sp.* for five days (GF + conditioned diet) has no effect on

wild-type but partially restores developmental rate of *Arc1^{E8}* animals. GF + heated diet:

1204 GF larvae reared on GF diet heated under the same conditions used to kill *Acetobacter*

1205 *sp.* after pre-conditioning. Each panel analyzed by one-way ANOVA with Tukey's post-

1206 hoc test. Throughout, conditions that share a letter are not statistically different from one

1207 another.

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

Figure 5. GF *Arc1* mutants exhibit phenotypes consistent with insulin signaling

defects. A. Schematic representing impacts of insulin signaling manipulations and Arc1

mutation-microbial condition interactions on wing organ properties. **B-D.** Wing

morphological parameters for wild-type and *Arc1^E8* females under *Acetobacter sp.*-

associated and GF conditions. Each point represents an individual wing. Each panel

analyzed by two-way ANOVA with Tukey's post-hoc test. **E.** Survival curves for adult

female wild-type and *Arc1* mutant *Acetobacter sp.*-associated and GF animals

transferred to starvation conditions 5-7 days post-eclosion. Data represent pooled

results from three biological replicates, ~50-120 animals per condition per replicate.

Statistical results represent Cox proportional-hazards model analysis. Conditions that

share a letter are not statistically different from one another. **F.** Representative images

showing GFP-tagged plextrin homology domain (tGPH) localization in the fat bodies of

pre-wandering third instar larvae of the indicated genotypes and microbial conditions.

Scale bar=100μm. Arrowheads indicate membrane-localized fluorescent signal.