Title: Mixing coniferous and deciduous trees (*Fagus sylvatica*): Site-specific response of soil
microorganisms

Authors: Jing-Zhong Lu[1*], Stefan Scheu[1,2]

[1] J. F. Blumenbach Institute of Zoology and Anthropology, University of Göttingen, Untere Karspüle
2, 37073 Göttingen, Germany

[2] Center of Biodiversity and Sustainable Land Use, University of Göttingen, Büsgenweg 1, 37077
Göttingen, Germany

[*] Corresponding author (E-mail: jlu@gwdg.de)

Keywords: Douglas-fir, Norway spruce, soil respiration, forest, ecosystem functioning, PLFA

Author contributions: SS and JZL designed the study. JZL collected the data, analyzed the data
and wrote the first draft of the manuscript. SS contributed to all stages of the manuscript.

Abstract

Coniferous and deciduous trees growing in pure and mixed forests likely supply different resources to soil microorganisms thereby shaping microbial communities. However, effects of forest types including non-native species and their variation with site conditions on microbial communities remain poorly understood. We quantified functional (substrate-induced respiration and stress indicators) and structural (phospholipid-derived fatty acids) responses of microbial communities to forest types at four nutrient-rich and four nutrient-poor sites. Each site comprised five forest types, including pure stands of native European beech (*Fagus sylvatica*), range-expanding Norway spruce (*Picea abies*) and non-native Douglas-fir (*Pseudotsuga menziesii*), as well as mixed species stands of beech with spruce or with Douglas-fir, i.e., 40 forest stands. Forest type only affected soil microorganisms at nutrient-poor sites, but not at nutrient-rich sites. At nutrient-poor sites, both pure and mixed coniferous forests, in particular pure Douglas-fir forests, detrimentally affected soil microorganisms compared to beech forests. By contrast, structural and functional indicators of microorganisms in beech forests at nutrient-poor sites resembled those at nutrient-rich sites, likely because beech maintained the provisioning of high amounts of root-derived resources. The results indicate that long-term effects of planting exotic tree species such as Douglas-fir either in pure or in mixed stands on ecosystem functioning need further attention at nutrient-poor sites, but planting Douglas-fir at nutrient-rich sites may be of little concern from the perspective of microbial communities. Overall, the results highlight the potential importance of root-derived resources in determining microbial community structure and functioning, and document the sensitivity of soil microorganisms to planting non-native tree species that may differ from native species in the provisioning of these resources.

1. Introduction

Studying the linkage between plants and soil microorganisms has great potential to improve our understanding of the factors that drive microbial community composition (Balvanera et al., 2006; van der Heijden et al., 2008; Anderson and Domsch, 2010; Bardgett and van der Putten, 2014; Graham et al., 2016). Trees as long-lived plants may affect microbial communities through several pathways, most notably by the quantity and quality of litter and root exudates (Wardle et al., 2004; Högberg and Read, 2006). These resources are likely to shape the community structure of soil microorganisms as dissimilar carbon resources favor different guilds of microorganisms (Ekblad and Nordgren, 2002; Fierer et al., 2003; Fanin et al., 2019). For example, Gram$^-$ bacteria are benefiting in particular from labile carbon compounds, whereas Gram$^+$ bacteria are adapted to use more recalcitrant carbon resources (Fierer et al., 2003; Fanin et al., 2019; Waller et al., 2020). Similarly, niches vary between fungal groups. While saprotrophic fungi are major decomposers of plant litter, mycorrhizal fungi are more abundant in deeper soil and receive carbon from plant roots (Lindahl et al., 2007; Nacke et al., 2016). Therefore, the composition of microbial communities may change in response to variations in tree species composition, associated with changes in litter quality, root-derived resources and mycorrhizal associations (L. Chen et al., 2019).

The accumulation of organic matter in coniferous forests has been attributed to the recalcitrance of coniferous needles, but roots and mycorrhizal fungi also affect the storage of soil carbon (Schmidt et al., 2011; Clemmensen et al., 2013; Averill et al., 2014). Below-ground processes, such as root exudation and the competition between saprotrophic and mycorrhizal fungi, maybe more important for structuring microbial communities than the input of above-ground litter (Scheu et al., 2003; Pollierer et al., 2007; Schmidt et al., 2011; Augusto et al., 2015; Yan et al., 2016; Bluhm et al., 2019; Jones et al., 2019). Despite that coniferous and deciduous trees likely differ in the supply of litter and root-derived resources to soil microorganisms, the response of microbial communities to mixed forests is little studied. Litter mixture may improve resource availability to soil microorganisms and facilitate litter decomposition (Hättenschwiler et al., 2005; Fujii et al., 2020), although tree species identity has been suggested to outweigh tree diversity as structuring force of soil microbial communities (Scheibe et al., 2015; Dawud et al., 2016; L. Chen et al., 2019). Mixing of tree species also has been shown to increase both above-ground and below-ground biomass due to interspecific facilitation and improved resource partitioning, resulting in higher complementarity in space and time in mixed forests (Cardinale et al., 2007; Ma and Chen, 2016; Emmett Duffy et al., 2017). These interactions likely affect the way forests shape soil microbial communities. Overall, however, the role of identity and diversity of tree species, either coniferous or deciduous, in shaping the structure and functioning of soil microorganisms is little studied.

The relationship between tree species and microbial communities strongly depends on environmental context (Ratcliffe et al., 2017; X. Chen et al., 2019), but to what extent effects of tree species composition on soil microorganisms vary with site conditions, such as nutrient status and water availability, is little known. Facilitation among plants is more pronounced in unfavorable environments according to the stress-gradient hypothesis, and such positive interspecific interactions are likely to be mediated by soil microorganisms (Defossez et al., 2011; Harpole et al., 2016; David et al., 2020). In fact, plants do not passively tolerate environmental stress, but respond in various ways to unfavorable growth conditions, such as by allocating more resources to roots than to shoots under nutrient-poor conditions (Callaway et al., 2003; Kawaletz et al., 2013; Yan et al., 2016). Further, de Vries et al. (2019) recently showed that under drought conditions herb species change the quality rather than quantity of root exudates and induce higher microbial respiration, presumably facilitating nutrient capture via stimulating microorganisms. Therefore, the role of tree species composition as driver of microbial community composition and functioning may be more pronounced under unfavorable than favorable environmental conditions, but this has not been investigated until today.

A better understanding of the effects of tree species composition and site conditions on soil microorganisms has important implications for forest management, especially in temperate and boreal regions where tree species richness is low and managed forests typically are dominated by only one or a few species (Knoke et al., 2008; Hu et al., 2009; Bauhus et al., 2010). While Norway spruce currently is the most popular high-yielding timber species in Central Europe, in recent years it has have been damaged severely by extreme weather conditions and the outbreak of bark beetles, events that are predicted to occur more frequently and at higher severity in future (Biedermann et al., 2019; Brandl et al., 2020; Pettit et al., 2020). For reducing the risk of damage and maintaining economic gains, planting of mixed forests including non-native species such as Douglas-fir is increasingly discussed (Pretzsch et al., 2017; Ammer et al., 2018). To date, the impact of planting Douglas-fir on biodiversity and functioning of forests is little studied, and this applies in particular to the belowground system, although e.g., Douglas-fir has been suggested to affect soil chemistry in a similar way than spruce (Schmid et al., 2014). Taken together, we lack a comprehensive evaluation of forest types on the structure and functioning of soil microbial communities across environmental gradients.

Here, we investigate the structure and functioning of microbial communities in litter and soil of five forest types at sites differing in environmental conditions, in particular in soil nutrients but also rainfall and pH. Forest types include pure stands of European beech (*Fagus sylvatica*), Norway spruce (*Picea abies*), Douglas-fir (*Pseudotsuga menziesii*), and the mixtures of the respective

coniferous species with beech. European beech forms the climax tree species in lowland and lower montane regions of Central Europe (Leuschner et al., 2017). Norway spruce originally only occurred at higher mountain ranges and boreal regions in Central Europe, but has been planted across wide areas of Central Europe including lowlands (Knoke et al., 2008; Schmid et al., 2014). Douglas-fir has been introduced into Germany from North America more than 150 years ago and has become the most abundant non-native tree species cultivated in Central Europe (Schmid et al., 2014). To understand the response of microorganisms to planting different tree species in single and mixed stands, physiological (respiration based) and community approaches (phospholipid fatty acid analysis) were combined. We quantified microbial basal respiration, microbial biomass, and community structure as well as stress indicators of microorganisms (Anderson and Domsch, 2010). Physiological response variables and stress indicators were used as functional indicators of microbial communities. Generally, we assumed the structure of microbial communities to be linked to their functioning. Specifically, we hypothesized (1) coniferous trees to detrimentally affect microbial basal respiration, microbial biomass and stress indicators, with the effects being similar in Norway spruce and Douglas-fir forests, but alleviated in mixed forests with beech. Further, we hypothesized (2) effects of forest types on microbial community structure and functioning to be present at both nutrient-rich and nutrient-poor sites, but to be stronger at nutrient-poor sites. Since the litter layer is less buffered against environmental conditions than the organic and mineral soil, we also hypothesized that (3) the effects of forest type on the structure and functioning of microbial communities are more pronounced in litter than in soil.

2. Methods

2.1. Field sites

The study was conducted in 40 forest stands with half of the stands located in four sites in the south and the other half in four sites in the north of Lower Saxony, Germany, covering a range of environmental conditions (eight sites x five forest types; Fig. S1). The four sites in the south stock on fertile soil and receive higher precipitation. Parental rock is either loess influenced Triassic sandstone or Paleozoic shares of greywacke, sandstone, quartzite and phyllite, resulting in soil types of partly podsolic Cambisol and Luvisol. Mean annual precipitation is 821 – 1029 mm (calculated from German weather service since 1981; Glatthorn 2017). The four sites in the north are located on out-washed sand with the soil type of Podzol. The mean annual precipitation is 672 – 746 mm. The southern sites are richer in nutrients than the northern sites as reflected by lower C/N ratio, higher total soil P, higher pH and higher cation exchange capacity (CEC) (Table 1; Foltran et al., 2020). Hereafter, we refer to the southern sites as nutrient-rich sites and to the northern sites as nutrient-poor sites.

Each site comprised three pure stands of European beech (*Fagus sylvatica* L.; Be), Douglas-fir

(*Pseudotsuga menziesii* [Mirbel] Franco.; Do), Norway spruce (*Picea abies* [L.] Karst.; Sp) and two

conifer-beech mixtures (Douglas-fir/European beech and Norway spruce/European beech; Do/Be

and Sp/Be, respectively). On average, trees were more than 50 years old in 2017. Within sites, the

distance between stands ranged from 76 m to 4600 m, and the distance between sites ranged from

5 to 190 km. Within each stand, plots of 2500 m^2 were established, mostly in rectangular shape (50

m x 50 m). Tree species in pure stands comprised more than 90% of the total basal area, and that

in mixed stands 33 – 53% for beech and 53 – 60% for conifers.

151   2.2. Soil sampling and properties

Samples were taken between November 2017 and January 2018. Soil cores of 5 cm diameter were

taken by a metal cylinder, and were separated into litter, 0 – 5 and 5 – 10 cm soil depth. In each

plot, three cores spaced by 5 m were taken. Samples from the same depth were pooled, resulting in

a total of 120 samples (40 plots × 3 depths). The soil was sieved through 2 mm mesh, and the litter

was fragmented into pieces using scissors (< 25 mm^2) (Maraun and Scheu, 1995). Roots > 2 mm in

diameter and stones were removed. Samples were frozen at -20°C and thawed at 4°C before

further analyses. The pH of litter and soil was determined using a ratio of sample to solution (KCl,

1M) of 1:10, 1:5 and 1:2.5 (g/v) for litter, 0 – 5 and 5 – 10 cm soil, respectively. Moisture content

was measured by drying samples at 105°C for 48 h. Dried samples were grinded in a ball mill and

then total carbon and nitrogen concentrations were determined by an elemental analyzer (NA 1110,

CE-instruments, Rodano, Milano, Italy). Biomass of litter was estimated by weighing samples that

had been dried at 50°C for > 48 h (Macfadyen, 1961). Water content did not differ consistently

between nutrient-rich and nutrient-poor sites across layers, with higher water content in litter but

lower water content in soil at nutrient-poor than at nutrient-rich sites (interaction of Site condition

and Depth; $F_{2,68}$ = 26.21, P < 0.001); further site characteristics including soil texture and soil

chemical properties (percentage of carbon and nitrogen, C/N ratio, cation exchange capacity, pH,

concentration of phosphorus and Ca^{2+}) are given in Table 1.

169   2.3. Microbial basal respiration and biomass

Field moist samples were weighed and preincubated at room temperature (20°C) overnight before

placement in an automated micro-respirometer system (Scheu, 1992). The system absorbs CO_2

produced by microorganisms using KOH solution, and O_2 consumed by microorganisms is replaced

through electrolytic release of O_2 from $CuSO_4$ solution. Microbial basal respiration was measured as

mean consumption of O_2 during 10 – 23 h after attachment of the vessels to the respirometer

system (μg O_2 g^{-1} h^{-1}). Microbial biomass was determined by substrate-induced respiration after the

addition of glucose to saturate microbial glycolytic enzymes based on the maximum initial

respiratory response 4 – 7 h after the addition of D-glucose ($MIRR$; μl O_2 g^{-1} h^{-1}) and converted to

microbial biomass (C_{mic}; μg C_{mic} g^{-1}) as 38 × $MIRR$ (Anderson and Domsch, 1978; Beck et al.,

1997). Basal respiration and substrate-induced respiration were measured at 22°C in a water bath.

Fresh litter and soil from 0 – 5 and 5 – 10 cm depth equivalent to approximately 0.3, 1 and 3.5 g dry

mass was supplemented with 80, 20 and 8 mg glucose solved in 400 μl H_2O, respectively. The ratio

between microbial basal respiration and microbial biomass was taken as microbial specific

respiration (qO_2; μg O_2 μg^{-1} C_{mic} h^{-1}).

2.4. Phospholipid fatty acid analysis

To quantify the composition of phospholipid-derived fatty acid (PLFA) composition, lipids from leaf

litter and soil were extracted using a modified Bligh and Dyer method (Frostegård et al., 1993) as

described in Pollierer et al. (2015). Lipids were fractionated into glycolipids, phospholipids and

neutral lipids by elution through silica acid columns using chloroform, acetone and methanol,

respectively (0.5 g silicic acid, 3 ml; HF BOND ELUT-SI, Varian, Inc., Darmstadt, Germany).

Phospholipids were subjected to mild alkaline methanolysis and fatty acid methyl esters (FAMEs)

were identified by chromatographic retention time compared to a general reference standard and a

bacteria-derived standard (FAME CRM47885, C11 to C24; BAME 47080-U, C11 to C20; Sigma-

Aldrich, Darmstadt, Germany) using a GC-FID Clarus 500 (PerkinElmer Corporation, Norwalk, USA)

equipped with an Elite 5 column (30 m × 0.32 mm inner diameter, film thickness 0.25 μm). The

temperature program started with 60 °C (hold time 1 min) and increased by 30 °C per min to

160 °C, and then by 3 °C per min to 280 °C. The injection temperature was 250 °C and helium was

used as carrier gas. Approximately 2 g of fresh litter and 4 g of fresh soil were used for the

extraction.

2.5. Stress indicators and fatty acid markers

The ratio of cyclopropyl PLFAs to their monoenoic precursors (cy/pre) and the ratio of saturated to

monounsaturated PLFAs (sat/mono) were used as indicators of physiological or nutritional stress

([(cy17:0 + cy19:0)/(16:1ω7 + 18:1ω7)]; [(14:0 + 15:0 + 16:0 + 17:0 + 18:0)/(16:1ω7 + 17:1 +

18:1ω9 + 18:1ω7)]; respectively) (Moore-Kucera and Dick, 2008; Pollierer et al., 2015). The ratio of

Gram+ to Gram- bacteria was used as indicator of carbon availability (Fanin et al., 2019). The

saturated fatty acids i15:0, a15:0, i16:0, i17:0 were used as markers for Gram+ bacteria, and the

fatty acids cy17:0, cy19:0, 16:1ω7 and 18:1ω7 were assigned as markers for Gram- bacteria

(Frostegård and Bååth, 1996; Zelles, 1999; Mitchell et al., 2010; Richter et al., 2018; Fanin et al.,

2019). Bacterial biomass was represented by the sum of Gram+ and Gram- bacteria. Linoleic acid

18:2ω6,9 was used as fungal marker (Frostegård and Bååth, 1996). Total amount of PLFAs were

summed up from all identified PLFAs (n = 40; nmol g^{-1} dry weight) and was used to calculate PLFA

proportions. All stress indicators and FA markers were analyzed from proportions (mole

percentage).

2.6. Statistical analyses

For univariate analysis, we first fitted linear mixed models (LMMs) to log-transformed response

variables and then applied planned contrasts to estimate the effect size of Forest type and Site

condition (Piovia-Scott et al., 2019). All LMMs included Forest type (European beech, Douglas-fir,

Douglas-fir/European beech, Norway spruce, Norway spruce/European beech), Site condition

(nutrient-rich and nutrient-poor sites) and Depth (litter, $0 – 5$ and $5 – 10$ cm) as fixed effects. Full

models were selected stepwise by likelihood ratio test, and minimal models included all main effects

and the interaction of Forest type and Site condition. The 40 forest plots were included as random

effects to account for non-independence of samples of different depth from the same plot by forcing

values from the same plot to have the same intercept (Harrison et al., 2018). Based on Akaike

Information Criterion (AIC), the eight sites were included as random effect, and the mean-variance

relationship was accounted for by a dispersion parameter to meet the assumption of homogeneity of

variance (Johnson and Omland, 2004; Zuur et al., 2009).

Contrasts for the effect size of forest type were designed to compare coniferous and mixed forests

to beech forests. European beech, the climax tree species in lowland and lower montane regions of

Central Europe, was used as reference (Leuschner et al., 2017). Due to the property of log-

transformed response variables, the planned contrasts are analog to log response ratios (Piovia-

Scott et al., 2019). Further, the log response ratios were back transformed into response ratios and

were defined as effect sizes. Effect sizes of forest type were estimated for nutrient-poor and

nutrient-rich sites based on minimal models. Additionally, we applied contrasts between nutrient-

poor vs. nutrient-rich sites in a similar way for the effect sizes of Site condition. Univariate response

variables included microbial basal respiration, microbial biomass, microbial specific respiration,

stress indicators and PLFA markers.

For multivariate analysis of variance (MANOVA), PLFAs first were arcsine root transformed and

then dimensions of fatty acids were reduced by principal component analysis (PCA). Separate

MANOVAs for each sample depth were calculated to inspect for effects of Forest type, Site

condition, and their interactions. Further, selected environmental variables were included into

MANOVAs as covariates to test whether they explain effects of main factors in the models

(Langerhans et al., 2007; Bennett et al., 2020). Significant principal components (PCs) were

determined by broken stick criterion (MacArthur, 1957). Covariates were selected according to

significance in permutation tests based on adjusted R^2 in redundancy analyses (RDA) (Oksanen et
al., 2019). Only fatty acids ≥0.2% mean mole proportion were included in the analyses.

All analyses were done in R with RStudio interface (R Core Team, 2019; RStudio Team, 2019). We
used the "nlme" package to fit LMMs (Pinheiro et al., 2019), and the "emmeans" package to conduct
planned contrasts (Lenth, 2020). All mixed models met the assumptions of normality of residuals
and homogeneity of variance.

3. Results

3.1. Functional indicators of microorganisms

Forest type consistently affected microbial basal respiration, microbial biomass and stress indicators
at nutrient-poor but not at nutrient-rich sites (Forest type x Site condition interactions; all P < 0.08;
Table 2, Figs 1 and 2), and this was generally true across soil layers (Fig. S2). At nutrient-poor sites,
microbial basal respiration and microbial biomass was lower in coniferous and mixed forests than in
beech forests. This was most pronounced in Douglas-fir, where basal respiration and microbial
biomass at nutrient-poor sites were 65% and 59% lower compared to pure beech, respectively;
similarly, basal respiration and microbial biomass in the two conifer-beech mixtures were 46 – 56%
lower than in beech. Microbial basal respiration and microbial biomass in spruce forests were also
lower than in beech forests, but the difference was less pronounced (-33 to -36%; Fig. 1). In
addition, in Douglas-fir forests at nutrient-poor sites microbial specific respiration was 19% lower
compared to beech forests.

In contrast to basal respiration and microbial biomass, at nutrient-poor sites stress indicators, the
cy/pre ratio, the sat/mono ratio as well as the Gram$^+$/Gram$^-$ bacteria ratio, all were lower in beech
forests than in the other forest types (Fig. 2). The difference in the cy/pre and sat/mono ratio was
most pronounced in Douglas-fir (+87% and +74%) and less in spruce forests (+39% and +43%).
Similar to spruce forests, the cy/pre and sat/mono ratio in conifer-beech mixtures were on average
37 – 46% higher than in beech forests. Effects of forest type on the Gram$^+$/Gram$^-$ bacteria ratio also
was strongest in Douglas-fir forests (+46 %) and less pronounced in the other forest types (+21 –
29%). Stress indicators generally did not differ at nutrient-rich sites (P > 0.31; Fig. 2, Fig. S2, Table
S1).

The response of microorganisms to changes in site conditions also varied between forest types.
Differences between nutrient-rich and nutrient-poor sites were most prominent in Douglas-fir forests
and not significant in beech forests (Figs 3 and 4, Table S2). At nutrient-poor sites, microbial basal
respiration in Douglas-fir forests was 57% lower and microbial biomass in Douglas-fir pure and

mixed forests both were 47% lower compared to nutrient-rich sites. Similarly, but less strong,

microbial biomass in spruce mixed forests was 36% lower at nutrient-poor sites than nutrient-rich

sites. Conversely, in Douglas-fir forests the stress indicator cy/pre ratio at nutrient-poor sites

exceeded that of nutrient-rich sites by 82% (Fig. 4, Table S2). Similarly, in Douglas-fir forests the

sat/mono and Gram$^+$/Gram$^-$ bacteria ratio at nutrient-poor sites exceeded that at nutrient-rich sites

by 38 – 40%. Further, in Douglas-fir mixed forests the Gram$^+$/Gram$^-$ bacteria ratio at nutrient-poor

sites exceeded that at nutrient-rich sites by 34%. Further, the response of microbial stress indicators

to changes in site conditions varied with soil depth; in particular, the responses of the sat/mono ratio

and the Gram$^+$/Gram$^-$ ratio were most pronounced in 0 – 5 cm soil depth (Fig. 4, Table 2).

3.2. Microbial community structure

As indicated by PLFA patterns, microbial community composition in beech forests differed from that

in Norway spruce and Douglas-fir forests, with mixed forests being intermediate between the

respective single species stands (Fig. 5). Across layers, effects of forest type were significant in litter

and 0 – 5 cm soil, and marginally significant in 5 – 10 cm soil (Table 3). Despite strong turnover

between nutrient-rich and nutrient-poor sites (Site condition effects; $p < 0.001$), beech maintained

similar microbial community structure between nutrient-poor and nutrient-rich sites, and this was

most apparent in 0 – 5 cm soil depth (Forest type × Site condition interaction; Table 3, Fig. 5).

A number of fatty acids contributed to the separation of microbial communities of the different forest

types and site conditions. Beech forests were characterized by unsaturated fatty acids at all soil

depths, in particular by 16:1ω7, 18:1ω7, 18:1ω9 (Fig. 5). Norway spruce forests were associated

with long-chain saturated fatty acids (22:0, 24:0), and Douglas-fir forests with bacterial marker fatty

acids (2OH-14:0, cy17:0). There also were clear differences in fatty acid patterns between nutrient-

rich and nutrient-poor sites. Litter at nutrient-rich sites was associated with unsaturated fatty acids

such as 18:1ω9 and the fungal marker 18:2ω6,9. By contrast, litter at nutrient-poor sites had higher

saturated and branched fatty acids, with many of them being bacterial markers, such as 14:0 and

cy17:0 and cy19:0.

Litter at nutrient-rich sites was generally richer in fungi than at nutrient-poor sites, but the opposite

was true for the soil (Site condition × Depth interaction; Table 2, Fig. S3). As indicated by PLFA

markers, the relative abundance of fungi and bacteria generally did not differ among forest types,

but the relative abundance of fungi consistently decreased, whereas the abundance of bacteria

increased with soil depth (Figs S3 and S4). Further, based on the contrast of Site condition, at

nutrient-poor sites the relative abundance of fungi in pure beech and spruce forests was higher than

at nutrient-rich sites. By contrast, in pure and mixed Douglas-fir forests at nutrient-poor sites the

relative abundance of fungi in soil was similar to that at nutrient-rich sites (Table S2, Fig. S3).

Nutrient-poor sites were more limited by nitrogen as indicated by higher C/N ratio but also had lower

soil pH (Fig. 5). Soil C/N ratio and pH as well as soil moisture were among the most important

environmental variables in explaining variations in PLFA composition, but their contribution varied

with soil depth (Table S3). Among the environmental variables studied (pH, C/N ratio, water content,

carbon content, litter mass), pH accounted for 8% out of the total of 13% explained by all

environmental variables in litter, whereas in 0 – 5 and 5 – 10 cm soil depth C/N ratio accounted for

22% and 13% out of the total of 37% and 15% of the variation explained by all environmental

variables (Table S3). Further, the main effects of Site condition became non-significant in soil after

including C/N ratio, pH and water content (0 – 5 cm soil depth) and C/N ratio (5 – 10 cm soil depth)

as covariates in the MANOVAs, but the effects of forest type remained marginally significant in both

0 – 5 and 5 – 10 cm soil depth (Table 3, Fig. 5). By contrast, in litter, including pH as covariate into

the MANOVA did not significantly affect the effects of Forest type and Site condition.

4. Discussion

Overall, the results only in part support our first and second hypotheses. Forest types only affected

the functioning of microbial communities at nutrient-poor sites, but not at nutrient-rich sites. At

nutrient-poor sites, microbial basal respiration, microbial biomass and stress indicators all were

detrimentally affected in pure and mixed coniferous forests, and, contrasting our first hypothesis the

effects differed between Douglas-fir and Norway spruce forests, with the effects being particularly

strong in Douglas-fir, but less pronounced in Norway spruce and mixed stands. By contrast, similar

basal respiration, microbial biomass and stress indicators at nutrient-rich sites across the studied

forest types contradict our second hypothesis and suggest that, at favorable environmental

conditions, soil microbial communities are rather irresponsive to changes in tree species. Notably,

microbial biomass and microbial stress indicators in beech forests did not differ between nutrient-

poor and nutrient-rich sites indicating that beech is able to maintain the functioning of soil microbial

communities regardless of site conditions. Admixture of beech in mixed stands with Douglas-fir or

Norway spruce both detrimentally affected microbial biomass and microbial stress indicators,

suggesting that caution is needed when mixing beech with conifers. However, results of the present

study also indicate that compared to pure Douglas-fir forests, mixed Douglas-fir forests with beech

mitigate the detrimental effects of pure Douglas-fir forests on soil microbial communities. Exotic

plants lack co-evolutionary history with native species and, therefore, complementary effects

between native and exotic plants are more likely to occur than between native species (Kawaletz et

al., 2013). This may explain the alleviation of microbial stress in mixtures of Douglas-fir with beech.

The site- and tree species-specific responses of microorganisms are likely caused by varying

energy supply in different forest types at nutrient-poor sites. In fact, at nutrient-poor sites, the higher

Gram+/Gram- bacteria ratio in coniferous and mixed forests suggests lower supply of labile carbon

than in beech forests (Fanin et al., 2019). Labile carbon compounds have been suggested to favor

Gram- bacteria, whereas Gram+ bacteria better cope with recalcitrant carbon resources (Kramer and

Gleixner, 2008; Fanin et al., 2019; Waller et al., 2020). Among the forest types studied, Douglas-fir

forests were particularly limited by labile carbon as further supported by its lower microbial specific

respiration, which was largely due to lower microbial basal respiration compared to beech stands

(Anderson and Domsch, 2010). Limited supply of labile carbon also has been suggested to favor

oligotrophic microbial communities characterized by low respiration rates and biomass (Potthast et

al., 2010; Fanin et al., 2019). This is consistent with our findings of strong carbon limitation as well

as low basal respiration and microbial biomass in single and mixed coniferous forests at nutrient-

poor sites. Moreover, at nutrient-poor sites, both the cy/pre and sat/mono ratios were higher in pure

coniferous as well as mixed forests than in pure beech forests, also indicating higher nutrient

(and/or water) stress in the former (Moore-Kucera and Dick, 2008; Pollierer et al., 2015). The Gram-

bacteria marker fatty acids 16:1ω7 and 18:1ω7 are precursors of cyclopropyl fatty acids and as

monounsaturated fatty acids they form part of the nominator of the calculated stress indicators. The

more pronounced stress response of microorganism in coniferous and mixed forests at nutrient-poor

sites largely was caused by lower proportions of these two ω7 fatty acids. Indeed, it has been

suggested that under stress conditions Gram- bacteria change the composition of their cell

membranes from 16:1ω7 and 18:1ω7 to cyclopropane fatty acids associated with slower growth

rates (Guckert et al., 1986; Kieft et al., 1994; Lundquist et al., 1999). The similarity of microbial

indicators in beech forests at nutrient-rich and nutrient-poor sites indicates that, compared to

coniferous stands, beech maintained microbial community functioning and avoided stress conditions

of microorganisms by providing more labile carbon resources at nutrient-poor conditions,

presumably by releasing high amounts of root-derived resources (Meier et al., 2020).

The suggestion that beech is able to alleviate microbial stress via maintaining or even increasing the

release of root-derived resources at nutrient-poor sites is supported by the following findings. First,

current studies at our study sites showed that at nutrient-poor sites the biomass of fine roots is

higher in beech forests than in pure and mixed coniferous forests while at nutrient-rich sites root

biomass varies little among forest types (A. Lwila, pers. comm.). Generally, it is well established that

plants allocate surplus carbon into roots if the availability of soil nutrients is declining and this is

associated with increased root exudation (Eisenhauer et al., 2017; Laliberté et al., 2017; Prescott et

al., 2020). Recently, this also was confirmed for beech forests where higher root exudation was

found in more acidic and nitrogen deficient stands (Meier et al., 2020). These findings support our

suggestion that beech indeed increased its root exudates at nutrient-poor sites, which in fact were

characterized by higher C/N ratio and more acidic conditions in soil. High root-derived resources

presumably also were responsible for the alleviated microbial resource deficiency and stress in

beech forests at nutrient-poor sites. Second, trenching and girdling experiments also confirmed that

reducing or omitting the flux of root-derived resources into the soil strongly reduces fungal and

bacterial biomass, demonstrating the importance of root-derived resources for maintaining soil

microbial biomass (Högberg et al., 2007; Kaiser et al., 2010; Bluhm et al., 2019). Overall, these

findings suggest that beech has the potential to facilitate soil microorganisms at environmental

stress conditions by increasing its root exudates contrasting both of the studied coniferous species.

This ability presumably contributed to the mitigation of the detrimental effects of Douglas-fir on

microorganisms at nutrient-poor / acidic sites in mixed forests. More broadly, it may explain the

observed overyielding in mixed stands of beech and conifers at nutrient-poor sites, highlighting the

potential role of microorganisms in the facilitation of plants under environment stress conditions

(Toïgo et al., 2015; Ammer, 2019; David et al., 2020).

Contrary to our third hypothesis, the effect of forest type was not stronger in litter than in soil. At

nutrient-poor sites, forest type affected microbial basal respiration, microbial biomass and microbial

stress in both litter and soil, with the strength of the effects generally being similar across layers.

This supports our suggestion that the effect of forest type was not mainly due to differences in litter

quantity and quality, but in particular to differences in root-derived resources. This contrasts earlier

suggestions that microorganisms in litter are more sensitive to changes in environmental conditions

than those in soil because litter is less buffered against environmental harshness (Pollierer et al.,

2015). Supporting our suggestion that microorganisms in soil sensitively respond to environmental

changes, microbial biomass generally decreased while microbial stress indicators generally

increased with soil depth. This sensitivity presumably is related to the fact that generally carbon

availability decreases with soil depth resulting in stronger resource limitation (Fierer et al., 2003). As

indicated by sat/mono and $Gram^+$/$Gram^-$ ratios, this increase in resource limitation is particularly

strong from litter to soil in pure coniferous and mixed forests at nutrient-poor sites.

We assumed that the functioning of microorganisms ultimately relies on its microbial community

structure, and this is supported by the consistent turnover in microbial community composition as

well as the change in microbial functions between nutrient-rich and nutrient-poor sites. Further, in

contrast to coniferous and mixed forests, microbial community structure and functional indices in

beech forests differed little between nutrient-rich and nutrient-poor sites, suggesting that among the

tree species studied only beech is able to maintain and stabilize microbial community structure and

functioning in litter and soil irrespective of site conditions. Notably, differences in microbial

community structure between forest types at nutrient-rich sites tended to be small in 5 – 10 cm soil

depth, further suggesting that the influence of tree species fades deeper in soil. By contrast, at

nutrient-poor sites, differences in microbial community structure between forest types were relatively

large in soil, again supporting our hypothesis that at nutrient-poor sites effects of tree species on

microorganisms in soil are predominantly due to differences in root-derived resources. Interestingly,

the differences in microbial community structure between beech and coniferous forests in 5 – 10 cm

soil depth at nutrient-poor sites were most pronounced in Douglas-fir, and carbon limitation also was

most prominent in 5 – 10 cm soil of Douglas-fir forests as indicated by lower microbial specific

respiration. This implies that among the tree species studied Douglas-fir provides the lowest amount

of root-derived resources to soil microorganisms at nutrient-poor sites resulting in pronounced

resource shortage and marked changes in microbial community composition. Such changes in

microbial community structure may significantly impact the functioning of soil microbial communities

as the availability of carbon resources and carbon limitation affect the vertical distribution of

saprotrophic fungi and decomposition processes, e.g. by shifting the competition between

saprotrophic and mycorrhizal fungi (Gadgil and Gadgil, 1971; Lindahl et al., 2007).

Our results are consistent with earlier findings indicating that regional factors more strongly shape

microbial community structure than tree species (Pollierer et al., 2015; Richter et al., 2018) as the

relative abundance of fungi and bacteria as well as the fungi/bacteria ratio did not vary with forest

types but with site conditions. As indicated by the relative abundance of the fungal marker FA

18:2ω6,9, fungi generally were favored in the soil of nutrient-poor sites, supporting earlier

suggestions that the fungal energy channel dominates in infertile systems (Wardle et al., 2004).

However, contrasting earlier suggestions, in Douglas-fir forests at nutrient-poor sites the fungal

channel did not dominate, although they were also characterized as infertile with high C/N ratio and

low pH (Fig. 5). In fact, at nutrient-poor sites the relative abundance of fungi in soil of Douglas-fir

forests was similar to that at nutrient-rich sites, resulting in lower abundance of fungi in Douglas-fir

compared to beech forests based on planned contrasts, particularly in 5 – 10 cm soil depth. The

relatively low abundance of fungi in soil of Douglas-fir forests may be due to a mismatch between

local and home (native range of Douglas-fir) fungal communities (Schmid et al., 2014), as in soil of

introduced plant species only a subset of fungal species present in their native range has been

recorded in soil of sites where they had been introduced (Schmid et al., 2014). Overall, the results of

our study support the sensitivity of PLFA analysis in detecting changes in microbial community

composition with forest types (Ramsey et al., 2006; Frostegård et al., 2011). However, future

studies using molecular techniques are needed to uncover the microbial taxa responsible for the

observed differences among forest types, including introduced and native tree species such as

Douglas-fir and beech in particular at nutrient-poor sites deeper in soil.

Notably, we sampled litter and soil during autumn / winter after beech had shed its leaves. The

hypothesis of high root-derived resources in beech forests implies that the impact of roots on

microorganisms still was strong in autumn / winter indicating that not only recent photosynthates but

also stored carbohydrates contribute to providing root-derived resources to soil microorganisms

(Druebert et al., 2009). However, it remains open whether the effects of forest type on microbial

communities get stronger or weaker with time hampering generalization of our finding across

seasons. Despite microbial communities may fluctuate considerably with season (Koranda et al.,

2013; Abramoff and Finzi, 2016; Nacke et al., 2016), comprehensive sampling in Douglas-fir forests

across seasons suggested that, except during summer drought, microbial stress indicators fluctuate

little (Moore-Kucera and Dick, 2008). In addition, as our sampling was limited to the upper 10 cm of

the soil, we may not have captured the full effects of roots, as e.g., roots of beech may extend

considerably deeper into the soil (Leuschner et al., 2004), although fine roots generally are

concentrated in upper soil layers (Schmid, 2002; Leuschner et al., 2004). To better understand the

linkage between roots and microbial communities, more information is needed on the vertical

distribution of fine roots as well as differences in root exudation between the studied tree species

across nutrient gradients.

In our study, among the soil chemical properties studied, C/N ratio and pH best explained the

variations in microbial community structure. Typically, soil pH correlates negatively with soil nutrient

availability (Meier et al., 2020), further supporting our conclusion on the importance of nutrients in

driving the responses of microorganisms to variations in site conditions. However, factors other than

soil nutrient status and soil pH may have contributed to the observed differences between nutrient-

rich and nutrient-poor sites. Despite moisture in litter and soil contributed little to microbial

community structure, nutrient-poor sites also are characterized by lower precipitation and this is

reflected by higher $\delta^{13}C$ values of litter at nutrient-poor compared to nutrient-rich sites (J.-Z. Lu,

unpubl. data), likely reflecting higher water stress at nutrient-poor compared to nutrient-rich sites

(Peuke et al., 2006). Presumably, in addition to nutrients, both pH and water stress contributed to

the observed differences in microbial community composition and functioning between nutrient-poor

and nutrient-rich sites. This is supported by the fact that soil water conditions are intricately linked to

soil nutrient dynamics and the uptake of nutrients by plants (de Vries et al., 2019). Future studies

need to disentangle the relative importance of nutrient availability, soil pH and water deficiency as

driving factors of the interrelationship between forest stands and soil microbial community

composition and functioning. Focusing on nutrient-poor / low pH stands is most promising for
resolving these interrelationships.

5. Conclusion

The present study documented that effects of forest type on soil microbial communities vary among
tree species and mixed stands, but this strongly depends on the site conditions, in particular soil
nutrients. Notably, variations between sites were least pronounced in native beech and most
pronounced in introduced Douglas-fir. Compared to beech, Douglas-fir strongly impacted microbial
community structure and functioning at nutrient-poor sites. Therefore, in the long-term planting non-
native tree species such as Douglas-fir may compromise ecosystem functioning at nutrient-poor
sites, and this needs to be studied in more detail as it may not only question the establishment of
pure Douglas-fir forests, but also the enrichment of beech forests by Douglas-fir at nutrient-poor
sites. However, as planting Douglas-fir in single species and mixed stands with beech hardly
impacted the structure and functioning of soil microbial communities at nutrient-rich sites, planting
Douglas-fir may provide an alternative option to Norway spruce at least at nutrient-rich sites. Such
options are urgently needed since Norway spruce forests are increasingly threatened due to global
climate change and associated outbreaks of bark beetles especially in lowland regions of Europe.
However, more information is needed for gaining a holistic understanding on the consequences of
planting single species or mixed stands of Douglas-fir on ecosystem properties and processes, such
as below- and aboveground biodiversity of other taxa, and the decomposition of litter. Future studies
need to focus on potential facilitation or inhibition of planting mixed forests of beech and Douglas-fir
on soil microorganisms and timber yield in particular at nutrient-poor sites. Overall, our findings
highlight that tree species differently shape the structure and functioning of soil microbial
communities with their effects varying strongly with site conditions.

Acknowledgements

This work was supported by the German Research Foundation [316045089] as part of the Research
Training Group (RTG) 2300: "Enrichment of European beech forests with conifers: impacts of
functional traits on ecosystem functioning" (DFG: RTG 2300). We thank C. Bluhm, T. Volovei, G.
Humpert, S. Zhou for assistance, and C. Ammer, H. Riebl, A. Polle, J. Glatthorn, X. Lu, N.
Lamersdorf, Z. Li, S. Bluhm, G. Schulz, T.-W. Chen, M. Pollierer, A. Lwila, K. Schumann, I. Bebre,
A. Rivera and other RTG members for helpful discussions. We are grateful to E. Foltran and J.
Glatthorn for providing data on site description. Special thanks go to S. Müller for coordination.

References

Abramoff, R.Z., Finzi, A.C., 2016. Seasonality and partitioning of root allocation to rhizosphere soils in a midlatitude forest. Ecosphere 7. doi:10.1002/ecs2.1547

Ammer, C., 2019. Diversity and forest productivity in a changing climate. New Phytologist 221, 50–66. doi:10.1111/nph.15263

Ammer, C., Fichtner, A., Fischer, A., Gossner, M.M., Meyer, P., Seidl, R., Thomas, F.M., Annighöfer, P., Kreyling, J., Ohse, B., Berger, U., Feldmann, E., Häberle, K.H., Heer, K., Heinrichs, S., Huth, F., Krämer-Klement, K., Mölder, A., Müller, J., Mund, M., Opgenoorth, L., Schall, P., Scherer-Lorenzen, M., Seidel, D., Vogt, J., Wagner, S., 2018. Key ecological research questions for Central European forests. Basic and Applied Ecology 32, 3–25. doi:10.1016/j.baae.2018.07.006

Anderson, J.P.E., Domsch, K.H., 1978. A physiological method for the quantitative measurement of microbial biomass in soils. Soil Biology and Biochemistry 10, 215–221. doi:10.1016/0038-0717(78)90099-8

Anderson, T.H., Domsch, K.H., 2010. Soil microbial biomass: The eco-physiological approach. Soil Biology and Biochemistry 42, 2039–2043. doi:10.1016/j.soilbio.2010.06.026

Augusto, L., De Schrijver, A., Vesterdal, L., Smolander, A., Prescott, C., Ranger, J., 2015. Influences of evergreen gymnosperm and deciduous angiosperm tree species on the functioning of temperate and boreal forests. Biological Reviews 90, 444–466. doi:10.1111/brv.12119

Averill, C., Turner, B.L., Finzi, A.C., 2014. Mycorrhiza-mediated competition between plants and decomposers drives soil carbon storage. Nature 505, 543–545. doi:10.1038/nature12901

Balvanera, P., Pfisterer, A.B., Buchmann, N., He, J.S., Nakashizuka, T., Raffaelli, D., Schmid, B., 2006. Quantifying the evidence for biodiversity effects on ecosystem functioning and services. Ecology Letters 9, 1146–1156. doi:10.1111/j.1461-0248.2006.00963.x

Bardgett, R.D., van der Putten, W.H., 2014. Belowground biodiversity and ecosystem functioning. Nature 515, 505–511. doi:10.1038/nature13855

Bauhus, J., van der Meer, P.J., Kanninen, M., 2010. Plantation forests: global perspectives, in: Ecosystem Goods and Services from Plantation Forests. Earthscan, pp. 1–16. doi:10.4324/9781849776417

Beck, T., Joergensen, R.G., Kandeler, E., Makeschin, F., Nuss, E., Oberholzer, H.R., Scheu, S., 1997. An inter-laboratory comparison of ten different ways of measuring soil microbial biomass C. Soil Biology and Biochemistry 29, 1023–1032. doi:10.1016/S0038-0717(97)00030-8

Bennett, J.A., Koch, A.M., Forsythe, J., Johnson, N.C., Tilman, D., Klironomos, J., 2020. Resistance of soil biota and plant growth to disturbance increases with plant diversity. Ecology Letters 23, 119–128. doi:10.1111/ele.13408

Biedermann, P.H.W., Müller, J., Grégoire, J.C., Gruppe, A., Hagge, J., Hammerbacher, A., Hofstetter, R.W., Kandasamy, D., Kolarik, M., Kostovcik, M., Krokene, P., Sallé, A., Six, D.L., Turrini, T., Vanderpool, D., Wingfield, M.J., Bässler, C., 2019. Bark beetle population dynamics in the Anthropocene: Challenges and solutions. Trends in Ecology and Evolution. doi:10.1016/j.tree.2019.06.002

Bluhm, S.L., Eitzinger, B., Ferlian, O., Bluhm, C., Schröter, K., Pena, R., Maraun, M., Scheu, S.,
2019. Deprivation of root-derived resources affects microbial biomass but not community
structure in litter and soil. PLoS ONE 14, e0214233. doi:10.1371/journal.pone.0214233

Brandl, S., Paul, C., Knoke, T., Falk, W., 2020. The influence of climate and management on
survival probability for Germany's most important tree species. Forest Ecology and
Management 458, 117652. doi:10.1016/j.foreco.2019.117652

Callaway, R.M., Pennings, S.C., Richards, C.L., 2003. Phenotypic plasticity and interactions among
plants. Ecology 84, 1115–1128. doi:10.1890/0012-9658(2003)084[1115:PPAIAP]2.0.CO;2

Cardinale, B.J., Wright, J.P., Cadotte, M.W., Carroll, I.T., Hector, A., Srivastava, D.S., Loreau, M.,
Weis, J.J., 2007. Impacts of plant diversity on biomass production increase through time
because of species complementarity. Proceedings of the National Academy of Sciences of the
United States of America 104, 18123–18128. doi:10.1073/pnas.0709069104

Chen, L., Xiang, W., Wu, H., Ouyang, S., Zhou, B., Zeng, Y., Chen, Y., Kuzyakov, Y., 2019. Tree
species identity surpasses richness in affecting soil microbial richness and community
composition in subtropical forests. Soil Biology and Biochemistry 130, 113–121.
doi:10.1016/j.soilbio.2018.12.008

Chen, X., Chen, H.Y.H., Chen, C., Peng, S., 2019. Water availability regulates negative effects of
species mixture on soil microbial biomass in boreal forests. Soil Biology and Biochemistry 139,
107634. doi:10.1016/j.soilbio.2019.107634

Clemmensen, K.E., Bahr, A., Ovaskainen, O., Dahlberg, A., Ekblad, A., Wallander, H., Stenlid, J.,
Finlay, R.D., Wardle, D.A., Lindahl, B.D., 2013. Roots and associated fungi drive long-term
carbon sequestration in boreal forest. Science 340, 1615–1618. doi:10.1126/science.1231923

David, A.S., Thapa-Magar, K.B., Menges, E.S., Searcy, C.A., Afkhami, M.E., Thapa-Magar, K.B.,
Menges, E.S., Searcy, C.A., Afkhami, M.E., Thapa-Magar, K.B., Menges, E.S., Searcy, C.A.,
Afkhami, M.E., Thapa-Magar, K.B., Menges, E.S., Searcy, C.A., Afkhami, M.E., 2020. Do
plant–microbe interactions support the Stress Gradient Hypothesis? Ecology 00, e03081.
doi:10.1002/ecy.3081

Dawud, S.M., Raulund-Rasmussen, K., Domisch, T., Finér, L., Jaroszewicz, B., Vesterdal, L., 2016.
Is tree species diversity or species identity the more important driver of soil carbon stocks, C/N
ratio, and pH? Ecosystems 19, 645–660. doi:10.1007/s10021-016-9958-1

de Vries, F.T., Williams, A., Stringer, F., Willcocks, R., McEwing, R., Langridge, H., Straathof, A.L.,
2019. Changes in root-exudate-induced respiration reveal a novel mechanism through which
drought affects ecosystem carbon cycling. New Phytologist 224, 132–145.
doi:10.1111/nph.16001

Defossez, E., Courbaud, B., Marcais, B., Thuiller, W., Granda, E., Kunstler, G., 2011. Do
interactions between plant and soil biota change with elevation? A study on *Fagus sylvatica*.
Biology Letters 7, 699–701. doi:10.1098/rsbl.2011.0236

Druebert, C., Lang, C., Valtanen, K., Polle, A., 2009. Beech carbon productivity as driver of
ectomycorrhizal abundance and diversity. Plant, Cell and Environment 32, 992–1003.
doi:10.1111/j.1365-3040.2009.01983.x

Eisenhauer, N., Lanoue, A., Strecker, T., Scheu, S., Steinauer, K., Thakur, M.P., Mommer, L., 2017.
Root biomass and exudates link plant diversity with soil bacterial and fungal biomass. Scientific
Reports 7. doi:10.1038/srep44641

Ekblad, A., Nordgren, A., 2002. Is growth of soil microorganisms in boreal forests limited by carbon
or nitrogen availability? Plant and Soil 242, 115–122. doi:10.1023/A:1019698108838

Emmett Duffy, J., Godwin, C.M., Cardinale, B.J., 2017. Biodiversity effects in the wild are common
and as strong as key drivers of productivity. Nature 549, 261–264. doi:10.1038/nature23886

Fanin, N., Kardol, P., Farrell, M., Nilsson, M.C., Gundale, M.J., Wardle, D.A., 2019. The ratio of
Gram-positive to Gram-negative bacterial PLFA markers as an indicator of carbon availability
in organic soils. Soil Biology and Biochemistry 128, 111–114. doi:10.1016/j.soilbio.2018.10.010

Fierer, N., Schimel, J.P., Holden, P.A., 2003. Variations in microbial community composition through
two soil depth profiles. Soil Biology and Biochemistry 35, 167–176. doi:10.1016/S0038-
0717(02)00251-1

Foltran, E.C., Ammer, C., Lamersdorf, N., 2020. Douglas fir and Norway spruce admixtures to
beech forests along a site gradient in Northern Germany – Are soil nutrient conditions
affected? BioRxiv. doi:https://doi.org/10.1101/2020.09.25.313213

Frostegård, A., Bååth, E., 1996. The use of phospholipid fatty acid analysis to estimate bacterial and
fungal biomass in soil. Biology and Fertility of Soils 22, 59–65. doi:10.1007/s003740050076

Frostegård, Å., Tunlid, A., Baath, E., 1993. Phospholipid fatty acid composition, biomass, and
activity of microbial communities from two soil types experimentally exposed to different heavy
metals. Applied and Environmental Microbiology 59, 3605–3617. doi:10.1128/aem.59.11.3605-
3617.1993

Frostegård, Å., Tunlid, A., Bååth, E., 2011. Use and misuse of PLFA measurements in soils. Soil
Biology and Biochemistry 43, 1621–1625. doi:10.1016/j.soilbio.2010.11.021

Fujii, S., Berg, M.P., Cornelissen, J.H.C., 2020. Living litter: Dynamic trait spectra predict fauna
composition. Trends in Ecology and Evolution 00. doi:10.1016/j.tree.2020.05.007

Gadgil, R.L., Gadgil, P.D., 1971. Mycorrhiza and litter decomposition. Nature 233, 133.
doi:10.1038/233133a0

Glatthorn, J., 2017. plot_general. v1.10.8. RTG2300-Information System. Dataset.
https://www.134.76.21.10/Data/ShowData.aspx?DatasetId=12540

Graham, E.B., Knelman, J.E., Schindlbacher, A., Siciliano, S., Breulmann, M., Yannarell, A.,
Beman, J.M., Abell, G., Philippot, L., Prosser, J., Foulquier, A., Yuste, J.C., Glanville, H.C.,
Jones, D.L., Angel, R., Salminen, J., Newton, R.J., Bürgmann, H., Ingram, L.J., Hamer, U.,
Siljanen, H.M.P., Peltoniemi, K., Potthast, K., Bañeras, L., Hartmann, M., Banerjee, S., Yu,
R.Q., Nogaro, G., Richter, A., Koranda, M., Castle, S.C., Goberna, M., Song, B., Chatterjee, A.,
Nunes, O.C., Lopes, A.R., Cao, Y., Kaisermann, A., Hallin, S., Strickland, M.S., Garcia-
Pausas, J., Barba, J., Kang, H., Isobe, K., Papaspyrou, S., Pastorelli, R., Lagomarsino, A.,
Lindström, E.S., Basiliko, N., Nemergut, D.R., 2016. Microbes as engines of ecosystem
function: When does community structure enhance predictions of ecosystem processes?
Frontiers in Microbiology 7, 214. doi:10.3389/fmicb.2016.00214

Guckert, J.B., Hood, M.A., White, D.C., 1986. Phospholipid ester-linked fatty acid profile changes
during nutrient deprivation of Vibrio cholerae: Increases in the trans/cis ratio and proportions of
cyclopropyl fatty acids. Applied and Environmental Microbiology 52, 794–801.
doi:10.1128/aem.52.4.794-801.1986

Harpole, W.S., Sullivan, L.L., Lind, E.M., Firn, J., Adler, P.B., Borer, E.T., Chase, J., Fay, P.A.,
Hautier, Y., Hillebrand, H., MacDougall, A.S., Seabloom, E.W., Williams, R., Bakker, J.D.,

Cadotte, M.W., Chaneton, E.J., Chu, C., Cleland, E.E., D'Antonio, C., Davies, K.F., Gruner,
635   D.S., Hagenah, N., Kirkman, K., Knops, J.M.H., La Pierre, K.J., McCulley, R.L., Moore, J.L.,
636   Morgan, J.W., Prober, S.M., Risch, A.C., Schuetz, M., Stevens, C.J., Wragg, P.D., 2016.
637   Addition of multiple limiting resources reduces grassland diversity. Nature 537, 93–96.
638   doi:10.1038/nature19324

639 Harrison, X.A., Donaldson, L., Correa-Cano, M.E., Evans, J., Fisher, D.N., Goodwin, C.E.D.,
640   Robinson, B.S., Hodgson, D.J., Inger, R., 2018. A brief introduction to mixed effects modelling
641   and multi-model inference in ecology. PeerJ 2018, 1–32. doi:10.7717/peerj.4794

642 Hättenschwiler, S., Tiunov, A. V., Scheu, S., 2005. Biodiversity and litter decomposition in terrestrial
643   ecosystems. Annual Review of Ecology, Evolution, and Systematics 36, 191–218.
644   doi:10.1146/annurev.ecolsys.36.112904.151932

645 Högberg, M.N., Högberg, P., Myrold, D.D., 2007. Is microbial community composition in boreal
646   forest soils determined by pH, C-to-N ratio, the trees, or all three? Oecologia 150, 590–601.
647   doi:10.1007/s00442-006-0562-5

648 Högberg, P., Read, D.J., 2006. Towards a more plant physiological perspective on soil ecology.
649   Trends in Ecology and Evolution 21, 548–554. doi:10.1016/j.tree.2006.06.004

650 Hu, J., Angeli, S., Schuetz, S., Luo, Y., Hajek, A.E., 2009. Ecology and management of exotic and
651   endemic Asian longhorned beetle Anoplophora glabripennis. Agricultural and Forest
652   Entomology 11, 359–375. doi:10.1111/j.1461-9563.2009.00443.x

653 Johnson, J.B., Omland, K.S., 2004. Model selection in ecology and evolution. Trends in Ecology
654   and Evolution 19, 101–108. doi:10.1016/j.tree.2003.10.013

655 Jones, G.L., Scullion, J., Allison, G., Stott, H., Johnson, D., Owen, R., Williams, G., Gwynn-Jones,
656   D., 2019. Reduced soil respiration beneath invasive Rhododendron ponticum persists after
657   cutting and is related to substrate quality rather than microbial community. Soil Biology and
658   Biochemistry 138. doi:10.1016/j.soilbio.2019.107577

659 Kaiser, C., Koranda, M., Kitzler, B., Fuchslueger, L., Schnecker, J., Schweiger, P., Rasche, F.,
660   Zechmeister-Boltenstern, S., Sessitsch, A., Richter, A., 2010. Belowground carbon allocation
661   by trees drives seasonal patterns of extracellular enzyme activities by altering microbial
662   community composition in a beech forest soil. New Phytologist 187, 843–858.
663   doi:10.1111/j.1469-8137.2010.03321.x

664 Kawaletz, H., Mölder, I., Zerbe, S., Annighöfer, P., Terwei, A., Ammer, C., 2013. Exotic tree
665   seedlings are much more competitive than natives but show underyielding when growing
666   together. Journal of Plant Ecology 6, 305–315. doi:10.1093/jpe/rts044

667 Kieft, T.L., Ringelberg, D.B., White, D.C., 1994. Changes in ester-linked phospholipid fatty acid
668   profiles of subsurface bacteria during starvation and desiccation in a porous medium. Applied
669   and Environmental Microbiology 60, 3292–3299. doi:10.1128/aem.60.9.3292-3299.1994

670 Knoke, T., Ammer, C., Stimm, B., Mosandl, R., 2008. Admixing broadleaved to coniferous tree
671   species: A review on yield, ecological stability and economics. European Journal of Forest
672   Research 127, 89–101. doi:10.1007/s10342-007-0186-2

673 Koranda, M., Kaiser, C., Fuchslueger, L., Kitzler, B., Sessitsch, A., Zechmeister-Boltenstern, S.,
674   Richter, A., 2013. Seasonal variation in functional properties of microbial communities in beech
675   forest soil. Soil Biology and Biochemistry 60, 95–104. doi:10.1016/j.soilbio.2013.01.025

Kramer, C., Gleixner, G., 2008. Soil organic matter in soil depth profiles: Distinct carbon preferences

of microbial groups during carbon transformation. Soil Biology and Biochemistry 40, 425–433.
doi:10.1016/j.soilbio.2007.09.016

Laliberté, E., Kardol, P., Didham, R.K., Teste, F.P., Turner, B.L., Wardle, D.A., 2017. Soil fertility
shapes belowground food webs across a regional climate gradient. Ecology Letters.
doi:10.1111/ele.12823

Langerhans, R.B., Gifford, M.E., Joseph, E.O., 2007. Ecological speciation in Gambusia fishes.
Evolution 61, 2056–2074. doi:10.1111/j.1558-5646.2007.00171.x

Lenth, R., 2020. Emmeans: estimated marginal means, aka Least-Squares Means. R Package
Version 1.4.4. https://CRAN.R-project.org/package=emmeans

Leuschner, C., Ellenberg, H., Leuschner, C., Ellenberg, H., 2017. Beech and mixed beech forests,
in: Ecology of Central European Forests. Springer, pp. 351–441. doi:10.1007/978-3-319-
43042-3_5

Leuschner, C., Hertel, D., Schmid, I., Koch, O., Muhs, A., Hölscher, D., 2004. Stand fine root
biomass and fine root morphology in old-growth beech forests as a function of precipitation and
soil fertility. Plant and Soil 258, 43–56. doi:10.1023/B:PLSO.0000016508.20173.80

Lindahl, B.D., Ihrmark, K., Boberg, J., Trumbore, S.E., Högberg, P., Stenlid, J., Finlay, R.D., 2007.
Spatial separation of litter decomposition and mycorrhizal nitrogen uptake in a boreal forest.
New Phytologist 173, 611–620. doi:10.1111/j.1469-8137.2006.01936.x

Lundquist, E.J., Scow, K.M., Jackson, L.E., Uesugi, S.L., Johnson, C.R., 1999. Rapid response of
soil microbial communities from conventional, low input, and organic farming systems to a
wet/dry cycle. Soil Biology and Biochemistry 31, 1661–1675. doi:10.1016/S0038-
0717(99)00080-2

Ma, Z., Chen, H.Y.H., 2016. Effects of species diversity on fine root productivity in diverse
ecosystems: a global meta-analysis. Global Ecology and Biogeography 25, 1387–1396.
doi:10.1111/geb.12488

MacArthur, R.H., 1957. On the relative abundance of bird species. Proceedings of the National
Academy of Sciences 43, 293–295. doi:10.1073/pnas.43.3.293

Macfadyen, A., 1961. Improved Funnel-Type Extractors for Soil Arthropods. Journal of Animal
Ecology 30, 171. doi:10.2307/2120

Maraun, M., Scheu, S., 1995. Influence of beech litter fragmentation and glucose concentration on
the microbial biomass in three different litter layers of a beechwood. Biology and Fertility of
Soils 19, 155–158. doi:10.1007/BF00336152

Meier, I.C., Tückmantel, T., Heitkötter, J., Müller, K., Preusser, S., Wrobel, T.J., Kandeler, E.,
Marschner, B., Leuschner, C., 2020. Root exudation of mature beech forests across a nutrient
availability gradient: the role of root morphology and fungal activity. New Phytologist 226, 583–
594. doi:10.1111/nph.16389

Mitchell, R.J., Campbell, C.D., Chapman, S.J., Cameron, C.M., 2010. The ecological engineering
impact of a single tree species on the soil microbial community. Journal of Ecology 98, 50–61.
doi:10.1111/j.1365-2745.2009.01601.x

Moore-Kucera, J., Dick, R.P., 2008. PLFA profiling of microbial community structure and seasonal
shifts in soils of a Douglas-fir chronosequence. Microbial Ecology 55, 500–511.
doi:10.1007/s00248-007-9295-1

Nacke, H., Goldmann, K., Schöning, I., Pfeiffer, B., Kaiser, K., Villamizar, G.A.C., Schrumpf, M.,
Buscot, F., Daniel, R., Wubet, T., 2016. Fine spatial scale variation of soil microbial
communities under European beech and Norway spruce. Frontiers in Microbiology 7.
doi:10.3389/fmicb.2016.02067

Oksanen, J., Blanchet, F.G., Michael, F., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R.,
O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., Friendly,
M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos,
P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2019. vegan: Community Ecology Package. R
Package Version 2.5-6. https://CRAN.R-project.org/package=vegan

Pettit, J., Voelker, S., DeRose, R.J., Burton, J., 2020. Spruce beetle outbreak was not driven by
drought stress: evidence from a tree-ring iso-demographic approach indicate temperatures
were more important. Global Change Biology. doi:10.1111/gcb.15274

Peuke, A.D., Gessler, A., Rennenberg, H., 2006. The effect of drought on C and N stable isotopes
in different fractions of leaves, stems and roots of sensitive and tolerant beech ecotypes. Plant,
Cell and Environment 29, 823–835. doi:10.1111/j.1365-3040.2005.01452.x

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., R Core Team, 2019. nlme: Linear and Nonlinear
Mixed Effects Models. R Package Version 3.1-142. https://CRAN.R-project.org/package=nlme

Piovia-Scott, J., Yang, L.H., Wright, A.N., Spiller, D.A., Schoener, T.W., 2019. Pulsed seaweed
subsidies drive sequential shifts in the effects of lizard predators on island food webs. Ecology
Letters 22, 1850–1859. doi:10.1111/ele.13377

Pollierer, M.M., Ferlian, O., Scheu, S., 2015. Temporal dynamics and variation with forest type of
phospholipid fatty acids in litter and soil of temperate forests across regions. Soil Biology and
Biochemistry 91, 248–257. doi:10.1016/j.soilbio.2015.08.035

Pollierer, M.M., Langel, R., Körner, C., Maraun, M., Scheu, S., 2007. The underestimated
importance of belowground carbon input for forest soil animal food webs. Ecology Letters 10,
729–736. doi:10.1111/j.1461-0248.2007.01064.x

Potthast, K., Hamer, U., Makeschin, F., 2010. Impact of litter quality on mineralization processes in
managed and abandoned pasture soils in Southern Ecuador. Soil Biology and Biochemistry 42,
56–64. doi:10.1016/j.soilbio.2009.09.025

Prescott, C.E., Grayston, S.J., Helmisaari, H.S., Kaštovská, E., Körner, C., Lambers, H., Meier, I.C.,
Millard, P., Ostonen, I., 2020. Surplus Carbon Drives Allocation and Plant–Soil Interactions.
Trends in Ecology and Evolution. doi:10.1016/j.tree.2020.08.007

Pretzsch, H., Knoke, T., Paul, C., Bauhus, J., Forrester, D.I., 2017. Perspectives for future research
on mixed-species systems, in: Mixed-Species Forests: Ecology and Management. Springer,
pp. 579–606. doi:10.1007/978-3-662-54553-9_12

R Core Team, 2019. R: A Language and Environment for Statistical Computing. R Fundation for
Statistical Computing, Vienna, Austria. Version 3.6.2. https://www.R-project.org

Ramsey, P.W., Rillig, M.C., Feris, K.P., Holben, W.E., Gannon, J.E., 2006. Choice of methods for
soil microbial community analysis: PLFA maximizes power compared to CLPP and PCR-based
approaches. Pedobiologia 50, 275–280. doi:10.1016/j.pedobi.2006.03.003

Ratcliffe, S., Wirth, C., Jucker, T., van der Plas, F., Scherer-Lorenzen, M., Verheyen, K., Allan, E.,
Benavides, R., Bruelheide, H., Ohse, B., Paquette, A., Ampoorter, E., Bastias, C.C., Bauhus,
J., Bonal, D., Bouriaud, O., Bussotti, F., Carnol, M., Castagneyrol, B., Checko, E., Dawud,

S.M., Wandeler, H. De, Domisch, T., Finér, L., Fischer, M., Fotelli, M., Gessler, A., Granier, A.,
Grossiord, C., Guyot, V., Haase, J., Hättenschwiler, S., Jactel, H., Jaroszewicz, B., Joly, F.X.,
Kambach, S., Kolb, S., Koricheva, J., Liebersgesell, M., Milligan, H., Müller, S., Muys, B.,
Nguyen, D., Nock, C., Pollastrini, M., Purschke, O., Radoglou, K., Raulund-Rasmussen, K.,
Roger, F., Ruiz-Benito, P., Seidl, R., Selvi, F., Seiferling, I., Stenlid, J., Valladares, F.,
Vesterdal, L., Baeten, L., 2017. Biodiversity and ecosystem functioning relations in European
forests depend on environmental context. Ecology Letters. doi:10.1111/ele.12849

Richter, A., Schöning, I., Kahl, T., Bauhus, J., Ruess, L., 2018. Regional environmental conditions
shape microbial community structure stronger than local forest management intensity. Forest
Ecology and Management 409, 250–259. doi:10.1016/j.foreco.2017.11.027

RStudio Team, 2019. RStudio: Integrated Development Environment for R. Rstudio, Inc., Boston,
MA. V.1.2.5033. https://www.rstudio.com

Scheibe, A., Steffens, C., Seven, J., Jacob, A., Hertel, D., Leuschner, C., Gleixner, G., 2015. Effects
of tree identity dominate over tree diversity on the soil microbial community structure. Soil
Biology and Biochemistry 81, 219–227. doi:10.1016/j.soilbio.2014.11.020

Scheu, S., 1992. Automated measurement of the respiratory response of soil microcompartments:
Active microbial biomass in earthworm faeces. Soil Biology and Biochemistry 24, 1113–1118.
doi:10.1016/0038-0717(92)90061-2

Scheu, S., Albers, D., Alphei, J., Buryn, R., Klages, U., Migge, S., Platner, C., Salamon, J.-A., 2003.
The soil fauna community in pure and mixed stands of beech and spruce of different age:
trophic structure and structuring forces. Oikos 101, 225–238. doi:DOI 10.1034/j.1600-
0706.2003.12131.x

Schmid, I., 2002. The influence of soil type and interspecific competition on the fine root system of
Norway spruce and European beech. Basic and Applied Ecology 3, 339–346.
doi:10.1078/1439-1791-00116

Schmid, M., Pautasso, M., Holdenrieder, O., 2014. Ecological consequences of Douglas fir
(Pseudotsuga menziesii) cultivation in Europe. European Journal of Forest Research 133, 13–
29. doi:10.1007/s10342-013-0745-7

Schmidt, M.W.I., Torn, M.S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I.A., Kleber, M.,
Kögel-Knabner, I., Lehmann, J., Manning, D.A.C., Nannipieri, P., Rasse, D.P., Weiner, S.,
Trumbore, S.E., 2011. Persistence of soil organic matter as an ecosystem property. Nature.
doi:10.1038/nature10386

Toïgo, M., Vallet, P., Perot, T., Bontemps, J.D., Piedallu, C., Courbaud, B., 2015. Overyielding in
mixed forests decreases with site productivity. Journal of Ecology 103, 502–512.
doi:10.1111/1365-2745.12353

van der Heijden, M.G.A., Bardgett, R.D., Van Straalen, N.M., 2008. The unseen majority: Soil
microbes as drivers of plant diversity and productivity in terrestrial ecosystems. Ecology Letters
11, 296–310. doi:10.1111/j.1461-0248.2007.01139.x

Waller, L.P., Allen, W.J., Barratt, B.I.P., Condron, L.M., França, F.M., Hunt, J.E., Koele, N., Orwin,
K.H., Steel, G.S., Tylianakis, J.M., Wakelin, S.A., Dickie, I.A., 2020. Biotic interactions drive
ecosystem responses to exotic plant invaders. Science 368, 967–972.
doi:10.1126/science.aba2225

Wardle, D.A., Bardgett, R.D., Klironomos, J.N., Setälä, H., Van Der Putten, W.H., Wall, D.H., 2004.

Ecological linkages between aboveground and belowground biota. Science.
doi:10.1126/science.1094875

Yan, B., Ji, Z., Fan, B., Wang, X., He, G., Shi, L., Liu, G., 2016. Plants adapted to nutrient limitation
allocate less biomass into stems in an arid-hot grassland. New Phytologist 211, 1232–1240.
doi:10.1111/nph.13970

Zelles, L., 1999. Fatty acid patterns of phospholipids and lipopolysaccharides in the characterisation
of microbial communities in soil: A review. Biology and Fertility of Soils 29, 111–129.
doi:10.1007/s003740050533

Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A., Smith, G.M., 2009. Dealing with heterogeneity,
in: Mixed Effects Models and Extensions in Ecology with R. Springer Science & Business
Media, pp. 71–100. doi:10.1007/978-0-387-87458-6_4

Figures and Tables

Figure 1. Effects of conifers and conifer-beech mixtures on microbial basal respiration (μg O_2 g^{-1} C h^{-1}), microbial biomass (μg C_{mic} g^{-1} C) and microbial specific respiration (μg O_2 μg^{-1} C_{mic} h^{-1}) at nutrient-poor and nutrient-rich sites (Douglas-fir [Do], Douglas-fir with beech [Do/Be], Norway spruce [Sp] and Norway spruce with beech [Sp/Be]). Effect sizes are given as back transformed log response ratios compared to beech forests [ln (value in coniferous or mixed forest types / values in beech)]. Effect sizes were averaged across three depths (litter, 0–5 and 5–10 cm soil). Values smaller than 1 indicate higher values in beech. Asterisks indicate significant effects (p < 0.05). Bars represent 95% confidence intervals (n=12).

Figure 2. Effects of conifers and conifer-beech mixtures on stress indicators: ratio of cyclopropyl PLFAs to its monoenoic precursors, ratio of saturated to monounsaturated PLFAs, and ratio of Gram$^+$ to Gram$^-$ bacteria (cy/pre, sat/mono and Gram$^+$/Gram$^-$, respectively) at nutrient-poor and nutrient-rich sites (Douglas-fir [Do], Douglas-fir with beech [Do/Be], Norway spruce [Sp] and Norway spruce with beech [Sp/Be]). Effect sizes are given as back transformed log response ratios compared to beech forests [ln (value in coniferous or mixed forest types / values in beech)]. Effect sizes were averaged across three depths (litter, 0–5 and 5–10 cm soil). Values larger than 1 indicate lower values in beech. Asterisks indicate significant effects (p < 0.05). Bars represent 95% confidence intervals (n=12).

Figure 3. Changes in microbial basal respiration (μg O_2 g^{-1} C h^{-1}), microbial biomass (μg C_{mic} g^{-1} C) and microbial specific respiration (μg O_2 μg^{-1} C_{mic} h^{-1}) with soil depth (litter and 0–5 and 5–10 cm soil) in five forest types (European beech [Be], Douglas-fir [Do], Douglas-fir with beech [Do/Be], Norway spruce [Sp] and Norway spruce with beech [Sp/Be]) at nutrient-poor and nutrient-rich sites. Points and horizontal bars represent means and standard errors (n=4). The grey vertical bars represent respective values in beech forests at nutrient-poor sites in litter, 0–5 and 5–10 cm soil depth. Note log scale for microbial biomass.

Figure 4. Changes in stress indicators (ratio of cyclopropyl fatty acids to its precursors, ratio of saturated to monounsaturated fatty acids and ratio of Gram$^+$ to Gram$^-$ bacteria [cy/pre, sat/mono and Gram$^+$/Gram$^-$, respectively]) with soil depth (litter and 0–5 and 5–10 cm soil) in five forest types (European beech [Be], Douglas-fir [Do], Douglas-fir with beech [Do/Be], Norway spruce [Sp] and Norway spruce with beech [Sp/Be]) at nutrient-poor and nutrient-rich sites. Points and horizontal bars represent means and standard errors (n=4). The grey vertical bars represent respective values in beech forests at nutrient-poor sites in litter, 0–5 and 5–10 cm depth.

Figure 5. Principal component analysis of the microbial community structure (as indicated by phospholipid-derived fatty acids) in five forest types (European beech [Be], Douglas-fir [Do], Douglas-fir with beech [Do/Be], Norway spruce [Sp] and Norway spruce with beech [Sp/Be]) at nutrient-poor and nutrient-rich sites in litter, 0–5 and 5–10 cm soil depth. Positions of forest types represent centroids (n=4). Nutrient-poor sites are given in red and nutrient-rich sites in green. C/N, pH, water content (water%), carbon content (C%) as well as litter mass were post-fitted into the plot. Significant variables selected by RDA permutation test were in bold (based adjusted R^2).

856   Figure 1

859   Figure 2

Figure 3

Figure 4

Figure 5

Table 1. General information as well as physical and chemical properties of the soil of the study
sites. Soil texture is based on 5–30 cm mineral soil, and soil chemical properties are based on 0–5
cm mineral soil. Soil moisture is expressed as percentage of dry mass, and was measured in litter,
0–5 and 5–10 cm soil. Ca^{2+} concentration was analyzed in exchangeable form by NH_4Cl extraction,
and total P was determined by pressure digestion (for detailed see Methods and Foltran et al.
2020).

	Site condition	Rich sites				Poor sites			
	Site number	1	2	3	4	5	6	7	8
General information	Site name	Harz	Dassel	Winnefeld	Nienover	Nienburg	Unterlüß	Göhrde II	Göhrde I
	Latitude (°N)	51.770	51.726	51.662	51.696	52.621	52.837	53.127	53.201
	Longitude (°E)	10.397	9.704	9.574	9.526	9.281	10.331	10.814	10.801
	MAT (°C)	7.63	8.59	8.90	9.01	9.70	9.03	9.19	9.20
	MAP (mm)	1029.24	821.29	822.43	874.97	733.34	746.56	681.68	672.63
	Slope (°)	15.52	2.96	4.97	6.56	1.85	1.19	1.43	2.05
	Elevation (m)	510.80	426.17	349.06	323.37	92.00	161.20	129.80	120.00
Texture & moisture	Sand%	16	26	20	20	80	79	79	73
	Silt%	16	53	57	57	13	15	15	24
	Clay%	68	21	23	23	7	6	6	3
	Water% (Litter)	237.2	221.2	235.5	232.4	245.9	303.1	328.5	265.7
	Water% (0–5)	135.5	106.6	84.7	62.7	71.4	120.2	88.1	76.2
	Water% (5–10)	59.1	39.6	37.8	29.6	22.4	26.9	23.9	22.9
Chemical properties	C %	7.09	6.19	7.50	5.30	7.97	4.03	9.31	5.36
	N %	0.37	0.34	0.37	0.29	0.28	0.17	0.34	0.24
	C/N	19.16	18.21	20.27	18.28	28.46	23.71	27.38	22.33
	CEC (mmol dm^{-3})	237.31	121.26	114.91	95.96	74.81	47.79	71.10	67.72
	pH (KCl)	3.42	3.15	3.34	3.47	3.00	3.00	2.71	3.01
	P (mg kg^{-1})	0.61	0.54	0.44	0.46	0.16	0.11	0.17	0.24
	Ca^{2+} (CEC; mmol dm^{-3})	28.35	13.77	32.56	27.88	20.38	12.91	7.90	11.28

MAT: mean annual temperature
MAP: mean annual precipitation
CEC: cation exchange capacity

Table 2. F- and P-values of linear mixed effects models on the effect of Forest type (European
beech, Douglas-fir, Norway spruce, mixture of European beech with Douglas-fir and mixture of
European beech with Norway spruce), Site condition (nutrient-poor and nutrient-rich sites) and
Depth (litter, 0–5 and 5–10 cm) on (a) basal respiration, microbial biomass and microbial specific
respiration, (b) stress indicators (cy/pre, sat/mono, Gram$^+$/Gram$^-$ ratios), and (c) percentages of
fungal and bacterial PLFAs (of total PLFAs), and fungi/bacteria ratio. Significant effects are given in
bold (p < 0.05).

(a)	Basal respiration			Microbial biomass			Specific respiration		
Factor	df	F	P	df	F	P	df	F	P
Forest type (F)	4,24	8.94	**<0.001**	4,30	5.95	**0.001**	4,24	0.92	0.470
Site condition (S)	1,6	3.90	0.096	1,30	16.93	**<0.001**	1,6	0.52	0.498
Depth (D)	2,78	168.37	**<0.001**	2,70	271.27	**<0.001**	2,78	17.37	**<0.001**
F × S	4,24	10.91	**<0.001**	4,30	4.61	**0.005**	4,24	2.36	0.082
F × D	-	-	**-**	8,70	2.15	**0.042**	-	-	-

(b)	cy/pre ratio			sat/mono ratio			Gram$^+$/Gram$^-$ ratio		
Factor	df	F	P	df	F	P	df	F	P
Forest type (F)	4,24	2.75	0.052	4,24	8.278	**<0.001**	4,24	3.39	**<0.001**
Site condition (S)	1,6	3.48	0.111	1,6	0.195	0.304	1,6	8.17	**0.028**
Depth (D)	2,68	238.82	**<0.001**	2,76	5.551	**<0.001**	2,76	228.51	**<0.001**
F × S	4,24	2.70	0.055	4,24	3.650	**0.002**	4,24	2.27	**0.012**
S × D	2,68	6.20	**0.003**	2,76	10.705	**<0.001**	2,76	7.51	**0.001**
F × D	8,68	3.72	**0.001**	-	-	-	-	-	-

(c)	% fungi			% bacteria			fungi/bacteria ratio		
Factor	df	F	P	df	F	P	df	F	P
Forest type (F)	4,30	1.77	0.161	4,30	2.19	0.094	4,24	0.83	0.520
Site condition (S)	1,30	6.46	**0.016**	1,30	4.34	**0.046**	1,6	4.98	**0.033**
Depth (D)	2,68	982.11	**<0.001**	2,78	149.46	**<0.001**	2,68	995.40	**<0.001**
F × S	4,30	1.99	0.121	4,30	0.73	0.577	4,24	1.60	0.207
S × D	2,68	18.54	**<0.001**	-	-	-	2,68	16.92	**<0.001**
F × D	8,68	3.61	**0.002**	-	-	-	8,68	2.84	**0.009**

Table 3. Results of two-way MANOVAs on PLFA composition (a) without covariates and (b) with
covariates (pH, C/N ratio, water content) in litter, 0–5 and 5–10 cm soil. Factors include Forest type
(European beech, Douglas-fir, Norway spruce and mixture of European beech with Douglas-fir and
mixture of European beech with Norway spruce), Site condition (nutrient-rich and nutrient-poor
sites) and their interactions. MANOVAs were based on principal components from each sample
depth (significant PCs were determined by broken stick criterion; see Methods). Type I sequential
sum of square was used for models with and without covariates. Significant effects are given in bold
($P < 0.05$).

| Depth | Factor | (a) Without covariates | | | | (b) With covariates | | | |
		df	Wilk's lambda	F	P	df	Wilk's lambda	F	P
						Covariates: pH			
Litter	Forest type (F)	4,30	0.32	3.29	**0.001**	4,29	0.39	2.53	**0.008**
	Site condition (S)	1,30	0.42	12.77	**<0.001**	1,29	0.61	5.72	**0.004**
	F × S	4,30	0.73	0.77	0.677	4,29	0.75	0.68	0.767
						Covariates: C/N + pH + water%			
0–5 cm	Forest type (F)	4,30	0.19	2.80	**<0.001**	4,27	0.30	1.67	0.057
	Site condition (S)	1,30	0.39	8.14	**<0.001**	1,27	0.75	1.54	0.218
	F × S	4,30	0.36	1.57	0.078	4,27	0.33	1.53	0.096
						Covariates: C/N			
5–10 cm	Forest type (F)	4,30	0.41	1.76	0.051	4,29	0.41	1.68	0.067
	Site condition (S)	1,30	0.45	8.15	**<0.001**	1,29	0.82	1.39	0.265
	F × S	4,30	0.69	0.68	0.805	4,29	0.73	0.54	0.915