Running title: Forest type effects on microbial communities

Journal: Soil Biology and Biochemistry (special issue: Plant-soil interactions in forests)

Title: Planting mixed forests of European beech and conifers: Variations in microbial responses with
site conditions

Authors: Jing-Zhong Lu[1*], Stefan Scheu[1,2]

[1] J. F. Blumenbach Institute of Zoology and Anthropology, University of Göttingen, Untere Karspüle
2, 37073 Göttingen, Germany

[2] Center of Biodiversity and Sustainable Land Use, University of Göttingen, Büsgenweg 1, 37077
Göttingen, Germany

[*] Corresponding author (E-mail: jlu@gwdg.de)

Keywords: Douglas-fir, Norway spruce, soil respiration, forest, ecosystem functioning, PLFA

Author contributions: SS and JZL designed the study. JZL collected the data, analyzed the data
and wrote the first draft of the manuscript. SS contributed to all stages of the manuscript.

Abstract

Tree - soil interactions depend on environmental context. Plantations of trees may impact soil microorganisms more strongly under unfavorable environmental conditions, compromising long-term ecosystem services. To contextually understand the effects of tree species composition on soil microorganisms, we quantified structural and functional responses of soil microorganisms to forest types across environmental gradients using substrate-induced respiration and phospholipid fatty acid analyses. Five forest types were studied including pure stands of native European beech (*Fagus sylvatica*), range expanding Norway spruce (*Picea abies*), and non-native Douglas-fir (*Pseudotsuga menziesii*), as well as the two conifer - beech mixtures. We found that microbial functioning strongly depends on environmental conditions, in particular on soil nutrients. At nutrient-poor sites, both pure and mixed coniferous forests, but especially Douglas-fir forests, stressed soil microorganisms compared to beech forests. By contrast, microbial structure and functional indicators in beech forests varied little with site conditions, likely because beech provided high amounts of root-derived resources for microbial growth. The results indicate that, at nutrient-poor sites, long-term effects of planting exotic Douglas-fir on ecosystem functioning need further attention, but planting Douglas-fir at nutrient-rich sites may be of little concern from the perspective of microbial communities. Overall, the results point to the importance of root-derived resources in determining the structure and functioning of soil microbial communities, and document the sensitivity of soil microorganisms to planting tree species that may differ in the provisioning of these resources.

1. Introduction

Soil microbes are the unseen majority of organisms in terrestrial ecosystems (van der Heijden et al., 2008). Trees affect soil microorganisms through several pathways, most importantly by litter and root exudates (Wardle et al., 2004; Högberg and Read, 2006). These resources are likely to shape microbial community composition as dissimilar carbon resources favor different guilds of microorganisms (Lindahl et al., 2007; Fanin et al., 2019). Despite that microbial community composition may change in response to variations in tree species composition, the relative role of litter and root exudates in structuring soil microbial communities remains controversial (Pollierer et al., 2007; Bluhm et al., 2019). The typically thicker organic layers in coniferous compared to deciduous forests have been attributed to the recalcitrance of needles (Augusto et al., 2015), but root and mycorrhizal fungi also affect soil carbon storage (Berger and Berger, 2012; Clemmensen et al., 2013; Averill et al., 2014). Studying the linkage between trees and soil microorganisms may allow uncovering the pathways by which trees drive microbial community composition.

Planting conifers beyond their native range has become common worldwide (Castro-Díez et al., 2019). Although conifers provide timber and other benefits to humans, they may also impact carbon and nutrient cycling, processes largely mediated by soil microorganisms (Bardgett and van der Putten, 2014; Castro-Díez et al., 2019). Admixing conifers to deciduous forests has been suggested to improve carbon and nutrient cycling via changing the functioning of microbial communities (Knoke et al., 2008; Cremer et al., 2016). Higher resource availability in litter mixtures may facilitate litter decomposition (Hättenschwiler et al., 2005). Further, broadleaf mixed forests are likely to increase both aboveground and belowground biomass due to interspecific facilitation and improved resource partitioning, resulting in higher resource availability to microorganisms (Cardinale et al., 2007; Emmett Duffy et al., 2017). The importance of biodiversity for ecosystem functioning is increasingly recognized, particularly under unfavorable environmental conditions (Wright et al., 2020). Overall, however, it is poorly understood how the composition of tree species affects soil microbial communities, especially across environmental gradients (Malchair and Carnol, 2009).

Tree - soil interactions depend on environmental context (Ratcliffe et al., 2017), but to what extent effects of tree species composition on soil microorganisms vary with site conditions, such as nutrient status and water availability, is little known. Facilitation among plants is more pronounced at unfavorable conditions according to the stress-gradient hypothesis, and such positive interspecific interactions are likely to be mediated by soil microorganisms (Defossez et al., 2011; David et al., 2020). In fact, plants do not passively tolerate environmental stress, but respond in various ways to unfavorable growth conditions, such as by shifting the allocation of resources towards roots under nutrient-poor conditions (Callaway et al., 2003; Augusto et al., 2015). Recently, de Vries et al. (2019) has shown that under drought conditions herbaceous species change the quality of root exudates

and induce higher microbial respiration, presumably facilitating nutrient capture by stimulating

microorganisms. Phenotypic plasticity or morphological changes induced by the environment may

differ between species, exacerbating differences in resource availability to soil microorganisms

under stress conditions (Meier and Leuschner, 2008; Schall et al., 2012). Thus, the impact of tree

species composition on microbial community composition and functioning may be more pronounced

under unfavorable than favorable environmental conditions.

A better understanding of the effects of tree species composition on soil microorganisms has

implications for forest management, especially in temperate and boreal regions where tree species

richness is low and managed forests are dominated by one or a few species (Knoke et al., 2008;

Bauhus et al., 2010). European beech is the climax species in lowland and lower montane regions

in Central Europe (Leuschner et al., 2017). The most popular timber species, Norway spruce

originally occurred in higher mountain ranges and boreal regions, but has been planted widely in

lowlands (Knoke et al., 2008). In recent years, Norway spruce has been damaged severely by

extreme weather and bark beetle outbreaks, events predicted to become more frequent in the future

(Biedermann et al., 2019; Pettit et al., 2020). Although the admixture of Norway spruce to native

European beech forests may reduce the risk of damage while maintaining economic gains,

Douglas-fir is increasingly planted (Schmid et al., 2014; Ammer, 2019). Since its introduction from

North America over 150 years ago, Douglas-fir has become the most abundant non-native

cultivated tree species in Central Europe (Schmid et al., 2014). To date, the impact of planting

Douglas-fir on biodiversity and functioning of forests is little studied, and this applies in particular to

the belowground system, although e.g., Douglas-fir has been suggested to affect soil chemistry in a

similar way than spruce (Schmid et al., 2014). Overall, we lack a comprehensive evaluation of forest

types on soil microbial communities across environmental gradients.

Here, we studied microbial community composition and functioning in litter and soil of five forest

types under different environmental conditions, in particular soil nutrients. Forest types included

pure stands of European beech (*Fagus sylvatica*), Norway spruce (*Picea abies*), Douglas-fir

(*Pseudotsuga menziesii*), and the two conifer - beech mixtures. We analyzed the structure of

microbial communities using phospholipid fatty acid patterns, and their functioning using microbial

basal respiration, biomass and stress indicators. In general, we assumed the structure of microbial

communities to be closely linked to their functioning. In particular, we hypothesized (1) microbial

structure and functioning to be detrimentally affected by coniferous trees, with the effects being

similar in Norway spruce and Douglas-fir forests, and intermediate in mixed beech - conifer forests.

Further, we hypothesized (2) microbial community structure and functioning to be strongly impacted

by forest type at nutrient-poor sites, but less at nutrient-rich sites. Since the litter layer is less

buffered against environmental harshness than the organic and mineral soil, we also hypothesized

that (3) the effects of forest types would be more pronounced in litter than in soil.

2. Methods

2.1. Field sites

The study included five forest types arranged as quintets at eight sites in northern Germany, covering a range of environmental conditions (5 forest types x 8 sites; Fig. S1). The four sites in the south stock on fertile soil and receive higher precipitation. Mean annual precipitation is 821 – 1029 mm (Ammer et al., 2020). Parental rock is either loess influenced Triassic sandstone or Paleozoic shares of greywacke, sandstone, quartzite and phyllite, resulting in soil types of partly podsolic Cambisol and Luvisol. The four sites in the north are located on out-washed sand with the soil type of Podzol. The mean annual precipitation is 672 – 746 mm. The southern sites are richer in nutrients than the northern sites as reflected by higher total soil P, cation exchange capacity and pH; more details on site characteristics and soil chemical properties are given in Table 1 (Ammer et al., 2020; Foltran et al., 2020). Hereafter, we refer to the southern sites as nutrient-rich sites and to the northern sites as nutrient-poor sites.

Each site comprised pure stands of European beech (*Fagus sylvatica* L.; Be), Douglas-fir (*Pseudotsuga menziesii* [Mirbel] Franco.; Do) and Norway spruce (*Picea abies* [L.] Karst.; Sp), as well as two conifer-beech mixtures (Douglas-fir/European beech and Norway spruce/European beech; Do/Be and Sp/Be). On average, trees were more than 50 years old. Within sites, the distance between stands ranged from 76 m to 4600 m, and the distance between sites ranged from 5 to 190 km. Within each stand, plots of 2500 m^2 were established, mostly in rectangular shape (50 × 50 m). Tree species in pure stands comprised more than 90% of the total basal area, and that in mixed stands 33 – 53% for beech and 53 – 60% for conifers.

2.2. Soil sampling and physio-chemical analyses

Samples were taken between November 2017 and January 2018. Soil cores of 5 cm diameter were taken by a metal cylinder, and were separated into litter, 0 – 5 and 5 – 10 cm soil depth. In each plot, three cores spaced by 5 m were taken. Samples from the same depth were pooled, resulting in 120 samples (40 plots × 3 depths). The soil was sieved through 2 mm mesh, and the litter was cut into pieces (< 25 mm^2) (Maraun and Scheu, 1995). Roots > 2 mm in diameter and stones were removed. Samples were stored at -20°C. The pH was determined using a ratio of sample to solution (g/ml; KCl, 1 M) of 1:10 for litter, 1:5 and 1:2.5 for 0 – 5 and 5 – 10 cm soil, respectively. Moisture content was measured by drying samples at 105°C for 48 h. Water content did not differ consistently between nutrient-rich and nutrient-poor sites across layers; it was higher in litter but lower in soil at nutrient-poor than at nutrient-rich sites (Site condition × Depth interaction; $F_{2,68} = 26.21$, $P < 0.001$). Dried samples were grinded in a ball mill, and total carbon and nitrogen concentrations were determined by an elemental analyzer (NA 1110, CE-instruments, Rodano, Milano, Italy). Litter mass was estimated by drying at 50°C for > 48 h (Macfadyen, 1961).

2.3. Microbial basal respiration and biomass

Microbial biomass was measured by substrate-induced respiration with respiration measured as O_2 consumption (Anderson and Domsch, 1978; Scheu, 1992). Samples stored at -20°C were thawed and incubated overnight at room temperature (20°C) before placing in an automated micro-respirometer. The system absorbs CO_2 using KOH solution and the O_2 consumed by microorganisms is replaced through electrolytic release of O_2 from $CuSO_4$ solution. Microbial basal respiration was measured as mean consumption of O_2 during 10 – 23 h after attachment of the vessels to the respirometer ($\mu g\ O_2\ g^{-1}\ h^{-1}$). Microbial biomass was determined after the addition of glucose to saturate microbial glycolytic enzymes based on the maximum initial respiratory response 4 – 7 h after the addition of D-glucose (*MIRR*; $\mu l\ O_2\ g^{-1}\ h^{-1}$) and converted to microbial biomass (C_{mic}; $\mu g\ C_{mic}\ g^{-1}$) as 38 × *MIRR* (Beck et al., 1997). Basal respiration and substrate-induced respiration were measured at 22°C in a water bath. Fresh litter and soil from 0 – 5 and 5 – 10 cm depth equivalent to approximately 0.3, 1.0 and 3.5 g dry mass was supplemented with 80, 20 and 8 mg glucose solved in 400 $\mu l\ H_2O$, respectively. The ratio between microbial basal respiration and microbial biomass was taken as microbial specific respiration (qO_2; $\mu g\ O_2\ \mu g^{-1}\ C_{mic}\ h^{-1}$).

2.4. Phospholipid fatty acid analysis

To quantify the composition of phospholipid fatty acid (PLFA), lipids were extracted using a modified Bligh and Dyer method (Frostegård et al., 1993; Pollierer et al., 2015). In short, lipids were fractionated into neutral lipids, glycolipids and phospholipids by elution through silica acid columns using chloroform, acetone and methanol, respectively (0.5 g silicic acid, 3 ml; HF BOND ELUT-SI, Varian, Inc., Darmstadt, Germany). Phospholipids were subjected to mild alkaline methanolysis and fatty acid methyl esters were identified by chromatographic retention time compared to standards (FAME CRM47885, C11 to C24; BAME 47080-U, C11 to C20; Sigma-Aldrich, Darmstadt, Germany) using a GC-FID Clarus 500 (PerkinElmer Corporation, Norwalk, USA) equipped with an Elite 5 column (30 m × 0.32 mm inner diameter, film thickness 0.25 μm). The temperature program started with 60°C (hold time 1 min) and increased by 30°C per min to 160°C, and then by 3°C per min to 280°C. The injection temperature was 250°C and helium was used as carrier gas. Approximately 2 g of fresh litter and 4 g of fresh soil were used for the extraction.

2.5. Stress indicators and fatty acid markers

The ratio of cyclopropyl PLFAs to their monoenoic precursors (cy/pre; (cy17:0 + cy19:0)/(16:1ω7 + 18:1ω7) and the ratio of saturated to monounsaturated PLFAs (sat/mono; (14:0 + 15:0 + 16:0 + 17:0 + 18:0)/(16:1ω7 + 17:1 + 18:1ω9 + 18:1ω7)) were used as indicators of physiological or nutritional stress (Moore-Kucera and Dick, 2008; Pollierer et al., 2015). The ratio of Gram[+] to Gram[-] bacteria was used as indicator of carbon availability (Fanin et al., 2019). The saturated fatty acids i15:0, a15:0, i16:0, i17:0 were used as markers for Gram[+] bacteria, and the fatty acids cy17:0, cy19:0,

16:1ω7 and 18:1ω7 were assigned as markers for Gram⁻ bacteria (Frostegård and Bååth, 1996;
Zelles, 1999; Fanin et al., 2019). Bacteria were represented by the sum of Gram⁺ and Gram⁻
bacteria. Linoleic acid 18:2ω6,9 was used as fungal marker (Frostegård and Bååth, 1996). Total
amount of PLFAs included all identified PLFAs (n = 40; nmol g⁻¹ dry weight) and was used to
calculate PLFA proportions. All stress indicators and FA markers were analyzed from proportions
(mole percentage).

2.6. Statistical analyses

To estimate the effect size of Forest type and Site condition, we first fitted linear mixed models
(LMMs) to log-transformed response variables and then applied planned contrasts (Piovia-Scott et
al., 2019). All LMMs included Forest type (European beech, Douglas-fir, Douglas-fir/European
beech, Norway spruce, Norway spruce/European beech), Site condition (nutrient-rich and nutrient-
poor sites) and Depth (litter, 0 – 5 and 5 – 10 cm) as fixed effects. Models were stepwise selected
by likelihood ratio test, and minimal models included all main effects and the interaction of Forest
type and Site condition. The 40 forest plots were included as random effects to account for non-
independence of samples from the same plot (Harrison et al., 2018). Based on Akaike Information
Criterion (AIC), the eight sites were included as random effect, and the mean-variance relationship
was accounted for by a dispersion parameter to meet the assumption of homogeneity of variance
(Zuur et al., 2009). Univariate response variables included microbial basal respiration, microbial
biomass, microbial specific respiration, stress indicators and PLFA markers.

Contrasts for the effect size of Forest type were designed to compare coniferous and mixed forests
to beech forests. European beech, the climax tree species in lowland and lower montane regions of
Central Europe, was used as reference (Leuschner et al., 2017). Due to the property of log-
transformed response variables, the planned contrasts are analog to log response ratios (Piovia-
Scott et al., 2019). To improve interpretation, we back transformed the log response ratio into
response ratio and defined it as effect size. Effect sizes of Forest type were estimated for nutrient-
poor and nutrient-rich sites. In addition, we applied contrasts between nutrient-poor vs. nutrient-rich
sites in a similar manner to estimate the effect sizes of Site condition.

For multivariate analysis of variance (MANOVA), PLFAs first were arcsine root transformed and
then dimensions of fatty acids were reduced by principal component analysis (PCA). Separate
MANOVAs for each sample depth were calculated to inspect for effects of Forest type, Site
condition, and their interactions. Further, selected environmental variables were included into
MANOVAs as covariates to test whether they explain main effects in the models (Fry et al., 2016;
Bennett et al., 2020). Significant principal components (PCs) were determined by broken stick
criterion (MacArthur, 1957). Covariates were selected according to permutation tests based on

adjusted R^2 in redundancy analyses (RDA). Only fatty acids ≥0.2% mean mole proportion were

included in the analyses.

All analyses were done in R 4.0.3 (R Core Team, 2019). We used the nlme package to fit LMMs

and the emmeans package to conduct planned contrasts. All mixed models met the assumptions of

normality of residuals and homogeneity of variance.

3. Results

3.1. Functional indicators of microorganisms

Forest type consistently affected microbial basal respiration, microbial biomass and stress indicators

at nutrient-poor but not at nutrient-rich sites (Forest type x Site condition interactions; all $P < 0.08$;

Table 2, Figs 1, 2), and this was generally true across soil layers (Fig. S2). At nutrient-poor sites,

microbial basal respiration and microbial biomass in Douglas-fir forests were 65% and 59% lower

compared to pure beech. Forest type effects were similar in the two conifer-beech mixtures (-46 to -

56%) and least pronounced in pure spruce forests (-33 to -36%; Fig. 1). In addition, at nutrient-poor

sites microbial specific respiration was 19% lower in Douglas-fir forests than in beech forests.

At nutrient-poor sites, stress indicators, the cy/pre ratio, the sat/mono ratio as well as the

Gram$^+$/Gram$^-$ bacteria ratio, all were lower in beech forests than in the other forest types (Fig. 2).

Forest type effects on the cy/pre and sat/mono ratio were more pronounced in Douglas-fir (+87%

and +74%) and less pronounced in spruce and conifer-beech mixtures (+37 to +46%). Effects of

Forest type on the Gram$^+$/Gram$^-$ bacteria ratio also were strongest in Douglas-fir forests (+46%) and

less strong in the other forest types (+21 to +29%). Stress indicators generally did not differ

significantly at nutrient-rich sites ($P > 0.31$; Figs. 2, S2, Table S1).

The response of microorganisms to changes in site conditions also varied with forest types.

Differences between nutrient-rich and nutrient-poor sites were largest in Douglas-fir forests and

smallest in beech forests (Figs 3, 4, Table S2). At nutrient-poor sites, microbial basal respiration in

Douglas-fir forests was 57% lower and microbial biomass in Douglas-fir pure and mixed forests both

were 47% lower compared to nutrient-rich sites. In parallel, but less strong, microbial biomass in

spruce mixed forests was 36% lower at nutrient-poor sites than that at nutrient-rich sites.

Likewise, in Douglas-fir forests the stress indicator cy/pre ratio at nutrient-poor sites exceeded that

at nutrient-rich sites by 82%, and similarly, the sat/mono and Gram$^+$/Gram$^-$ bacteria ratio at nutrient-

poor sites exceeded that at nutrient-rich sites by 38 – 40% (Fig. 4, Table S2). In Douglas-fir mixed

forests the Gram$^+$/Gram$^-$ bacteria ratio at nutrient-poor sites exceeded that at nutrient-rich sites by

34%. Further, the response of microbial stress to changes in site conditions varied with soil depth. In

particular, the responses of the sat/mono ratio and the Gram$^+$/Gram$^-$ ratio to site conditions were

most pronounced at 0 – 5 cm soil depth.

3.2. Microbial community structure

As indicated by the PLFA patterns, microbial community composition in beech forests differed from

that in Norway spruce and Douglas-fir forests, with mixed forests being intermediate between pure

beech and coniferous forests (Fig. 5). Across layers, effects of Forest type on microbial community

composition were significant in litter and 0 – 5 cm soil, and marginally significant in 5 – 10 cm soil

(Table 3). Despite strong turnover of microbial community composition between nutrient-rich and

nutrient-poor sites (Site condition effects; all p < 0.001), in beech forests microbial community

structure in nutrient-poor and nutrient-rich sites was similar, and this was most apparent in 0 – 5 cm

soil (Forest type × Site condition interaction; Fig. 5, Table 3).

A number of fatty acids varied among forest types and with site conditions. Beech forests were

characterized by unsaturated fatty acids at all depths, in particular by 16:1ω7, 18:1ω7, 18:1ω9 (Fig.

5). Norway spruce forests were associated with long-chain saturated fatty acids (22:0, 24:0), and

Douglas-fir forests with bacterial marker fatty acids (2OH-14:0, cy17:0). Further, nutrient-rich sites

were associated with unsaturated fatty acids in litter, such as 18:1ω9 and 18:2ω6,9.

Litter at nutrient-rich sites was generally richer in fungi than at nutrient-poor sites, but the opposite

was true for the soil (Site condition × Depth interaction; Table 2, Fig. S3). The relative abundance of

fungi and bacteria generally did not differ among forest types, but the relative abundance of fungi

consistently decreased, whereas the relative abundance of bacteria increased with soil depth (Figs

S3, S4). In addition, based on the contrast of Site condition, at nutrient-poor sites the relative

abundance of fungi in pure beech and spruce forests was higher than at nutrient-rich sites. By

contrast, in pure and mixed Douglas-fir forests at nutrient-poor sites the relative abundance of fungi

in soil was similar to that at nutrient-rich sites (Table S2, Fig. S3).

Litter mass did not differ significantly among forest types, but was 58% higher at nutrient-poor than

at nutrient-rich sites (Forest type: $F_{4,30}$ = 0.55, P = 0.70; Site condition: $F_{1,30}$ = 13.21, P = 0.001; Fig.

S5). Nutrient-poor sites also were more nitrogen-limited and acidic than nutrient-rich sites (Fig. 5).

Soil C/N ratio, pH and soil moisture were among the most important environmental variables in

explaining variations in PLFA composition (Table S3). In litter, pH accounted for most (8%) of the

variation (13%) explained by all environmental variables. By contrast, in 0 – 5 cm as well as 5 – 10

cm soil depth, C/N ratio accounted for most of the variation explained by all environmental variables

(22% out of the total of 37% and 13% out of the total of 15%, respectively; Table S3). Underlining

the importance of environmental factors in structuring microbial communities at the study sites, the

main effect of Site condition became non-significant in soil after including C/N ratio, pH, water

content (0 – 5 cm soil depth) and C/N ratio (5 – 10 cm soil depth) as covariates in MANOVAs. By

contrast, in litter, including pH as covariate did not significantly affect the effects of Site condition.

The effects of Forest type remained marginally significant in both depths of soil and significant in

litter when including covariates (Fig. 5, Table 3).

284 4. Discussion

The present study evaluated effects of two coniferous trees in pure and mixed forests on soil

microbial community composition and functioning covering different site conditions, with native

European beech as reference. Overall, the results in part support our first and second hypotheses.

Forest type only affected the functioning of microbial communities at nutrient-poor sites, but not at

nutrient-rich sites. At nutrient-poor sites, microbial basal respiration, biomass and stress indicators

all were detrimentally affected in pure and mixed coniferous forests, and, contrasting our first

hypothesis the effects differed between Douglas-fir and spruce forests, with the effects being

particularly strong in Douglas-fir, but less pronounced in Norway spruce and mixed stands.

Enrichment of beech with either Douglas-fir or spruce in mixed forests both compromised microbial

community functioning, suggesting that caution is needed when admixing conifers to European

beech forests at nutrient-poor sites. By contrast, similar basal respiration, microbial biomass and

stress indicators at nutrient-rich sites across the studied forest types contradict our second

hypothesis and suggest that, at favorable environmental conditions, soil microbial communities are

rather irresponsive to changes in tree species. This implies that under favorable site conditions

planting Douglas-fir in pure or mixed forests may provide an alternative to planting Norway spruce.

300 4.1. Functional indicators

The site- and tree species-specific responses of microorganisms are likely caused by differences in

the provisioning of carbon resources. Gram$^+$ bacteria better cope with recalcitrant carbon resources,

whereas Gram$^-$ bacteria favor labile carbon compounds (Kramer and Gleixner, 2008; Fanin et al.,

2019; Waller et al., 2020). High Gram$^+$/Gram$^-$ ratio in Douglas-fir and low Gram$^+$/Gram$^-$ ratio in

European beech at nutrient-poor sites therefore indicate more pronounced shortage of labile carbon

resources in Douglas-fir compared to beech forests. This is further supported by the lower microbial

specific respiration in Douglas-fir compared to beech forests, which was largely due to lower

microbial basal respiration (Anderson and Domsch, 2010). Low availability of labile carbon favors

oligotrophic microbial communities characterized by low respiration rates and biomass (Potthast et

al., 2010; Fanin et al., 2019), and this is consistent with our findings of low basal respiration and

microbial biomass in pure and mixed coniferous forests at nutrient-poor sites.

Moreover, higher ratio of cyclopropyl PLFAs to their monoenoic precursors and higher ratio of

saturated to monounsaturated PLFAs in pure and mixed coniferous forests compared to pure beech

forests at nutrient-poor sites indicates that microorganisms in pure and mixed coniferous forests

were more nutrient (and/or water) stressed (Moore-Kucera and Dick, 2008; Pollierer et al., 2015).

More stressed microorganisms in coniferous and mixed forests compared to pure beech forests at

nutrient-poor sites also is indicated by lower proportions of the presumably Gram[-] bacteria marker

fatty acids 16:1ω7 and 18:1ω7 in the former. Under stress, Gram[-] bacteria change the composition

of their cell membranes from 16:1ω7 and 18:1ω7 to cyclopropane fatty acids associated with slower

growth rates (Guckert et al., 1986; Kieft et al., 1994; Lundquist et al., 1999). The similarity of

microbial indicators in beech forests at nutrient-rich and nutrient-poor sites indicates that, compared

to coniferous stands, beech maintained microbial community functioning and prevented stress

conditions for microorganisms by providing more labile carbon resources at nutrient-poor conditions,

presumably by releasing high amounts of root-derived resources (Meier et al., 2020).

The suggestion that beech is able to alleviate microbial stress via maintaining or even increasing the

release of root-derived resources at nutrient-poor sites is supported by higher fine root biomass in

beech than in pure and mixed coniferous forests at nutrient-poor sites, whereas at nutrient-rich sites

fine roots biomass varies little among forest types (A. Lwila, pers. comm.). If soil nutrient availability

declines, plants allocate surplus carbon into roots, associated with increased root exudation

(Eisenhauer et al., 2017; Laliberté et al., 2017; Prescott et al., 2020). Recently, higher root

exudation has been confirmed for beech forests at more acidic and nitrogen deficient stands (Meier

et al., 2020), conditions similar to our nutrient-poor sites characterized by higher C/N ratio and more

acidic soil. High amounts of root-derived resources presumably also were responsible for the

alleviated microbial resource deficiency and stress in beech forests at nutrient-poor sites. Trenching

and girdling experiments also confirmed that reducing or omitting the flux of root-derived resources

into the soil strongly reduces fungal and bacterial biomass, demonstrating the importance of root-

derived resources for maintaining soil microbial biomass (Högberg et al., 2007; Kaiser et al., 2010;

Bluhm et al., 2019). Thus, beech may facilitate soil microorganisms at environmental stress

conditions by increasing root exudation contrasting both of the studied coniferous species. This may

also mitigate detrimental effects of Douglas-fir on microorganisms at nutrient-poor or acidic sites in

mixed forests. Generally, high amounts of root-derived resources in beech may be responsible for

the facilitation of neighboring coniferous trees by beech, resulting in overyielding in mixed stands at

nutrient-poor sites (Toïgo et al., 2015; Ammer, 2019; David et al., 2020).

Notably, we sampled litter and soil during autumn / winter after beech had shed its leaves, which

might have affected microbial communities. However, any litter effect on microbial communities

should have been similar in nutrient-rich and nutrient-poor sites, as litter input in beech forests

varies little with site conditions (Meier et al., 2005). By contrast, we found microbial community

characteristics to only differ among forest types at nutrient-poor sites, and litter mass on the forest

floor did not differ among forest types (Fig. S5). Supporting our suggestion on the importance of

root-derived resources, at nutrient-poor sites fine root biomass in beech forests exceeded that in
coniferous forests by more than 1.6 times, whereas fine root biomass did not differ significantly
among forest types at nutrient-rich sites (A. Lwila, pers. comm.). The hypothesis of high root-derived
resources in beech forests implies that the impact of roots on microorganisms still was strong in
autumn / winter indicating that not only recent photosynthates but also stored carbohydrates
contribute to providing root-derived resources to soil microorganisms (Druebert et al., 2009).

However, it remains open whether the effects of Forest type on microbial communities get stronger
or weaker with time, hampering generalization of our finding across seasons. Although microbial
communities fluctuate considerably with season (Koranda et al., 2013; Abramoff and Finzi, 2016;
Nacke et al., 2016), comprehensive sampling in Douglas-fir forests across seasons suggests that,
except during summer drought, microbial stress indicators fluctuate little (Moore-Kucera and Dick,
2008). Further, as our sampling was limited to the upper 10 cm of the soil, we may not have
captured the full effects of roots, as e.g., beech roots may extend considerably deeper into the soil
(Leuschner et al., 2004), although fine roots generally are concentrated in upper soil layers (Schmid,
2002; Leuschner et al., 2004). Experimental manipulations of root-derived resources in the field are
needed to fully resolve the role of root-derived resources in driving microbial community structure
and functioning (Scheu, 2002; Kaiser et al., 2010; Bluhm et al., 2019).

Contrary to our third hypothesis, the effects of Forest type were not stronger in litter than in soil.
Forest types impacted microbial respiration, biomass and stress indicators similarly across layers at
nutrient-poor sites. This uniformity further supports our conclusion that Forest type effects were not
the result of different litter quantity and quality, but were due to root-derived resources. This
contrasts earlier suggestions that, because litter is not buffered against environmental conditions,
microbial responses in litter are more sensitive to environmental change than those in soil (Pollierer
et al., 2015). Reflecting environmental changes, microbial biomass decreased while microbial stress
indicators increased with soil depth. This sensitivity may be related to lower carbon availability with
soil depth, resulting in stronger resource limitation in deeper soil (Fierer et al., 2003). As indicated
by Gram$^+$/Gram$^-$ ratios, the difference in resource limitation between litter and soil was greatest in
pure coniferous and mixed forests at nutrient-poor sites.

378 4.2. Community structure

We assumed that the microbial functioning ultimately relies on its community structure, and this is
supported by the consistent turnover in microbial community composition as well as the differences
in microbial functioning between nutrient-rich and poor sites. Despite increased resource limitation
with soil depth, microbial community structure in 5 – 10 cm soil differed little between forest types at
nutrient-rich sites. This suggests that the influence of forest types fades deeper in soil. By contrast,
at nutrient-poor sites, microbial community structure differed between beech and coniferous forests

in soil, particularly at 5 – 10 cm depth, supporting our conclusion that effects of tree species on

microorganisms in nutrient-poor soil are predominantly due to differences in root-derived resources.

Unlike Douglas-fir and spruce, European beech maintained and stabilized microbial community

structure and functioning in litter and soil irrespective of site conditions. Interestingly, at 5 – 10 cm

depth in nutrient-poor sites, microbial community structure and carbon limitation under Douglas-fir

differed most strongly from beech. This implies that, among the tree species studied, root-derived

resources are most limited under Douglas-fir at nutrient-poor sites, resulting in pronounced resource

shortage and marked changes in microbial community composition. Such changes in microbial

community structure may significantly impact the functioning of soil microbial communities as the

availability of carbon resources and carbon limitation affect the vertical distribution of saprotrophic

fungi and decomposition processes, e.g. by shifting the competition between saprotrophic and

mycorrhizal fungi (Gadgil and Gadgil, 1971; Lindahl et al., 2007).

Our results support previous findings that regional factors more strongly shape microbial community

structure than tree species (Pollierer et al., 2015; Richter et al., 2018). The relative abundance of

fungi and bacteria as well as the fungi/bacteria ratio did not vary with forest types but rather with site

conditions. The relative abundance of fungi was favored at nutrient-poor sites, supporting earlier

suggestions that the fungal energy channel dominates in low nutrient systems (Wardle et al., 2004).

However, at nutrient-poor sites, the fungal energy channel did not dominate in Douglas-fir forests,

although they were also characterized as infertile with high C/N ratio and low pH. In fact, the relative

abundance of fungi in Douglas-fir forest soil was similar at nutrient-rich and poor sites, resulting in

lower fungal abundance in Douglas-fir than beech forests, particularly at 5 – 10 cm depth. In

addition to low amounts of root-derived resources, low fungal abundance in Douglas-fir forest soil

may be due to a mismatch between local and home (native range of Douglas-fir) fungal

communities, as only a subset of native-range fungal species has been recorded in the soils to

which Douglas-fir has been introduced (Schmid et al., 2014). Although molecular techniques are

needed to identify the microbial taxa responsible for differences among forest types, particularly

deeper in soil between Douglas-fir and beech at nutrient-poor sites, our study supports the

sensitivity of PLFA analysis in detecting changes in microbial community composition (Ramsey et

al., 2006; Frostegård et al., 2011).

Forest type effects on microbial community composition were marginally significant after including

covariates, indicating that forest types affected microbial community composition independently of

the studied environmental variables. By contrast, the measured environmental variables well

captured differences in soils between nutrient-poor and nutrient-rich sites, and explained much of

the variance in site conditions. Among the environmental variables studied, C/N ratio and pH best

explained variations in microbial community structure. Soil pH typically correlates negatively with

soil nutrient availability (Meier et al., 2020), further supporting the importance of nutrients in driving

microbial responses to variations in site conditions. However, factors other than soil nutrient status

and soil pH may have contributed to the observed differences between nutrient-rich and nutrient-

poor sites. Despite no consistent effects of site conditions on moisture content across layers, and

little effects of moisture on litter and soil microbial community structure, nutrient-poor sites also are

characterized by lower precipitation. This is supported by higher δ^{13}C values of litter at nutrient-poor

sites (J.-Z. Lu, unpubl. data), reflecting higher water stress at nutrient-poor than nutrient-rich sites

(Peuke et al., 2006). Presumably, in addition to nutrients and pH, water stress contributed to the

observed differences in microbial community composition and functioning between nutrient-poor

and nutrient-rich sites. This is supported by the fact that soil water conditions are intricately linked to

soil nutrient dynamics and the uptake of nutrients by plants (de Vries et al., 2019). To disentangle

pathways linking forest stands and soil microbial communities, it is most promising to focus on

nutrient-poor, water deficient or low pH stands.

4.3. Management implications

Due to increasing drought and associated outbreaks of bark beetles, Norway spruce forests are

increasingly threatened in lowland regions of Europe. Facing such challenge, mixed forests and

planting alternative tree species such Douglas-fir may provide better options for provisioning long-

term ecosystem services. We studied microbial communities in response to pure and mixed forests

covering different site conditions, and we found that effects of forest types on soil microorganisms

vary among tree species and mixed stands, but this strongly depends on site conditions, in

particular soil nutrients. At nutrient-poor sites, Douglas-fir strongly impacted microbial community

structure and functioning compared to beech. Therefore, in the long-term planting non-native tree

species such as Douglas-fir may compromise ecosystem functioning at nutrient-poor sites. This may

not only question the establishment of pure Douglas-fir forests, but also the enrichment of beech

forests by Douglas-fir at nutrient-poor sites. However, as planting Douglas-fir in single species and

mixed stands with beech hardly impacted the structure and functioning of soil microbial communities

at nutrient-rich sites, planting Douglas-fir may provide an alternative to Norway spruce at least at

nutrient-rich sites. Although more information on ecosystem properties and processes is needed to

gain a holistic understanding on the consequence of forest plantation on ecosystem functioning, we

conclude that microbial stress is intensified in nutrient-poor soil by planting Douglas-fir.

Acknowledgements

This work was supported by the German Research Foundation through the Research Training

Group 2300: "Enrichment of European beech forests with conifers: impacts of functional traits on

ecosystem functioning" [316045089]. We thank C. Bluhm, T. Volovei, G. Humpert, M. Unger, J.

Meyer and S.-X. Zhou for assistance. We are indebted to C. Ammer, H. Riebl, T. Hartke, A. Polle,

X.-M. Lu, N. Lamersdorf, Z.-P. Li, S. Bluhm, G. Schulz, T.-W. Chen, M. Pollierer, A. Lwila, K.

Schumann, I. Bebre and A. Rivera for helpful discussions. Special thanks to S. Müller for
coordinating the project.

References

Abramoff, R.Z., Finzi, A.C., 2016. Seasonality and partitioning of root allocation to rhizosphere soils in a
midlatitude forest. Ecosphere 7. doi:10.1002/ecs2.1547

Ammer, C., 2019. Diversity and forest productivity in a changing climate. New Phytologist 221, 50–66.
doi:10.1111/nph.15263

Ammer, C., Annighöfer, P., Balkenhol, N., Hertel, D., Leuschner, C., Polle, A., Lamersdorf, N., Scheu, S.,
Glatthorn, J., 2020. RTG 2300 - study design, location, topography and climatic conditions of research
plots in 2020. PANGAEA. doi:doi.pangaea.de/10.1594/PANGAEA.923125

Anderson, J.P.E., Domsch, K.H., 1978. A physiological method for the quantitative measurement of microbial
biomass in soils. Soil Biology and Biochemistry 10, 215–221. doi:10.1016/0038-0717(78)90099-8

Anderson, T.H., Domsch, K.H., 2010. Soil microbial biomass: The eco-physiological approach. Soil Biology and
Biochemistry 42, 2039–2043. doi:10.1016/j.soilbio.2010.06.026

Augusto, L., De Schrijver, A., Vesterdal, L., Smolander, A., Prescott, C., Ranger, J., 2015. Influences of
evergreen gymnosperm and deciduous angiosperm tree species on the functioning of temperate and
boreal forests. Biological Reviews 90, 444–466. doi:10.1111/brv.12119

Averill, C., Turner, B.L., Finzi, A.C., 2014. Mycorrhiza-mediated competition between plants and decomposers
drives soil carbon storage. Nature 505, 543–545. doi:10.1038/nature12901

Bardgett, R.D., van der Putten, W.H., 2014. Belowground biodiversity and ecosystem functioning. Nature 515,
505–511. doi:10.1038/nature13855

Bauhus, J., van der Meer, P.J., Kanninen, M., 2010. Plantation forests: global perspectives, in: Ecosystem
Goods and Services from Plantation Forests. Earthscan, pp. 1–16. doi:10.4324/9781849776417

Beck, T., Joergensen, R.G., Kandeler, E., Makeschin, F., Nuss, E., Oberholzer, H.R., Scheu, S., 1997. An inter-
laboratory comparison of ten different ways of measuring soil microbial biomass C. Soil Biology and
Biochemistry 29, 1023–1032. doi:10.1016/S0038-0717(97)00030-8

Bennett, J.A., Koch, A.M., Forsythe, J., Johnson, N.C., Tilman, D., Klironomos, J., 2020. Resistance of soil biota
and plant growth to disturbance increases with plant diversity. Ecology Letters 23, 119–128.
doi:10.1111/ele.13408

Berger, T.W., Berger, P., 2012. Greater accumulation of litter in spruce (Picea abies) compared to beech
(Fagus sylvatica) stands is not a consequence of the inherent recalcitrance of needles. Plant and Soil
358, 349–369. doi:10.1007/s11104-012-1165-z

Biedermann, P.H.W., Müller, J., Grégoire, J.C., Gruppe, A., Hagge, J., Hammerbacher, A., Hofstetter, R.W.,
Kandasamy, D., Kolarik, M., Kostovcik, M., Krokene, P., Sallé, A., Six, D.L., Turrini, T., Vanderpool, D.,
Wingfield, M.J., Bässler, C., 2019. Bark beetle population dynamics in the Anthropocene: Challenges
and solutions. Trends in Ecology and Evolution. doi:10.1016/j.tree.2019.06.002

Bluhm, S.L., Eitzinger, B., Ferlian, O., Bluhm, C., Schröter, K., Pena, R., Maraun, M., Scheu, S., 2019.
Deprivation of root-derived resources affects microbial biomass but not community structure in litter
and soil. PLoS ONE 14, e0214233. doi:10.1371/journal.pone.0214233

Callaway, R.M., Pennings, S.C., Richards, C.L., 2003. Phenotypic plasticity and interactions among plants.
Ecology 84, 1115–1128. doi:10.1890/0012-9658(2003)084[1115:PPAIAP]2.0.CO;2

Cardinale, B.J., Wright, J.P., Cadotte, M.W., Carroll, I.T., Hector, A., Srivastava, D.S., Loreau, M., Weis, J.J.,
2007. Impacts of plant diversity on biomass production increase through time because of species
complementarity. Proceedings of the National Academy of Sciences of the United States of America
104, 18123–18128. doi:10.1073/pnas.0709069104

Castro-Díez, P., Vaz, A.S., Silva, J.S., van Loo, M., Alonso, Á., Aponte, C., Bayón, Á., Bellingham, P.J., Chiuffo,
M.C., DiManno, N., Julian, K., Kandert, S., La Porta, N., Marchante, H., Maule, H.G., Mayfield, M.M.,
Metcalfe, D., Monteverdi, M.C., Núñez, M.A., Ostertag, R., Parker, I.M., Peltzer, D.A., Potgieter, L.J.,
Raymundo, M., Rayome, D., Reisman-Berman, O., Richardson, D.M., Roos, R.E., Saldaña, A., Shackleton,
R.T., Torres, A., Trudgen, M., Urban, J., Vicente, J.R., Vilà, M., Ylioja, T., Zenni, R.D., Godoy, O., 2019.
Global effects of non-native tree species on multiple ecosystem services. Biological Reviews 94, 1477–
1501. doi:10.1111/brv.12511

Clemmensen, K.E., Bahr, A., Ovaskainen, O., Dahlberg, A., Ekblad, A., Wallander, H., Stenlid, J., Finlay, R.D.,
Wardle, D.A., Lindahl, B.D., 2013. Roots and associated fungi drive long-term carbon sequestration in
boreal forest. Science 340, 1615–1618. doi:10.1126/science.1231923

Cremer, M., Kern, N.V., Prietzel, J., 2016. Soil organic carbon and nitrogen stocks under pure and mixed
stands of European beech, Douglas fir and Norway spruce. Forest Ecology and Management 367, 30–40.
doi:10.1016/j.foreco.2016.02.020

David, A.S., Thapa-Magar, K.B., Menges, E.S., Searcy, C.A., Afkhami, M.E., Thapa-Magar, K.B., Menges, E.S.,
Searcy, C.A., Afkhami, M.E., Thapa-Magar, K.B., Menges, E.S., Searcy, C.A., Afkhami, M.E., Thapa-Magar,
K.B., Menges, E.S., Searcy, C.A., Afkhami, M.E., 2020. Do plant–microbe interactions support the Stress
Gradient Hypothesis? Ecology 00, e03081. doi:10.1002/ecy.3081

de Vries, F.T., Williams, A., Stringer, F., Willcocks, R., McEwing, R., Langridge, H., Straathof, A.L., 2019.
Changes in root-exudate-induced respiration reveal a novel mechanism through which drought affects
ecosystem carbon cycling. New Phytologist 224, 132–145. doi:10.1111/nph.16001

Defossez, E., Courbaud, B., Marcais, B., Thuiller, W., Granda, E., Kunstler, G., 2011. Do interactions between
plant and soil biota change with elevation? A study on Fagus sylvatica. Biology Letters 7, 699–701.
doi:10.1098/rsbl.2011.0236

Druebert, C., Lang, C., Valtanen, K., Polle, A., 2009. Beech carbon productivity as driver of ectomycorrhizal
abundance and diversity. Plant, Cell and Environment 32, 992–1003. doi:10.1111/j.1365-
3040.2009.01983.x

Eisenhauer, N., Lanoue, A., Strecker, T., Scheu, S., Steinauer, K., Thakur, M.P., Mommer, L., 2017. Root
biomass and exudates link plant diversity with soil bacterial and fungal biomass. Scientific Reports 7.
doi:10.1038/srep44641

Emmett Duffy, J., Godwin, C.M., Cardinale, B.J., 2017. Biodiversity effects in the wild are common and as
strong as key drivers of productivity. Nature 549, 261–264. doi:10.1038/nature23886

Fanin, N., Kardol, P., Farrell, M., Nilsson, M.C., Gundale, M.J., Wardle, D.A., 2019. The ratio of Gram-positive
to Gram-negative bacterial PLFA markers as an indicator of carbon availability in organic soils. Soil
Biology and Biochemistry 128, 111–114. doi:10.1016/j.soilbio.2018.10.010

Fierer, N., Schimel, J.P., Holden, P.A., 2003. Variations in microbial community composition through two soil
depth profiles. Soil Biology and Biochemistry 35, 167–176. doi:10.1016/S0038-0717(02)00251-1

Foltran, E.C., Ammer, C., Lamersdorf, N., 2020. Douglas fir and Norway spruce admixtures to beech forests
along a site gradient in Northern Germany − Are soil nutrient conditions affected? BioRxiv
2020.09.25.313213. doi:https://doi.org/10.1101/2020.09.25.313213

Frostegård, Å., Bååth, E., 1996. The use of phospholipid fatty acid analysis to estimate bacterial and fungal
biomass in soil. Biology and Fertility of Soils 22, 59−65. doi:10.1007/s003740050076

Frostegård, Å., Tunlid, A., Baath, E., 1993. Phospholipid fatty acid composition, biomass, and activity of
microbial communities from two soil types experimentally exposed to different heavy metals. Applied
and Environmental Microbiology 59, 3605−3617. doi:10.1128/aem.59.11.3605-3617.1993

Frostegård, Å., Tunlid, A., Bååth, E., 2011. Use and misuse of PLFA measurements in soils. Soil Biology and
Biochemistry 43, 1621−1625. doi:10.1016/j.soilbio.2010.11.021

Fry, E.L., Manning, P., Macdonald, C., Hasegawa, S., De Palma, A., Power, S.A., Singh, B.K., 2016. Shifts in
microbial communities do not explain the response of grassland ecosystem function to plant functional
composition and rainfall change. Soil Biology and Biochemistry 92, 199−210.
doi:10.1016/j.soilbio.2015.10.006

Gadgil, R.L., Gadgil, P.D., 1971. Mycorrhiza and litter decomposition. Nature 233, 133. doi:10.1038/233133a0

Guckert, J.B., Hood, M.A., White, D.C., 1986. Phospholipid ester-linked fatty acid profile changes during
nutrient deprivation of Vibrio cholerae: Increases in the trans/cis ratio and proportions of cyclopropyl
fatty acids. Applied and Environmental Microbiology 52, 794−801. doi:10.1128/aem.52.4.794-801.1986

Harrison, X.A., Donaldson, L., Correa-Cano, M.E., Evans, J., Fisher, D.N., Goodwin, C.E.D., Robinson, B.S.,
Hodgson, D.J., Inger, R., 2018. A brief introduction to mixed effects modelling and multi-model
inference in ecology. PeerJ 2018, 1−32. doi:10.7717/peerj.4794

Hättenschwiler, S., Tiunov, A. V., Scheu, S., 2005. Biodiversity and litter decomposition in terrestrial
ecosystems. Annual Review of Ecology, Evolution, and Systematics 36, 191−218.
doi:10.1146/annurev.ecolsys.36.112904.151932

Högberg, M.N., Högberg, P., Myrold, D.D., 2007. Is microbial community composition in boreal forest soils
determined by pH, C-to-N ratio, the trees, or all three? Oecologia 150, 590−601. doi:10.1007/s00442-
006-0562-5

Högberg, P., Read, D.J., 2006. Towards a more plant physiological perspective on soil ecology. Trends in
Ecology and Evolution 21, 548−554. doi:10.1016/j.tree.2006.06.004

Kaiser, C., Koranda, M., Kitzler, B., Fuchslueger, L., Schnecker, J., Schweiger, P., Rasche, F., Zechmeister-
Boltenstern, S., Sessitsch, A., Richter, A., 2010. Belowground carbon allocation by trees drives seasonal
patterns of extracellular enzyme activities by altering microbial community composition in a beech
forest soil. New Phytologist 187, 843−858. doi:10.1111/j.1469-8137.2010.03321.x

Kieft, T.L., Ringelberg, D.B., White, D.C., 1994. Changes in ester-linked phospholipid fatty acid profiles of
subsurface bacteria during starvation and desiccation in a porous medium. Applied and Environmental
Microbiology 60, 3292−3299. doi:10.1128/aem.60.9.3292-3299.1994

Knoke, T., Ammer, C., Stimm, B., Mosandl, R., 2008. Admixing broadleaved to coniferous tree species: A
review on yield, ecological stability and economics. European Journal of Forest Research 127, 89−101.
doi:10.1007/s10342-007-0186-2

Koranda, M., Kaiser, C., Fuchslueger, L., Kitzler, B., Sessitsch, A., Zechmeister-Boltenstern, S., Richter, A.,
2013. Seasonal variation in functional properties of microbial communities in beech forest soil. Soil

Biology and Biochemistry 60, 95–104. doi:10.1016/j.soilbio.2013.01.025

Kramer, C., Gleixner, G., 2008. Soil organic matter in soil depth profiles: Distinct carbon preferences of
microbial groups during carbon transformation. Soil Biology and Biochemistry 40, 425–433.
doi:10.1016/j.soilbio.2007.09.016

Laliberté, E., Kardol, P., Didham, R.K., Teste, F.P., Turner, B.L., Wardle, D.A., 2017. Soil fertility shapes
belowground food webs across a regional climate gradient. Ecology Letters. doi:10.1111/ele.12823

Leuschner, C., Ellenberg, H., Leuschner, C., Ellenberg, H., 2017. Beech and mixed beech forests, in: Ecology of
Central European Forests. Springer, pp. 351–441. doi:10.1007/978-3-319-43042-3_5

Leuschner, C., Hertel, D., Schmid, I., Koch, O., Muhs, A., Hölscher, D., 2004. Stand fine root biomass and fine
root morphology in old-growth beech forests as a function of precipitation and soil fertility. Plant and
Soil 258, 43–56. doi:10.1023/B:PLSO.0000016508.20173.80

Lindahl, B.D., Ihrmark, K., Boberg, J., Trumbore, S.E., Högberg, P., Stenlid, J., Finlay, R.D., 2007. Spatial
separation of litter decomposition and mycorrhizal nitrogen uptake in a boreal forest. New Phytologist
173, 611–620. doi:10.1111/j.1469-8137.2006.01936.x

Lundquist, E.J., Scow, K.M., Jackson, L.E., Uesugi, S.L., Johnson, C.R., 1999. Rapid response of soil microbial
communities from conventional, low input, and organic farming systems to a wet/dry cycle. Soil Biology
and Biochemistry 31, 1661–1675. doi:10.1016/S0038-0717(99)00080-2

MacArthur, R.H., 1957. On the relative abundance of bird species. Proceedings of the National Academy of
Sciences 43, 293–295. doi:10.1073/pnas.43.3.293

Macfadyen, A., 1961. Improved funnel-type extractors for soil arthropods. Journal of Animal Ecology 30, 171.
doi:10.2307/2120

Malchair, S., Carnol, M., 2009. Microbial biomass and C and N transformations in forest floors under
European beech, sessile oak, Norway spruce and Douglas-fir at four temperate forest sites. Soil Biology
and Biochemistry 41, 831–839. doi:10.1016/j.soilbio.2009.02.004

Maraun, M., Scheu, S., 1995. Influence of beech litter fragmentation and glucose concentration on the
microbial biomass in three different litter layers of a beechwood. Biology and Fertility of Soils 19, 155–
158. doi:10.1007/BF00336152

Meier, I.C., Leuschner, C., 2008. Genotypic variation and phenotypic plasticity in the drought response of fine
roots of European beech. Tree Physiology 28, 297–309. doi:10.1093/treephys/28.2.297

Meier, I.C., Leuschner, C., Hertel, D., 2005. Nutrient return with leaf litter fall in Fagus sylvatica forests across
a soil fertility gradient. Plant Ecology 177, 99–112. doi:10.1007/s11258-005-2221-z

Meier, I.C., Tückmantel, T., Heitkötter, J., Müller, K., Preusser, S., Wrobel, T.J., Kandeler, E., Marschner, B.,
Leuschner, C., 2020. Root exudation of mature beech forests across a nutrient availability gradient: the
role of root morphology and fungal activity. New Phytologist 226, 583–594. doi:10.1111/nph.16389

Moore-Kucera, J., Dick, R.P., 2008. PLFA profiling of microbial community structure and seasonal shifts in
soils of a Douglas-fir chronosequence. Microbial Ecology 55, 500–511. doi:10.1007/s00248-007-9295-1

Nacke, H., Goldmann, K., Schöning, I., Pfeiffer, B., Kaiser, K., Villamizar, G.A.C., Schrumpf, M., Buscot, F.,
Daniel, R., Wubet, T., 2016. Fine spatial scale variation of soil microbial communities under European
beech and Norway spruce. Frontiers in Microbiology 7. doi:10.3389/fmicb.2016.02067

Pettit, J., Voelker, S., DeRose, R.J., Burton, J., 2020. Spruce beetle outbreak was not driven by drought stress:

evidence from a tree-ring iso-demographic approach indicate temperatures were more important.
Global Change Biology. doi:10.1111/gcb.15274

Peuke, A.D., Gessler, A., Rennenberg, H., 2006. The effect of drought on C and N stable isotopes in different
fractions of leaves, stems and roots of sensitive and tolerant beech ecotypes. Plant, Cell and
Environment 29, 823–835. doi:10.1111/j.1365-3040.2005.01452.x

Piovia-Scott, J., Yang, L.H., Wright, A.N., Spiller, D.A., Schoener, T.W., 2019. Pulsed seaweed subsidies drive
sequential shifts in the effects of lizard predators on island food webs. Ecology Letters 22, 1850–1859.
doi:10.1111/ele.13377

Pollierer, M.M., Ferlian, O., Scheu, S., 2015. Temporal dynamics and variation with forest type of
phospholipid fatty acids in litter and soil of temperate forests across regions. Soil Biology and
Biochemistry 91, 248–257. doi:10.1016/j.soilbio.2015.08.035

Pollierer, M.M., Langel, R., Körner, C., Maraun, M., Scheu, S., 2007. The underestimated importance of
belowground carbon input for forest soil animal food webs. Ecology Letters 10, 729–736.
doi:10.1111/j.1461-0248.2007.01064.x

Potthast, K., Hamer, U., Makeschin, F., 2010. Impact of litter quality on mineralization processes in managed
and abandoned pasture soils in Southern Ecuador. Soil Biology and Biochemistry 42, 56–64.
doi:10.1016/j.soilbio.2009.09.025

Prescott, C.E., Grayston, S.J., Helmisaari, H.S., Kaštovská, E., Körner, C., Lambers, H., Meier, I.C., Millard, P.,
Ostonen, I., 2020. Surplus Carbon Drives Allocation and Plant–Soil Interactions. Trends in Ecology and
Evolution. doi:10.1016/j.tree.2020.08.007

R Core Team, 2019. R: A Language and Environment for Statistical Computing. R Fundation for Statistical
Computing, Vienna, Austria. Version 3.6.2. URL https://www.R-project.org/.

Ramsey, P.W., Rillig, M.C., Feris, K.P., Holben, W.E., Gannon, J.E., 2006. Choice of methods for soil microbial
community analysis: PLFA maximizes power compared to CLPP and PCR-based approaches.
Pedobiologia 50, 275–280. doi:10.1016/j.pedobi.2006.03.003

Ratcliffe, S., Wirth, C., Jucker, T., van der Plas, F., Scherer-Lorenzen, M., Verheyen, K., Allan, E., Benavides, R.,
Bruelheide, H., Ohse, B., Paquette, A., Ampoorter, E., Bastias, C.C., Bauhus, J., Bonal, D., Bouriaud, O.,
Bussotti, F., Carnol, M., Castagneyrol, B., Ćhećko, E., Dawud, S.M., Wandeler, H. De, Domisch, T., Finér,
L., Fischer, M., Fotelli, M., Gessler, A., Granier, A., Grossiord, C., Guyot, V., Haase, J., Hättenschwiler, S.,
Jactel, H., Jaroszewicz, B., Joly, F.X., Kambach, S., Kolb, S., Koricheva, J., Liebersgesell, M., Milligan, H.,
Müller, S., Muys, B., Nguyen, D., Nock, C., Pollastrini, M., Purschke, O., Radoglou, K., Raulund-
Rasmussen, K., Roger, F., Ruiz-Benito, P., Seidl, R., Selvi, F., Seiferling, I., Stenlid, J., Valladares, F.,
Vesterdal, L., Baeten, L., 2017. Biodiversity and ecosystem functioning relations in European forests
depend on environmental context. Ecology Letters 20, 1414–1426. doi:10.1111/ele.12849

Richter, A., Schöning, I., Kahl, T., Bauhus, J., Ruess, L., 2018. Regional environmental conditions shape
microbial community structure stronger than local forest management intensity. Forest Ecology and
Management 409, 250–259. doi:10.1016/j.foreco.2017.11.027

Schall, P., Lödige, C., Beck, M., Ammer, C., 2012. Biomass allocation to roots and shoots is more sensitive to
shade and drought in European beech than in Norway spruce seedlings. Forest Ecology and
Management 266, 246–253. doi:10.1016/j.foreco.2011.11.017

Scheu, S., 2002. The soil food web: structure and perspectives. European Journal of Soil Biology 38, 11–20.
doi:10.1016/S1164-5563(01)01117-7

Scheu, S., 1992. Automated measurement of the respiratory response of soil microcompartments: Active
microbial biomass in earthworm faeces. Soil Biology and Biochemistry 24, 1113–1118.
doi:10.1016/0038-0717(92)90061-2

Schmid, I., 2002. The influence of soil type and interspecific competition on the fine root system of Norway
spruce and European beech. Basic and Applied Ecology 3, 339–346. doi:10.1078/1439-1791-00116

Schmid, M., Pautasso, M., Holdenrieder, O., 2014. Ecological consequences of Douglas fir (Pseudotsuga
menziesii) cultivation in Europe. European Journal of Forest Research 133, 13–29. doi:10.1007/s10342-
013-0745-7

Toïgo, M., Vallet, P., Perot, T., Bontemps, J.D., Piedallu, C., Courbaud, B., 2015. Overyielding in mixed forests
decreases with site productivity. Journal of Ecology 103, 502–512. doi:10.1111/1365-2745.12353

van der Heijden, M.G.A., Bardgett, R.D., Van Straalen, N.M., 2008. The unseen majority: Soil microbes as
drivers of plant diversity and productivity in terrestrial ecosystems. Ecology Letters 11, 296–310.
doi:10.1111/j.1461-0248.2007.01139.x

Waller, L.P., Allen, W.J., Barratt, B.I.P., Condron, L.M., França, F.M., Hunt, J.E., Koele, N., Orwin, K.H., Steel,
G.S., Tylianakis, J.M., Wakelin, S.A., Dickie, I.A., 2020. Biotic interactions drive ecosystem responses to
exotic plant invaders. Science 368, 967–972. doi:10.1126/science.aba2225

Wardle, D.A., Bardgett, R.D., Klironomos, J.N., Setälä, H., Van Der Putten, W.H., Wall, D.H., 2004. Ecological
linkages between aboveground and belowground biota. Science. doi:10.1126/science.1094875

Wright, A.J., Mommer, L., Barry, K., Ruijven, J., 2020. Stress gradients and biodiversity: monoculture
vulnerability drives stronger biodiversity effects during drought years. Ecology. doi:10.1002/ecy.3193

Zelles, L., 1999. Fatty acid patterns of phospholipids and lipopolysaccharides in the characterisation of
microbial communities in soil: A review. Biology and Fertility of Soils 29, 111–129.
doi:10.1007/s003740050533

Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A., Smith, G.M., 2009. Dealing with heterogeneity, in: Mixed
Effects Models and Extensions in Ecology with R. Springer Science & Business Media, pp. 71–100.
doi:10.1007/978-0-387-87458-6_4

Figures and Tables

Figure 1. Effects of conifers and conifer-beech mixtures on microbial basal respiration (μg O_2 g^{-1} C h^{-1}), microbial biomass (μg C_{mic} g^{-1} C) and microbial specific respiration (μg O_2 μg^{-1} C_{mic} h^{-1}) at nutrient-poor and nutrient-rich sites (Douglas-fir [Do], Douglas-fir with beech [Do/Be], Norway spruce [Sp] and Norway spruce with beech [Sp/Be]). Effect sizes are given as back transformed log response ratios compared to beech forests [ln (value in coniferous or mixed forest types / values in beech)]. Effect sizes were averaged across three depths (litter, 0–5 and 5–10 cm soil). Values smaller than 1 indicate higher values in beech. Asterisks indicate significant effects ($p < 0.05$). Bars represent 95% confidence intervals (n=12).

Figure 2. Effects of conifers and conifer-beech mixtures on stress indicators: ratio of cyclopropyl PLFAs to its monoenoic precursors (cy/pre), ratio of saturated to monounsaturated PLFAs (sat/mono), and ratio of Gram$^+$ to Gram$^-$ bacteria (Gram$^+$/Gram$^-$) at nutrient-poor and nutrient-rich sites (Douglas-fir [Do], Douglas-fir with beech [Do/Be], Norway spruce [Sp] and Norway spruce with beech [Sp/Be]). Effect sizes are given as back transformed log response ratios compared to beech forests [ln (value in coniferous or mixed forest types / values in beech)]. Effect sizes were averaged across three depths (litter, 0–5 and 5–10 cm soil). Values larger than 1 indicate lower values in beech. Asterisks indicate significant effects ($p < 0.05$). Bars represent 95% confidence intervals (n=12).

Figure 3. Changes in microbial basal respiration (μg O_2 g^{-1} C h^{-1}), microbial biomass (μg C_{mic} g^{-1} C) and microbial specific respiration (μg O_2 μg^{-1} C_{mic} h^{-1}) with soil depth (litter and 0–5 and 5–10 cm soil) in five forest types (European beech [Be], Douglas-fir [Do], Douglas-fir with beech [Do/Be], Norway spruce [Sp] and Norway spruce with beech [Sp/Be]) at nutrient-poor and nutrient-rich sites. Points and horizontal bars represent means and standard errors (n=4). The grey vertical bars represent respective values in beech forests at nutrient-poor sites in litter, 0–5 and 5–10 cm soil. Note log scale for microbial biomass.

Figure 4. Changes in stress indicators (ratio of cyclopropyl fatty acids to its precursors [cy/pre], ratio of saturated to monounsaturated fatty acids [sat/mono] and ratio of Gram$^+$ to Gram$^-$ bacteria [Gram$^+$/Gram$^-$]) with soil depth (litter and 0–5 and 5–10 cm soil) in five forest types (European beech [Be], Douglas-fir [Do], Douglas-fir with beech [Do/Be], Norway spruce [Sp] and Norway spruce with beech [Sp/Be]) at nutrient-poor and nutrient-rich sites. Points and horizontal bars represent means and standard errors (n=4). The grey vertical bars represent respective values in beech forests at nutrient-poor sites in litter, 0–5 and 5–10 cm soil.

Figure 5. Principal component analysis of the microbial community structure (as indicated by phospholipid-derived fatty acids) in five forest types (European beech [Be], Douglas-fir [Do], Douglas-fir with beech [Do/Be], Norway spruce [Sp] and Norway spruce with beech [Sp/Be]) at nutrient-poor and nutrient-rich sites in litter, 0–5 and 5–10 cm soil. Positions of forest types represent centroids (n=4). Nutrient-poor sites are given in red and nutrient-rich sites in green. C/N, pH, water content (water%), carbon content (C%) as well as litter mass were post-fitted into the plot. Significant variables selected by RDA permutation test were in bold (based adjusted R^2).

Table 1. General information as well as soil physical and chemical properties of the study sites. Soil texture is based on 5–30 cm mineral soil, and soil chemical properties are based on 0–5 cm mineral soil. Soil moisture is expressed as percentage of dry mass, and was measured in litter, 0–5 and 5–10 cm soil. Ca^{2+} concentration was analyzed in exchangeable form by NH_4Cl extraction, and total P was determined by pressure digestion (for detailed see Methods and Foltran et al. 2020).

	Site condition	Rich sites				Poor sites			
	Site number	1	2	3	4	5	6	7	8
	Site name	Harz	Dassel	Winnefeld	Nienover	Nienburg	Unterlüß	Göhrde II	Göhrde I
General information	Latitude (°N)	51.770	51.726	51.662	51.696	52.621	52.837	53.127	53.201
	Longitude (°E)	10.397	9.704	9.574	9.526	9.281	10.331	10.814	10.801
	MAT (°C)	7.63	8.59	8.90	9.01	9.70	9.03	9.19	9.20
	MAP (mm)	1029.24	821.29	822.43	874.97	733.34	746.56	681.68	672.63
	Slope (°)	15.52	2.96	4.97	6.56	1.85	1.19	1.43	2.05
	Elevation (m)	510.80	426.17	349.06	323.37	92.00	161.20	129.80	120.00
Texture & moisture	Sand%	16	26	20	20	80	79	79	73
	Silt%	16	53	57	57	13	15	15	24
	Clay%	68	21	23	23	7	6	6	3
	Water% (Litter)	237.2	221.2	235.5	232.4	245.9	303.1	328.5	265.7
	Water% (0–5)	135.5	106.6	84.7	62.7	71.4	120.2	88.1	76.2
	Water% (5–10)	59.1	39.6	37.8	29.6	22.4	26.9	23.9	22.9
Chemical properties	C %	7.09	6.19	7.50	5.30	7.97	4.03	9.31	5.36
	N %	0.37	0.34	0.37	0.29	0.28	0.17	0.34	0.24
	C/N	19.16	18.21	20.27	18.28	28.46	23.71	27.38	22.33
	CEC (mmol dm^{-3})	237.31	121.26	114.91	95.96	74.81	47.79	71.10	67.72
	pH (KCl)	3.42	3.15	3.34	3.47	3.00	3.00	2.71	3.01
	P (mg kg^{-1})	0.61	0.54	0.44	0.46	0.16	0.11	0.17	0.24
	Ca^{2+} (CEC; mmol dm^{-3})	28.35	13.77	32.56	27.88	20.38	12.91	7.90	11.28

MAT: mean annual temperature
MAP: mean annual precipitation
CEC: cation exchange capacity

Table 2. F- and P-values of linear mixed effects models on the effect of Forest type (European
beech, Douglas-fir, Norway spruce, mixture of European beech with Douglas-fir and mixture of
European beech with Norway spruce), Site condition (nutrient-poor and nutrient-rich sites) and
Depth (litter, 0–5 and 5–10 cm) on (a) basal respiration, microbial biomass and microbial specific
respiration, (b) stress indicators (ratio of cyclopropyl PLFAs to its monoenoic precursors [cy/pre],
ratio of saturated to monounsaturated PLFAs [sat/mono], Gram$^+$/Gram$^-$ ratio), and (c) percentages
of fungal and bacterial PLFAs (of total PLFAs), and fungi/bacteria ratio. Significant effects are given
in bold (p < 0.05).

(a)	Basal respiration			Microbial biomass			Specific respiration		
Factor	df	F	P	df	F	P	df	F	P
Forest type (F)	4,24	8.94	**<0.001**	4,30	5.95	**0.001**	4,24	0.92	0.470
Site condition (S)	1,6	3.90	0.096	1,30	16.93	**<0.001**	1,6	0.52	0.498
Depth (D)	2,78	168.37	**<0.001**	2,70	271.27	**<0.001**	2,78	17.37	**<0.001**
F × S	4,24	10.91	**<0.001**	4,30	4.61	**0.005**	4,24	2.36	0.082
F × D	-	-	**-**	8,70	2.15	**0.042**	-	-	-

(b)	cy/pre ratio			sat/mono ratio			Gram$^+$/Gram$^-$ ratio		
Factor	df	F	P	df	F	P	df	F	P
Forest type (F)	4,24	2.75	0.052	4,24	8.278	**<0.001**	4,24	3.39	**<0.001**
Site condition (S)	1,6	3.48	0.111	1,6	0.195	0.304	1,6	8.17	**0.028**
Depth (D)	2,68	238.82	**<0.001**	2,76	5.551	**<0.001**	2,76	228.51	**<0.001**
F × S	4,24	2.70	0.055	4,24	3.650	**0.002**	4,24	2.27	**0.012**
S × D	2,68	6.20	**0.003**	2,76	10.705	**<0.001**	2,76	7.51	**0.001**
F × D	8,68	3.72	**0.001**	-	-	-	-	-	-

(c)	% fungi			% bacteria			fungi/bacteria ratio		
Factor	df	F	P	df	F	P	df	F	P
Forest type (F)	4,30	1.77	0.161	4,30	2.19	0.094	4,24	0.83	0.520
Site condition (S)	1,30	6.46	**0.016**	1,30	4.34	**0.046**	1,6	4.98	**0.033**
Depth (D)	2,68	982.11	**<0.001**	2,78	149.46	**<0.001**	2,68	995.40	**<0.001**
F × S	4,30	1.99	0.121	4,30	0.73	0.577	4,24	1.60	0.207
S × D	2,68	18.54	**<0.001**	-	-	-	2,68	16.92	**<0.001**
F × D	8,68	3.61	**0.002**	-	-	-	8,68	2.84	**0.009**

Table 3. Results of two-way MANOVAs on PLFA composition (a) without covariates and (b) with
covariates (pH, C/N ratio, water content) in litter, 0–5 and 5–10 cm soil. Factors include Forest type
(European beech, Douglas-fir, Norway spruce and mixture of European beech with Douglas-fir and
mixture of European beech with Norway spruce), Site condition (nutrient-rich and nutrient-poor sites)
and their interactions. MANOVAs were based on principal components from each sample depth
(significant PCs were determined by broken stick criterion; see Methods). Type I sequential sum of
square was used for models with and without covariates. Significant effects are given in bold (P <
0.05).

| Depth | Factor | (a) Without covariates | | | | (b) With covariates | | | |
		df	Wilk's lambda	F	P	df	Wilk's lambda	F	P
						Covariates: pH			
Litter	Forest type (F)	4,30	0.32	3.29	**0.001**	4,29	0.39	2.53	**0.008**
	Site condition (S)	1,30	0.42	12.77	**<0.001**	1,29	0.61	5.72	**0.004**
	F × S	4,30	0.73	0.77	0.677	4,29	0.75	0.68	0.767
						Covariates: C/N + pH + water%			
0–5 cm	Forest type (F)	4,30	0.19	2.80	**<0.001**	4,27	0.30	1.67	0.057
	Site condition (S)	1,30	0.39	8.14	**<0.001**	1,27	0.75	1.54	0.218
	F × S	4,30	0.36	1.57	0.078	4,27	0.33	1.53	0.096
						Covariates: C/N			
5–10 cm	Forest type (F)	4,30	0.41	1.76	0.051	4,29	0.41	1.68	0.067
	Site condition (S)	1,30	0.45	8.15	**<0.001**	1,29	0.82	1.39	0.265
	F × S	4,30	0.69	0.68	0.805	4,29	0.73	0.54	0.915