Running title: Forest type effects on microbial communities

Title: Mixing coniferous and deciduous trees (*Fagus sylvatica*): Site-specific response of soil microorganisms

Authors: Jing-Zhong Lu[1*], Stefan Scheu[1,2]

[1] J. F. Blumenbach Institute of Zoology and Anthropology, University of Göttingen, Untere Karspüle 2, 37073

Göttingen, Germany

[2] Center of Biodiversity and Sustainable Land Use, University of Göttingen, Büsgenweg 1, 37077 Göttingen,

Germany

[*] Corresponding author (E-mail: jlu@gwdg.de)

Keywords: Douglas-fir, Norway spruce, soil respiration, forest, ecosystem functioning, PLFA

Abstract

Coniferous and deciduous trees growing in pure and mixed forests supply different resources to soil microorganisms thereby shaping microbial communities. However, effects of different forest types including non-native species on microbial communities and their variation with site conditions such as soil type and nutritional status remain poorly understood. We quantified functional (substrate-induced respiration and stress indicators) and structural (phospholipid-derived fatty acids) responses of microbial communities to forest types at four nutrient-rich and four nutrient-poor sites. Five types of forests were investigated at each site, including pure stands of native European beech (*Fagus sylvatica*), range-expanding Norway spruce (*Picea abies*) and non-native Douglas-fir (*Pseudotsuga menziesii*), as well as mixed species stands of beech with spruce or with Douglas-fir, i.e., 40 forest plots. Forest type only affected soil microorganisms at nutrient-poor sites, but not at nutrient-rich sites. At nutrient-poor sites, both pure and mixed coniferous forests, in particular pure Douglas-fir forests, detrimentally affected soil microorganisms compared to beech forests. By contrast, structural and functional indicators of microorganisms in beech forests at nutrient-poor sites resembled those at nutrient-rich sites, likely because beech maintained the provisioning of high amounts of root-derived resources. The results indicate that long-term effects of planting exotic tree species such as Douglas-fir either in pure or in mixed stands on ecosystem functioning need further attention especially at nutrient-poor sites, but planting Douglas-fir at nutrient-rich sites appear of little concern from the perspective of microbial communities. Overall, the results highlight the potential importance of root-derived resources in determining microbial community structure and functioning, and document the sensitivity of soil microorganisms to planting non-native tree species that may differ in the provisioning of these resources from native species.

1. Introduction

Studying the linkage between plants and soil microorganisms has great potential to improve our understanding of the factors that drive microbial community composition (Balvanera et al., 2006; van der Heijden et al., 2008; Anderson and Domsch, 2010; Bardgett and van der Putten, 2014; Graham et al., 2016). Trees as long-lived plants may affect microbial communities through several pathways, most notably by the quantity and quality of litter and root exudates (Wardle et al., 2004; Högberg and Read, 2006). These resources are likely to shape the community structure of soil microorganisms as dissimilar carbon resources favor different guilds of microorganisms (Ekblad and Nordgren, 2002; Fierer et al., 2003; Fanin et al., 2019). For example, Gram[-] bacteria are benefiting in particular from labile carbon compounds, whereas Gram[+] bacteria are adapted to use more recalcitrant carbon resources (Fierer et al., 2003; Fanin et al., 2019; Waller et al., 2020). Similarly, niches vary between fungal groups. While saprotrophic fungi are major decomposers of plant litter, mycorrhizal fungi are more abundant in deeper soil and receive carbon from plant roots (Lindahl et al., 2007; Nacke et al., 2016). Therefore, the composition of microbial communities may change in response to variations in tree species composition, associated with changes in litter quality, root-derived resources and mycorrhizal associations (L. Chen et al., 2019).

Conifers and deciduous trees growing in pure and mixed stands likely differ in the supply of resources to soil microorganisms. Accumulation of organic matter in the floor of coniferous forests has been attributed to the recalcitrance of conifer needles, but roots and mycorrhizal fungi also affect the storage of soil carbon (Schmidt et al., 2011; Clemmensen et al., 2013; Averill et al., 2014). Below-ground processes, such as root exudation and the competition of saprotrophic and mycorrhizal fungi, maybe more important for structuring microbial communities than the amount and quality of above-ground litter (Scheu et al., 2003; Pollierer et al., 2007; Schmidt et al., 2011; Augusto et al., 2015; Wang et al., 2016; Bluhm et al., 2019; Jones et al., 2019). Further, the response of microbial communities to mixed forests is poorly understood. Litter mixture may improve resource availability for soil microorganisms and facilitate litter decomposition (Hättenschwiler et al., 2005; Fujii et al., 2020), although tree identity has been suggested to outweigh tree diversity as structuring force of soil microbial communities (Scheibe et al., 2015; Dawud et al., 2016; L. Chen et al., 2019). Mixing of tree species has also been shown to increase both above-ground and below-ground biomass due to interspecific facilitation and improved resource partitioning, resulting in higher complementarity in space and time in mixed forests (Cardinale et al., 2007; Ma and Chen, 2016; Emmett Duffy et al., 2017). These interactions likely affect the way forests shaping soil microbial communities. Overall, however, the role of identity and diversity of tree species, either coniferous or deciduous, in shaping the structure and functioning of soil microorganisms is little studied.

The relationship between tree species and microbial communities strongly depends on environmental context (Ratcliffe et al., 2017; X. Chen et al., 2019), but little is known to what extent effects of tree species composition on soil microorganisms vary with site conditions, such as soil type and nutritional status. Facilitation among plants is more pronounced in nutrient-limited environments according to the stress-gradient hypothesis, and such positive interspecific interactions at nutrient-poor conditions are likely to be mediated by soil microorganisms (Defossez et al., 2011; Harpole et al., 2016; David et al., 2020). In fact, plants do not passively tolerate environmental stress, but adapt in various ways to unfavorable growth conditions by induced plasticity (Callaway et al., 2003; de Vries et al., 2019). As a trade-off, plants generally allocate more resources to roots than to shoots under nutrient-poor conditions (Kawaletz et al., 2013; Yan et al., 2016), and this likely modifies soil microbial communities. Indeed, de Vries et al. (2019) recently showed that under drought conditions herb species change the quality rather than quantity of root exudates and induce higher microbial respiration, presumably facilitating nutrient capture via stimulating microorganisms. Therefore, the role of tree species composition as driver of microbial community composition and functioning may be more pronounced under nutrient-poor than nutrient-rich conditions, but this has not been investigated until today.

A better understanding of the effects of tree species composition and site conditions on soil microorganisms has important implications for forest management, especially in temperate and boreal regions where tree species richness is low and managed forests typically are dominated by only one or a few species (Knoke et al., 2008; Hu et al., 2009; Bauhus et al., 2010). While Norway spruce currently is the most popular high-yielding timber species in Central Europe, in recent years it has have been severely damaged by extreme conditions of weather and the outbreak of bark beetles, and such events are predicted to occur more frequently and at higher severity in the future (Biedermann et al., 2019; Brandl et al., 2020; Pettit et al., 2020). For reducing the risk of damage and maintaining economic gains, planting of mixed forests including non-native species such as Douglas-fir is increasingly discussed (Pretzsch et al., 2017; Ammer et al., 2018). To date, the impact of planting Douglas-fir on biodiversity and functioning of forests is little studied, and this applies in particular to the belowground system, although e.g., Douglas-fir has been suggested to affect soil chemistry in a similar way than spruce (Schmid et al., 2014). Taken together, we lack a comprehensive evaluation of forest types on the structure and functioning of soil microbial communities across nutritional gradients.

Here, we investigate the structure and functioning of microbial communities in litter and soil of five forest types at nutrient-rich and nutrient-poor sites. Forest types include pure stands of European beech (*Fagus sylvatica*), Norway spruce (*Picea abies*), Douglas-fir (*Pseudotsuga menziesii*), and the mixtures of the two coniferous species with beech. European beech forms the climax tree species in lowland and lower montane

regions of Central Europe (Leuschner et al., 2017). Norway spruce originally only occurred at higher

mountain ranges and boreal regions in Central Europe, but it has been planted across wide areas of Central

Europe including lowlands (Knoke et al., 2008; Schmid et al., 2014). Douglas-fir was introduced into Germany

from North America more than 150 years ago and has become the most abundant non-native tree species

cultivated in Central Europe (Schmid et al., 2014). To understand the response of microorganisms to planting

different tree species in single and mixed stands, physiological (respiration based) and community

approaches (phospholipid fatty acid analysis) were combined. We quantified microbial basal respiration,

microbial biomass, and community structure as well as stress indicators of microorganisms (Anderson and

Domsch, 2010). Physiological response variables and stress indicators were used as functional indicators of

microbial communities. Generally, we assumed the structure of microbial communities to be linked to their

functioning. Specifically, we hypothesized (1) coniferous trees to detrimentally affect microbial basal

respiration, microbial biomass and stress indicators, with the effects being similar in Norway spruce and

Douglas-fir forests, but alleviated in mixed forests with beech. Further, we hypothesized (2) effects of forest

types on microbial community structure and functioning to be present at both nutrient-rich and nutrient-

poor sites, but to be stronger at nutrient-poor sites. Since the litter layer is less buffered against

environmental conditions than the organic and mineral soil, we also hypothesized that (3) the effects of

forest type on the structure and functioning of microbial communities are more pronounced in litter than in

soil.

2. Methods

2.1. Field sites

The study was conducted in 40 forest stands located in the south and north of Lower Saxony in Germany

(eight sites x five forest types; Fig. S1). Eight sites were selected to cover a range of soil nutrients and water

conditions based on prior knowledge of soil types and annual total rainfall. Four sites in the south stock on

fertile soil and receive higher precipitation. Parental rock material is either loess influenced Triassic

sandstone or Paleozoic shares of greywacke, sandstone, quartzite and phyllite, leading to soil types of partly

podsolic Cambisol and Luvisol. Mean annual precipitation is 821 – 1029 mm (calculated from German

weather service since 1981; Glatthorn 2017). The four northern sites are located on out-washed sand with

the soil type of Podzol. The mean annual precipitation is 672 – 746 mm (Table S1). Each site (quintet)

comprised three pure stands of European beech (*Fagus sylvatica* L.; Be), Douglas-fir (*Pseudotsuga menziesii*

[Mirbel] Franco.; Do), Norway spruce (*Picea abies* [L.] Karst.; Sp) and two conifer-beech mixtures (Douglas-

fir/European beech and Norway spruce/European beech; Do/Be and Sp/Be, respectively). On average trees

were more than 50 years old in 2017. Within quintets, the distance between plots ranged from 76 m to 4600

m, and the distance of quintets ranged from 5 to 190 km. Within each stand, plots of 2500 m^2 were

133 established, mostly in rectangular shape (50 x 50 m). Tree species in pure stands comprised more than 90%

134 of the total basal area, and that in mixed stands 33 – 53% for beech and 53 – 60% for conifers.

2.2. Soil sampling and properties

Samples were taken in the eight quintets between November 2017 and January 2018. Soil cores of 5 cm

diameters were taken by a metal cylinder, and were separated into litter, 0 – 5 and 5 – 10 cm soil depth. In

each plot, three cores spaced by 5 m were taken. Samples from the same depth were pooled, resulting in a

total of 120 samples (40 plots × 3 depths). The soil was sieved through 2 mm mesh, and the litter was

fragmented into pieces using scissors (approximately < 25 mm^2) (Maraun and Scheu, 1995). Roots and

stones were removed. Samples were frozen at -20°C and thawed at 4°C before the analyses. The pH (KCl) of

litter and soil was determined using a ratio of sample to solution of 1:10, 1:5 and 1:2.5 (g/v) for litter, 0 – 5

and 5 – 10 cm soil, respectively. Dried soil (70°C, 48 h) was grinded in a ball mill and then total C and N

concentrations were determined by an elemental analyzer (NA 1110, CE-instruments, Rodano, Milano, Italy).

2.3. Microbial basal respiration and biomass.

Microbial respiration was measured by O_2 consumption using an automated micro-respirometer. The system

absorbs CO_2 produced by microorganisms using KOH solution, and the O_2 consumption by microorganisms is

replaced through electrolytic release of O_2 from $CuSO_4$ solution (Scheu, 1992). Microbial basal respiration

was measured as mean consumption of O_2 during 10 – 23 h after attachment of the vessels to the

respirometer system (BR; µg O_2 g^{-1} h^{-1}). Microbial biomass was determined by substrate-induced respiration

after the addition of glucose to saturate microbial glycolytic enzymes based on the maximum initial

respiratory response 4 – 7 h after the addition of D-glucose (*MIRR*; µl O_2 g^{-1} h^{-1}) and converted to microbial

biomass (C_{mic}; µg C_{mic} g^{-1}) as 38 × *MIRR* (Anderson and Domsch, 1978; Beck et al., 1997). Fresh litter and soil

from 0 – 5 and 5 – 10 cm depth equivalent to approximately 0.3, 1 and 3.5 g dry mass was supplemented

with 80, 20 and 8 mg glucose solved in 400 µl H_2O, respectively. The ratio between microbial basal

respiration and microbial biomass was taken as microbial specific respiration (qO_2; µg O_2 $µg^{-1}$ C_{mic} h^{-1}).

2.4. Phospholipid fatty acid analysis

To quantify the composition of phospholipid-derived fatty acid (PLFA) composition, lipids from leaf litter and

soil were extracted using a modified Bligh and Dyer method (Frostegård et al., 1993b) as described in

Pollierer et al. (2015). Lipids were fractionated into glycolipids, phospholipids and neutral lipids by elution

through silica acid columns using chloroform, acetone and methanol, respectively (0.5 g silicic acid, 3 ml; HF

BOND ELUT-SI, Varian, Inc., Darmstadt, Germany). Phospholipids were subjected to mild alkaline

methanolysis, and fatty acid methyl esters (FAMEs) of phospholipids were identified by chromatographic

retention time compared to a general reference standard and a bacteria-derived standard (FAME CRM47885,

C11 to C24; BAME 47080-U, C11 to C20; Sigma-Aldrich, Darmstadt, Germany) using a GC-FID Clarus 500

(PerkinElmer Corporation, Norwalk, USA) equipped with an Elite 5 column (30 m × 0.32 mm inner diameter,

film thickness 0.25 μm). The temperature program started with 60 °C (hold time 1 min) and increased with

30 °C per min to 160 °C, and then by 3 °C per min to 280 °C. The injection temperature was 250 °C and

helium was used as carrier gas. About 2 g of fresh litter and 4 g of fresh soil were used for the extraction.

2.5. Stress indicators and fatty acid markers

The ratio of cyclopropyl PLFAs to their monoenoic precursors (cy/pre) and the ratio of saturated to

monounsaturated PLFAs (sat/mono) were used as indicators of physiological or nutritional stress ([(cy17:0 +

cy19:0)/(16:1ω7 + 18:1ω7)]; [(14:0 + 15:0 + 16:0 + 17:0 + 18:0)/(16:1ω7 + 17:1 + 18:1ω9 + 18:1ω7)];

respectively) (Moore-Kucera and Dick, 2008; Pollierer et al., 2015). The ratio of Gram$^+$ to Gram$^-$ bacteria was

used as indicator of carbon availability (Fanin et al., 2019). Linoleic acid 18:2ω6,9 was used as fungal marker

(Frostegård and Bååth, 1996). The saturated fatty acids i15:0, a15:0, i16:0, i17:0 were used as markers for

Gram$^+$ bacteria, and the fatty acids cy17:0, cy19:0, 16:1ω7 and 18:1ω7 were assigned as markers for Gram$^-$

bacteria (Frostegård and Bååth, 1996; Zelles, 1999; Mitchell et al., 2010; Richter et al., 2018; Fanin et al.,

2019). Bacterial biomass was represented by the sum of Gram$^+$ and Gram$^-$ bacteria. Total amount of PLFAs

were summed up from all identified PLFAs (n = 40; nmol g^{-1} dry weight) and was used to calculate PLFA

proportions. All stress indicators and FA markers were analyzed from proportions (mole percentage).

2.6. Statistical analyses

For univariate analysis, we first fitted linear mixed models (LMMs) to log-transformed response variables

and then applied planned contrasts to estimate the effect size of forest type and quintet group (Piovia-Scott

et al., 2019). All LMMs included Forest type (European beech, Douglas-fir, Douglas-fir/European beech,

Norway spruce, Norway spruce/European beech), Quintet group (nutrient-rich and nutrient-poor sites) and

Depth (litter, 0 – 5 and 5 – 10 cm) as fixed effects. Backwards model selection was applied to full models,

and minimal models included all main effects and the interaction of forest type and quintet group. The 40

forest plots were included as random effects to account for non-independence of samples of different depth

from the same plot by forcing values from the same plot to have the same intercept (Harrison et al., 2018).

Based on Akaike Information Criterion (AIC), the eight quintets were included as random effect, and the

mean - variance relationship was also accounted for by a dispersion parameter to meet the assumption of

homogeneity of variance (Johnson and Omland, 2004; Zuur et al., 2009). Further, fixed effects were selected

based on likelihood ratio test.

Contrasts for the effect size of forest type were designed to compare coniferous and mixed forests to beech

forests. European beech, the climax tree species in lowland and lower montane regions of Central Europe,

was used as reference (Leuschner et al., 2017). Due to the property of log-transformed response variables,

the planned contrasts are analog to log response ratios (Piovia-Scott et al., 2019). Further, the log response

ratios were back transformed into response ratios and were used as effect size. Effect sizes of forest type

were estimated for nutrient-poor and nutrient-rich sites. Additionally, we applied contrasts between

nutrient-poor vs. nutrient-rich sites in a similar way for the effect size of quintet group. Univariate response

variables included microbial basal respiration, microbial biomass, microbial specific respiration, stress

indicators and PLFA markers.

For multivariate analysis of variance (MANOVA), PLFAs first were arcsine root transformed and then

dimensions of fatty acids were reduced by principal component analysis (PCA). Separate MANOVAs for each

sample depth were calculated to inspect for effects of Forest type, Quintet group and their interactions. Soil

properties (pH, C/N ratio) were included as covariates in multivariate analysis of covariance (MANCOVA)

(Langerhans et al., 2007). Significant principal components (PCs) were determined by broken stick criterion

(MacArthur, 1957). Only fatty acids ≥ 0.2% mean mole proportion were included in the analyses.

All analyses were done in R with RStudio interface (R Core Team, 2019; RStudio Team, 2019). We used the

"nlme" package to fit the linear mixed-effects models (Pinheiro et al., 2019), and the "emmeans" package to

conduct planned contrasts (Lenth, 2020). All mixed models met the assumptions of normality of residuals

and homogeneity of variance.

214 3. Results

215 3.1. Functional indicators of microorganisms

Forest type consistently affected microbial basal respiration, microbial biomass and stress indicators at

nutrient-poor but not at nutrient-rich sites (Forest type x Quintet group interactions; all $P < 0.08$; Table 1,

Figs 1 and 2), and this was generally true across the studied soil layers (Fig. S2). At nutrient-poor sites,

microbial basal respiration and microbial biomass was lower in coniferous and mixed forests than in beech

forests. This was most pronounced in Douglas-fir, where at nutrient-poor sites basal respiration and

microbial biomass were 65% and 59% lower compared to pure beech, respectively; similarly, basal

respiration and microbial biomass in the two conifer-beech mixtures were 46 – 56% lower than in beech.

Microbial basal respiration and microbial biomass in spruce forests were also lower than in beech forests,

but the difference was less pronounced (-33 to -36%; Fig. 1). In addition, in Douglas-fir forests at the

nutrient-poor sites microbial specific respiration was 19% lower compared to beech.

In contrast to basal respiration and microbial biomass, at nutrient-poor sites stress indicators, the cy/pre

ratio, the sat/mono ratio as well as the Gram$^+$/Gram$^-$ bacteria ratio, all were lower in beech forests than in

the other forest types (Fig. 2). The difference in the cy/pre and sat/mono ratio was most pronounced in

Douglas-fir (+87% and +74%) and less in spruce forests (+39% and +43%). Similar to spruce forest, the cy/pre

and sat/mono ratio in conifer-beech mixtures was on average 37 – 46% higher than in beech forests. Effects

of forest type on the Gram[+]/Gram[-] bacteria ratio also was strongest in Douglas-fir forests (+46 %) and less

pronounced in the other forest types (+21 – 29%). Stress indicators generally did not differ at nutrient-rich

sites (P > 0.31; Fig. 2, Fig. S2, Table S2).

The response of microorganisms to changes in site conditions also varied between forest types. Differences

between nutrient-rich and nutrient-poor sites were most prominent in Douglas-fir forests and not significant

in beech forests (Figs 3 and 4, Table S3). At nutrient-poor sites, microbial basal respiration in Douglas-fir

forests was 57% lower and microbial biomass in Douglas-fir pure and mixed forests both were 47% lower

compared to nutrient-rich sites. Similarly, but less strong, microbial biomass in spruce mixed forests was 36%

lower at nutrient-poor sites than nutrient-rich sites. Conversely, in Douglas-fir forests the stress indicator

cy/pre ratio at nutrient-poor sites exceeded that of nutrient-rich sites by 82% (Fig. 4, Table S3). Similarly, in

Douglas-fir forests the sat/mono and Gram[+]/Gram[-] bacteria ratio at nutrient-poor sites exceeded that at

nutrient-rich sites by 38 – 40%. Further, in Douglas-fir mixed forests the Gram[+]/Gram[-] bacteria ratio at

nutrient-poor sites exceeded that at nutrient-poor sites by 34%.

3.2. Microbial community structure

As indicated by PLFA patterns, microbial community composition in beech forest differed from that in

Norway spruce and Douglas-fir forests, with mixed forests being intermediate between the respective single

species stands (Fig. 5). However, effects of forest type varied among soil layers and were most pronounced

in litter and 0 – 5 cm soil, but only marginally significant in 5 – 10 cm soil (Table 2). Despite strong turnover

between nutrient-rich and nutrient-poor sites (Quintet group effects p < 0.001), beech maintained similar

microbial community structure between nutrient-poor and nutrient-rich sites, and this was most

pronounced in 0 – 5 cm soil depth (Forest type × Quintet group interaction; Table 2, Fig. 5).

A number of fatty acids contributed to the separation of microbial communities of the different forest types

and quintet groups. Beech forests were characterized by unsaturated fatty acids at all soil depths, in

particular by 16:1ω7, 18:1ω7, 18:1ω9 (Fig. 5). Norway spruce forests were associated with long-chain

saturated fatty acids (22:0, 24:0), and Douglas-fir forests with bacterial marker fatty acids (2OH-14:0, cy17:0).

There also were clear differences in fatty acid patterns between nutrient-rich and nutrient-poor sites. Litter

at nutrient-rich sites was associated with unsaturated fatty acids such as 18:1ω9 and the fungal marker

18:2ω6,9. By contrast, nutrient-poor sites had higher saturated and branched fatty acids, with many of them

being bacterial markers, such as 14:0 and cy17:0 and cy19:0. Whereas the litter layer at nutrient-rich sites

was richer in fungi than at nutrient-poor sites, the opposite was true for the soil (Quintet group × Depth

interaction; Table 1, Figs S3 and S4).

As indicated by PLFA markers, the relative abundance of fungi and bacteria generally did not differ among

forest types, but consistently the relative abundance of fungi decreased, whereas the abundance of bacteria

increased with soil depth (Table 1, Fig. S3). Further, the abundance of fungi in the soil of Douglas-fir forests

was lower than in beech forests and the difference was more pronounced in 5 – 10 cm soil compared to

other depths at nutrient-poor sites (Fig. S4).

The northern nutrient-poor sites also were more acidic and more limited by nitrogen as indicated by lower

soil pH and higher C/N ratio, and the main effects of Quintet group and Forest type became non-significant

when pH and C/N ratio were included as covariates in the multivariate analysis in soil (Table 2, Fig. 5). In

addition, soil pH significantly explained the structure of microbial communities particularly in 0 – 5 cm depth

(Table 2).

272 4. Discussion

Overall, the results only in part support our first and second hypotheses. Forest types only affected the

functioning of microbial communities at nutrient-poor sites, but not at nutrient-rich sites. At nutrient-poor

sites, microbial basal respiration, microbial biomass and stress indicators all were detrimentally affected in

pure and mixed coniferous forests, and, contrasting our first hypothesis the effects differed between

Douglas-fir and Norway spruce, with the effects being particularly strong in Douglas-fir, but less pronounced

in Norway spruce and mixed stands. By contrast, similar basal respiration, microbial biomass and stress

indicators at nutrient-rich sites across the studied forest types contradict our second hypothesis and suggest

that, at favorable environmental conditions, soil microbial communities are rather irresponsive to changes in

tree species. Notably, microbial biomass and microbial stress indicators in beech forests did not differ

between nutrient-poor and nutrient-rich sites indicating that beech is able to maintain the functioning of soil

microbial communities regardless of site conditions. Admixture of beech in mixed stands with Douglas-fir or

Norway spruce both detrimentally affected microbial biomass and microbial stress indicators, suggesting

that caution is needed when mixing beech with conifers. However, results of the present study also indicate

that compared to pure Douglas-fir forests, mixed Douglas-fir forests with beech mitigate the detrimental

effects of pure Douglas-fir forests on soil microbial communities. More broadly, exotic plants lack co-

evolutionary history with native species and therefore, complementary effects between native and exotic

plants may be stronger and more likely to occur than between native species (Kawaletz et al., 2013). This

may also explain the more pronounced alleviation of microbial stress in mixtures of Douglas-fir with beech

than in those of Norway spruce with beech.

The site- and tree species-specific responses of microorganisms are likely caused by varying energy supply in

different forest types at nutrient-poor sites. In fact, at nutrient-poor sites, the higher Gram$^+$/Gram$^-$ bacteria

ratio in coniferous and mixed forests suggests lower supply of labile carbon than in beech forests (Fanin et al., 2019). Labile carbon compounds have been suggested to favor Gram$^-$ bacteria, whereas Gram$^+$ bacteria better cope with recalcitrant carbon resources (Kramer and Gleixner, 2008; Fanin et al., 2019; Waller et al., 2020). Among the forest types studied, Douglas-fir forests were particularly limited by labile carbon as further supported by its lower microbial specific respiration, which was largely due to lower microbial basal respiration compared to beech stands (Anderson and Domsch, 2010). Limited supply of labile carbon also has been suggested to favor oligotrophic microbial communities characterized by low respiration rates and biomass (Potthast et al., 2010; Fanin et al., 2019). This is consistent with our findings of strong carbon limitation as well as low basal respiration and microbial biomass in single and mixed coniferous forests at nutrient-poor sites. Moreover, at nutrient-poor sites, both the cy/pre and sat/mono ratios were higher in pure species coniferous as well as mixed forests than in pure beech forests, also indicating higher nutrient (and/or water) stress in the former (Moore-Kucera and Dick, 2008; Pollierer et al., 2015). The Gram$^-$ bacteria marker fatty acids 16:1ω7 and 18:1ω7 are precursors of cyclopropyl fatty acids and as monounsaturated fatty acids. They form part of the nominator of the calculated stress indicators. The more pronounced stress response of microorganism in coniferous and mixed forests at nutrient-poor sites was largely caused by lower concentration of these two ω7 fatty acids. Indeed, it has been suggested that under stress conditions Gram$^-$ bacteria change the composition of their cell membranes from 16:1ω7 and 18:1ω7 to cyclopropane fatty acids associated with slower growth rates (Guckert et al., 1986; Kieft et al., 1994; Lundquist et al., 1999). The similarity of microbial indicators in beech forests at nutrient-rich and nutrient-poor sites indicates that, compared to coniferous stands, beech maintained microbial community functioning and avoided stress conditions of microorganisms by providing more labile carbon resources at nutrient-poor conditions, presumably by releasing high amounts of root-derived resources.

The conclusion that beech is able to alleviate microbial stress via maintaining or even increasing the release of root-derived resources at nutrient-poor sites is supported by the following findings. First, current studies at our study sites showed that at nutrient-poor sites the biomass of fine roots is higher in beech forests than in pure and mixed coniferous forests while at nutrient-rich sites root biomass varies little among forest types (A. Lwila, pers. comm.). Generally, it is well established that plants allocate more resources into roots if the availability of soil nutrients is declining and this is associated with increased root exudation (Eisenhauer et al., 2017; Laliberté et al., 2017). Recently, this also was confirmed for beech forests where higher root exudation was found in more acidic and nitrogen deficient stands (Meier et al., 2020). These findings support our conclusion that beech indeed maintained or even increased its root exudates at nutrient-poor sites which in fact were characterized by more acidic conditions and higher C/N ratio in soil. High root-derived resources presumably also were responsible for the alleviated microbial resource deficiency and stress in beech forests at nutrient-poor sites. Second, trenching and girdling experiments confirmed that reducing or omitting the

flux of root-derived resources into the soil strongly reduces fungal and bacterial biomass, demonstrating the

importance of root-derived resources for maintaining soil microbial biomass (Högberg et al., 2007; Kaiser et

al., 2010; Bluhm et al., 2019). Notably, we sampled litter and soil during autumn / winter after beech had

shed its leaves. Nevertheless, the evidence provided above suggests that the impact of roots on

microorganisms still was strong indicating that not only recent photosynthates but also stored carbohydrates

contribute to providing root-derived resources to soil microorganisms (Druebert et al., 2009). Further, these

findings suggest that beech has the potential to facilitate soil microorganisms at environmental stress

conditions by maintaining or even increasing its root exudates contrasting both of the studied coniferous

species. This ability presumably contributed to the mitigation of the detrimental effects of Douglas-fir on

microorganisms at nutrient-poor / acidic sites in mixed forests. More broadly, it may even explain the

observed overyielding in mixed stands of beech and conifers at nutrient-poor sites, highlighting the potential

role of microorganisms in the facilitation of plants under environment stress conditions (Toïgo et al., 2015;

Ammer, 2019; David et al., 2020).

Contrary to our third hypothesis, the effect of forest types was not stronger in litter than in soil. At nutrient-

poor sites, forest type affected microbial basal respiration, microbial biomass and microbial stress in both

litter and soil, with the strength of the effects generally being similar across layers. This supports our

conclusion that the effect of forest type was not mainly due to differences in litter quantity and quality, but

in particular to differences in root-derived resources. This contrasts earlier suggestions that microorganisms

in litter are more sensitive to changes in environmental conditions than those in soil because litter is less

buffered against environmental harshness (Pollierer et al., 2015). Supporting our conclusion that

microorganisms in soil sensitively respond to environmental changes, microbial stress indicators generally

increased while microbial biomass generally decreased with soil depth. This sensitivity presumably is related

to the fact that carbon availability generally decreases with soil depth resulting in stronger resource

limitation (Fierer et al., 2003).

We assumed that the functioning of microorganisms ultimately relies on its microbial community structure,

and this is supported by the consistent turnover in microbial community composition as well as the change

in microbial functions between nutrient-rich and nutrient-poor sites. Further, in contrast to coniferous and

mixed forests, microbial community structure and functional indices in beech forests differed little between

nutrient-rich and nutrient-poor sites, suggesting that among the tree species studied only beech is able to

maintain and stabilize microbial community structure and functioning in litter and soil irrespective of site

conditions. Notably, differences in microbial community structure between forest types at nutrient-rich sites

tended to be small in 5 – 10 cm soil depth, further suggesting that the influence of tree species fades deeper

in soil. By contrast, at nutrient-poor sites, differences in microbial community structure between forest types

were relatively large in soil, again supporting our conclusion that at nutrient-poor sites effects of tree species
on microorganisms in soil are predominantly due to differences in root-derived resources. Interestingly, the
differences in microbial community structure between beech and coniferous forests in 5 – 10 cm soil depth
at nutrient-poor sites were most pronounced in Douglas-fir, and carbon limitation also was most prominent
in 5 – 10 cm soil of Douglas-fir forests as indicated by lower microbial specific respiration. This implies that
among the tree species studied Douglas-fir provides the lowest amount of root-derived resources to soil
microorganisms at nutrient-poor sites resulting in pronounced resource shortage and marked changes in
microbial community composition. Such changes in microbial community structure may significantly impact
the functioning of soil microbial communities as the availability of carbon resources and carbon limitation
affect the vertical distribution of saprotrophic fungi and decomposition processes, e.g. by shifting the
competition between saprotrophic and mycorrhizal fungi (Gadgil and Gadgil, 1971; Lindahl et al., 2007).

The relative abundance of fungi and bacteria differed between nutrient-rich and nutrient-poor sites, but less
among forest types. As indicated by the relative abundance of the fungal marker FA 18:2ω6,9, fungi were
favored at nutrient-poor sites in spruce and beech forests, supporting earlier suggestions that the fungal
energy channel dominates in low fertile systems (Wardle et al., 2004). However, in contrast to beech and
spruce, the relative abundance of fungi in pure and mixed forests of Douglas-fir was not higher at nutrient-
poor sites than that at nutrient-rich sites. Further, at nutrient-poor sites the relative abundance of fungi was
lower in Douglas-fir than in beech forests and this was most pronounced in 5 – 10 cm depth. Potentially, the
lower relative abundance of fungi in soil of Douglas-fir forests is due to a mismatch between local and home
(native range of Douglas-fir) fungal communities (Schmid et al., 2014). In fact, in soil of introduced plant
species only a subset of fungal species present in their native range has been recorded in soil of sites where
they had been introduced (Schmid et al., 2014). Future studies need to examine in more detail the
taxonomic composition of microbial communities in different forest types, including introduced and native
tree species such as Douglas-fir and beech in particular at nutrient-poor sites deeper in soil.

In addition to root-derived resources, tree species may affect microbial community structure in soil by
modifying chemical properties. In our study, soil properties, especially pH, well explained changes in the
community structure of microorganisms in soil. This finding are consistent with results of continental scale
studies as well as local liming experiments suggesting that soil pH is among the most prominent driver of
microbial community composition (Frostegård et al., 1993a; Fierer and Jackson, 2006). Further, soil pH
generally correlates negatively with soil nutrient availability, supporting the importance of nutrients in the
mediation of microbial responses (Meier et al., 2020). However, factors other than soil pH and soil nutrient
status may have contributed to the observed differences between nutrient-rich and nutrient-poor sites. In
fact, nutrient-poor sites also were characterized by lower precipitation and this is reflected by higher $\delta^{13}C$

values of litter at nutrient-poor compared to nutrient-rich sites (J.-Z. Lu, unpubl. data), likely reflecting higher

water stress at nutrient-poor compared to nutrient-rich sites (Peuke et al., 2006). Presumably, in addition to

pH, both nutrients and water stress contributed to the observed differences in microbial community

composition and functioning between nutrient-poor and nutrient-rich sites. This is supported by the fact that

soil water conditions are intricately linked to soil nutrient dynamics and the uptake of nutrients by plants (de

Vries et al., 2019). Future studies need to disentangle the relative importance of water deficiency, nutrient

availability and soil pH as driving factors of the interrelationship between tree species and soil microbial

community composition and functioning. Focusing on nutrient-poor / low pH stands is most promising for

resolving these interrelationships.

5. Conclusion

The present study documented that effects of forest type on soil microbial communities vary among tree

species and mixed stands, but this strongly depends on the nutrient status of the sites. Notably, variations

between sites were least pronounced in native beech and most pronounced in introduced Douglas-fir.

Compared to beech, Douglas-fir strongly impacted microbial community structure and functioning at

nutrient-poor sites. Therefore, in the long-term planting non-native tree species such as Douglas-fir may

compromise ecosystem functioning at nutrient-poor sites, and this needs to be studied in more detail as it

may not only question the establishment of pure Douglas-fir forests, but also the enrichment of beech

forests by Douglas-fir at nutrient-poor sites. However, as planting Douglas-fir in single species and mixed

stands with beech hardly impacted the structure and functioning of soil microbial communities at nutrient-

rich sites, planting Douglas-fir may provide an alternative option to Norway spruce at least at nutrient-rich

sites. Such options are urgently needed since Norway spruce forests are increasingly threatened due to

global climate change and associated outbreaks of bark beetles especially in lowland regions of Europe.

However, more information is needed for gaining a holistic understanding on the consequences of planting

single species or mixed stands of Douglas-fir on ecosystem properties and processes, such as below- and

aboveground biodiversity of other taxa, and the decomposition of litter. Future studies need to focus on

potential facilitation or inhibition of planting mixed forests of beech and Douglas-fir on soil microorganisms

and timber yield in particular at nutrient-poor sites. Overall, our findings highlight that tree species

differently shape the structure and functioning of soil microbial communities with their effects varying

strongly with soil conditions.

Acknowledgements

We thank C. Bluhm, T. Volovei, G. Humpert, S. Zhou for assistance, and C. Ammer, H. Riebl, A. Polle, J.

Glatthorn, X. Lu, N. Lamersdorf, Z. Li, S. Bluhm, G. Schulz, T.-W. Chen, M. Maraun, A. Lwila, K. Schumann, I.

Bebre, A. Rivera and other RTG members for helpful discussions. Special thanks go to S. Müller for

coordination. The study formed part of the Research Training Group 2300: "Enrichment of European beech

forests with conifers: impacts of functional traits on ecosystem functioning" (DFG: GRK 2300). This work was

supported by the German Research Foundation [316045089].

430 References

Ammer, C., 2019. Diversity and forest productivity in a changing climate. New Phytologist 221, 50–66.
doi:10.1111/nph.15263

Ammer, C., Fichtner, A., Fischer, A., Gossner, M.M., Meyer, P., Seidl, R., Thomas, F.M., Annighöfer, P.,
Kreyling, J., Ohse, B., Berger, U., Feldmann, E., Häberle, K.H., Heer, K., Heinrichs, S., Huth, F., Krämer-
Klement, K., Mölder, A., Müller, J., Mund, M., Opgenoorth, L., Schall, P., Scherer-Lorenzen, M., Seidel,
D., Vogt, J., Wagner, S., 2018. Key ecological research questions for Central European forests. Basic and
Applied Ecology 32, 3–25. doi:10.1016/j.baae.2018.07.006

Anderson, J.P.E., Domsch, K.H., 1978. A physiological method for the quantitative measurement of microbial
biomass in soils. Soil Biology and Biochemistry 10, 215–221. doi:10.1016/0038-0717(78)90099-8

Anderson, T.H., Domsch, K.H., 2010. Soil microbial biomass: The eco-physiological approach. Soil Biology and
Biochemistry 42, 2039–2043. doi:10.1016/j.soilbio.2010.06.026

Augusto, L., De Schrijver, A., Vesterdal, L., Smolander, A., Prescott, C., Ranger, J., 2015. Influences of
evergreen gymnosperm and deciduous angiosperm tree species on the functioning of temperate and
boreal forests. Biological Reviews 90, 444–466. doi:10.1111/brv.12119

Averill, C., Turner, B.L., Finzi, A.C., 2014. Mycorrhiza-mediated competition between plants and decomposers
drives soil carbon storage. Nature 505, 543–545. doi:10.1038/nature12901

Balvanera, P., Pfisterer, A.B., Buchmann, N., He, J.S., Nakashizuka, T., Raffaelli, D., Schmid, B., 2006.
Quantifying the evidence for biodiversity effects on ecosystem functioning and services. Ecology Letters
9, 1146–1156. doi:10.1111/j.1461-0248.2006.00963.x

Bardgett, R.D., van der Putten, W.H., 2014. Belowground biodiversity and ecosystem functioning. Nature 515,
505–511. doi:10.1038/nature13855

Bauhus, J., van der Meer, P.J., Kanninen, M., 2010. Plantation forests: global perspectives, in: Ecosystem
Goods and Services from Plantation Forests. Earthscan, pp. 1–16. doi:10.4324/9781849776417

Beck, T., Joergensen, R.G., Kandeler, E., Makeschin, F., Nuss, E., Oberholzer, H.R., Scheu, S., 1997. An inter-
laboratory comparison of ten different ways of measuring soil microbial biomass C. Soil Biology and
Biochemistry 29, 1023–1032. doi:10.1016/S0038-0717(97)00030-8

Biedermann, P.H.W., Müller, J., Grégoire, J.C., Gruppe, A., Hagge, J., Hammerbacher, A., Hofstetter, R.W.,
Kandasamy, D., Kolarik, M., Kostovcik, M., Krokene, P., Sallé, A., Six, D.L., Turrini, T., Vanderpool, D.,
Wingfield, M.J., Bässler, C., 2019. Bark beetle population dynamics in the Anthropocene: Challenges
and solutions. Trends in Ecology and Evolution. doi:10.1016/j.tree.2019.06.002

Bluhm, S.L., Eitzinger, B., Ferlian, O., Bluhm, C., Schröter, K., Pena, R., Maraun, M., Scheu, S., 2019.
Deprivation of root-derived resources affects microbial biomass but not community structure in litter
and soil. PLoS ONE 14, e0214233. doi:10.1371/journal.pone.0214233

Brandl, S., Paul, C., Knoke, T., Falk, W., 2020. The influence of climate and management on survival
probability for Germany's most important tree species. Forest Ecology and Management 458, 117652.
doi:10.1016/j.foreco.2019.117652

Callaway, R.M., Pennings, S.C., Richards, C.L., 2003. Phenotypic plasticity and interactions among plants.
Ecology 84, 1115–1128. doi:10.1890/0012-9658(2003)084[1115:PPAIAP]2.0.CO;2

Cardinale, B.J., Wright, J.P., Cadotte, M.W., Carroll, I.T., Hector, A., Srivastava, D.S., Loreau, M., Weis, J.J.,
2007. Impacts of plant diversity on biomass production increase through time because of species
complementarity. Proceedings of the National Academy of Sciences of the United States of America
104, 18123–18128. doi:10.1073/pnas.0709069104

Chen, L., Xiang, W., Wu, H., Ouyang, S., Zhou, B., Zeng, Y., Chen, Y., Kuzyakov, Y., 2019. Tree species identity
surpasses richness in affecting soil microbial richness and community composition in subtropical forests.
Soil Biology and Biochemistry 130, 113–121. doi:10.1016/j.soilbio.2018.12.008

Chen, X., Chen, H.Y.H., Chen, C., Peng, S., 2019. Water availability regulates negative effects of species
mixture on soil microbial biomass in boreal forests. Soil Biology and Biochemistry 139, 107634.
doi:10.1016/j.soilbio.2019.107634

Clemmensen, K.E., Bahr, A., Ovaskainen, O., Dahlberg, A., Ekblad, A., Wallander, H., Stenlid, J., Finlay, R.D.,
Wardle, D.A., Lindahl, B.D., 2013. Roots and associated fungi drive long-term carbon sequestration in
boreal forest. Science 340, 1615–1618. doi:10.1126/science.1231923

David, A.S., Thapa-Magar, K.B., Menges, E.S., Searcy, C.A., Afkhami, M.E., Thapa-Magar, K.B., Menges, E.S.,
Searcy, C.A., Afkhami, M.E., Thapa-Magar, K.B., Menges, E.S., Searcy, C.A., Afkhami, M.E., Thapa-Magar,
K.B., Menges, E.S., Searcy, C.A., Afkhami, M.E., 2020. Do plant–microbe interactions support the Stress
Gradient Hypothesis? Ecology 00, e03081. doi:10.1002/ecy.3081

Dawud, S.M., Raulund-Rasmussen, K., Domisch, T., Finér, L., Jaroszewicz, B., Vesterdal, L., 2016. Is tree
species diversity or species identity the more important driver of soil carbon stocks, C/N ratio, and pH?
Ecosystems 19, 645–660. doi:10.1007/s10021-016-9958-1

de Vries, F.T., Williams, A., Stringer, F., Willcocks, R., McEwing, R., Langridge, H., Straathof, A.L., 2019.
Changes in root-exudate-induced respiration reveal a novel mechanism through which drought affects
ecosystem carbon cycling. New Phytologist 224, 132–145. doi:10.1111/nph.16001

Defossez, E., Courbaud, B., Marcais, B., Thuiller, W., Granda, E., Kunstler, G., 2011. Do interactions between
plant and soil biota change with elevation? A study on *Fagus sylvatica*. Biology Letters 7, 699–701.
doi:10.1098/rsbl.2011.0236

Druebert, C., Lang, C., Valtanen, K., Polle, A., 2009. Beech carbon productivity as driver of ectomycorrhizal
abundance and diversity. Plant, Cell and Environment 32, 992–1003. doi:10.1111/j.1365-
3040.2009.01983.x

Eisenhauer, N., Lanoue, A., Strecker, T., Scheu, S., Steinauer, K., Thakur, M.P., Mommer, L., 2017. Root
biomass and exudates link plant diversity with soil bacterial and fungal biomass. Scientific Reports 7.
doi:10.1038/srep44641

Ekblad, A., Nordgren, A., 2002. Is growth of soil microorganisms in boreal forests limited by carbon or
nitrogen availability? Plant and Soil 242, 115–122. doi:10.1023/A:1019698108838

Emmett Duffy, J., Godwin, C.M., Cardinale, B.J., 2017. Biodiversity effects in the wild are common and as
strong as key drivers of productivity. Nature 549, 261–264. doi:10.1038/nature23886

Fanin, N., Kardol, P., Farrell, M., Nilsson, M.C., Gundale, M.J., Wardle, D.A., 2019. The ratio of Gram-positive
to Gram-negative bacterial PLFA markers as an indicator of carbon availability in organic soils. Soil
Biology and Biochemistry 128, 111–114. doi:10.1016/j.soilbio.2018.10.010

Fierer, N., Jackson, R.B., 2006. The diversity and biogeography of soil bacterial communities. Proceedings of
the National Academy of Sciences of the United States of America 103, 626–631.
doi:10.1073/pnas.0507535103

Fierer, N., Schimel, J.P., Holden, P.A., 2003. Variations in microbial community composition through two soil
depth profiles. Soil Biology and Biochemistry 35, 167–176. doi:10.1016/S0038-0717(02)00251-1

Frostegård, A., Bååth, E., 1996. The use of phospholipid fatty acid analysis to estimate bacterial and fungal
biomass in soil. Biology and Fertility of Soils 22, 59–65. doi:10.1007/s003740050076

Frostegård, Å., Bååth, E., Tunlio, A., 1993a. Shifts in the structure of soil microbial communities in limed
forests as revealed by phospholipid fatty acid analysis. Soil Biology and Biochemistry 25, 723–730.
doi:10.1016/0038-0717(93)90113-P

Frostegård, Å., Tunlid, A., Baath, E., 1993b. Phospholipid fatty acid composition, biomass, and activity of
microbial communities from two soil types experimentally exposed to different heavy metals. Applied
and Environmental Microbiology 59, 3605–3617. doi:10.1128/aem.59.11.3605-3617.1993

Fujii, S., Berg, M.P., Cornelissen, J.H.C., 2020. Living litter: Dynamic trait spectra predict fauna composition.
Trends in Ecology and Evolution 00. doi:10.1016/j.tree.2020.05.007

Gadgil, R.L., Gadgil, P.D., 1971. Mycorrhiza and litter decomposition. Nature 233, 133. doi:10.1038/233133a0

Glatthorn, Jonas (2017): plot_general. v1.10.8. RTG2300-Information System. Dataset.
https://www.134.76.21.10/Data/ShowData.aspx?DatasetId=12540

Graham, E.B., Knelman, J.E., Schindlbacher, A., Siciliano, S., Breulmann, M., Yannarell, A., Beman, J.M., Abell,
G., Philippot, L., Prosser, J., Foulquier, A., Yuste, J.C., Glanville, H.C., Jones, D.L., Angel, R., Salminen, J.,
Newton, R.J., Bürgmann, H., Ingram, L.J., Hamer, U., Siljanen, H.M.P., Peltoniemi, K., Potthast, K.,
Bañeras, L., Hartmann, M., Banerjee, S., Yu, R.Q., Nogaro, G., Richter, A., Koranda, M., Castle, S.C.,
Goberna, M., Song, B., Chatterjee, A., Nunes, O.C., Lopes, A.R., Cao, Y., Kaisermann, A., Hallin, S.,
Strickland, M.S., Garcia-Pausas, J., Barba, J., Kang, H., Isobe, K., Papaspyrou, S., Pastorelli, R.,
Lagomarsino, A., Lindström, E.S., Basiliko, N., Nemergut, D.R., 2016. Microbes as engines of ecosystem
function: When does community structure enhance predictions of ecosystem processes? Frontiers in
Microbiology 7, 214. doi:10.3389/fmicb.2016.00214

Guckert, J.B., Hood, M.A., White, D.C., 1986. Phospholipid ester-linked fatty acid profile changes during
nutrient deprivation of Vibrio cholerae: Increases in the trans/cis ratio and proportions of cyclopropyl
fatty acids. Applied and Environmental Microbiology 52, 794–801. doi:10.1128/aem.52.4.794-801.1986

Harpole, W.S., Sullivan, L.L., Lind, E.M., Firn, J., Adler, P.B., Borer, E.T., Chase, J., Fay, P.A., Hautier, Y.,
Hillebrand, H., MacDougall, A.S., Seabloom, E.W., Williams, R., Bakker, J.D., Cadotte, M.W., Chaneton,
E.J., Chu, C., Cleland, E.E., D'Antonio, C., Davies, K.F., Gruner, D.S., Hagenah, N., Kirkman, K., Knops,
J.M.H., La Pierre, K.J., McCulley, R.L., Moore, J.L., Morgan, J.W., Prober, S.M., Risch, A.C., Schuetz, M.,
Stevens, C.J., Wragg, P.D., 2016. Addition of multiple limiting resources reduces grassland diversity.
Nature 537, 93–96. doi:10.1038/nature19324

Harrison, X.A., Donaldson, L., Correa-Cano, M.E., Evans, J., Fisher, D.N., Goodwin, C.E.D., Robinson, B.S.,
Hodgson, D.J., Inger, R., 2018. A brief introduction to mixed effects modelling and multi-model
inference in ecology. PeerJ 2018, 1–32. doi:10.7717/peerj.4794

Hättenschwiler, S., Tiunov, A. V., Scheu, S., 2005. Biodiversity and litter decomposition in terrestrial
ecosystems. Annual Review of Ecology, Evolution, and Systematics 36, 191–218.
doi:10.1146/annurev.ecolsys.36.112904.151932

Högberg, M.N., Högberg, P., Myrold, D.D., 2007. Is microbial community composition in boreal forest soils
determined by pH, C-to-N ratio, the trees, or all three? Oecologia 150, 590–601. doi:10.1007/s00442-
006-0562-5

Högberg, P., Read, D.J., 2006. Towards a more plant physiological perspective on soil ecology. Trends in
Ecology and Evolution 21, 548–554. doi:10.1016/j.tree.2006.06.004

Hu, J., Angeli, S., Schuetz, S., Luo, Y., Hajek, A.E., 2009. Ecology and management of exotic and endemic
Asian longhorned beetle Anoplophora glabripennis. Agricultural and Forest Entomology 11, 359–375.
doi:10.1111/j.1461-9563.2009.00443.x

Johnson, J.B., Omland, K.S., 2004. Model selection in ecology and evolution. Trends in Ecology and Evolution
19, 101–108. doi:10.1016/j.tree.2003.10.013

Jones, G.L., Scullion, J., Allison, G., Stott, H., Johnson, D., Owen, R., Williams, G., Gwynn-Jones, D., 2019.
Reduced soil respiration beneath invasive Rhododendron ponticum persists after cutting and is related
to substrate quality rather than microbial community. Soil Biology and Biochemistry 138.
doi:10.1016/j.soilbio.2019.107577

Kaiser, C., Koranda, M., Kitzler, B., Fuchslueger, L., Schnecker, J., Schweiger, P., Rasche, F., Zechmeister-
Boltenstern, S., Sessitsch, A., Richter, A., 2010. Belowground carbon allocation by trees drives seasonal
patterns of extracellular enzyme activities by altering microbial community composition in a beech
forest soil. New Phytologist 187, 843–858. doi:10.1111/j.1469-8137.2010.03321.x

Kawaletz, H., Mölder, I., Zerbe, S., Annighöfer, P., Terwei, A., Ammer, C., 2013. Exotic tree seedlings are
much more competitive than natives but show underyielding when growing together. Journal of Plant
Ecology 6, 305–315. doi:10.1093/jpe/rts044

Kieft, T.L., Ringelberg, D.B., White, D.C., 1994. Changes in ester-linked phospholipid fatty acid profiles of
subsurface bacteria during starvation and desiccation in a porous medium. Applied and Environmental
Microbiology 60, 3292–3299. doi:10.1128/aem.60.9.3292-3299.1994

Knoke, T., Ammer, C., Stimm, B., Mosandl, R., 2008. Admixing broadleaved to coniferous tree species: A
review on yield, ecological stability and economics. European Journal of Forest Research 127, 89–101.
doi:10.1007/s10342-007-0186-2

Kramer, C., Gleixner, G., 2008. Soil organic matter in soil depth profiles: Distinct carbon preferences of
microbial groups during carbon transformation. Soil Biology and Biochemistry 40, 425–433.
doi:10.1016/j.soilbio.2007.09.016

Laliberté, E., Kardol, P., Didham, R.K., Teste, F.P., Turner, B.L., Wardle, D.A., 2017. Soil fertility shapes
belowground food webs across a regional climate gradient. Ecology Letters. doi:10.1111/ele.12823

Langerhans, R.B., Gifford, M.E., Joseph, E.O., 2007. Ecological speciation in Gambusia fishes. Evolution 61,
2056–2074. doi:10.1111/j.1558-5646.2007.00171.x

Lenth, R., 2020. Emmeans: estimated marginal means, aka Least-Squares Means. R Package Version 1.4.4.
https://CRAN.R-project.org/package=emmeans

Leuschner, C., Ellenberg, H., Leuschner, C., Ellenberg, H., 2017. Beech and mixed beech forests, in: Ecology of
Central European Forests. Springer, pp. 351–441. doi:10.1007/978-3-319-43042-3_5

Lindahl, B.D., Ihrmark, K., Boberg, J., Trumbore, S.E., Högberg, P., Stenlid, J., Finlay, R.D., 2007. Spatial
separation of litter decomposition and mycorrhizal nitrogen uptake in a boreal forest. New Phytologist
173, 611–620. doi:10.1111/j.1469-8137.2006.01936.x

Lundquist, E.J., Scow, K.M., Jackson, L.E., Uesugi, S.L., Johnson, C.R., 1999. Rapid response of soil microbial
communities from conventional, low input, and organic farming systems to a wet/dry cycle. Soil Biology
and Biochemistry 31, 1661–1675. doi:10.1016/S0038-0717(99)00080-2

Ma, Z., Chen, H.Y.H., 2016. Effects of species diversity on fine root productivity in diverse ecosystems: a
global meta-analysis. Global Ecology and Biogeography 25, 1387–1396. doi:10.1111/geb.12488

MacArthur, R.H., 1957. On the relative abundance of bird species. Proceedings of the National Academy of
Sciences 43, 293–295. doi:10.1073/pnas.43.3.293

Maraun, M., Scheu, S., 1995. Influence of beech litter fragmentation and glucose concentration on the
microbial biomass in three different litter layers of a beechwood. Biology and Fertility of Soils 19, 155–
158. doi:10.1007/BF00336152

Meier, I.C., Tückmantel, T., Heitkötter, J., Müller, K., Preusser, S., Wrobel, T.J., Kandeler, E., Marschner, B.,
Leuschner, C., 2020. Root exudation of mature beech forests across a nutrient availability gradient: the
role of root morphology and fungal activity. New Phytologist 226, 583–594. doi:10.1111/nph.16389

Mitchell, R.J., Campbell, C.D., Chapman, S.J., Cameron, C.M., 2010. The ecological engineering impact of a
single tree species on the soil microbial community. Journal of Ecology 98, 50–61. doi:10.1111/j.1365-
2745.2009.01601.x

Moore-Kucera, J., Dick, R.P., 2008. PLFA profiling of microbial community structure and seasonal shifts in
soils of a Douglas-fir chronosequence. Microbial Ecology 55, 500–511. doi:10.1007/s00248-007-9295-1

Nacke, H., Goldmann, K., Schöning, I., Pfeiffer, B., Kaiser, K., Villamizar, G.A.C., Schrumpf, M., Buscot, F.,
Daniel, R., Wubet, T., 2016. Fine spatial scale variation of soil microbial communities under European
beech and Norway spruce. Frontiers in Microbiology 7. doi:10.3389/fmicb.2016.02067

Pettit, J., Voelker, S., DeRose, R.J., Burton, J., 2020. Spruce beetle outbreak was not driven by drought stress:
evidence from a tree-ring iso-demographic approach indicate temperatures were more important.
Global Change Biology. doi:10.1111/gcb.15274

Peuke, A.D., Gessler, A., Rennenberg, H., 2006. The effect of drought on C and N stable isotopes in different
fractions of leaves, stems and roots of sensitive and tolerant beech ecotypes. Plant, Cell and
Environment 29, 823–835. doi:10.1111/j.1365-3040.2005.01452.x

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., R Core Team, 2019. nlme: Linear and nonlinear mixed effects
models. R Package Version 3.1-142. https://CRAN.R-project.org/package=nlme

Piovia-Scott, J., Yang, L.H., Wright, A.N., Spiller, D.A., Schoener, T.W., 2019. Pulsed seaweed subsidies drive
sequential shifts in the effects of lizard predators on island food webs. Ecology Letters 22, 1850–1859.
doi:10.1111/ele.13377

Pollierer, M.M., Ferlian, O., Scheu, S., 2015. Temporal dynamics and variation with forest type of
phospholipid fatty acids in litter and soil of temperate forests across regions. Soil Biology and
Biochemistry 91, 248–257. doi:10.1016/j.soilbio.2015.08.035

Pollierer, M.M., Langel, R., Körner, C., Maraun, M., Scheu, S., 2007. The underestimated importance of
belowground carbon input for forest soil animal food webs. Ecology Letters 10, 729–736.
doi:10.1111/j.1461-0248.2007.01064.x

Potthast, K., Hamer, U., Makeschin, F., 2010. Impact of litter quality on mineralization processes in managed
and abandoned pasture soils in Southern Ecuador. Soil Biology and Biochemistry 42, 56–64.
doi:10.1016/j.soilbio.2009.09.025

Pretzsch, H., Knoke, T., Paul, C., Bauhus, J., Forrester, D.I., 2017. Perspectives for future research on mixed-
species systems, in: Mixed-Species Forests: Ecology and Management. Springer, pp. 579–606.
doi:10.1007/978-3-662-54553-9_12

R Core Team, 2019. R: A Language and Environment for Statistical Computing. R Fundation for Statistical
Computing, Vienna, Austria. Version 3.6.2. https://www.R-project.org/.

Ratcliffe, S., Wirth, C., Jucker, T., van der Plas, F., Scherer-Lorenzen, M., Verheyen, K., Allan, E., Benavides, R.,
Bruelheide, H., Ohse, B., Paquette, A., Ampoorter, E., Bastias, C.C., Bauhus, J., Bonal, D., Bouriaud, O.,
Bussotti, F., Carnol, M., Castagneyrol, B., Chećko, E., Dawud, S.M., Wandeler, H. De, Domisch, T., Finér,
L., Fischer, M., Fotelli, M., Gessler, A., Granier, A., Grossiord, C., Guyot, V., Haase, J., Hättenschwiler, S.,
Jactel, H., Jaroszewicz, B., Joly, F.X., Kambach, S., Kolb, S., Koricheva, J., Liebersgesell, M., Milligan, H.,
Müller, S., Muys, B., Nguyen, D., Nock, C., Pollastrini, M., Purschke, O., Radoglou, K., Raulund-
Rasmussen, K., Roger, F., Ruiz-Benito, P., Seidl, R., Selvi, F., Seiferling, I., Stenlid, J., Valladares, F.,
Vesterdal, L., Baeten, L., 2017. Biodiversity and ecosystem functioning relations in European forests
depend on environmental context. Ecology Letters. doi:10.1111/ele.12849

Richter, A., Schöning, I., Kahl, T., Bauhus, J., Ruess, L., 2018. Regional environmental conditions shape
microbial community structure stronger than local forest management intensity. Forest Ecology and
Management 409, 250–259. doi:10.1016/j.foreco.2017.11.027

RStudio Team, 2019. RStudio: Integrated Development Environment for R. Rstudio, Inc., Boston, MA.
V.1.2.5033. https://www.rstudio.com/.

Scheibe, A., Steffens, C., Seven, J., Jacob, A., Hertel, D., Leuschner, C., Gleixner, G., 2015. Effects of tree
identity dominate over tree diversity on the soil microbial community structure. Soil Biology and
Biochemistry 81, 219–227. doi:10.1016/j.soilbio.2014.11.020

Scheu, S., 1992. Automated measurement of the respiratory response of soil microcompartments: Active
microbial biomass in earthworm faeces. Soil Biology and Biochemistry 24, 1113–1118.
doi:10.1016/0038-0717(92)90061-2

Scheu, S., Albers, D., Alphei, J., Buryn, R., Klages, U., Migge, S., Platner, C., Salamon, J.-A., 2003. The soil
fauna community in pure and mixed stands of beech and spruce of different age: trophic structure and
structuring forces. Oikos 101, 225–238. doi:DOI 10.1034/j.1600-0706.2003.12131.x

Schmid, M., Pautasso, M., Holdenrieder, O., 2014. Ecological consequences of Douglas fir (Pseudotsuga
menziesii) cultivation in Europe. European Journal of Forest Research 133, 13–29. doi:10.1007/s10342-
013-0745-7

Schmidt, M.W.I., Torn, M.S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I.A., Kleber, M., Kögel-
Knabner, I., Lehmann, J., Manning, D.A.C., Nannipieri, P., Rasse, D.P., Weiner, S., Trumbore, S.E., 2011.
Persistence of soil organic matter as an ecosystem property. Nature. doi:10.1038/nature10386

Toïgo, M., Vallet, P., Perot, T., Bontemps, J.D., Piedallu, C., Courbaud, B., 2015. Overyielding in mixed forests
decreases with site productivity. Journal of Ecology 103, 502–512. doi:10.1111/1365-2745.12353

van der Heijden, M.G.A., Bardgett, R.D., Van Straalen, N.M., 2008. The unseen majority: Soil microbes as
drivers of plant diversity and productivity in terrestrial ecosystems. Ecology Letters 11, 296–310.
doi:10.1111/j.1461-0248.2007.01139.x

Waller, L.P., Allen, W.J., Barratt, B.I.P., Condron, L.M., França, F.M., Hunt, J.E., Koele, N., Orwin, K.H., Steel,
G.S., Tylianakis, J.M., Wakelin, S.A., Dickie, I.A., 2020. Biotic interactions drive ecosystem responses to
exotic plant invaders. Science 368, 967–972. doi:10.1126/science.aba2225

Wang, H., Liu, S.R., Wang, J.X., Shi, Z.M., Xu, J., Hong, P.Z., Ming, A.G., Yu, H.L., Chen, L., Lu, L.H., Cai, D.X.,
2016. Differential effects of conifer and broadleaf litter inputs on soil organic carbon chemical
composition through altered soil microbial community composition. Scientific Reports 6, 1–9.
doi:10.1038/srep27097

Wardle, D.A., Bardgett, R.D., Klironomos, J.N., Setälä, H., Van Der Putten, W.H., Wall, D.H., 2004. Ecological
linkages between aboveground and belowground biota. Science. doi:10.1126/science.1094875

Yan, B., Ji, Z., Fan, B., Wang, X., He, G., Shi, L., Liu, G., 2016. Plants adapted to nutrient limitation allocate less
biomass into stems in an arid-hot grassland. New Phytologist 211, 1232–1240. doi:10.1111/nph.13970

Zelles, L., 1999. Fatty acid patterns of phospholipids and lipopolysaccharides in the characterisation of
microbial communities in soil: A review. Biology and Fertility of Soils 29, 111–129.
doi:10.1007/s003740050533

Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A., Smith, G.M., 2009. Dealing with heterogeneity, in: Mixed
Effects Models and Extensions in Ecology with R. Springer Science & Business Media, pp. 71–100.
doi:10.1007/978-0-387-87458-6_4

Figures and Tables

Figure 1. Effects of conifers and conifer-beech mixtures (Douglas-fir [Do], Douglas-fir with beech [Do/Be], Norway spruce [Sp] and Norway spruce with beech [Sp/Be]) on microbial basal respiration ($\mu g\ O_2\ g^{-1}\ C\ h^{-1}$), microbial biomass ($\mu g\ C_{mic}\ g^{-1}\ C$) and microbial specific respiration ($\mu g\ O_2\ \mu g^{-1}\ C_{mic}\ h^{-1}$) at nutrient-poor and nutrient-rich sites. Effect sizes are given as back transformed log response ratios compared to beech forests [ln (value in coniferous or mixed forest types / values in beech)]. Effect sizes were averaged across three depths (litter, 0–5 and 5–10 cm soil). Values smaller than 1 indicate higher values in beech. Asterisks indicate significant effects ($p < 0.05$). Bars represent 95% confidence intervals (n=12).

Figure 2. Effects of conifers and conifer-beech mixtures (Douglas-fir [Do], Douglas-fir with beech [Do/Be], Norway spruce [Sp] and Norway spruce with beech [Sp/Be]) on stress indicators: ratio of cyclopropyl PLFAs to its monoenoic precursors, ratio of saturated to monounsaturated PLFAs, and ratio of Gram$^+$ to Gram$^-$ bacteria (cy/pre, sat/mono and Gram$^+$/Gram$^-$, respectively) at nutrient-poor and nutrient-rich sites. Effect sizes are given as back transformed log response ratios compared to beech forests [ln (value in coniferous or mixed forest types / values in beech)]. Effect sizes were averaged across three depths (litter, 0–5 and 5–10 cm soil). Values larger than 1 indicate lower values in beech. Asterisks indicate significant effects ($p < 0.05$). Bars represent 95% confidence intervals (n=12).

Figure 3. Changes in microbial basal respiration ($\mu g\ O_2\ g^{-1}\ C\ h^{-1}$), microbial biomass ($\mu g\ C_{mic}\ g^{-1}\ C$) and microbial specific respiration ($\mu g\ O_2\ \mu g^{-1}\ C_{mic}\ h^{-1}$) with soil depth (litter and 0–5 and 5–10 cm soil) in five forest types (European beech [Be], Douglas-fir [Do], Douglas-fir with beech [Do/Be], Norway spruce [Sp] and Norway spruce with beech [Sp/Be]) at nutrient-poor and nutrient-rich sites. Points and horizontal bars represent means and standard errors (n=4). The grey vertical bars represent respective values in beech forests at nutrient-poor sites in litter, 0–5 and 5–10 cm soil depth. Note log scale for microbial biomass.

Figure 4. Changes in stress indicators (ratio of cyclopropyl fatty acids to its precursors, ratio of saturated to monounsaturated fatty acids and ratio of Gram$^+$ to Gram$^-$ bacteria [cy/pre, sat/mono and Gram$^+$/Gram$^-$, respectively]) with soil depth (litter and 0–5 and 5–10 cm soil) in five forest types (European beech [Be], Douglas-fir [Do], Douglas-fir with beech [Do/Be], Norway spruce [Sp] and Norway spruce with beech [Sp/Be]) at nutrient-poor and nutrient-rich sites. Points and horizontal bars represent means and standard errors (n=4). The grey vertical bars represent respective values in beech forests at nutrient-poor sites in litter, 0–5 and 5–10 cm depth.

Figure 5. Principal component analysis of the microbial community structure (as indicated by phospholipid-derived fatty acids) in five forest types (European beech [Be], Douglas-fir [Do], Douglas-fir with beech [Do/Be], Norway spruce [Sp] and Norway spruce with beech [Sp/Be]) at nutrient-poor and nutrient-rich sites in litter, 0–5 and 5–10 cm soil depth. Positions of forest types represent centroids (n=4). Nutrient-poor sites are given in red and nutrient-rich sites in green. C/N and pH were post-fitted into the plot.

724   Figure 1

727   Figure 2

730   Figure 3

733   Figure 4

736   Figure 5

Table 1. F- and P-values of linear mixed effects models on the effect of Forest type (European beech, Douglas-fir, Norway spruce, mixture of European beech with Douglas-fir and mixture of European beech with Norway spruce), Quintet group (nutrient-poor and nutrient-rich sites) and Depth (litter, 0–5 and 5–10 cm) on (a) basal respiration, microbial biomass and microbial specific respiration, (b) stress indicators (cy/pre, sat/mono, $Gram^+/Gram^-$ ratios) and (c) percentages of fungal, bacteria, and fungi/bacteria ratio. Significant effects are given in bold (p < 0.05).

(a)	Basal respiration			Microbial biomass			Specific respiration		
Factor	df	F	P	df	F	P	df	F	P
Forest type (F)	4,24	7.51	**<0.001**	4,30	5.95	**0.001**	4,24	0.92	0.47
Quintet group (Q)	1,6	3.27	0.12	1,30	16.93	**<0.001**	1,6	0.52	0.49
Depth (D)	2,78	181.58	**<0.001**	2,70	271.27	**<0.001**	2,78	17.37	**<0.001**
F × Q	4,24	8.83	**<0.001**	4,30	4.61	**0.005**	4,24	2.36	0.08
F × D	-	-	**-**	8,70	2.15	**0.042**	-	-	-

(b)	cy/pre ratio			sat/mono ratio			$Gram^+/Gram^-$ ratio		
Factor	df	F	P	df	F	P	df	F	P
Forest type (F)	4,24	2.75	0.052	4,24	8.278	**<0.001**	4,24	3.39	**<0.001**
Quintet group (Q)	1,6	3.48	0.11	1,6	0.195	0.67	1,6	8.17	**0.030**
Depth (D)	2,68	238.82	**<0.001**	2,76	5.551	**0.006**	2,76	228.51	**<0.001**
F × Q	4,24	2.70	0.055	4,24	3.650	**0.019**	4,24	2.27	**0.012**
Q × D	2,68	6.20	**0.003**	2,76	10.71	**<0.001**	2,76	7.51	**0.001**
F × D	8,68	3.72	**0.001**	-	-	-	-	-	-

(c)	% Fungi			% Bacteria			Fungi/Bacteria ratio		
Factor	df	F	P	df	F	P	df	F	P
Forest type (F)	4,30	1.77	0.16	4,30	2.19	0.094	4,30	0.83	0.52
Quintet group (Q)	1,30	6.46	**0.02**	1,30	4.34	**0.046**	1,30	4.98	**0.033**
Depth (D)	2,68	982.11	**<0.001**	2,78	149.47	**<0.001**	2,68	995.4	**<0.001**
F × Q	4,30	1.99	0.12	4,30	0.733	0.58	4,30	1.60	0.20
Q × D	2,68	3.61	**0.002**	-	-	-	2,68	16.92	**<0.001**
F × D	8,68	18.54	**<0.001**	-	-	-	8,68	2.84	**0.009**

Table 2. Results of (a) two-way MANOVAs and (b) MANCOVA of PLFA composition in litter, 0–5 and 5–10 cm
soil. Factors include Forest type (European beech, Douglas-fir, Norway spruce and mixture of European
beech with Douglas-fir and mixture of European beech with Norway spruce), Quintet group (nutrient-rich
and nutrient-poor sites) and their interactions. MANOVA was based on principal components from each
sample depth (significant PCs were determined by broken stick criterion; see Methods). Type I sum of square
was used for models without soil parameters, and type II sum of squares for those with soil parameters.
Significant effects are given in bold ($p < 0.05$).

Soil depths	Factors	(a) Without soil parameters				(b) With soil parameters			
		df	Wilk's lambda	F	P	df	Wilk's lambda	F	P
Litter	Forest type (F)	4,30	0.32	3.29	**0.001**	4,28	0.42	2.23	**0.019**
	Quintet group (Q)	1,30	0.42	12.77	**<0.001**	1,28	0.61	5.57	**0.004**
	F × Q	4,30	0.73	0.77	0.677	4,28	0.74	0.69	0.757
	pH	-	-	-	-	1,28	0.75	2.85	0.057
	C/N	-	-	-	-	1,28	0.82	1.95	0.146
0–5 cm	Forest type (F)	4,30	0.19	2.80	**<0.001**	4,28	0.34	1.57	0.081
	Quintet group (Q)	1,30	0.39	8.14	**<0.001**	1,28	0.80	1.18	0.350
	F × Q	4,30	0.36	1.57	0.078	4,28	0.31	1.68	0.054
	pH	-	-	-	-	1,28	0.47	5.41	**0.002**
	C/N	-	-	-	-	1,28	0.66	2.44	0.063
5–10 cm	Forest type (F)	4,30	0.41	1.76	0.051	4,28	0.45	1.45	0.142
	Quintet group (Q)	1,30	0.45	8.15	**<0.001**	1,28	0.82	1.34	0.285
	F × Q	4,30	0.69	0.68	0.805	4,28	0.74	0.51	0.936
	pH	-	-	-	-	1,28	0.84	1.22	0.326
	C/N	-	-	-	-	1,28	0.75	2.08	0.114