Article

Effects of early life stress on bone homeostasis in mice and humans

Karin Wuertz-Kozak[1,2,3]▲*, Martin Roszkowski[3,4]▲, Elena Cambria[3], Andrea Block[12], Gisela A. Kuhn[3], Thea Abele[6], Wolfgang Hitzl[7,8,9], David Drießlein[10], Ralph Müller[3], Michael A. Rapp[5], Isabelle M. Mansuy[3,4]▼, Eva M. J. Peters[6,11]▼, Pia M. Wippert[3,5,12]▼

[1] Department of Biomedical Engineering, Rochester Institute of Technology (RIT), Rochester (NY), USA; kwbme@rit.edu

[2] Schön Clinic Munich Harlaching, Spine Center, Academic Teaching Hospital and Spine Research Institute of the Paracelsus Medical University Salzburg (AU), Munich, Germany; kwuertz@schoen-kliniken.de

[3] Department of Health Sciences and Technology, ETH Zurich, Zurich, Switzerland; kwuertz@ethz.ch; martin.roszkowski@hest.ethz.ch; mansuy@hifo.uzh.ch; wippert@uni-potsdam.de; gakuhn@ethz.ch; ram@ethz.ch; elena.cambria@hest.ethz.ch

[4] Laboratory of Neuroepigenetics, University of Zurich and Swiss Federal Institute of Technology, Brain Research Institute, Neuroscience Center Zürich, 8057 Zurich, Switzerland; martin.roszkowski@hest.ethz.ch; mansuy@hifo.uzh.ch

[5] Faculty of Health Brandenburg, University of Potsdam, Potsdam, Germany; wippert@uni-potsdam.de; michael.rapp@uni-potsdam.de

[6] Psychoneuroimmunology Laboratory, Department of Psychosomatics and Psychotherapy, University of Gießen, Gießen, Germany; thea-abele@gmx.de; eva.peters@eva-peters.com

[7.] Research Office (Biostatistics), Paracelsus Medical University, Salzburg, Austria; wolfgang.hitzl@pmu.ac.at

[8.] Research Program Experimental Ophthalmology and Glaucoma Research, Paracelsus Medical University, Salzburg, Austria; wolfgang.hitzl@pmu.ac.at

[9.] Paracelsus Medical University Salzburg, Department of Ophthalmology and Optometry, Austria; wolfgang.hitzl@pmu.ac.at

[10] Statistical Consulting Unit StaBLab, Ludwig-Maximilians-University Munich, Munich, Germany; david.driesslein@gmail.com

[11] Charité Center 12 Internal Medicine and Dermatology, Division for General Internal Medicine, Psychosomatics and Psychotherapy, Universitätsmedizin Berlin, Berlin, Germany; eva.peters@eva-peters.com

[12.] Sociology of Health and Physical Activity, University of Potsdam, Potsdam, Germany; wippert@uni-potsdam.de; anblock@uni-potsdam.de

* Correspondence: kwbme@rit.edu; Tel.: 001 585 475-7355

Abstract: Bone pathology is frequent in stressed individuals. A comprehensive examination of mechanisms linking life stress, depression and disturbed bone homeostasis is missing. In this translational study, mice exposed to early life stress (MSUS) were examined on bone microarchitecture (μCT), metabolism (qPCR), and neuronal stress mediator expression (qPCR) and contrasted with a sample of depressive patients with or without early life stress by analyzing bone mineral density [BMD] (DXA) and metabolic changes in serum (osteocalcin, PINP, CTX-I). MSUS mice showed a significant decrease in NGF, NPYR1, VIPR1 and TACR1 expression, higher innervation density in bone and increased serum levels of CTX-I, suggesting a milieu in favor of catabolic bone turnover. MSUS mice had a significantly lower body weight compared to control mice, and this caused minor effects on bone micro-CT. Depressive patients with experiences of childhood neglect also showed a catabolic pattern. A significant reduction in BMD was observed in depressive patients with childhood abuse and stressful life events during childhood. Therefore, future studies on prevention and treatment strategies for both mental and bone disease should consider early life stress as a risk factor for bone pathologies.

Keywords: Psychosocial stress; bone pathologies; osteoporosis; bone mineral density; childhood; neuroendocrine

1. Introduction

Bones are essential components of the musculoskeletal system and subjected to continuous remodeling as an adaptation mechanism to environmental changes. Disturbances in bone development and remodeling by formation/resorption of the extracellular matrix (ECM) or differentiation of osteoblasts into osteocytes and apoptosis of osteocytes could result in reduced bone mass and increased fracture risk.

Beside well-documented mechanisms like menopause-associated hormonal changes, aging-related factors, changes in physical activity, as well as drugs and diseases, recent research suggested that psychosocial stress and stress-linked neuronal processes are associated with an increased risk for bone pathologies [1-5]. Autonomic and sensory nerve fibers densely innervate bone, predominantly in the periosteum and in areas with high metabolic activity and bone turnover. Neuronal structures can make direct contact with osteoblasts and osteoclasts, which in turn express receptors for neuronal markers on their surface and can produce neuronal mediators in an auto- and paracrine fashion [6-10]. Importantly, sympathetic and peptidergic neuronal factors can act as stress mediators and appear to play a key role in bone turnover [11-13]. Of specific relevance are tyrosine hydroxylase (TH), a rate-limiting enzyme for noradrenaline synthesis, neuropeptide Y (NPY), often co-expressed with noradrenaline, vasoactive intestinal peptide (VIP), often co-expressed with acetylcholine, and the neuropeptides calcitonin gene related peptide (CGRP) and substance P (SP) that are expressed in sensory nerve fibers. In addition, neurotrophic growth factors such as nerve growth factor (NGF) and brain derived neurotrophic factor (BDNF) that guide outgrowing nerve fibers to bony areas requiring innervation are of interest. Stress alters their expression, and they can also act as direct and indirect growth factors for osteoblasts [14-26]. Latest studies indicate that psychosocial stress can lead to structural and functional changes in neuronal plasticity, neuronal marker expression, mitochondria and inflammation [4, 27, 28], possibly resulting in downstream alterations of bone homeostasis [29-33].

In humans, major early life stress like childhood maltreatment strongly affects the stress
response, as well as behavioral and cognitive functions throughout life [34, 35]. Strong associations
were shown between childhood maltreatment and higher risk for psychological and physiological
disorders including depression, hypersensitivity of the neuroendocrine, autonomic and peptidergic
stress response leading to altered dynamics of the hypothalamic pituitary adrenal (HPA) axis, and
changes in the neuronal mediators that can influence bone health [36]. Childhood maltreatment is
one of the best-documented risk factors for depression [37-40] and depression in turn has been
associated with a higher risk for reduced bone mineral density (BMD), osteoporosis and bone
fractures [41-45].

However, existing knowledge about underlying mechanisms between early life stress,
neurogenic stress response elements, and detrimental changes in bone metabolism and
microstructure/BMD has been limited by several reasons. First, although early life stress like
childhood maltreatment is a frequent phenomenon, it is hardly accessible to retrospective and
longitudinal studies due to its rare diagnosis and ethical requirements for case-control studies.
Second, access to biological material and strict control of confounding factors, such as physical
activity and eating habits, are challenging in human studies [46, 47]. Animal models offer an obvious
way to investigate major stress experiences early in life despite these obstacles. Animal models are
helpful counterparts of human studies to investigate underlying mechanisms in a highly controlled
environment that allows extensive tissue sampling, strict definition of adverse events and
confounding factors, and lower subject variability.

Unpredictable maternal separation and maternal stress during early life (MSUS paradigm [48])
is an experimental stress paradigm in rodents that can mimic aspects of childhood maltreatment and
induce long-lasting health effects, such as increased depression-like behavior (anhedonia) and altered
brain activity in adulthood [49]. The MSUS paradigm severely impacts physiology and behavior in
the exposed animals but also in their descendants across several generations via mechanisms
involving epigenetic factors in the germline [48, 50-52].

To address the knowledge gap between early life stress, stress-dependent neurogenic markers
that mark neuronal plasticity and detrimental changes in bone metabolism and bone
microstructure/BMD, we first examined the impact of early life stress on neuronal mediator
expression (qPCR) and bone microarchitecture (μCT), metabolism (qPCR on bone, ELISA on serum
for osteocalcin OC, PINP, CTX-I), and in mice using the MSUS paradigm (**Aim 1**). Then we translated
the findings to humans by analyzing metabolic changes in serum (OC, PINP, CTX-I) and
microarchitecture (DXA) in a sample of depressive patients with or without retrospectively reported
experiences of early life stress, like childhood maltreatment and stressful life events during childhood
(**Aim 2**).

2. Results

2.1. Mouse Study

2.1.1. Bone innervation and neuronal mediators

We first assessed the effects of early life stress on neuronal signaling and neurogenic factors in
bone metabolism by measuring neuronal density and expression patterns of neurogenic markers in
the bone of MSUS and control mice (*n* = 6-8 each).

To study changes in neuronal marker gene expression, qPCR was performed. Gene expression
analysis revealed that key receptors for nerve growth factors, neurotransmitters and neuropeptides
(***Figure 1A***) and selected receptors ligands (***Figure 1B***) were changed in MSUS mice in a pattern that
may promote higher bone turnover and lower bone density. In detail, mRNA expression of the
receptors NPYR1 ($p < 0.05$, ***Figure 1A***), VIPR1 ($p < 0.001$, ***Figure 1A***) and TACR1 ($p < 0.01$, ***Figure 1A***)
were significantly reduced in MSUS mice whole bone homogenates (after removal of bone marrow),
suggesting their presence on osteocytes. These three receptors are involved in the regulation of
proliferation and differentiation in bone and their concomitant downregulation may lead to higher
remodelling activity with less proliferation and more differentiation of bone cells. Furthermore, a
significant reduction of the neurotrophin NGF ($p < 0.001$, ***Figure 1B***) was revealed, while NPY showed
a downtrend in MSUS ($p = 0.068$, ***Figure 1B***). This would hamper both neuronal plasticity and bone
remodelling in the face of a challenge such as a wound or aging leading to osteoporosis. Other
markers of bone innervation and metabolism such as NGFR, ADRB2, CHRNA7, RAMP1, and TAC1
were similar in control and MSUS mice, whereas TRKA, TRKB, TH, ChAT and VIP were below the
detection limit.

Figure 1A: Differential gene expression of neuronal receptors in bone between control and MSUS mice. *Shown are*

the gene expression of neurotrophin receptors (A), noradrenergic receptors (B), cholinergic receptors (C) and receptors
involved in sensory innervation (D). Data (n = 8 per group) are expressed as Min to Max of 2^{-dCt}. Significance level: ***p <
0.001, **p < 0.01 and *p < 0.05 between indicated groups.

Figure 1B: Differential gene expression of neuronal ligands in bone between control and MSUS mice. *Shown are*
the gene expression of neurotrophins (A), noradrenergic ligands (B), cholinergic ligands (C) and ligands involved in sensory
innervation (D). Data (n = 8 per group) are expressed as Min to Max of 2^{-dCt}. Significance level: ***p < 0.001 between
indicated groups.

We subsequently investigated bone innervation by immunohistomorphometry. The density of
nerve fibers labeled with the pan-neuronal marker PGP 9.5 was significantly increased in the bone of
MSUS mice (***Figure 2***), whereas nerve fibers labeled with the neuronal plasticity marker GAP 43 were
unaffected (not shown), indicating that higher innervation was not due to increased neuronal
plasticity, but was likely a stable feature. Higher innervation density potentially leads to higher
sensitivity of bone neurons to stimuli like injury. Noteworthy, the neuropeptides NPY, VIP and SP
were undetectable by immunohistochemistry, most likely due to insufficient bone preservation. This
likely results from the post-mortem fixation method used for ethical requirements, which does not

preserve small peptides (the eleven amino-acid sequence of SP for example is rapidly digested by
endogenous enzymes after termination of circulation) [53-55].

Figure 2: Increased bone innervation in MSUS mice. Representative images of the third tail bone from control and
MSUS mice labeled with PGP 9.5 pan-neuronal marker for immunofluorescence histomorphometry of nerve fibers (NF -
red) in bone. Nuclei are counterstained with DAPI (blue), mast cells with FITC-avidin (green) (A). Shown are the increased
mean numbers of immune-positive profiles per compartment unit of single NF in bone (B), NF bundles at entry to bone (C)
and single NF in bone marrow (D). Data (n = 8 Control, n = 6-7 MSUS) are expressed as Min to Max of number of positive
*profiles per compartment unit. Significance level: **p < 0.01 and *p < 0.05 between indicated groups.*

2.1.2. Bone metabolic parameters

We then examined if changes in neuronal markers were associated with bone metabolic changes
by measuring markers in serum and bone. Serum levels of osteocalcin and PINP, two markers
associated with bone formation, did not change after MSUS (*Figure 3A-B*). In contrast, the level of
CTX-I, which is associated with bone resorption, was significantly increased in MSUS mice compared
to control mice (p < 0.01, n=14 each; *Figure 3C*).

Figure 3: Bone formation and resorption markers in serum of mice. Shown are serum concentration measurements of
osteocalcin (A), procollagen type 1 N-terminal propeptide (PINP) (B) and c-terminal telopeptide of type I collagen (CTX-
I) (C) in mice. Data (n=14 per group) are expressed as Min to Max of concentration in serum [ng/ml]. Significance level:
***p < 0.01 between indicated groups.*

However, this catabolic shift was not reflected by altered gene expression of tissue-specific
extracellular matrix markers (osteocalcin, osteoprotegerin, osteopontin, sclerostin) in bone samples
of MSUS (*Figure 4A-D*) (*n=8* each).

Figure 4: **Gene expression of extracellular matrix in bone of mice.** *Shown are the gene expressions of tissue-specific extracellular matrix markers in bone of controls and MSUS. Genes include osteocalcin **(A)**, osteoprotegerin **(B)**, osteopontin **(C)** and sclerostin **(D)**. Data (n=8 per group) are expressed as Min to Max of 2^{-dCt}. Significance level: all p > 0.05*

2.1.3. Bone microarchitecture

Next, it was examined if MSUS affects bone microarchitecture in mice using micro-CT (n=14 MSUS, n=13 Control). As bone microarchitecture is sensitive to mechanical loading, we also measured body weight of experimental animals over time. Body weight was significantly lower in MSUS mice compared to control mice starting at postnatal day 7 (PND7) and remained lower in adulthood (p< 0.001 for PND 7-28 and p < 0.05 for 10 months, **Supplementary Figure 1**), and this caused minor effects on bone microarchitecture.

After adjusting micro-CT data for body weight, a comparison of means between control and MSUS mice did not reveal any difference in full, cortical or trabecular microarchitecture (all values p > 0.05) (**Supplementary Table 1**). We also used blocking in tertiles for body weight and conducted a two-factorial multivariate analysis of variance, which revealed that indeed the effects of MSUS versus controls were driven by body weight. To specify this effect, we conducted sets of linear regression analyses to test for simple mediation effects and found that body weight mediated the effect of MSUS on full length, TV, BV, MV and BS (all p's for direct effects > 0.05).

2.2. Human Study

2.2.1. Descriptive results

N=193 out of *n*=208 patients under study suffered from depression with minimal (35%) up to
moderate/ severe (65%) severity (BDI-II: *M* = 23.8, *SD* = 10.7) and were included in further analysis.
For the analysis of bone metabolic parameters, blood samples from n=145 patients were available, of
whom *n* = 145 filled in the *Inventory of Stressful Life Events* (ILE) and *n*=54 the *Childhood Trauma Screener*
(CTS). For the BMD analysis, n=17 patients took part in DXA bone densitometry measurements. Out
of these participants, *n*=15 patients filled in the *Inventory of Stressful Life Events* (ILE) and *n*=17 the
Childhood Trauma Screener (CTS). Missing data is due to incomplete questionnaires or drop out. Table
1 provides descriptive data on age, gender, antidepressant use, weight and early life stress for the
subsamples. Average bone mineral density (BMD) in the sample was *M*= 0.942 g/cm^2 (*SD* = 0.186
g/cm^2) in the trochanter region of the proximal femur.

Table 1. *Descriptive data on sociodemographic and clinical characteristics of the human sample*

	N	age		depression (BDI-II score)		weight		sex		Antidepressant use		early life stress			
												ILE $_{child}$ [c]		CTS [d]	
								N		N		N		N	
		M	(SD)	M	(SD)	M	(SD)	(female)	(%)	(yes)	(%)	(yes)	(%)	(yes)	(%)
Sample of bone metabolic parameters [a]	**145**	47.9	(10.1)	24.0	(10.1)	78.1	(17.4)	118	(81.4)	100	(69.0)	128	(88.3)	47	(32.4)
➥ *life events during childhood (ILE $_{child}$)*	128	48.1	(10.1)	24.0	(10.0)	78.2	(17.3)	104	(81.3)	86	(67.2)				
➥ *childhood maltreatment (CTS)*	47	47.9	(10.4)	24.6	(10.7)	75.8	(17.3)	35	(74.5)	32	(68.1)				
DXA sample [b]	**17**	51.9	(6.6)	23.7	(11.7)	74.3	(17.5)	12	(70.6)	13	(76.5)	15	(88.2)	17	(100.0)
➥ *life events during childhood (ILE $_{child}$)*	15	52.2	(7.0)	24.9	(12.2)	71.2	(16.3)	11	(73.3)	11	(73.3)				
➥ *childhood maltreatment (CTS)*	17	51.9	(6.6)	23.7	(11.7)	74.3	(17.5)	12	(70.6)	13	(76.5)				

[a] *bone serum markers OC, PINP, CTX-I*

[b] *bone mineral density (BMD) measured by Dual Energy X-ray Absorptiometry (DXA)*

[c] *Inventory of Stressful Life Events; stressful life events during childhood (≤12 years)*

[d] *Childhood Trauma Screener (CTS) at least at risk for one type of neglect or abuse (sexual, physical, emotional abuse, physical, emotional neglect)*

2.2.2. Bone metabolic parameters

Regarding changes in human bone metabolism following early life stress, bone serum markers
OC, PINP, CTX-I were analyzed. It was shown that depressed patients with childhood trauma (CTS,
CTS abuse) had an increased anabolic bone metabolism in comparison to depressed patients without
early life stress (CTS: *p* = 0.05* for CTX; CTS abuse: *p* = 0.03* for P1PN, **Table 2**). This anabolic bone

metabolism was likewise found for depressed patients with stressful life events during childhood
(ILE child) in comparison to depressed patients without early life stress (*p* = 0.04* for P1PN, *Table 2*).
Depressed patients with experiences of childhood neglect (CTS neglect) lack such anabolic reaction

(*Table 2*).

2.2.3. Bone mineral density
An explorative analysis on bone mineral density in a subsample of DXA measurements suggests
that depressive patients *with* stressful life events during childhood (ILE child) had reduced BMD in the
trochanter major (p = 0.03* for BMD_T) and as a tendency a reduction in general BMD (p = 0.08# for
BMD_G, *Table 2*). There are indications that depressed patients with a history of childhood abuse
showed specifically a BMD reduction in the trochanter major (p = 0.05* for BMD_T), and a tendency
in reduced BMD in general (p = 0.07# for BMD_G), giving notice of long-term health consequences of
people with maltreatment and stressful life events during childhood in comparison to people with
depression lacking experiences of early life stress (**Table 2, Figure 5**).

***Figure 5**: **Bone mineral density measurements in human patients.** Shown is the BMD in G (total), H (collum femoris),*
S (corpus femoris), and T (trochanter major) as mean of the right and left site and L1-L4 (lumbar vertebralis bodies) stratified
for at risk and not at risk for childhood maltreatment (not controlled for gender, age, body weight). Data (n=18) are expressed
as mean and standard deviation of BMD, stratified for CTS (at risk for abuse or neglect). Significance level: *p < 0.05, and
#p < 0.1 between indicated groups.

Table 2. *Main effects (regression coefficient b) of childhood maltreatment (abuse or neglect) and stressful life events during childhood (≤12 years; ILE child) on bone serum marker (osteocalcin (OC),*
procollagen type 1 N-terminal propeptide (PINP), and C-terminal telopeptide of type I collagen (CTX-I)) and bone mineral density (BMD)

	P1NP			OC			CTX-I				
	b	95%CI	p	b	95%CI	p	b	95%CI	p		
CTS [a]	9.27	-1.43; 19.97	0.10[#]	2.87	-0.56; 6.30	0.11	**0.10**	0.01; 0.18	0.05*		
CTS abuse	**11.42**	1.86; 20.99	0.03*	2.51	-0.65; 5.67	0.13	0.06	0.02; 0.14	0.16		
CTS neglect	-6.26	-16.45; 3.93	0.24	0.32	-3.63; 2.98	0.85	0.03	-0.06; 0.12	0.50		
ILE child [b]	**8.37**	0.71; 16.03	0.04*	2.28	-0.10; 4.65	0.06[#]	0.06	-0.01; 0.11	0.06[#]		

	BMD_L1L4 [c]			BMD_T			BMD_S			BMD_G			BMD_H		
	b	95%CI	p	b	95%CI	p	b	95%CI	p	b	95%CI	p	b	95%CI	p
CTS	-0.07	-0.23; 0.08	0.40	-0.09	-0.22; 0.04	0.20	-0.09	-0.25; 0.08	0.33	-0.10	-0.23; 0.03	0.17	-0.10	-0.22; 0.02	0.12
CTS abuse	-0.07	-0.26, 0.12	0.48	**-0.15**	-0.28; -0.02	0.05*	-0.13	-0.32, 0.06	0.22	-0.15	-0.29, -0.01	0.07[#]	-0.12	-0.26, 0.01	0.10
CTS neglect	-0.06	-0.21, 0.10	0.51	-0.06	-0.20, 0.07	0.39	-0.07	-0.24, 0.10	0.45	-0.07	-0.21, 0.07	0.38	-0.03	-0.16, 0.10	0.68
ILE child	-0.18	-0.46, 0.10	0.25	**-0.26**	-0.44, -0.07	0.03*	-0.18	-0.48, 0.11	0.26	-0.22	-0.43, -0.01	0.08[#]	-0.17	-0.36, 0.03	0.14

Multiple regression models, adjusted for age, gender, and study sides. Significant Regression coefficients b are bold. Significance level: *p < 0.05, and #p < 0.1.

[a] Childhood Trauma Screener (CTS) at least at risk for one type of neglect or abuse (sexual, physical, emotional abuse, physical, emotional neglect)

[b] Inventory of Stressful Life Events; stressful life events during childhood (≤12 years)

[c] Bone mineral density (BMD) was measured as mean value (M) in the lumbar spine L1-L4 (lumbar vertebralis bodies), T (trochanter major), S (corpus femoris), G (total), and H (collum femoris)

3. Discussion

The first aim of this study was to investigate the effects of early life stress on bone health and associated biological factors in a mouse model of early life stress (MSUS). The second study aim was the translation of the gained knowledge in a human model using a sample of depressive patients with and without early life stress.

Aim 1: In summary, we found that early life stress in the MSUS model altered bone innervation and neuronal mediator expression as well as bone metabolism, as evidenced by increased CTX-I serum concentrations, but did not affect expression of bone markers within the bone or bone microarchitecture. In detail, we found significantly decreased expression of NGF, NPYR1, VIPR1 and TACR1 in MSUS bones. This response pattern is suggestive for a higher rate of bone turnover and lower bone healing capacities as a consequence of early life stress, suggesting a concurrent joint downregulation, reduced bone remodeling and long-term destabilization. The neurotrophin NGF is important for the expression of cholinergic and sensory neuronal factors and hence for the development and the repair of bone, as demonstrated by reduced bone length when absent [14-26]. The neuropeptide NPY promotes osteoblast differentiation and balances the effects of stress-induced bone-loss through beta-adrenergic stimulation, both with consequences for bone remodeling during healing [10, 53, 56-63]. The neuropeptide VIPR1, finally, promotes bone mineralization [64-68] while TACR1 regulates osteoblasts, osteoclasts and mesenchymal stem cell functionality associated with protection from osteoporosis [2, 62, 69-85].

While MSUS mice showed a clear neuroendocrine response pattern, gene expression of typical bone markers (in bone) was unaffected. Similarly, we did not observe increased osteocalcin in the serum of MSUS mice. However, CTX-I serum levels were significantly increased in MSUS mice, which is an increasingly promoted and robust biochemical factor for bone resorption in clinical settings and further giving notice about correlations to bone mineral density [86-89]. Additional analyses, such as bone histomorphometry to determine whether MSUS leads to alterations in the numbers of osteoblasts (type 1 procollagen staining) and/or osteoclasts (TRAP staining) [90] could provide useful insights into bone adaptation processes.

Regarding bone microarchitecture, no significant changes were observed after normalization to body weight. Body weight itself, which was affected by MSUS, did cause minor changes in bone microarchitecture, likely due to differences in mechanical loading of the skeleton. It is unclear, if altered body weight might be a post stress consequence for MSUS or an incidental finding. This should be considered in future studies, as it can be a confounding factor in micro-CT analyses. To test whether the observed lack of differences in bone microarchitecture (despite an increase in serum CTX-I) might be due to divergent timelines of these events [91] or might be sex-dependent (different susceptibility to bone alterations and/or sensitivity to stress) [92, 93]. We additionally conducted micro-CT scans of the femurs of female mice over a more extended time range (up to 16 months). We did not find any differences in bone microarchitecture in the female mice, even not at later time points (data not shown). It is important to keep in mind that changes in bone microarchitecture may be site specific and may hence have been overlooked by restricting the micro-CT analysis to the femur only.

Perhaps, the use of models of aging (e.g., the senescence-accelerated mouse = SAM) [94], models that
include bone injury requiring repair (such as osteotomy) [95], or a comparison of stress-resilient and
stress-prone subpopulations [96] might potentially have led to observable changes in bone structure
within a reasonable time frame [97], but further investigations will be needed to allow for more
definite statements.

Aim 2: In summary, we found that people that experienced childhood maltreatment or stressful
life events during childhood showed an overshooting anabolic bone metabolism during a depressive
episode, rather than the expected minor, short-term metabolic adaptation. For example, people in our
study showed a threefold higher metabolic reaction in PINP during the depressive episode, in
comparison to depressed adults without childhood maltreatment [5]. This was evidenced by
increased serum levels in bone metabolism markers (distinctly in PINP and CTX-I, less in OC).
Further, our data suggests that depressive adults with experiences of childhood maltreatment
(childhood abuse) and stressful life events during childhood showed a significantly decreased BMD
in the long term. In contrast, for depressive adults with no experiences of childhood neglect, these
patterns were not observed. In other words and regarding latest studies: an overshooting or lacking
metabolic adaptation during a depressive episode may serve as a link for the often observed
development of BMD in depressed patients in the long term [5]. These mechanisms fit the hypotheses
of the destructive effects of a hypersensitive central stress response (overshooting) or a blunted
response (lacking adaptation) on organs and bodily tissues.
Other human studies have linked childhood maltreatment to neuroendocrinological and
neuroanatomic alterations that overlap with depressive symptoms and result in a vulnerable
phenotype that is affected by a hypersensitive central stress response, blunted stress reactivity and
lower stress resilience [98-100]. It was suggested that further severe distress, such as additional
traumatization, or the development of a depression, may promote an over-adaptive reaction. The
response pattern in our data seems to confirm this assumption, which can result from various aspects,
such as an altered mitochondrial function, gene dysregulation, accumulation of oxidative stress,
inflammation and senescence or changes in neuronal plasticity [27, 28]. The associations between
stressful events early in life and an early decrease of noradrenergic, cholinergic and sensory neuronal
factors was also observed in our animal model. It is also comprehensible that no such pattern could
be found for childhood neglect, as it represents acts of omissions that are typically temporary, such
as lacks of nutrition, medical or emotional care. Hence, these might have a lower impact on a person's
psychological integrity, identity and health than childhood abuse. Further, the data matches well
with further results within this study sample: patients that suffer from chronic stress (measured by
the allostatic load index) showed a catabolic shift in bone metabolism during a depression episode
[5]. This can be explained by the fact that other mechanisms are expected for allostatic load as an
accumulating damage over time with the consequence of neuroendocrinological and neuroanatomic
alterations [36, 99, 101]. The observed reduced bone mineral density can be expected in both response
pattern: within a hypersensitive stress response and within an increased allostatic load. Both
mechanisms were demonstrated in data of people of the presented DEPREHA-study.

In summary, we found that early life stress affects bone metabolism in mice and humans. In
MSUS mice, this was evidenced by a significant shift towards a neurogenic and osteogenic catabolic
milieu (decreased NGF, NPYR1, VIPR1, TACR1; higher innervation density in bone; increased serum
CTX-I), but no alterations in bone microarchitecture in relatively young mice not exposed to

additional life stresses. In humans however, that suffered both from early life stress and depression
a significant reduction in BMD was observed. Taking the age range and morbidity (depression) of
the human sample under consideration, the differences between mice and human could be
interpreted as a patterns of vulnerability and acute consequences (mouse model) compared to long-
term burden of early life stress (human). Otherwise, species-related differences in bone metabolism
between mice and humans have been described [102]. Our data supports the notion that the
mechanisms of bone loss seem different between mice and humans, as mice did not show any
decrease in trabecular thickness, which is considered as typical for ageing humans [103]. However,
this contrasts Halloran et al. [103], who showed that cancellous bone volume fraction is reduced in
24 month old male C57BL/6J mice and is hence comparable to humans. Furthermore, the observed
differences in the response of mice and humans are likely caused by differences in types of stress
early in life. In fact, we were able to demonstrate clear differences between different types of early
life stress in the human sample, further supporting this theory. Although the results obtained in the
mouse study do not overlap entirely with the human results, it provides relevant and promising
potential for translation. Furthermore, the animal model can be used to identify probable
mechanisms underlying osteoporosis development following early life stress (i.e. related to
innervation patterns) that could not be investigated easily in humans, thereby allowing to develop
ideas for future studies or even intervention and prevention strategies.

In conclusion, this study demonstrates that stressful life events early in life leads to (mal-) adaptive
responses in bone metabolism in both mice and humans. However, the major conclusion to be drawn
from the presented data is that different types of early life stress lead to different response patterns
concerning bone health and have to be consider separately and possibly consecutively: MSUS mice
responded with enhanced bone resorption, which however did not cause changes in bone
microarchitecture 8-10 months after the stress exposure when later kept under stress-free conditions.
In humans, a reduction in BMD was evident only in depressive patients with severe stress exposure
in early life like childhood abuse and stressful life events during childhood. Similarly, the changes in
bone serum markers were also dependent on the severity of stress experiences early in life. People
with childhood abuse and stressful life events during childhood showed an increased bone
metabolism while people with childhood neglect showed a counter regulated reaction. Hence, the
consequences of the observed changes in the context of challenges to bone health (fractures, aging
etc.) should be studied in a more personalized manner and future prevention strategies should
respect the severity of the stress exposure to be treated (e.g. specific type of stress experiences during
childhood, double hit paradigms) to achieve the optimal benefit for patients with increased risk for
bone pathologies, such as people with depressive disorders.

4. Materials and Methods

4.1. Mouse Study

4.1.1. Animal model
Animal experiments were conducted in strict adherence to the Swiss Law for Animal Protection
and were approved by the local authority (Veterinäramt des Kantons Zürich, Switzerland) under
license number 57/2015. C57Bl/6J mice were obtained from Janvier (France) and bred in-house to
generate the male mice (n=47 MSUS, n=46 Control) used for experiments.

The MSUS paradigm was conducted as previously described [48]. As shown in *Figure 6*, 2-3
months old C57Bl/6J naïve females were mated with naïve males for 1 week. Mothers were assigned
to control or MSUS group based on the number of male pups born on that day to balance the groups.
The MSUS paradigm started on postnatal day 1 (PND1): Pups were separated unpredictably from
their mother for 3 hours per day from PND1 until PND14. Separation occurred at any time during
the dark cycle. In addition, mothers were subjected to forced swim in cold water (18 °C for 5 min) or
restraint in a plastic tube (20 min) at unpredictable times during the 3 hours of separation. Control
mice were left undisturbed except for weekly cage changes and weight measurements. Pups were
weaned at PND21 and assigned to sex and treatment matched cages (4 to 5 mice) between PND22
and PND28. Co-assignment of siblings was avoided to exclude litter effects. At the age of 8-10
months, control (*n*=24) and MSUS (*n*=24) males were deeply anesthetized by Isoflurane (Attane,
Piramal Enteprise Limited), followed by decapitation. Males were single-housed the day before
sacrifice to reduce potential acute stress effects [104]. Body weight was monitored throughout the
experiment and on the day of single housing. Adult mice were housed in groups of 3 to 5 animals
in individually ventilated cages (SealSafe PLUS, Tecniplast). Animals were kept in a temperature-
and humidity-controlled facility on a 12h reversed light/dark cycle (light on at 20:00) with food (M/R
Haltung Extrudat, Provimi Kliba SA) and water *ad libitum*. Cages were changed weekly.

Figure 6*: **MSUS is a mouse model of early life stress**. Unpredictable maternal separation and unpredictable maternal*
stress is an animal model for early life stress. Naïve males are mated with naïve females for 1-7 days, then males are removed.
Dams gestate for about 21 days in normal conditions until delivery. The MSUS paradigm (red bar in the timeline) starts
at postnatal day 1 (PND1) and lasts until PND14. During MSUS, pups are unpredictably separated from their mothers
for 3 hours each day at different times of the dark cycle (lights off from 8am to 8pm). During separation, the mother is
stressed unpredictably by forced swim or restraint as shown in the timeline at the bottom. Pups and dam are left undisturbed
from PND15 until weaning at PND21. Gray mice are naïve without any previous stress exposure. Blue pups are controls
without any stress exposure. Red pups are exposed to MSUS.

4.1.2. Collection of serum samples

Trunk blood was collected in 2 ml tubes immediately after decapitation and incubated at 4°C.
Clotted blood was centrifuged at 2000 rcf at 4°C for 10 minutes. The collected serum was stored at -
80°C.

4.1.3. Collection of bone samples

After euthanasia and blood collection, the left femur was dissected on ice under aseptic
conditions and cleaned from muscle tissue. The ends were cut off to allow for removal of bone
marrow (flushed out with sterile phosphate buffered saline, PBS) and the entire flushed femur
(including the ends) was subsequently shock-frozen in liquid nitrogen and stored at -80°C until
further processing for qPCR. Right femurs were dissected, cleaned and stored in 70% ethanol at 4°C
until μCT analysis. The third tail bone was dissected and immersed in a mixture of 4%
paraformaldehyde and 14% saturated picric acid overnight at 4°C, cryoembedded and stored at -80°C
until further processing for histomorphometry was done.

4.1.4. Histomorphometry of neuronal markers

For bone innervation morphometry, immunohistochemistry of pan-neuronal marker and
exemplary markers for adrenergic, cholinergic and peptidergic nerve fiber subsets were performed
with undecalcified 14 μm thick transverse mouse tail cryostat sections analog to previous
descriptions [55]. The sections were done through the center of the third tail bone and contained the
bone with the adherent muscle, tendons and skin for internal positive control. Primary antibodies
used in this study were: anti-PGP 9.5 (Biotrend, Cologne, Germany), anti-GAP 43 (Millipore,
Darmstadt, Germany), anti-ChAT (Serotec, Kidlington, UK), anti-TH (Abcam, Cambridge, MA, USA)
and anti-SP (BMA Biomedicals, Augst, Switzerland). Only PGP 9.5 and GAP 43 gave satisfactory
staining results, most likely due to insufficient fixation of more digestion prone markers in bone
compared to the skin (positive controls were all positive), which commonly require transcardial
fixation [105]. Standard controls were performed by omission of the primary antibodies and by
incubation with mouse IgG1 instead of primary antibodies. No labeling was observed in negative
controls. Nerve fiber profiles were counted per area with the help of a grid equipped ocular at x200
magnification using a Leica inverted automated fluorescence-microscope (Leica, Wetzlar, Germany).
Briefly, numbers of positively stained fibers were counted per grid field in at least 30 consecutive
fields per experimental mouse and in at least *n=8* mice per experimental group by two blinded
independent researchers. The data were then pooled and expressed as means per group +/- standard
error of the mean.

4.1.5. qPCR for gene expression in bone samples

Mouse femurs were pulverized using a metal ball mill (Retsch, Germany) and total RNA was
isolated with Trizol/chloroform extraction, followed by column-based purification using RNeasy
Mini Kit (Qiagen, Germany) according to the manufacturer's protocol, but including proteinase K
incubation (55°C, 15 min). Contaminating DNA was removed by using the RNAse-free DNAse kit
(Qiagen, Germany) and 32 μl of RNA with an average yield of 48 ng/μl were reverse transcribed with
Superscript II reverse transcriptase (Life Technologies, Germany) to obtain 80 μl of cDNA. qPCR
analysis was performed in duplicate, using the TaqMan Fast PCR Master Mix with Taqman assays
for bone markers (Life Technologies, Germany) (*Supplementary Table 2*) or the QuantiTect PCR Kit

with self-designed primers for neuronal factors (Qiagen, Germany) (***Supplementary Table 3***). The
amount of cDNA per reaction depended on the respective target gene and its expression level and
was 1-18 ng. Gene expression was normalized to the housekeeping gene (dCt) and 2^{-dCt} were
computed.

4.1.6. ELISA of serum samples
Commercially available ELISA kits were used to determine the serum levels of OC (60-1305,
Immunotopics/Teca, Switzerland), PINP (AC-33F1, Immunodiagnosticsystem, Germany) and CTX-I
(AC-06F1, Immunodiagnosticsystem, Germany) according to the manufacturers' instructions. Serum
was used either undiluted (CTX-I) or at a 1/11 (osteocalcin) or 1/10 (PINP) dilution.

4.1.6. Micro-CT of bone samples
Analysis of bone microarchitecture by μCT was conducted as previously described [106]. Briefly,
bones were measured on a μCT40 (Scanco Medical AG, Brüttisellen, Switzerland), operated at 55 kVp
and 145 μA with an integration time of 200 ms and 2-fold frame averaging. Images were
reconstructed at an isotropic nominal resolution of 10 μm and filtered using a constrained 3D
Gaussian filter (sigma 0.8, support 1) to suppress noise. Masks of the full bone, cortical (shaft at 56%
bone length, 1 mm long) and trabecular region of interest (from 66% to 88% of bone length) were
created automatically. Morphometric parameters were calculated according to standard guidelines
such as total volume (TV), bone volume (BV), marrow volume (MV), apparent volume density
(AVD), femur length, cortical area fraction (Ct.Ar/Tt.Ar), cortical thickness (Ct.Th), trabecular bone
volume fraction (BV/TV), specific bone surface (BS/BV), trabecular thickness (Tb.Th), trabecular
spacing (Tb.Sp) and trabecular number (Tb.N) [107]. Due to differences in body weight between the
control mice and the MSUS mice, data was normalized for body weight and body weight adjusted
means were compared.

4.2 Human Study

4.2.1. Participants
n=240 patients with depressive disorder (ICD-10 F32.x, F33.x, F34.1, F43.21) were recruited of
which *n*=208 completed the initial examination (baseline (t0), for more details see [5]. A subsample of
n=54 patients took part at the fourth follow up measurement (t4) including DXA imaging. Patients
who fulfilled the following criteria were included: 18 to 65 years of age, depressive episode (ICD-10
F32.x or F33.x), dysthymia (F34.1) or an adjustment disorder with prolonged depressive reaction
(F43.21), ≥ 21 days absenteeism within the last year. Exclusion criteria were: pregnancy,
hormonotherapy (expect contraceptive and thyroid hormone therapy), inability to fill in a
questionnaire, intellectual disabilities (ICD-10 F70-79) or one of the following diseases: acute
infection, endocrine and metabolic disorders, neurological diseases, dementia (ICD-10 F00-F03),
schizophrenia (ICD-10 F20), emotional-unstable personality disorders (ICD-10 F60.3x), disease of the
immune system, substance abuse and dependency (expect nicotine).
All participants were fully informed in verbal and written form about the intent and content of
the study, and gave their written informed consent. The clinical investigations were conducted
according to the principles of the Declaration of Helsinki. Final ethical approval was provided on
11/12/2017 from the Ethics Review Board of the University of Potsdam, Germany (number 15/2017).

4.2.2. Study Procedure

Study objectives were investigated at two assessments (t_0 and t_4) of an 8-month observational

multicenter study with four measurement points: baseline (t_0), after 5 weeks (t_1), 5 months (t_2), 8

months (t_3) and an additional follow-up measurement (t_4: M= 15.4 month, SD = 4.5 month after

baseline). At each measurement point, study participants answered a comprehensive questionnaire

($t_0 - t_4$) under the supervision of trained study nurses. The questionnaire comprised demographic

characteristics, psychological symptoms (e.g. depression) and constructs (e.g. stress, early life events,

affect, coping), physical ailments, and information regarding alcohol consumption and medication

intake. Blood samples were collected at baseline (t_0) and 5 months follow-up (t_2) and DXA bone

densitometry measurements took place at the last follow up (t_4).

4.2.3. Psychometric Measures

Early life stress was measured retrospectively comprising experiences of childhood

maltreatment and stressful life events during childhood. The Childhood Trauma Screener (CTS)

[108], a 5-item screening tool of the 28-item Childhood Trauma Questionnaire (CTQ) [109, 110],

assesses five types of childhood maltreatment such as emotional, physical, and sexual abuse plus

emotional and physical neglect. Each item is scored on a five-point rating scale from "never true" (1)

to "very often true" (5). In accordance to Glaesmer et al. [111], we classified participants at risk if they

rated at least mild forms of childhood abuse or neglect. We additionally controlled for response bias

by the 3-item Minimization-Denial subscale from the CTQ and excluded participants when indicated.

Cronbach´s Alpha was previously specified with 0.76 [112].

Stressful life events were assessed by a modified version of the Inventory of Stressful Life Events

(ILE; [113]). Participants rated 34 adverse life events regarding occurrence, frequency and year of

occurrence. The accumulation of stressful, life-changing events at different stages of development

was counted for every participant, in this analysis focusing on childhood (ILE $_{child}$ ≤ 12 years). The

scores ranged between 0 (no strain) and 4 (high strain). Cronbach´s Alpha for ILE was 0.83.

Depressive symptoms and severity were assessed using the Beck Depression Inventory-II (BDI-II)

[114, 115]. Internal consistency in the sample was Cronbach's Alpha 0.89.

Furthermore, confounding factors like sociodemographic and biometric characteristics and

study site were assessed.

4.2.4. Serum bone marker measurement

Blood samples were drawn from the arm in the morning (7-9 am), collected in plain blood

collection tubes, allowed to clot at room temperature for 30 min, followed by centrifugation, isolation

of serum and freeze-storage until further analysis. Bone-related blood markers (OC, PINP, CTX-I)

were analyzed in serum samples by electrochemiluminescence immunoassays "ECLIA" (12149133

122 for Osteocalcin, 03141071 190 for PINP, 11972308 122 for CTX-I, all from F. Hoffmann-La Roche,

Ltd., Basel, Switzerland)

4.2.5. DXA measurement

Bone mineral density (BMD) was measured by DXA bone densitometry measurement (Lunar,

Prodigy Advance, GE Healthcare, Illinois, USA) in the lumbar spine (lumbar vertebral bodies L1-L4)

and both hips. Parameters are given as mean values (M) of the left and right site: G (total), H (collum

femoris), S (corpus femoris), T (trochanter major), and L1L4 (lumbar vertebral bodies L1-L4).

4.3. Statistical Analysis

Data consistency was checked, screened for outliers and analyzed descriptively. Continuous
variables were also tested for normality by using skewness, kurtosis, omnibus test; variance
homogeneity was proven by Variance Ratio and Levene test.

For the examination of **Aim 1** (mouse model), student t-test for equal or Aspin-Welch Unequal
Variance test were performed, using multiple t-test and adjusted for multiple comparison by Holm-
Sidak method (alpha = 0.05). Since body weight was found to be significantly different, ANCOVA
models with body weight as covariate were applied for micro-CT data to compute corresponding
means at their covariate means (i.e. least square means at covariate mean for body weight of 16.41).
In addition, blocking on total body weight was used as a means of accounting for collinearity in
analyses of variance and mediation analyses were conducted using linear regression models to
identify mediation effects between body weight and bone microstructure. All reported tests are two-
sided, and p-values < 0.05 are considered as statistically significant.

Regarding **Aim 2** (human model) multiple regression models were applied. Statistical models were
cross-sectional and controlled for age, gender, study sites and additionally body weight for DXA
measurements.

Statistical analyses were conducted using NCSS (NCSS 10, NCSS, LLC. Kaysville, UT), STATISTICA
13, IBM SPSS 25 and the statistical software R [116, 117].

Supplementary Materials: Supplementary materials can be found at www.mdpi.com/xxx/s1.

Author Contributions: KW-K planned and coordinated the project, participated in tissue collection, supervised
experiments and drafted the manuscript; MR conducted the animal experiments, participated in tissue collection
and drafted the manuscript; EC participated in tissue collection, analyzed samples and drafted the manuscript;
AB helped to coordinate the human study, analyzed sample and proof-read the manuscript; GAK participated
in tissue collection, analyzed samples and drafted the manuscript; TA analyzed samples and proof-read the
manuscript; WH conducted statistical analysis of all animal experiments and proof-read the manuscript; DD
conducted statistical analysis of all human experiments and proof-read the manuscript; RM provided input in
micro-CT analysis and proof-read the manuscript; MAR helped in the coordination of the human study and
proof-read the manuscript; IMM planned and coordinated the animal study, supervised experiments, acquired
funding and proof-read the manuscript; EMJP helped to coordinate the mouse study, analyzed samples and
drafted the manuscript; PMW planned and coordinated the project, conducted human study as PI, supervised
experiments, acquired funding and drafted the manuscript.

Funding: This study was supported by the Deutsche Forschungsgemeinschaft, the Central Research Grant for
Innovative Ideas from the University of Potsdam, the German Pension Fund Association (10-R-40.07.05.07.008),
the Swiss National Science Foundation (31003A_153147/1) and the ETHZ grant (ETH-10 15-2).

Acknowledgments: In this section you can acknowledge any support given which is not covered by the author
contribution or funding sections. This may include administrative and technical support, or donations in kind
(e.g., materials used for experiments).

Conflicts of Interest: None of the authors has any conflicts of interest or financial disclosures.

Abbreviations

ADRB2	adrenoceptor beta 2
ANCOVA	analysis of covariance
AVD	apparent volume density
BDI-II	Beck Depression Inventory Revision
BDNF	brain derived neurotrophic factor
BMD	bone mineral density
BMD_G	bone mineral density general
BMD_H	bone mineral density collum femoris
BMD_L1L4	bone mineral density lumbar vertebralis bodies 1-4
BMD_S	bone mineral density corpus femoris
BMD_T	bone mineral density trochanter major
BV	bone volume
BS	specific bone surface
CGRP	calcitonin gene related peptide
ChAT	choline Acetyltransferase
CHRNA7	cholinergic Receptor Nicotinic Alpha 7 Subunit
CTQ	Childhood Trauma Questionnaire
CTS	Childhood Trauma Screener
CTX-I	c-terminal telopeptide of type I collagen
Ct.Ar	cortical area fraction
Ct.Th	cortical thickness
DAPI	4′,6-diamidino-2-phenylindole
dCt	delta Ct
DEPREHA	[Der Einfluss unterschiedlicher Behandlungs-settings auf den Therapieerfolg bei Patienten mit depressiven Erkrankungen in der Rehabilitation]
DXA	Dual Energy X-ray Absorptiometry
ECLIA	electrochemiluminescence immunoassays
ECM	extracellular matrix
ELISA	enzyme-linked immunosorbent assay
FITC	Fluorescein isothiocyanate
GAP 43	Growth Associated Protein 43
HPA	hypothalamic pituitary adrenal
ICD-10	International Classification of Diseases Version 10
IgG1	Immunoglobulin G
ILE	Inventory of Stressful Life Events
ILE child	Inventory of Stressful Life Events; life events during childhood ≤12 years
μCT	micro-computed tomography
M	mean
MV	marrow volume
MSUS	maternal separation and unpredictable stress
mRNA	messenger ribonucleic acid
NF	nerve fibers
NGF	nerve growth factor
NPY	neuropeptide Y
NPYR1	neuropeptide Y receptor 1
OC	osteocalcin
qPCR	quantitative polymerase chain reaction
PBS	phosphate buffered saline
PINP	procollagen type 1 N-terminal propeptide
PND	postnatal day
RAMP1	receptor activity modifying protein 1
SAM	senescence-accelerated mouse
SD	standard deviation
SP	substance P

TAC1	tachykinin 1
TACR1	tachykinin receptor 1
Tb.N	trabecular number
Tb.Sp	trabecular spacing
Tb.Th	trabecular thickness
TH	tyrosine hydroxylase
TRKA	tropomyosin receptor kinase A
Tt.Ar	total area
TRKB	tropomyosin receptor kinase B
TV	total volume
VIP	vasoactive intestinal peptide
VIPR1	vasoactive intestinal peptide receptor 1

References

1. Wippert, P. M.; Rector, M.; Kuhn, G.; Wuertz-Kozak, K., Stress and Alterations in Bones: An interdisciplinary Perspective. *Front Endocrinol* **2017,** 8, doi:ARTN 96 10.3389/fendo.2017.00096.

2. Grassel, S., The role of peripheral nerve fibers and their neurotransmitters in cartilage and bone physiology and pathophysiology. *Arthritis Res Ther* **2014,** 16, (6), doi:ARTN 485 10.1186/s13075-014-0485-1.

3. Mattia, C.; Coluzzi, F.; Celidonio, L.; Vellucci, R., Bone pain mechanism in osteoporosis: a narrative review. *Clin Cases Miner Bon* **2016,** 13, (2), 97-100, doi:10.11138/ccmbm/2016.13.2.097.

4. McEwen, B. S.; Nasca, C.; Gray, J. D., Stress Effects on Neuronal Structure: Hippocampus, Amygdala, and Prefrontal Cortex. *Neuropsychopharmacol* **2016,** 41, (1), 3-23, doi:10.1038/npp.2015.171.

5. Wippert, P. M.; Block, A.; Mansuy, I. M.; Peters, E. M. J.; Rose, M.; Rapp, M. A.; Huppertz, A.; Wuertz-Kozak, K., Alterations in Bone Homeostasis and Microstructure Related to Depression and Allostatic Load. *Psychother Psychosom* **2019,** 88, (6), 383-385, doi:10.1159/000503640.

6. Asada, K.; Obata, K.; Horiguchi, K.; Takaki, M., Age-related changes in afferent responses in sensory neurons to mechanical stimulation of osteoblasts in coculture system. *Am J Physiol Cell Physiol* **2012,** 302, (5), C757-65, doi:10.1152/ajpcell.00362.2011.

7. Takahashi, N.; Matsuda, Y.; Sato, K.; de Jong, P. R.; Bertin, S.; Tabeta, K.; Yamazaki, K., Neuronal TRPV1 activation regulates alveolar bone resorption by suppressing osteoclastogenesis via CGRP. *Sci Rep* **2016,** 6, 29294, doi:10.1038/srep29294.

8. Mach, D. B.; Rogers, S. D.; Sabino, M. C.; Luger, N. M.; Schwei, M. J.; Pomonis, J. D.; Keyser, C. P.; Clohisy, D. R.; Adams, D. J.; O'Leary, P.; Mantyh, P. W., Origins of skeletal pain: sensory and sympathetic innervation of the mouse femur. *Neuroscience* **2002,** 113, (1), 155-66.

9. Chartier, S. R.; Mitchell, S. A. T.; Majuta, L. A.; Mantyh, P. W., The Changing Sensory and Sympathetic Innervation of the Young, Adult and Aging Mouse Femur. *Neuroscience* **2018,** doi:10.1016/j.neuroscience.2018.01.047.

10. Alves, C. J.; Alencastre, I. S.; Neto, E.; Ribas, J.; Ferreira, S.; Vasconcelos, D. M.; Sousa, D.
M.; Summavielle, T.; Lamghari, M., Bone Injury and Repair Trigger Central and Peripheral
NPY Neuronal Pathways. *PLoS One* **2016,** 11, (11), e0165465,
doi:10.1371/journal.pone.0165465.
11. Jiao, K.; Niu, L.; Xu, X.; Liu, Y.; Li, X.; Tay, F. R.; Wang, M., Norepinephrine Regulates
Condylar Bone Loss via Comorbid Factors. *J Dent Res* **2015,** 94, (6), 813-20,
doi:10.1177/0022034515577677.
12. Yirmiya, R.; Goshen, I.; Bajayo, A.; Kreisel, T.; Feldman, S.; Tam, J.; Trembovier, V.;
Csernus, V.; Shohami, E.; Bab, I., Depression induces bone loss through stimulation of the
sympathetic nervous system. *P Natl Acad Sci USA* **2006,** 103, (45), 16876-16881,
doi:10.1073/pnas.0604234103.
13. Furuzawa, M.; Chen, H. Y.; Fujiwara, S.; Yamada, K.; Kubo, K., Chewing ameliorates
chronic mild stress-induced bone loss in senescence-accelerated mouse (SAMP8), a murine
model of senile osteoporosis. *Exp Gerontol* **2014,** 55, 12-18, doi:10.1016/j.exger.2014.03.003.
14. Tomlinson, R. E.; Li, Z.; Zhang, Q.; Goh, B. C.; Li, Z.; Thorek, D. L. J.; Rajbhandari, L.;
Brushart, T. M.; Minichiello, L.; Zhou, F.; Venkatesan, A.; Clemens, T. L., NGF-TrkA
Signaling by Sensory Nerves Coordinates the Vascularization and Ossification of
Developing Endochondral Bone. *Cell Rep* **2016,** 16, (10), 2723-2735,
doi:10.1016/j.celrep.2016.08.002.
15. Chen, W. H.; Mao, C. Q.; Zhuo, L. L.; Ong, J. L., Beta-nerve growth factor promotes
neurogenesis and angiogenesis during the repair of bone defects. *Neural Regen Res* **2015,** 10,
(7), 1159-65, doi:10.4103/1673-5374.160114.
16. Zhuang, Y. F.; Li, J., Serum EGF and NGF levels of patients with brain injury and limb
fracture. *Asian Pac J Trop Med* **2013,** 6, (5), 383-6, doi:10.1016/S1995-7645(13)60043-7.
17. Sabsovich, I.; Wei, T.; Guo, T. Z.; Zhao, R.; Shi, X.; Li, X.; Yeomans, D. C.; Klyukinov, M.;
Kingery, W. S.; Clark, J. D., Effect of anti-NGF antibodies in a rat tibia fracture model of
complex regional pain syndrome type I. *Pain* **2008,** 138, (1), 47-60,
doi:10.1016/j.pain.2007.11.004.
18. Guo, T. Z.; Wei, T. P.; Shi, X. Y.; Li, W. W.; Hou, S. Y.; Wang, L. P.; Tsujikawa, K.; Rice, K.
C.; Cheng, K. J.; Clark, D. J.; Kingery, W. S., Neuropeptide deficient mice have attenuated
nociceptive, vascular, and inflammatory changes in a tibia fracture model of complex
regional pain syndrome. *Mol Pain* **2012,** 8, doi:Artn 85
10.1186/1744-8069-8-85.
19. Sang, X. G.; Wang, Z. Y.; Cheng, L.; Liu, Y. H.; Li, Y. G.; Qin, T.; Di, K., Analysis of the
mechanism by which nerve growth factor promotes callus formation in mice with tibial
fracture. *Exp Ther Med* **2017,** 13, (4), 1376-1380, doi:10.3892/etm.2017.4108.
20. Grills, B. L.; Schuijers, J. A.; Ward, A. R., Topical application of nerve growth factor
improves fracture healing in rats. *J Orthop Res* **1997,** 15, (2), 235-42,
doi:10.1002/jor.1100150212.
21. Hemingway, F.; Taylor, R.; Knowles, H. J.; Athanasou, N. A., RANKL-independent human
osteoclast formation with APRIL, BAFF, NGF, IGF I and IGF II. *Bone* **2011,** 48, (4), 938-44,
doi:10.1016/j.bone.2010.12.023.

22.   Camerino, C.; Conte, E.; Caloiero, R.; Fonzino, A.; Carratu, M.; Lograno, M. D.; Tricarico,
625   D., Evaluation of Short and Long Term Cold Stress Challenge of Nerve Grow Factor, Brain-
626   Derived Neurotrophic Factor, Osteocalcin and Oxytocin mRNA Expression in BAT, Brain,
627   Bone and Reproductive Tissue of Male Mice Using Real-Time PCR and Linear Correlation
628   Analysis. *Front Physiol* **2017,** 8, 1101, doi:10.3389/fphys.2017.01101.

629   23.   Liu, Q.; Lei, L.; Yu, T.; Jiang, T.; Kang, Y., Effect of Brain-Derived Neurotrophic Factor on
630   the Neurogenesis and Osteogenesis in Bone Engineering. *Tissue Eng Part A* **2018**,
631   doi:10.1089/ten.TEA.2017.0462.

632   24.   Yang, M.; Lu, B. J.; Duan, Y. Y.; Chen, X. F.; Ma, J. G.; Guo, Y., Genetics association study
633   and functional analysis on osteoporosis susceptibility gene BDNF. *Yi Chuan* **2017,** 39, (8),
634   726-736, doi:10.16288/j.yczz.17-145.

635   25.   Kauschke, V.; Gebert, A.; Calin, M.; Eckert, J.; Scheich, S.; Heiss, C.; Lips, K. S., Effects of
636   new beta-type Ti-40Nb implant materials, brain-derived neurotrophic factor, acetylcholine
637   and nicotine on human mesenchymal stem cells of osteoporotic and non osteoporotic
638   donors. *PLoS One* **2018,** 13, (2), e0193468, doi:10.1371/journal.pone.0193468.

639   26.   Camerino, C.; Conte, E.; Cannone, M.; Caloiero, R.; Fonzino, A.; Tricarico, D., Nerve
640   Growth Factor, Brain-Derived Neurotrophic Factor and Osteocalcin Gene Relationship in
641   Energy Regulation, Bone Homeostasis and Reproductive Organs Analyzed by mRNA
642   Quantitative Evaluation and Linear Correlation Analysis. *Front Physiol* **2016,** 7, 456,
643   doi:10.3389/fphys.2016.00456.

644   27.   Picard, M.; Juster, R. P.; McEwen, B. S., Mitochondrial allostatic load puts the 'gluc' back in
645   glucocorticoids. *Nat Rev Endocrinol* **2014,** 10, (5), 303-10, doi:10.1038/nrendo.2014.22.

646   28.   Picard, M.; McEwen, B. S., Psychological Stress and Mitochondria: A Conceptual
647   Framework. *Psychosom Med* **2018,** 80, (2), 126-140, doi:10.1097/PSY.0000000000000544.

648   29.   Kelly, R. R.; McDonald, L. T.; Jensen, N. R.; Sidles, S. J.; LaRue, A. C., Impacts of
649   Psychological Stress on Osteoporosis: Clinical Implications and Treatment Interactions.
650   *Front Psychiatry* **2019,** 10, 200, doi:10.3389/fpsyt.2019.00200.

651   30.   Lee, S. H.; Mastronardi, C. A.; Li, R. W.; Paz-Filho, G.; Dutcher, E. G.; Lewis, M. D.; Vincent,
652   A. D.; Smith, P. N.; Bornstein, S. R.; Licinio, J.; Wong, M. L., Short-term antidepressant
653   treatment has long-lasting effects, and reverses stress-induced decreases in bone features in
654   rats. *Transl Psychiatry* **2019,** 9, (1), 10, doi:10.1038/s41398-018-0351-z.

655   31.   Ji-Ye, H.; Xin-Feng, Z.; Lei-Sheng, J., Autonomic control of bone formation: its clinical
656   relevance. *Handb Clin Neurol* **2013,** 117, 161-71, doi:10.1016/B978-0-444-53491-0.00014-6.

657   32.   Schiavone, S.; Morgese, M. G.; Mhillaj, E.; Bove, M.; De Giorgi, A.; Cantatore, F. P.;
658   Camerino, C.; Tucci, P.; Maffulli, N.; Cuomo, V.; Trabace, L., Chronic Psychosocial Stress
659   Impairs Bone Homeostasis: A Study in the Social Isolation Reared Rat. *Front Pharmacol* **2016,**
660   7, 152, doi:10.3389/fphar.2016.00152.

661   33.   Huang, W. S.; Hsu, J. W.; Huang, K. L.; Bai, Y. M.; Su, T. P.; Li, C. T.; Lin, W. C.; Chen, T. J.;
662   Tsai, S. J.; Liou, Y. J.; Chen, M. H., Post-traumatic stress disorder and risk of osteoporosis: A
663   nationwide longitudinal study. *Stress Health* **2018,** 34, (3), 440-445, doi:10.1002/smi.2806.

664   34.   Gershon, A.; Sudheimer, K.; Tirouvanziam, R.; Williams, L. M.; O'Hara, R., The long-term
665   impact of early adversity on late-life psychiatric disorders. *Curr Psychiatry Rep* **2013,** 15, (4),
666   352, doi:10.1007/s11920-013-0352-9.

667   35.   Raymond, C.; Marin, M. F.; Majeur, D.; Lupien, S., Early child adversity and
668   psychopathology in adulthood: HPA axis and cognitive dysregulations as potential
669   mechanisms. *Prog Neuro-Psychoph* **2018,** 85, 152-160, doi:10.1016/j.pnpbp.2017.07.015.
670   36.   Shonkoff, J. P.; Boyce, W. T.; McEwen, B. S., Neuroscience, molecular biology, and the
671   childhood roots of health disparities: building a new framework for health promotion and
672   disease prevention. *JAMA* **2009,** 301, (21), 2252-9, doi:10.1001/jama.2009.754.
673   37.   Falgares, G.; Marchetti, D.; Manna, G.; Musso, P.; Oasi, O.; Kopala-Sibley, D. C.; De Santis,
674   S.; Verrocchio, M. C., Childhood Maltreatment, Pathological Personality Dimensions, and
675   Suicide Risk in Young Adults. *Front Psychol* **2018,** 9, 806, doi:10.3389/fpsyg.2018.00806.
676   38.   Lindert, J.; von Ehrenstein, O. S.; Grashow, R.; Gal, G.; Braehler, E.; Weisskopf, M. G.,
677   Sexual and physical abuse in childhood is associated with depression and anxiety over the
678   life course: systematic review and meta-analysis. *Int J Public Health* **2014,** 59, (2), 359-372,
679   doi:10.1007/s00038-013-0519-5.
680   39.   Infurna, M. R.; Reichl, C.; Parzer, P.; Schimmenti, A.; Bifulco, A.; Kaess, M., Associations
681   between depression and specific childhood experiences of abuse and neglect: A meta-
682   analysis. *J Affect Disorders* **2016,** 190, 47-55, doi:10.1016/j.jad.2015.09.006.
683   40.   Nanni, V.; Uher, R.; Danese, A., Childhood Maltreatment Predicts Unfavorable Course of
684   Illness and Treatment Outcome in Depression: A Meta-Analysis. *Am J Psychiat* **2012,** 169,
685   (2), 141-151, doi:10.1176/appi.ajp.2011.11020335.
686   41.   Mezuk, B.; Eaton, W. W.; Golden, S. H., Depression and osteoporosis: epidemiology and
687   potential mediating pathways. *Osteoporosis Int* **2008,** 19, (1), 1-12, doi:10.1007/s00198-007-
688   0449-2.
689   42.   Cizza, G.; Primma, S.; Coyle, M.; Gourgiotis, L.; Csako, G., Depression and Osteoporosis: A
690   Research Synthesis with Meta-Analysis. *Horm Metab Res* **2010,** 42, (7), 467-482, doi:10.1055/s-
691   0030-1252020.
692   43.   Azuma K; Adachi Y; Hayashi H; KY., K., Chronic Psychological Stress as a Risk Factor of
693   Osteoporosis. *J UOEH.* **2015,** 37, (4), 245-53.
694   44.   Williams, L. J.; Pasco, J. A.; Jackson, H.; Kiropoulos, L.; Stuart, A. L.; Jacka, F. N.; Berk, M.,
695   Depression as a risk factor for fracture in women: A 10 year longitudinal study. *J Affect
696   Disorders* **2016,** 192, 34-40, doi:10.1016/j.jad.2015.11.048.
697   45.   Furlan, P. M.; Ten Have, T.; Cary, M.; Zemel, B.; Wehrli, F.; Katz, I. R.; Gettes, D. R.; Evans,
698   D. L., The role of stress-induced cortisol in the relationship between depression and
699   decreased bone mineral density. *Biol Psychiat* **2005,** 57, (8), 911-917,
700   doi:10.1016/j.biopsych.2004.12.033.
701   46.   Stults-Kolehmainen, M. A.; Sinha, R., The effects of stress on physical activity and exercise.
702   *Sports Med* **2014,** 44, (1), 81-121, doi:10.1007/s40279-013-0090-5.
703   47.   Yau, Y. H.; Potenza, M. N., Stress and eating behaviors. *Minerva Endocrinol* **2013,** 38, (3),
704   255-67.
705   48.   Franklin, T. B.; Russig, H.; Weiss, I. C.; Graff, J.; Linder, N.; Michalon, A.; Vizi, S.; Mansuy,
706   I. M., Epigenetic Transmission of the Impact of Early Stress Across Generations. *Biol
707   Psychiat* **2010,** 68, (5), 408-415, doi:10.1016/j.biopsych.2010.05.036.
708   49.   Schmidt, M. V., Molecular mechanisms of early life stress--lessons from mouse models.
709   *Neurosci Biobehav Rev* **2010,** 34, (6), 845-52, doi:10.1016/j.neubiorev.2009.05.002.

50. McGowan, P. O.; Szyf, M., The epigenetics of social adversity in early life: implications for
mental health outcomes. *Neurobiol Dis* **2010,** 39, (1), 66-72, doi:10.1016/j.nbd.2009.12.026.

51. Bohacek, J.; Mansuy, I. M., Epigenetic inheritance of disease and disease risk.
Neuropsychopharmacol **2013,** 38, (1), 220-36, doi:10.1038/npp.2012.110.

52. Gapp, K.; Jawaid, A.; Sarkies, P.; Bohacek, J.; Pelczar, P.; Prados, J.; Farinelli, L.; Miska, E.;
Mansuy, I. M., Implication of sperm RNAs in transgenerational inheritance of the effects of
early trauma in mice. *Nat Neurosci* **2014,** 17, (5), 667-9, doi:10.1038/nn.3695.

53. Baldock, P. A.; Lin, S.; Zhang, L.; Karl, T.; Shi, Y.; Driessler, F.; Zengin, A.; Hormer, B.; Lee,
N. J.; Wong, I. P. L.; Lin, E. J. D.; Enriquez, R. F.; Stehrer, B.; During, M. J.; Yulyaningsih, E.;
Zolotukhin, S.; Ruohonen, S. T.; Savontaus, E.; Sainsbury, A.; Herzog, H., Neuropeptide Y
Attenuates Stress-Induced Bone Loss Through Suppression of Noradrenaline Circuits. *J
Bone Miner Res* **2014,** 29, (10), 2238-2249, doi:10.1002/jbmr.2205.

54. Ballica, R.; Valentijn, K.; Khachatryan, A.; Guerder, S.; Kapadia, S.; Gundberg, C.; Gilligan,
J.; Flavell, R. A.; Vignery, A., Targeted expression of calcitonin gene-related peptide to
osteoblasts increases bone density in mice. *J Bone Miner Res* **1999,** 14, (7), 1067-74,
doi:10.1359/jbmr.1999.14.7.1067.

55. Rommel, F. R.; Raghavan, B.; Paddenberg, R.; Kummer, W.; Tumala, S.; Lochnit, G.; Gieler,
U.; Peters, E. M., Suitability of Nicotinic Acetylcholine Receptor alpha7 and Muscarinic
Acetylcholine Receptor 3 Antibodies for Immune Detection: Evaluation in Murine Skin. *J
Histochem Cytochem* **2015,** 63, (5), 329-39, doi:10.1369/0022155415575028.

56. Sousa, D. M.; McDonald, M. M.; Mikulec, K.; Peacock, L.; Herzog, H.; Lamghari, M.; Little,
D. G.; Baldock, P. A., Neuropeptide Y modulates fracture healing through Y1 receptor
signaling. *J Orthop Res* **2013,** 31, (10), 1570-8, doi:10.1002/jor.22400.

57. Teixeira, L.; Sousa, D. M.; Nunes, A. F.; Sousa, M. M.; Herzog, H.; Lamghari, M., NPY
Revealed as a Critical Modulator of Osteoblast Function In Vitro: New Insights Into the
Role of Y1 and Y2 Receptors. *J Cell Biochem* **2009,** 107, (5), 908-916, doi:10.1002/jcb.22194.

58. Tang, P.; Duan, C. G.; Wang, Z.; Wang, C. M.; Meng, G. L.; Lin, K. F.; Yang, Q.; Yuan, Z.,
NPY and CGRP Inhibitor Influence on ERK Pathway and Macrophage Aggregation during
Fracture Healing. *Cell Physiol Biochem* **2017,** 41, (4), 1457-1467, doi:10.1159/000468405.

59. Igwe, J. C.; Jiag, X.; Paic, F.; Ma, L.; Adams, D. J.; Baldock, P. A.; Pilbeam, C. C.; Kalajzic, I.,
Neuropeptide Y Is Expressed by Osteocytes and Can Inhibit Osteoblastic Activity. *J Cell
Biochem* **2009,** 108, (3), 621-630, doi:10.1002/jcb.22294.

60. Allison, S. J.; Baldock, P. A.; Enriquez, R. F.; Lin, E. J.; During, M.; Gardiner, E. M.; Eisman,
J. A.; Sainsbury, A.; Herzog, H., Critical Interplay Between Neuropeptide Y and Sex Steroid
Pathways in Bone and Adipose Tissue Homeostasis. *J Bone Miner Res* **2009,** 24, (2), 294-304,
doi:10.1359/Jbmr.081013.

61. Sousa, D. M.; Baldock, P. A.; Enriquez, R. F.; Zhang, L.; Sainsbury, A.; Lamghari, M.;
Herzog, H., Neuropeptide Y Y1 receptor antagonism increases bone mass in mice. *Bone
748* **2012,** 51, (1), 8-16, doi:10.1016/j.bone.2012.03.020.

749 62. Ma, W. H.; Liu, Y. J.; Wang, W.; Zhang, Y. Z., Neuropeptide Y, substance P, and human
750 bone morphogenetic protein 2 stimulate human osteoblast osteogenic activity by enhancing
751 gap junction intercellular communication. *Braz J Med Biol Res* **2015,** 48, (4), 299-307,
752 doi:10.1590/1414-431x20144226.

753   63.   Long, H.; Ahmed, M.; Ackermann, P.; Stark, A.; Li, J. A., Neuropeptide Y innervation
754   during fracture healing and remodeling. *Acta Orthop* **2010,** 81, (5), 639-646,
755   doi:10.3109/17453674.2010.504609.

756   64.   Cherruau, M.; Morvan, F. O.; Schirar, A.; Saffar, J. L., Chemical sympathectomy-induced
757   changes in TH-, VIP-, and CGRP-immunoreactive fibers in the rat mandible periosteum:
758   Influence on bone resorption. *J Cell Physiol* **2003,** 194, (3), 341-348, doi:10.1002/jcp.10209.

759   65.   Lundberg, P.; Lundgren, I.; Mukohyama, H.; Lehenkari, P. P.; Horton, M. A.; Lerner, U. H.,
760   Vasoactive intestinal peptide (VIP)/pituitary adenylate cyclase-activating peptide receptor
761   subtypes in mouse calvarial osteoblasts: Presence of VIP-2 receptors and differentiation-
762   induced expression of VIP-1 receptors. *Endocrinology* **2001,** 142, (1), 339-347, doi:DOI
763   10.1210/en.142.1.339.

764   66.   Ransjo, M.; Lie, A.; Mukohyama, H.; Lundberg, P.; Lerner, U. H., Microisolated mouse
765   osteoclasts express VIP-1 and PACAP receptors. *Biochem Bioph Res Co* **2000,** 274, (2), 400-404,
766   doi:DOI 10.1006/bbrc.2000.3151.

767   67.   Nagata, A.; Tanaka, T.; Minezawa, A.; Poyurovsky, M.; Mayama, T.; Suzuki, S.; Hashimoto,
768   N.; Yoshida, T.; Suyama, K.; Miyata, A.; Hosokawa, H.; Nakayama, T.; Tatsuno, I., cAMP
769   Activation by PACAP/VIP Stimulates IL-6 Release and Inhibits Osteoblastic Differentiation
770   Through VPAC2 Receptor in Osteoblastic MC3T3 Cells. *J Cell Physiol* **2009,** 221, (1), 75-83,
771   doi:10.1002/jcp.21831.

772   68.   Persson, E.; Lerner, U. H., The Neuropeptide VIP Regulates the Expression of
773   Osteoclastogenic Factors in Osteoblasts. *J Cell Biochem* **2011,** 112, (12), 3732-3741,
774   doi:10.1002/jcb.23304.

775   69.   Wei, T. P.; Li, W. W.; Guo, T. Z.; Zhao, R.; Wang, L. P.; Clark, D. J.; Oaklander, A. L.;
776   Schmelz, M.; Kingery, W. S., Post-junctional facilitation of Substance P signaling in a tibia
777   fracture rat model of complex regional pain syndrome type I. *Pain* **2009,** 144, (3), 278-286,
778   doi:10.1016/j.pain.2009.04.020.

779   70.   Aoki, M.; Tamai, K.; Saotome, K., Substance P-Related and Calcitonin-Gene-Related
780   Peptide-Immunofluorescent Nerves in the Repair of Experimental Bone Defects. *Int Orthop*
781   **1994,** 18, (5), 317-324.

782   71.   Goto, T.; Yamaza, T.; Kido, M. A.; Tanaka, T., Light- and electron-microscopic study of the
783   distribution of axons containing substance P and the localization of neurokinin-1 receptor
784   in bone. *Cell Tissue Res* **1998,** 293, (1), 87-93, doi:DOI 10.1007/s004410051100.

785   72.   Wang, L. P.; Zhao, R.; Shi, X. Y.; Wei, T. P.; Halloran, B. P.; Clark, D. J.; Jacobs, C. R.;
786   Kingery, W. S., Substance P stimulates bone marrow stromal cell osteogenic activity,
787   osteoclast differentiation, and resorption activity in vitro. *Bone* **2009,** 45, (2), 309-320,
788   doi:10.1016/j.bone.2009.04.203.

789   73.   Mori, T.; Ogata, T.; Okumura, H.; Shibata, T.; Nakamura, Y.; Kataoka, K., Substance p
790   regulates the function of rabbit cultured osteoclast; Increase of intracellular free calcium
791   concentration and enhancement of bone resorption. *Biochem Bioph Res Co* **1999,** 262, (2), 418-
792   422, doi:DOI 10.1006/bbrc.1999.1220.

793   74.   Sohn, S. J., Substance P upregulates osteoclastogenesis by activating nuclear factor kappa B
794   in osteoclast precursors. *Acta Oto-Laryngol* **2005,** 125, (2), 130-133,
795   doi:10.1080/00016480410017710.

75. Matayoshi, T.; Goto, T.; Fukuhara, E.; Takano, H.; Kobayashi, S.; Takahashi, T.,
Neuropeptide substance P stimulates the formation of osteoclasts via synovial fibroblastic
cells. *Biochem Bioph Res Co* **2005,** 327, (3), 756-764, doi:DOI 10.1016/j.bbrc.2004.12.055.

76. Adamus, M. A.; Dabrowski, Z. J., Effect of the neuropeptide substance P on the rat bone
marrow-derived osteogenic cells in vitro. *J Cell Biochem* **2001,** 81, (3), 499-506.

77. Liu, H. J.; Yan, H.; Yan, J.; Li, H.; Chen, L.; Han, L. R.; Yang, X. F., Substance P Promotes the
Proliferation, but Inhibits Differentiation and Mineralization of Osteoblasts from Rats with
Spinal Cord Injury via RANKL/OPG System. *PLoS One* **2016,** 11, (10), e0165063,
doi:10.1371/journal.pone.0165063.

78. Shih, C.; Bernard, G. W., Neurogenic substance P stimulates osteogenesis in vitro. *Peptides*
1997, 18, (2), 323-6.

79. Goto, T.; Nakao, K.; Gunjigake, K. K.; Kido, M. A.; Kobayashi, S.; Tanaka, T., Substance P
stimulates late-stage rat osteoblastic bone formation through neurokinin-1 receptors.
Neuropeptides **2007,** 41, (1), 25-31, doi:10.1016/j.npep.2006.11.002.

80. Kook, Y. A.; Lee, S. K.; Son, D. H.; Kim, Y.; Kang, K. H.; Cho, J. H.; Kim, S. C.; Kim, Y. S.;
Lee, H. J.; Lee, S. K.; Kim, E. C., Effects of substance P on osteoblastic differentiation and
heme oxygenase-1 in human periodontal ligament cells. *Cell Biol Int* **2009,** 33, (3), 424-428,
doi:10.1016/j.cellbi.2008.12.007.

81. Azuma, H.; Kido, J.; Ikedo, D.; Kataoka, M.; Nagata, T., Substance P enhances the inhibition
of osteoblastic cell differentiation induced by lipopolysaccharide from Porphyromonas
gingivalis. *J Periodontol* **2004,** 75, (7), 974-981, doi:DOI 10.1902/jop.2004.75.7.974.

82. Kingery, W. S.; Offley, S. C.; Guo, T. Z.; Davies, M. F.; Clark, J. D.; Jacobs, C. R., A substance
P receptor (NK1) antagonist enhances the widespread osteoporotic effects of sciatic nerve
section. *Bone* **2003,** 33, (6), 927-936, doi:10.1016/j.bone.2003.07.003.

83. Zheng, X. F.; Zhao, E. D.; He, J. Y.; Zhang, Y. H.; Jiang, S. D.; Jiang, L. S., Inhibition of
substance P signaling aggravates the bone loss in ovariectomy-induced osteoporosis. *Prog*
Biophys Mol Bio **2016,** 122, (2), 112-121, doi:10.1016/j.pbiomolbio.2016.05.011.

84. Ding, W. G.; Zhang, Z. M.; Zhang, Y. H.; Jiang, S. D.; Jiang, L. S.; Dai, L. Y., Changes of
substance P during fracture healing in ovariectomized mice. *Regul Peptides* **2010,** 159, (1-3),
28-34, doi:10.1016/j.regpep.2009.11.004.

85. Lerner, U. H.; Persson, E., Osteotropic effects by the neuropeptides calcitonin gene-related
peptide, substance P and vasoactive intestinal peptide. *J Musculoskelet Neuronal Interact*
2008, 8, (2), 154-65.

86. Terreni, A.; Pezzati, P., Biochemical markers in the follow-up of the osteoporotic patients.
Clin Cases Miner Bone Metab **2012,** 9, (2), 80-4.

87. Vasikaran, S.; Eastell, R.; Bruyere, O.; Foldes, A. J.; Garnero, P.; Griesmacher, A.; McClung,
M.; Morris, H. A.; Silverman, S.; Trenti, T.; Wahl, D. A.; Cooper, C.; Kanis, J. A.; Group, I.-I.
B. M. S. W., Markers of bone turnover for the prediction of fracture risk and monitoring of
osteoporosis treatment: a need for international reference standards. *Osteoporos Int* **2011,** 22,
(2), 391-420, doi:10.1007/s00198-010-1501-1.

88. Yoshimura, N.; Muraki, S.; Oka, H.; Kawaguchi, H.; Nakamura, K.; Akune, T., Biochemical
markers of bone turnover as predictors of osteoporosis and osteoporotic fractures in men

and women: 10-year follow-up of the Taiji cohort. *Mod Rheumatol* **2011,** 21, (6), 608-20,
doi:10.1007/s10165-011-0455-2.

89.   Garnero, P.; Sornay-Rendu, E.; Claustrat, B.; Delmas, P. D., Biochemical markers of bone
turnover, endogenous hormones and the risk of fractures in postmenopausal women: the
OFELY study. *J Bone Miner Res* **2000,** 15, (8), 1526-36, doi:10.1359/jbmr.2000.15.8.1526.

90.   Blue, M. E.; Boskey, A. L.; Doty, S. B.; Fedarko, N. S.; Hossain, M. A.; Shapiro, J. R.,
Osteoblast function and bone histomorphometry in a murine model of Rett syndrome. *Bone*
2015, 76, 23-30, doi:10.1016/j.bone.2015.01.024.

91.   Eastell, R.; Barton, I.; Hannon, R. A.; Chines, A.; Garnero, P.; Delmas, P. D., Relationship of
early changes in bone resorption to the reduction in fracture risk with risedronate. *J Bone*
Miner Res **2003,** 18, (6), 1051-1056, doi:DOI 10.1359/jbmr.2003.18.6.1051.

92.   Callewaert, F.; Venken, K.; Kopchick, J. J.; Torcasio, A.; van Lenthe, G. H.; Boonen, S.;
Vanderschueren, D., Sexual dimorphism in cortical bone size and strength but not density
is determined by independent and time-specific actions of sex steroids and IGF-1: evidence
from pubertal mouse models. *J Bone Miner Res* **2010,** 25, (3), 617-26, doi:10.1359/jbmr.090828.

93.   Goel, N.; Bale, T. L., Sex differences in the serotonergic influence on the hypothalamic-
pituitary-adrenal stress axis. *Endocrinology* **2010,** 151, (4), 1784-94, doi:10.1210/en.2009-1180.

94.   Takeda, T.; Hosokawa, M.; Higuchi, K., Senescence-accelerated mouse (SAM): a novel
murine model of accelerated senescence. *J Am Geriatr Soc* **1991,** 39, (9), 911-9.

95.   Paccione, M. F.; Warren, S. M.; Spector, J. A.; Greenwald, J. A.; Bouletreau, P. J.; Longaker,
M. T., A mouse model of mandibular osteotomy healing. *J Craniofac Surg* **2001,** 12, (5), 444-
450, doi:Doi 10.1097/00001665-200109000-00008.

96.   Dang, R.; Guo, Y. Y.; Zhang, K.; Jiang, P.; Zhao, M. G., Predictable chronic mild stress
promotes recovery from LPS-induced depression. *Mol Brain* **2019,** 12, (1), 42,
doi:10.1186/s13041-019-0463-2.

97.   Furuzawa, M.; Chen, H.; Fujiwara, S.; Yamada, K.; Kubo, K. Y., Chewing ameliorates
chronic mild stress-induced bone loss in senescence-accelerated mouse (SAMP8), a murine
model of senile osteoporosis. *Exp Gerontol* **2014,** 55, 12-8, doi:10.1016/j.exger.2014.03.003.

98.   Heim, C.; Mletzko, T.; Purselle, D.; Musselman, D. L.; Nemeroff, C. B., The
dexamethasone/corticotropin-releasing factor test in men with major depression: role of
childhood trauma. *Biol Psychiatry* **2008,** 63, (4), 398-405, doi:10.1016/j.biopsych.2007.07.002.

99.   Heim, C.; Shugart, M.; Craighead, W. E.; Nemeroff, C. B., Neurobiological and psychiatric
consequences of child abuse and neglect. *Dev Psychobiol* **2010,** 52, (7), 671-90,
doi:10.1002/dev.20494.

100.   Saveanu, R. V.; Nemeroff, C. B., Etiology of Depression: Genetic and Environmental
Factors. *Psychiat Clin N Am* **2012,** 35, (1), 51-+, doi:10.1016/j.psc.2011.12.001.

101.   Picard, M.; McEwen, B. S.; Epel, E. S.; Sandi, C., An energetic view of stress: Focus on
mitochondria. *Front Neuroendocrinol* **2018,** 49, 72-85, doi:10.1016/j.yfrne.2018.01.001.

102.   Sophocleous, A.; Idris, A. I., Rodent models of osteoporosis. *Bonekey Rep* **2014,** 3, 614,
doi:10.1038/bonekey.2014.109.

103.   Halloran, B. P.; Ferguson, V. L.; Simske, S. J.; Burghardt, A.; Venton, L. L.; Majumdar, S.,
Changes in bone structure and mass with advancing age in the male C57BL/6J mouse. *J*
Bone Miner Res **2002,** 17, (6), 1044-1050, doi:DOI 10.1359/jbmr.2002.17.6.1044.

104. Bohacek, J.; Manuella, F.; Roszkowski, M.; Mansuy, I. M., Hippocampal gene expression
induced by cold swim stress depends on sex and handling. *Psychoneuroendocrinology* **2015,**
52, 1-12, doi:10.1016/j.psyneuen.2014.10.026.

105. Hendrix, S.; Picker, B.; Liezmann, C.; Peters, E. M., Skin and hair follicle innervation in
experimental models: a guide for the exact and reproducible evaluation of neuronal
plasticity. *Exp Dermatol* **2008,** 17, (3), 214-27, doi:10.1111/j.1600-0625.2007.00653.x.

106. Kohler, T.; Beyeler, M.; Webster, D.; Muller, R., Compartmental bone morphometry in the
mouse femur: Reproducibility and resolution dependence of microtomographic
measurements. *Calcified Tissue Int* **2005,** 77, (5), 281-290, doi:10.1007/s00223-005-0039-2.

107. Bouxsein, M. L.; Boyd, S. K.; Christiansen, B. A.; Guldberg, R. E.; Jepsen, K. J.; Muller, R.,
Guidelines for Assessment of Bone Microstructure in Rodents Using Micro-Computed
Tomography. *J Bone Miner Res* **2010,** 25, (7), 1468-1486, doi:10.1002/jbmr.141.

108. Glaesmer, H.; Schulz, A.; Häuser, W.; Freyberger, H. J.; Brähler, E.; Grabe, H.-J., Der
Childhood Trauma Screener (CTS) – Entwicklung und Validierung von Schwellenwerten
zur Klassifikation. [The Childhood Trauma Screener (CTS) – Development and Validation
of Cut-Off-Scores for Classificatory Diagnostics]. *Psychiatrische Praxis* **2013,** 40, (04), 220-226,
doi:10.1055/s-0033-1343116.

109. Wingenfeld, K.; Spitzer, C.; Mensebach, C.; Grabe, H. J.; Hill, A.; Gast, U.; Schlosser, N.;
Höpp, H.; Beblo, T.; Driessen, M., Die deutsche Version des Childhood Trauma
Questionnaire (CTQ): Erste Befunde zu den psychometrischen Kennwerten. [The German
version of the Childhood Trauma Questionnaire (CTQ): Preliminary psychometric
properties]. *Psychotherapie, Psychosomatik, und Medizinische Psychologie* **2010,** 60, (11), 442-450.

110. Bernstein, D.; Fink, L., Childhood Trauma Questionnaire (CTQ): A retrospective self-Report
questionnaire and manual. In Corporation, T. P., Ed. San Antonio, 1998.

111. Glaesmer, H.; Schulz, A.; Hauser, W.; Freyberger, H. J.; Brahler, E.; Grabe, H. J., [The
childhood trauma screener (CTS) - development and validation of cut-off-scores for
classificatory diagnostics]. *Psychiatr Prax* **2013,** 40, (4), 220-6, doi:10.1055/s-0033-1343116.

112. Grabe, H. J.; Schulz, A.; Schmidt, C. O.; Appel, K.; Driessen, M.; Wingenfeld, K.; Barnow, S.;
Spitzer, C.; John, U.; Berger, K.; Wersching, H.; Freyberger, H. J., [A brief instrument for the
assessment of childhood abuse and neglect: the childhood trauma screener (CTS)]. *Psychiatr*
Prax **2012,** 39, (3), 109-15, doi:10.1055/s-0031-1298984.

113. Siegrist, J.; Geyer, S., ILE–Inventar zur Erfassung lebensverändernder Ereignisse. In
Diagnostische Verfahren in der Psychotherapie, Hogrefe, Ed. Göttingen, 2002; pp 211-213.

114. Beck, A. T.; Steer, R. A.; Brown, G. K., *Manual for the Beck Depression Inventory-II*. T.P.
Corporation: San Antonio, 1996.

115. Wintjen, L.; Petermann, F., Beck-Depressions-Inventar Revision (BDI-II). *Z Psychiatr Psych*
Ps **2010,** 58, (3), 243-245, doi:10.1024/1661-4747.a000033.

116. Muche, R.; Lanzinger, S.; Rau, M., *Medizinische Statistik mit R und Excel*. Springer: 2011.

117. Team, R. C., R foundation for Statistical Computing. In Vienna, Austria, 2014.

Supplementary Data

Supplementary Figures

S-Figure 1: Body weight development. *Shown are body weights of male pups, which are equal at PND1 and decreased following MSUS at PND7 (n=45 Control, n=46 MSUS), PND14 (n=45 Control, n=46 MSUS), PND21 (n=46 Control, n=44 MSUS) and in adults (n=14 Control, n=14 MSUS). Data are expressed as Min to Max of body weight in [g] and considered significant at *p < 0.05 and ***p < 0.001.*

Supplementary Tables

S-Table 1: *Micro-CT data obtained for full, cortical (Ct) and trabecular (Tb) bone in control mice and MSUS mice with regard to length, total volume (TV), bone volume (BV), marrow volume (MV), apparent volume density (AVD), area (Ar), thickness (Th), bone surface (BS), spacing (Sp) and number (N) and thereof calculated characteristics.*

		Control	MSUS	
	Variables	**Adjusted Mean**	**Adjusted Mean**	**p-value (adjusted)[1]**
Full Bone	**Length (mm)**	16.45	16.37	0.263
	TV (mm³)	44.23	42.88	0.129
	BV (mm³)	21.31	20.48	0.128
	MV (mm³)	22.92	22.40	0.457
	AVD (%)	48.19	47.77	0.675
Cortical Bone	**Tt.Ar (mm²)**	0.92	0.90	0.478
	Ct.Ar/Tt.Ar (%)	40.31	40.36	0.959
	Ct.Th (μm)	0.18	0.18	0.844
Trabecular Bone	**TV (mm³)**	5.19	5.10	0.558
	BV/TV (%)	6.99	7.26	0.792
	MV/TV (%)	93.01	92.74	0.792
	BS/BV (mm/mm³)	45.10	47.37	0.203
	Tb.Th (mm)	0.07	0.07	0.282
	Tb.Sp (mm)	0.49	0.44	0.319
	Tb.N (1/mm)	1.98	2.14	0.452

[1] *adjusted for body weight at a covariate mean of 16.41. Significance level: all $p > 0.05$*

943   **S-Table 2:** Taqman gene expression assays

Target	Identifier
GAPDH	Mm99999915_g1
Osteocalcin	Mm03413826_mH
Osteoprotegerin	Mm00435452_m1
Osteopontin	Mm00436767_m1
Sclerostin	Mm00470479_m1

947   **S-Table 3:** Self-designed primers

NCBI accession Nr	Target identifier / Primer Sequence
NM_007420	ADRB2
muAdrb2 - Forward	gATTgCAgTggATCgCTATgTTg
muAdrb2 - Reverse	gACCACTCgggCCTTATTCTTg
muAdrb2 -TM	F-ATCACATCgCCCTTCAAgTACCAgAgC-Q
NM_001033124	NtTRK
muNtrk1 - Forward	gggAgTTgAgAAgCCTAACCAT
muNtrk1 - Reverse	CgCATTggAggACAgATTCA
muNtrk1 -TM	F-CAgATgCCTTCCgTTTCACCCCT-Q
NM_033217	NGFR
muNgfr - Forward	gTggAgAgTgCTgCAAAgC
muNgfr - Reverse	CgCTCACCACgTCAgAg
muNgfr -TM	F-CCTTgCggAgCCAACCAgACC-Q
NM_001025074	NRTK2
muNtrk2 - Forward	CTCAgCAAATCgCAgCAgg
muNtrk2 - Reverse	AgTAgTCggTgCTgTACA
muNtrk2 -TM	F-TTCACCAgCAggTTCTCTCCCACC-Q
NM_016894	RAMP1
muRAMP1- Forward	gTggggCTCTgCTTgC
muRAMP1-Reverse	gCACAgCTCCTggATgAgAgT
muRAMP1 -TM	F-ATCTCTTCATggTCACTgCCTgCCg-Q
NM_011703	VIPR1
muVipr1 - Forward	gCCTgTTCAggAAgCTgC
muVipr1 - Reverse	ggTCTgTCTCCCCgTTgT
muVipr1 -TM	F-CgAAACTACATCCACATgCATCTCTTCATg-Q
NM_01093	NPY1r
muNpy1r - Forward	CACgACTCTCCTCCTggTg
muNpy1r - Reverse	gCgAATgTATATCTTgAAgTAg
muNpy1r -TM	F-AgCAgAgTgggCCgAAATACTgCA--—Q
NM_009313	TACR1
Tacr1 - Forward	gACTTACgAgAAAgCgTACCACAT
Tacr1 - Reverse	CTggCCCACAgTgTAATCCCTA
Tacr1 -TM	FAM-TgTgTgACTgTgCTgATCTACTTCCTgCCT—BBQ
NM_007390	NAChRα7
nAChRα7 - Forward	GCCCTTGATAGCACAGTACTTCG
nAChRα7 - Reverse	GATCCTGGTCCACTTAGGCATTT
nAChRα7TM	6FAM-CAGTGGTCGTGACAGTGATTGTGCTGC-BBQ
NM_009311.2	TAC1
Tac1- Forward	CAACTGAGGAATCAGCATCC
Tac1 - Reverse	GCAGAGAATCGCCCGAA

Tac1-TM	6FAM-AgCCTCAgCAgTTCTTTggATTAATggg-BBQ
NM_007540	BDNF ex9
Bdnf- Forward	gTCTCTgCTTCCTTCCCACAg
Bdnf - Reverse	CCTTgTCCgTggACgTTTACT
TM	6FAM-CgCCTTCATgCAACCgAAgTATgAAA-BBQ
NM_009891.2	ChAT
Chat - Forward	CCAggACggTCCTCTTAAAAgAC
Chat - Reverse	CCCTgTgTgTgTCACTgAggT
Chat -TM	6FAM-CgggACTCCCTggACATgATCgAgC-BBQ
NM_013609.3	NGF
Ngf- Forward	gCAgTgAggTgCATAgCg
Ngf - Reverse	gTggAgTCTCCgTTTCTTAAACAgTC
Ngf-TM	6FAM-CCgCAgTgCCCCTACTgCACCAATA--—BBQ
NM_011702	VIP
Vip - Forward	TgAATggAAAgAggAgCAgTg
Vip - Reverse	CgTggTTgTTTTCCTTCgAg
TM	6FAM- TTgAAgAgCTggAgAAATgATgggAAgA--—BBQ
NM_023456	NPY
Npy - Forward	gCAgAggACATggCCAgATAC
Npy - Reverse	TgTgCTTTCCTTCATTAAgAggTC
TM	6FAM-CAAgAgATCCAgCCCTgAgACACTgA--—BBQ
NM_013684.3	TBP
Tbp - Forward	gTgAATCTTggCTgTAAACTTgACCT
Tbp - Reverse	gCAgTTgTCCgTggCTC
Tbf -TM	6FAM-AAATgCTgAATATAATCCCAAgCgATTTgC—BBQ

950