Effects of early life stress on bone homeostasis in mice and humans

Karin Wuertz-Kozak[a,b,c*], Martin Roszkowski[d*], Elena Cambria[c], Andrea Block[e,] Gisela A. Kuhn[c], Thea Abele[f], Wolfgang Hitzl[g,h,i], David Drießlein[j], Ralph Müller[c], Michael A. Rapp[k], Isabelle M. Mansuy[d#], Eva M. J. Peters[f,l#], Pia M. Wippert[c, e#]

[a] Department of Biomedical Engineering, Rochester Institute of Technology (RIT), Rochester (NY), USA

[b] Schön Clinic Munich Harlaching, Spine Center, Academic Teaching Hospital and Spine Research Institute of the Paracelsus Medical University Salzburg (AU), Munich, Germany

[c] Institute for Biomechanics, Department of Health Sciences and Technology, ETH Zurich, Zurich, Switzerland

[d] Laboratory of Neuroepigenetics, University of Zurich and Swiss Federal Institute of Technology, Brain Research Institute, Neuroscience Center Zürich, 8057 Zurich, Switzerland

[e] Department of Health Sciences, University of Potsdam, Potsdam, Germany

[f] Psychoneuroimmunology Laboratory, Department of Psychosomatics and Psychotherapy, University of Gießen, Gießen, Germany

[g.] Research Office (Biostatistics), Paracelsus Medical University, Salzburg, Austria

[h.] Research Program Experimental Ophthalmology and Glaucoma Research, Paracelsus Medical University, Salzburg, Austria

[i.] Paracelsus Medical University Salzburg, Department of Ophthalmology and Optometry, Austria

[j] Statistical Consulting Unit StaBLab, Ludwig-Maximilians-University Munich, Munich, Germany

[k] Social and Preventive Medicine, Department of Health Sciences, University of Potsdam, Potsdam, Germany

[l] Charité Center 12 Internal Medicine and Dermatology, Division for General Internal Medicine, Psychosomatics and Psychotherapy, Universitätsmedizin Berlin, Berlin, Germany

* Shared first authorship   # Shared last authorship

Corresponding Author
Prof. Dr. Karin Wuertz-Kozak (Orcid 0000-0002-3281-4629)
Department of Biomedical Engineering, Rochester Institute of Technology (RIT)
160 Lomb Memorial Drive Bldg. 73
Rochester, NY 14623
kwbme@rit.edu; 001 585 475-7355

Short title: Early life stress and bone homeostasis

Key Words: Psychosocial stress; bone pathologies; osteoporosis; bone mineral density; childhood; neuroendocrine

Impact Statement

Specific types of stress in early life lead to maladaptive responses in bone metabolism and bone mineral density and hence possibly to bone pathologies.

Abstract

A comprehensive examination of mechanisms linking early life stress, depression and disturbed bone homeostasis is missing. In this translational study, mice exposed to early life stress (MSUS) were examined on bone microarchitecture (μCT), metabolism (qPCR), and neuronal mediator expression (qPCR) and contrasted with a sample of depressive patients with or without early life stress by analyzing bone mineral density [BMD] (DXA) and metabolic changes in serum (osteocalcin, PINP, CTX-I). MSUS mice showed a significant decrease in NGF, NPYR1, VIPR1 and TACR expression, higher innervation density in bone and increased serum levels of CTX-I, suggesting a milieu in favor of catabolic bone turnover. Depressive patients with experiences of childhood neglect also showed a catabolic pattern. A significant reduction in BMD was observed in depressive patients with childhood abuse and stressful life events during childhood. Therefore, future prevention strategies should consider early life stress as a risk factor for bone pathologies.

Introduction

Bones are essential components of the musculoskeletal system and subjected to continuous remodeling as an adaptation mechanism to environmental changes. Disturbances in bone development and remodeling by formation/resorption of the extracellular matrix (ECM) could result in reduced bone mass and increased fracture risk.

Beside well-documented mechanisms like menopause-associated hormonal changes, aging-related factors, changes in physical activity, as well as drugs and diseases, recent research suggested that psychosocial stress and stress-linked neuronal processes are associated with an increased risk for bone pathologies [1-5]. Autonomic and sensory nerve fibers densely innervate bone, predominantly in the periosteum and in areas with high metabolic activity and bone turnover. Neuronal structures can make direct contact with osteoblasts and osteoclasts, which in turn express receptors for neuronal markers on their surface and can produce neuronal mediators in an auto- and paracrine fashion [6-10]. Importantly, sympathetic and peptidergic neuronal factors appear to play a key role in bone turnover [11-13]. Of specific relevance are tyrosine hydroxylase (TH), a rate-limiting enzyme for noradrenaline synthesis, neuropeptide Y (NPY), often co-expressed with noradrenaline, vasoactive intestinal peptide (VIP), often co-expressed with acetylcholine, and the neuropeptides calcitonin gene related peptide (CGRP) and substance P (SP) that are expressed in sensory nerve fibers. In addition, neurotrophic growth factors such as nerve growth factor (NGF) and BDNF, reduced by stress, guide outgrowing nerve fibers to bony areas requiring innervation, and can act as direct and indirect growth factors for osteoblasts [14-26]. Latest studies indicate that mediators of psychosocial stress can lead to structural and functional changes in mitochondria, neuronal plasticity and inflammation [4, 27, 28], possibly resulting in downstream alterations of bone homeostasis [29-33].

In humans, major early life stress like childhood maltreatment strongly affects the stress response, as well as behavioral and cognitive functions throughout life [34, 35]. Strong associations were shown between childhood maltreatment and higher risk for psychological

and physiological disorders including depression, hypersensitivity of the neuroendocrinological

and autonomic stress response, altered dynamics of the hypothalamic pituitary adrenal (HPA)

axis, and changes in neuronal mediators (e.g. brain derived neurotrophic factor, BDNF) that

can influence bone health [36]. Childhood maltreatment is one of the best documented risk

factors for depression [37-40] and depression in turn has been associated with a higher risk

for reduced bone mineral density (BMD), osteoporosis and bone fractures [41-45].

However, existing knowledge about underlying mechanisms between early life stress,

neurogenic stress response elements, and detrimental changes in bone metabolism and

microstructure/BMD has been limited by several reasons. First, although early life stress like

childhood maltreatment is an ubiquitous and frequent phenomenon, it is hardly accessible to

retrospective and longitudinal studies due to denial, concealment or memory bias and ethical

obsolete for case-control studies. Second, access to biological material and strict control of

confounding factors, such as reduced physical activity and eating habits, are also challenging

in humans [46, 47]. An obvious way out of the mentioned hardship to investigate major stress

experiences early in life offer animal models. Animal models are helpful counterparts of human

studies to investigate underlying mechanisms in a manageable environment by allowing

extensive tissue sampling, strict control of adverse events and confounding factors, and lower

subject variability.

Unpredictable maternal separation and maternal stress during early life (MSUS paradigm [48])

is an experimental stress paradigm in rodents that can mimic aspects of childhood

maltreatment and induce long-lasting health effects, such as increased depression-like

behavior (anhedonia) and altered brain activity in adulthood [49]. The MSUS paradigm

severely impacts physiology and behavior in the exposed animals but also in their descendants

across several generations via mechanisms involving epigenetic factors in the germline [48,

50-52].

To address the knowledge gap between early life stress, stress-dependent neuronal plasticity

and detrimental changes in bone metabolism and bone microstructure/BMD, we first examined

the impact of early life stress on bone microarchitecture (μCT), metabolism (qPCR on bone,

ELISA on serum for osteocalcin OC, PINP, CTX-I), and neuronal mediator expression (qPCR)

in mice using the MSUS paradigm (**Aim 1**). Then we translated the findings to humans by

analyzing microarchitecture (DXA) and metabolic changes in serum (OC, PINP, CTX-I) in a

sample of depressive patients with or without retrospectively reported experiences of early life

stress, like childhood maltreatment and stressful life events during childhood (**Aim 2**).

Results

[A] Mouse Study

Bone innervation and neuronal mediators

We first assessed the effects of early life stress on neuronal signaling and neurogenic factors

in bone metabolism by measuring neuronal density and expression patterns of neurogenic

markers in the bone of MSUS and control mice (*n=6-8* each). The density of nerve fibers

labeled with the pan-neuronal marker PGP 9.5 was significantly increased in the bone of

MSUS mice (***Figure 1***), whereas nerve fibers labeled with the neuronal plasticity marker GAP

43 were unaffected (not shown), indicating that higher innervation was not due to increased

neuronal plasticity, but was likely a stable feature. Higher innervation density potentially leads

to higher sensitivity of bone neurons to stimuli like injury. Noteworthy, the neuropeptides NPY,

VIP and SP were undetectable by immunohistochemistry, most likely due to insufficient bone

preservation. This likely results from the post-mortem fixation method used for ethical

requirements, which does not preserve small peptides (the eleven amino-acid sequence of SP

for example is rapidly digested by endogenous enzymes after termination of circulation).

To study changes in neuronal markers, qPCR was performed in addition to

immunohistomorphometry. Gene expression analysis revealed that key receptors for nerve

growth factors, neurotransmitters and neuropeptides (***Figure 2A***) and selected receptors

ligands (***Figure 2B***) were changed in MSUS mice in a pattern that may promote higher bone

turnover and lower bone density. In detail, mRNA expression of the receptors NPYR1 ($p <$

0.05, ***Figure 2A***), VIPR1 ($p < 0.001$, ***Figure 2A***) and TACR1 ($p < 0.01$, ***Figure 2A***) were

significantly reduced in MSUS mice. These three receptors are involved in the regulation of proliferation and differentiation in bone and their concomitant downregulation may lead to higher remodelling activity with less proliferation and more differentiation of bone cells. Furthermore, a significant reduction of the neurotrophin NGF ($p < 0.001$, **Figure 2B**) was revealed, while NPY showed a downtrend in MSUS ($p = 0.068$, **Figure 2B**). This would hamper both neuronal plasticity and bone remodelling in the face of a challenge such as a wound or aging leading to osteoporosis. Other markers of bone innervation and metabolism such as NGFR, ADRB2, CHRNA7, RAMP1, TACR1 and TAC1 were similar in control and MSUS mice, whereas TRKA, TRKB, TH, ChAT and VIP were below the detection limit.

Bone metabolic parameters

We then examined if changes in neuronal markers were associated with bone metabolic changes by measuring markers in serum and bone. Serum levels of osteocalcin and PINP, two markers associated with bone formation, did not change after MSUS (**Figure 3A-B**). In contrast, the level of CTX-I, which is associated with bone resorption, was significantly increased in MSUS mice compared to control mice ($p < 0.01$, n=14 each; **Figure 3C**). However, this catabolic shift was not reflected by altered gene expression of tissue-specific extracellular matrix markers (osteocalcin, osteoprotegerin, osteopontin, sclerostin) in bone samples of MSUS (**Figure 4A-D**) (*n=8* each).

Bone microarchitecture

Next, it was examined if MSUS affects bone microarchitecture in mice using micro-CT (n=14 MSUS, n=13 Control). As bone microarchitecture is sensitive to mechanical loading, we also measured body weight of experimental animals over time. Body weight was significantly lower in MSUS mice compared to control mice starting at postnatal day 7 (PND7) and remained lower in adulthood ($p < 0.001$ for PND 7-28 and $p < 0.05$ for 10 months, **Supplementary Figure 1**), and this caused minor effects on bone microarchitecture.

After adjusting micro-CT data for body weight, a comparison of means between control and MSUS mice did not reveal any difference in full, cortical or trabecular microarchitecture (all

values p > 0.05) (**Supplementary Table 1**). We also used blocking in tertiles for body weight

and conducted a two-factorial multivariate analysis of variance, which revealed that indeed the

effects of MSUS versus controls were driven by body weight. To specify this effect, we

conducted sets of linear regression analyses to test for simple mediation effects and found that

body weight mediated the effect of MSUS on full length, TV, BV, MV and BS (all p's for direct

effects > 0.05).

[B] Human Study

Descriptive results

N=193 out of n=208 patients under study suffered from depression with minimal (35%) up to

moderate/ severe (65%) severity (BDI-II: $M = 23.8$, $SD = 10.7$). A subset of $n = 165$ (age $M =$

47.6, $SD = 10.8$; $female = 77.6\%$) filled in the *Inventory of Stressful Life Events* (ILE) and n=54

(age $M = 48.5$, $SD = 10.2$; $female = 75.4\%$) the *Childhood Trauma Screener* (CTS). N=32

patients (age: $M = 47.4$, $SD = 9.9$, $f = 68.8\%$, BDI-II: $M = 25.4$, $SD = 11.1$) were identified at

risk by the CTS and experienced at least one type of childhood maltreatment (abuse or

neglect). Out of these participants, n=18 were available for DXA bone densitometry

measurements. Stressful life events at childhood (≤12 years; ILE $_{child}$) were identified for n=34

patients (age: $M = 42.7$ $SD = 11.9$, f = 76.5%) of which n=15 participated in DXA

measurements. Average bone mineral density (BMD) in the sample was M= 0.942 g/cm^2 (SD

= 0.186 g/cm^2) in the trochanter region of the proximal femur. Blood samples regarding bone

turnover markers were available for n=193 participants.

Bone metabolic parameters

Regarding changes in human bone metabolism following early life stress, bone serum markers

OC, PINP, CTX-I were analyzed. It was shown that depressed patients with childhood trauma

(CTS) had an increased anabolic bone metabolism in comparison to depressed patients

without early life stress (p = 0.03* for P1PN, p = 0.05* for CTX, **Table 1**). This anabolic bone

metabolism was likewise found for depressed patients with stressful life events during

childhood (ILE $_{child}$) in comparison to depressed patients without early life stress (p = 0.04* for

P1PN, *Table 1*). Depressed patients with experiences of childhood neglect (CTS neglect) lack

such anabolic reaction (*Table 1*).

Bone mineral density

Depressive patients *with* stressful life events during childhood (ILE $_{child}$) had reduced BMD in

the trochanter major (p = 0.03* for BMD_T) and as a tendency a reduction in general BMD (p

= 0.08# for BMD_G, *Table 1*). Depressed patients with a history of childhood abuse showed

specifically a BMD reduction in the trochanter major (p = 0.05* for BMD_T) and a tendency in

reduced BMD in general (p = 0.07# for BMD_G), giving notice of a disadvantage of people with

maltreatment and stressful life events during childhood in comparison to people with

depression lacking experiences of early life stress (**Table 1, Figure 5**).

Discussion

The first aim of this study was to investigate the effects of early life stress on bone health and

associated biological factors in a mouse model of early life stress (MSUS). The second study

aim was the translation of the gained knowledge in a human model using a sample of

depressive patients with and without early life stress.

Aim 1: In summary, we found that early life stress in the MSUS model altered bone innervation

and neuronal mediator expression as well as bone metabolism, as evidenced by increased

CTX-I serum concentrations, but did not affect expression of bone markers within the bone or

bone microarchitecture. In detail, we found significantly decreased expression of NGF,

NPYR1, VIPR1 and TACR1 in MSUS bones. This response pattern is suggestive for a higher

rate of bone turnover and lower bone healing capacities as a consequence of early life stress,

suggesting a concurrent joint downregulation, reduced bone remodeling and long-term

destabilization. Neurotrophin NGF is important for the expression of cholinergic and sensory

neuronal factors and hence for the development and the repair of bone, as demonstrated by

reduced bone length when absent [14-26]. The neuropeptide NPY promotes osteoblast

differentiation and balances the effects of stress-induced bone-loss through beta-adrenergic stimulation, both with consequences for bone remodeling during healing [10, 53-61]. The neuropeptide VIPR1, finally, promotes bone mineralization [62-66] while TACR1 regulates osteoblasts, osteoclasts and mesenchymal stem cell functionality associated with protection from osteoporosis [2, 60, 67-83].

While MSUS mice showed a clear neuroendocrine response pattern, gene expression of typical bone markers (in bone) was unaffected. Similarly, we did not observe increased osteocalcin in the serum of MSUS mice. However, CTX-I serum levels were significantly increased in MSUS mice, which is an increasingly promoted and robust biochemical factor for bone resorption in clinical settings and further giving notice about correlations to bone mineral density [84-87].

Regarding bone microarchitecture, no significant changes were observed after normalization to body weight. Body weight itself, which was affected by MSUS, did cause minor changes in bone microarchitecture, likely due to differences in mechanical loading of the skeleton. It is unclear, if altered body weight might be a post stress consequence for MSUS or an incidental finding. This should be considered in future studies, as it can be a confounding factor in micro-CT analyses. To test whether the observed lack of changes in bone microarchitecture (despite an increase in serum CTX-I) might be due to divergent timelines of these events [88] or might be sex-dependent (different susceptibility to bone alterations and/or sensitivity to stress) [89, 90]. We additionally conducted micro-CT scans of the femurs of female mice over a more extended time range (up to 16 months). We did not find any differences in bone microarchitecture in the female mice, even not at later time points (data not shown). It is important to keep in mind that changes in bone microarchitecture may be site specific and may hence have been overlooked by restricting the micro-CT analysis to the femur only. Perhaps, the use of models of aging (e.g., the senescence-accelerated mouse = SAM) [91], models that include additional stress exposures (such as osteotomy) [92], or a comparison of stress-resilient and stress-prone subpopulations [93] might potentially have led to observable

changes in bone structure within a reasonable time frame [94], but further investigations will

be needed to allow for more definite statements.

Aim 2: In summary, we found that people that experienced childhood maltreatment or stressful

life events during childhood showed increased anabolic bone metabolism during a depression

episode. This was evidenced by increased serum levels in bone metabolism markers (distinctly

in PINP and CTX-I, less in OC). Further, depressive adults with experiences of childhood

maltreatment (childhood abuse) and stressful life events during childhood showed a

significantly decreased BMD. This indicates that people affected by stressful life events during

childhood suffer from a reduced BMD in adulthood and further, showed a threefold higher

metabolic reaction in PINP, for example, during a depressive episode in comparison to

depressed adults without childhood maltreatment [5]. In contrast, for depressive adults with

experiences of childhood neglect this pattern was not observed.

Other human studies have linked childhood maltreatment to neuroendocrinological and

neuroanatomic alterations that overlap with depressive symptoms and result in a vulnerable

phenotype that is affected by a hypersensitive central stress response, blunted stress reactivity

and lower stress resistance [95-97]. It was suggested that further severe distress, such as

additional traumatization, or the development of a depression, may promote an overreaction

of adaption. The response pattern in our data seems to confirm this assumption, which can

result from various aspects, such as an altered mitochondrial function, gene dysregulation,

accumulation of oxidative stress, inflammation and senescence or changes in neuronal

plasticity [27, 28]. The associations between stressful events early in life and an early decrease

of noradrenergic, cholinergic and sensory neuronal factors was also observed in our animal

model. It is also comprehensible that no such pattern could be found for childhood neglect, as

it represents acts of omissions that are typically temporary, such as lacks of nutrition, medical

or emotional care. Hence, these might have a lower impact on a person's psychological

integrity, identity and health than childhood abuse. Further, the data matches well with further

results within this study sample: patients that suffer from chronic stress (measured by the

allostatic load index) showed a catabolic shift in bone metabolism during a depression episode

[5]. This can be explained by the fact that other mechanisms are expected for allostatic load

as an accumulating damage over time with the consequence of neuroendocrinological and

neuroanatomic alterations [36, 96, 98]. The observed reduced bone mineral density can be

expected in both response pattern: within a hypersensitive stress response and within an

increased allostatic load. Both mechanisms were demonstrated in data of people of the

presented DERPEHA-study.

In summary, we found that early life stress affects bone metabolism in mice and humans. In

MSUS mice, this was evidenced by a significant shift towards a catabolic milieu (decreased

NGF, NPYR1, VIPR1, TACR; higher innervation density in bone; increased serum CTX-I), but

no alterations in bone microarchitecture. In humans however, a significant reduction in BMD

was observed as a result of early life stress. Species-related differences in bone metabolism

between mice and humans have been described [99]. Our data supports the notion that the

mechanisms of bone loss seem different between mice and humans, as mice did not show

any decrease in trabecular thickness, which is considered as typical for ageing humans [100].

However, this contrasts Halloran et al. [100], who showed that cancellous bone volume fraction

is reduced in 24 month old male C57BL/6J mice and is hence comparable to humans. The

observed differences in the response of mice and humans are likely caused by differences in

types of stress early in life. In fact, we were able to demonstrate clear differences between

different types of early life stress in the human sample, further supporting this theory. Although

the results obtained in the mouse study do not overlap entirely with the human results, it

provides relevant and promising potential for translation. Furthermore, the animal model can

be used to identify probable mechanisms underlying osteoporosis development following early

life stress (i.e. related to innervation patterns) that could not be investigated easily in humans,

thereby allowing to develop ideas for future studies or even intervention and prevention

strategies.

In conclusion, this study demonstrates that stressful life events early in life leads to (mal-)

adaptive responses in bone metabolism in both mice and humans. However, the major

conclusion to be drawn from the presented data is that different types of early life stress lead

to different response patterns concerning bone health and have to be consider separately:

MSUS mice responded with enhanced bone resorption, which however did not cause changes

in bone microarchitecture 8-10 months after the stress exposure. In humans, a reduction in

BMD was evident in depressive patients with severe stress exposure in early life like childhood

abuse and stressful life events during childhood. Similarly, the changes in bone serum markers

were also dependent on the severity of stress experiences early in life. People with childhood

abuse and stressful life events during childhood showed an increased bone metabolism while

people with childhood neglect showed a counter regulated reaction. Hence, the consequences

of the observed changes in the context of challenges to bone health (fractures, aging etc.)

should be studied in a more personalized manner and future prevention strategies should

respect the severity of the stress exposure to be treated (e.g. specific type of stress

experiences during childhood, double hit paradigms) to achieve the optimal benefit for patients

with increased risk for bone pathologies, such as people with depressive disorders.

Methods

[A] Mouse Study

Animal model

Animal experiments were conducted in strict adherence to the Swiss Law for Animal Protection

and were approved by the local authority (Veterinäramt des Kantons Zürich, Switzerland)

under license number 57/2015. C57Bl/6J mice were obtained from Janvier (France) and bred

in-house to generate the male mice (n=47 MSUS, n=46 Control) used for experiments.

The MSUS paradigm was conducted as previously described [48]. In brief, 2-3 months old

C57Bl/6J naïve females were mated with naïve males for 1 week. Mothers were assigned to

control or MSUS group based on the number of male pups born on that day to balance the

groups. The MSUS paradigm started on postnatal day 1 (PND1): Pups were separated unpredictably from their mother for 3 hours per day from PND1 until PND14. Separation occurred at any time during the dark cycle. In addition, mothers were subjected to forced swim in cold water (18 °C for 5 min) or restraint in a plastic tube (20 min) at unpredictable times during the 3 hours of separation. Control mice were left undisturbed except for weekly cage changes and weight measurements. Pups were weaned at PND21 and assigned to sex and treatment matched cages (4 to 5 mice) between PND22 and PND28. Co-assignment of siblings was avoided to exclude litter effects. At the age of 8-10 months, control (*n=24*) and MSUS (*n=24*) males were deeply anesthetized by Isoflurane (Attane, Piramal Enteprise Limited), followed by decapitation. Males were single-housed the day before sacrifice to reduce potential acute stress effects [101]. Body weight was monitored throughout the experiment and on the day of single housing. Adult mice were housed in groups of 3 to 5 animals in individually ventilated cages (SealSafe PLUS, Tecniplast). Animals were kept in a temperature- and humidity-controlled facility on a 12h reversed light/dark cycle (light on at 20:00) with food (M/R Haltung Extrudat, Provimi Kliba SA) and water *ad libitum*. Cages were changed weekly.

Collection of serum samples

Trunk blood was collected in 2 ml tubes immediately after decapitation and incubated at 4°C. Clotted blood was centrifuged at 2000 rcf at 4°C for 10 minutes. The collected serum was stored at -80°C.

Collection of bone samples

After euthanasia and blood collection, the left femur was dissected on ice under aseptic conditions and cleaned from muscle tissue. The ends were cut off to allow for removal of bone marrow (flushed out with sterile phosphate buffered saline, PBS) and the entire flushed femur (including the ends) was subsequently shock-frozen in liquid nitrogen and stored at -80°C until further processing for qPCR. Right femurs were dissected, cleaned and stored in 70% ethanol at 4°C until μCT analysis. The third tail bone was dissected and immersed in a mixture of 4%

paraformaldehyde and 14% saturated picric acid overnight at 4°C, cryoembedded and stored at -80°C until further processing for histomorphometry was done.

Histomorphometry of neuronal markers

For bone innervation morphometry, immunohistochemistry of pan-neuronal marker and exemplary markers for adrenergic, cholinergic and peptidergic nerve fiber subsets were performed with undecalcified 14 μm thick transverse mouse tail cryostat sections analog to previous descriptions [102]. The sections were done through the center of the third tail bone and contained the bone with the adherent muscle, tendons and skin for internal positive control. Primary antibodies used in this study were: anti-PGP 9.5 (Biotrend, Cologne, Germany), anti-GAP 43 (Millipore, Darmstadt, Germany), anti-ChAT (Serotec, Kidlington, UK), anti-TH (Abcam, Cambridge, MA, USA) and anti-SP (BMA Biomedicals, Augst, Switzerland). Only PGP 9.5 and GAP 43 gave satisfactory staining results, most likely due to insufficient fixation of more digestion prone markers in bone compared to the skin (positive controls were all positive), which commonly require transcardial fixation [103]. Standard controls were performed by omission of the primary antibodies and by incubation with mouse IgG1 instead of primary antibodies. No labeling was observed in negative controls. Nerve fiber profiles were counted per area with the help of a grid equipped ocular at x200 magnification using a Leica inverted automated fluorescence-microscope (Leica, Wetzlar, Germany). Briefly, numbers of positively stained fibers were counted per grid field in at least 30 consecutive fields per experimental mouse and in at least *n=8* mice per experimental group by two blinded independent researchers. The data were then pooled and expressed as means per group +/- standard error of the mean.

qPCR for gene expression in bone samples

Mouse femurs were pulverized using a metal ball mill (Retsch, Germany) and total RNA was isolated with Trizol/chloroform extraction, followed by column-based purification using RNeasy Mini Kit (Qiagen, Germany) according to the manufacturer's protocol, but including proteinase K incubation (55°C, 15 min). Contaminating DNA was removed by using the RNAse-free

DNAse kit (Qiagen, Germany) and 32 µl of RNA with an average yield of 48 ng/µl were reverse

transcribed with Superscript II reverse transcriptase (Life Technologies, Germany) to obtain

80 µl of cDNA. qPCR analysis was performed in duplicate, using the TaqMan Fast PCR Master

Mix with Taqman assays for bone markers (Life Technologies, Germany) (**Supplementary**

Table 2) or the QuantiTect PCR Kit with self-designed primers for neuronal factors (Qiagen,

Germany) (**Supplementary Table 3**). The amount of cDNA per reaction depended on the

respective target gene and its expression level and was 1-18 ng. Gene expression was

normalized to the housekeeping gene (dCt) and 2^{-dCt} were computed.

ELISA of serum samples

Commercially available ELISA kits were used to determine the serum levels of OC (60-1305,

Immunotopics/Teca, Switzerland), PINP (AC-33F1, Immunodiagnosticsystem, Germany) and

CTX-I (AC-06F1, Immunodiagnosticsystem, Germany) according to the manufacturers'

instructions. Serum was used either undiluted (CTX-I) or at a 1/11 (osteocalcin) or 1/10 (PINP)

dilution.

Micro-CT of bone samples

Analysis of bone microarchitecture by µCT was conducted as previously described [104].

Briefly, bones were measured on a µCT40 (Scanco Medical AG, Brüttisellen, Switzerland),

operated at 55 kVp and 145 µA with an integration time of 200 ms and 2-fold frame averaging.

Images were reconstructed at an isotropic nominal resolution of 10 µm and filtered using a

constrained 3D Gaussian filter (sigma 0.8, support 1) to suppress noise. Masks of the full bone,

cortical (shaft at 56% bone length, 1 mm long) and trabecular region of interest (from 66% to

88% of bone length) were created automatically. Morphometric parameters were calculated

according to standard guidelines such as total volume (TV), bone volume (BV), marrow volume

(MV), apparent volume density (AVD), femur length, cortical area fraction (Ct.Ar/Tt.Ar), cortical

thickness (Ct.Th), trabecular bone volume fraction (BV/TV), specific bone surface (BS/BV),

trabecular thickness (Tb.Th), trabecular spacing (Tb.Sp) and trabecular number (Tb.N) [105].

Due to differences in body weight between the control mice and the MSUS mice, data was

normalized for body weight and body weight adjusted means were compared.

[B] Human Study

Participants

n=240 patients with depressive disorder (ICD-10 F32.x, F33.x, F34.1, F43.21) were recruited

of which *n*=208 completed the initial examination (baseline (t_0), for more details see [5]. A

subsample of *n*=54 patients took part at the fourth follow up measurement (t_4) including DXA

imaging. Patients who fulfilled the following criteria were included: 18 to 65 years of age,

depressive episode (ICD-10 F32.x or F33.x), dysthymia (F34.1) or an adjustment disorder with

prolonged depressive reaction (F43.21), $\geq$ 21 days absenteeism within the last year. Exclusion

criteria were: pregnancy, hormonotherapy (expect contraceptive and thyroid hormone

therapy), inability to fill in a questionnaire, intellectual disabilities (ICD-10 F70-79) or one of the

following diseases: acute infection, endocrine and metabolic disorders, neurological diseases,

dementia (ICD-10 F00-F03), schizophrenia (ICD-10 F20), emotional-unstable personality

disorders (ICD-10 F60.3x), disease of the immune system, substance abuse and dependency

(expect nicotine).

All participants were fully informed in verbal and written form about the intent and content of

the study, and gave their written informed consent. The clinical investigations were conducted

according to the principles of the Declaration of Helsinki. Final ethical approval was provided

on 11/12/2017 from the Ethics Review Board of the University of Potsdam, Germany (number

15/2017).

Study Procedure

Study objectives were investigated at two assessments (t_0 and t_4) of an 8-month observational

multicenter study with four measurement points: baseline (t_0), after 5 weeks (t_1), 5 months (t_2),

8 months (t_3) and an additional follow-up measurement (t_4: M= 15.4 month, SD = 4.5 month

after baseline). At each measurement point, study participants answered a comprehensive

questionnaire (t_0 – t_4) under the supervision of trained study nurses. The questionnaire

comprised demographic characteristics, psychological symptoms (e.g. depression) and

constructs (e.g. stress, early life events, affect, coping), physical ailments, and information

regarding alcohol consumption and medication intake. Blood samples were collected at

baseline (t_0) and 5 months follow-up (t_2) and DXA bone densitometry measurements took place

at the last follow up (t_4).

Psychometric Measures

Early life stress was measured retrospectively comprising experiences of childhood

maltreatment and stressful life events during childhood. The Childhood Trauma Screener

(CTS) [106], a 5-item screening tool of the 28-item Childhood Trauma Questionnaire (CTQ)

[107, 108], assesses five types of childhood maltreatment such as emotional, physical, and

sexual abuse plus emotional and physical neglect. Each item is scored on a five-point rating

scale from "never true" (1) to "very often true" (5). In accordance to Glaesmer et al. [109], we

classified participants at risk if they rated at least mild forms of childhood abuse or neglect. We

additionally controlled for response bias by the 3-item Minimization-Denial subscale from the

CTQ and excluded participants when indicated. Cronbach´s Alpha was previously specified

with 0.76 [110].

Stressful life events were assessed by a modified version of the Inventory of Stressful Life

Events (ILE; [111]). Participants rated 34 adverse life events regarding occurrence, frequency

and year of occurrence. The accumulation of stressful, life-changing events at different stages

of development was counted for every participant, in this analysis focusing on childhood (ILE

$_{child}$ ≤12 years). The scores ranged between 0 (no strain) and 4 (high strain). Cronbach´s Alpha

for ILE was 0.83.

Depressive symptoms and severity were assessed using the Beck Depression Inventory-II

(BDI-II) [112, 113]. Internal consistency in the sample was Cronbach's Alpha 0.89.

Furthermore, confounding factors like sociodemographic and biometric characteristics and

study site were assessed.

Serum bone marker measurement

Blood samples were drawn from the arm in the morning (7-9 am), collected in plain blood collection tubes, allowed to clot at room temperature for 30 min, followed by centrifugation, isolation of serum and freeze-storage until further analysis. Bone-related blood markers (OC, PINP, CTX-I) were analyzed in serum samples by electrochemiluminescence immunoassays "ECLIA" (12149133 122 for Osteocalcin, 03141071 190 for PINP, 11972308 122 for CTX-I, all from F. Hoffmann-La Roche, Ltd., Basel, Switzerland)

DXA measurement

Bone mineral density (BMD) was measured by DXA bone densitometry measurement (Lunar, Prodigy Advance, GE Healthcare, Illinois, USA) in the lumbar spine (lumbar vertebral bodies L1-L4) and both hips. Parameters are given as mean values (M): G (total), H (collum femoris), S (corpus femoris), T (trochanter major), and L1L4 (lumbar vertebral bodies L1-L4).

[A/B] Statistical Analysis

Data consistency was checked, screened for outliers and analyzed descriptively. Continuous variables were also tested for normality by using skewness, kurtosis, omnibus test; variance homogeneity was proven by Variance Ratio and Levene test.

For the examination of **objective 1** (mouse model), student t-test for equal or Aspin-Welch Unequal Variance test were performed, using multiple t-test and adjusted for multiple comparison by Holm-Sidak method (alpha = 0.05). Since body weight was found to be significantly different, ANCOVA models with body weight as covariate were applied for micro-CT data to compute corresponding means at their covariate means (i.e. least square means at covariate mean for body weight of 16.41). In addition, blocking on total body weight was used as a means of accounting for collinearity in analyses of variance and mediation analyses were conducted using linear regression models to identify mediation effects between body weight and bone microstructure. All reported tests are two-sided, and p-values < 0.05 are considered as statistically significant.

Regarding **objective 2** (human model) in bone of depressed patients with and without early life stress) multiple regression models were applied. Statistical models were cross-sectional and controlled for age, gender, study sites and additionally body weight for DXA measurements.

Statistical analyses were conducted using NCSS (NCSS 10, NCSS, LLC. Kaysville, UT), STATISTICA 13, IBM SPSS 25 and the statistical software R [114, 115].

Author Contribution

KW-K planned and coordinated the project, participated in tissue collection, supervised experiments and drafted the manuscript; MR conducted the animal experiments, participated in tissue collection and drafted the manuscript; EC participated in tissue collection, analyzed samples and drafted the manuscript; AB helped to coordinate the human study, analyzed sample and proof-read the manuscript; GAK participated in tissue collection, analyzed samples and drafted the manuscript; TA analyzed samples and proof-read the manuscript; WH conducted statistical analysis of all animal experiments and proof-read the manuscript; DD conducted statistical analysis of all human experiments and proof-read the manuscript; RM provided input in micro-CT analysis and proof-read the manuscript; MAR helped in the coordination of the human study and proof-read the manuscript; IMM planned and coordinated the animal study, supervised experiments, acquired funding and proof-read the manuscript; EMJP helped to coordinate the mouse study, analyzed samples and drafted the manuscript; PMW planned and coordinated the project, conducted human study as PI, supervised experiments, acquired funding and drafted the manuscript.

Acknowledgement

This study was supported by the Deutsche Forschungsgemeinschaft, the Central Research Grant for Innovative Ideas from the University of Potsdam, the German Pension Fund Association (10-R-40.07.05.07.008), the Swiss National Science Foundation (31003A_153147/1) and the ETHZ grant (ETH-10 15-2).

Disclosures

None of the authors has any conflicts of interest or financial disclosures.

References

1. Wippert, P.M., et al., *Stress and Alterations in Bones: An interdisciplinary Perspective.* Frontiers in Endocrinology, 2017. **8**.

2. Grassel, S., *The role of peripheral nerve fibers and their neurotransmitters in cartilage and bone physiology and pathophysiology.* Arthritis Research & Therapy, 2014. **16**(6).

3. Mattia, C., et al., *Bone pain mechanism in osteoporosis: a narrative review.* Clinical Cases in Mineral and Bone Metabolism, 2016. **13**(2): p. 97-100.

4. McEwen, B.S., C. Nasca, and J.D. Gray, *Stress Effects on Neuronal Structure: Hippocampus, Amygdala, and Prefrontal Cortex.* Neuropsychopharmacology, 2016. **41**(1): p. 3-23.

5. Wippert, P.M., et al., *Alterations in Bone Homeostasis and Microstructure Related to Depression and Allostatic Load.* Psychother Psychosom, 2019. **88**(6): p. 383-385.

6. Asada, K., et al., *Age-related changes in afferent responses in sensory neurons to mechanical stimulation of osteoblasts in coculture system.* Am J Physiol Cell Physiol, 2012. **302**(5): p. C757-65.

7. Takahashi, N., et al., *Neuronal TRPV1 activation regulates alveolar bone resorption by suppressing osteoclastogenesis via CGRP.* Sci Rep, 2016. **6**: p. 29294.

8. Mach, D.B., et al., *Origins of skeletal pain: sensory and sympathetic innervation of the mouse femur.* Neuroscience, 2002. **113**(1): p. 155-66.

9. Chartier, S.R., et al., *The Changing Sensory and Sympathetic Innervation of the Young, Adult and Aging Mouse Femur.* Neuroscience, 2018.

10. Alves, C.J., et al., *Bone Injury and Repair Trigger Central and Peripheral NPY Neuronal Pathways.* PLoS One, 2016. **11**(11): p. e0165465.

11. Jiao, K., et al., *Norepinephrine Regulates Condylar Bone Loss via Comorbid Factors.* J Dent Res, 2015. **94**(6): p. 813-20.

12. Yirmiya, R., et al., *Depression induces bone loss through stimulation of the sympathetic nervous system.* Proceedings of the National Academy of Sciences of the United States of America, 2006. **103**(45): p. 16876-16881.

13. Furuzawa, M., et al., *Chewing ameliorates chronic mild stress-induced bone loss in senescence-accelerated mouse (SAMP8), a murine model of senile osteoporosis.* Experimental Gerontology, 2014. **55**: p. 12-18.

14. Tomlinson, R.E., et al., *NGF-TrkA Signaling by Sensory Nerves Coordinates the Vascularization and Ossification of Developing Endochondral Bone.* Cell Rep, 2016. **16**(10): p. 2723-2735.

15. Chen, W.H., et al., *Beta-nerve growth factor promotes neurogenesis and angiogenesis during the repair of bone defects.* Neural Regen Res, 2015. **10**(7): p. 1159-65.

16. Zhuang, Y.F. and J. Li, *Serum EGF and NGF levels of patients with brain injury and limb fracture.* Asian Pac J Trop Med, 2013. **6**(5): p. 383-6.

17. Sabsovich, I., et al., *Effect of anti-NGF antibodies in a rat tibia fracture model of complex regional pain syndrome type I.* Pain, 2008. **138**(1): p. 47-60.

18. Guo, T.Z., et al., *Neuropeptide deficient mice have attenuated nociceptive, vascular, and inflammatory changes in a tibia fracture model of complex regional pain syndrome.* Molecular Pain, 2012. **8**.

19. Sang, X.G., et al., *Analysis of the mechanism by which nerve growth factor promotes callus formation in mice with tibial fracture.* Experimental and Therapeutic Medicine, 2017. **13**(4): p. 1376-1380.

20. Grills, B.L., J.A. Schuijers, and A.R. Ward, *Topical application of nerve growth factor improves fracture healing in rats.* J Orthop Res, 1997. **15**(2): p. 235-42.

21. Hemingway, F., et al., *RANKL-independent human osteoclast formation with APRIL,*
BAFF, NGF, IGF I and IGF II. Bone, 2011. **48**(4): p. 938-44.
22. Camerino, C., et al., *Evaluation of Short and Long Term Cold Stress Challenge of Nerve*
Grow Factor, Brain-Derived Neurotrophic Factor, Osteocalcin and Oxytocin mRNA
Expression in BAT, Brain, Bone and Reproductive Tissue of Male Mice Using Real-
Time PCR and Linear Correlation Analysis. Front Physiol, 2017. **8**: p. 1101.
23. Liu, Q., et al., *Effect of Brain-Derived Neurotrophic Factor on the Neurogenesis and*
Osteogenesis in Bone Engineering. Tissue Eng Part A, 2018.
24. Yang, M., et al., *Genetics association study and functional analysis on osteoporosis*
susceptibility gene BDNF. Yi Chuan, 2017. **39**(8): p. 726-736.
25. Kauschke, V., et al., *Effects of new beta-type Ti-40Nb implant materials, brain-derived*
neurotrophic factor, acetylcholine and nicotine on human mesenchymal stem cells of
osteoporotic and non osteoporotic donors. PLoS One, 2018. **13**(2): p. e0193468.
26. Camerino, C., et al., *Nerve Growth Factor, Brain-Derived Neurotrophic Factor and*
Osteocalcin Gene Relationship in Energy Regulation, Bone Homeostasis and
Reproductive Organs Analyzed by mRNA Quantitative Evaluation and Linear
Correlation Analysis. Front Physiol, 2016. **7**: p. 456.
27. Picard, M., R.P. Juster, and B.S. McEwen, *Mitochondrial allostatic load puts the 'gluc'*
back in glucocorticoids. Nat Rev Endocrinol, 2014. **10**(5): p. 303-10.
28. Picard, M. and B.S. McEwen, *Psychological Stress and Mitochondria: A Conceptual*
Framework. Psychosom Med, 2018. **80**(2): p. 126-140.
29. Kelly, R.R., et al., *Impacts of Psychological Stress on Osteoporosis: Clinical*
Implications and Treatment Interactions. Front Psychiatry, 2019. **10**: p. 200.
30. Lee, S.H., et al., *Short-term antidepressant treatment has long-lasting effects, and*
reverses stress-induced decreases in bone features in rats. Transl Psychiatry, 2019.
9(1): p. 10.
31. Ji-Ye, H., Z. Xin-Feng, and J. Lei-Sheng, *Autonomic control of bone formation: its*
clinical relevance. Handb Clin Neurol, 2013. **117**: p. 161-71.
32. Schiavone, S., et al., *Chronic Psychosocial Stress Impairs Bone Homeostasis: A Study*
in the Social Isolation Reared Rat. Front Pharmacol, 2016. **7**: p. 152.
33. Huang, W.S., et al., *Post-traumatic stress disorder and risk of osteoporosis: A*
nationwide longitudinal study. Stress Health, 2018. **34**(3): p. 440-445.
34. Gershon, A., et al., *The long-term impact of early adversity on late-life psychiatric*
disorders. Curr Psychiatry Rep, 2013. **15**(4): p. 352.
35. Raymond, C., et al., *Early child adversity and psychopathology in adulthood: HPA axis*
and cognitive dysregulations as potential mechanisms. Progress in Neuro-
Psychopharmacology & Biological Psychiatry, 2018. **85**: p. 152-160.
36. Shonkoff, J.P., W.T. Boyce, and B.S. McEwen, *Neuroscience, molecular biology, and*
the childhood roots of health disparities: building a new framework for health
promotion and disease prevention. JAMA, 2009. **301**(21): p. 2252-9.
37. Falgares, G., et al., *Childhood Maltreatment, Pathological Personality Dimensions, and*
Suicide Risk in Young Adults. Front Psychol, 2018. **9**: p. 806.
38. Lindert, J., et al., *Sexual and physical abuse in childhood is associated with depression*
and anxiety over the life course: systematic review and meta-analysis. International
Journal of Public Health, 2014. **59**(2): p. 359-372.
39. Infurna, M.R., et al., *Associations between depression and specific childhood*
experiences of abuse and neglect: A meta-analysis. Journal of Affective Disorders,
2016. **190**: p. 47-55.

40. Nanni, V., R. Uher, and A. Danese, *Childhood Maltreatment Predicts Unfavorable*
Course of Illness and Treatment Outcome in Depression: A Meta-Analysis. American
Journal of Psychiatry, 2012. **169**(2): p. 141-151.
41. Mezuk, B., W.W. Eaton, and S.H. Golden, *Depression and osteoporosis: epidemiology*
and potential mediating pathways. Osteoporosis International, 2008. **19**(1): p. 1-12.
42. Cizza, G., et al., *Depression and Osteoporosis: A Research Synthesis with Meta-*
Analysis. Hormone and Metabolic Research, 2010. **42**(7): p. 467-482.
43. Azuma K, et al., *Chronic Psychological Stress as a Risk Factor of Osteoporosis.* J
UOEH., 2015. **37**(4): p. 245-53.
44. Williams, L.J., et al., *Depression as a risk factor for fracture in women: A 10 year*
longitudinal study. Journal of Affective Disorders, 2016. **192**: p. 34-40.
45. Furlan, P.M., et al., *The role of stress-induced cortisol in the relationship between*
depression and decreased bone mineral density. Biological Psychiatry, 2005. **57**(8): p.
911-917.
46. Stults-Kolehmainen, M.A. and R. Sinha, *The effects of stress on physical activity and*
exercise. Sports Med, 2014. **44**(1): p. 81-121.
47. Yau, Y.H. and M.N. Potenza, *Stress and eating behaviors.* Minerva Endocrinol, 2013.
38(3): p. 255-67.
48. Franklin, T.B., et al., *Epigenetic Transmission of the Impact of Early Stress Across*
Generations. Biological Psychiatry, 2010. **68**(5): p. 408-415.
49. Schmidt, M.V., *Molecular mechanisms of early life stress--lessons from mouse models.*
Neurosci Biobehav Rev, 2010. **34**(6): p. 845-52.
50. McGowan, P.O. and M. Szyf, *The epigenetics of social adversity in early life:*
implications for mental health outcomes. Neurobiol Dis, 2010. **39**(1): p. 66-72.
51. Bohacek, J. and I.M. Mansuy, *Epigenetic inheritance of disease and disease risk.*
Neuropsychopharmacology, 2013. **38**(1): p. 220-36.
52. Gapp, K., et al., *Implication of sperm RNAs in transgenerational inheritance of the*
effects of early trauma in mice. Nat Neurosci, 2014. **17**(5): p. 667-9.
53. Baldock, P.A., et al., *Neuropeptide Y Attenuates Stress-Induced Bone Loss Through*
Suppression of Noradrenaline Circuits. Journal of Bone and Mineral Research, 2014.
29(10): p. 2238-2249.
54. Sousa, D.M., et al., *Neuropeptide Y modulates fracture healing through Y1 receptor*
signaling. J Orthop Res, 2013. **31**(10): p. 1570-8.
55. Teixeira, L., et al., *NPY Revealed as a Critical Modulator of Osteoblast Function In*
Vitro: New Insights Into the Role of Y1 and Y2 Receptors. Journal of Cellular
Biochemistry, 2009. **107**(5): p. 908-916.
56. Tang, P., et al., *NPY and CGRP Inhibitor Influence on ERK Pathway and Macrophage*
Aggregation during Fracture Healing. Cellular Physiology and Biochemistry, 2017.
41(4): p. 1457-1467.
57. Igwe, J.C., et al., *Neuropeptide Y Is Expressed by Osteocytes and Can Inhibit*
Osteoblastic Activity. Journal of Cellular Biochemistry, 2009. **108**(3): p. 621-630.
58. Allison, S.J., et al., *Critical Interplay Between Neuropeptide Y and Sex Steroid*
Pathways in Bone and Adipose Tissue Homeostasis. Journal of Bone and Mineral
Research, 2009. **24**(2): p. 294-304.
59. Sousa, D.M., et al., *Neuropeptide Y Y1 receptor antagonism increases bone mass in*
mice. Bone, 2012. **51**(1): p. 8-16.
60. Ma, W.H., et al., *Neuropeptide Y, substance P, and human bone morphogenetic protein*
2 stimulate human osteoblast osteogenic activity by enhancing gap junction
intercellular communication. Brazilian Journal of Medical and Biological Research,
2015. **48**(4): p. 299-307.

61. Long, H., et al., *Neuropeptide Y innervation during fracture healing and remodeling.* Acta Orthopaedica, 2010. **81**(5): p. 639-646.

62. Cherruau, M., et al., *Chemical sympathectomy-induced changes in TH-, VIP-, and CGRP-immunoreactive fibers in the rat mandible periosteum: Influence on bone resorption.* Journal of Cellular Physiology, 2003. **194**(3): p. 341-348.

63. Lundberg, P., et al., *Vasoactive intestinal peptide (VIP)/pituitary adenylate cyclase-activating peptide receptor subtypes in mouse calvarial osteoblasts: Presence of VIP-2 receptors and differentiation-induced expression of VIP-1 receptors.* Endocrinology, 2001. **142**(1): p. 339-347.

64. Ransjo, M., et al., *Microisolated mouse osteoclasts express VIP-1 and PACAP receptors.* Biochemical and Biophysical Research Communications, 2000. **274**(2): p. 400-404.

65. Nagata, A., et al., *cAMP Activation by PACAP/VIP Stimulates IL-6 Release and Inhibits Osteoblastic Differentiation Through VPAC2 Receptor in Osteoblastic MC3T3 Cells.* Journal of Cellular Physiology, 2009. **221**(1): p. 75-83.

66. Persson, E. and U.H. Lerner, *The Neuropeptide VIP Regulates the Expression of Osteoclastogenic Factors in Osteoblasts.* Journal of Cellular Biochemistry, 2011. **112**(12): p. 3732-3741.

67. Wei, T.P., et al., *Post-junctional facilitation of Substance P signaling in a tibia fracture rat model of complex regional pain syndrome type I.* Pain, 2009. **144**(3): p. 278-286.

68. Aoki, M., K. Tamai, and K. Saotome, *Substance P-Related and Calcitonin-Gene-Related Peptide-Immunofluorescent Nerves in the Repair of Experimental Bone Defects.* International Orthopaedics, 1994. **18**(5): p. 317-324.

69. Goto, T., et al., *Light- and electron-microscopic study of the distribution of axons containing substance P and the localization of neurokinin-1 receptor in bone.* Cell and Tissue Research, 1998. **293**(1): p. 87-93.

70. Wang, L.P., et al., *Substance P stimulates bone marrow stromal cell osteogenic activity, osteoclast differentiation, and resorption activity in vitro.* Bone, 2009. **45**(2): p. 309-320.

71. Mori, T., et al., *Substance p regulates the function of rabbit cultured osteoclast; Increase of intracellular free calcium concentration and enhancement of bone resorption.* Biochemical and Biophysical Research Communications, 1999. **262**(2): p. 418-422.

72. Sohn, S.J., *Substance P upregulates osteoclastogenesis by activating nuclear factor kappa B in osteoclast precursors.* Acta Oto-Laryngologica, 2005. **125**(2): p. 130-133.

73. Matayoshi, T., et al., *Neuropeptide substance P stimulates the formation of osteoclasts via synovial fibroblastic cells.* Biochemical and Biophysical Research Communications, 2005. **327**(3): p. 756-764.

74. Adamus, M.A. and Z.J. Dabrowski, *Effect of the neuropeptide substance P on the rat bone marrow-derived osteogenic cells in vitro.* J Cell Biochem, 2001. **81**(3): p. 499-506.

75. Liu, H.J., et al., *Substance P Promotes the Proliferation, but Inhibits Differentiation and Mineralization of Osteoblasts from Rats with Spinal Cord Injury via RANKL/OPG System.* PLoS One, 2016. **11**(10): p. e0165063.

76. Shih, C. and G.W. Bernard, *Neurogenic substance P stimulates osteogenesis in vitro.* Peptides, 1997. **18**(2): p. 323-6.

77. Goto, T., et al., *Substance P stimulates late-stage rat osteoblastic bone formation through neurokinin-1 receptors.* Neuropeptides, 2007. **41**(1): p. 25-31.

78. Kook, Y.A., et al., *Effects of substance P on osteoblastic differentiation and heme oxygenase-1 in human periodontal ligament cells.* Cell Biology International, 2009. **33**(3): p. 424-428.

79. Azuma, H., et al., *Substance P enhances the inhibition of osteoblastic cell differentiation induced by lipopolysaccharide from Porphyromonas gingivalis.* Journal of Periodontology, 2004. **75**(7): p. 974-981.

80. Kingery, W.S., et al., *A substance P receptor (NK1) antagonist enhances the widespread osteoporotic effects of sciatic nerve section.* Bone, 2003. **33**(6): p. 927-936.

81. Zheng, X.F., et al., *Inhibition of substance P signaling aggravates the bone loss in ovariectomy-induced osteoporosis.* Progress in Biophysics & Molecular Biology, 2016. **122**(2): p. 112-121.

82. Ding, W.G., et al., *Changes of substance P during fracture healing in ovariectomized mice.* Regulatory Peptides, 2010. **159**(1-3): p. 28-34.

83. Lerner, U.H. and E. Persson, *Osteotropic effects by the neuropeptides calcitonin gene-related peptide, substance P and vasoactive intestinal peptide.* J Musculoskelet Neuronal Interact, 2008. **8**(2): p. 154-65.

84. Terreni, A. and P. Pezzati, *Biochemical markers in the follow-up of the osteoporotic patients.* Clin Cases Miner Bone Metab, 2012. **9**(2): p. 80-4.

85. Vasikaran, S., et al., *Markers of bone turnover for the prediction of fracture risk and monitoring of osteoporosis treatment: a need for international reference standards.* Osteoporos Int, 2011. **22**(2): p. 391-420.

86. Yoshimura, N., et al., *Biochemical markers of bone turnover as predictors of osteoporosis and osteoporotic fractures in men and women: 10-year follow-up of the Taiji cohort.* Mod Rheumatol, 2011. **21**(6): p. 608-20.

87. Garnero, P., et al., *Biochemical markers of bone turnover, endogenous hormones and the risk of fractures in postmenopausal women: the OFELY study.* J Bone Miner Res, 2000. **15**(8): p. 1526-36.

88. Eastell, R., et al., *Relationship of early changes in bone resorption to the reduction in fracture risk with risedronate.* Journal of Bone and Mineral Research, 2003. **18**(6): p. 1051-1056.

89. Callewaert, F., et al., *Sexual dimorphism in cortical bone size and strength but not density is determined by independent and time-specific actions of sex steroids and IGF-1: evidence from pubertal mouse models.* J Bone Miner Res, 2010. **25**(3): p. 617-26.

90. Goel, N. and T.L. Bale, *Sex differences in the serotonergic influence on the hypothalamic-pituitary-adrenal stress axis.* Endocrinology, 2010. **151**(4): p. 1784-94.

91. Takeda, T., M. Hosokawa, and K. Higuchi, *Senescence-accelerated mouse (SAM): a novel murine model of accelerated senescence.* J Am Geriatr Soc, 1991. **39**(9): p. 911-9.

92. Paccione, M.F., et al., *A mouse model of mandibular osteotomy healing.* Journal of Craniofacial Surgery, 2001. **12**(5): p. 444-450.

93. Dang, R., et al., *Predictable chronic mild stress promotes recovery from LPS-induced depression.* Mol Brain, 2019. **12**(1): p. 42.

94. Furuzawa, M., et al., *Chewing ameliorates chronic mild stress-induced bone loss in senescence-accelerated mouse (SAMP8), a murine model of senile osteoporosis.* Exp Gerontol, 2014. **55**: p. 12-8.

95. Heim, C., et al., *The dexamethasone/corticotropin-releasing factor test in men with major depression: role of childhood trauma.* Biol Psychiatry, 2008. **63**(4): p. 398-405.

96. Heim, C., et al., *Neurobiological and psychiatric consequences of child abuse and neglect.* Dev Psychobiol, 2010. **52**(7): p. 671-90.

97. Saveanu, R.V. and C.B. Nemeroff, *Etiology of Depression: Genetic and Environmental Factors.* Psychiatric Clinics of North America, 2012. **35**(1): p. 51-+.

98. Picard, M., et al., *An energetic view of stress: Focus on mitochondria.* Front Neuroendocrinol, 2018. **49**: p. 72-85.

99. Sophocleous, A. and A.I. Idris, *Rodent models of osteoporosis.* Bonekey Rep, 2014. **3**: p. 614.

100. Halloran, B.P., et al., *Changes in bone structure and mass with advancing age in the male C57BL/6J mouse.* Journal of Bone and Mineral Research, 2002. **17**(6): p. 1044-1050.

101. Bohacek, J., et al., *Hippocampal gene expression induced by cold swim stress depends on sex and handling.* Psychoneuroendocrinology, 2015. **52**: p. 1-12.

102. Rommel, F.R., et al., *Suitability of Nicotinic Acetylcholine Receptor alpha7 and Muscarinic Acetylcholine Receptor 3 Antibodies for Immune Detection: Evaluation in Murine Skin.* J Histochem Cytochem, 2015. **63**(5): p. 329-39.

103. Hendrix, S., et al., *Skin and hair follicle innervation in experimental models: a guide for the exact and reproducible evaluation of neuronal plasticity.* Exp Dermatol, 2008. **17**(3): p. 214-27.

104. Kohler, T., et al., *Compartmental bone morphometry in the mouse femur: Reproducibility and resolution dependence of microtomographic measurements.* Calcified Tissue International, 2005. **77**(5): p. 281-290.

105. Bouxsein, M.L., et al., *Guidelines for Assessment of Bone Microstructure in Rodents Using Micro-Computed Tomography.* Journal of Bone and Mineral Research, 2010. **25**(7): p. 1468-1486.

106. Glaesmer, H., et al., *Der Childhood Trauma Screener (CTS) – Entwicklung und Validierung von Schwellenwerten zur Klassifikation. [The Childhood Trauma Screener (CTS) – Development and Validation of Cut-Off-Scores for Classificatory Diagnostics].* Psychiatrische Praxis, 2013. **40**(04): p. 220-226.

107. Wingenfeld, K., et al., *Die deutsche Version des Childhood Trauma Questionnaire (CTQ): Erste Befunde zu den psychometrischen Kennwerten. [The German version of the Childhood Trauma Questionnaire (CTQ): Preliminary psychometric properties].* Psychotherapie, Psychosomatik, und Medizinische Psychologie, 2010. **60**(11): p. 442-450.

108. Bernstein, D. and L. Fink, *Childhood Trauma Questionnaire (CTQ): A retrospective self-Report questionnaire and manual*, T.P. Corporation, Editor. 1998: San Antonio.

109. Glaesmer, H., et al., *[The childhood trauma screener (CTS) - development and validation of cut-off-scores for classificatory diagnostics].* Psychiatr Prax, 2013. **40**(4): p. 220-6.

110. Grabe, H.J., et al., *[A brief instrument for the assessment of childhood abuse and neglect: the childhood trauma screener (CTS)].* Psychiatr Prax, 2012. **39**(3): p. 109-15.

111. Siegrist, J. and S. Geyer, *ILE–Inventar zur Erfassung lebensverändernder Ereignisse*, in *Diagnostische Verfahren in der Psychotherapie*, Hogrefe, Editor. 2002: Göttingen. p. 211-213.

112. Beck, A.T., R.A. Steer, and G.K. Brown, *Manual for the Beck Depression Inventory-II.* 1996, San Antonio: T.P. Corporation.

113. Wintjen, L. and F. Petermann, *Beck-Depressions-Inventar Revision (BDI-II).* Zeitschrift Fur Psychiatrie Psychologie Und Psychotherapie, 2010. **58**(3): p. 243-245.

114. Muche, R., S. Lanzinger, and M. Rau, *Medizinische Statistik mit R und Excel.* 2011: Springer.

115. Team, R.C. *R foundation for Statistical Computing.* 2014.

Figures with Legends

*Figure 1: **Increased bone innervation in MSUS mice.** Representative images of the third tail*

bone from control and MSUS mice labeled with PGP 9.5 pan-neuronal marker for

immunofluorescence histomorphometry of nerve fibers (NF - red) in bone. Nuclei are

*counterstained with DAPI (blue), mast cells with FITC-avidin (green) **(A)**. Shown are the*

increased mean numbers of immune-positive profiles per compartment unit of single NF in

*bone **(B)**, NF bundles at entry to bone **(C)** and single NF in bone marrow **(D)**. Data (n=8*

Control, n=6-7 MSUS) are expressed as Min to Max of number of positive profiles per

*compartment unit. Significance level: **p < 0.01 and *p < 0.05 between indicated groups.*

Figure 2A*: **Differential gene expression of neuronal receptors in bone between control***

and MSUS mice*. Shown are the gene expression of neurotrophin receptors **(A)**, noradrenergic*

*receptors **(B)**, cholinergic receptors **(C)** and receptors involved in sensory innervation **(D)**. Data*

(n=8 per group) are expressed as Min to Max of 2^{-dCt}. Significance level: ***p < 0.001, **p <

0.01 and *p < 0.05 between indicated groups.

Figure 2B: Differential gene expression of neuronal ligands in bone between control and MSUS mice. Shown are the gene expression of neurotrophins (A), noradrenergic ligands (B), cholinergic ligands (C) and ligands involved in sensory innervation (D). Data (n=8 per group) are expressed as Min to Max of 2^{-dCt}. Significance level: ***p < 0.001 between indicated groups.

*Figure 3: **Bone formation and resorption markers in serum of mice**. Shown are serum*

*concentration measurements of osteocalcin **(A)**, procollagen type 1 N-terminal propeptide*

*(PINP) **(B)** and c-terminal telopeptide of type I collagen (CTX-I) **(C)** in mice. Data (n=14 per*

group) are expressed as Min to Max of concentration in serum [ng/ml]. Significance level: **p

< 0.01 between indicated groups.

Figure 4*: **Gene expression of extracellular matrix in bone of mice.** Shown are the gene*

expressions of tissue-specific extracellular matrix markers in bone of controls and MSUS.

*Genes include osteocalcin **(A)**, osteoprotegerin **(B)**, osteopontin **(C)** and sclerostin **(D)**. Data*

(n=8 per group) are expressed as Min to Max of 2^{-dCt}. Significance level: all p > 0.05

Figure 5*: **Bone mineral density measurements in human patients.** Shown is the BMD in*

G (total), H (collum femoris), L1-L4 (lumbar vertebralis bodies), S (corpus femoris), and T

(trochanter major), stratified for at risk and not at risk for childhood maltreatment (not controlled

for gender, age, body weight). Data (n=18) are expressed as mean and standard deviation of

BMD, stratified for CTS (at risk for abuse or neglect). Significance level: *p < 0.05, and #p <

0.1 between indicated groups.

Figure 6: **MSUS is a mouse model of early life stress**. *Unpredictable maternal separation and unpredictable maternal stress is an animal model for early life stress. Naïve males are mated with naïve females for 1-7 days, then males are removed. Dams gestate for about 21 days in normal conditions until delivery. The MSUS paradigm (red bar in the timeline) starts at postnatal day 1 (PND1) and lasts until PND14. During MSUS, pups are unpredictably separated from their mothers for 3 hours each day at different times of the dark cycle (lights off from 8am to 8pm). During separation, the mother is stressed unpredictably by forced swim or restraint as shown in the timeline at the bottom. Pups and dam are left undisturbed from PND15 until weaning at PND21. Gray mice are naïve without any previous stress exposure. Blue pups are controls without any stress exposure. Red pups are exposed to MSUS.*

865 **Tables**

867 ***Table 1***. *Main effects (regression coefficient b) of childhood maltreatment (abuse or neglect) and stressful life events during childhood (≤12 years;*

868 *ILE $_{child}$) on bone serum marker (osteocalcin (OC), procollagen type 1 N-terminal propeptide (PINP), and C-terminal telopeptide of type I collagen*

869 *(CTX-I)) and bone mineral density (BMD)*

	P1NP			OC			CTX-I				
	b	*95%CI*	*p*	*b*	*95%CI*	*p*	*b*	*95%CI*	*p*		
CTS [a]	9.27	-1.43; 19.97	0.10#	2.87	-0.56; 6.30	0.11	**0.10**	0.01; 0.18	0.05*		
CTS abuse	**11.42**	1.86; 20.99	0.03*	**2.51**	-0.65; 5.67	0.13	0.06	0.02; 0.14	0.16		
CTS neglect	-6.26	-16.45; 3.93	0.24	0.32	-3.63; 2.98	0.85	0.03	-0.06; 0.12	0.50		
ILE $_{child}$ [b]	**8.37**	0.71; 16.03	0.04*	**2.28**	-0.10; 4.65	0.06#	**0.06**	-0.01; 0.11	0.06#		

	BMD_ L1L4 [c]			BMD_T			BMD_S			BMD_G			BMD_H		
	b	*95%CI*	*p*	*b*	*95%CI*	*p*	*b*	*95%CI*	*p*	*b*	*95%CI*	*p*	*b*	*95%CI*	*p*
CTS	-0.07	-0.23; 0.08	0.40	-0.09	-0.22; 0.04	0.20	-0.09	-0.25; 0.08	0.33	-0.10	-0.23; 0.03	0.17	**-0.10**	-0.22; 0.02	0.12
CTS abuse	-0.07	-0.26, 0.12	0.48	**-0.15**	-0.28; -0.02	0.05*	-0.13	-0.32, 0.06	0.22	**-0.15**	-0.29, -0.01	0.07#	**-0.12**	-0.26, 0.01	0.10
CTS neglect	-0.06	-0.21, 0.10	0.51	-0.06	-0.20, 0.07	0.39	-0.07	-0.24, 0.10	0.45	-0.07	-0.21, 0.07	0.38	-0.03	-0.16, 0.10	0.68
ILE $_{child}$	-0.18	-0.46, 0.10	0.25	**-0.26**	-0.44, -0.07	0.03*	-0.18	-0.48, 0.11	0.26	**-0.22**	-0.43, -0.01	0.08#	**-0.17**	-0.36, 0.03	0.14

Multiple regression models, adjusted for age, gender, and study sides. Significant Regression coefficients b are bold. Significance level: *p < 0.05, and #p < 0.1.
[a] Childhood Trauma Screener (CTS) at least at risk for one type of neglect or abuse (sexual, physical, emotional abuse, physical, emotional neglect)
[b] Inventory of Stressful Life Events; stressful life events during childhood (≤12 years)
[c] Bone mineral density (BMD) was measured as mean value (M) in the lumbar spine L1-L4 (lumbar vertebralis bodies), T (trochanter major), S (corpus femoris), G (total), and H (collum femoris)