Predicting cancer prognosis and drug response from the tumor microbiome

Leandro C. Hermida[1,2†], E. Michael Gertz[1†], Eytan Ruppin[1*].

[1] Cancer Data Science Laboratory (CDSL), National Cancer Institute (NCI), National Institutes

of Health (NIH), Bethesda, MD, USA.

[2] Department of Computer Science, University of Maryland, College Park, MD, USA.

[†] Equally contributing first authors

[*] Corresponding author (eytan.ruppin@nih.gov)

Abstract

Tumor gene expression is predictive of patient prognosis in some cancers. However, RNA-seq and whole genome sequencing data contain not only reads from host tumor and normal tissue, but also reads from the tumor microbiome, which can be used to infer the microbial abundances in each tumor. Here, we show that tumor microbial abundances, alone or in combination with tumor gene expression data, can predict cancer prognosis and drug response to some extent – microbial abundances are significantly less predictive of prognosis than gene expression, although remarkably, similarly as predictive of drug response, but in mostly different cancer-drug combinations. Thus, it appears possible to leverage existing sequencing technology, or develop new protocols, to obtain more non-redundant information about prognosis and drug response from RNA-seq and whole genome sequencing experiments than could be obtained from gene expression or mutation data alone.

Introduction

The Cancer Genome Atlas (TCGA), available from the NCI Genomic Data Commons[1], provides RNA-seq and whole genomic sequencing (WGS) data for thousands of cases across dozens of cancer types. RNA-seq data is typically used to measure the expression of human genes, and there is a long history linking tumor gene expression to cancer outcomes[2-8]. Milanez-Almeida et al.[9] recently showed that gene expression from TCGA RNA-seq data could predict overall survival (OS) or progression-free interval (PFI) better than classical clinical prognostic covariates – age at diagnosis, gender, and tumor stage. Importantly, Milanez-Almeida et al. used a data-driven machine learning (ML) based approach that selected features that were predictive of and correlated with prognosis, rather than an approach based on classical statistics or biological knowledge that chose features *a priori*. Though RNA-seq and WGS data not only

contain human sequencing reads, but also reads from the local microbiome that are typically

filtered out from the data when analyzing human gene expression. Our central research questions

then were, 1) does a data-driven ML approach reveal that tumor microbial abundances in TCGA

data, quantified from these reads, are predictive of cancer prognosis or drug response, 2) what

microbial genera are potentially predictive biomarkers of prognosis or drug response, 3) how do

these models compare to equivalent models based on tumor gene expression data, and 4) does

combining both microbial abundance and gene expression features produce models and select

combinations of gene and microbial genera features that are more predictive of prognosis or drug

response than models from each individual data type?

Poore et al.[10] recently published estimated, decontaminated, and normalized microbial

abundances for the TCGA cohort, derived from either WGS or RNA-seq data. We used these

abundances, taken directly from the Poore et al. dataset, to build predictive models of prognosis

and drug response for TCGA. As a positive control, we also showed that our prognosis ML

modeling methods, which differed somewhat from Milanez-Almeida et al.[9], identified a similar

set cancers and outcomes for which gene expression was predictive of prognosis.

Results

Tumor microbial abundances are substantially less predictive of prognosis than gene

expression

We built OS and PFI gene expression ML models of 32 TCGA tumor types (cohort

information in **Supplementary Data 1**) using the Coxnet[11] algorithm, which jointly selects the

most predictive subset of features via cross-validation (CV) while simultaneously being able to

control for prognostic clinical covariates. In our models, we included and controlled for the

clinical covariates age at diagnosis, gender, and tumor stage. For comparison, we also built

standard Cox regression models based on the clinical covariates alone. We evaluated the

predictive performance of our models using Harrell's concordance index (C-index), which is a

metric of survival model predictive accuracy. Each model analysis generated 100 model

instances and C-index scores from different randomly shuffled train-test CV splits on the data.

We found 33 OS and PFI models for 21 tumor types that had a mean C-index score ≥ 0.6 and

significantly outperformed their corresponding clinical covariate-only models (**Fig. 1a & c,**

Supplementary Fig. 1a, 2a). Our models were predictive of prognosis in 11 of the same 13

tumor types that were reported by Milanez-Almeida et al.[9] (**Supplementary Table 1**). We did

not analyze one tumor type that Milanez-Almeida did, acute myeloid leukemia (LAML), because

Poore et al. excluded it from their analysis. Among the cancers and outcomes that Milanez-

Almeida et al. analyzed, our methodology produced predictive models for four additional tumor

types: breast cancer (BRCA), cervical squamous cell carcinoma (CESC), sarcoma (SARC), and

uterine corpus endometrial carcinoma (UCEC), as well as quite a few predictive models for

additional cancers and outcomes that were not analyzed in their study (**Supplementary Table 1**).

We also evaluated prognosis model performance by calculating the time-dependent,

cumulative/dynamic area under the curve $(AUC^{C/D}(t))$[12,13], which is an extension of the area

under the receiver operating characteristic (AUROC) for continuous outcomes and can provide a

more detailed resolution picture of predictive performance throughout the test outcome time

range compared to the C-index score. Although 33 of our OS and PFI gene expression models

had a statistically significant C-index score improvement compared to clinical covariates alone,

only 22 of these models showed an improvement in $AUC^{C/D}(t)$, where the improvement in mean

$AUC^{C/D}(t)$ over the entire test time range after diagnosis was ≥ 0.025 (**Supplementary Fig. 1b,**

2b).

We applied Coxnet[11] using the same methodology to build prognosis models using the

microbial abundance estimates provided by Poore et al.[10]. We found six microbial abundance

models that had a mean C-index score ≥ 0.6 and significantly outperformed their corresponding

clinical covariate-only models (**Fig. 1b & c, Supplementary Fig. 3a**). We found that in only two

of the six models, microbial abundances outperformed clinical covariates alone in terms of

$AUC^{C/D}(t)$, where the improvement in mean $AUC^{C/D}(t)$ over the entire test time range after

diagnosis was ≥ 0.025 (**Supplementary Fig. 3b**). In adrenocortical carcinoma (ACC), microbial

features predicted OS significantly better than clinical prognostic covariates starting at

approximately 6 years after diagnosis. In CESC, microbial abundances predicted OS better than

clinical covariates from approximately 6 months to 10 years after diagnosis. In contrast to a

recent report[14] on intratumor microbes predicting prognosis in TCGA thyroid carcinoma

(THCA), we did not find that Poore et al. tumor microbial abundances estimated from TCGA

were predictive of OS or PFI using our computational methods. A few possible reasons for this

are that Gnanasekar et al. analyzed the THCA cohort by tumor subtype and they applied different

methods, including 1) they used harmonized and normalized GDC TCGA data instead of legacy

TCGA followed by normalization and batch effect correction, 2) they applied different methods

for extraction of microbial reads and decontamination compared to Poore et al., and 3) they used

classical statistics on all of their data to infer correlations between features and prognosis instead

of a regularized ML model approach focused on fitting models on training data that have the

potential to generalize better and discover features that are predictive of prognosis on unseen

data. Overall, we found that tumor microbial abundances from Poore et al. were only marginally

predictive of prognosis across the TCGA cohort, and that gene expression was a significantly

more powerful predictor of prognosis (**Fig. 1, Supplementary Figs. 1-3**).

Figure 1. Performance of gene expression and microbial abundance prognosis prediction models where features add predictive power to clinical covariates. Mean C-index scores for **(a)** gene expression with clinical covariate models (orange) and **(b)** microbial abundance with clinical covariate models (blue) vs clinical covariate-only models (grey). Error bars denote standard deviations. Significance: * ≤ 0.01, ** ≤ 0.001, *** ≤ 0.0001. **(c)** C-index score violin density plots for the six models where microbial abundance with clinical covariate features outperform clinical covariate-only models. Corresponding gene expression models shown for comparison. Lines connecting points (light grey) represent score pairs from same train-test split on the data. Mean C-index scores shown as red dots with red lines connecting the means. Significance for the prediction improvement over clinical covariate-only models was calculated using a two-sided Wilcoxon signed-rank test and adjusted for multiple testing using the Benjamini-Hochberg method with adjusted p-values shown at top. Adjusted p-values colored in red signify difference where clinical covariate-only model is better.

Tumor microbial abundances are predictive of chemotherapy drug response in some

cancers and in mostly different cancer-drug combinations than gene expression

We next asked whether tumor microbial abundances from pre-treatment biopsies could

predict drug response better than the clinical covariates age at diagnosis, gender and tumor stage

alone. Drug response data were obtained from the TCGA Research Network as described in

Methods (cohorts shown in **Supplementary Data 1**). Cases with complete response (CR) or

partial response (PR) were labeled as responders and those with stable disease (SD) or

progressive disease (PD) as non-responders. Thirty TCGA cancer-drug combinations met our

minimum dataset size thresholds. We built drug response models using three different ML

methods: 1) a variant of the linear support vector machine recursive feature elimination (SVM-

RFE) algorithm[15] that we developed, 2) logistic regression (LGR) with elastic net[16] (L1 + L2)

penalties and embedded feature selection, and 3) logistic regression with an L2 penalty and

limma[17] (for tumor microbial and combination datasets) or edgeR[18,19] (for RNA-seq count

datasets) differential abundance/expression feature scoring and selection methods (see Methods

for details). All three ML methods unconditionally included the clinical covariates – age at

diagnosis, gender, and tumor stage – in the model (bypassing feature selection) while selecting

the most predictive subset of microbial abundance or gene expression features. For comparison,

we built standard linear SVM or LGR models using the clinical covariates alone. We evaluated

the predictive performance of drug response models using AUROC. Each analysis generated 100

model instances, AUROC, and area under the precision-recall curve (AUPRC) scores from

different random train-test CV splits.

We found five cancer-drug microbial abundance models had a mean AUROC score ≥ 0.6 and

performed better than clinical covariates alone in at least two out of three ML methods (**Fig. 2**).

Three of these cancer-drug combinations involved stomach adenocarcinoma (STAD). We

performed the same drug response modeling using TCGA gene expression data and here we

found six cancer-drug combinations had a mean AUROC score ≥ 0.6 and significantly

outperformed their corresponding clinical covariate-only models in at least two out of three ML

methods (**Fig. 3**). Only one cancer-drug combination, SARC docetaxel, overlapped between the

microbial abundance and gene expression drug response model results, suggesting that tumor

microbial abundances have independent predictive power. Overall, our results support the notion

that the tumor microbiome may contain information that is predictive of drug response in some

cancers, consistent with recent reports[20,21]. We also found there was considerable overlap in the

selected microbial abundance and gene expression features reported by each ML method (**Fig. 4a**

& c). We also frequently found a significant correlation between the feature importance rankings

reported by each ML method when comparing the two most significant methods in each cancer-

drug combination (**Fig. 4b & d**).

Figure 2. Performance of microbial abundance drug response prediction models in the five cancer-drug combinations where models performed better than clinical covariates alone. (a) Mean AUROC scores for microbial abundance with clinical covariate models (blue) vs clinical covariate-only models (grey) and **(b)** mean AUROC scores for each ML method. In both **(a)** and **(b)** error bars denote standard deviations. Significance: * ≤ 0.01, ** ≤ 0.001, *** ≤ 0.0001. **(c)** Violin density plots of AUROC scores (top row) for microbial abundance with clinical covariate models vs clinical covariate-only models. Lines connecting points (light grey) represent score pairs from same train-test split on the data. Mean AUROC scores are shown as red dots connected by red lines. **(d)** Mean ROC (middle row, blue) and **(e)** precision-recall (PR) curves (bottom row, purple) for microbial abundance with clinical covariate models vs clinical covariate-only models (grey). Mean AUROC and AUPRC scores shown in legends and shaded areas denote standard deviations. Significance for the prediction improvement over clinical covariate-only models was calculated using a two-sided Wilcoxon signed-rank test and adjusted for multiple testing using the Benjamini-Hochberg method with adjusted p-values shown at top of violin plots in **(c)**.

Figure 3. Performance of gene expression drug response prediction models in the six cancer-drug combinations where models performed better than clinical covariates alone. (a) Mean AUROC scores for gene expression with clinical covariate models (orange) vs clinical covariate-only models (grey) and **(b)** mean AUROC scores for each ML method. In both **(a)** and **(b)** error bars denote standard deviations. Significance: * ≤ 0.01, ** ≤ 0.001, *** ≤ 0.0001. **(c)** Violin density plots of AUROC scores (top row) for gene expression with clinical covariate models vs clinical covariate-only models. Lines connecting points (light grey) represent score pairs from same train-test split on the data. Mean AUROC scores are shown as red dots connected by red lines. **(d)** Mean ROC (middle row, orange) and **(e)** precision-recall (PR) curves (bottom row, green) for gene expression with clinical covariate models vs clinical covariate-only models (grey). Mean AUROC and AUPRC scores shown in legends and shaded areas denote standard deviations. Significance for the prediction improvement over clinical covariate-only models was calculated using a two-sided Wilcoxon signed-rank test and adjusted for multiple testing using the Benjamini-Hochberg method with adjusted p-values shown at top of violin plots in **(c)**.

Figure 4. Comparison of top-ranked selected drug response model features by each ML method. For each drug response model, we selected the two best ML methods by significance for the prediction improvement over their respective clinical covariate-only model. **(a, c)** Venn diagrams for microbiome **(a)** or gene expression **(c)** models comparing the number of features individually selected by each ML method, and the intersection of the two ML methods. **(b, d)** Spearman rank correlation plots microbiome **(b)** or gene expression **(d)** models showing that the median rank of features (among the 100 model instances in which the feature was selected) often correlated between the two most significant ML methods. The best method is shown on the x-axis, the second best on the y-axis.

180 **Combining tumor microbial abundance and gene expression features add a modest**

181 **predictive improvement in some cancers**

182   Finally, we investigated if models built from combining microbial abundance and gene

183 expression features would result in an improvement in predictive power over their corresponding

184 single data type models. Combining data types resulted in a modest predictive improvement in

185 only three prognosis models: SARC OS, STAD PFI, and THYM OS (**Supplementary Fig. 4a**).

186 Although this improvement was not statistically significant in terms of C-index score, the

187 $AUC^{C/D}(t)$ metric showed a clear improvement in prognostic predictive power for the combined

188 data type models, where the improvement in mean $AUC^{C/D}(t)$ over the entire time range after

189 diagnosis was ≥ 0.025 compared to their respective single data type models. We also found five

190 combined data type drug response models which performed significantly better than clinical

191 covariates alone, although none of these models reached statistical significance when compared

192 to the their respective single data type model in terms of improvement in AUROC score, but one

193 of these models, for BLCA cisplatin, did show an improvement in AUROC ≥ 0.025 compared to

194 its corresponding single data type models (**Supplementary Fig. 4b-c**).

195 **Feature analysis reveals a wide range of predictive microbial genera**

196   To learn more about the most predictive features, we determined the top microbial genera

197 and top genes (**Supplementary Data 2**) selected by each of the significantly predictive microbial

198 abundance and gene expression models, respectively, according to their selection frequency and

199 model coefficients across the 100 model instances from each analysis. There were 428 distinct

200 microbial genera appearing in at least one prognosis or drug response model. Of these 428

201 genera, 160 were individually significantly predictive of prognosis or drug response by a

202 Wilcoxon test, indicating that the other genera were significantly predictive in combination. The

203 median number of genera selected per model was 52, with a minimum of 3 (BRCA docetaxel)

204   and a maximum of 78 (STAD cisplatin). Of the 428 genera, 95 were selected in more than one

205   model and only 13 were selected in more than two models. This is consistent with the

206   observation of Nejman et al.[22] that the tumor microbiome is tumor type specific. The predictive

207   genera we found span all non-eukaryotic domains of life, in total encompassing 365 bacterial, 17

208   archaeal, and 46 viral genera (**Supplementary Data 2**).

Discussion

In summary, we find that the microbial abundance estimates generated by Poore et al.[10] are

predictive of cancer patient prognosis and response to chemotherapy in a subset of tumor types,

survival outcomes, and treatments. Machine learning methods, such as those applied in this

study, are not able to infer causality, but only inform on the positive or negative predictive

associations covariates have with the response variable. The potential causal role that those may

play in determining patient prognosis or drug response can only be ascertained via dedicated

mechanistic studies. Overall, in terms of the number of significant models, based on their cross-

validated C-index or AUROC scores and improvement over clinical covariates alone, the tumor

microbiome is considerably less predictive than the tumor human transcriptome at predicting

patient prognosis, but notably, performs similarly to gene expression at predicting chemotherapy

response and in mostly different cancer-drug combinations. Our investigation motivates future

studies investigating the role of the tumor microbiome in predicting the response to targeted

therapies and immunotherapies.

There are also limitations to our current study. With respect to the drug response models, the

relatively small cohort size of most TCGA cancers where drug response data were available and

also the often small responder or non-responder case numbers within these cohorts does have an

impact on the interpretability of the results (see **Supplementary Data 1** for cohort size

information). We have attempted to overcome this limitation as best as possible by applying

robust and advanced ML techniques at every level of model building and evaluation, including

fitting and evaluating 100 randomly shuffled model instances, performing an exhaustive grid

search for hyperparameter optimization and model selection, using a nested cross-validation

(CV) strategy, utilizing advanced stratified and replicate grouping CV iterators, and applying

both sample and class weighting to the learning algorithm and scoring methods (see Methods for

full details).

There are further limitations to this study inherited from limitations in the data, originally

raised by Poore et al.[10] First, the study was retrospective, using existing data from the TCGA. As

such, it did not involve any specific protocols to capture microbial reads or to control for

contamination. Second, decontamination of such retrospective data is a highly involved and

dataset-specific process, which they made great effort to validate. Poore et al. conclude from this

validation that the retrospective study of TCGA was successful, and that similar retrospective

studies would be valuable. A third point, which they touch on briefly, is that the protocols that

were used have limitations with respect to capturing microbial reads and cannot distinguish if the

source of microbial reads is intracellular or extracellular, or alive or dead when the sample was

taken. Poore et al. suggest, correctly we believe, that additional protocols need to be developed

for prospective studies.

Accepting the limitations of the study, we observed certain trends. Proteobacteria and

Firmicutes were the most frequent phyla identified as predictive features (**Supplementary Data**

2), followed by Actinobacteria and Bacteroidetes. Among viruses, Herpesvirales were the most

frequent. More microbial genera were negatively predictive of drug response or prognosis than

those positively predictive (negative for 306/537 features; two-sided binomial test p-value =

0.0014). Firmicutes reversed this trend, being more often positively predictive (positive for 49/82

features, two-sided Fisher's exact test p-value = 0. 0.0036; **Supplementary Table 2**).

Further examining the predictive features of specific cancer types, we noted that several

genera of Firmicutes were predictive of OS in CESC, including genera of *Lactobacillales*.

Notably, though CESC is known to often arise from HPV infection, the presence of other

microbial species, in particular the Firmicutes genus *Lactobacillus* has been reported in the

literature to be positively associated with the risk of developing CESC[23].

In STAD, tumor microbial abundances were predictive of response to three different drugs:

cisplatin, leucovorin, and oxaliplatin. The genus *Helicobacter* was a quantified microbial

abundance feature in the Poore et al. dataset although notably, even though it is well established

that patients infected with *H. pylori* have an increased risk of developing gastric cancer[24],

Helicobacter was not identified as a predictive feature of drug response in our STAD models.

This findng is in line with recent research indicating reduced microbial diversity, decreased

abundance of *H. pylori*, and enrichment of other mostly commensal bacterial genera in gastric

carcinoma[25]. Instead, in STAD we found that known opportunistic bacteria *Cedecea* and

Sphingobacterium were both strongly negatively predictive of leucovorin response,

Sphingobacterium was strongly negatively predictive of cisplatin response, and the opportunistic

bacteria *Rouxiella* was strongly negative predictive of oxaliplatin response. *Cedecea* and

Sphingobacterium have been implicated in bacteremia in immunocompromised individuals in

rare cases, including cancer[26,27,28,29]. As dysbiosis is frequent in stomach cancer[30,31], and in light

of the mechanism of action of leucovorin, it may be of interest to study whether organisms from

these two genera may sequester or prevent the bacterial production of folinic acid[32].

We found three microbial genera whose abundances were strongly associated with breast

cancer response to doxetaxel. Indeed, the involvement of the microbiome in breast cancer

(BRCA)[22,33] has recently received considerable attention. In BRCA, we found that the genus

containing Epstein-Barr virus (EBV) was negatively associated with response to docetaxel,

which is concordant with previous findings that EBV is associated with chemoresistance to

docetaxel in gastric cancer[34]. Interestingly, Cyanobacteria were predictive features in several

cancers in our study and we identified a genus of Cyanobacteria as predictive of response to

docetaxel in BRCA. Notably, the presence of Cyanobacteria in BRCA was recently confirmed

by Nejman et al.[22] by 16S-rRNA sequencing. While the genus we identified, *Raphidiopsis*, a

planktonic Cyanobacteria that produces toxins harmful to human health and found in freshwater,

is possibly a taxonomic identification error in the original microbial abundance estimates, our

findings may point to a related genus under the recently discovered clade Melainabacteria of

Cyanobacteria[35], which is present in humans. Though Melainabacteria are difficult to culture, we

believe that confirmation of the relationship between BRCA to response to docetaxel and

Melainabacteria should be tested, and a first step would be to confirm our computationally

derived findings in a dedicated 16S-rRNA analysis.

Interestingly, in sarcoma (SARC), among the most predictive microbial features we found

the genus *Lactococcus* to be positively associated with response to docetaxel. *Lactococcus*

contains species that can sometimes cause opportunistic infections in humans, as *Lactococcus*

are similar to *Streptococcus* and formerly belonged to that genus. The result that this genus was

positively associated with response in our model initially appeared counterintuitive, although

while the use of therapeutic bacteria as antitumor agents has not been an extensively studied

field, there have been some limited findings in the literature that suggest the use of

bacteriotherapy as anticancer agents[36]. Historically, the intentional use of the toxins of various

Streptococcus species showing significant antitumor activity in SARC has been

documented[37,38,39]. One possible testable explanation for some microbes being strongly positive

predictive of docetaxel response in our model is that they might produce some extracellular

products or toxins that could work as an adjuvant to the chemotherapy.

In summary, while these findings and others reported in the paper are computationally

derived associations, we believe that they can serve as leads for further experimental studies of

the role of microbial species in modulating patients survival and drug response, potentially by

metabolizing drug levels in the tumor microenvironment as suggested above, or by altering the

immune response, either by changing the levels of specific immunometabolites or by having the

tumors present specific bacterial antigens[40].

Methods
306

Data retrieval and processing

Normalized and batch effect corrected microbial abundance data for 32 TCGA tumor types

were downloaded from the online data repository referenced in Poore et al.[10]

(ftp://ftp.microbio.me/pub/cancer_microbiome_analysis). Specifically, the "Kraken-TCGA-

Voom-SNM-Plate-Center-Filtering-Data.csv" microbial abundance data file and adjoining

"Metadata-TCGA-Kraken-17625-Samples.csv" metadata file were used as the starting input for

further data processing.

We first filtered the data for primary tumor samples (TCGA "Primary Tumor" or "Additional

- New Primary" sample types). Poore et al. generated microbial abundances from all the

available WGS and RNA-seq data in legacy TCGA (after some quality filters), which frequently

contained replicate WGS and RNA-seq data for the same case and sample type. It was common

in legacy TCGA to increase WGS sequencing coverage by performing an additional sequencing

run from the same sample and these secondary runs typically had a much lower number of reads

and coverage compared to their corresponding primary sequencing runs. When comparing the

normalized and batch effect corrected read counts between these WGS runs, we found that

microbial abundance data which came from lower coverage secondary runs could be

substantially different from abundances derived from the larger primary sequencing runs.

Therefore, we excluded microbial abundance data which came from secondary runs. In addition,

legacy TCGA commonly contained data for the same samples analyzed using different

computational pipeline versions. We excluded replicate microbial abundance data from older

TCGA analysis pipeline versions if a replicate from a newer version existed. After the above

filters, the Poore et al. data went from 17,625 samples and 10,183 unique cases to 12,111

samples and 9,812 unique cases (comprising of 1,944 WGS samples from 1,904 unique cases

and 10,167 RNA-seq samples from 9,745 unique cases).

TCGA gender, age at diagnosis, and tumor stage demographic and clinical data and as well

as primary tumor RNA-seq read count data for the 32 TCGA tumor types included in our study

were obtained from the NCI Genomic Data Commons (GDC Data Release v29.0) using the R

Bioconductor package GenomicDataCommons. TCGA GENCODE v22 gene annotations were

obtained from the GDC data portal and Ensembl Gene v98 using the R package rtracklayer and

R Bioconductor packages AnnotationHub and ensembldb. The downloaded GDC primary tumor

cohort with RNA-seq read count data comprised of 9,735 samples from 9,680 unique cases.

There were 68 cases at the GDC which had missing age of diagnosis but existing values in the

Poore et al. data and we chose not to exclude these data and used the Poore et al. age of diagnosis

values for these cases. TCGA curated survival phenotypic data[41] were obtained from UCSC

Xena. Cases which had both missing overall survival (OS) and progressive-free interval (PFI)

outcome data were excluded from survival modeling.

Drug response data were compiled from the TCGA Research Network. Our drug response

models used the following binary classification targets: complete response (CR) and partial

response (PR) were labeled as responders and stable disease (SD) and progressive disease (PD)

as non-responders. All TCGA samples with drug response phenotypic data were from pre-

treatment biopsies. Due to the limited cancer-drug combination cohort sizes in TCGA, we

modeled each drug individually, even if a patient received multiple drugs concurrently. If the

same drug was given at multiple timepoints to a patient, we only considered their first drug

response. We considered cancer-drug combinations that contained a minimum of 18 cases and at

least 4 cases per response binary class, except for STAD oxaliplatin, where we allowed a

minimum of 14 cases so that the gene expression dataset could be included. In total, we analyzed

30 cancer-drug combinations which had paired microbial abundance and gene expression data

that met the above thresholds. Combined feature microbial abundance and gene expression

datasets were created by joining data from each individual dataset which had matching TCGA

sample UUIDs. For some TCGA cases, data existed from multiple different aliquots per sample

or multiple technical runs per aliquot, therefore in these cases all combinations were joined at the

sample UUID level. Cross-validation sampling probability weights as well as model and scoring

sample weights were applied to account and adjust for any imbalance caused by the process.

ML modeling

Machine learning (ML) models were built using the scikit-learn[42] and scikit-survival

libraries[43,44,45]. Custom extensions to scikit-learn and scikit-survival were developed to add new

methods and functionalities required by this project. Survival models were built using Coxnet –

regularized Cox regression with elastic net penalties[11]. Coxnet models controlled for gender, age

at diagnosis, and tumor stage clinical prognostic covariates by including them as unpenalized

features in the model (Coxnet penalty factor = 0). Drug response classification models were built

using three different ML methods: 1) a variant of the linear support vector machine recursive

feature elimination (SVM-RFE) algorithm[15] that we developed with a number of additional

features and better performance than the scikit-learn built-in version, 2) logistic regression

(LGR) with elastic net[16] (L1 + L2) penalties and embedded feature selection, and 3) LGR with

an L2 penalty and limma[17] (for tumor microbial and combination datasets) or edgeR[18,19] (for

RNA-seq count datasets) differential abundance/expression feature scoring inside a k-best

wrapper feature selection method around the learning algorithm. limma differential abundance

analysis was run inside the ML pipeline with default parameters except for fitting an intensity-

dependent trend to the prior variances and running a robust empirical Bayes procedure (eBayes

function parameters trend = TRUE and robust = TRUE). edgeR differential expression analysis

was run inside the ML pipeline with default parameters except for enabling robust estimation of

the negative binomial dispersion (calcDispersions function robust = TRUE) and robust

estimation of the prior quasi-likelihood (QL) dispersion (glmQLFit function robust = TRUE).

Both limma and edgeR methods scored and ranked features by differential abundance/expression

p-value.

All three drug response ML methods unconditionally included the same three clinical

covariates in the model as in the prognosis models by having them bypass feature selection in the

ML pipeline, thought in drug response models clinical covariates where modeled as L2 penalized

features. In SVM-RFE, clinical covariate features bypassed recursive feature elimination but

were always included at each RFE recursive feature elimination model fitting step as well as

final model refitting. To the best of our knowledge, no available comprehensive ML library in

python or R currently provides an elastic net LGR algorithm with the functionality to specify

features that can bypass embedded feature selection and be modeled with an L2 penalty (setting

the R glmnet penalty factor, for example, does not provide this functionality as it isn't a penalty

factor per regularization term but a factor applied to the sum of both L1 and L2 terms). In order

to develop this functionality for our study, our elastic net LGR model pipeline was designed as a

two-level LGR, 1) an elastic net LGR and embedded feature selection on only microbial

abundance or gene expression features with clinical covariates bypassing this step, followed by

2) an L2 penalized LGR on features selected by the elastic net LGR step and the clinical

covariates. We know this design doesn't likely produce the exact same model settings and results

of a single-level elastic net LGR algorithm with the functionality we needed, if such a

implementation it existed, though we tested every drug response model through an ML pipeline

with elastic net LGR and no clinical feature selection bypass and found that model predictive

performance, feature coefficients and signs, and feature importance rankings were similar to our

two-level ML pipeline setup.

Gender was one-hot encoded and tumor stage ordinal encoded by major stage. In the final

cohort included in our prognosis and drug response models, 3363 out of 9708 tumor microbial

abundance cases (34.64%) and 3244 out of 9484 gene expression cases (34.21%) had tumor

stage "not reported" or BRCA stage "X". Since missing tumor stage metadata is so prevalent in

TCGA, we took the approach of including these in our study and modeled missing tumor stage

with as neutral an ordinal encoding as possible. Looking at the distribution of reported major

tumor stages in our cohort, we determined that encoding missing data as an ordinal between

tumor stage II and III was as close to the middle of the distribution of stages in TCGA as we

could possibly achieve with ordinal encoding.

All prognosis and drug response models included the previously described feature selection as well as normalization and transformation steps integrated into the ML modeling pipeline using an extended version of the scikit-learn Pipeline framework. Training data splits always underwent feature selection, normalization, and transformation through the ML pipeline independently from held-out test and validation data splits before learning. Models built using gene expression read count data included edgeR low count filtering, weighted trimmed mean of M-values (TMM) normalization, and log counts per million (CPM) transformation steps within the ML pipeline. These were developed and integrated into our scikit-learn-based framework via R and rpy2. All models also included standardization of features within the ML pipeline just before learning. During prediction, held-out test or validation data were feature selected, normalized and transformed through the ML pipeline using the parameters learned from the training data at each pipeline step before model prediction and scoring. Each cancer, data type, and survival or drug response target type combination was modeled individually using a nested cross-validation (CV) strategy to perform model selection and evaluation on held-out test data. All cross-validation iterators kept replicate sample data per case grouped together such that data would only reside in either the train or test split during each CV iteration.

Survival models used a stratified, randomly shuffled outer CV with 75% train and 25% test split sizes that was repeated 100 times. The CV procedure stratified the splits on event status. Each training set from the outer CV was used to perform hyperparameter tuning and model selection by optimizing C-index over a stratified, randomly shuffled, 4-fold inner CV on the training set, repeated 5 times. A few cancer datasets contained fewer than four uncensored cases which required reducing the number of inner CV folds for these models such that at least one case per fold was uncensored. The data derived from Poore et al. often included more than one sample per case, and an unequal number of samples between cases, therefore requiring either ML

model sample weighting or CV random sampling per case. The Coxnet implementation in scikit-

survival does not currently support sample weights, therefore our custom outer CV iterator

randomly sampled one replicate sample per case during each iteration, using a sampling

procedure with probability weights that balanced the probability that a replicate WGS- or RNA-

seq-based sample was selected during each CV iteration. Model selection grid search was

performed on the following hyperparameters: elastic net penalty L1 ratios 0.1, 0.3, 0.5, 0.7, 0.8,

0.9, 0.95, 0.99, and 1, and for each L1 ratio a default alpha path of 100 alphas using an alpha min

ratio of 10^{-2}. Alpha is the constant multiplier of the penalty terms in the Coxnet objective

function. Optimal alpha and L1 ratio settings were determined via inner CV and a model with

these settings was then refit on the entire outer CV train data split. Model performance was

evaluated in both inner and outer CV on the held-out validation or test data split, respectively, by

generating model test predicted risk scores and using these scores to directly calculate a Harrell's

concordance index (C-index). We also evaluated and compared model predictive performance

for each test data split survival time period by calculating time-dependent cumulative/dynamic

AUCs[12,13].

Drug response models used a stratified, randomly shuffled, 4-fold outer CV that was repeated

25 times (i.e. 100 model instances). Each training set from the outer CV was used to perform

hyperparameter tuning and model selection by optimizing the area under receiver-operator curve

(AUROC) over a stratified, randomly shuffled, 3-fold inner CV repeated 5 times. Case replicate

sample weights were provided to SVM-RFE and LGR learning algorithms and all model

selection and evaluation scoring methods. Class weights were provided to SVM-RFE and LGR

learning algorithms to adjust for any class imbalance. Model selection grid search was performed

on the following hyperparameters: SVM and L2 penalized LGR C regularization parameter from

a range of 10^{-5} to 10^3, elastic net LGR L1 ratios of 0.1, 0.3, 0.5, 0.7, 0.8, 0.9, 0.95, 0.99 and 1,

elastic net LGR C regularization parameter from a range of 10^{-2} to 10^3 (microbial abundance) or

from 10^{-2} to 10^1 (gene expression), and finally RFE, elastic net LGR, and limma and edgeR

feature scorer k-best feature selection search range from 1 to 400 top scoring microbial

abundance or gene expression features. Optimized hyperparameter settings were determined via

inner CV and a model with the optimized settings was then refit on the entire outer CV train data

split. Model performance was evaluated in both inner and outer CV on the held-out validation or

test data split, respectively, by AUROC, area under precision-recall curve (AUPRC), and

balanced accuracy (BCR). AUROC was used to evaluate and select the best model and

optimized hyperparameter settings from the grid search. Average precision and BCR scores were

also reported for each model instance.

Gender, age at diagnosis, and tumor stage clinical covariate-only survival models were built

using standard unpenalized Cox regression. Clinical covariate-only drug response models were

built using linear SVM or LGR. Models included standardization of features as part of the ML

pipeline. Models were trained and tested using the same outer CV iterators and train/test data

splits as their corresponding microbial abundance, gene expression, or combination data type

models. To test whether a Coxnet, SVM-RFE, or LGR microbial abundance or gene expression

model was significantly better than their corresponding Cox, linear SVM, or LGR clinical

covariate-only model, respectively, a two-sided Wilcoxon signed-rank test was performed

between the 100 pairs of C-index or AUROC scores between both models. All raw p-values

generated from the signed-rank test across survival or drug response analyses from the same data

type were adjusted for multiple testing using the Benjamini-Hochberg (BH) procedure to control

the false discovery rate (FDR), and a threshold FDR ≤ 0.01 was used to determine statistical

significance. To test whether a combined data type model was significantly better than its

corresponding microbial abundance or gene expression model, a two-sided Dunn test was

performed between all three groups of data type model scores. Each Dunn test raw p-value was

adjusted for multiple testing using the Benjamini-Hochberg (BH) procedure to control the false

discovery rate (FDR), and a threshold FDR ≤ 0.05 was used to determine statistical significance.

Microbial abundance model feature analysis

For each analysis, 100 prognosis or drug response model instances were generated from the

outer CV procedure. Each model instance selected a subset of features that performed best

during CV and the model algorithm learned coefficients (or weights) for each feature. To select

microbial genera for downstream investigation from the feature results across all these model

instances, we proceeded as follows. First, we applied a two-sided Wilcoxon signed-rank test that

the mean feature coefficient rank generated by the model is shifted away from zero, and thus that

the genus is identifiably positively or negatively associated with survival or drug response. For

all Wilcoxon tests, we used the package coin[46], which allows exact calculation of p-values.

Coefficients were ignored when a genus was assigned a zero coefficient or absent from a model.

Second, within each model, all coefficients, ignoring the results of the Wilcoxon test, were

ranked by absolute magnitude. We then kept genera that were among the top 50 features in at

least 20% of the models and for which the Holm-adjusted, two-sided Wilcoxon signed-rank test

p-value was ≤ 0.01. Having a Coxnet feature coefficient equal to zero or feature being absent

from an SVM-RFE or LGR model was not strong enough evidence that the genus has no effect,

but rather that one or more features with stronger effect were chosen. Thus, we ignored genera

with a zero coefficient or absent from a model when computing mean coefficient weight and

Wilcoxon statistics on the means.

For the drug response models, where three ML methods were tested, we noted the features

selected by individual models and the median rank the feature attained in the instances in which

it appeared, but further filtered the features to account for the consensus between ML models.

We kept features selected in any two ML model methods that individually met our criteria for

inclusion, ignoring features in ML models that did not meet these criteria. We then computed the

Spearman correlation between the median ranks attained by the features.

For each selected microbial feature, we tested whether it was a significantly univariate

feature of survival or drug response. This is a strictly different question than whether the

coefficient of a feature has consistent sign – sign may be consistent when used in combination

with other features, but the feature may not be individually predictive. For drug response models,

we divided individuals into responders and non-responders, and for survival data we divided

individuals whose survival time was greater or less than the censored median, ignoring those

who were lost to follow up before median time. For each cancer-test type pair, we applied a two-

sided Wilcoxon rank-sum test. We applied a Benjamini-Hochberg multiple hypothesis correction

for each cancer-test type pair and report the false discovery rate in **Supplementary Data 2**.

We analyzed the distribution of features, selected by the rules described above, that had

positive or negative signs for their mean coefficient. We used a two-sided binomial test to show

that selected features had significantly more negative the positive mean coefficients. We used a

two-sided Fisher's exact test to determine if selected genera belonging to Firmicutes had a

statistically significant difference in the breakdown between positive and negative mean

coefficients than selected features as a whole.

Data availability

526   All results generated from this work are available under https://doi.org/10.5281/zenodo.5221525.

Code availability

528 All code and data used to produce this work are available under

529 https://github.com/ruppinlab/tcga-microbiome-prediction.

References

1. Grossman, R. L. et al. Toward a Shared Vision for Cancer Genomic Data. *New England*
Journal of Medicine **375**, 1109–1112 (2016).

2. Ahluwalia, P., Kolhe, R. & Gahlay, G. K. The clinical relevance of gene expression based
prognostic signatures in colorectal cancer. *Biochimica et Biophysica Acta (BBA) - Reviews*
on Cancer **1875**, 188513 (2021).

3. Brodsky, A. S. *et al.* Expression profiling of primary and metastatic ovarian tumors reveals
differences indicative of aggressive disease. *PLoS One* **9**, e94476 (2014).

4. Liu, Y. *et al.* Pan-cancer analysis of clinical significance and associated molecular features of
glycolysis. *Bioengineered* **12**, 4233–4246 (2021).

5. Selfors, L. M., Stover, D. G., Harris, I. S., Brugge, J. S. & Coloff, J. L. Identification of
cancer genes that are independent of dominant proliferation and lineage programs. *Proc Natl*
Acad Sci U S A **114**, E11276–E11284 (2017).

6. Shimoni, Y. Association between expression of random gene sets and survival is evident in
multiple cancer types and may be explained by sub-classification. *PLoS Comput Biol* **14**,
e1006026 (2018).

7. Shukla, S. *et al.* Development of a RNA-Seq Based Prognostic Signature in Lung
Adenocarcinoma. *J Natl Cancer Inst* **109**, (2017).

8. Venet, D., Dumont, J. E. & Detours, V. Most random gene expression signatures are
significantly associated with breast cancer outcome. *PLoS Comput Biol* **7**, e1002240 (2011).

9. Milanez-Almeida, P., Martins, A. J., Germain, R. N. & Tsang, J. S. Cancer prognosis with
shallow tumor RNA sequencing. *Nature Medicine* **26**, 188–192 (2020).

10. Poore, G. D. *et al.* Microbiome analyses of blood and tissues suggest cancer diagnostic
approach. *Nature* **579**, 567–574 (2020).

11. Simon, N., Friedman, J., Hastie, T., & Tibshirani, R. Regularization Paths for Cox's
Proportional Hazards Model via Coordinate Descent. *Journal of Statistical Software* **39** (5),
1-13 (2011).

12. Hung, H. & Chiang, C.T. Estimation methods for time-dependent AUC models with survival
data. *Canadian Journal of Statistics* **38** (1), 8–26 (2010).

13. Lambert, J. & Chevret, S. Summary measure of discrimination in survival models based on
cumulative/dynamic time-dependent ROC curves. *Statistical methods in medical research* **25**
(5), 2088–2102 (2016).

14. Gnanasekar A. The intratumor microbiome predicts prognosis across gender and subtypes in
papillary thyroid carcinoma. *Comput Struct Biotechnol J* **19**, 1986-1997 (2021).

15. Guyon, I., Weston, J., Barnhill, S. & Vapnik, V. Gene Selection for Cancer Classification
using Support Vector Machines. *Machine Learning* **46**, 389–422 (2002).

16. Zou H. & Hastie T. Regularization and variable selection via the elastic net. *J R Statist Soc B*
67 (2), 301–320 (2005).

17. Ritchie, M.E. *et al. limma* powers differential expression analyses for RNA-sequencing and
microarray studies. *Nucleic Acids Res* **43** (7), e47 (2015).

18. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140
(2010).

19. McCarthy, D. J., Chen, Y. & Smyth, G. K. Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res* **40**, 4288–4297
(2012).

20. Geller, L. T. *et al.* Potential role of intratumor bacteria in mediating tumor resistance to the
chemotherapeutic drug gemcitabine. *Science* **357**, 1156–1160 (2017).

21. Pushalkar, S. *et al.* The Pancreatic Cancer Microbiome Promotes Oncogenesis by Induction
of Innate and Adaptive Immune Suppression. *Cancer Discov* **8**, 403–416 (2018).

22. Nejman, D. *et al.* The human tumor microbiome is composed of tumor type–specific
intracellular bacteria. *Science* **368**, 973–980 (2020).

23. Lin, D. *et al.* Microbiome factors in HPV-driven carcinogenesis and cancers. *PLoS Pathog*
16, (2020).

24. Parsonnet, J. et al. Helicobacter pylori infection and the risk of gastric carcinoma. *N Engl J*
Med **325**, 1127–1131 (1991).

25. Ferreira R.M. et al. Gastric microbial community profiling reveals a dysbiotic cancer-
associated microbiota. *Gut* **67** (2): 226-236 (2018).

26. Abate, G., Qureshi, S. & Mazumder, S. A. Cedecea davisae bacteremia in a neutropenic
patient with acute myeloid leukemia. *J Infect* **63**, 83–85 (2011).

27. Akinosoglou, K. *et al.* Bacteraemia due to Cedecea davisae in a patient with sigmoid colon
cancer: a case report and brief review of the literature. *Diagn Microbiol Infect Dis* **74**, 303–
306 (2012).

28. Koh, Y. R. *et al.* The first Korean case of Sphingobacterium spiritivorum bacteremia in a
patient with acute myeloid leukemia. *Ann Lab Med* **33**, 283–287 (2013).

29. Wu, P. *et al.* Profiling the Urinary Microbiota in Male Patients with Bladder Cancer in
China. *Front Cell Infect Microbiol* **8**, 167 (2018).

30. Coker, O. O. *et al.* Mucosal microbiome dysbiosis in gastric carcinogenesis. *Gut* **67**, 1024–
1032 (2018).

31. Castaño-Rodriguez N. *et al*. Dysbiosis of the microbiome in gastric carcinogenesis. *Sci Rep* **7**
(1), 15957 (2015).

32. Ogwang, S. et al. Bacterial Conversion of Folinic Acid Is Required for Antifolate Resistance.
Journal of Biological Chemistry **286**, 15377–15390 (2011).

33. Eslami-S, Z., Majidzadeh-A, K., Halvaei, S., Babapirali, F. & Esmaeili, R. Microbiome and
Breast Cancer: New Role for an Ancient Population. *Front Oncol* **10**, (2020).

34. Shin, H. J., Kim, D. N. & Lee, S. K. Association between Epstein-Barr virus infection and
chemoresistance to docetaxel in gastric carcinoma. *Mol. Cells* **32**, 173–179 (2011).

35. Di Rienzi, S. C. *et al.* The human gut and groundwater harbor non-photosynthetic bacteria
belonging to a new candidate phylum sibling to Cyanobacteria. *eLife* **2**, e01102 (2013).

36. Sedighi M. *et al*. Therapeutic bacteria to combat cancer; current advances, challenges, and
opportunities. *Cancer Med* **8** (6), 3167-3181 (2019).

37. Coley WB. Late results of the treatment of inoperable sarcoma by the mixed toxins of
erysipelas and bacillus prodigiosus. *Trans Southern Surg Gynecol Ass* **18**, 197 (1906).

38. Fehleisen F. Ueber die Züchtung der Erysipelkokken auf künstlichem Nährboden und ihre
Übertragbarkeit auf den Menschen. *Dtsch Med Wochenschr* **8**, 553-554 (1882).

39. Busch W. Aus der Sitzung der medicinischen Section vom 13 November 1867. *Berl Klin*
Wochenschr **5**, 137 (1868).

40. Kalaora S. *et al*. Identification of bacteria-derived HLA-bound peptides in melanoma. *Nature*
592 (7852), 138-143 (2021).

41. Liu, J. *et al.* An Integrated TCGA Pan-Cancer Clinical Data Resource to Drive High-Quality
Survival Outcome Analytics. *Cell* **173**, 400-416.e11 (2018).

42. Pedregosa *et al*. Scikit-learn: Machine Learning in Python. *J Mach Learn Res* **12**, 2825-2830
(2011).

43. Pölsterl, S., Navab, N. & Katouzian, A. Fast Training of Support Vector Machines for
Survival Analysis. in *Machine Learning and Knowledge Discovery in Databases* (eds.
Appice, A. et al.) 243–259 (Springer International Publishing, 2015).

44. Pölsterl, S., Navab, N., & Katouzian, A., An Efficient Training Algorithm for Kernel
Survival Support Vector Machines. *4th Workshop on Machine Learning in Life Sciences*, 23
September 2016, Riva del Garda, Italy.

45. Pölsterl, S. *et al.* Heterogeneous ensembles for predicting survival of metastatic, castrate-
resistant prostate cancer patients. *F1000Res* **5**, 2676 (2017).

46. Hothorn, T., Hornik, K., Wiel, M. A. van de & Zeileis, A. Implementing a Class of
Permutation Tests: The coin Package. *Journal of Statistical Software* **28**, 1–23 (2008).

Acknowledgements

The results shown here are in whole or part based upon data generated by the TCGA Research

Network (https://www.cancer.gov/tcga). This research was supported by the Intramural Research

Program of the National Institutes of Health, National Cancer Institute and by NIH grant

1ZIABC011803-03. The authors would like to personally thank Christopher Buck from the NCI,

Pedro Milanez-Almeida and John Tsang from NIH NIAID, and Alejandro Schäffer, Welles

Robinson, Fiorella Schischlik, Sanju Sinha, and Sanna Madan from NCI CDSL for their

assistance in this project. This work utilized the computational resources of the NIH HPC

Biowulf cluster (http://hpc.nih.gov). The authors would like to thank Richard Lehr, Tim Miller,

Wolfgand Resch, and Steve Fellini for their assistance in running this analysis on the NIH HPC

Biowulf cluster. The authors would also like to thank Joel Nothman, Andreas Mueller, and Adrin

Jalali from the scikit-learn core development team and scikit-survival author Sebastian Pölsterl

for their assistance with developing extensions to their libraries.

Author information

Affiliations

Cancer Data Science Laboratory (CDSL), National Cancer Institute (NCI), National
Institutes of Health (NIH), Bethesda, MD, USA

Leandro Cruz Hermida, E. Michael Gertz & Eytan Ruppin

Department of Computer Science, University of Maryland, College Park, MD, USA

Leandro Cruz Hermida

Contributions

L.C.H., E.M.G., and E.R. designed the study. L.C.H. and E.M.G. performed all computational

analyses and results interpretation. L.C.H., E.M.G., and E.R. wrote the paper.

Corresponding authors

Correspondence to Eytan Ruppin.

659 Ethics declaration

660 **Competing interests**

661 All authors declare that they have no competing interests.

Supplementary Tables

Supplementary Table 1. Prognosis model results comparison to Milanez-Almeida et al.

| | Milanez-Almeida | | Hermida & Gertz | | | | | |
Cancer	Tested	Hit	Tested	Hit	Common Hit	New Hit	M-A Tested New Hit	Missing M-A Hit
ACC	OS	OS	OS, PFI	OS, PFI	OS	PFI		
BLCA	OS	OS	OS, PFI	OS, PFI	OS	PFI		
BRCA	PFI		OS, PFI	PFI		PFI	PFI	
CESC	OS		OS, PFI	OS, PFI		OS, PFI	OS	
CHOL	*EXCL*		OS, PFI					
COAD	OS		OS, PFI					
DLBC	*EXCL*		OS, PFI					
ESCA	OS		OS, PFI					
GBM	OS		OS, PFI					
HNSC	OS	OS	OS, PFI	OS	OS			
KICH	*EXCL*		OS, PFI					
KIRC	OS	OS	OS, PFI	OS, PFI	OS	PFI		
KIRP	OS	OS	OS, PFI	OS, PFI	OS	PFI		
LAML	OS	OS	*EXCL*					OS
LGG	PFI	PFI	OS, PFI	OS, PFI	PFI	OS		
LIHC	OS	OS	OS, PFI	OS, PFI	OS	PFI		
LUAD	OS	OS	OS, PFI	OS	OS			
LUSC	OS		OS, PFI	PFI		PFI		
MESO	OS	OS	OS, PFI	OS, PFI	OS	PFI		
OV	OS		OS, PFI					
PAAD	OS	OS	OS, PFI	PFI		PFI		OS
PCPG	*EXCL*		OS, PFI	OS		OS		
PRAD	PFI	PFI	OS, PFI	PFI	PFI			
READ	PFI		OS, PFI					
SARC	OS		OS, PFI	OS, PFI		OS, PFI	OS	
SKCM	*EXCL*		OS, PFI	OS, PFI		OS, PFI		

STAD	OS		OS, PFI	PFI		PFI		
TGCT	OS		OS, PFI					
THCA	PFI		OS, PFI					
THYM	PFI		OS, PFI	OS		OS		
UCEC	OS		OS, PFI	OS, PFI		OS, PFI	OS	
UCS	OS		OS, PFI					
UVM	OS	OS	OS, PFI	OS, PFI	OS	PFI		

Supplementary Table 2. Per cancer, the number of times genera from the phylum Firmicutes were found among the selected features, whether positively or negatively associated with drug response or prognosis.

Cancer	Selected Negatively	Selected Positively	Total
ACC	8	13	21
BRCA	1	0	1
CESC	4	7	11
ESCA	0	4	4
KIRC	4	7	11
LGG	1	6	7
SARC	2	2	4
STAD	13	10	23

669   Supplementary Figures

Supplementary Figure 1. Performance of gene expression overall survival (OS) models in the 16 tumor types where gene expression adds predictive power to clinical covariates. (a) C-index score density distributions (top rows) for gene expression with clinical covariate models vs clinical covariate-only models. Lines connecting points (light grey) represent score pairs from same train-test split on the data. Mean C-index scores and connecting lines shown in red. Significance for the prediction improvement over clinical covariate-only models was calculated using a two-sided Wilcoxon signed-rank test and adjusted for multiple testing using the Benjamini-Hochberg method with adjusted p-values shown at top. **(b)** Time-dependent, cumulative/dynamic AUCs (bottom rows) for gene expression with clinical covariate models (orange) vs clinical covariate-only models (grey) following years after diagnosis. Mean AUCs across entire test time range after diagnosis shown as a horizontal dotted line and in legends and shaded areas denote standard deviations.

Supplementary Figure 2. Performance of gene expression progression-free interval (PFI) models in the 17 tumor types where gene expression adds predictive power to clinical covariates. (a) C-index score density distributions (top rows) for gene expression with clinical covariate models vs clinical covariate-only models. Lines connecting points (light grey) represent score pairs from same train-test split on the data. Mean C-index scores and connecting lines shown in red. Significance for the prediction improvement over clinical covariate-only models was calculated using a two-sided Wilcoxon signed-rank test and adjusted for multiple testing using the Benjamini-Hochberg method with adjusted p-values shown at top. **(b)** Time-dependent, cumulative/dynamic AUCs (bottom rows) for gene expression with clinical covariate models (orange) vs clinical covariate-only models (grey) following years after diagnosis. Mean AUCs across entire test time range shown as a horizontal dotted line and in legends and shaded areas denote standard deviations.

Supplementary Figure 3. Performance of microbial abundance prognosis models in the five tumor types
where microbial abundance features add predictive power to clinical covariates. (a) C-index score density
distributions (top row) for microbial abundance with clinical covariate models vs clinical covariate-only models.
Lines connecting points (light grey) represent score pairs from same train-test split on the data. Mean C-index scores
and connecting lines shown in red. **(b)** Time-dependent, cumulative/dynamic AUCs (bottom row) for microbial
abundance with clinical covariate models (blue) vs clinical covariate-only models (grey) following years after
diagnosis. Mean AUCs across entire test time range shown as a horizontal dotted line and in legends and shaded
areas denote standard deviations. Significance was calculated using a two-sided Wilcoxon signed-rank test and
adjusted for multiple testing using the Benjamini-Hochberg method with adjusted p-values shown at top of violin
plots in **(a)**.

Supplementary Figure 4. Performance of combined microbial abundance and gene expression models where combining both data types adds to predictive power. **(a)** Prognosis model C-index score violin density distributions (left) between microbial abundance, gene expression, and combined data type models. Mean scores shown in red. Significance was calculated using a two-sided Dunn test and p-values were adjusted for multiple testing using the Benjamini-Hochberg method. Model C-index score violin density distributions (middle) for combined data type with clinical covariate models vs clinical covariate-only models. Time-dependent, cumulative/dynamic AUCs (right) for combined data type (purple), microbial abundance (blue), and gene expression (orange) models following years after diagnosis. **(b)** ML method drug response model mean AUROC scores where combining data types performed better than clinical covariates alone. For the combined data type drug response model closest to reach significant improvement over respective single data type models **(c)** AUROC score violin density distributions (left) comparing each data type model, AUROC score violin density distributions (middle left) comparing combined data type and clinical covariate model to clinical covariate-only model, and mean ROC and PR curves (right) for each data type model. Mean AUROC and AUPRC scores shown in panel legends and shaded areas denote standard deviations.

Description of Supplementary Data Files

Supplementary Data 1.xlsx

Worksheet #1 Prognosis model cohort sizes: number of cases in each prognosis model dataset.

Worksheet #1 Drug response model cohort sizes: number of cases in each drug response

model dataset, including the breakdown between responder (R) and non-responder (NR) cases.

Supplementary Data 2.xlsx

Worksheet #1 Microbial genera features: by cancer and by comparator, the genera selected as

features by the criteria outline in Methods, and the ML methods that selected each genus.

Further we show the number of times each genus was seen with rank at most 50 over the number

of times the genus could be seen: 100, 200 or 300 if one, two, or three ML methods chose the

feature, respectively. The column conditional direction shows the direction assigned to the

feature for each ML method; an asterisk indicates that there was more than one relevant ML

method, but all methods chose the same direction. The conditional direction is the typical sign of

the coefficient within a model, which is conditional on other features in the model. Also shown

is the Univariate FDR, which is the FDR that the feature is univariately predictive (n.s. is not

significant), and the direction in which the feature is univariately predicative, which may differ

from the conditional direction.

Worksheet #2 Gene expression features: by cancer and by comparator, the genes selected as

features by the criteria outline in Methods, and the ML methods that selected each gene. Further

we show the number of times each gene was seen with rank at most 50 over the number of times

the gene could be seen: 100, 200 or 300 if one, two, or three ML methods chose the feature,

respectively. The column conditional direction shows the direction assigned to the feature for

739   each ML method; an asterisk indicates that there was more than one relevant ML method, but all

740   methods chose the same direction. The conditional direction is the typical sign of the coefficient

741   within a model.