Oestrogen promotes innate immune evasion of *Candida albicans*

through inactivation of the alternative complement system

Pizga Kumwenda[1], Fabien Cottier[1], Ben Keevan[1], Hannah Gallagher[1] and Rebecca A. Hall*[1,2]

[1] Institute of Microbiology and Infection, School of Biosciences, University to Birmingham,

Birmingham, UK, B15 2TT

[2] Kent Fungal Group, School of Biosciences, University of Kent, Canterbury, UK, CT2 7NJ

r.a.hall@kent.ac.uk

Abstract

Gender is a risk factor for several infections that, for many pathogens, has been linked to sex hormones impacting host immunity and directly affecting microbial virulence. *Candida albicans* is a commensal of the urogenital tract and the predominant cause of vulvovaginal candidiasis (VVC). Factors that increase circulatory oestrogen levels like pregnancy, the use of oral contraceptives, and hormone replacement therapy predispose women to VVC, but the reasons for this are largely unknown. Here, we investigate how adaptation of *C. albicans* to oestrogen impacts the fungal host-pathogen. Physiologically relevant concentrations of oestrogen promoted fungal virulence by enabling *C. albicans* to avoid the actions of the innate immune system. Oestrogen-induced innate immune evasion was mediated via inhibition of opsonophagocytosis through enhanced acquisition of Factor H on the fungal cell surface and was dependent on the *C. albicans* moonlighting protein Gpd2. Oestrogen dependent derepression of *GPD2* was mediated via a non-canonical signalling pathway involving Ebp1 and Bcr1. Therefore, we propose that, in addition to affecting the antifungal potential of vaginal epithelial cells, elevated oestrogen levels predispose women to VVC by directly enhancing fungal pathogenicity. The discovery of this new hormone sensing pathway might pave the way in explaining gender biases associated with fungal infections and may provide an alternative approach to improving women's health.

Introduction

Microbial infections exhibiting gender bias are common. This bias may predispose one sex to infection over the other, or result in one sex exhibiting severer infection outcomes [1-3]. Sex hormones like oestrogen, testosterone and progesterone regulate many functions of the immune system, and generally males are more prone to infection than females, as overall immune responses are lower in the male population [4-6]. Besides the impact of sex hormones on immune cell function, sex hormones also have a direct effect on microbial pathogenicity increasing microbial persistence, metabolism and virulence gene expression [7, 8].

The opportunistic fungal pathogen *Candida albicans* is the predominant fungal coloniser of the female reproductive tract and the major cause of genital thrush (vulvovaginal candidiasis, VVC). Approximately 75% of the female population will encounter at least one episode of VVC in their life-time, while up to 15% experience recurrent infection (RVVC) defined as four or more episodes in a twelve-month period [9]. Although not life threatening, mucosal infections are expensive to treat, impact quality of life, and increase population morbidity.

One of the major risk factors associated with the development of VVC is elevated levels of oestrogen which occur as a result of pregnancy, the use of high oestrogen-containing oral contraceptives and hormone replacement therapy [10, 11]. Therefore, oestrogen plays a key role in predisposing women to VVC, but the precise mechanism(s) for this are unknown. Oestrogen promotes glycogen production at the vaginal mucosa which might provide a nutrient rich environment to promote the expansion of *C. albicans* [10]. In mouse models of VVC, were pseudoestus is induced to maintain fungal colonisation, vaginal epithelial cells have a diminished ability to control the growth of *C. albicans* [12]. In addition, oestrogen decreases the infiltration of phagocytes and suppresses cell mediated immunity [13]. However, in rats, preincubation of *C. albicans* with oestrogen prior to vaginal infection enhances fungal survival [14], suggesting that in addition to affecting host immunity, oestrogen

may directly affect the virulence of *C. albicans*. In agreement with this idea, oestrogen has

been shown to promote hyphal morphogenesis of *C. albicans* [15], a key virulence factor of

the fungus. Furthermore, an oestrogen binding protein (Ebp1) has been identified in *C.*

albicans [16], although the importance of this protein in VVC is not known. Therefore, how *C.*

albicans adapts to oestrogen is still unclear.

The *C. albicans* cell wall is a multi-layered structure consisting of an inner layer of chitin and

beta-glucan, and an outer layer of heavily glycosylated proteins [17]. The fungal cell wall is a

highly dynamic structure providing rigidity, strength and protection from the environment. In

addition, many components of the cell wall act as pathogen associated molecular patterns

(PAMPs) and are recognised by the innate immune system [17, 18]. Recently, remodelling of

the cell wall in response to adaptation to host environments has been shown to regulate the

host-pathogen interaction [19-30]. In addition to direct recognition of cell wall PAMPs

mediating phagocytosis, *C. albicans* also activates the alternative complement system,

resulting in the deposition of complement proteins (i.e. C3) on the fungal cell surface, resulting

in opsonophagocytosis [31]. However, *C. albicans* can evade opsonophagocytosis through

the binding of complement regulatory proteins to its cell surface that inactivate the complement

cascade [32]. Here we show that *C. albicans* does adapt to oestrogen, and that this adaptation

perturbs the host-pathogen interaction, inhibiting phagocytosis of the fungal pathogen.

Avoidance of the innate immune system was mediated via the fungal cell surface protein,

Gpd2, recruiting the human complement regulatory protein, Factor H, to the fungal cell surface

inactivating the alternative complement system.

Results

Adaptation of *C. albicans* to oestrogen promotes immune evasion

Pregnant women and women taking high oestrogen-containing oral contraceptives are more

prone to vulvovaginal candidiasis (VVC) [33], while VVC occurs less frequently in

postmenopausal women, indicating that oestrogen may play a role in promoting the virulence

of *C. albicans*. There are four main forms of oestrogen, estrone (E1), 17β-estradiol (E2), estriol

(E3) and 17α-ethynylestradiol (EE2). Estrone is the weakest form of oestrogen produced by

the ovaries and adipose tissue, and is only found in menopausal women, while 17β-estradiol

is the strongest form of oestrogen produced by the ovaries and has been associated with

many gynaecological disorders. Estriol is a by-product from the metabolism of estradiol, and

as such is found in high concentrations during pregnancy. Finally, 17α-ethynylestradiol is a

synthetic oestrogen used in oral contraceptive pills. To ascertain whether adaptation of *C.*

albicans to oestrogen affects the host-pathogen interaction, *C. albicans* cells were grown in

the presence of physiological (0.0001 μM) and super-physiological (0.1 μM, 10 μM)

concentrations of 17β-estradiol, estriol, or 17α-ethynylestradiol, and phagocytosis rates

quantified. Physiological and super-physiological concentrations of all three forms of

oestrogen significantly inhibited macrophage and neutrophil phagocytosis of *C. albicans,*

resulting in a 50% drop in phagocytosis compared to the ethanol vehicle control (Fig 1A-C, S1

Fig A, B). The reduced rates of *C. albicans* phagocytosis were independent of any impact of

oestrogen on fungal growth (S1 Fig C) or morphology (S1 Fig D). Given that all tested forms

of oestrogen elicited similar results, all subsequent experiments were performed only with

17β-estradiol.

To determine whether the observed inhibition of phagocytosis is truly mediated by adaptation

of *C. albicans* to oestrogen, and not a result of residual oestrogen coating the surface of *C.*

albicans inhibiting phagocyte function, macrophages were pre-treated with oestrogen prior to

the addition of *C. albicans* or latex beads. Macrophages pre-incubated with oestrogen

phagocytosed *C. albicans* at rates comparable to non-treated cells (S1 Fig E), suggesting that

under the tested conditions oestrogen does not directly impact the ability of macrophages to

phagocytose target particles. Pre-incubating inert particles like latex beads with oestrogen had

no effect on the ability of macrophages to phagocytose the particles (S1 Fig F), suggesting

that any residual oestrogen on the surface of particles does not interfere with the ability of

macrophages to phagocytose them. In agreement with this, incubation of *C. albicans* with

oestrogen on ice, which inhibits fungal growth and metabolism, did not affect phagocytosis

rates (S1 Fig G). Furthermore, re-inoculation of oestrogen-adapted *C. albicans* cells into fresh

YPD media restored *C. albicans* phagocytosis rates (S1 Fig H). Therefore, in response to

oestrogen, *C. albicans* undergoes some form of adaptation that alters the host-pathogen

interaction.

Finally, we tested several vaginal *C. albicans* isolates and observed the all isolates displayed

reduced phagocytosis after adaptation to oestrogen, confirming that oestrogen-induced

immune evasion is a general trait of clinically relevant *C. albicans* (Fig 1D). Taking this data

together we conclude that the reduction in *C. albicans* phagocytosis is due to the fungus

reversibly adapting to the oestrogen, and that this adaptation interferes with the host

mechanism of fungal clearance.

Fig 1. Oestrogen promotes innate immune evasion of *C. albicans* *C. albicans* cells were grown in YPD with or without oestrogen supplementation. Cells were harvested, washed in PBS and co-incubated with (**A**) J774A.1 cells (**B**) human macrophages (**C**), human neutrophils. (**D**) J774A.1 phagocytosis rates of *C. albicans* clinical isolates after exposure to 10 μM 17β-estradiol. (**E**) J774A.1 phagocytosis rates of *C. albicans* clinical isolates after exposure to progesterone (**F**) J774A.1 phagocytosis rates of *C. albicans* clinical isolates after exposure to progesterone and oestrogen. Cells were fixed with 4% PFA, imaged by microscopy, scored using ImageJ software and phagocytic index determined. All data represent the mean ± SEM from at least three independent experiments. (*$p < 0.05$, **$p < 0.01$, ****$p < 0.0001$).

Hormone induced innate immune evasion is specific to Oestrogen

Progesterone been shown to affect phagocytosis rates [34], and is structurally quite similar to oestrogen. Therefore, we hypothesised that *C. albicans* may adapt to progesterone in a similar way to oestrogen to evade the innate immune system. *C. albicans* cells pre-exposed to physiological and super-physiological concentrations of progesterone exhibited comparable phagocytosis rates as *C. albicans* cells grown in YPD (Fig 1E), while *C. albicans* treated with both oestrogen and progesterone still exhibited innate immune evasion (Fig 1F). Therefore, the promotion of innate immune evasion appears to be a specific and dominant attribute of oestrogen.

Altered phagocytosis rates does not affect pro-inflammatory cytokine production.

Previous studies have shown that oestrogen can interfere with NF-κB signalling to attenuate inflammatory responses [35, 36]. Therefore, we investigated whether adaptation of *C. albicans* to oestrogen dampens the pro-inflammatory innate immune response. Human peripheral blood monocytes (PBMCs) were exposed to oestrogen-adapted *C. albicans* cells and secretion of TNFα, IFN-γ, IL-6 and IL-1β was evaluated. Adaptation to oestrogen did not affect the cytokine response (Fig 2). Therefore, although significantly fewer oestrogen-adapted cells

are phagocytosed by the innate immune system, the extracellular *C. albicans* cells still

stimulate significant cytokine production, likely through non-phagocytic receptors like TLR4.

Fig 2. Adaptation of *C. albicans* to oestrogen does not affect cytokine secretion. *C.*

albicans cells were grown in YPD with or without 10 μM 17β-estradiol for 4 h. Cells were

washed, fixed with 4% PFA and co-incubated with PBMCs for 24h. Secretion of (**A**) TNFα, (**B**)

IFNγ (**C**) IL-1β and (**D**) IL-16 by PBMCs was quantified by ELISA. Data represent the mean ±

SEM from four independent experiments using different donors.

Oestrogen dependent innate immune evasion is not dependent on fungal carbohydrate cell wall remodelling.

Recently, modulation of the *C. albicans* cell wall carbohydrates has been shown to influence the host-pathogen interaction [19-30]. Therefore, the ability of oestrogen to promote remodelling of the fungal cell wall was quantified by flow cytometry. However, initial studies confirmed that adaptation of *C. albicans* to oestrogen does not result in increased incorporation of mannan, β-glucan or chitin into the cell wall (Fig 3A-C).

Dectin-1 is a phagocytic receptor important for controlling fungal infections through the recognition of surface exposed beta-glucan [37]. Host environmental cues that reduce *C. albicans* phagocytosis do so through masking surface exposed beta-glucan inhibiting Dectin-1 dependent recognition [21-23, 30]. To ascertain whether the observed reduction in phagocytosis was dependent on Dectin-1, the attachment of oestrogen-adapted *C. albicans* cells to Dectin-1 expressing fibroblasts was quantified. Dectin-1 recognised and bound oestrogen-adapted cells at rates comparable to control cells, suggesting that the altered phagocytosis rates are not due to reduced Dectin-1 recognition (Fig 3D). In agreement with this, adaptation of *C. albicans* to oestrogen did not affect the surface exposure of either β-glucan or chitin (Fig 3E, F). Taken together, these data indicate that oestrogen does not affect the overall structure of the *C. albicans* cell wall, and that the oestrogen-dependent reduction in phagocytosis is not dependent on fungal carbohydrate cell wall remodelling.

Fig 3. Reduced phagocytosis rates are not due to gross changes in cell wall

carbohydrates. *C. albicans* cells were grown in YPD with or without 10 μM 17β-estradiol.

Cells were harvested, washed in PBS, fixed with 4% PFA and stained for (**A**) total mannan

(**B**) total glucan (**C**) total chitin. (**D**) To assess the role of Dectin-1 in *C. albicans* phagocytosis,

Dectin-1 was expressed in NIH 3T3 fibroblasts. *C. albicans* cells were stained for (**E**) exposed

β1,3-glucan and (**F**) exposed chitin levels. Staining was quantified by flow cytometry and

median fluorescence intensities (MFI) determined. All data represent the mean ± SEM from

at least three independent experiments.

Oestrogen-induced innate immune evasion is mediated via inhibition of

opsonophagocytosis.

In addition to direct detection of cell wall PAMPs leading to phagocytosis, *C. albicans* can be

opsonised. *C. albicans* activates the alternative complement cascade resulting in the

deposition of complement (C3) on its surface inducing opsonophagocytosis via the

complement receptors CR1 and CR3. To confirm whether oestrogen-induce innate immune

evasion was mediated via inhibition of complement, we assessed phagocytosis rates in heat-

inactivated serum, where the major complement proteins (i.e. C3) are denatured. Overall, the

phagocytosis rates of *C. albicans* in heat-inactivated serum were lower than in live serum (Fig

4A), confirming the importance of opsonophagocytosis in the recognition of *C. albicans*.

However, adaptation of *C. albicans* to oestrogen did not result in further evasion of

phagocytosis in heat-inactivated serum (Fig 4A), suggesting that pre-exposure of *C. albicans*

to oestrogen impacts complement activation. Supplementation of heat-inactivated serum with

purified C3 restored oestrogen-induced immune evasion (Fig 4B), confirming that oestrogen-

induced immune evasion likely occurs through the avoidance of opsonophagocytosis.

Fig 4. Oestrogen promotes innate immune evasion of *C. albicans* through the inhibition

of opsonophagocytosis. A) J774A.1 macrophages were maintained in either complete

serum or heat-inactivated (HI) serum and infected with *C. albicans* pre-exposed to YPD, 0.3%

ethanol or 10 μM 17β-estradiol at a MOI of 5. **B)** J774A.1 macrophages were maintained in

either complete serum, heat-inactivated (HI) serum, or heat-inactivated serum supplemented

with purified C3 and infected with *C. albicans* pre-exposed to YPD, 0.3% ethanol or 10 μM

17β-estradiol at a MOI of 5. **C)** *C. albicans* was grown in YPD with or without 10 μM 17β-

estradiol, incubated in human serum for 20 minutes, fixed with PFA and C3 and C3b binding

quantified by FACS. **(D)**. *C. albicans* was grown in YPD with or without 10 μM 17β-estradiol,

incubated in human serum for 20 minutes, fixed with PFA and Factor H binding quantified by

FACS. Data represent the mean ± SEM from at least three independent experiments. (*p <

0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001).

Next, we tested whether adaptation of *C. albicans* to oestrogen affected the deposition of C3

and C3b on the fungal cell surface. Oestrogen-adapted cells exhibited enhanced C3 and C3b

deposition compared to non-adapted cells (Fig 4C), suggesting that adaptation to oestrogen

promotes C3 and C3b binding on the *C. albicans* cell surface. Increased deposition of

complement is normally associated with increased phagocytosis. However, processing of C3

can be inhibited by the recruitment of host regulatory proteins like Factor H, and several

pathogens are known to avoid opsonophagocytosis through enhanced recruitment of Factor

H [38, 39]. Therefore, the ability of oestrogen adapted *C. albicans* cells to bind Factor H was

quantified. Oestrogen-adapted cells bound significantly more Factor H, compared to the

solvent control (Fig 4D), suggesting that oestrogen-adapted *C. albicans* evade the innate

immune system through enhanced Factor H recruitment and decreased opsonophagocytosis.

Oestrogen-induced innate immune evasion is mediated through Gpd2

GPD2 is predicted to encode a cytosolic protein with glycerol-3-phosphate dehydrogenase

activity [40]. However, more recently, Gpd2 has been shown to localise to the fungal cell

surface in response to serum [41], and has been classed as a moonlighting protein (a protein

that has a biological function not predicted by its amino acid sequence). The moonlighting

activities of Gpd2 include binding Factor H, Factor H like protein-1 (FHL-1) and plasminogen

[40], key components of the alternative complement system. Binding of these host

complement regulatory proteins to microbial surfaces has been shown to promote immune

evasion via inhibition of complement-mediated phagocytosis [42]. Thus, we hypothesised that

adaptation of *C. albicans* to oestrogen promotes binding of these complement regulatory

proteins to Gpd2 thereby inhibiting opsonophagocytosis. To test this hypothesis, *GPD2* was

deleted in *C. albicans* and phagocytosis rates in the presence and absence of oestrogen

determined. Deletion of *GPD2* prevented oestrogen-dependent inhibition of macrophage

phagocytosis rates (Fig 5A), confirming that *GPD2* is required for oestrogen dependent innate

immune evasion.

Fig 5. Inhibition of opsonophagocytosis is dependent on Gpd2. A) J77A.1 macrophages

were infected with *C. albicans* strains grown in YPD with or without 10 μM 17β-estradiol and

phagocytosis rates quantified. **B)** J774A.1 macrophages were infected with CAI4-pSM2 and

CAI4-pSM2-GPD2 (*GPD2* over-expression strain) grown in YPD with or without 10 μM 17β-

estradiol and phagocytosis rates quantified. **C)** *C. albicans* strains were grown in YPD with or

without 10 μM 17β-estradiol, incubated in human serum for 20 minutes, fixed with PFA and

Factor H binding quantified by FACS. All data represent the mean ± SEM from at least three

independent experiments. (*p < 0.05, **p < 0.01, ***p < 0.001).

To determine whether increased expression of *GPD2* is sufficient to promote innate immune

evasion, we over-expressed *GPD2* in *C. albicans*. Over-expression of *GPD2* (2.75-fold

increased mRNA expression compared to wild type cells) resulted in reduced phagocytosis

rates irrespective of oestrogen treatment (Fig 5B), thereby confirming that enhanced

expression of *GPD2* is sufficient to promote *C. albicans* innate immune evasion. To determine

whether the expression levels of *GPD2* correlate with Factor H binding, we quantified Factor

H binding in the *gpd2Δ* mutant. As predicted, the *gpd2Δ* mutant did not bind more Factor H

than wild type cells in the presence of oestrogen (Fig 5C). Therefore, the expression of Gpd2

plays a key role in regulating the host-pathogen interaction.

Oestrogen elicits a mild transcriptional response in *C. albicans*.

To elucidate how quickly *C. albicans* adapts to oestrogen the fungus was grown in the

presence of 10 μM 17β-estradiol for varying lengths of time before macrophage phagocytosis

rates were quantified. This time course analysis confirmed that phagocytosis rates gradually

reduced after 60 minutes of growth in the presence of oestrogen (S2 Fig A), suggesting that

C. albicans undergoes some form of transcriptional or translational response to oestrogen. To

identify how oestrogen affects the global transcriptional profile of *C. albicans*, RNA-Seq was

performed on *C. albicans* cells that had been adapted to 10 μM 17β-estradiol for 4 hours.

Despite having a strong impact on the host-pathogen interaction, oestrogen only had a mild

effect on the transcriptome of *C. albicans*, (S2 Fig B, Table 1) with 5 genes being significantly

up-regulated and 23 genes being significantly down-regulated in the presence of oestrogen

(fold change > 2, FDR < 0.01). Genes that were upregulated were largely involved in cellular

redox or efflux, suggesting that oestrogen imposes oxidative stress on *C. albicans*. The most

significantly down regulated genes were *FRE7*, *SAP2*, *HWP1* and *ECE1*. *FRE7* is a cupric

reductase that has previously been confirmed to be repressed by oestrogen [43]. *ECE1* and *HWP1* are strongly associated with hyphal formation, suggesting that in YPD oestrogen represses hyphal formation. As phagocytosis occurs through interactions with the fungal cell surface we further analysed the expression of cell surface proteins in response to oestrogen stimulation to see if other surface proteins contributed to the observed innate immune evasion. In addition to *GPD2*, transcripts corresponding to *CDR1*, *EBP1, SOD6, ALS2, UTR2* and *FGR23* were moderately up regulated in the presence of oestrogen (Table 2). Oestrogen is a substrate of Cdr1, and it is thought that *CDR1* is upregulated in the presence of oestrogen to increase the efflux of oestrogen thereby reducing toxicity [43]. To determine whether Cdr1 is involved in oestrogen-dependent innate immune evasion, phagocytosis rates of the *cdr1Δ* and *cdr1Δ/cdr2Δ* mutants were quantified. Deletion of *CDR1* alone or in combination with *CDR2* did not affect oestrogen dependent inhibition of *C. albicans* phagocytosis (S2 Fig C).

Table 1 RNA Seq analysis

Gene name	Fold change	FDR p-value	Function
AOX2	4.25	5.04E-11	Alternative oxidase
orf19.3120	3.16	2.36E-07	PDR-subfamily ABC transporter
orf19.355	2.46	0.03	Predicted oxidoreductase activity
orf19.3337	2.18	4.62E-03	Protein of unknown function
GST1	2.17	0.04	Putative glutathione S-transferase
GAP2	-2.02	1.12E-03	Broad specificity amino acid permease
PST1	-2.03	4.12E-08	Flavodoxin-like protein
FOX2	-2.03	6.50E-11	3-hydroxyacyl-CoA epimerase
orf19.94	-2.07	2.09E-06	Protein of unknown function
MLS1	-2.11	8.93E-08	Malate synthase
ICL1	-2.15	1.71E-03	Isocitrate lyase
orf19.7310	-2.15	1.53E-10	Uncharacterized
CAT1	-2.33	4.45E-10	Catalase
ASR1	-2.33	4.69E-11	Heat shock protein
orf19.6688	-2.37	6.53E-03	Protein of unknown function
orf19.2515	-2.4	3.86E-10	ZZ-type zinc finger protein
IHD1	-2.4	1.72E-12	GPI-anchored protein
FDH1	-2.43	3.22E-09	Formate dehydrogenase
HSP30	-2.55	0.04	Heat shock protein
orf19.7029	-2.64	7.66E-03	Putative guanine deaminase
HPD1	-2.69	0.03	3-hydroxypropionate dehydrogenase

Gene name	Fold change	p-value	Function
GIT1	-2.73	1.85E-06	Glycerophosphoinositol permease
HMX1	-2.91	0	Heme oxygenase
FRE30	-3.17	3.71E-03	Similarity to ferric reductases
FRE7	-3.23	2.25E-03	Copper-regulated cupric reductase
SAP2	-3.49	1.20E-03	Secreted aspartyl proteinase
HWP1	-3.72	0.03	Host transglutaminase substrate
ECE1	-6.66	1.34E-03	Candidalysin cytolytic peptide toxin

323 **Table 2 Analysis of gene expression for cell surface proteins**

Gene name	Fold change	p-value	Function
EBP1	1.86	7.85E-09	NADPH oxidoreductase
CDR1	1.65	0.0000396	Multidrug transporter of ABC superfamily
SOD6	1.56	0.000749	Superoxide dismutase
ALS2	1.31	0.01	ALS family protein
UTR2	1.27	0.02	Cell wall glycosidase
FGR23	1.26	0.03	Protein of unknown function
GPD2	1.21	0.06	Glycerol 3-P dehydrogenase
ECM331	-1.21	0.03	GPI-anchored protein
HSP21	-1.22	0.07	Small heat shock protein
PGA4	-1.23	0.03	beta-1,3-glucanosyltransferase
CSA1	-1.25	0.03	Surface antigen
SAP10	-1.27	0.03	Secreted aspartyl protease
PLB4.5	-1.27	0.02	Phospholipase B
PGA23	-1.31	0.02	GPI-anchored protein of unknown function
PGA10	-1.32	0.01	GPI anchored membrane protein
RBT4	-1.32	0.01	Pry family protein
HSP104	-1.35	0.00253	Heat-shock protein
PHR1	-1.36	0.03	Cell surface glycosidase
IFF4	-1.37	0.02	Adhesin-like cell surface protein
CSP37	-1.37	0.00229	Hyphal cell wall protein
GPH1	-1.43	0.000767	Putative glycogen phosphorylase
XOG1	-1.46	0.000325	Exo-1,3-beta-glucanase
HSP70	-1.47	0.000249	Putative Hsp70 chaperone
CSH1	-1.55	0.0000623	Aldo-keto reductase
MET6	-1.62	3.51E-08	5-methyltetrahydropteroyltriglutamate-homocysteine methyltransferase
PGA26	-1.65	0.07	GPI-anchored adhesin-like protein
PHO100	-1.77	0.05	Putative inducible acid phosphatase
DDR48	-1.77	0.000000264	Immunogenic stress-associated protein
IFF9	-1.89	0.02	Predicted GPI-linked cell-wall protein

IHD1	-2.4	5.55E-16	GPI-anchored protein
PGA15	-2.43	0.06	Putative GPI-anchored protein
PGA46	-2.44	0.06	Putative GPI-anchored protein
HWP1	-3.72	0.000928	Host transglutaminase substrate

Ebp1 is a negative regulator of innate immune evasion.

The most significantly differentially regulated cell surface gene was oestrogen binding protein 1 (Ebp1). Ebp1 is homologous to old yellow enzyme 2 (*OYE2*) of *S. cerevisiae* and is an NADPH oxidoreductase [16]. Oestrogen inhibits the NADPH oxidoreductase activity of recombinant Ebp1, yet the function of this enzyme in the *C. albicans* oestrogen response is unknown. To determine whether Ebp1 plays a role in oestrogen induced innate immune evasion, *EBP1* was deleted in *C. albicans* and phagocytosis rates of the mutant in the presence and absence of oestrogen quantified. Deletion of *EBP1* resulted in consistently low phagocytosis even in the absence of hormone stimulation (Fig 6A). To determine whether this enhanced innate immune evasion was linked to *GPD2*, the expression of *GPD2* in the *ebp1Δ* mutant was quantified by RT-PCR. Deletion of *EBP1* resulted in 2.74-fold higher expression of *GPD2* (comparable levels of *GPD2* expression to the over-expression strain) compared to wild type cells (Fig 6B), suggesting Ebp1 is a negative regulator of *GPD2* expression and therefore innate immune evasion.

Fig 6. Adaptation of *C. albicans* to oestrogen is mediated via Ebp1 and Bcr1 A) The

*ebp1*Δ mutant and the parental control strain were grown with or without 10 μM 17β-estradiol

for 4 hrs and then exposed to J774A.1 macrophages for 45 minutes and phagocytosis rates

quantified. **B)** The *ebp1Δ* mutant and the parental control strain were grown in YPD to mid-log phase, total RNA extracted and *GPD2* gene expression quantified by RT-PCR relative to *ACT1*. **C)** The *bcr1Δ* mutant and the parental control strain were grown in YPD to mid-log phase, total RNA extracted and *GPD2* gene expression quantified by RT-PCR relative to *ACT1*. **D)** The *bcr1Δ* mutant and parental control strain were grown with or without 10 μM 17β-estradiol for 4 hrs and then exposed to J774A.1 macrophages for 45 minutes and phagocytosis rates quantified. **E)** The *rob1Δ* mutant and the parental control strain were grown in YPD with or without 10 μM 17β-estradiol for 4 h and then co-incubated with J774A.1 macrophages for 45 min and phagocytosis rates quantified. All data represent the mean and SEM from three biological replicates (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$).

Oestrogen dependent innate immune evasion requires a threshold level of *GPD2* expression.

To identify transcription factors involved in the expression of *GPD2,* we analysed the promoter region of *GPD2*. The Pathoyeastract program [44] identified 17 transcription factors associated with the promoter region of *GPD2.* Of these 17 transcription factors, 10 have been deleted in *C. albicans* and are available in the transcription factor knockout library [45]. To investigate the role of these 10 transcription factors in the regulation of *GPD2*, the expression of *GPD2* was quantified by qPCR. Of the 10 transcription factor mutants tested, deletion of *BCR1* and *ROB1* had a significant impact on the expression of *GPD2* (Fig 6C, S3 Fig), reducing the expression of *GPD2* to 25% of the wildtype control. To assess whether the reduced expression levels of *GPD2* in the *bcr1Δ* and *rob1Δ* mutants are sufficient to perturb the response to oestrogen, the phagocytosis rates of the *bcr1Δ* and *rob1Δ* mutants in the presence or absence of oestrogen were quantified. Deletion of *BCR1* resulted in a loss of oestrogen-mediated innate immune evasion (Fig 6D), while deletion of *ROB1* partially blocked innate immune evasion (Fig 6E), confirming that a threshold level of *GPD2* expression is required for innate immune evasion.

Oestrogen-induced immune evasion plays a key role in *C. albicans* pathogenicity.

Having established that adaptation of *C. albicans* to oestrogen inhibits phagocytosis, we explored whether this phenomenon could influence virulence *in vivo* using a zebrafish larval model for disseminated disease. Previously, it was shown that exposing zebrafish larvae (3 hours post fertilisation) to media containing 1 µM oestrogen results in an *in vivo* oestrogen concentration of 0.0057 µM, equivalent to physiological levels during pregnancy in humans [46, 47]. Taking advantage of this observation, zebrafish larvae were infected with *C. albicans* SC5314 and maintained in E3 media supplemented with 1 µM oestrogen and survival rate monitored for up to 5 days post fertilisation. Compared to zebrafish incubated in E3 media alone, or E3 media supplemented with ethanol, supplementation of E3 media with 1 µM oestrogen enhanced the virulence of *C. albicans,* resulting in an 63% reduction in zebrafish survival 60 hrs post infection (Fig 7A).

To investigate whether Gpd2 is required for virulence in this model, zebrafish were infected with the parental control strain, the *gpd2*Δ mutant, or reconstituted control strains of *C. albicans*. Deletion of *GPD2* led to attenuation of *C. albicans* virulence, which was restored to parental control levels via complementation with a single copy of *GPD2* (Fig 7B). To assess whether increased expression of Gpd2 is sufficient to drive fungal virulence, *GPD2* was over expressed in the CAI4 background. Infection of zebrafish larvae with *C. albicans* over-expressing *GPD2* enhanced *C. albicans* virulence in the absence of oestrogen compared to the respective control strain (Fig 7C). Therefore, Gpd2 plays a key role in promoting *C. albicans* virulence *in vivo*.

A

B

C

Fig 7. **Oestrogen promotes the virulence of *C. albicans* in a zebrafish larval infection model. A)** *C. albicans* (SC5314) were microinjected into the hindbrain ventricle of zebrafish larvae in the Prim25 stage. Infected larvae were maintained in E3 media, or E3 media supplemented with 0.3% ethanol, or 1 µM 17β-estradiol and larval survival monitored every 24 h until day 5 post fertilisation. **B**) SN250-CIP30, *gpd2Δ-CIP30* and *gpd2Δ-CIP30-GPD2* were microinjected into the hindbrain ventricle of zebrafish larvae in the Prim25 stage dpf. Larvae were maintained in E3 media and larval survival monitored every 24 h until day 5 post fertilisation. **C)** CAI4-pSM2 or CAI4-pSM2-GPD2 were microinjected into the hindbrain ventricle of zebrafish larvae in the Prim25 stage dpf. Larvae were maintained in E3 media and larval survival monitored every 24 h until day 5 post fertilisation. The survival curves represent data pooled from three independent experiments. Statistically significant differences were determined by Log-rank (Mantel-Cox) test. (*$p < 0.05$, **$p < 0.01$; ***$p < 0.001$).

Discussion

Vulvovaginal candidiasis (VVC) is a mucosal infection affecting 75% of the female population of reproductive age [11]. Oestrogen is known to govern susceptibility to VVC infections, with women with low circulatory oestrogen levels (i.e. postmenopausal women) having a low risk of developing VVC, and women with high oestrogen levels (i.e. during pregnancy, or women taking high oestrogen containing oral contracepts) having a high risk of VVC. Elevated oestrogen levels increase glycogen at the vaginal mucosa, reduce leukocyte infiltration, and reduce antifungal activity of epithelial cells, promoting infection [10, 12, 13]. However, here we show that, in addition to these effects on the host, oestrogen promotes adaptation responses in *C. albicans* which induce evasion of the innate immune system through Gpd2-dependent inhibition of complement mediated opsonophagocytosis. Unlike the induction of fungal morphogenesis, which appears to be limited to 17β-estradiol [43], all tested forms of oestrogen promoted innate immune evasion, suggesting that *C. albicans* has evolved at least two signalling pathways that are responsive to oestrogen.

Complement evasion is a successful mechanism employed by viruses, bacteria, parasites and fungi to escape the innate immune system [48]. Evasion of the complement system can be mediated through the secretion of degradative proteins, or by recruitment of host regulatory proteins [49]. One of the most common methods pathogens use to evade complement is through enhanced recruitment of Factor H, a key regulatory protein in the complement system, to the microbial surface. Bound Factor H prevents further activation of the alternative complement system through both the destabilisation of the C3 convertase and enhancement of Factor I mediated degradation of C3b to iC3b, reducing opsonisation and inhibiting the formation of the membrane attack complex [49]. *C. albicans* has been shown by others to recruit Factor H to its surface through the expression of moonlighting proteins. Moonlighting proteins are proteins that perform a biological function not predicted from the amino acid sequence. So far, four moonlighting proteins (Phr1, Gpd2, Hgt1 and Gpm1) have been identified in *C. albicans* [40, 50-52]. Although Gpd2 is predicted to be a cytoplasmic protein involved in cellular redox, Gpd2 has also been identified in cell wall proteomic studies [41], suggesting that Gpd2 functions in the cell wall. Interestingly, Gpd2 was only identified in the cell wall proteome of *C. albicans* grown in complete serum, and not heat-inactivated serum where complement proteins have been inactivated [41]. Therefore, deposition of complement on the fungal cell surface may drive localisation of Gpd2 to the cell surface.

Purified Gpd2 binds all three complement regulatory proteins [40], although the biological significance of these interactions in infections is not known. Factor H and FHL1 bound to Gpd2 remain active, cleaving C3b and thereby inhibiting the alternative complement system [40]. In addition, Gpd2 can also bind plasminogen, which is then processed into plasmin, contributing to inactivation of the alternative complement system [40]. FACS analysis of the deposition of complement proteins on the surface of *C. albicans* in response to oestrogen suggested that oestrogen-adapted cells bound more Factor H, indicative of oestrogen promoting complement evasion. However, oestrogen-adapted *C. albicans* also bound more C3 and C3b, which is

surprising as binding of Factor H should prevent C3 deposition. Factor H is a glycoprotein that

is made up of 20 complement control proteins (CCP). Microbes bind Factor H at various

positions, with *Neisseria* species binding Factor H at CCP6-8 [53]. However, multiple microbes

bind Factor H at CCP20, and this has been termed the Common Microbial Binding site [54].

Binding of Factor H to some of these microbial cell surface proteins at CCP20 results in

increased affinity of Factor H for C3b [54]. The enhanced affinity of Factor H for C3b might

explain why we observe elevated binding of both C3b and Factor H, and would suggest that

Gpd2 binds to CCP20. The formation of this stable tri-partite complex (microbial protein,

Factor H, C3b) results in enhanced activity of Factor H and therefore rapid inactivation of the

alternative complement cascade [54], which would explain why we observe reduced levels of

phagocytosis.

Innate immune evasion was only observed after *C. albicans* had been grown in the presence

of oestrogen for 60 minutes, suggesting that the response required either transcriptional or

post-transcriptional regulation. However, in line with previous studies [43], global

transcriptional analysis confirmed that only a small proportion of *C. albicans* genes were

differentially regulated by oestrogen. In comparison to the work of Cheng *et al.,* where most

of the differentially regulated genes are involved in fungal morphogenesis [43], our analysis

mainly identified genes involved in metabolism and oxidation-reduction processes. This lack

of overlap is likely reflective of the growth conditions used, as Cheng *et al.,* used RPMI where

oestrogen promotes hyphal formation [13, 43], while under our experimental conditions (YPD)

oestrogen did not promote hyphal formation. Although innate immune evasion was dependent

on the presence of Gpd2, *GPD2* mRNA levels were only moderately increased in response to

oestrogen, suggesting that other factors (i.e. post-translational modifications, protein

localisation) may be involved. Given that Gpd2 has only been identified as a cell wall protein

under specific conditions [41], it might be that in response to oestrogen a higher level of Gpd2

is recruited to the cell wall, enabling the fungus to bind more Factor H. In *S. cerevisiae*, the

activity of both Gpd1 and Gpd2 is regulated via post-translational modification [55]. Snf1-

dependent phosphorylation of *Sc*Gpd1 (orthologue of Ca*GPD2*) results in decreased enzyme

activity [55]. Although *Ca*Gpd2 has several predicted phosphorylation sites, these are located

in different regions of the protein compared to *Sc*Gpd1, and therefore it is possible that

phosphorylation/dephosphorylation at any of these sites in *Ca*Gpd2 could affect the

localisation, or potential to bind Factor H.

Ebp1, despite showing no homology to mammalian hormone binding proteins, has been

described as an oestrogen binding protein. Ebp1 is homologous to old yellow enzyme 2

(*OYE2*) of *S. cerevisiae*, an NADPH oxidoreductase. Biochemical characterisation of

recombinant Ebp1 confirms that Ebp1 is an NADPH oxidoreductase, and that oestrogen

inhibits this enzymatic activity [16, 56]. The importance of Ebp1 enzymatic activity in *C.*

albicans pathogenicity is not known, but here we demonstrate that Ebp1 may serve as a

negative regulator of innate immune evasion through the regulation of *GPD2*. Interaction of

oestrogen with Ebp1, through an as yet unidentified pathway likely involving Bcr1, results in

elevated expression of *GPD2* and, potentially in association with other post-translational

modifications, promotes innate immune evasion and pathogenicity (Fig 8).

Fig 8 Proposed mechanism of oestrogen-induced innate immune evasion. Under standard laboratory conditions the C3 is deposited on the surface of *C. albicans* resulting in effective phagocytosis. However, in the presence of oestrogen the NADPH activity of Ebp1 is reduced resulting in derepression (through an as yet to be identified signalling pathway) and surface localisation of Gpd2. Gpd2 recruits Factor H to the fungal cell surface, preventing the formation of the C3 convertase and disposition of C3 on the fungal cell surface. Reduced opsonisation results in innate immune evasion and insufficient clearance of *C. albicans*, promoting fungal pathogenicity.

In recent years zebrafish larvae have become an excellent model for studying pathogenicity mechanisms. Zebrafish have a complement system that is structurally and functionally similar to the mammalian complement system [57], and an orthologue of Factor H has been identified and cloned [58]. The presence of oestrogen in the zebrafish larval model of disseminated *C. albicans* infection, confirmed that oestrogen promotes the virulence of *C. albicans*, suggesting that oestrogen dependent inactivation of the alternative complement system promotes the virulence of *C. albicans in vivo*. In support of this, deletion of *GPD2* attenuated fungal virulence, while over expression of *GPD2* enhanced virulence in the zebrafish model of infection, confirming that Gpd2 is an important fungal virulence factor. Given the strong association of elevated oestrogen levels with the development of VVC, it is important to understand the role of Gpd2 in promoting innate immune evasion, and may lead to the development of alternative treatment options for VVC and RVVC.

Materials and Methods

Ethics

Protocols for human blood collection and isolation of neutrophils and peripheral blood mononuclear cells (PBMCs) were approved by ethical review board of the School of

Biosciences at the University of Birmingham. Blood was collected anonymously and on

voluntary basis after getting written informed consent.

Zebrafish care and experiment protocols were performed in accordance with the Home Office

project license 40/3681 and personal license IE905E215 as per Animal Scientific Procedures

Act 1986.

Strains and media

Unless indicated otherwise, all media and chemicals were purchased from Sigma-Aldrich UK.

All strains used in the study are listed in S1 Table. *Candida* strains were routinely maintained

on YPD agar (1% yeast extract, 1% peptone, 2% glucose and 2% agar). For broth cultures all

strains were cultured in YPD (1% yeast extract, 1% bacto-peptone, 2% glucose) buffered to

pH6 with 3.57% HEPES. Oestrogen was diluted in 10% ethanol to a stock concentration of

100 μg/ml and diluted into YPD at the required concentrations, maintaining the final ethanol

concentration at 0.3%.

Phagocytosis experiments

Phagocytosis assay was performed as previously described [59]. Briefly, overnight cultures of

C. albicans were sub-cultured 1:100 in fresh YPD media, media supplemented with 0.3%

ethanol, or media supplemented with oestrogen (0.0001 μM, 0.01 μM or 10 μM) and incubated

at 37ºC, 200 rpm for 4 h. *C. albicans* cells were washed three times in PBS and 1 x10^5 J774A.1

macrophages (Sigma, UK) were infected with 5 x 10^5 yeast cells (multiplicity of infection [MOI]

of 5) for 45 min at 37°C, 5% CO_2. Cells were washed with PBS to remove non-phagocytosed

yeast cells, and phagocytosis stopped by fixing with 4% paraformaldehyde (PFA) for 20

minutes. Samples were stained for 30 minutes with 50 μg/ml ConA-FITC to stain non-

phagocytosed fungal cells, washed and imaged. Phagocytosis events were scored from

multiple fields of view using imageJ. When required, J774A.1 cells were maintained in DMEM

supplemented with heat inactivated serum and the assay was complemented with 1 µg/mL

C3 (Sigma, C2910) or C3b (Sigma, 204860). All experiments were performed in technical

duplicates, with a minimum of three independent biological repeats. Data were analysed by

Kruskal-Wallis test followed by a post-hoc Dunn's multiple comparisons test at 95%

confidence.

Human macrophages and neutrophils

PBMCs and neutrophils were isolated as previously described [20]. Neutrophils were seeded

at 2×10^5 cells/mL in 24-well plates in serum free RPMI supplemented with 100 mM L-

glutamine, incubated for 1 h at 37°C, 5% CO_2 and then co-incubated with 1×10^6 *C. albicans*

cells (MOI = 5) for 45 min at 37°C, 5% CO_2. Cells were immediately fixed with 4% PFA and

stained with 50 μg/ml ConA-FITC for 30 minutes to stain non-phagocytosed fungal cells,

washed and imaged. Phagocytosis events were scored from multiple fields of view using

imageJ. To assess primary macrophage phagocytosis rates PBMCs (0.5×10^6) were seeded

into 24 well plates in differentiation media (RPMI 1640 supplemented with 100 mM L-

glutamine, 10% human AB serum, 100 mM Pen/Strep and 20 ng/ml M-CSF) for 7 days

replacing the media every 2-3 days, and then phagocytosis rates determined as described

above. To assess cytokine production 2.5×10^4 PBMCs were stimulated with 5×10^4 PFA fixed

C. albicans for 24 hours, supernatants collected and stored at -20°C and cytokine

concentrations quantified by ELISA. All experiments were performed in technical duplicates,

with a minimum of three independent biological repeats using different donors. Data were

analysed by Kruskal-Wallis test followed by a post-hoc Dunn's multiple comparisons test at

95% confidence.

Genetic manipulation of *C. albicans*

All primers used in genetic manipulation of *C. albicans* are listed in S2 Table. To reintroduce

GPD2 into *C. albicans* *gpd2*Δ mutant, the *GPD2* locus was PCR-amplified from *C. albicans*

SC5314 genomic DNA using primers GPD2-SacI-F and GPD2-NotI-R. The PCR product was

cloned into CIp30 plasmid[60] restricted with *SacI* and *NotI* using T4 DNA ligase. The

generated CIp30-*GPD2* was linearized with *StuI* and transformed into *C. albicans gpd2*Δ

mutant by standard heat-shock transformation [61].

To generate a *C. albicans* strain that over expresses *GPD2*, the open reading frame of *GPD2*

was cloned into pSM2[62] using primers GPD2-OE-F and GPD2-OE-R under the control of

the *TEF2* promoter. The resulting plasmid was linearized with *PacI*, and integrated into CAI4

at the *URA3* locus by standard heat-shock transformation.

To generate *ebp1*Δ, 500 bp of the 5' and 3' UTR were amplified from genomic DNA using

primers EBP1-5F, EBP1-5R, and EBP1-3F and EBP1-3R. The resulting PCR products were

purified, digested with *EcoRV* and *SacI* or *HindIII* with *KpnI* and cloned into the mini URA

blaster cassette pDDB57 [63]. The resulting disruption cassette was digested with *EcoRV* and

KpnI and transformed in CAI4 by standard heat-shock transformation. Resulting colonies were

screened by PCR and positive colonies were plated onto YNB supplemented with 5-

fluoroorotic acid (5-FOA) and uridine to select for spontaneous homologous recombination

and loss of *URA3*. Resulting colonies were then re-transformed with the *EBP1* knockout

cassette, and loss of both alleles was confirmed by PCR, and lack of expression was

confirmed by qPCR. The URA baster cassette was recycled and *URA3* replaced at its native

locus by transforming the strain with pSM2.

Immunofluorescent staining of *C. albicans* cell wall components

C. albicans cells were stained as previously described [20]. Briefly, *C. albicans* cells from

overnight culture were sub-cultured in YPD broth with or without oestrogen supplementation

and incubated at 37°C, 200 rpm for 4 h. Cells were harvested by centrifugation, washed in

PBS and fixed with 4% PFA. To quantify total mannan, glucan and chitin levels in the cell wall,

cells were stained with 50 μg/ml TRITC-conjugated concanavalin A (Molecular Probes, Life

Technologies), 33.3 μg/ml Aniline Blue fluorochrome (Bioscience supplies) and 3 μg/ml

Calcofluor White for 30 minutes. To quantify surface exposure of β1,3-glucan and chitin, cells

were stained with 3 μg/ml Fc-Dectin-1 (a gift from G. Brown, University of Aberdeen) and 50

μg/ml TRITC-conjugated wheat germ agglutinin (Molecular Probes, Life Technologies). Cells

were analysed by flow cytometry and median fluorescence intensity (MFI) quantified. FACS

data were analysed by Kruskal-Wallis test followed by a post-hoc Dunn's multiple comparisons

test at 95% confidence.

RNA sequencing

C. albicans cells were grown for 4 h in YPD broth with or without 10 μM 17β-estradiol at 37ºC,

200 rpm. Cells were harvested by centrifugation, washed three times in PBS, and snapped

frozen in liquid nitrogen. Total RNA was extracted as per manufacturer's instructions using the

RNeasy Plus mini kit (Qiagen). Total RNA was quantified using NanoDrop 8000

spectrophotometer (ND-8000-GL; Thermo Fisher). RNA samples were assessed for genomic

DNA contamination by PCR and agarose gel electrophoresis. Samples were then processed

as previously reported [59]. Sequencing reads are available at the Gene Expression Omnibus

(GEO) database (http://www.ncbi.nlm.nih.gov/geo/) at the following accession number

GSE145240.

Reads were analysed following a previous published method[59] and using CLC Genomic

workbench 11.0.1 software (Qiagen). In summary, adapter and quality trimming was

performed before reads were mapped to *C. albicans* reference genome (Assembly 21, version

s02-m09-r10). Transcript Per Kilobase Million (TPM) were reported for each open reading

frame (ORFs). Statistical analysis was performed after addition to all values of the lowest TPM

measurement, then data were log10 transformed and differential expression between

condition were considered significant if the absolute value Fold Change >2 and FDR < 0.01.

CGD GO term finder [64] was used to perform Gene ontology (GO) analysis with P-values

corresponding to Bonferroni-corrected hypergeometric test P-values.

RT-PCR

C. albicans cells were grown for 4 h in YPD broth with or without 10 μM 17β-estradiol at 37ºC, 200 rpm. Cells were harvested by centrifugation, and snapped frozen in liquid nitrogen. RNA was extracted using the RNeasy Plus mini kit (Qiagen) as per manufacturer's instructions. RNA quality and quantity were checked by electrophoresis and spectroscopy. The qRT-PCR was performed using the 2x qPCRBIO SyGreen mix kit (PCRbiosystem) according to manufacturer's recommendations with 50 ng of total RNA (primers shown in S2 Table). Relative quantification of gene expression was determined by the Delta Delta Ct method with *ACT1* as an endogenous control. mRNA expression was performed in technical triplicate, and data represent the mean and SEM from three independent biological repeats, and were analysed using a paired T-test with 95% confidence.

Complement binding

C. albicans cells were grown for 4 h in YPD broth with or without 10 μM 17β-estradiol at 37ºC, 200 rpm. Cells were harvested, washed three times in PBS, fixed on ice for 30 min with 4% PFA. About 2 x 10^6 yeast cells were labelled with 400 μL 10% normal human serum for 20 min at 37ºC, 200 rpm. Cells were washed thrice in PBS and incubated on ice with 100 μL of either 10 µg/mL Anti-Factor H Goat pAb (Sigma, 341276-1ML) or 1 µg/mL Goat anti Chicken IgY (H+L) diluted in 1% BSA/PBS. Cells were washed thrice in PBS and incubated in dark with 100 µL of either Rabbit anti Goat IgG (H+L) Secondary Antibody, Alexa Fluor 488 (Invitrogen, A11078) or Goat anti Chicken IgY (H+L) Secondary Antibody, Alexa Fluor 594, (Invitrogen, A11042) diluted 1:200 in 1% BSA/PBS. Cells were analysed by flow cytometry and median fluorescence intensity determined. FACS data were analysed by Kruskal-Wallis test followed by a post-hoc Dunn's multiple comparisons test at 95% confidence.

Zebrafish infection

Wild type (AB) *Danio rerio* zebrafish used in the study were housed in a recirculating system

of gallon tanks at the University of Birmingham Zebrafish Facility. To obtain embryos, 4 male

and 5 female fish were transferred into a breeding tank and maintained at 28°C, 14 h light/10

h dark cycle. Embryos were collected the following day, sorted and incubated at 30°C for 24

h in E3 media (5 mM NaCl, 0.17 mM KCl, 0.33 mM $CaCl_2$, 0.33 mM $MgSO_4$, 0.00003%

methylene blue, pH 7). Embryos were maintained at a density of 100 per 14.5 cm dish

containing 150 mL E3 media supplemented with 0.02 mg/mL Phenylthiourea.

Hind brain infections were performed as previously described [65]. Briefly, zebrafish at the

prim25 stage were manually dechorionated, and anesthetized in 160 μg/mL Tricaine.

Approximately 5 nL of injection buffer (10% PVP-40 in PBS, 0.05% phenol red) or *C. albicans*

suspension at 5×10^7 cells/mL in injection buffer was microinjected into the hindbrain ventricle

via the otic vesicle to achieve a dose of 20-50 yeast/larva. Within 1 h of infection, larvae were

screened by microscopy to remove fish noticeably traumatised from microinjection and to

ascertain correct injection site and inoculum size. At least 15 larvae per condition were

transferred into a 6-well plate, incubated at 28°C in E3 media either with or without 1 μM

oestrogen and observed for survival every 24 h until day 5 post fertilisation. Data from three

independent biological replicates were pooled together to determine percent survival. Data

were analysed by Log-rank Mantel-Cox test and Gehan-Breslow-Wilcoxon test (extra weight

for early time points).

Statistics

Unless indicated otherwise, data were analysed in Prism (version 8) using a Kruskal-Wallis

test followed with a post-hoc Dunn's multiple comparisons test at 95% confidence. Data

represent the mean +/- SEM from at least three independent biological experiments.

Acknowledgements

We would like to thank the fungal community for the donation of fungal strains, Prof G. Brown (University of Exeter) for FC-Dectin-1, Prof G. Ramage for clinical isolates, the Environmental Omics facility at the University of Birmingham for help with the RNA sequencing, and Elizabeth Ballou (University of Birmingham) for critical reading of the manuscript. PK is funded by the Wellcome trust strategic award for medical mycology (097377/Z/11/Z), FC and RAH are funded by the MRC (MR/L00903X/1) and BBSRC (BB/R00966X/1).

References

1.   Ghazeeri G, Abdullah L, Abbas O. Immunological differences in women compared with men: overview and contributing factors. American journal of reproductive immunology (New York, NY : 1989). 2011;66(3):163-9. Epub 2011/07/14. doi: 10.1111/j.1600-0897.2011.01052.x. PubMed PMID: 21749547.

2.   García-Gómez E, González-Pedrajo B, Camacho-Arroyo I. Role of sex steroid hormones in bacterial-host interactions. BioMed research international. 2013;2013:928290. Epub 2013/03/20. doi: 10.1155/2013/928290. PubMed PMID: 23509808; PubMed Central PMCID: PMCPMC3591248.

3.   van Lunzen J, Altfeld M. Sex differences in infectious diseases-common but neglected. J Infect Dis. 2014;209 Suppl 3:S79-80. Epub 2014/06/27. doi: 10.1093/infdis/jiu159. PubMed PMID: 24966193.

4.   Klein SL. The effects of hormones on sex differences in infection: from genes to behavior. Neuroscience & Biobehavioral Reviews. 2000;24(6):627-38. doi: https://doi.org/10.1016/S0149-7634(00)00027-0.

5.   Taneja V. Sex Hormones Determine Immune Response. Front Immunol. 2018;9(1931). doi: 10.3389/fimmu.2018.01931.

6.   McClelland EE, Smith JM. Gender specific differences in the immune response to infection. Archivum immunologiae et therapiae experimentalis. 2011;59(3):203-13. Epub 2011/03/29. doi: 10.1007/s00005-011-0124-3. PubMed PMID: 21442309.

7.   Amirshahi A, Wan C, Beagley K, Latter J, Symonds I, Timms P. Modulation of the

Chlamydia trachomatis in vitro transcriptome response by the sex hormones estradiol and

progesterone. BMC Microbiol. 2011;11:150. Epub 2011/06/28. doi: 10.1186/1471-2180-11-

150. PubMed PMID: 21702997; PubMed Central PMCID: PMCPMC3224131.

8.   Chotirmall SH, Smith SG, Gunaratnam C, Cosgrove S, Dimitrov BD, O'Neill SJ, et al.

Effect of estrogen on *Pseudomonas* mucoidy and exacerbations in cystic fibrosis. N Engl J

Med. 2012;366(21):1978-86. Epub 2012/05/23. doi: 10.1056/NEJMoa1106126. PubMed

PMID: 22607135.

9.   Sobel JD. Vaginitis. New England Journal Medicine. 1997;337:1896-903.

10.   Dennerstein GJ, Ellis DH. Oestrogen, glycogen and vaginal candidiasis. Australian and

New Zealand Journal of Obstetrics and Gynaecology. 2001;41(3):326-8. doi: 10.1111/j.1479-

828X.2001.tb01238.x.

11.   SOBEL JD. Pathogenesis and Epidemiology of Vulvovaginal Candidiasis. Annals of

the   New   York   Academy   of   Sciences.   1988;544(1):547-57.   doi: 10.1111/j.1749-

6632.1988.tb40450.x.

12.   Fidel PL, Jr., Cutright J, Steele C. Effects of reproductive hormones on experimental

vaginal   candidiasis.   Infection   and   immunity.   2000;68(2):651-7.   doi: 10.1128/iai.68.2.651-

657.2000. PubMed PMID: 10639429.

13.   Salinas-Muñoz L, Campos-Fernández R, Mercader E, Olivera-Valle I, Fernández-

Pacheco C, Matilla L, et al. Estrogen Receptor-Alpha (ESR1) Governs the Lower Female

Reproductive Tract Vulnerability to *Candida albicans*. Front Immunol. 2018;9:1033-. doi:

10.3389/fimmu.2018.01033. PubMed PMID: 29881378.

14.   Kinsman OS, Collard AE. Hormonal factors in vaginal candidiasis in rats. Infection and

immunity. 1986;53(3):498-504. PubMed PMID: 3527984.

15.   White S, Larsen B. *Candida albicans* morphogenesis is influenced by estrogen.

Cellular and Molecular Life Sciences CMLS. 1997;53(9):744-9. doi: 10.1007/s000180050094.

16. Madani ND, Malloy PJ, Rodriguez-Pombo P, Krishnan AV, Feldman D. *Candida*

albicans estrogen-binding protein gene encodes an oxidoreductase that is inhibited by

estradiol. PNAS. 1994;91(3):922-6. doi: 10.1073/pnas.91.3.922. PubMed PMID: 8302868.

17. Netea MG, Brown GD, Kullberg BJ, Gow NAR. An integrated model of the recognition

of *Candida albicans* by the innate immune system. Nature Reviews: Microbiology.

2008;6(1):67-78.

18. Hall RA, Gow NAR. Mannosylation in *Candida albicans*: role in cell wall function and

immune recognition. Molecular Microbiology. 2013.

19. Hall RA. Dressed to impress: impact of environmental adaptation on the

Candida albicans cell wall. Molecular Microbiology. 2015;97(1):7-17. doi: 10.1111/mmi.13020.

20. Sherrington SL, Sorsby E, Mahtey N, Kumwenda P, Lenardon MD, Brown I, et al.

Adaptation of *Candida albicans* to environmental pH induces cell wall remodelling and

enhances innate immune recognition. PLOS Pathogens. 2017;13(5):e1006403. doi:

10.1371/journal.ppat.1006403.

21. Pradhan A, Avelar GM, Bain JM, Childers DS, Larcombe DE, Netea MG, et al. Hypoxia

Promotes Immune Evasion by Triggering β-Glucan Masking on the *Candida albicans* Cell

Surface via Mitochondrial and cAMP-Protein Kinase A Signaling. mBio. 2018;9(6):e01318-18.

doi: 10.1128/mBio.01318-18. PubMed PMID: 30401773.

22. Pradhan A, Avelar GM, Bain JM, Childers D, Pelletier C, Larcombe DE, et al. Non-

canonical signalling mediates changes in fungal cell wall PAMPs that drive immune evasion.

Nature communications. 2019;10(1):5315-. doi: 10.1038/s41467-019-13298-9. PubMed

PMID: 31757950.

23. Lopes JP, Stylianou M, Backman E, Holmberg S, Jass J, Claesson R, et al. Evasion

of Immune Surveillance in Low Oxygen Environments Enhances *Candida albicans* Virulence.

mBio. 2018;9(6). Epub 2018/11/08. doi: 10.1128/mBio.02120-18. PubMed PMID: 30401781;

PubMed Central PMCID: PMCPMC6222133.

24. Williams RB, Lorenz MC. Multiple Alternative Carbon Pathways Combine To Promote

Candida albicans Stress Resistance, Immune Interactions, and Virulence. mBio. 2020;11(1).

Epub 2020/01/16. doi: 10.1128/mBio.03070-19. PubMed PMID: 31937647; PubMed Central

PMCID: PMCPMC6960290.

25. Tripathi A, Liverani E, Tsygankov A, Puri S. Iron alters the cell wall composition and

intracellular lactate to affect *Candida albicans* susceptibility to antifungals and host immune

response. J Biol Chem. 2020. Epub 2020/06/07. doi: 10.1074/jbc.RA120.013413. PubMed

PMID: 32503842.

26. Pericolini E, Perito S, Castagnoli A, Gabrielli E, Mencacci A, Blasi E, et al. Epitope

unmasking in vulvovaginal candidiasis is associated with hyphal growth and neutrophilic

infiltration.   PLoS   One.   2018;13(7):e0201436.   Epub   2018/08/01.   doi:

10.1371/journal.pone.0201436.   PubMed   PMID: 30063729;   PubMed   Central   PMCID:

PMCPMC6067721.

27. Hopke A, Nicke N, Hidu EE, Degani G, Popolo L, Wheeler RT. Neutrophil Attack

Triggers Extracellular Trap-Dependent *Candida* Cell Wall Remodeling and Altered Immune

Recognition. PLoS Pathogens. 2016;12(5):e1005644. doi: 10.1371/journal.ppat.1005644.

PubMed PMID: PMC4880299.

28. Wheeler RT, Fink GR. A Drug-Sensitive Genetic Network Masks Fungi from the

Immune System. PLoS Pathog. 2006;2(4):e35.

29. Wheeler RT, Kombe D, Agarwala SD, Fink GR. Dynamic, Morphotype-Specific

Candida albicans β-Glucan Exposure during Infection and Drug Treatment. PLoS Pathogens.

2008;4(12):e1000227. doi: 10.1371/journal.ppat.1000227. PubMed PMID: PMC2587227.

30. Ballou ER, Avelar GM, Childers DS, Mackie J, Bain JM, Wagener J, et al. Lactate

signalling regulates fungal β-glucan masking and immune evasion. Nature Microbiology.

2016;2:16238. doi: 10.1038/nmicrobiol.2016.238.

31. Kozel TR, Weinhold LC, Lupan DM. Distinct characteristics of initiation of the classical

and alternative complement pathways by *Candida albicans*. Infect Immun. 1996;64(8):3360-

8. Epub 1996/08/01. PubMed PMID: 8757876; PubMed Central PMCID: PMCPMC174230.

32.   Meri T, Hartmann A, Lenk D, Eck R, Würzner R, Hellwage J, et al. The yeast *Candida*

albicans binds complement regulators factor H and FHL-1. Infection and immunity.

2002;70(9):5185-92. doi: 10.1128/iai.70.9.5185-5192.2002. PubMed PMID: 12183569.

33.   Gonçalves B, Ferreira C, Alves CT, Henriques M, Azeredo J, Silva S. Vulvovaginal

candidiasis: Epidemiology, microbiology and risk factors. . Critical reviews in microbiology.

2016;42: 905-27.

34.   György C. Is there a hormonal regulation of phagocytosis at unicellular and

multicellular levels? A critical review. Acta Microbiologica et Immunologica Hungarica.

2017;64(4):357-72. doi: 10.1556/030.64.2017.024. PubMed PMID: 28859501.

35.   Murphy AJ, Guyre PM, Pioli PA. Estradiol suppresses NF-κB activation through

coordinated regulation of let-7a and miR-125b in primary human macrophages. The Journal

of Immunology. 2010;184(9):5029-37.

36.   Xing D, Oparil S, Yu H, Gong K, Feng W, Black J, et al. Estrogen modulates NFκB

signaling by enhancing IκBα levels and blocking p65 binding at the promoters of inflammatory

genes via estrogen receptor-β. PloS one. 2012;7(6):e36890.

37.   Brown GD, Gordon S. Immune recognition: A new receptor for [beta]-glucans. Nature.

2001;413(6851):36-7.

38.   Dasari P, Shopova IA, Stroe M, Wartenberg D, Martin-Dahse H, Beyersdorf N, et al.

Aspf2 From *Aspergillus fumigatus* Recruits Human Immune Regulators for Immune Evasion

and Cell Damage. Front Immunol. 2018;9:1635-. doi: 10.3389/fimmu.2018.01635. PubMed

PMID: 30166981.

39.   van der Maten E, van den Broek B, de Jonge MI, Rensen KJW, Eleveld MJ, Zomer

AL, et al. *Streptococcus pneumoniae* PspC Subgroup Prevalence in Invasive Disease and

Differences in Contribution to Complement Evasion. Infection and immunity.

2018;86(4):e00010-18. doi: 10.1128/IAI.00010-18. PubMed PMID: 29378798.

40.   Luo S, Hoffmann R, Skerka C, Zipfel PF. Glycerol-3-Phosphate Dehydrogenase 2 Is a

Novel Factor H–, Factor H–like Protein 1–, and Plasminogen-Binding Surface Protein of

Candida albicans. The Journal of Infectious Diseases. 2012;207(4):594-603. doi: 10.1093/infdis/jis718.

41. Marín E, Parra-Giraldo CM, Hernández-Haro C, Hernáez ML, Nombela C, Monteoliva L, et al. *Candida albicans* Shaving to Profile Human Serum Proteins on Hyphal Surface. Frontiers in microbiology. 2015;6:1343-. doi: 10.3389/fmicb.2015.01343. PubMed PMID: 26696967.

42. Luo S, Skerka C, Kurzai O, Zipfel PF. Complement and innate immune evasion strategies of the human pathogenic fungus Candida albicans. Molecular immunology. 2013;56(3):161-9.

43. Cheng G, Yeater KM, Hoyer LL. Cellular and molecular biology of *Candida albicans* estrogen response. Eukaryotic Cell. 2006;5(1):180-91. doi: 10.1128/ec.5.1.180-191.2006.

44. Monteiro PT, Oliveira J, Pais P, Antunes M, Palma M, Cavalheiro M, et al. YEASTRACT+: a portal for cross-species comparative genomics of transcription regulation in yeasts. Nucleic Acids Research. 2019;48(D1):D642-D9. doi: 10.1093/nar/gkz859.

45. Homann OR, Dea J, Noble SM, Johnson AD. A Phenotypic Profile of the *Candida albicans* Regulatory Network. PLoS Genet. 2009;5(12):e1000783. doi: 10.1371/journal.pgen.1000783.

46. Hao R, Bondesson M, Singh AV, Riu A, McCollum CW, Knudsen TB, et al. Identification of estrogen target genes during zebrafish embryonic development through transcriptomic analysis. PloS one. 2013;8(11):e79020-e. doi: 10.1371/journal.pone.0079020. PubMed PMID: 24223173.

47. Souder JP, Gorelick DA. Quantification of Estradiol Uptake in Zebrafish Embryos and Larvae. Toxicol Sci. 2017;158(2):465-74. doi: 10.1093/toxsci/kfx107. PubMed PMID: 28535311.

48. Lambris JD, Ricklin D, Geisbrecht BV. Complement evasion by human pathogens. Nature Reviews Microbiology. 2008;6(2):132-42. doi: 10.1038/nrmicro1824.

49.   Merle NS, Noe R, Halbwachs-Mecarelli L, Fremeaux-Bacchi V, Roumenina LT.

Complement System Part II: Role in Immunity. Front Immunol. 2015;6:257-. doi:

10.3389/fimmu.2015.00257. PubMed PMID: 26074922.

50.   Kenno S, Speth C, Rambach G, Binder U, Chatterjee S, Caramalho R, et al. *Candida*

albicans factor H binding molecule Hgt1p–a low glucose-induced transmembrane protein is

trafficked to the cell wall and impairs phagocytosis and killing by human neutrophils. Frontiers

in microbiology. 2018;9:3319.

51.   Luo S, Hartmann A, Dahse H-M, Skerka C, Zipfel PF. Secreted pH-regulated antigen

1 of *Candida albicans* blocks activation and conversion of complement C3. The Journal of

Immunology. 2010;185(4):2164-73.

52.   Poltermann S, Kunert A, von der Heide M, Eck R, Hartmann A, Zipfel PF. Gpm1p is a

factor H-, FHL-1-, and plasminogen-binding surface protein of *Candida albicans*. Journal of

biological chemistry. 2007;282(52):37537-44.

53.   Ngampasutadol J, Ram S, Gulati S, Agarwal S, Li C, Visintin A, et al. Human Factor H

Interacts Selectively with *Neisseria gonorrhoeae* and Results in Species-Specific Complement

Evasion. The Journal of Immunology. 2008;180(5):3426. doi: 10.4049/jimmunol.180.5.3426.

54.   Meri T, Amdahl H, Lehtinen MJ, Hyvärinen S, McDowell JV, Bhattacharjee A, et al.

Microbes Bind Complement Inhibitor Factor H via a Common Site. PLOS Pathogens.

2013;9(4):e1003308. doi: 10.1371/journal.ppat.1003308.

55.   Lee YJ, Jeschke GR, Roelants FM, Thorner J, Turk BE. Reciprocal phosphorylation of

yeast glycerol-3-phosphate dehydrogenases in adaptation to distinct types of stress.

Molecular and cellular biology. 2012;32(22):4705-17. Epub 2012/09/17. doi:

10.1128/MCB.00897-12. PubMed PMID: 22988299.

56.   Buckman J, Miller SM. Binding and reactivity of *Candida albicans* estrogen binding

protein with steroid and other substrates. Biochemistry. 1998;37(40):14326-36. Epub

1998/10/07. doi: 10.1021/bi981106y. PubMed PMID: 9760270.

57.   Zhang S, Cui P. Complement system in zebrafish. Developmental & Comparative

Immunology. 2014;46(1):3-10. doi: https://doi.org/10.1016/j.dci.2014.01.010.

58.   Sun G, Li H, Wang Y, Zhang B, Zhang S. Zebrafish complement factor H and its related

genes: Identification, evolution, and expression. Functional and Integrative Genomics.

2010;10(4):577-87. doi: 10.1007/s10142-010-0182-3.

59.   Cottier F, Sherrington S, Cockerill S, del Olmo Toledo V, Kissane S, Tournu H, et al.

Remasking of Candida albicans β-Glucan in Response to Environmental pH Is Regulated by

Quorum Sensing. mBio. 2019;10(5):e02347-19.

60.   Dennison PMJ, Ramsdale M, Manson CL, Brown AJP. Gene disruption in *Candida*

albicans using a synthetic, codon-optimised Cre-loxP system. Fungal Genetics and Biology.

2005;42(9):737-48. doi: https://doi.org/10.1016/j.fgb.2005.05.006.

61.   Walther A, Wendland J. An improved transformation protocol for the human fungal

pathogen *Candida albicans*. Curr Genetics. 2003;42(6):339-43.

62.   Barkani AE, Kurzai O, Fonzi WA, Porta RA, Frosch M, Muhlschlegel FA. Dominant

Active Alleles of *RIM101 (PRR2)* Bypass the pH Restriction on Filamentation of *Candida*

albicans. Mol and Cell Biol. 2000;20(13):4635-47.

63.   Wilson RB, Davis D, Enloe BM, Mitchell AP. A recyclable *Candida albicans URA3*

cassette for PCR product-directed gene disruptions. Yeast. 2000;16(1):65-70. doi:

10.1002/(sici)1097-0061(20000115)16:1<65::aid-yea508>3.0.co;2-m.

64.   Inglis DO, Arnaud MB, Binkley J, Shah P, Skrzypek MS, Wymore F, et al. The *Candida*

genome database incorporates multiple *Candida* species: multispecies search and analysis

tools with curated gene and protein information for *Candida albicans* and *Candida glabrata*.

Nucleic acids research. 2012;40(Database issue):D667-D74. Epub 2011/11/07. doi:

10.1093/nar/gkr945. PubMed PMID: 22064862.

65.   Mallick EM, Bergeron AC, Jones SK, Newman ZR, Brothers KM, Creton R, et al.

Phenotypic Plasticity Regulates *Candida albicans* Interactions and Virulence in the Vertebrate

Host. Frontiers in Microbiology. 2016;7(780). doi: 10.3389/fmicb.2016.00780.

66.   Gillum AM, Tsay EYH, Kirsch DR. Isolation of the *Candida albicans* gene for orotidine-

5′-phosphate decarboxylase by complementation of *S. cerevisiae ura3* and *E. coli* pyrF

mutations. Mol and Gen Genet. 1984;198(1):179-82.

67.   Noble SM, French S, Kohn LA, Chen V, Johnson AD. Systematic screens of a *Candida*

albicans homozygous deletion library decouple morphogenetic switching and pathogenicity.

Nature genetics. 2010;42(7):590-8. Epub 2010/06/13. doi: 10.1038/ng.605. PubMed PMID:

20543849.

68.   Fonzi WA, Irwin MY. Isogenic Strain Construction and Gene Mapping in *Candida*

albicans. Genetics. 1993;134(3):717-28. PubMed PMID: PMC1205510.

69.   Hall RA, De Sordi L, MacCallum DM, Topal H, Eaton R, Bloor JW, et al. CO2 acts as

a signalling molecule in populations of the fungal pathogen *Candida albicans*. PLoS Pathog.

2010;6(11):e1001193.

70.   Sanglard D, Ischer F, Monod M, Bille J. Cloning of *Candida albicans* genes conferring

resistance to azole antifungal agents: characterization of *CDR2*, a new multidrug ABC

transporter gene. Microbiology. 1997;143 (Pt 2):405-16. Epub 1997/02/01. doi:

10.1099/00221287-143-2-405. PubMed PMID: 9043118.

927 **Supporting information**

A

B

C

D

E

F

G

H

S1 Fig Oestrogen inhibits *C. albicans* phagocytosis, but does not affect fungal growth

or morphology. A) J774A.1 phagocytosis rates of *C. albicans* (SC5314) grown in the

presence of estriol. **B)** J774A.1 phagocytosis rates of *C. albicans* grown in the presence of

17α-ethynylestradiol. **C)** *C. albicans* cells were grown in 96-welled plate in YPD broth

supplemented with 10 µM 17β-estradiol, 10 µM 17α-ethynylestradiol or 10 µM estriol. Optical

densities (OD) of the cultures were recorded every 30 min for 18 h. **D)** *C. albicans* was grown

in YPD broth supplemented with 17β-estradiol 10 µM for 4 h. Cells were washed with PBS,

stained with concanavalin A, imaged by microscopy and analysed for shape. **E)** J774A.1

macrophages preincubated with 10 µM 17β-estradiol were infected with *C. albicans* cells

grown in YPD and phagocytosis rates quantified. **F)** Beads were incubated at room

temperature in PBS with or without 10 µM 17β-estradiol. Beads were washed and co-

incubated with J774A.1 macrophages and phagocytosis rates quantified. **G)** *C. albicans* cells

were incubated in PBS at 4°C for 4 h with or without 10 µM 17β-estradiol. Cells were washed

and co-incubated with J774A.1 macrophages and phagocytosis rates quantified **H)** *C. albicans*

cells previously grown in YPD with or without 10 µM 17β-estradiol were harvested, washed

and re-incubated in fresh YPD for 24 h. Cells were co-incubated with J774A.1 macrophages

and phagocytosis rates quantified. All data represent the mean ± SEM from at least three

independent experiments. (**$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$).

S2 Fig Oestrogen induces a mild transcriptional response in *C. albicans* A) *C. albicans*

(SC5314) was grown with or without 10 μM 17β-estradiol for the indicated time and then

exposed to J774A.1 macrophages for 45 minutes and phagocytosis rates quantified. **B)** PCA

of RNA Seq data. **C)** The *cdr1Δ*, *cdr1Δ/cdr2Δ* mutants and the parental control strain were

grown with or without 10 μM 17β-estradiol for 4 hr, and then exposed to J774A.1 macrophages

for 45 minutes and phagocytosis rates quantified. All data represent the mean ± SEM from at

least three independent experiments. (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$).

S3 Fig Bcr1 and Rob1 regulate the expression of *GPD2*. Transcription factor mutants and

the parental control strain were grown in YPD to mid-log phase, total RNA extracted and *GPD2*

gene expression quantified by RT-PCR relative to *ACT1*.

975 **S1 Table *Candida* strains used in this study**

Strain	Genotype	Source/reference
SC5314	Wild type	[66]
SVS006B	Clinical isolate	Prof Ramage, Glasgow University
SVS062A	Clinical isolate	Prof Ramage, Glasgow University
SN152	*arg4Δ/arg4Δ leu2Δ/leu2Δ his1Δ/his1Δ URA3/ura3Δ:: λimm434 IRO1/iro1Δ:: λimm434*	[67]
SN250	*his1Δ/his1Δ, leu2Δ::C. dubliniensis HIS1 /leu2Δ::C. maltosa LEU2-arg4Δ/arg4Δ, URA3/ura3Δ::imm434-IRO1/iro1Δ::imm434*	[67]
SN250-CIP30	As SN250 but *RPS1/rps1::*CIP30	This study
rob1Δ	As SN152 but *rob1Δ::C. dubliniensisHIS1/rob1Δ::C. maltose LEU2*	[67]
bcr1Δ	*arg4Δ/arg4Δ leu2Δ/leu2Δ his1Δ/his1Δ URA3/ura3Δ::λimm434 IRO1/iro1Δ:: λimm434 bcr1::LEU/bcr1::HIS1*	[67]
gpd2Δ	As SN152 but *gpd2Δ::C. dubliniensisHIS1/gpd2Δ::C. maltose LEU2*	[67]
gpd2Δ–CIP30	As SN152 but *gpd2Δ::C. dubliniensisHIS1/gpd2Δ::C. maltose LEU2, RPS1/rps1::CIP30*	This study
gpd2Δ-CIP30-GPD2	As SN152 but *gpd2Δ::C. dubliniensisHIS1/gpd2Δ::C. maltose LEU2RPS1/rps1::CIP30-GPD2*	This study
CAI4	*ura3::imm434/ura3::imm434 iro1/iro1::imm434*	[68]
CAI4-pSM2	*ura3::imm434/ura3::imm434 iro1/iro1::imm434::pSM2*	[69]

CAI4-pSM2- *GPD2*	*ura3::imm434/ura3::imm434 iro1/iro1::imm434::pSM2- pTef2-GPD2*	This study
ebp1Δ	As CAI4 but *ebp1::*dlp200/*ebp1::*dlp200	This study
ebp1Δ-pSM2	As CAI4 but *ebp1::*dlp200/*ebp1::*dlp200-pSM2	This study
CAF2-1	*URA3/ura3:: λimm434*	[68]
cdr1Δ	As CAF2-1, but *crd1::hisG/cdr1:hisG-URA3-HisG*	[70]
cdr1/2Δ	As CAF2-1, but *crd1::hisg/cdr1:hisG, cdr2::hisG-URA3- hisG/cdr2::hisG*	[70]

978 **S2 Table Primers used in this study**

Primer	Sequence
GPD2-SacI-F	CTCCGAGCTCGGTGATGGTGATGGTGATGG
GPD2-NotI-R	GGAGAGCGGCCGCTGGTAAATTGGACAACGAGTGG
EBP1-5F:	CTCCGATATCATCGCATGAG
EBP1-5R:	GGAGAGGAGCTCctgatgatatgataatttgc
EBP1-3F:	GGAGAGAAGCTTGGGAATGAAGTTCTATTAGC
EBP1-3R:	GGAGAGGGTACCgttacatctactactacagg
RT-ACT1-F	CCTACGTGTACTTGTGCAAGGCAA
RT-ACT1-R	TAGTTGTGTGCACTGAGCGTCGAA
RT-GDP2-F	GCCAACGAAGTTGCCAAAGGT
RT-GPD2-R	AGGCACCAGCAATAGAGGCA