Prion cross-seeding between fungal and mammalian amyloid signaling motifs

Thierry Bardin[1], Asen Daskalov[1,], Sophie Barrouilhet[1#a], Alexandra Granger-Farbos[1],

Bénédicte Salin[1], Corinne Blancard[1], Sven J. Saupe[1*] and Virginie Coustou[1*]

[1] Non-Self Recognition in Fungi, Institut de Biochimie et de Génétique Cellulaire (CNRS UMR

5095, Université de Bordeaux), 33077 Bordeaux, France.

[#a] Current Address: Universite de Pau et des Pays de l'Adour, E2S UPPA, CNRS, IPREM UMR

5254, Hélioparc, 64053 Pau, France

*Corresponding authors

E-mail : virginie.coustou@ibgc.cnrs.fr (VC); sven.saupe@ibgc.cnrs.fr (SJS)

Running title: Amyloid signaling pathways in *P. anserina*

Key words: amyloid; prion; programmed cell death; RHIM; RIP1, RIP3

Abstract

In fungi, NLR-based signalosomes activate downstream membrane-targeting cell-death inducing proteins by a mechanism of amyloid templating. In the species *Podospora anserina*, two such signalosomes, NWD2/HET-S and FNT1/HELLF have been described as well as an analogous system, involving a distinct amyloid signaling motif termed PP in the species *Chaetomium globosum*. The PP-motif bears resemblance to the RHIM and RHIM-like motifs controlling necroptosis in mammals and innate immunity in Drosophila. We identified here, a third NLR signalosome in *Podospora anserina* comprising a PP-motif and organized as a two-gene cluster encoding a NLR and a HELL-domain cell-death execution protein termed HELLP. We show that the PP-motif region of HELLP forms a prions we term [π] and that [π] prions trigger the cell-death inducing activity of full length HELLP. We detect no prion cross-seeding between HET-S, HELLF and HELLP amyloid motifs. In addition, we find that akin to PP-motifs, RHIM motifs from human RIP1 and RIP3 kinases are able to form prions in Podospora, and that [π] and [Rhim] prions cross-seed. Our study shows Podospora display three independent amyloid signalosomes. Based on the described functional similarity between RHIM and PP it appears likely that these amyloid motifs constitute evolutionary related cell-death signaling modules.

Author summary

Amyloids are fibrillar protein polymers assembled from cross β-sheets. Pathogenic amyloids are responsible for a range of neuronal and systemic protein misfolding diseases. In addition, functional amyloids fulfill a variety of biological functions in microbes and animals. In fungi, amyloids control prion-like signaling pathways with roles in host defense and immune

programmed cell death. These signaling pathways involve a Nod-like receptor and a cell-death

execution protein activated through amyloid templating. In *P. anserina* two such amyloid

signaling systems were identified so far. Here we identified and characterized a third amyloid

signaling system comprising a different amyloid motif termed PP. We showed a total absence

of cross-induction between the three *P. anserina* systems allowing for the co-existence of

three independent amyloid signaling cascades in the same organism. An evolutionary

relationship between fungal amyloid motifs and mammalian RHIM amyloid motif involved in

the necroptosis pathway was previously suggested. We validated this assumption by showing

that the human RHIM motif could not only propagate as a prion in *P. anserina* but also

functionally cross-interact with the fungal PP-motif prions. These results point to a long term

evolutionary conservation of amyloid signaling mechanisms from fungi to mammals.

Introduction

Prions are protein polymers that behave as infectious entities by propagating their amyloid structural state [1,2]. In addition to disease causing prions in mammals, prions have been identified in fungi as infectious, cytoplasmic, non-Mendelian genetic elements [3]. One such prion is [Het-s] of the filamentous fungus *Podospora anserina.* [Het-s] functions in a non-self recognition process known as heterokaryon incompatibility. The *het-s* gene exists as two incompatible allelic variants termed *het-s* and *het-S*. The HET-s protein when assembled into the self-propagating prion form confers incompatibility to the HET-S variant, meaning that co-expression of the [Het-s] prion with HET-S leads to cell death. The soluble non-prion form confers a phenotype termed [Het-s*] compatible with HET-S [4,5]. [Het-s*] strains acquire the prion state either spontaneously at a low frequency or systematically after fusion with a prion infected strain. HET-s and HET-S are highly homologous allelic variants that display two domains, a C-terminal Prion-Forming Domain (PFD) necessary and sufficient for prion propagation and a N-terminal α-helical globular domain named HeLo responsible for cell-death inducing activity [6,7]. The HET-s PFD is natively unfolded in the soluble state and upon prion formation assembles into to a specific cross-β amyloid structure [6,8–10]. The HET-s β-solenoid fold is composed of 2 rungs of β-strands per monomer, comprising two 21 residues imperfect repeats (R1 and R2) connected by a flexible loop [9,11]. In [Het-s]/HET-S incompatibility cell death is triggered when the [Het-s] PFD templates conversion of the HET-S PFD region into the β-solenoid fold which in turn induces refolding of the HET-S HeLo domain that acquires pore-forming activity by exposing an N-terminal transmembrane helix targeting the cell membrane [7,12]. In this incompatibility system, HET-S acts as cell-death execution protein and [Het-s] triggers its activation.

Further work on the [Het-s] system revealed that this incompatibility system is

evolutionary derived from a programmed cell death pathway in which HET-S is activated by a

specific NLR receptor [13]. NLR receptors are intracellular receptors that control immune

defense and programmed cell death pathways in animals and plants and also in fungi and

function by ligand induced oligomerization [14,15]. The NWD2 NLR receptor is encoded by the

gene immediately adjacent to *het-S* in the *P. anserina* genome and displays a central NACHT

nucleotide binding and oligomerization domain and a C-terminal WD40 repeat domain. The

N-terminal region of NWD2 displays a short region of homology to the R1 and R2 motifs of

HET-s. This region designated R0 is able to adopt a *het-s*-like fold. Ligand induced

oligomerization of NWD2 allows for spatial clustering of R0 motifs, cooperative nucleation of

the β-solenoid fold templating activation of the HET-S HeLo domain [13,16,17]. More

generally, it is proposed that a fraction of the NLR receptors in fungi activate their cognate

effector domains by a mechanism of prion amyloid templating by which the amyloid fold

formed after oligomerization of the NLR receptor is transmitted through heterotypic

interactions to the amyloid motif found in the C-terminal of the effector protein [18].

NWD2/HET-S pair was used as a paradigm for extensive bioinformatic screening to identify

additional NLR/effector pairs functioning through amyloid templating in fungi [16,19]. This

search resulted in the identification in *Podospora anserina* of a distant homolog of HET-S

termed HELLF [20]. HELLF encodes a protein with a HeLo-like N-terminal domain and a C-

terminal PFD with two repeated submotifs that share less than 15% identity to the HET-s PFD.

The HeLo-like domain shows distant homology to the HET-S HeLo domain but also to the 4HB

pore-forming domain of the MLKL protein [21], the terminal effector protein of necroptotic

death in mammals, which led to the proposition that this form of fungal programmed cell

death is related to mammalian necroptosis [22]. The gene adjacent to *hellf* encodes a NLR with

a NB-ARC domain and TPR repeats termed FNT1, which bears a N-terminal R0 region

homologous to the HELLF R1 and R2 PFD repeats. HELLF behaves analogously to HET-S in every

aspect that was analyzed. The C-terminal region of HELLF forms a prion termed [Φ] and [Φ] is

incompatible with full length HELLF. Upon interaction with [Φ], HELLF relocates to cell

membrane region and causes cell death and the N-terminal region of FNT1 forms fibrils and is

able to induce [Φ] prion formation. The solid-state NMR structure of the HELLF PFD structure

revealed that HET-s and HELLF prions have an almost identical backbone structure in spite of

extensive sequence divergence. The FNT1/HELLF system thus constitutes a second cell death-

inducing amyloid signaling pathway of *Podospora anserina*. Noteworthy is the fact that in spite

of their almost identical backbone structure [Het-s] and [Φ] prions do not cross-seed

indicating that NWD2/HET-S and FNT1/HELLF constitute two parallel signaling pathways.

In bioinformatics surveys, additional amyloid signaling motifs were identified in fungi

[14,16]. In particular, a prion motif termed PP (and also associated to NLR signaling) was

characterized in *Chaetomium globosum* in a three-gene cluster. In this case, the NLR protein

PNT1 regulates activation of two effector proteins, a HeLo-like domain cell death inducing

protein, HELLP (herein renamed CgHELLP for clarity) and a putative lipase, SBP. Remarkably,

the PP-motif is related in sequence to the RHIM amyloid motif that controls assembly of the

RIP1 and RIP3 kinases in the necroptosis pathway in mammals [13,22,23]. RHIM-like amyloid

motifs also occurs in the PGRP-LC, PGRP-LE, and Imd proteins controlling anti-bacterial

immunity in Drosophila [24]. The structure of the mammalian RIP1-RIP3 core (forming an

hetero-amyloid signaling complex) has recently been established [25]. RHIM amyloids

comprise two protein units per cross-β layer that run antiparallel and are interdigitated in a

compact hydrophobic interface formed by the G-(I,V)-Q-(I,V)-G motif. This central core motif

is common to the PP-motif but structural characterization of PP-amyloids will be required to

determine whether this short sequence homology reflect structural similarity between RHIM

and PP-amyloids.

Herein, we mine the genome of *P. anserina* and identify HELLP, a novel protein PP-motif

protein that defines a third cell death inducing amyloid signaling pathway in that species. We

show that the C-terminal PP-motif of HELLP forms a prion termed [π] in *Podospora anserina*

and assembles into fibrils *in vitro.* We find that akin to HET-S and HELLF, membrane-targeting

cell death inducing activity of HELLP can be triggered by [π] prions or by the prion form of the

N-terminal region of the HELLP-associated NLR, termed PNT1. We analyze in *Podospora*

anserina the functional interactions of the HET-s, HELLF and HELLP PFD prions and find that

they form three independent signaling systems. We further analyze prion formation by

mammalian RHIM motifs and find that [Rhim] and [π] prions partially cross-seed underlying

the functional similarity between these mammalian and fungal amyloid motifs.

Results

A PP-motif gene cluster in *Podospora anserina*

During the course of a survey of NLR encoding genes in fungi, we identified, in *Podospora*

anserina a gene (*Pa_5_8060*) encoding a NLR with a NB-ARC domain and TPR repeats (we term

PNT1) [14]. We found that the adjacent gene (*Pa_5_8070*) encodes a HeLo-like-domain (HELL)

protein, we term HELLP (Fig 1A). HELLP shows homology to the CgHELLP cell-death execution

protein of *Chaetomium globosum* (Fig 1B and S1 Fig) [22]. As previously reported for CgHELLP

and HELLF, the HELL domain of HELLP shows homology to the 4HB membrane-targeting

domain of the mammalian MLKL necroptosis execution protein (S1 Fig). The N-terminal region

of PNT1 and the C-terminal region of HELLP share a region of homology encompassing a

predicted PP amyloid motif (Fig 1B) [22]. This region is also homologous to the RHIM motif

found in the RIP1 and RIP3 kinase in humans [23,25]. Based on the homology between HELLP

and CgHELLP, we reasoned that analogously to *nwd2/het-s* and *fnt1/hellf*, the *pnt1/hellp* gene

pair might encode the components of a third amyloid NLR signalosome in Podospora (Fig 1A).

We thus engaged into the functional characterization of this gene cluster.

The C-terminal region of HELLP is a prion-forming domain

In order to analyze whether the C-terminal region of HELLP displays the anticipated prion

forming ability, we generated different constructs with the C-terminal domain of HELLP (214-

271) fused either to GFP or RFP (in a C-terminal position) or to GFP in an N-terminal position.

These constructs were expressed in a *Δhellp* strain of *P. anserina.* For all constructs, the

fluorescence signal was initially diffuse but cytoplasmic foci formed upon further growth

(Table 1 and Fig 2A). The number of transformants spontaneously exhibiting the foci

phenotype was determined after 5, 11 and 19 days of growth (Table 1). The highest rate of

spontaneous foci formation was observed for the fusion protein bearing the GFP in N-terminal

position. For all fusion constructs, strains with diffuse fluorescence were systematically

converted to the foci phenotype after cytoplasmic contact with strains expressing foci,

indicating that the foci state is transmitted by cytoplasmic contact and infectious. By analogy

with the [Het-s] system, we term the phenotypic state with diffuse fluorescence [π*] and the

foci state [π] (Fig 2A). We conclude from these experiments that the PP-motif region of

HELLP(214-271) allows for prion formation and propagation. As previously described for [Het-

s] [26], the [π] state could be reversed to [π*] in meiotic progeny. When a [π] strain is used as

male parent, the meiotic progeny display the [π*] phenotype.

The same experiments were also carried out with a larger construct HELLP(171-271) fused

to GFP either in N- or C-terminus (S2A Fig). The larger construct also led to foci formation but

in contrast to analogous HET-S and CgHELLP constructs (HET-S(157-289)-GFP and

CgHELLP(170-278)-GFP), HELLP(171-271) fusions did not form elongated aggregates [22,27].

HELLP(214-271) PFD region was expressed in *E. coli* with a C-terminal histidine tag and

purified under denaturing conditions. Upon removal of the denaturant by dialysis, the protein

spontaneously formed fibrils that most often associated laterally as bundles (Fig 2B). These

results indicate that the HELLP(214-271) PFD region is able to form fibrils *in vitro.*

Table 1. Rate of spontaneous prion formation and propagation

transgene(s)	days after transfection								
	5			11			19		
	Soluble	Aggregated		Soluble	Aggregated		Soluble	Aggregated	
	n	n	%	n	n	%	n	n	%
HELLP(214-271)-RFP	27	7	20.6	13	21	61.8	2	32	94.1
HELLP(214-271)-GFP	40	7	14.9	24	23	48.9	8	39	83.0
GFP-HELLP(214-271)	11	22	66.7	1	32	97.0	0	33	100.0
RIP3(444-469)-RFP	18	21	53.8	8	31	79.5	2	37	94.9
RIP3(444-469)-GFP	30	16	34.8	15	31	67.4	4	42	91.3
GFP-RIP3(524-551)	8	17	68.0	3	22	88.0	0	25	100.0
RIP1(524-551)-RFP	23	3	11.5	14	12	46.2	10	16	61.5
RIP1(524-551)-GFP	27	0	0.0	25	2	7.4	23	4	14.8
GFP-RIP1(524-551)	38	0	0.0	36	2	5.3	35	3	7.9

HELLP relocates to the cell membrane and triggers cell death upon interaction with [π] prions

It was shown that HET-S, HELLF and CgHELLP cell-death inducing activity is induced by interaction with the prion form of their respective PFDs [7,12,20,22]. As a result confrontation of strains expressing PFDs in the prion state with strains expressing the corresponding full length protein leads to an incompatibility reaction which results in the death of the fusion cells [20,22,27]. This reaction manifests through the formation of a demarcation line visible at the macroscopic level and termed "barrage". To determine whether HELLP could also lead to incompatibility, strains expressing GFP-HELLP(214-271) were confronted to strains expressing full length HELLP, either from the wild-type resident copy or from HELLP-GFP or HELLP-RFP transgene copies in a *Δhellp* background (Fig 3A). A barrage reaction was observed specifically in confrontations with strains expressing the [π] phenotype. [π*] strains did not produce a barrage reaction. The barrage reaction was stronger with strains expressing HELLP-GFP or -RFP fusions from the strong *gpd* promoter than with strains expressing HELLP from the resident copy (Fig 3A).

Propagation of the [π] phenotype was then assayed using barrage test as typically done for monitoring [Het-s] propagation. Upon previous contact with [π] strains, [π*] strains became capable of forming a barrage with strains expressing full length HELLP (S1 Table and S2 Fig). Based on these experiments the definition of the [π] phenotype can be expanded to represent both the foci state of the PP-motif GFP and RFP fusions and the ability to produce a barrage reaction to full-length HELLP. The [π*]/[π] phenotypes of HELLP(214-271) are in that sense analogous to [Het-s*]/[Het-s] phenotypes of HET-s and [Φ*]/[Φ] phenotypes of HELLF(209-277).

It was shown that prion conversion of the HET-S PFD leads to the refolding of the HET-S

HeLo domain that acquires pore-forming activity by exposing its N-terminal transmembrane

helix, relocating it to the cell membrane and inducing cell death [7,12]. The same relocalization

process was observed for HELLF and CgHELLP in the incompatibility reaction leading to cell

death [20,22]. We thus determined whether HELLP relocates to the cell membrane upon

interaction with [π] prions. We examined the contact zone between incompatible [π] and

HELLP-GFP or -RFP expressing strains (Fig 3B). While full length HELLP-GFP or -RFP alone

produced diffuse fluorescence signals that remain stable over time (S3 Fig), in fusion cells co-

expressing these proteins and the prion form of HELLP(214-271)-RFP or GFP-HELLP(214-271),

we observed a localization of HELLP at the cell membrane region. This localization is associated

to cell death as observed using the vital dye Evans Blue (Fig 3C). These observations indicate

that HELLP functions analogously to HET-S, HELLF and CgHELLP and that the HELL membrane-

targeting domain is regulated by prion conversion of the C-terminal PFD region.

218   The PNT1(1-31) region forms a prion and induces HELLP activation

It has been shown that the N-terminal region of NWD2 adopts an amyloid fold, displays prion

infectivity and is able to activate HET-S pore forming activity [13]. Similarly, N-terminal regions

of the FNT1 and CgPNT1 NLRs display prion forming activity and trigger the cell-death inducing

activity of their effectors proteins, HELLF and CgHELLP respectively [22]. To determine

whether the N-terminal region of PNT1 displays similar properties we expressed PNT1(1-31)

fused to GFP or RFP in *Δhellp* strains. Two phenotypic states were observed, a diffuse

cytoplasmic fluorescence state, and foci state comparable to the [π*] and [π] phenotypes

observed previously with HELLP(214-271) (Fig 4A). Spontaneous conversion to the aggregated

form occurred after few days and cytoplasmic contact with [π] strains induced systematic

conversion.

PNT1(1-31)-GFP [π] strains systematically convert [π*] HELLP(214-271)-RFP strains to the

[π] state. Conversely, [π] HELLP(214-271)-RFP systematically convert [π*] PNT1(1-31)-GFP to

the [π] state.

PNT1(1-31) and HELLP(214-271) were then co-expressed in the same strain (Fig 4B).

Diffuse fluorescence was observed very briefly in the case of HELLP(214-271)-RFP and PNT1(1-

31)-GFP co-expression, indicating that these two proteins were in a soluble state just after

transfection. Then, dot-like aggregates formed concomitantly for the two proteins and co-

localized in the cell, as expected if they could interact and cross-seed *in vivo*. The same was

true during co-expression of PNT1(1-31)-RFP and GFP-HELLP(214-271). An initial diffuse state

could however not be observed in this setting presumably as the result of the high

spontaneous conversion rate of the N-terminal GFP-HELLP(214-271) fusion (Table 1).

Finally, we tested the ability of the PNT1(1-31) to activate the HELLP effector protein. We

observed fusion cells between strains expressing PNT1(1-31) in the aggregated [π] state and

strains expressing full length HELLP (Fig 4C and 4D). We observed the localization of HELLP to

the plasma membrane region and cell death of the fusion cells. Thus, consistent with the

proposed role of HELLP as cell death execution protein activated by the PNT1 NLR, we find

that the PP-motif region of PNT1 forms a prion and is able to activate HELLP.

HET-S, HELLF and HELLP are components of three independent cell-death

inducing pathways

We previously reported a complete lack of cross-seeding between [Het-s] and [Φ] prions

despite identical backbone structures leading to the existence of two independent parallel

amyloid signaling pathways in *P. anserina* [20]. Here, we identified HELLP as an effector in a

third amyloid signaling pathway in *P. anserina*. To analyze HELLP/HELLF and HELLP/HET-s

interactions, we co-expressed HELLP(214-271) and HET-s or HELLF(209-271) respectively in a

ΔhellpΔhet-sΔhellf strain. We observed the diffuse and foci states co-existing in all possible

combinations (Fig 5 and S4). Namely strains expressing HELLP(214-271) and HET-s could

display four different phenotypes : [Het-s]/[π] and [Het-s*]/[π*] but also [Het-s]/[π*] and [Het-

s*]/[π]. The same was true for HELLP(214-271) and HELLF(209-277 co-expression. These

results show that [Φ] or [Het-s] prions do not to convert [π*] and conversely that [π] prions

do not to convert [Φ*] or [Het-s*]. We also noted an absence of co-localization between

HELLP(214-271) and HET-s or HELLP(214-271) and HELLF(209-277) prion forms. Together,

these results demonstrated a lack of *in vivo* cross-seeding between the three *P. anserina* prion

forming domains.

We also monitored [π*] conversion by [Het-s] and [Φ] prions through barrage tests (S2

Table). Strains expressing HELLP(214-271)-GFP or -RFP and displaying [π*] phenotype were

confronted to strains expressing [π], [Φ] or [Het-s] prion forms and then tested for their ability

to form a barrage with tester strains expressing full length HELLP. Barrage were observed only

when the prion donor strain express [π] prions indicating that [π*] conversion do not occur in

presence of [Het-s] or [Φ] prions.

Finally, we analyzed the cross-activation of the cell death effectors of the three systems

by the different prions (S3 Table). Again, we used incompatibility assays and confronted [π],

[Φ] or [Het-s] prions expressing strains with strains expressing either HELLP, HELLF or HET-S

full-length proteins. We observed barrage formation solely when [π] strains were confronted

with HELLP expressing strains, [Φ] with HELLF and [Het-s] with HET-S, that is, when the prion

and the effector bear the same PFD. These results confirm that there is no cross-interaction

between these three amyloid signaling pathways.

277 *P. anserina* and *C. globosum* [π] prions cross-seed and co-localize

It was shown previously that HET-S PFDs homologs from *P. anserina* and *Fusarium*

graminearum which share about 38% identity do cross-seed [28]. We wondered whether the

45% similarity in the PFD region between HELLP and CgHELLP from *Chaetomium globosum*

would also allow for cross-seeding. To examine if the two PFD regions could co-aggregate *in*

vivo, strains co-expressing HELLP(214-271)-RFP and GFP-CgHELLP(215-278) were obtained

and analyzed by fluorescence microscopy (Fig 6A). As expected, we initially observed diffuse

fluorescence corresponding to the [π*] state and after a few days, foci appearing

concomitantly for the two proteins. There was no co-existence of soluble and aggregated

forms suggesting cross-conversion of HELLP and CgHELLP. Moreover, we observed a

significant co-localization indicating co-aggregation of the two prions *in vivo*. Upon

confrontation, [π] GFP-HELLP(214-271) donor converted GFP-CgHELLP(215-278) [π*] recipient

strains and conversely [π] CgHELLP(215-278) strains converted GFP-HELLP(214-271) [π*]

recipients (S4 Table).

We also used barrage assays to analyze the cross-induction of cell death activity of the

full-length CgHELLP and HELLP proteins (Fig 6B and S5 Table). GFP-CgHELLP(215-278) [π]

strains produced a barrage reaction to strain expressing HELLP and conversely GFP-HELLP(214-

271) [π] produced a barrage reaction to strain expressing CgHELLP. However, this was only

true when HELLP was highly expressed as a transgene. No barrage was observed in

confrontation to wild type. This result suggest that heterologous activation of HELLP by GFP-

CgHELLP(215-278) is less efficient than homotypic activation by GFP-HELLP(214-271). We

conclude that the "species-barrier" for prion seeding can be crossed for these PP-motifs but

that nevertheless homotypic interaction appear more efficient to activate the cell death

activity of the full-length protein.

The RHIM regions of human RIP3 and RIP1 kinases form prions in *P. anserina*

The PP-motif as found in HELLP shows sequence homology to the mammalian RHIM amyloid

motif. We thus envisioned that akin to the PP-motif, mammalian RHIMs might behave as PFDs

when expressed in Podospora. We expressed the RIP3(444-469) and the RIP1(524-551)

regions fused to GFP or RFP in *ΔhellpΔhet-sΔhellf* strains (Fig 7). Initially fluorescence was

diffuse and cytoplasmic and foci appeared spontaneously after few days of growth. The rate

of spontaneous transition to the aggregated state was monitored as previously for HELLP

(Table 1). For RIP3(444-469), the rates were comparable to the PP-motif with the highest rate

observed when GFP is in N-terminal position. For the RIP1 RHIM motif, the spontaneous rate

of foci formation was much lower especially for the two GFP fusions. However, this difference

relates only to spontaneous foci formation as the induced conversion through cytoplasmic

infection (with the corresponding foci form) is equally effective for the RIP1 and RIP3 motifs

(Fig 8). In addition, strains expressing RIP1(524-551) foci induced formation of RIP3(444-469)

foci and the converse was true. To test the ability of the [Rhim] prions observed in *P. anserina*

to interact, we co-expressed RIP1(524-551) and RIP3(444-469) fused to GFP or RFP in the same

strain (S5 Fig). Diffuse fluorescence was observed for several days during which these two

proteins were in the soluble [Rhim*] state. Then, foci formed concomitantly and co-localized

in the cell, often in the close vicinity of the septa where they appeared as aligned dots. The

concomitant emergence of RIP1 and RIP3 aggregates and their co-localization showed that

the [Rhim] prions co-aggregate and confirm RIP1(524-551) and RIP3(444-469) cross-seeding

in vivo in *P. anserina*.

We conclude that, the RHIM containing RIP3(444-469) and RIP1(524-551) regions behave

as PFDs *in vivo* in *P. anserina* and lead to the expression of two alternate phenotypes we

termed [Rhim*] and [Rhim]. In addition, the RIP1 and RIP3 [Rhim] prions cross-seed and co-

localize consistent with the fact that in mammalian cells and *in vitro*, RIP1 and RIP3 interact

through their RHIM regions to form amyloid heteropolymers [25].

329 The [Rhim] and [π] prions partially cross-seed

To examine possible interaction between [Rhim] and [π] prions, we analyzed the ability of the

prion containing strains to cross convert prion-free strains. For this purpose, strains expressing

HELLP(214-271)-RFP [π*], GFP-RIP3(444-496) and GFP-RIP1(524-551) [Rhim*] strains were

confronted with [π] and [Rhim] strains (Fig 8). After four days, the recipient strains were

sampled and subcultured and analyzed for the presence of foci by fluorescence microscopy.

In these conditions spontaneous prion formation was in the range of 10 to 20% for all

constructs. Homotypic interactions led to 80 to 90% prion conversion. High conversion rates

were observed for heterotypic conversion of GFP-RIP3(444-496) [Rhim*] by GFP-RIP1(524-

551) [Rhim]; and GFP-RIP1(524-551) [Rhim*] by GFP-RIP3(444-496) [Rhim]). Importantly, [π]

prions induced formation of both [Rhim] prions at a rate of about 60%. Conversely, both

[Rhim] prions induced formation of [π] prions at a similar rate (Fig 8). The results clearly

showed the ability of cross conversion between [Rhim] and [π] prions. The conversion is

however less efficient as in the case of homotypic interactions.

We also tested the ability of the two [Rhim] prions to induce the conversion of [Het-

s*] strains to the [Het-s] phenotype and observed a total absence of induction of this

conversion. This result is in line with the previous results concerning the absence of cross-

interaction between [π] and the other amyloid signaling pathways of *P. anserina* and rather

expected.

To explore interactions between [π] and [Rhim] further, HELLP and RIP1 (or RIP3) PFDs

fused to GFP or RFP were coexpressed in the same strain and transformants were analyzed by

fluorescence microscopy (Fig 9 and S6). Just after transfection, we observed diffuse signals for

both proteins (top panels in Fig 9A-B and S6A-B). It is possible to observe transformants where

one of the protein has already shifted towards the aggregated form while the other remained

soluble, with for instance, the coexistence of [π*] and [Rhim] phenotypes (Fig 9A and S6A) or

of [π] and [Rhim*] phenotypes (Fig 9B and S6B). In the combination RIP3(444-469)/HELLP(214-

271), most often spontaneous conversion to the [Rhim] phenotype is observed first,

consistent with the higher aggregation rate measured for RIP3(444-469) (Table 1). On the

contrary, in the RIP1(524-551)/HELLP(214-271) combination, the [π] prion appeared first,

again consistent with the measured lower spontaneous conversion rate of RIP1(524-551). In

both cases, upon further subculture, we observed foci for the two proteins that partially co-

localized (Fig 9A-B and S6A-B, bottom lines). Most often, the majority of the foci co-localized

while few others corresponded only to GFP or RFP-only foci. In addition, even in mixed foci

the co-localization is not absolute with RFP and GFP fluorescence that only partially

overlapped. Strains co-expressing RIP3(444-469) fused with GFP and RFP, or RIP1(524-551)

fused with GFP and RFP were used as positive control (S6C Fig) and showed co-localization

with almost perfectly overlapping foci. This phenomenon of patchy co-localization has already

been observed during coexpression of the HET-s proteins of *F. graminearum* and *P. anserina*

[28]. The same experiments were also carried out with CgHELLP(215-278) and the same results

were obtained (S7 Fig). These results indicate partial co-localization of [Rhim] and [π] prions,

a situation that is distinct both from the total lack of interaction seen between [π] and [Het-s]

and [Φ] prions and from the strong interactions seen between RIP1 and RIP3 [Rhim] prions on

one hand and HELLP and CgHELLP [π] prions on the other hand.

[Rhim] prions induce toxicity of full length HELLP

Confrontation of [Rhim] strains and full-length HELLP expressing strains did not lead to

barrage formation but microscopic observation of the confrontation zone revealed cell death

in few mixed cells (S8 Fig). We thus co-expressed in the same strain GFP-RIP3(444-469) or GFP-

RIP1(524-551) and HELLP-RFP to determine whether [Rhim] prions could induce HELLP

toxicity. Fluorescence signals were diffuse and cytoplasmic for both of the proteins and

remained stable for weeks indicating that [Rhim*] state remained much more stable in these

transformants than in GFP-RIP3(444-469) single transformants showing high rates of

spontaneous prion acquisition. To force transition to the [Rhim] state, these transformants

were confronted with [Rhim] strains and subsequently subcultured. After contact, about 60%

of the GFP-RIP3(444-469) [Rhim]/HELLP transformants displayed altered phenotypes with

reduced growth, 20% displayed sub-lethal phenotypes and only 20% showed normal growth.

Similarly in the case of GFP-RIP1(524-551) [Rhim]/HELLP coexpression 56% of the

transformants displayed altered phenotypes, 25% sub-lethal phenotypes and 19% appeared

normal. A representative sample of the abnormal transformants is shown (Fig 10A).

Fluorescence microscopy on these abnormal strains (Fig 10B) revealed not only the presence

of [Rhim] dots but also HELLP relocalization to the membrane associated with cell death,

swollen cells that were associated with sub-lethal growth, and HELLP aggregation in dots that

co-localized with [Rhim] dots in strains with altered growth phenotypes. On the contrary, strains presenting normal growth rates either showed a [Rhim*] phenotype or showed an important loss of fluorescence intensity for one or both of the co-expressed proteins. In the same way, upon further sub-culturing transformants presenting growth defects often "escaped" and presented normal growth associated with loss of fluorescence levels or of the [Rhim] phenotype. These results suggest that [Rhim] prions are able to interact with HELLP and to trigger trans-conformation of the full-length protein leading to cell membrane relocalization and cell death. These results are in agreement with the partial [Rhim] and [π] cross-seeding observed before. Similar results were obtained for CgHELLP, strains co-expressing CgHELLP and GFP-RIP3(444-469) or GFP-RIP1(524-551) displayed altered growth after contact with strains [Rhim] strains (S9 Fig).

Discussion

A number of innate immune pathways function following a principle of "signaling by cooperative assembly formation" (SCAF) [29]. Upon recognition of PAMPs or MAMPs, receptors assemble into higher-order complexes or "supramolecular organizing centres" (SMOCs) [30]. Following ligand recognition, the sensor modules oligomerize and recruit adaptor or effector proteins -often via homotypic interactions- that further amplify the oligomerization process and form cooperative, open-ended polymers (typically filaments). This general scheme applies to many immune signaling modules including the RIG-I-like receptors, the AIM2-like receptors and to NLRs. Formation of higher-order assemblies entitles such signaling cascade with signal amplification, sharp threshold behavior and noise reduction. In fact, this so-called prion-like polymerization process can involve nucleation of

either folded globular domains such as the TIR or Death domains [31,32] or short amyloid

motifs. The first such motif is RHIM identified in the necroptosis pathway in mammals and

allows for assembly of the necrosome formed by the RIP1 and RIP3 kinases [23]. RHIM-like

motif also regulate amyloid formation in innate immune cascades in Drosophila [24]. In

addition, several NLR-associated amyloid signaling motifs were identified in filamentous fungi

and more recently in multicellular bacteria [18,33]. Among them is the PP-motif that shows

similarity to the animal RHIM and RHIM-like motifs [22]. We identify here in the species

Podospora anserina, a PP-motif two-component system comprising a NLR termed PNT1 and a

cell-death execution protein termed HELLP. We showed that the HELLP PP-motif supports

prion propagation *in vivo* and leads to amyloid formation *in vitro*. We generated an artificial

incompatibility system based on HELLP, analogous to the natural [Het-s]/HET-S system, where

a truncated form of HELLP containing the PP motif is equivalent to [Het-s] and the full length

HELLP to HET-S. The N-terminal HELL domain of HELLP shows homology to the 4HB domain of

MLKL responsible for membrane permeation in mammalian necroptosis. HELLP induced

programmed cell death thus appears to be related to a family of evolutionary widespread form

of immune PCD which encompasses fungal cell death, necroptosis and cell death associated

with the hypersensitive response in plants [22,34,35]. We propose that PNT1 and HELLP

constitute a third amyloid immune-related signaling cascade controlling cell fate in *P.

anserina*. We show a total absence of cross-induction between the three *P. anserina* systems

allowing for the co-existence of three independent amyloid signaling cascades in the same

species. We find that in addition to the originally described NWD2/HET-S pathway, Podospora

possesses two additional PCD pathways. It is believed that fungal NLRs akin to their plant and

animal counterparts represent innate immune receptors that regulate response to pathogen

and symbiotic interactions in fungi. The three amyloid-associated NLRs identified in Podospora

share the fact that their WD or TPR repeat ligand binding domain is highly polymorphic in

natural populations consistent with the proposed role as immune receptor [13,14]. Yet, except

in the specific case of NLRs involved in incompatibility systems the nature of the ligands

activating these NLRs are currently unknown. Compared to the two other systems, the

NWD2/HET-S is characterized by the fact that this system has been exaptated as an

allorecognition system in [Het-s]/HET-S incompatibility [36]. It has been possible

experimentally, to derive incompatibility systems analogous to [Het-s]/HET-S from both HELLP

and HELLF. It will be of interest in the future to survey, natural Podospora strains to determine

whether exaptation of allorecognition incompatibility might have also occurred from HELLP

and HELLF. It is also of note, that the *Chaetomium globosum* and *Podospora anserina* PP-based

signalosome differ in the sense that the *Chaetomium* system has three components and

involves also CgSBP, a lipase also comprising a PP-motif and that is targeted to the membrane

region by CgHELLP. HELLP and CgHELLP should thus not be consider to be orthologs, as

supported by the fact that the two proteins share 39.6% identify while PNT1 and CgPNT1 show

only 25.2% identity.

So far, four mammalian proteins involved in necroptosis and five drosophila proteins also

involved in immune signaling have been described for RHIM-based amyloid signal

transduction [23,24]. The fact that a range of viruses express RHIM containing proteins and

that pathogenic bacteria express an enzyme that specifically cleave host RHIMs to prevent

necrosome assembly highlight the crucial role of RHIM-based interactions in the host innate

immune response against microbial infection [37]. As PP-motifs of Cg and HELLP resemble

RHIM motifs. This similarity is concentrate in the central G-X-Q-X-G core of the motif.

Mompeán and co-workers resolved the structure of human RIP1-RIP3 core as a serpentine

fold with turns and kinks resulting in an enclosed oblong-shaped hydrophobic core stabilized

by N and Q ladders and Y stacking [25]. RHIM-related motif appear to represents a new class

of amyloid fold distinct from the HRAM β-solenoid fold of HET-s but nonetheless presenting

common features with them as with other amyloid forming domains such as conservation of

G/N/Q residues. N and Q allow for formation of H-bonded N/Q ladders and G connect short

adjacent β-strands [11,38–40]. The fungal PP-motifs show strong conservation of the

pseudopalindromic structure centered on the Q residue along with conservation of flanking N

and G residues. We find here that the RHIM motifs of RIP1 and RIP3 behave as PFDs when

expressed in Podospora and thus behave analogously to PP-motifs in that respect.

Importantly, we find that [Rhim] and [π] prions cross-seed, albeit not as efficiently as RHIM

motifs or fungal PP-motifs. These results suggest both a structural similarity between RHIM

and PP-amyloids and substantial differences since the efficiency of prion seeding is clearly

distinct from that of homotypic or intraspecific seeding. Extensive structural analyses of the

amyloid fibers of HELLP or CgHELLP will be required to determine to which extend PP-based

amyloid polymers share the same fold as RIP1-RIP3. It appears however that the partial cross

seeding between PP and RHIM support a model of a common evolutionary origin of the two

motifs. In this context, it is interesting to note that in addition to the RHIM-like motifs

described in Drosophila a RHIM-related amyloid motif termed BASS3 was also identified in

multicellular bacteria [33]. It might be that the RHIM-related motif represent an archetypal

amyloid motif largely shared through evolution while other motifs such as HRAMs or other

bacterial signaling motifs have a more restricted phylogenetic distribution.

Materials and Methods

Gene annotation

The Pa_5_8070 and Pa_5_8060 genes were annotated manually using two intron prediction

softwares, the GENSCAN Web server at MIT (genes.mit.edu/GENSCAN.html) and the

NetGene2 Web server (www.cbs.dtu.dk/services/NetGene2/), and using blastn searches

against EST sequences at ncbi and RNASEQ data from [41]. A single 47pb intron was identified

in nucleotide position 429 to 475 of the ORF of Pa_5_8070, this newly defined ORF encodes a

271 aa protein named HELLP. A single intron of 54 pb was identified in nucleotide position 95

to 148 of the ORF of Pa_5_8060, repeats number in the 3' part of the ORF is variable in *P.*

anserina strains, we described here the ORF of the S Orsay reference strain encoding a 971 aa

protein named PNT1.

Strains and plasmids

The *P. anserina* strains used in this study were wild-type *het-s* or *het-S* and the strains *Δhellp*

(*ΔPa_5_8070*) *het-s°* [22] and *Δhellp (ΔPa_5_8070) Δhet-s (ΔPa_3_620) Δhellf (ΔPa_3_9900)*

obtained by crossing the *Δhet-s Δhellf* strain [20] with the *Δhellp het-s°* strain. The *Δhellp het-*

s° strain was used as recipient strain for the expression of molecular fusions of full length

HELLP or PP motif-containing regions of HELLP and the GFP (green fluorescent protein) or RFP

(red fluorescent protein). These fusions were expressed from plasmids based on the pGEM-T

backbone (Promega) named pOP plasmids [22], containing either the GFP or RFP, or in a

derivative of the pAN52.1 GFP vector [27], named pGB6-GFP plasmid. In both cases, the

molecular fusions were under the control of the constitutive *P. anserina* gpd (glyceraldehyde-

3-phosphate dehydrogenase) promoter. The *Δhellp het-s°* strain was transformed as described

[42] with a fusion construct along with a second vector carrying a phleomycin-resistance gene

ble, pPaBle (using a 10:1 molar ratio). Phleomycin-resistant transformants were selected,

grown for 30 h at 26°C and screened for the expression of the transgenes using a fluorescence

microscope. The *hellp* gene and the *hellp(171-271)* and *hellp(214-271)* gene fragments were

amplified with the respective 5' forward oligonucleotides 5'

ggcttaattaaATGGATCCTCTCAGTATCACAGC 3', 5' ggcttaattaaATGATCACAGCGACAAACGATCAG

3' and 5' ggcttaattaaATGAAAGTACTGCATGAATCGCGC 3' and a same 3' reverse

oligonucleotide 5' ggcagatcttgctccCCCCCTTCGGCCAAATGTAG 3' (capital letters correspond to

P. anserina genomic DNA sequences). The PCR products were cloned upstream of the GFP or

RFP coding sequence in the pOP plasmids using PacI/BglII restriction enzymes to generate the

pOPhellp-GFP, pOPhellp-RFP, pOPhellp(171-271)-GFP, pOPhellp(171-271)-RFP, pOPhellp(214-

271)-GFP and pOPhellp(214-271)-RFP vectors in which in addition to the BglII site, a two aa

linker (GA) was introduced between the sequences encoding HELLP and GFP or RFP. *hellp(171-*

271) and *hellp(214-271)* were also amplified with the respective 5' forward oligonucleotides

5'   ggcgcgcggccgcATCACAGCGACAAACGATCAG   3',   5'

ggcgcgcggccgcATCACAGCGACAAACGATCAG 3' and the same 3' reverse oligonucleotide 5'

ggcggatccCTACCCCCTTCGGCCAAATG 3' and cloned downstream of the GFP using NotI/BamHI

restriction enzymes to generate the plasmids pGB6-GFP-hellp(171-271) and pGB6-GFP-

hellp(214-271).

To investigate the co-localization/cross-seeding of HELLP with CgHELLP or of HELLP with

HELLF or HET-s, the *Δhellp Δhet-s Δhellf* strain was transformed both with vectors expressing

fluorescently tagged versions of full length or partial HELLP described above and with

fluorescently tagged versions of full length or partial HELLF, HET-s, HET-S or CgHELLP

previously described [20,22,27]: pOPhellf-GFP, pOPhellf(209-277)-RFP, pOPhellf(209-277)-

GFP, pOPhellf(L52K)-GFP, pOPhet-s-RFP, pGPD-het-S-GFP, pOPhet-S-RFP, pGB6-GFP-

Cghellp(215-278).

In the same way the sequence encoding the first 31 aa of *pnt1 (Pa_5_8060)* was amplified

with oligonucleotides 5' ggcttaattaaATGTCAGACAGTTATCGTTTCGGC 3' and 5'

ggcagatcttgctccTGGCGCCTGCAGGAAAGTATTG 3' and cloned in the pOP plasmids using

PacI/BglII restriction enzymes to generate the pOPpnt1(1-31)-GFP and the pOPpnt1(1-31)-

RFP. The *Δhellp Δhet-s Δhellf* strain was transformed with these vectors (and pPaBle) either

alone or with pOPhellp(214-271)-RFP or pGB6-GFP-hellp(214-271).

For heterologous expression in *E. coli*, we cloned *hellp(214-271)* in pET24 (Novagen) using

the NdeI/XhoI restrictions sites. The gene fragment *hellp(214-271)* was amplified with

oligonucleotides 5' agccatatgAAAGTACTGCATGAATCGCGC 3' and 5'

agcctcgagCCCCCTTCGGCCAAATGTAG 3' and cloned in front of a polyhistidine-tag to generate

pET24-hellp(214-271)-6xhis plasmid.

For the expression of RIP3(444-469) and RIP1(524-551), PCR products were amplified with

5' ggcttaattaaATGGTTACCGGTCGTCCGCTG 3' and 5'

ggcagatcttgctccCTGCATGGTCAGGTAGTTGTTG 3' or 5' ggcgcggccgcGTTACCGGTCGTCCGCTGG

3' and 5' ggcGGATCCTTACTGCATGGTCAGGTAGTTG 3' for RIP3 and with 5'

ggcttaattaaATGACTGACGAATCCATCAAATACACC 3' and 5'

ggcagatcttgctccGCCGCCGATTTCCATGTAGTT 3' or 5'

ggcgcggccgcACTGACGAATCCATCAAATACACC 3' and 5'

ggcGGATCCTTAGCCGCCGATTTCCATGTAGTT 3' for RIP1 and cloned as described above in the

pOP plasmids using PacI/BglII restriction enzymes, or in pGB6-GFP plasmid using NotI/BamHI

restriction enzymes.

Prion propagation and incompatibility assays.

Methods for determination of incompatibility phenotypes, prion formation and prion

propagation were as previously described [20,43]. In brief, incompatibility phenotypes were

determined by confronting strains on solid corn meal agar medium and a 'barrage' reaction

(abnormal contact lines forming upon confrontation of incompatible strains) was assessed 3

days post-contact. The [π] phenotype (acquisition of the [π] prion) was assessed as the ability

of a strain to form a barrage with a wild type strain or with a *Δhellp* bearing a transgene

encoding full length HELLP in fusion with GFP or RFP, named tester strain. [π] prion formation

can be measured as the rate of spontaneous acquired [π] prion phenotype in [π*] subculture

after 4 to 15 days of growth at 26°C on corn meal agar or as the ability to propagate [π] prion

form a donor strain to a [π*] prion-free strain. In the latter case, the donor strain can contain

the [π] prion (strains expressing HELLP or CgHELLP PFD) or another prion containing strain like

[Het-s] or [Φ] and is placed in contact for 3 days with the [π*] prion-free strain consecutively

tested in barrage assay with a tester strain. Cross reaction between the different prion systems

was also assessed in barrage tests as the ability of a prion containing strain (either [π], [Het-s]

or [Φ]) to form a barrage with a strain expressing one of the full length HeLo or HELL containing

proteins HELLP, CgHELLP, HELLF or HET-S. In these experiments strains previously described

by Daskalov and colleagues were used [20,22].

Microscopy

P. anserina hyphae were inoculated on solid medium and cultivated for 24 to 72 h at 26°C.

The medium was then cut out, placed on a glass slide and examined with a Leica DMRXA

microscope equipped with a Micromax CCD (Princeton Instruments) controlled by the

Metamorph 5.06 software (Roper Scientific). The microscope was fitted with a Leica PL APO

100X immersion lens. To observe cell death reaction and HELLP plasma membrane re-

localization, HELLP-GFP or HELLP-RFP expressing strains were inoculated at a distance of 2 cm

from a strain expressing one of the HELLP PFD containing fusion or the PNT1(1-31)-GFP or -

RFP. The confrontation zone between the two strains was observed hours after contact. For

Evans blue staining, an aqueous 0.001% solution was put directly on the mycelium for 5 min

followed by washing with distilled water before observation as described above.

For HELLP fibrils observations, negative staining was performed: aggregated proteins

were adsorbed onto Formvar-coated copper grids (400 mesh) and allowed to dry for 15 min

in air, grids were then negatively stained 1 min with 10 µL of freshly prepared 2% uranyl

acetate in water, dried with filter paper, and examined with a Hitachi H7650 transmission

electron microscope (Hitachi, Krefeld, Germany) at an accelerating voltage of 120 kV. TEM was

performed at the Pôle Imagerie Électronique of the Bordeaux Imaging Center using a Gatan

USC1000 2k x 2k camera.

Protein preparation and fibril formation

HELLP(214-271) protein was expressed in *E. coli* BL21-CodonPlus®-RP competent cells as

insoluble proteins and purified under denaturing conditions using its terminal 6 histidine tag

as previously described [44]. Briefly, cells were grown at 37°C in DYT medium to 0.6 OD_{600} and

expression was induced with 1 mM isopropyl β-D-1-thiogalactopyranoside. After, 4 h, cells

were harvested by centrifugation, frozen at -80°C sonicated on ice in a lysis buffer (Tris 50

mM, 150 mM NaCl, pH 8) and centrifuged for 20 min at 20,000 g to remove *E. coli*

contaminants. The pellet containing HELLP(214-271) in inclusion bodies was washed in the

same buffer and resuspended in denaturing buffer (8M guanidinium HCl, 150 mM NaCl, and

100 mM Tris-HCl, pH 8) until complete solubilization. The lysate was incubated with Talon

Resin (CLONTECH) for 1 h at 20°C, and the resin was extensively washed with 8 M urea, 150

mM NaCl, and 100 mM Tris-HCl, pH 8. The protein was eluted from the resin in the same buffer

containing 200 mM imidazole. The protein HELLP(214-271) was pure as judged by sodium-

dodecyl-sulfate polyacrylamide-gel electrophoreses (SDS-PAGE) followed by Coomassie-Blue

staining and yield was in the range of ~2-4 mg of protein per liter of culture. To eliminate urea,

elution buffer was replaced by overnight dialysis at 4°C against Milli-Q water. Fibrils formation

resulted spontaneously from dialysis process followed by sample storage at 4°C for 7 days.

Other conditions were used for sample preparation to test the influence of salt or acidic buffer

on the fibrils formation: presence or absence of 500 mM NaCl in the dialysis buffer,

replacement of dialysis by 50 times dilution in Milli-Q water or in ammonium acetate buffer

100 mM pH 4.5, they all resulted in spontaneous appearance of fibrils.

Bioinformatic methods

Blast analyses on the *P. anserina* genome were achieved on the *Podospora anserina* Genome

Project site (podospora.i2bc.paris-saclay.fr/). Sequence alignments were performed with

Clustal Omega or MAFFT at (www.ebi.ac.uk) and edited with Jalview (www.jalview.org/). To

generate the HMM profile signatures for PP and RHIM, psi-blast searches were performed at

(blast.ncbi.nlm.nih.gov) using the HELLP PP sequence and the RHIM motif of human RIP3 and

matching sequences were aligned with Clustal Omega and the alignment was analyzed with

Skylign (skylign.org). Secondary structure predictions were performed with PSIPRED

(bioinf.cs.ucl.ac.uk/psipred/). Hidden Markov Model searches were performed using HHPred

(toolkit.tuebingen.mpg.de) and Jackhhmer (www.ebi.ac.uk/Tools/hmmer/search/phmmer),

both with default settings. The prediction of transmembrane helix was performed with the

TMHMM server (www.cbs.dtu.dk/services/TMHMM).

Author Contributions

Conceptualization: Thierry Bardin, Asen Daskalov, Sven J. Saupe, Virginie Coustou

Data curation: Thierry Bardin, Asen Daskalov, Sophie Barrouilhet, Sven J. Saupe, Virginie Coustou

Formal analysis: Thierry Bardin, Asen Daskalov, Sophie Barrouilhet, Sven J. Saupe, Virginie Coustou

Funding acquisition: Sven J. Saupe

Investigation: Thierry Bardin, Asen Daskalov, Sophie Barrouilhet, Alexandra Granger-Farbos, Bénédicte Salin, Corinne Blancard, Virginie Coustou

Methodology: Thierry Bardin, Asen Daskalov, Alexandra Granger-Farbos, Bénédicte Salin, Corinne Blancard, Sven J. Saupe, Virginie Coustou

Project administration: Sven J. Saupe, Virginie Coustou

Resources: Alexandra Granger-Farbos, Bénédicte Salin, Corinne Blancard, Sven J. Saupe, Virginie Coustou

Supervision: Sven J. Saupe, Virginie Coustou

Validation: Thierry Bardin, Asen Daskalov, Sven J. Saupe, Virginie Coustou

Visualization: Thierry Bardin, Asen Daskalov, Sven J. Saupe, Virginie Coustou

Writing: Sven J. Saupe, Virginie Coustou

References

1. Prusiner S. Novel proteinaceous infectious particles cause scrapie. Science. 1982;216: 136. doi:10.1126/science.6801762

2. Colby DW, Prusiner SB. Prions. Cold Spring Harbor Perspectives in Biology. 2011;3: a006833–a006833. doi:10.1101/cshperspect.a006833

3. Wickner RB, Edskes HK, Son M, Bezsonov EE, DeWilde M, Ducatez M. Yeast Prions Compared to Functional Prions and Amyloids. Journal of Molecular Biology. 2018;430: 3707–3719. doi:10.1016/j.jmb.2018.04.022

4. Coustou V, Deleu C, Saupe S, Begueret J. The protein product of the het-s heterokaryon incompatibility gene of the fungus Podospora anserina behaves as a prion analog. Proc Natl Acad Sci U S A. 1997;94: 9773–9778.

5. Saupe SJ. The [Het-s] prion of Podospora anserina and its role in heterokaryon incompatibility. Semin Cell Dev Biol. 2011;22: 460–468. doi:10.1016/j.semcdb.2011.02.019

6. Balguerie A, Dos Reis S, Ritter C, Chaignepain S, Coulary-Salin B, Forge V, et al. Domain organization and structure-function relationship of the HET-s prion protein of Podospora anserina. EMBO J. 2003;22: 2071–2081. doi:10.1093/emboj/cdg213

7. Seuring C, Greenwald J, Wasmer C, Wepf R, Saupe SJ, Meier BH, et al. The mechanism of toxicity in HET-S/HET-s prion incompatibility. PLoS Biol. 2012;10: e1001451. doi:10.1371/journal.pbio.1001451

8. Nazabal A, Dos Reis S, Bonneu M, Saupe SJ, Schmitter J-M. Conformational transition occurring upon amyloid aggregation of the HET-s prion protein of Podospora anserina analyzed by hydrogen/deuterium exchange and mass spectrometry. Biochemistry. 2003;42: 8852–8861. doi:10.1021/bi0344275

9. Ritter C, Maddelein M-L, Siemer AB, Lührs T, Ernst M, Meier BH, et al. Correlation of structural elements and infectivity of the HET-s prion. Nature. 2005;435: 844–848. doi:10.1038/nature03793

10. Sen A, Baxa U, Simon MN, Wall JS, Sabate R, Saupe SJ, et al. Mass analysis by scanning transmission electron microscopy and electron diffraction validate predictions of stacked beta-solenoid model of HET-s prion fibrils. J Biol Chem. 2007;282: 5545–5550. doi:10.1074/jbc.M611464200

11. Wasmer C, Lange A, Van Melckebeke H, Siemer AB, Riek R, Meier BH. Amyloid fibrils of the HET-s(218-289) prion form a beta solenoid with a triangular hydrophobic core. Science. 2008;319: 1523–1526. doi:10.1126/science.1151839

12. Mathur V, Seuring C, Riek R, Saupe SJ, Liebman SW. Localization of HET-S to the cell periphery, not to [Het-s] aggregates, is associated with [Het-s]-HET-S toxicity. Mol Cell Biol. 2012;32: 139–153. doi:10.1128/MCB.06125-11

13. Daskalov A, Habenstein B, Martinez D, Debets AJM, Sabaté R, Loquet A, et al. Signal transduction by a fungal NOD-like receptor based on propagation of a prion amyloid fold. PLoS Biol. 2015;13: e1002059. doi:10.1371/journal.pbio.1002059

685   14.   Dyrka W, Lamacchia M, Durrens P, Kobe B, Daskalov A, Paoletti M, et al. Diversity and variability
686   of NOD-like receptors in fungi. Genome Biol Evol. 2014;6: 3137–3158. doi:10.1093/gbe/evu251

687   15.   Jones JDG, Vance RE, Dangl JL. Intracellular innate immune surveillance devices in plants and
688   animals. Science. 2016;354. doi:10.1126/science.aaf6395

689   16.   Daskalov A, Paoletti M, Ness F, Saupe SJ. Genomic clustering and homology between HET-S and
690   the NWD2 STAND protein in various fungal genomes. PLoS ONE. 2012;7: e34854.
691   doi:10.1371/journal.pone.0034854

692   17.   Cai X, Chen J, Xu H, Liu S, Jiang Q-X, Halfmann R, et al. Prion-like polymerization underlies signal
693   transduction in antiviral immune defense and inflammasome activation. Cell. 2014;156: 1207–
694   1222. doi:10.1016/j.cell.2014.01.063

695   18.   Loquet A, Saupe SJ. Diversity of Amyloid Motifs in NLR Signaling in Fungi. Biomolecules. 2017;7.
696   doi:10.3390/biom7020038

697   19.   Daskalov A, Dyrka W, Saupe SJ. Theme and variations: evolutionary diversification of the HET-s
698   functional amyloid motif. Sci Rep. 2015;5: 12494. doi:10.1038/srep12494

699   20.   Daskalov A, Martinez D, Coustou V, El Mammeri N, Berbon M, Andreas LB, et al. Structural and
700   molecular basis of cross-seeding barriers in amyloids. bioRxiv. 2020; 2020.07.06.188508.
701   Available   from: https://www.biorxiv.org/content/10.1101/2020.07.06.188508v1.
702   doi:10.1101/2020.07.06.188508

703   21.   Murphy JM, Czabotar PE, Hildebrand JM, Lucet IS, Zhang J-G, Alvarez-Diaz S, et al. The
704   Pseudokinase MLKL Mediates Necroptosis via a Molecular Switch Mechanism. Immunity.
705   2013;39: 443–453. doi:10.1016/j.immuni.2013.06.018

706   22.   Daskalov A, Habenstein B, Sabaté R, Berbon M, Martinez D, Chaignepain S, et al. Identification of
707   a novel cell death-inducing domain reveals that fungal amyloid-controlled programmed cell
708   death is related to necroptosis. Proc Natl Acad Sci USA. 2016;113: 2720–2725.
709   doi:10.1073/pnas.1522361113

710   23.   Li J, McQuade T, Siemer AB, Napetschnig J, Moriwaki K, Hsiao Y-S, et al. The RIP1/RIP3 Necrosome
711   Forms a Functional Amyloid Signaling Complex Required for Programmed Necrosis. Cell.
712   2012;150: 339–350. doi:10.1016/j.cell.2012.06.019

713   24.   Kleino A, Ramia NF, Bozkurt G, Shen Y, Nailwal H, Huang J, et al. Peptidoglycan-Sensing Receptors
714   Trigger the Formation of Functional Amyloids of the Adaptor Protein Imd to Initiate Drosophila
715   NF-κB Signaling. Immunity. 2017;47: 635-647.e6. doi:10.1016/j.immuni.2017.09.011

716   25.   Mompeán M, Li W, Li J, Laage S, Siemer AB, Bozkurt G, et al. The Structure of the Necrosome
717   RIPK1-RIPK3 Core, a Human Hetero-Amyloid Signaling Complex. Cell. 2018;173: 1244-1253.e10.
718   doi:10.1016/j.cell.2018.03.032

719   26.   Beisson-Schecroun J. [Cellular incompatibility and nucleo-cytoplasmic interactions in the
720   "barrage" phenomena in Podospora anserina]. Ann Genet. 1962;4: 4–50.

721   27.   Balguerie A, Dos Reis S, Coulary-Salin B, Chaignepain S, Sabourin M, Schmitter J-M, et al. The
722   sequences appended to the amyloid core region of the HET-s prion protein determine higher-
723   order aggregate organization in vivo. J Cell Sci. 2004;117: 2599–2610. doi:10.1242/jcs.01116

28. Benkemoun L, Ness F, Sabaté R, Ceschin J, Breton A, Clavé C, et al. Two structurally similar fungal
prions efficiently cross-seed in vivo but form distinct polymers when coexpressed. Mol Microbiol.
2011;82: 1392–1405. doi:10.1111/j.1365-2958.2011.07893.x

29. Nanson JD, Kobe B, Ve T. Death, TIR, and RHIM: Self-assembling domains involved in innate
immunity and cell-death signaling. J Leukoc Biol. 2019;105: 363–375. doi:10.1002/JLB.MR0318-
123R

30. Kagan JC, Magupalli VG, Wu H. SMOCs: supramolecular organizing centres that control innate
immunity. Nat Rev Immunol. 2014;14: 821–826. doi:10.1038/nri3757

31. Wu H. Higher-order assemblies in a new paradigm of signal transduction. Cell. 2013;153: 287–
292. doi:10.1016/j.cell.2013.03.013

32. Cai X, Chen ZJ. Prion-like polymerization as a signaling mechanism. Trends Immunol. 2014;35:
622–630. doi:10.1016/j.it.2014.10.003

33. Dyrka W, Coustou V, Daskalov A, Lends A, Bardin T, Berbon M, et al. Identification of NLR-
associated amyloid signaling motifs in filamentous bacteria. bioRxiv; 2020 Jan. Available from:
https://www.biorxiv.org/content/10.1101/2020.01.06.895854v1.
doi:10.1101/2020.01.06.895854

34. Burdett H, Bentham AR, Williams SJ, Dodds PN, Anderson PA, Banfield MJ, et al. The Plant
"Resistosome": Structural Insights into Immune Signaling. Cell Host & Microbe. 2019;26: 193–
201. doi:10.1016/j.chom.2019.07.020

35. Hofmann K. The Evolutionary Origins of Programmed Cell Death Signaling. Cold Spring Harb
Perspect Biol. 2019; a036442. doi:10.1101/cshperspect.a036442

36. Daskalov A, Saupe SJ. As a toxin dies a prion comes to life: A tentative natural history of the [Het-
s] prion. Prion. 2015;9: 184–189. doi:10.1080/19336896.2015.1038018

37. Baker MODG, Shanmugam N, Pham CLL, Strange M, Steain M, Sunde M. RHIM-based
protein:protein interactions in microbial defence against programmed cell death by necroptosis.
Seminars in Cell & Developmental Biology. 2018 [cited 11 Sep 2018].
doi:10.1016/j.semcdb.2018.05.004

38. Plumley JA, Dannenberg JJ. The importance of hydrogen bonding between the glutamine side
chains to the formation of amyloid VQIVYK parallel beta-sheets: an ONIOM DFT/AM1 study. J Am
Chem Soc. 2010;132: 1758–1759. doi:10.1021/ja909690a

39. Halfmann R, Alberti S, Krishnan R, Lyle N, O'Donnell CW, King OD, et al. Opposing effects of
glutamine and asparagine govern prion formation by intrinsically disordered proteins. Mol Cell.
2011;43: 72–84. doi:10.1016/j.molcel.2011.05.013

40. Tuttle MD, Comellas G, Nieuwkoop AJ, Covell DJ, Berthold DA, Kloepper KD, et al. Solid-state NMR
structure of a pathogenic fibril of full-length human α-synuclein. Nat Struct Mol Biol. 2016;23:
409–415. doi:10.1038/nsmb.3194

41. Lamacchia M, Dyrka W, Breton A, Saupe SJ, Paoletti M. Overlapping Podospora anserina
Transcriptional Responses to Bacterial and Fungal Non Self Indicate a Multilayered Innate
Immune Response. Front Microbiol. 2016;7: 471. doi:10.3389/fmicb.2016.00471

42. Bergès T, Barreau C. Heat shock at an elevated temperature improves transformation efficiency
of protoplasts from Podospora anserina. J Gen Microbiol. 1989;135: 601–604.
doi:10.1099/00221287-135-3-601

43. Benkemoun L, Sabaté R, Malato L, Dos Reis S, Dalstra H, Saupe SJ, et al. Methods for the in vivo
and in vitro analysis of [Het-s] prion infectivity. Methods. 2006;39: 61–67.
doi:10.1016/j.ymeth.2006.04.006

44. Dos Reis S, Coulary-Salin B, Forge V, Lascu I, Bégueret J, Saupe SJ. The HET-s prion protein of the
filamentous fungus Podospora anserina aggregates in vitro into amyloid-like fibrils. J Biol Chem.
2002;277: 5703–5706. doi:10.1074/jbc.M110183200

FIGURES

Fig 1. The PP motif and the two HRAM motif gene clusters of *P. anserina*.

(A) Genome organization and domain architecture of the *het-s/nwd2*, *hellf/fnt1* and *hellp/pnt1* gene clusters of *P. anserina*. These loci are composed of two adjacent genes encoding an effector protein and a NLR receptor. The effector proteins are composed of a PFD and a HeLo or HELL domain. The NLRs are composed of a PFD, a NB-ARC or NACHT nucleotide binding domain and of WD or TPR repeats (repetition numbers is variable as indicated). PFDs of the HRAM family are represented as a turquoise line while those of the PP family as a light-

782   brown line. (B) Alignment of the PP motifs of HELLP, PNT1, CgHELLP, CgSBP, and CgPNT1 along

783   with the RHIM motif of human RIP1 and RIP3.

Fig 2. HELLP PP region behaves as a prion *in vivo* and forms fibrils *in vitro*

(A) Micrographs of *P. anserina* strains expressing different molecular fusions (HELLP(214-271-RFP/GFP, GFP-HELLP(214-271)) as indicated above each micrograph (Scale bar: 5 μm). Transformants present initially a diffuse fluorescence in the non-prion state designated [π*] (left) and acquire dot-like fluorescent aggregates after contact with a strain already expressing the infectious prion state designated [π] (right). (B) Electron micrograph of negatively stained HELLP(214-271) fibrils, scale bar is 100 nm.

Fig 3. Activation of HELLP cell death activity by [π] prion.

(A) Observation of incompatibility reaction between [π] prion and HELLP in barrage tests.

Confrontation on solid medium of transformants expressing GFP-HELLP(214-271) in soluble

[π*] or aggregated [π] states with strains expressing full length HELLP from a transgene

(written in white) or from the wild type *hellp* gene or with the *Δhellp* strain as a negative

control. Formation of a barrage indicative of an incompatibility reaction is marked by a white

arrow. The thinner arrow indicates an attenuated barrage reaction. (B) Micrographs of fusion

cells coexpressing full-length HELLP (HELLP-GFP (upper panel) or HELLP-RFP (lower panel)),

and HELLP(214-271)-RFP (upper panel) or GFP-HELLP(214-271) (lower panel) in the [π]

aggregated state. Note that HELLP relocalizes to the cell membrane region in presence of [π].

(C) Micrographs showing confrontations between HELLP-GFP and GFP-HELLP(214-271)

806   expressing strains after Evans blue staining. Note that heterocaryotic fusion cells are stained

807   indicating their death. Scale bar (2 μm).

Fig 4. PNT1(1-31) aggregates *in vivo*, co-localizes with HELLP(214-271) and targets HELLP to the plasma membrane.

(A) Micrographs of PNT1(1-31) fusions with GFP or RFP either in the soluble state (left) or aggregated state (right). (B) Micrographs showing co-localization of PNT1(1-31) and HELLP(214-271) in soluble (top picture) or aggregated states. (C) In fusion cells coexpressing full-length HELLP (fused with RFP (upper panel) or GFP (lower panel)), and aggregated forms of PaPNT1(1-31) (fused with GFP (lower panel), or RFP, lower panel), HELLP relocalized to the plasma membrane. Scale bar (2 μm). (D) Micrographs showing confrontations between HELLP-GFP and GFP-PNT1(1-31) expressing strains after Evans blue staining. Note that heterocaryotic fusion cells are stained indicating their death. Scale bar (2 μm).

Fig 5. Lack of cross-seeding *in vivo* between [π] and [Het-s] or [Φ] prions.

(A) Micrographs of strains co-expressing HET-s-RFP ([Het-s] or [Het-s*] state) and GFP-HELLP(214-271) ([π*] or [π] state). Note an absence of co-localization of the two aggregated prion forms and the absence of conversion of [Het-s*] state by [π] prion and of [π*] by [Het-s] prion. Scale bar (5 µm). (B) Micrographs of strains co-expressing HELLF(209-277)-RFP ([Φ]

829   or [Φ*] state) and GFP-HELLP(214-271) ([π*] or [π] state). Note an absence of co-localization

830   of the two aggregated prion forms and the absence of conversion of [Φ *] state by [π] prion

831   and of [π*] by [Φ] prion. Scale bar (5 μm).

Fig 6. Co-localization of HELLP and CgHELLP PFDs and cross induction of cell death activity.

(A) Micrographs of strains co-expressing HELLP(214-271)-RFP and GFP-CgHELLP(215-278) in soluble [π*] state (upper panels) or aggregated [π] state (lower panels). Note the co-localization of the two aggregated prion forms. Scale bar (5 μm). (B) Confrontation on solid medium of strains expressing GFP-HELLP(214-271) and GFP-CgHELLP(215-278) [π] prions forms with strains expressing full length HELLP from a transgene (written in white) or from the wild type *hellp* gene or with the *Δhellp* strain as a negative control. Formation of a barrage indicative of an incompatibility reaction is marked by a white arrow. The thinner arrow indicates an attenuated barrage reaction.

Fig 7. *In vivo* prion propagation of the human RIP3 and RIP1 RHIM regions.

(A) Micrographs of *P. anserina* strains expressing the molecular fusions RIP3(444-469)-RFP, RIP3(444-469)-GFP or GFP-RIP3(444-469). (B) Micrographs of *P. anserina* strains expressing the molecular fusions RIP1(524-551)-RFP, RIP1(524-551)-GFP or GFP-RIP1(524-551). (A and B) Transformants present initially a diffuse fluorescence in the non-prion state designated [Rhim*] (left) and acquire dot-like fluorescent aggregates after contact with a strain already expressing the infectious prion state designated [Rhim] (right). (Scale bar: 5 µm).

Fig 8. Cross conversion between [π] and [Rhim] prions.

Histograms representing the percentage of [π] and [Rhim] prion containing strains obtained

after contact of induced strains (initially displaying [π*] or [Rhim*] phenotype) with a

ΔhellpΔhet-sΔhellf prion free strain (control), a GFP-HELLP(214-271) [π] containing strain or

GFP-RIP3(444-469) or GFP-RIP1(524-551) [Rhim] containing strains (indicated on top).

Phenotype after induction was determined by monitoring the acquisition of dot-like

aggregates by fluorescence microscopy as an indicator of the prion propagation. Percentages

of prion formation were expressed as the mean value ± standard deviation for 3 to 5

independent experiments on 4 different transformants. P-values for cross conversion were

determined using two tails Fisher's test by comparison of the number of prion free and prion

containing strains obtained after induction by the prion free control or by the heterotypic

prion containing strain.

: p-values for cross conversion of HELLP(214-271) [π] strains to [π] strains by GFP-RIP3(444-

496) [Rhim] or by GFP-RIP1(524-551) [Rhim] inducers were respectively 8.5×10^{-8} and 1.6×10^{-8}.

**: p-values for cross conversion of GFP-RIP3(444-496) [Rhim*] or GFP-RIP1(524-551) [Rhim*]

strains to [Rhim] strains by HELLP(214-271) [π] inducer were respectively 3.8×10^{-8} and $7.2 \times 10^{-}$

869 [8].

870 ***: p-values for cross conversion of GFP-RIP3(444-496) [Rhim*] to [Rhim] strains by GFP-

871 RIP1(524-551) [Rhim] inducer or GFP-RIP1(524-551) [Rhim*] to [Rhim] strains by GFP-

872 RIP3(444-496) [Rhim] inducer were respectively 1.5×10^{-16} and 3.2×10^{-16}.

Fig 9. Partial cross-seeding *in vivo* between [π] and [Rhim] prions.

(A) Micrographs of strains co-expressing RIP3(444-469)-RFP ([Rhim*] or [Rhim] states) and

GFP-HELLP(214-271) ([π*] or [π] states). (B) Micrographs of strains co-expressing RIP1(524-

551)-RFP ([Rhim*] or [Rhim] states) and GFP-HELLP(214-271) ([π*] or [π] states). In both cases,

note the co-existence in the second lane of [Rhim] and [π*] (A) or [Rhim*] and [π] (B), these

880   situations are only observable for a short period of time showing a low conversion rate of [π*]

881   by [Rhim] prion or of [Rhim*] by [π]. Note also the partial co-localization of the two prions,

882   some of the dots were zoomed to show incomplete overlapping in the case of heteropolymers

883   (Scale bar: 5 µm).

Fig 10. Presence of [Rhim] prion elicits HELLP cell death activity *in vivo*.

(A) Comparison of growth on solid medium of strains co-expressing HELLP-RFP and GFP-
RIP3(444-469) or RIP1(524-551) before and after contact with a [Rhim] prion containing strain.

Note that induction of [Rhim] spreading in transformants leads to altered growth phenotypes:

some of the transformants displayed drastic sublethal phenotypes and others only stunted

growth. (B) Micrographs of strains co-expressing full length HELLP-RFP and GFP-RIP3(444-469)

or GFP-RIP1(524-551) in the [Rhim] state. Note that HELLP relocalizes to the cell membrane in

presence of [Rhim] in dying cells or aggregates as dots partially co-localizing with [Rhim]

polymers in apparently more healthy cells.

Supporting information

Tables

S1 Table. Monitoring of [π] prion propagation using barrage tests.

Tested strain		Previous contact with		Tester strains		
				ΔhellpΔhet-sΔhellf		Recipient strain
Recipient strain	Transgene	Recipient strain	Transgene	HELLP-GFP	HELLP-RFP	Transgene
Δhellp het-s°	GFP-HELLP (214-271) **[π*]**	*Δhellp het-s°*	-	-	-	
			GFP-HELLP(214-271) **[π]**	+	+	
			HELLP (214-271)-GFP **[π]**	+	+	
			HELLP (214-271)-RFP **[π∗]**	-	-	
			HELLP (214-271)-RFP **[π]**	+	+	
			HELLP (171-271)-GFP **[π]**	+	+	
			GFP-HELLP (171-271) **[π]**	+	+	

+ indicates the presence of a barrage between the tested and the tester strains, - indicates the absence of barrage. The phenotype conferred by the transgene is written in bold. Tested strains were previously in contact with strains displaying the [π∗] or [π] phenotype or with a negative control lacking transgene before confrontation with the tester strains.

904 **S2 Table. Test of the [Het-s] and [Φ] prions ability to convert [π∗].**

Tested strain		Previous contact with		Tester strains		Recipient strain
				ΔhellpΔhet-sΔhellf		
Recipient strain	Transgene	Recipient strain	Transgene	HELLP-RFP	HELLP-GFP	Transgene
Δhellp Δhet-sΔhellf	HELLP (214-271)-GFP **[π*]**	*Δhellp Δhet-sΔhellf*	HET-s-RFP **[Het-s]**	-	-	
			GFP-HELLF(209-277) **[Φ]**	-	-	
			GFP-HELLP (214-271) **[π]**	+	+	
Δhellp Δhet-sΔhellf	HELLP (214-271)-RFP **[π*]**	*Δhellp Δhet-sΔhellf*	HET-s-RFP **[Het-s]**	-	-	
			GFP-HELLF(209-277) **[Φ]**	-	-	
			GFP-HELLP (214-271) **[π]**	+	+	

906 + indicates the presence of a barrage between the tested and the tester strains, - indicates the absence
907 of barrage. The phenotype conferred by the transgene is written in bold. Tested strains were previously
908 in contact with strains displaying the prion phenotype associated to the transgene: either [Het-s], [Φ]
909 or [π] before confrontation with the tester strains.
910
911

912 **S3 Table. Test of cross-interactions between the three *P. anserina* amyloid signaling systems**

913 **by incompatibility assays.**

Tested strains		Tester strain *ΔhellpΔhetsΔhellf*			Recipient strain
Recipient strain	Transgene	HET-S-GFP	HELLF-GFP	HELLP-GFP	Transgene
ΔhellpΔhet-sΔhellf	HET-s-RFP	+	-	-	
Δhet-s	HET-s(218-289)-GFP	+	-	-	
ΔhellpΔhet-sΔhellf	HELLF(209-277)-RFP	-	+	-	
ΔhellpΔhet-sΔhellf	GFP-HELLF(209-277)	-	+	-	
ΔhellpΔhet-sΔhellf	HELLF(L52K)-GFP	-	+	-	
ΔhellpΔhet-sΔhellf	HELLP(214-271)-GFP	-	-	+	
ΔhellpΔhet-sΔhellf	GFP-HELLP(214-271)	-	-	+	
Δhellp het-s°	GFP-CgHELLP(215-278)	-	-	+	

914 + indicates the presence of a barrage between the tested and the tester strains, - indicates the absence
915 of barrage. All the tested strains used in these barrage tests displayed the prion phenotype associated
916 to the transgene: either [Het-s], [Φ] or [π].

919 **S4 Table. [π] prion transmission and activation of HELLP or CgHELLP cell death activity across**

920 **the species barrier.**

Tested strain		Previous contact with		Tester strains		
				Δhellp het-s°	*Δhellp Δhet-sΔhellf*	Recipient strain
Recipient strain	Transgene	Recipient strain	Transgene	CgHELLP-RFP	HELLP-RFP	Transgene
Δhellp Δhet-sΔhellf	GFP-HELLP (214-271) **[π*]**	*Δhellp Δhet-sΔhellf*	-	-	-	
		Δhellp Δhet-sΔhellf	GFP-HELLP (214-271) **[π]**	+	+	
		Δhellp het-s°	GFP-CgHELLP (215-278) **[π]**	+	+	
Δhellp het-s°	GFP-CgHELLP (215-278) **[π*]**	*Δhellp Δhet-sΔhellf*	-	-	-	
		Δhellp Δhet-sΔhellf	GFP-HELLP (214-271) **[π]**	+	+	
		Δhellp het-s°	GFP-CgHELLP (215-278) **[π]**	+	+	

+ indicates the presence of a barrage between the tested and the tester strains, - indicates the absence
of barrage. The phenotype conferred by the transgene is written in bold. Tested strains were previously
in contact with strains displaying the [π] phenotype or with a negative control lacking transgene before
confrontation with the tester strains. Same results were obtained with HELLP(214-271)-GFP,
HELLP(171-271)-GFP, GFP-HELLP(171-271) or HELLP(214-271)-RFP transgenes instead of the GFP-
HELLP(214-271) transgene and with CgHELLP(170-278)-RFP instead of GFP-CgHELLP(215-278).

S5 Table. Cross-induction of HELLP and CgHELLP cell death activity by [π] prions.

Tested strains		Tester strains					Recipient strain
		Δhellp Δhet-sΔhellf	*Δhellp Δhet-sΔhellf*	*Δhellp Δhet-sΔhellf*	*wt*	*Δhellp het-s°*	Recipient strain
Recipient strain	Transgene	-	HELLP-RFP	HELLP-GFP	-	CgHELLP-RFP	Transgene
Δhellp Δhet-sΔhellf	-	-	-	-	-	-	
Δhellp Δhet-sΔhellf	GFP-HELLP (214-271)	-	+	+	+[a]	+	
Δhellp Δhet-sΔhellf	GFP-HELLP (171-271)	-	+	+	+[a]	+	
Δhellp Δhet-sΔhellf	HELLP (214-271)-RFP	-	+	+	+[a]	+	
Δhellp het-s°	GFP-CgHELLP (215-278)	-	+	+	-	+	
Δhellp het-s°	CgHELLP (170-278)-RFP	-	+	+	-	+	

+ indicates the presence of a barrage between the tested and the tester strains, - indicates the absence of barrage. [a] the barrage is barely perceptible. All the tested strains bearing a transgene used in these incompatibility barrage tests displayed the [π] phenotype. *wt* is usually the wild type Cs strain, but the same results were obtained with other wild type strains.

Figures

S1 Fig. Homology between HeLo or HELL domains of *P. anserina* and pore forming domain of human MLKL.

(A) Alignment of the HET-S HeLo domain and the HELLF and HELLP HELL domains of *P. anserina* with the 4HB region of the human MLKL. Alignment was generated with clustal omega with default

943   settings. As for CgHELLP, the first hit to HELLP is MLKL in HHPred searches with a p-value score of

944   1.6×10^{-6} *versus* only 1.5×10^{-4} for HET-S again emphasing its significant homology with the MLKL

945   4HB necroptotic factor domain. (B) Secondary structure prediction for HELLP HELL domain and

946   MLKL 4HB domain using PSIPRED.

948
S2 Fig. Observation of [π] prion propagation.

(A) Micrographs of *P. anserina* strains expressing HELLP(171)271-GFP or GFP-HELLP(171-271) as

indicated above each micrograph (Scale bar: 5 μm). Transformants present initially a diffuse

fluorescence in the non-prion state designated [π*] (left) and acquire dot-like fluorescent

aggregates after contact with a strain already expressing the infectious prion state designated [π]

(right). (B) Barrage tests showing conversion of [π*] strains (expressing either HELLP(214-271)-GFP

or HELLP(171-271)-GFP) to [π] phenotype by contact with strains expressing GFP-HELLP(214-271)

in the [π] state. The acquired ability to form a barrage (marked by a white arrow) with a tester

strain expressing HELLP-GFP indicates conversion. [π] prion propagation is allowed by the position

of the inocula, as [π] and [π*] strains are closer and made contact before the tester strain reach

the tested strain. A *Δhellp* strain was used as a negative control of conversion.

961
962 **S3 Fig. HELLP full length diffuse and cytoplasmic localization.**

963 Micrographs of strains expressing HELLP-GFP or HELLP-RFP fusion proteins. The fluorescence

964 signal is diffuse and cytoplasmic and remains stable.

S4 Fig. Lack of cross-seeding and co-localization *in vivo* between [π] and [Φ] prions.

Micrographs of strains co-expressing HELLP(214-271)-RFP ([π*] or [π] state) and HELLF(209-277)-GFP ([Φ*] or [Φ] state). Note an absence of co-localization of the two aggregated prion forms ([Φ] and [π]) and the absence of conversion of [Φ*] state by [π] prion and of [π*] by [Φ] prion.

972
973 **S5 Fig. Cross-seeding of [Rhim] prions in *P. anserina*.**

974 Micrographs of *P. anserina* strains coexpressing the molecular fusions RIP3(444-469)-RFP and GFP-

975 RIP1(524-551) or GFP-RIP3(444-469) and RIP1(524-551)-RFP. (Scale bar: 5 μm). Note the strong

976 colocalization between [Rhim] prions and dots alignment nearby the septa (zoomed regions).

978
S6 Fig. Partial cross-seeding of [π] and [Rhim] prions *in vivo*.

(A) Micrographs of strains co-expressing GFP-RIP3(444-469) ([Rhim*] or [Rhim] states) and

HELLP(214-271)-RFP ([π*] or [π] states). (B) Micrographs of strains co-expressing GFP-RIP1(524-

551) ([Rhim*] or [Rhim] states) and HELLP(214-271)-RFP ([π*] or [π] states). In both cases, note

the co-existence on the second lane of [Rhim] and [π*] (A) or [Rhim*] and [π] (B), these situations

were only observable for a short period of time and showed a low conversion rate of [π*] by [Rhim]

prion or of [Rhim*] by [π]. Note the partial co-localization of the two prions; some of the dots were

zoomed to show incomplete overlapping. (C) Micrographs of strains co-expressing GFP-RIP3(444-

469) and RIP3(444-469)-RFP or GFP-RIP1(524-551) and RIP1(524-551)-RFP of [Rhim] phenotypes.

Note the almost complete overlapping of dots of [Rhim] homopolymers (Scale bar: 5 µm).

S7 Fig. Interaction between human RIP3 or RIP1 RHIM region and CgHELLP PFD.

(A) Induction of [Rhim] prion by cross conversion of [Rhim*] by [π] Cg or [Rhim] prions. Histogram

representing the percentage of [Rhim*] strains displaying [Rhim] phenotype after contact with a

ΔhellpΔhet-sΔhellf prion free strain (control), a GFP-CgHELLP(215-278) [π] containing strain or a

GFP-RIP3(444-469) or GFP-RIP1(524-551) [Rhim] containing strains (indicated on top). Strains

phenotype after induction was determined by monitoring the acquisition of dot-like aggregates

by fluorescence microscopy. Percentages of prion formation were expressed as the mean value ±

standard deviation for 3 to 5 independent experiments on 4 different transformants. P-values for

cross conversion were determined using two tails Fisher's test by comparison of the number of

prion free and prion containing strains obtained after induction by the prion free control or by the

heterotypic prion containing strain. (B) Partial cross-seeding of [π] Cg and [Rhim] prions *in vivo*.

Micrographs of strains co-expressing RIP3(444-469)-RFP or RIP1(524-551)-RFP ([Rhim*] or [Rhim]

states) and GFP-CgHELLP(215-278) ([π*] and [π] state). Note the co-existence of [Rhim*] and [π],

this situation is only observable for a short period of time, showing a low conversion rate of

[Rhim*] by [π] prion. Note the partial co-localization of the two prions, some of the dots were

zoomed to show incomplete overlapping (Scale bar: 5 μm).

1008

1009

1010 **S8 Fig. Cell death occurring throughout confrontation of [Rhim] and HELLP expressing strains.**

1011 Micrographs showing confrontations between HELLP-GFP and GFP-RIP3(444-469) (Upper panel)

1012 or GFP-RIP1(524-551) (lower panel) expressing strains after Evans blue staining. Note that

1013 heterocaryotic fusion cells are stained indicating their death. Scale bar (2 μm).

1014

1015

S9 Fig. Triggering of CgHELLP cell death activity *in vivo* by [Rhim] prion

(A) Growth of strains co-expressing CgHELLP-RFP and GFP-RIP3(444-469) or GFP-RIP1(524-551)

before and after contact with a [Rhim] prion containing strain were compared on solid medium.

1019   Note that induction of [Rhim] in transformants leads to growth alteration to varying extent ranging

1020   from sublethal phenotype to more or less stunted growth. (B) Micrographs of strains co-expressing

1021   full length CgHELLP-RFP and GFP-RIP3(444-469) or GFP-RIP1(524-551) in the [Rhim] state. CgHELLP

1022   relocalizes to the cell membrane or aggregates as dots partially co-localizing with [Rhim] polymers

1023   in damaged cells (Scale bar: 5 μm).

1024

1025