**Title:** Distinct networks coupled with parietal cortex for spatial representations

inside and outside the visual field

**Authors:** Bo Zhang [1, 3, 4], Fan Wang [7,8,9], Qi Zhang [1, 10], Naya Yuji [1, 2, 5, 6, *]

**Affiliations:**

[1] School of Psychological and Cognitive Sciences, Peking University, No. 52, Haidian Road,

Haidian District, Beijing 100805, China

[2] IDG/McGovern Institute for Brain Research at Peking University, No. 52, Haidian Road,

Haidian District, Beijing 100805, China

[3] Beijing Academy of Artificial Intelligence, Beijing, 100084, China

[4] Tsinghua Laboratory of Brain and Intelligence, 160 Chengfu Rd., SanCaiTang Building,

Haidian District, Beijing, 100084, China

[5] Center for Life Sciences, Peking University, No. 52, Haidian Road, Haidian District, Beijing

100805, China

[6] Beijing Key Laboratory of Behavior and Mental Health, Peking University, No. 52, Haidian

Road, Haidian District, Beijing 100805, China

[7] State Key Laboratory of Brain and Cognitive Science, Institute of Biophysics, Chinese

Academy of Sciences, 15 Datun Road, Beijing 100101, China

[8] CAS Center for Excellence in Brain Science and Intelligence Technology

[9] University of Chinese Academy of Sciences, 19A Yuquan Road, Beijing 100049, China

[10] Fujian Key Laboratory of Applied Cognition and Personality, School of Educational

Science, Minnan Normal University, 36 Xianqianzhi Street, Zhangzhou 363000, China

**Corresponding author:** Yuji Naya

Address: School of Psychological and Cognitive Sciences, Peking University, No. 52,

Haidian Road, Wang Kezhen Building, Room 1707, Haidian District, Beijing 100805, China

E-mail: yujin@pku.edu.cn

**Conflict of interest statement**: No competing financial interests

**Acknowledgments:** This work was supported by the National Natural Science Foundation of

China Grant 31421003 (to Y.N.) and by the Fundamental Research Funds for the Central

Universities, PKU (7100602954 to Y.N.). We thank the National Center for Protein Sciences

at Peking University for assistance with MRI data collection, and the State Key Laboratory of

Brain and Cognitive Science at the Chinese Academy of Sciences for assistance with MEG

scanning, which was supported in part by grants from the Ministry of Science and

Technology of China (2019YFA0707103), National Nature Science Foundation of China

(31730039), and Strategic Priority Research Program of the Chinese Academy of Science

(XDB32010300). Computational work was supported by resources provided by the High-

performance Computing Platform of Peking University.

**Abstract**

Our mental representation of egocentric space is influenced by the disproportionate sensory

perception of the body. Previous studies have focused on the neural architecture for

egocentric representations within the visual field. However, the space representation

underlying the body is still unclear. To address this problem, we applied both fMRI and MEG

to a spatial-memory paradigm by using a virtual environment in which human participants

remembered a target location left, right, or back relative to their own body. Both experiments

showed larger involvement of the frontoparietal network in representing a retrieved target on

the left/right side than on the back. Conversely, the medial temporal lobe (MTL)-parietal

network was more involved in retrieving a target behind the participants. The MEG data

showed preferential connectivity in the alpha-band frequency in both networks. These

findings suggest that the parietal cortex may represent the entire space around the self-body

by coordinating two distinct brain networks.

## Introduction

While planning to reach a target, it is necessary to obtain its location information in the body-centered reference frame or the so-called "egocentric" spatial coordinates[1-4]. Previous studies using humans and nonhuman primates indicated the crucial involvement of the parietal cortex in the representation of egocentric location for sensory perception, motor action, and their coordination[4-10]. Anatomically, the parietal cortex is located at the late stage of the dorsal pathway, which is often named the "where" or "how" pathway[11-12]. Perceptual signals, including visual, somatosensory, and vestibular information, converge on the parietal cortex, which interacts with the frontal eye field (FEF), the region associated with attention and eye movement toward a target in the external space around the self-body[13-16]. These anatomical connection patterns are consistent with those reported by neuropsychological studies, that is, damage to the parietal cortex results in a form of impaired egocentric spatial awareness known as hemispherical neglect, which leads to neglect of a target on one side of the field of vision[17]. Hemispherical neglect appears not only in perception but also in the memory field, where it is known as "representational neglect"[18]. On the basis of data accumulated from human imaging studies examining spatial navigation and episodic recollection[19-23], the parietal cortex is considered to represent the egocentric space for both perception and memory.

Mnemonic representations of the egocentric space raise the question of whether the spatial representation differs between the inside and outside of a visual field. While the former can be represented in both vision and memory, the latter can be represented only in memory. Previous human behavioral studies reported decreased performance[24] or prolonged response latency[25] when participants located a target behind them. This phenomenon, known as the "alignment effect"[26] or "front facilitation,"[27] suggests that the space surrounding our body is coded heterogeneously by different neural systems depending on the target location

along the anterior-posterior axis of the self-body[27]. To explore the neural architectures

responsible for the representation of egocentric space, previous studies mostly examined the

target representation within the visual field; however, one neuroimaging study, which

examined target representation retrieved from memory, suggested that the parietal cortex

codes the whole space around the self-body[28]. However, the representation properties of the

inside/outside of the visual field in the parietal cortex and its interactions with associated

brain areas remain unsolved.

To characterize the neural architectures supporting the mental representations of

egocentric space inside and outside the visual field, we applied functional magnetic

resonance imaging (fMRI) and magnetoencephalography (MEG) to a spatial-memory

paradigm using a 3D virtual environment for human participants (Fig. 1a), which we recently

devised[29]. In this paradigm, participants encoded a spatial relationship among three objects

(walking period) in each trial. While the same three objects were used across trials, their

spatial relationships differed pseudorandomly. After the walking period, one of the three

objects was presented in the center of the display with the environmental background, which

prompted the participants to feel as if they were facing the objects in the virtual environment

(facing period)[29]. Then, one of the other two objects was presented on a scrambled

background as the target object (targeting period). The participants remembered the location

of the target object relative to the body. The to-be-remembered targeting locations could be in

the left, right, or back positions of the participants' egocentric spaces (Fig. 1b). This spatial-

memory paradigm allowed us to compare the neural representations of egocentric space

between the inside and outside of the visual field when the participants remembered a target

location, which had been encoded in each trial.

# Results

**Behavioral performance**

Nineteen and 12 healthy volunteers participated in the fMRI and MEG experiments,

respectively. The participants performed the spatial-memory task with high accuracy rates in

both the experiments (MRI: 93.6% ± 1.5%; MEG: 90.4% ± 1.9%). The performances did not

differ among the three target locations (i.e., right, left, and back) in both fMRI [$F$ (2,54) =

0.82, $P$ = 0.44] and MEG [$F$ (2,33) = 0.08, $P$ = 0.93] experiments (Fig. 2a), indicating that

the participants solved the spatial memory task accurately regardless of the target location.

**MRI contrast analysis**

The neural activity in the fMRI experiment was examined using the 4-s time-window of the

targeting period in which the participants remembered the location of the target object (i.e.,

human character) relative to their self-body in the virtual environment. A contrast analysis

showed a significantly stronger blood-oxygen-level-dependent (BOLD) signal for the

left/right-target condition than for the back-target condition in the parietal cortex, precuneus,

FEF, and SMA (Fig. 3a, $P$ < 0.01; initial threshold, $P$ < 0.05; cluster-corrected for multiple

comparisons). In contrast, no brain areas showed a significantly stronger BOLD signal for the

back-target condition than for the left/right-target condition in the same statistical condition

with multiple comparisons, although we found a significant cluster for the back-target

location in the right rhinal cortex (entorhinal cortex [ERC] and perirhinal cortex [PRC]) of

the MTL when we used a more liberal threshold (Fig. 3b, $P$ < 0.05, uncorrected).

Regarding the larger activation of the dorsal brain regions under the left/right-target

conditions than the back-target condition, one question might be whether the elevated

activities can be explained by eye movements under different target conditions. To test this

possibility, we examined the participants' gaze positions during the 4-s time window of the

targeting period. The gaze positions did not differ among the egocentric locations of the

retrieved targets in the horizontal direction [$F$ (2, 51) = 1.7, $P$ = 0.19] or the vertical direction

[$F$ (2, 51) = 0.084, $P$ = 0.92] (Fig. 2b). In addition to the average gaze position, we examined

the frequencies of saccadic eye movements (see Methods), but no significant difference was

found between the left/right and back-target conditions during the targeting period [$F$ (2, 51)

= 0.007, $P$ = 0.99] (Fig. 2c). These results indicate that the increased activity in the dorsal

brain regions was not due to eye movements that may have been induced by the retrieved

egocentric locations. Taken together, these findings reasonably suggest that the frontoparietal

cortical areas were involved in an additional mental process to represent a retrieved target to

the left/right in comparison with that behind the participants' self-body. Similarly,

remembering a target behind the self-body may employ the right MTL more than when the

participants remembered a target to their left/right.

**MRI functional-connectivity analysis**

We subsequently investigated the brain regions that interacted with the parietal cortex and

precuneus when the participants remembered the target location to their left, right, and back.

To examine the interactions, we conducted whole-brain functional connectivity analysis

using the anatomical ROIs of the parietal cortex and precuneus as the seeds based on the

AAL (see Methods for details). The results showed that, in addition to a mutual connectivity

between these brain areas, both the parietal cortex and precuneus exhibited a significantly

higher connectivity to the FEF and SMA under the left/right-target condition than under the

back-target condition (Fig. 4a, $P < 0.01$, initial threshold, $P < 0.05$, cluster-corrected for

multiple comparison). This result indicates the cooperative involvement of the FEF, SMA,

parietal cortex, and precuneus while processing a target object in the visual field, even when

the egocentric object location was not perceived. In contrast, the parietal cortex, but not the

precuneus, showed increased functional connectivity to the right ERC when a retrieved target

location was behind the participants' body (Fig. 4b, $P < 0.01$, initial threshold, $P < 0.05$,

cluster-corrected for multiple comparisons via small-volume correction using the bilateral

MTL mask). We also examined the functional connectivity using FEF and SMA seeds and

found that these two brain regions exhibited strong connectivity to the parietal cortex and

precuneus in the left/right-target condition but did not show increased functional connectivity

to any other brain regions, including the ERC, in the back-target condition. These results

suggest that our egocentric space is represented by two distinct brain networks: the

frontoparietal network for the target within the visual field, and the MTL-parietal network for

the target outside the visual field.

**MEG contrast analysis**

The fMRI experiments showed preferential involvement of the frontoparietal network in the

left/right-target condition and those of the MTL-parietal network in the back-target condition.

To examine the temporal dynamics of these brain networks, we subsequently conducted an

MEG study using the same spatial memory task that was used for the fMRI study, except for

the time parameters (Fig. 1a).

163           Figure 5a shows the results of the contrast analysis that compared the activity strength

at each sensor between the left/right-target and back-target conditions. We found a cluster of

sensors in the left-posterior area, which showed significantly stronger activity under the

left/right-target condition than under the back-target condition during 0.67–0.85 s after the

onset of the targeting period ($P < 0.05$, initial threshold, two tailed; $P = 0.04$, spatial-temporal

cluster-corrected for multiple comparison) (Fig. 5b). Conversely, no cluster of sensors

showed stronger activity under the back-target condition than under the left/right-target

condition. In addition to the fMRI study, we examined the participants' gaze positions and

frequencies of saccadic eye movements in MEG experiments. There was no significant

difference among the left, right, and back target conditions during the 1-s time-window of the

targeting period in the eye positions [$F (2, 27) = 0.32, 0.42, P = 0.73$ and $0.66$, respectively,

for the horizontal and vertical positions, respectively] (Fig. 2b) nor the saccade frequencies

[$F$ (2, 27) = 0.14, $P$ = 0.87] (Fig. 2c). These results were consistent with the results of the

fMRI experiments, indicating predominant spatial representation processing within the visual

field relative to that outside the visual field.

To examine the changes in neural activity required for the retrieval of a target

location, we compared the activity strength under back-target, left-target, and right-target

conditions with the control trials, in which a targeting-character was not presented and the

participants were not required to remember a target location (Fig. 5c). Figure 5d shows the

time course of the topographic activity map under each condition relative to that under

control conditions. Increased activity was observed in the left posterior area of the head. This

left-posterior cluster showed a significant increase in activity during 0.5–0.6 s from the onset

of the targeting period under the left-target and right-target conditions ($P < 0.05$, initial

threshold; $P < 0.05$, cluster-corrected for multiple comparison) although the same trend of

increase in activity was observed under the back-target condition [$t(11) = 2.42$, $P = 0.03$,

uncorrected]. To localize the brain regions contributing to the significant activity increase in

the left-posterior cluster, we conducted a source analysis of the MEG signal[30]. The source

powers were distributed largely in the parietal cortex and precuneus in the left- and right-

target conditions (Fig. 5e). We also found that the source power for the back-target condition

was distributed in the parietal cortex and precuneus, although the level of source power was

smaller than those under the left -and right-target conditions.

To explore the brain regions exhibiting larger neural activity under the back-target

condition than under the left/right-target condition, we conducted a whole-brain analysis to

compare the source power between the two conditions at intervals of 0.2 s after the onset of

the targeting period (Fig. 6a). We found a strong source-power for the back-target condition

in the right MTL, including the ERC, during 0.2–0.4 s after the onset of the targeting period.

Using anatomical ROIs of each hemisphere of the whole MTL, we examined the precise time

courses of the source power for the back -and left/right-target conditions relative to the

control trials. The results indicated elevations in the source power after the onset of the

targeting period under both back-target and left/right-target conditions in both hemispheres of

the MTL. However, only the right MTL exhibited a significantly larger source power for the

back-target condition than for the left/right-target condition in the early phase (0.25–0.37 s)

after the onset of the targeting period (Fig. 6b, $P < 0.05$,  initial threshold; $P < 0.05$, spatial-

temporal cluster correction for multiple comparison, two-tailed). We further examined the

source power in the right MTL by using the anatomical masks of its subregions and found

that the source power was larger under the back-target condition than under the left/right-

target condition in all the subregions [HPC: $t(11) = 3.00$, $P = 0.048$; PHC: $t(11) = 2.98$, $P =$

0.049; PRC: $t(11) = 3.22$, $P = 0.032$; ERC: $t(11) = 3.39$, $P = 0.024$, Bonferroni-corrected for

multiple comparisons (n = 4)] (Fig. 6c). Collectively, the MEG contrast analyses in source-

power between the back- and left/right-target conditions indicate that the right MTL,

including the ERC, was involved more in the back-target condition than in the left/right-

target condition in the early phase (0.25–0.37 s) after the onset of the targeting period, while

the parietal cortex and precuneus were involved more in the left/right-target condition than in

the back-target condition in the late phase (Fig. 5b, 0.67–0.85 s).

**MEG functional-connectivity analysis**

Next, we examined the connectivity of the parietal cortex with the FEF, SMA, and MTL

areas by calculating the phase lag index (PLI) for the MEG data[31-32]. We chose the two time-

windows of interest (0.08–0.48 s and 0.56–0.96 s after the onset of targeting period, see

method for details) since they could include at least three cycles of alpha-band waves and

cover the early and late phases, which were revealed from the MEG contrast analysis (Figs.

5&6). Figure 7 shows the differences in connectivity between the back- and left/right-target

conditions. The connectivity patterns differed significantly between the two time windows in

the alpha band (8–13 Hz) [$F(1, 132) = 8.24$, $P = 0.02$, repeated-measures two-way ANOVA,

Bonferroni-corrected for multiple comparisons, n = 4 for frequency bands], but not in the

other bands. During the early time window, the parietal cortex showed a larger connectivity

with the right ERC and PRC of the MTL under the back-target condition than under the

left/right-target condition, although the difference was statistically marginal [ERC: $t(11) =$

2.16, $P = 0.06$; PRC: $t(11) = 1.91$, $P = 0.08$, uncorrected]. Conversely, we found a larger

connectivity of the parietal cortex with the FEF [$t(11) = 2.61$, $P = 0.02$, uncorrected] and

SMA [$t(11) = 1.73$, $P = 0.11$, uncorrected] under the left/right-target condition than under the

back-target condition during the late time window. These results were consistent with those

obtained with functional connectivity analysis using fMRI (Fig. 4). In contrast to the time-

dependent alpha-band connectivity of the parietal cortex to the MTL and frontal areas, we did

not find a change in the connectivity of the precuneus with any of the ROIs in any frequency

band across the two time windows of interest.

# Discussion

The present combined fMRI and MEG study showed greater involvement of the fronto-parietal network when a participant remembered a target location on the left/right side than on the back of the self-body. Conversely, larger involvement of the MTL-parietal network was revealed when a target object was behind a participant. These results suggest that the parietal cortex represents the external space surrounding the self-body in mind by interacting with the dorsal frontal regions and the MTL.

A previous human fMRI study using multi-voxel pattern analysis (MVPA) reported that the parietal cortex codes the egocentric space both inside and outside the visual field[28], which is consistent with the findings of our previous MVPA-based study that employed the same spatial-memory task as the present one[29]. However, Schindler and Barteles[28] did not find the relationship between BOLD signal strength and egocentric space demonstrated by the present study. One reason for this discrepancy might be the different types of memory required for the two studies. Schindler and Barteles[28] intensively trained participants to learn a fixed spatial arrangement of eight objects for several days before the MRI scans. During the scans, the participants remembered a target object location based on the knowledge that they stored as long-term memory in a different room. In contrast, the present study prompted the participants to encode a trial-specific spatial relationship among the three objects in each trial (see Methods) and remember a target location from their short-term memory in the same 3D virtual environment as they encoded during the scan. These different task conditions may have resulted in different participant retrieval strategies between the two studies, which may have affected space representations in the parietal cortex.

The present study exhibited disproportional spatial representations around the self-body, with a bias toward the left/right-target location in comparison with the back-target location. The preferential processing of the left/right-target location relative to the back by

the frontoparietal network is consistent with the results of previous behavioral studies that

reported "front facilitation," i.e., the location of a target in the anterior part of a participant is

more efficiently detected than a target located behind them[24,25,27]. On the other hand, we

found greater involvement of the MTL-parietal network in representing the back target

location than the left/right target location. Interestingly, the present MEG experiment

comparing each target-location condition with the control condition showed that the source

power increased in the MTL not only for the back-target location but also for the left/right-

target location (Figs. 6b). These results suggest that the retrieved target location information

is transmitted from the MTL to the parietal cortex regardless of whether the location was

inside or outside the visual field. These findings are consistent with the results of previous

studies that suggested the involvement of both the MTL and parietal cortex as members of

the core brain system in the recollection of episodic memory[33,34]. We hypothesized that the

MTL-parietal network was involved in the back-target condition more than the left/right-

target condition to generate a mental representation of a retrieved target location because the

representation in the left/right-target condition was also supported by the frontoparietal

network, which might reduce the demand on the MTL-parietal network in comparison with

the back-target condition.

While the fMRI experiment suggested the preferential involvement of only the rhinal

cortex, particularly the ERC in the back-target condition, the MEG experiment suggested the

involvement of all MTL subareas. This inconsistency might be due to the ill-posed nature of

the MEG inverse problem (e.g., the "source leakage"), since a limited number of magnetic-

field sensors yielded insufficient activity to discriminate among thousands of source points,

particularly for neighboring regions[35-37]. The MEG source power in the other MTL

subregions might thus be caused by signal leakage from the ERC. Another possibility might

be that the MEG signal reflected synchronized activity at each instantaneous time point and

would be more sensitive to transient neuronal operations than fMRI analysis, which is based

on the averaged BOLD signal[38] in each TR (2 s). In either case, the ERC might play a key

role in the MTL-parietal network for the representation of a retrieved target location, which is

consistent with the results of previous studies that examined the spatial properties of the ERC

neurons (e.g., grid cells, head direction cells) in both rodents[22,39-42] and primates[43-45].

Importantly, the primate ERC reportedly represents the external space according to the gaze

position and even in imagined navigation[46]. In contrast to the ERC, other MTL subregions

may be involved in the retrieval of the target location[47-49]. Considering the high performance

of the present spatial memory task and the fact that post-scan interviews of the participants

showed no strategic efforts for retrieval, the retrieval process might be only transient in the

present experimental paradigm, which could be detected more efficiently by MEG than by

fMRI.

The MEG connectivity analysis using PLI revealed a preferential increase in

synchronization at alpha-band frequency (8–13 Hz) for both frontoparietal and MTL-parietal

networks for the left/right-target and back-target conditions, respectively. In contrast, we did

not observe a differential increase in synchronization in the other frequency bands. These

findings were consistent with the results of previous MEG and EEG studies, which suggested

functional roles of alpha-band phase synchrony in long-range communications across distant

brain regions, including the parietal cortex[8,50-53]. One remaining question concerns the

functional significance of the MTL-parietal network in daily life. In the present spatial-

memory paradigm, the participants obtained the egocentric target location from their short-

term memory, either inside or outside the visual field[29]. Instead, we usually locate a target

inside the visual field for a subsequent action by looking at it[54,55], while we occasionally

retrieve a target behind us from short-term memory. Therefore, the visual modality of

information on the back is always encoded in the past, and the preceding spatiotemporal

information is retained in the MTL[56,57]. Collectively, the parietal cortex may represent the

whole space around the self-body by coordinating the MTL-parietal network and the

frontoparietal network, which may equip us with the mental representation of the present

external world interacting with our past (retrieval) and future (action planning).

## Materials and Methods

**Participants**

Nineteen and 12 right-handed university students with normal or corrected-to-normal vision
were recruited from Peking University for fMRI and MEG experiments, respectively (fMRI:
12 women, 7 men; MEG: 4 women, 8 men). The average age of the participants recruited for
the fMRI and MEG experiments was 24.9 years (range: 18–30 years) and 22.5 years (range:
19–25 years), respectively. None of the participants had a history of psychiatric or
neurological disorders; all of them provided written informed consent prior to the start of the
experiment, which was approved by the Research Ethics Committee of Peking University.

**Experimental design**

*Experimental design:* The details of the study design have been described previously[29]. A 3D
virtual environment was programmed using the Unity platform (Unity Technologies, San
Francisco). Three animated 3D human characters (Mixamo, San Francisco,
https://www.mixamo.com) were placed on three out of four locations pseudo-randomly
across trials (Fig. 1a). Participants performed the task using a first-person perspective with a
90° field of view (aspect ratio = 4:3) and had never seen a top-down view of the virtual
environment. Experimental stimuli were presented through an LCD projector with a
resolution of 1024 × 768 pixels.

The spatial memory task included 144 and 72 trials in the MRI and MEG
experiments, respectively. In each trial, participants walked from one of the four starting
locations toward the human characters and stopped on the wood plate in the center of the
environment. After the walking period, participants experienced a "facing period" and a
"targeting period" sequentially. In the facing period, one of the human characters was

presented as a facing-character in the center of the display with the environmental

background for 2.0 s (MRI) or 1.0 s (MEG), with the other two characters being invisible. In

the targeting period, a photo of one of the remaining characters (named as the "targeting-

character") was presented as a target on a scrambled background for 2.0 s (MRI) or 1.0 s

(MEG). Each of the three periods was followed by a 2.0-s (MRI) or 1.0-s (MEG) delay (noise

screen). At the end of each trial, participants indicated the direction of the target relative to

their self-body by pressing a button when a cue was presented.

We added head-nodding detection (HND) trials (16 trials for MRI and 36 trials for

MEG) to the spatial memory task. In the HND trials, a photo of one of the human characters

was presented after the walking period, and the participants were then asked to indicate

whether the human character nodded its head during the walking period. Each human

character nodded its head at a probability of 20.6% at a random time point between the start

and end of walking. Because the trial types were indistinguishable during the walking period,

the participants were required to pay attention to the head-nodding of the human characters

during the walking period, which would reduce the possibility of voluntary memorization of

the spatial relationship of the three objects. Post-scanning interviews showed that none of the

participants made efforts to memorize the spatial relationship[29].

*MEG control conditions:* Two control conditions (36 trials for each) were added to the spatial

memory task in the MEG experiment. In the control conditions, after the walking period, a

white cross was presented instead of a human character during both the facing and targeting

periods or during the targeting period (Fig. 4c). The participants were instructed to rest with

their eyes open and fixate on the white cross. After the targeting period, the participants

pressed a button corresponding to the number presented on the screen during the response

period. The number was determined randomly from one to four in each trial of the control

conditions.

**fMRI acquisition and analysis**

*MRI scanning parameters:* BOLD MRI images were acquired using a 3T Siemens Prisma

scanner equipped with a 20-channel receiver head coil. Functional data were acquired with a

multi-band echo planar imaging sequence (TR: 2000 ms; TE: 30 ms; matrix size: 112 × 112 ×

62; flip angle: 90°; gap: 0.3 mm; resolution: 2 × 2 × 2.3 mm$^3$; number of slices: 62; slice

thickness: 2 mm; gap between slices: 0.3 mm; slice orientation: transversal). The signals of

the original voxels (i.e., 2 × 2 × 2 mm$^3$) were assigned to the corresponding voxels without a

gap (2 × 2 × 2.3 mm$^3$) to construct participants' native space images. Four experimental

sessions were conducted with 478, 476, 473, and 475 of TRs on average. A high-resolution

T1-weighted three-dimensional anatomical data set was collected to facilitate registration

(MPRAGE: TR: 2530 ms; TE: 2.98 ms; matrix size: 448 × 512 × 192; flip angle: 7°;

resolution: 0.5 × 0.5 × 1 mm$^3$; number of slices: 192; slice thickness: 1 mm; slice orientation:

sagittal).

*fMRI preprocessing:* BOLD images of each scanning session were preprocessed

independently using FSL FEAT (FMRIB's Software Library, version 6.00,

https://fsl.fmrib.ox.ac.uk/fsl/fslwiki)[58,59]. For each session, the first three functional volumes

were discarded to allow for T1 equilibration, and the remaining functional volumes were

slice-time corrected, realigned to the first image, high-pass filtered at 100 s, and smoothed

using a 5-mm FWHM Gaussian filter.

*Univariate analysis:* The 4-s BOLD signals of the targeting period were modeled using

univariate general linear models (GLM) with the three egocentric directions (left, right, and

back) included as regressors in each experimental session (40 trials) for each participant. The

BOLD signals in the other task periods were modeled by additional regressors, including 12

visual patterns for the 8-s walking period (three spatial arrangements of human characters ×

four walking directions), the four types of body rotation for the 4-s facing period (135° and

45° in the clockwise and anti-clockwise directions), the three buttons that participants pressed

during their response for the response period, the time period for the HND trials, and the six

motion parameters. A canonical hemodynamic response function was used for each task

period in GLM. This procedure generated a parameter map for each of the three egocentric

directions (left, right, and back). The parameter maps were averaged across four scanning

sessions for each egocentric direction and registered to a T1-weighted standard image

(MNI152) using FSL FLIRT[60,61] before group-level statistical analyses. The voxel size was

converted into a $2 \times 2 \times 2$ mm$^3$ resolution during the registration process.

*Connectivity analysis:* To conduct whole-brain analyses examining the functional

connectivity of the parietal cortex and precuneus, we first removed several sources of

spurious variance along with their temporal derivatives from the preprocessed functional data

by performing a GLM, which included (1) the mean time courses of the BOLD signal across

voxels within the lateral ventricles; (2) white matter; (3) whole brain; and (4) the six motion

parameters as regressors. The residual signals were bandpass-filtered, leaving signals within

the frequency range of 0.01–0.1 Hz and shifted by two TR intervals (4 s)[62-64]. The signals in

the participant's native space were registered to a T1-weighted standard image (MNI152)

using FSL FLIRT[60,61]. Then, we extracted the TRs during the 4-s targeting period from each

trial and concatenated them with those in the next trial for each of the three egocentric

directions and each session. The BOLD signals in each voxel were z-transformed along the

time course for each direction. Next, we averaged the z-transformed BOLD signals over the

anatomical masks of the brain areas (i.e., the seeds of the parietal cortex and precuneus) on

the basis of the automated anatomical labeling (AAL) template[65], which generated a regional

time course across trials for each egocentric direction in a session[66]. The regional time course

of each seed was then correlated with the time course of each voxel in the whole brain, which

yielded a whole-brain correlation map for the left-, right-, and back-target conditions

separately. The correlation maps were averaged across four scanning sessions (and left-and

right-target conditions) before submitting them to a group-level statistical analysis.

**MEG acquisition and analysis**

*MEG scanning parameters:* Neuromagnetic signals were recorded with a 275-channel whole-

head axial gradiometer DSQ-3500 MEG system (CTF MEG, Canada) at a sampling rate of

1200 Hz. A third-order synthetic gradiometer and linear drift corrections were applied to

remove far-field noise. To measure the head position within the MEG helmet, three head-

position indicator (HPI) coils were attached to the nasion and two preauricular points of each

participant to coregister their head position with the sensor coordinate system. During

scanning, customized chin-rest equipment compatible with MEG was prepared to ensure that

head movements did not exceed 2 mm. After MEG recording, each participant underwent

anatomical MRI scans on a 3T Siemens Prisma scanner (voxel size: 1 mm$^3$; flip angle: 9°;

TE: 1.97 ms; TR: 2,300 ms; field of view: $256 \times 256 \times 176$ mm$^3$), and three MRI markers

were attached to the locations of the HPI coils to align each participant's anatomical image to

the MEG sensor positions.

*MEG preprocessing:* Raw signals from the MEG experiment were preprocessed and analyzed

using the MNE Python toolbox (v0.19; available at: https://mne.tools/stable/index.html)[67,68].

The time courses of the MEG signal were bandpass-filtered between 1 and 100 Hz offline.

Artifacts (e.g., eye movements, eye blinks, and cardiac movements) were removed by

performing independent component analysis (ICA). Then, the MEG signals were visually

inspected and downsampled to 200 Hz to increase the processing speed in later analyses[69].

*Sensor space analysis:* The baseline MEG signal for each trial was calculated by averaging

the signals during the 0.2-s period preceding the onset of the targeting period, and the

baseline was subtracted from the signal at each time point during the targeting period in that

trial. Then, the time series of the MEG signal was averaged across trials for each target and

control condition. Contrast analyses were performed to compare the difference between the

left/right-target and back-target conditions, and to test the effects of the recollection in each

of the three target conditions relative to the control conditions. These analyses were

performed on each sensor, and a sensor space was generated for each session. The sensor

spaces were averaged across sessions before submitting them to the statistical test for group-

level statistical analysis.

*Source space analysis:* To reconstruct the spatial-temporal activity from the sensor space to

the anatomical space, the forward model was created using a single-compartment (inner

skull) boundary-element method on the basis of each participant's anatomical image, and the

spatial-temporal activity was then inversely modeled using the dynamic statistical parameter

map at each source point and time[30]. The source space was estimated using a subsampling

strategy, which involved subdividing a polygon (oct6) using the spherical coordinate system

provided by FreeSurfer, producing 4098 source points per hemisphere with an average source

spacing of 4.9 mm (assuming a surface area of 1000 cm$^2$/hemisphere)[67,70]. The source space

of each participant was morphed to an average surface. The percentile ranks of source-power

strength from the top 5% to 1% was calculated for each egocentric location in comparison

with the control condition or the contrast between the left/right and back conditions. For ROI

analysis, we manually delineated each of the medial temporal lobe (MTL) subareas

(hippocampus [HPC], parahippocampal cortex [PHC], perirhinal cortex [PRC], and

entorhinal cortex [ERC]) on the participant's native space using established protocols[71-74] as

well as the delineating software ITK-SNAP (www.itksnap.org). The mean source power

within the MTL subareas was calculated by averaging the source power within each mask for

each egocentric location before submitting it to a group-level statistical analysis.

*Connectivity analysis:* To assess the connectivity of the parietal cortex and precuneus with

the frontal lobe (FEF/supplementary motor area [SMA]) and MTL subareas in each temporal

period revealed by the MEG contrast analysis (i.e., "early" [0.25–0.37 s after the onset of

targeting period] for the back-target condition and "late" [0.67–0.85 s] for the left/right-target

condition), we examined the phase synchronization of the MEG time series for the following

four frequency bands: alpha (8–13 Hz), beta (13–30 Hz), low-gamma (30–60 Hz), and high-

gamma (60–99 Hz) bands. For this purpose, we first calculated the powers of the MEG signal

in a time window of 400 ms centering iteratively at each of the time points within the two

time periods. The 400-ms time window was applied in this analysis to ensure that at least

three cycles of the source time series could be covered in the alpha band[75]. The powers at

each time point were averaged across trials including all target locations and a control

condition for each participant, and the grand-mean of power across participants was

calculated. We selected the centering time point of the time window that had the maximum

grand-mean power for each time period. This procedure yielded two time-series (the time

window of 0.08–0.48 s corresponding to the centering point of 0.28 s and the time window of

0.56–0.96 s corresponding to the centering point of 0.76 s for the "early" and "late" periods,

respectively). Phase synchronization was tested among the eight ROIs (parietal cortex,

precuneus, FEF, SMA, and right MTL subareas) for each time series, condition (left, right, or

back), and frequency band using PLI[31,32] and the built-in function of the MNE Python

toolbox[67,68]. In each frequency band, we calculated the difference in the connectivity of each

time series for each ROI between the back- and left/right-target conditions before group-level

statistical analysis.

**Statistical analysis**

For the MRI univariate and connectivity analyses, an initial threshold of $P < 0.01$ was applied

to each voxel, and the reliability of clusters was tested using a nonparametric statistical

inference that did not make assumptions about the distribution of the data[22,66,67]. The test was

conducted with the FSL randomize package (version v2.9,

http://fsl.fmrib.ox.ac.uk/fsl/fslwiki/Randomise) with 5000 random sign-flips, and clusters

with a size higher than 95% of the maximal supra-threshold clusters in the permutation

distribution were then reported. Data obtained by ROI analysis of the MRI BOLD signal,

MEG source power, and MEG connectivity analysis were tested using either a $t$-test with

Bonferroni correction or repeated-measures two-way ANOVA. The MEG sensor space

analysis used either a spatial-temporal cluster permutation test or a spatial-cluster

permutation test. All statistical tests were two-sided unless otherwise noted, and significance

was determined when the corrected $P$ value was less than 0.05.

**Data and code availability**

The datasets and code supporting the current study are available from the

corresponding author (Yuji Naya, yujin@pku.edu.cn) upon reasonable request.

## References

1. Stein JF (1989) Representation of egocentric space in the posterior parietal cortex. Q J Exp Physiol 74:583–606.

2. Shelton AL, McNamara TP (1997) Multiple views of spatial memory. Psychon Bull Rev 4:102–106.

3. Burgess N (2006) Spatial memory: how egocentric and allocentric combine. Trends Cogn Sci 10:551–557.

4. Crawford JD, Henriques DY, Medendorp WP (2011) Three-dimensional transformations for goal-directed action. Annu Rev Neurosci 34:309–331.

5. Khan AZ, Blangero A, Rossetti Y, Salemme R, Luauté J, Deubel H, Schneider WX, Laverdure N, Rode G, Boisson D, Pisella L (2009) Parietal damage dissociates saccade planning from presaccadic perceptual facilitation. Cereb Cortex 19:383–387.

6. Ciaramelli E, Rosenbaum RS, Solcz S, Levine B, Moscovitch M (2010) Mental space travel: damage to posterior parietal cortex prevents egocentric navigation and reexperiencing of remote spatial memories. J Exp Psychol Learn Mem Cogn 36:619–634.

7. Iriki A, Taoka M (2012) Triadic (ecological, neural, cognitive) niche construction: a scenario of human brain evolution extrapolating tool use and language from the control of reaching actions. Philos Trans R Soc Lond B Biol Sci 367:10–23.

8. Verhagen L, Dijkerman HC, Medendorp WP, Toni I (2013) Hierarchical organization of parietofrontal circuits during goal-directed action. J Neurosci 33:6492–6503.

9. Wang X, Fung CC, Guan S, Wu S, Goldberg ME, Zhang M (2016) Perisaccadic receptive field expansion in the lateral intraparietal area. Neuron 90:400–409.

10. Zhou Y, Liu Y, Lu H, Wu S, Zhang M (2016) Neuronal representation of saccadic error

in macaque posterior parietal cortex (PPC). eLife 5:e10912.

11. Goodale MA, Milner AD (1992) Separate visual pathways for perception and action.

Trends Neurosci 15:20–25.

12. Ungerleider LG, Pessoa L (2008) What and where pathways. Scholarpedia 3:5342.

13. Stein JF (1992) The representation of egocentric space in the posterior parietal

cortex. Behav Brain Sci 15 Spec No 4:691–700.

14. Desmurget M, Epstein CM, Turner RS, Prablanc C, Alexander GE, Grafton ST (1999)

Role of the posterior parietal cortex in updating reaching movements to a visual

target. Nat Neurosci 2:563–567.

15. Donner T, Kettermann A, Diesch E, Ostendorf F, Villringer A, Brandt SA (2000)

Involvement of the human frontal eye field and multiple parietal areas in covert

visual selection during conjunction search. Eur J Neurosci 12:3407–3414.

16. Bisley JW, Mirpour K (2019) The neural instantiation of a priority map. Curr Opin

Psychol 29:108–112.

17. Vallar G (1998) Spatial hemineglect in humans. Trends Cogn Sci 2:87–97.

18. Bisiach E, Luzzatti C (1978) Unilateral neglect of representational space. Cortex

14:129–133.

19. Farrer C, Frith CD (2002) Experiencing oneself vs another person as being the cause

of an action: the neural correlates of the experience of agency. Neuroimage 15:596–

603.

20. Spiers HJ, Maguire EA (2007) A navigational guidance system in the human brain.

Hippocampus 17:618–626.

21. Gaesser B, Spreng RN, McLelland VC, Addis DR, Schacter DL (2013) Imagining the

future: evidence for a hippocampal contribution to constructive processing.

Hippocampus 23:1150–1161.

22. Chadwick MJ, Jolly AE, Amos DP, Hassabis D, Spiers HJ (2015) A goal direction signal

in the human entorhinal/subicular region. Curr Biol 25:87–92.

23. Suarez-Jimenez B, Bisby JA, Horner AJ, King JA, Pine DS, Burgess N (2018) Linked

networks for learning and expressing location-specific threat. Proc Natl Acad Sci U S

A 115:E1032–E1040.

24. Attneave F, Farrar P (1977) The visual world behind the head. Am J Psychol 90:549–

563.

25. Sholl MJ (1987) Cognitive maps as orienting schemata. J Exp Psychol Learn Mem

Cogn 13:615–628.

26. Presson CC, Hazelrigg MD (1984) Building spatial representations through primary

and secondary learning. J Exp Psychol Learn Mem Cogn 10:716–722.

27. Kelly JW, McNamara TP (2009) Facilitated pointing to remembered objects in front:

evidence for egocentric retrieval or for spatial priming? Psychon Bull Rev 16:295–

300.

28. Schindler A, Bartels A (2013) Parietal cortex codes for egocentric space beyond the

field of view. Curr Biol 23:177–182.

29. Zhang B, Naya Y (2020) Medial prefrontal cortex represents the object-based

cognitive map when remembering an egocentric target location. Cereb Cortex

30:5356–5371.

30. Dale AM, Liu AK, Fischl BR, Buckner RL, Belliveau JW, Lewine JD, Halgren E (2000)

Dynamic statistical parametric mapping: combining fMRI and MEG for high-

resolution imaging of cortical activity. Neuron 26:55–67.

31. Stam CJ, Nolte G, Daffertshofer A (2007) Phase lag index: assessment of functional

connectivity from multi channel EEG and MEG with diminished bias from common

sources. Hum Brain Mapp 28:1178–1193.

32. Colclough GL, Woolrich MW, Tewarie PK, Brookes MJ, Quinn AJ, Smith SM (2016)

How reliable are MEG resting-state connectivity metrics? Neuroimage 138:284–293.

33. Schacter DL, Addis DR, Buckner RL (2007) Remembering the past to imagine the

future: the prospective brain. Nat Rev Neurosci 8:657–661.

34. Byrne P, Becker S, Burgess N (2007) Remembering the past and imagining the future:

a neural model of spatial memory and imagery. Psychol Rev 114:340–375.

35. Grave de Peralta-Menendez RG, Gonzalez-Andino SL (1998) A critical analysis of

linear inverse solutions to the neuroelectromagnetic inverse problem. IEEE Trans Bio

Med Eng 45:440–448.

36. Sato MA, Yoshioka T, Kajihara S, Toyama K, Goda N, Doya K, Kawato M (2004)

Hierarchical Bayesian estimation for MEG inverse problem. Neuroimage 23:806–826.

37. Colclough GL, Brookes MJ, Smith SM, Woolrich MW (2015) A symmetric multivariate

leakage correction for MEG connectomes. Neuroimage 117:439–448.

38. Logothetis NK, Pauls J, Augath M, Trinath T, Oeltermann A (2001) Neurophysiological

investigation of the basis of the fMRI signal. Nature 412:150–157.

39. Hafting T, Fyhn M, Molden S, Moser MB, Moser EI (2005) Microstructure of a spatial

map in the entorhinal cortex. Nature 436:801–806.

40. Hargreaves EL, Rao G, Lee I, Knierim JJ (2005) Major dissociation between medial

and lateral entorhinal input to dorsal hippocampus. Science 308:1792–1794.

41. Howard LR, Javadi AH, Yu Y, Mill RD, Morrison LC, Knight R, Loftus MM, Staskute L,

Spiers HJ (2014) The hippocampus and entorhinal cortex encode the path and

Euclidean distances to goals during navigation. Curr Biol 24:1331–1340.

42. Wang C, Chen X, Lee H, Deshmukh SS, Yoganarasimha D, Savelli F, Knierim JJ

(2018) Egocentric coding of external items in the lateral entorhinal cortex. Science

362:945–949.

43. Doeller CF, Barry C, Burgess N (2010) Evidence for grid cells in a human memory

network. Nature 463:657–661.

44. Killian NJ, Jutras MJ, Buffalo EA (2012) A map of visual space in the primate

entorhinal cortex. Nature 491:761–764.

45. Wilming N, König P, König S, Buffalo EA (2018) Entorhinal cortex receptive fields are

modulated by spatial attention, even without movement. eLife 7:e31745.

46. Horner AJ, Bisby JA, Zotow E, Bush D, Burgess N (2016) Grid-like processing of

imagined navigation. Curr Biol 26:842–847.

47. Tambini A, Nee DE, D'Esposito M (2018) Hippocampal-targeted theta-burst

stimulation enhances associative memory formation. J Cogn Neurosci 30:1452–1472.

48. Tambini A, Davachi L (2019) Awake reactivation of prior experiences consolidates

memories and biases cognition. Trends Cogn Sci 23:876–890.

49. Yang C, Naya Y (2020) Hippocampal cells integrate past memory and present

perception for the future. PLOS Biol 18:e3000876.

50. Doesburg SM, Green JJ, McDonald JJ, Ward LM (2009) From local inhibition to long-

range integration: a functional dissociation of alpha-band synchronization across

cortical scales in visuospatial attention. Brain Res 1303:97–110.

51. Sadaghiani S, Scheeringa R, Lehongre K, Morillon B, Giraud AL, D'Esposito M,

Kleinschmidt A (2012) Alpha-band phase synchrony is related to activity in the

fronto-parietal adaptive control network. J Neurosci 32:14305–14310.

52. Takahashi T, Kitazawa S (2017) Modulation of illusory reversal in tactile temporal

order by the phase of posterior α rhythm. J Neurosci 37:5298–5308.

53. Lobier M, Palva JM, Palva S (2018) High-alpha band synchronization across frontal,

parietal and visual cortex mediates behavioral and neuronal effects of visuospatial

attention. Neuroimage 165:222–237.

54. Chen H, Naya Y (2020a) Forward processing of object–location association from the

ventral stream to medial temporal lobe in nonhuman primates. Cereb Cortex

30:1260–1271.

55. Chen H, Naya Y (2020b) Automatic encoding of a view-centered background image in

the macaque temporal lobe. Cereb Cortex 30:6270–6283.

56. Naya Y, Suzuki WA (2011) Integrating what and when across the primate medial

temporal lobe. Science 333:773–776.

57. Eichenbaum H, Cohen NJ (2014) Can we reconcile the declarative memory and

spatial navigation views on hippocampal function? Neuron 83:764–770.

58. Woolrich MW, Ripley BD, Brady M, Smith SM (2001) Temporal autocorrelation in

univariate linear modeling of FMRI data. NeuroImage 14:1370–1386.

59. Woolrich MW, Behrens TEJ, Smith SM (2004) Constrained linear basis sets for HRF

modelling using Variational Bayes. NeuroImage 21:1748–1761.

60. Jenkinson M, Smith S (2001) A global optimisation method for robust affine

registration of brain images. Med Image Anal 5:143–156.

61. Jenkinson M, Bannister P, Brady M, Smith S (2002) Improved optimization for the

robust and accurate linear registration and motion correction of brain images.

Neuroimage 17:825–841.

62. Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME (2005) The

human brain is intrinsically organized into dynamic, anticorrelated functional

networks. Proc Natl Acad Sci U S A 102:9673–9678.

63. Vincent JL, Snyder AZ, Fox MD, Shannon BJ, Andrews JR, Raichle ME, Buckner RL

(2006) Coherent spontaneous activity identifies a hippocampal-parietal memory

network. J Neurophysiol 96:3517–3531.

64. Tompary A, Davachi L (2017) Consolidation promotes the emergence of

representational overlap in the hippocampus and medial prefrontal cortex. Neuron

96:228–241.e5.

65. Rolls ET, Joliot M, Tzourio-Mazoyer N (2015) Implementation of a new parcellation of

the orbitofrontal cortex in the automated anatomical labeling atlas. Neuroimage

122:1–5.

66. Ranganath C, Heller A, Cohen MX, Brozinsky CJ, Rissman J (2005) Functional

connectivity with the hippocampus during successful memory formation.

Hippocampus 15:997–1005.

67. Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, Goj R, Jas

M, Brooks T, Parkkonen L, Hämäläinen M (2013) MEG and EEG data analysis with

MNE-Python. Front Neurosci 7:267.

68. Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, Parkkonen

L, Hämäläinen MS (2014) MNE software for processing MEG and EEG data.

Neuroimage 86:446–460.

69. Grenander U (1959) Probability and Statistics: the Harald Cramér Volume. New York,

NY: Almqvist & Wiksell.

70. Fischl B, Sereno MI, Tootell RB, Dale AM (1999) High-resolution intersubject

averaging and a coordinate system for the cortical surface. Hum Brain Mapp 8:272–

284.

71. Duvernoy HM (2005) The Human Hippocampus: Functional Anatomy, Vascularization

and Serial Sections with MRI. New York, NY: Springer Science and Business Media.

72. Insausti R, Juottonen K, Soininen H, Insausti AM, Partanen K, Vainio P, Laakso MP,

Pitkänen A (1998) MR volumetric analysis of the human entorhinal, perirhinal, and

temporopolar cortices. AJNR Am J Neuroradiol 19:659–671.

73. Pruessner JC, Li LM, Serles W, Pruessner M, Collins DL, Kabani N, Lupien S, Evans AC

(2000) Volumetry of hippocampus and amygdala with high-resolution MRI and

three-dimensional analysis software: minimizing the discrepancies between

laboratories. Cereb Cortex 10:433–442.

74. Pruessner JC, Köhler S, Crane J, Pruessner M, Lord C, Byrne A, Kabani N, Collins DL,

Evans AC (2002) Volumetry of temporopolar, perirhinal, entorhinal and

parahippocampal cortex from high-resolution MR images: considering the variability

of the collateral sulcus. Cereb Cortex 12:1342–1353.

75. Hu L, Zhang Z (2019) EEG Signal Processing and Feature Extraction. Singapore:

Springer Singapore.

76. Nichols TE, Holmes AP (2002) Nonparametric permutation tests for functional

neuroimaging: a primer with examples. Hum Brain Mapp 15:1–25.

77. Winkler AM, Ridgway GR, Webster MA, Smith SM, Nichols TE (2014) Permutation

inference for the general linear model. Neuroimage 92:381–397.

**Figures**

Figure 1. a) Spatial-memory task paradigm. Each trial consisted of four periods. Walking

period: participants walked toward three human characters using the first-person perspective

and stopped on a wood plate in the center. Facing period: one of the human characters was

presented, indicating the participant's current self-orientation. Targeting period: a photo of

the target character was presented on a scrambled background. Choice period: the

participants chose the direction of the target character relative to their body upon the

presentation of a response cue. b) The spatial relationship between the participant and the

human characters in the example trial of Fig. 1a, with the target location behind the

participant's self-body (*black circle*).

Figure 2. a) Performance of participants for the three egocentric target locations in the MRI

(n = 19) and MEG (n = 12) experiments. A statistically significant difference was not

detected in the performances of the target locations by repeated-measures one-way ANOVA

in the MRI [$F(2,54) = 0.82$, $P = 0.44$] or MEG [$F(2,33) = 0.08$, $P = 0.93$] experiments. b)

Gaze positions of the participants during the targeting period (4 s for MRI, 1 s for MEG)

were shown for each target condition. No significant difference was found among the left,

right, and back-target conditions in the MRI experiments [horizontal: $F(2, 51) = 1.7$, $P =$

0.19, vertical: $F(2, 51) = 0.08$, P = 0.92) or the MEG experiments [horizontal: $F(2, 27) =$

0.32, $P = 0.73$, vertical: $F(2, 27) = 0.42$, $P = 0.66$]. c) No significant difference in the

frequencies of saccadic eye movements was found among the three conditions during the

targeting period [MRI: $F(2, 51) = 0.01$, $P = 0.99$; MEG: $F(2, 27) = 0.14$, $P = 0.87$]. Error bars

indicate standard deviations.

720                                                        34

Figure 3. a) MRI contrast for the "left/right-target" – "back-target" condition: significant clusters were observed in the parietal cortex, precuneus, FEF, and SMA ($P < 0.01$, initial threshold, $P < 0.05$, cluster-corrected for multiple comparison). b) MRI contrast for the "back-target" – "left/right-target" condition: a cluster was observed in the right rhinal cortex, including the entorhinal cortex (ERC) and perirhinal cortex (PRC) ($P < 0.05$ uncorrected).

a

MRI connectivity contrast: Left/right - back

b

MRI connectivity contrast: back - Left/right

Figure 4. MRI connectivity analysis using the parietal cortex and precuneus as seeds. a)

Connectivity contrast for the "left/right-target" – "back-target" condition: significant

connections were found in the parietal cortex, precuneus, FEF, and SMA for both seeds ($P <$

0.01, initial threshold, $P < 0.05$, cluster-corrected for multiple comparison). b) Connectivity

contrast for the "back-target" – "left/right-target" condition: a significant connection was

observed between the parietal cortex and right ERC ($P < 0.01$, initial threshold, $P < 0.05$,

cluster-corrected for multiple comparisons via small-volume correction using the bilateral

MTL mask); no connection was found between the precuneus and MTL regions.

Figure 5. a) Mean topographic map for the MEG contrast of "left/right-target" – "back-

target" condition for every 0.2 s during the targeting period. b) Time courses of the signal

strength on the left-posterior cluster of sensors in the back-target and left/right-target

conditions. A significantly higher activity was found for the left/right-target condition

relative to the back-target condition during 0.67–0.85 s after the onset of the targeting period

($P < 0.05$, initial threshold, $P < 0.05$, spatial-temporal cluster correction for multiple

comparison, two tailed). c) A comparison between the spatial-memory-task (SMT) trials and

the control-condition trials in the MEG experiment. d) Mean topographic map for the left-

target, right-target, and back-target conditions relative to the control condition for every 0.1 s

during the targeting period. Significant clusters were found in the left-posterior from 0.5 to

0.6 s for the left-target and right-target conditions ($P < 0.05$, initial threshold; $P < 0.05$,

cluster-corrected for multiple comparison, two tailed), but not for the back-target condition.

e) Source-power distribution on the brain surface for each of the three conditions relative to

the control condition within 0.5–0.6 s. Color bar represents the percentile rank of source-

power strength.

Figure 6. a) MEG contrast for the "back-target" – "left/right-target" condition in source-power for every 0.2 s during the targeting period. The color bar represents the percentile rank of source-power strength. b) Time courses of the source-power in the anatomical ROIs for both hemispheres of the MTL. The shaded area (right MTL, 0.25–0.37 s after the onset of targeting period in the top panel) indicates a significantly higher source-power in the back-target condition than in the left/right-target condition ($P < 0.05$, spatial-temporal cluster correction for multiple comparison, two tailed). c) ROI analysis of the source-power in each of the right MTL subregions for each condition. *$P < 0.05$, $t(11) = 3.00$, 2.98, 3.22, and 3.39 for HPC, PHC, PRC, and ERC, respectively, "back" vs. "left/right," two-tailed, Bonferroni-corrected for multiple comparisons (n = 4). Error bars represent SEMs.

Figure 7. MEG connectivity contrast for the "back-target" – "left/right-target" condition

using the parietal cortex as a seed. The connectivity with each of the six anatomical ROIs was

estimated for alpha, beta, and gamma frequency bands using PLI during the early phase

(0.08–0.48 s) and late phase (0.56–0.96 s). $*P = 0.02$, $F$ (2, 132) = 8.24, a main effect of

time-windows (early vs. late), repeated-measures two-way ANOVA with brain areas as

another main effect, Bonferroni-corrected for multiple comparisons of frequency bands (n =

4). Error bars represent SEMs.