Title: Distinct brain networks coupled with the parietal cortex represent target

location inside and outside the visual field

Abbreviated title: Brain networks represent target locations in visual field

Authors: Bo Zhang [1], Fan Wang [5,6,7], Naya Yuji [1, 2, 3, 4, *]

Affiliations:

[1] School of Psychological and Cognitive Sciences, Peking University, No. 52, Haidian Road,

Haidian District, Beijing 100805, China

[2] IDG/McGovern Institute for Brain Research at Peking University, No. 52, Haidian Road,

Haidian District, Beijing 100805, China

[3] Center for Life Sciences, Peking University, No. 52, Haidian Road, Haidian District,

Beijing 100805, China

[4] Beijing Key Laboratory of Behavior and Mental Health, Peking University, No. 52, Haidian

Road, Haidian District, Beijing 100805, China

[5] State Key Laboratory of Brain and Cognitive Science, Institute of Biophysics, Chinese

Academy of Sciences, 15 Datun Road, Beijing 100101, China

[6] CAS Center for Excellence in Brain Science and Intelligence Technology

[7] University of Chinese Academy of Sciences, 19A Yuquan Road, Beijing 100049, China

Corresponding author: Yuji Naya

Address: School of Psychological and Cognitive Sciences, Peking University, No. 52,

Haidian Road, Wang Kezhen Building, Room 1707, Haidian District, Beijing 100805, China

E-mail: yujin@pku.edu.cn

Conflict of interest statement: no competing financial interests

Acknowledgments: This work was supported by the National Natural Science Foundation of

China Grant 31421003 (to Y.N.). We thank the National Center for Protein Sciences at

Peking University for assistance with MRI data collection, and the State Key Laboratory of

Brain and Cognitive Science at Chinese Academy of Sciences for assistance with MEG

scanning, which was supported in part by the Ministry of Science and Technology of China

grant (2019YFA0707103), the National Nature Science Foundation of China grant

(31730039), and the Strategic Priority Research Program of Chinese Academy of Science

(XDB32010300). Computational work was supported by resources provided by the High-

performance Computing Platform of Peking University.

Abstract

Our mental representation of the egocentric space is influenced by the disproportionate sensory perception along the ventral-dorsal (i.e., front-back) axis of the body. Previous studies typically investigated the neural architectures involved in the egocentric representations within the visual field and revealed a crucial involvement of the parietal cortex and its interaction with the frontal lobe. However, neural architectures involved in the space representation behind self-body are still unclear. To address this problem, we applied both functional magnetic resonance imaging (fMRI) and magnetoencephalography (MEG) to a 3D spatial-memory paradigm for human participants, in which the participants remembered a target location within (left or right) or outside (back) their visual field. Both fMRI and MEG experiments showed that the involvements of the fronto-parietal network including the frontal eye field and supplementary motor area were larger in the representations of retrieved target object within the visual field than outside. Conversely, the medial temporal lobe (MTL)-parietal network, including the right entorhinal cortex, was more involved in the target representation when the target was outside the visual field in both the experiments. In the MEG experiment, the connectivity increased in alpha-band frequency (8–13 Hz) in both fronto-parietal and MTL-parietal networks, and the preferential enhancement of the MTL-parietal network for the back preceded that of the fronto-parietal network for left/right. These findings suggest that the parietal cortex may represent whole space around self-body by coordinating the two distinct brain networks to process the egocentric spatial representation inside and outside the visual field.

Significance Statement (120 words maximum)

Representations of external space around self-body are necessary for coordinated actions, and our sensory-motor system is strongly biased toward processing the space in front. However,

space exists behind us too. We conducted both fMRI and MEG studies and found that the

parietal cortex coupled with the frontal lobe (FEF and SMA) and medial temporal lobe (MTL)

represents a target inside and outside the visual field, respectively, although the target

location was retrieved from short-term memory in both conditions. These results suggest that

the parietal cortex may represent the whole of the present external space by interacting (alpha

rhythm) with the frontal lobe for subsequent actions and the MTL to inquire the back scene.

Key words: egocentric direction, fMRI, MEG, parietal cortex, entorhinal cortex

Introduction

When we plan to reach for a target, it is necessary to attain its location information in the body-centered reference frame or the so-called "egocentric" spatial coordinate (Stein, 1989; Shelton and McNamara, 1997; Burgess, 2006). Previous studies using humans and nonhuman primates as subjects indicated crucial involvements of the parietal cortex in the representations of egocentric location for sensory perception, motor action, and their coordination (Khan, 2008; Ciaramelli et al., 2010; Iriki and Taoka, 2012; Wang et al., 2016; Zhou et al., 2018). Anatomically, the parietal cortex is located at the final stage of the dorsal pathway, which is often named either "where" or "how" pathway (Goodale and Milner 1992; Ungerleider et al., 2008). Perceptual signals including visual and somatosensory information thus converge on the parietal cortex, which interacts with the frontal eye field (FEF)—the region associated with attention and eye movement toward a target in the external space around the self (Stein, 1992; Desmurget et al., 1999; Donner, 2000). These anatomical connection patterns are consistent with those reported by neuropsychological studies, i.e., a damaged parietal cortex shows impaired egocentric spatial awareness known as hemispherical neglect, which leads to neglect of a target on one side of a field of vision (Vallar, 1998). Hemispherical neglect appears not only in perception but also in the memory field, known as "representational neglect" (Bisiach and Luzzatti, 1978). With accumulating human imaging studies examining spatial navigation and episodic recollection (Farrer and Frith, 2002; Spiers and Maguire, 2007; Gaesser et al., 2013; Chadwick et al., 2015; Suarez-Jimenez et al., 2018), the parietal cortex is considered to represent the egocentric space for both perception and memory.

Mnemonic representations of the egocentric space bring the question of whether the spatial representation differs between the inside and outside of a visual field. While the former can be represented in both vision and memory, the latter can be represented only in

91 memory. Previous human behavioral studies reported a decreased performance (Attneave,

92 1977) or prolonged response latency (Sholl, 1987) when participants located a target behind

93 them. This phenomenon, known as "alignment effect" (Presson, 1984) or "front facilitation"

94 (Kelly, 2009), suggests that the space surrounding our body is coded heterogeneously by

95 different neural system depending on whether a target is inside or outside the visual field

96 (Kelly, 2009). To explore the neural architectures responsible for the representation of

97 egocentric space, previous studies mostly examined the target representation within the visual

98 field; however, one previous neuroimaging study, which examined target representation

99 outside the visual field, suggested that the parietal cortex codes unseen egocentric spaces

100 (Schindler, 2013). However, the target representation of inside/outside of the visual field on

101 the parietal cortex and its interactions with associated brain areas are still unsolved.

102   To characterize neural architectures supporting the mental representations of

103 egocentric space inside and outside the visual field in the present study, we applied both

104 functional magnetic resonance imaging (fMRI) and magnetoencephalography (MEG) to a

105 spatial-memory paradigm using a 3D virtual environment for human participants, which we

106 recently devised (Zhang and Naya, 2020) (Fig. 1a). In this paradigm, participants encoded a

107 spatial relationship among three objects (walking period) in each trial. While the same three

108 objects were used across trials, their spatial relationship was different in each trial. After the

109 walking period, the participants faced one object (facing period), following a presentation of

110 a target object (targeting period). The participants remembered the location of the target

111 object relative to self-body. To-be-remembered targeting locations could be on the left, right,

112 or back position of the participants' egocentric spaces. This spatial-memory paradigm

113 allowed us to compare the neural representations of egocentric space between inside (i.e.,

114 left/right condition) and outside (i.e., back condition) of the visual field (Knapp, 1938).

Materials and Methods

Participants

Nineteen and twelve right-handed university students with normal or corrected-to-normal vision were recruited from Peking University for fMRI and MEG experiments, respectively (fMRI: 12 women, 7 men; MEG: 4 women, 8 men). The average ages of the participants recruited for fMRI and MEG experiments were 24.9 years (range: 18–30 years) and 22.5 years (range: 19–25 years), respectively. None of the participants had a history of psychiatric or neurological disorders; all of them provided written informed consent prior to the start of the experiment, which was approved by the Research Ethics Committee of Peking University.

Experimental design

Experimental design: The details of the present design were described previously (Zhang and Naya, 2020). A 3D virtual environment was programmed using Unity platform (Unity Technologies, San Francisco). In the environment, three animate 3D human characters (Mixamo, San Francisco, https://www.mixamo.com) were placed on three out of four locations pseudo-randomly across trials (Fig. 1a). Participants performed the task using first-person perspective with a 90° field of view (aspect ratio = 4:3) and had never seen a top-down view of the virtual environment. Experimental stimuli were presented through an LCD projector with a resolution of 1024 × 768 pixels.

Spatial-memory task: The task included 144 and 72 trials in the MRI and MEG experiments, respectively. In each trial, participants walked from one of four starting locations toward the human characters and stopped at the center of a wood plate. After the walking period, participants experienced a "facing period" and a "targeting period" sequentially. In the facing period, one of the human characters was presented in the center of

the display with the environmental background for 2.0 s (MRI) or 1.0 s (MEG) as a facing-

character with the other two characters being invisible. In the targeting period, a photo of one

of the remaining characters (named as "targeting-character") was presented as a target on a

scrabbled background for 2.0 s (MRI) or 1.0 s (MEG). Each of the three periods was

followed by a 2.0 s (MRI) or 1.0 s (MEG) delay (noise screen). At the end of each trial,

participants indicated the direction of the target relative to their self-body by pressing a

button when a cue was presented.

We inserted head-nodding detection (HND) trials in the spatial-memory task (16 trials

for MRI and 36 trials for MEG). In the HND trials, a photo of one of the human characters

was presented after the walking period, and then the participants were asked to indicate

whether the human character nodded its head or not during the walking period. Each human

character nodded its head in a probability of 20.6% at a random time point between the start

and end of walking in the HND trials. Because the trial-types were indistinguishable during

the walking period, the participants were required to pay attention to the head-nodding of the

human characters during the walking period, which would reduce the possibility of voluntary

memorization of the spatial relationship of the three objects. Post-scanning interviews

showed that none of the participants made efforts to memorize the spatial relationship (Zhang

and Naya, 2020).

MEG control conditions: Two control conditions (36 trials for each) were added to the

spatial-memory task in the MEG experiment. In the control conditions, after the walking

period, a white cross was presented instead of a human character during both facing and

targeting periods or during the targeting period (Fig. 4c). The participants were instructed to

rest with eyes opened and fixate on the white cross. After the targeting period, the

participants pressed the button corresponding to a number presented on the screen during the

response period. The number was determined randomly from 1 to 4 in each trial of the

control conditions.

fMRI acquisition and analysis

MRI scanning parameters: Blood-oxygen-level-dependent (BOLD) MRI images were

acquired using a 3T Siemens Prisma scanner equipped with a 20-channel receiver head coil.

Functional data were acquired with a Multi-band Echo Planer imaging (EPI) sequence (TR:

2000 ms; TE: 30 ms; matrix size: $112 \times 112 \times 62$; flip angle: 90°; gap: 0.3 mm; resolution: 2

$\times 2 \times 2.3$ mm^3; number of slices: 62; slice thickness: 2 mm; gap between slices: 0.3 mm; slice

orientation: transversal). The signals of the original voxels (i.e., $2 \times 2 \times 2$ mm^3) were

assigned to the corresponding voxels without gap ($2 \times 2 \times 2.3$ mm^3) to construct participants'

native space images. Four experimental sessions were conducted with average 478, 476, 473,

and 475 TRs. A high-resolution T1-weighted three-dimensional anatomical data set was

collected to aid registration (MPRAGE, TR: 2530 ms; TE: 2.98 ms; matrix size: $448 \times 512 \times$

192; flip angle: 7°; resolution: $0.5 \times 0.5 \times 1$ mm^3; number of slices: 192; slice thickness: 1

mm; slice orientation: sagittal).

fMRI preprocessing: BOLD images of each experimental session were preprocessed

independently using FSL FEAT (FMRIB's Software Library, version 6.00,

https://fsl.fmrib.ox.ac.uk/fsl/fslwiki; Woolrich et al., 2001; Woolrich et al., 2004). For each

session, the first three functional volumes were discarded to allow for T1 equilibration, and

the remaining functional volumes were slice-time corrected, realigned to the first image,

high-pass filtered at 100 s, and smoothed using a 5-mm FWHM Gaussian filter.

Univariate analysis: The 4-s BOLD signals of the targeting period were modeled using

univariate general linear models (GLM) with the three egocentric directions (left, right, back)

included as regressors. To remove nuisance effects, an additional twenty-six regressors were

included: twelve for modeling the visual patterns of the walking period (3 spatial

arrangements of human characters × 4 walking directions), four for the body turning-

direction (turn left or turn right) and turning-angle (135° or 45) during the facing period,

three for the response key pressed in the response period, one for HND trials, and six for

motion parameters. This procedure generated three parameter maps of egocentric directions

in participant's native space (2 × 2 × 2.3 mm voxels). For group-level statistical analysis, the

activity parameter maps were averaged across four scanning sessions and registered to a T1-

weighted standard image (MNI152) using FSL FLIRT (Jenkinson and Smith 2001; Jenkinson

et al. 2002); this procedure also resampled the voxel size into the $2 \times 2 \times 2$ mm^3 resolution

before subjecting the data to two-tailed t-test.

Connectivity analysis: To examine the functional connectivity of the parietal cortex and

precuneus inside and outside the visual field, a GLM was created to remove the nuisance

covariates from the preprocessed functional data. The signal averaged over the lateral

ventricles, white matter, and whole brain and six motion parameters, and their derivatives

were specified as regressors. The residual signals were bandpass-filtered leaving signals

within the frequency range of 0.01–0.1 Hz and shifted by two TR intervals (4 s) (Tompary

and Davachi, 2017). The residual signals were registered to a T1-weighted standard image

(MNI152) using FSL FLIRT (Jenkinson and Smith 2001; Jenkinson et al. 2002) with the

voxel size resampled to a resolution of $2 \times 2 \times 2$ mm^3. The regional time courses of targeting

period were then extracted from the anatomical masks of the parietal cortex and precuneus on

the basis of the AAL template (Rolls et al., 2015). To do this, we averaged signals over the

mask at each TR within the period and then concatenated the two TRs in a trial with those in

the next trial for each of the egocentric directions in a session (Ranganath et al., 2005, 24 TRs

in total: 2 TRs × 12 trials). The regional time course was then correlated with the time course

of each voxel in the rest of the brain, which yielded a whole-brain correlation map for inside

(left and right) and outside (back) of visual field separately. For group-level statistical

analysis, the correlation maps were averaged across four scanning sessions before subjecting

them to a two-tailed t-test.

MEG acquisition and analysis

MEG scanning parameter: Neuromagnetic signals were recorded with a 275-channel whole-

head axial gradiometer MEG system DSQ-3500 (CTF MEG, Canada) at the sampling rate of

1200 Hz. A third-order synthetic gradiometer and linear drift corrections were applied to

remove far-field noise. To measure head position within the MEG helmet, three head position

indicator (HPI) coils were attached to the nasion and two preauricular points of each

participant to coregister their head position with the sensor coordinate system. During

scanning, a customized chin-rest equipment compatible with MEG was prepared to ensure

that head movements did not exceed 2 mm. After MEG recording, each participant

underwent anatomical MRI scans on a 3T Siemens Prisma scanner (voxel size: 1 mm^3; flip

angle: 9; TE: 1.97 ms; TR: 2,300 ms; field of view: $256 \times 256 \times 176$ mm^3); three MRI

markers were attached to the same location of HPI coils to align each participant's

anatomical image to the MEG sensor positions.

MEG preprocessing: The raw MEG data were preprocessed and analyzed using the MNE

Python toolbox (v0.19; available at: https://mne.tools/stable/index.html)(Gramfort et al, 2013;

Gramfort et al, 2014). The time series were bandpass-filtered between 1 and 100 Hz offline.

To verify the data quality and remove artifacts, such as eye movements, eye blinks, and

cardiac and environmental noises, independent component analysis (ICA) was performed,

and visually inspection was applied after ICA artifact removal. The preprocessed time series

were epoched from 0.2 s before the onset of targeting period to the end of subsequent noise

period (2.2 s duration) for each of left, right, back, and control conditions; the epochs were

downsampled to 200 Hz to increase the processing speed in later steps (Grenander, 1959).

Sensor-space analysis: From the neural activity indexed by the strength of magnetic field on

sensor space, the mean activity during the 0.2-s period preceding the onset of targeting period

in each trial was subtracted. To extract the difference in neural activity between the

egocentric location inside (left and right) and outside (back) the visual field of view, a

contrast analysis was performed between the mean of left/right condition and back condition

for each sensor. Another contrast analysis was performed to examine the main effect of the

three egocentric locations relative to the control conditions. The contrasted neural activity on

sensor space was then subjected to two-tailed t-test for group-level statistical analysis.

Source space analysis: To reconstruct the spatial-temporal activity from sensor space to

anatomical space, the forward model was created using single-compartment (inner skull)

boundary-element method (BEM) on the basis of each participant's anatomical image, and

the spatial-temporal activity was then inversely modeled using the dynamic statistical

parameter map in each source point and time (Dale et al., 2000). The source space was

estimated using a subsampling strategy, which involved subdividing a polygon (oct6) using

the spherical coordinate system provided by FreeSurfer, producing 4098 source points per

hemisphere with an average source spacing of 4.9 mm (assuming a surface area of 1000

cm^2/hemisphere)(Fischl et al., 1999; Gramfort et al., 2013). Noise covariance matrix was

derived from the baseline period (-0.2 s to 0 s relative to targeting period). The source space

of each participant was morphed to an fsaverage surface before submitting to group-level

statistical analysis. The percentile rank of source-power strength from top 5% to 1% was

calculated for either each of egocentric locations or the contrast between the left/right and

back. For ROI analysis, we manually delineated each of the medial temporal lobe (MTL)

subareas (HPC, PHC, PRC, ERC) on participant's native space using established protocols

(Insausti et al., 1998; Pruessner et al., 2000; Pruessner et al., 2002; Duvernoy, 2005) as well

as the delineating software ITK-SNAP (www.itksnap.org). The mean source-power within

the MTL subareas was calculated by averaging the source-power within each mask for each

of egocentric locations before subjecting the data to group-level statistical analysis.

Connectivity analysis: To access the connectivity of the parietal cortex and precuneus with

the frontal lobe (FEF/SMA) and MTL subareas in each temporal domain revealed by the

MEG contrast analysis [i.e., 'early' (0.25-0.37 s after the onset of targeting period) for the

back and 'late' (0.67-0.85 s) for the left/right], we examined the phase synchronization of

MEG time series for the following four frequency bands: alpha (8-13 Hz), beta (13-30 Hz),

low-gamma (30-60 Hz), and high-gamma (60-99 Hz) bands. We extracted the MEG time

series using a 400 ms time windows for each of the "early" and "late" periods to ensure that

at least three cycles of source time series could be covered in alpha band (Hu and Zhang,

2019). To do so, we first calculated powers of the MEG signal using all frequency bands in

the time windows of 400 ms centering on the time points within the two time periods. We

then selected a time window that had the maximum power for each time period. This

procedure resulted in two time series (0.08-0.48 s and 0.56-0.96 s for the "early" and "late"

periods, respectively). The phase synchronization was tested among the eight ROIs (parietal

cortex, precuneus, FEF, SMA, and right MTL subareas) for each time series, condition (left,

right or back), and frequency band using phase lag index (PLI) (Stam et al., 2007; Colclough

et al., 2016) and the built-in function of MNE Python toolbox (Gramfort et al, 2013;

Gramfort et al, 2014). Subsequently, the connectivity contrast analysis ("back"–"left/right")

was performed before subjecting the data to group-level statistical analysis.

Statistical analysis

An initial threshold of $P < 0.01$ was applied to the whole-brain statistics of MRI univariate

and connectivity analyses. The reliability of clusters was tested using a nonparametric

statistical inference that does not make assumptions about the distribution of the data

(Nichols and Holmes, 2002; Winkler et al., 2014; Chadwick et al., 2015), The test was

conducted with the FSL randomize package (version v2.9,

http://fsl.fmrib.ox.ac.uk/fsl/fslwiki/Randomise), with 5000 random sign-flips; we then

reported clusters with a size higher than 95% of the maximal suprathreshold clusters in

permutation distribution. Data obtained by ROI analysis of MRI BOLD signal, MEG source-

power, and MEG connectivity analysis were tested using either repeated-measures ANOVA

or t-test. The MEG sensor space analysis used either spatial-temporal cluster permutation test

or spatial cluster permutation test. All statistical tests were two-sided unless otherwise noted

and the significance was determined according to whether the corrected P value was smaller

than 0.05.

Results

Behavioral performance in the spatial-memory task

Nineteen and twelve healthy volunteers participated in the fMRI and MEG experiments, respectively. The performances in the spatial-memory task exhibited a ceiling effect in both the experiments (MRI: 93.6% ± 1.5%; MEG: 90.4% ± 1.9%) (Fig. 1c). The performances did not differ among the three target locations (i.e., right, left, and back) in the fMRI [$F(2,54) = 0.82$, $P = 0.44$] and MEG [$F(2,33) = 0.08$, $P = 0.93$] experiments. These behavioral results indicated that the participants solved the spatial-memory task accurately in both fMRI and MEG experiments regardless of the target location in the present study.

MRI contrast analysis between left/right and back-target location

The neural activity in the fMRI experiment was examined using the 2 s time-window of the targeting period in which the participants remembered the location of a targeting object (i.e., human character) relative to their self-body in the virtual environment. We first investigated the brain regions that represented the egocentric location of the target object, when it was inside and outside the visual field, by comparing the mean BOLD signal strengths for the left and right-target locations (left/right condition) with that for the back-target location (back condition). A contrast analysis showed a significantly stronger BOLD signal for the left/right-target location than for the back-target location in the parietal cortex, precuneus, frontal eye field (FEF), and supplementary motor area (SMA) (Fig. 2a, $P < 0.01$, initial threshold, $P < 0.05$, cluster-corrected for multiple comparison). In contrast, no brain region showed a BOLD signal for the back-target location that was significantly larger than that shown by any brain region for the left/right-target location even with the use of a more liberal threshold of statistical significance ($P < 0.05$, uncorrected).

The BOLD signal during the targeting period was further examined for each of the

right, left, and back-target locations in each hemisphere of the parietal cortex, precuneus, FEF,

and SMA separately using the eight functional ROIs derived from the contrast analysis in Fig.

2a (i.e., 2 hemispheres, 4 regions, and 3 egocentric locations). The results showed significant

BOLD signal changes across the three target locations for all ROIs ($P < 0.05$, repeated-

measures one-way ANOVA, Bonferroni-corrected for multiple comparisons among the brain

regions, n = 2 × 4 = 8). In addition, for all ROIs, a significantly weaker BOLD signal was

found for the back-target condition than for either the left-target or right-target condition (Fig.

2b, $P < 0.05$, paired t-test, Bonferroni-corrected for multiple comparison among combination

of the conditions, n = 3). The elevated activities in the left and right conditions indicate that

involvements of those dorsal brain areas in processing a target in the visual field were greater

than those involved in processing the target outside of it although the participants did not

perceive the target location directly under the left/right-target (inside of the visual field) or

back-target condition (outside of the visual field). In addition to the strong activity bias for

the target inside the visual field, the left FEF and bilateral SMA showed significantly

different strengths of BOLD signals between the left-target and right-target conditions [Fig.

2b, the left FEF: $t(18) = 3.79$, $P = 0.01$, the left SMA: $t(18) = 5.50$, $P = 0.0002$, the right

SMA: $t(18) = 4.46$, $P = 0.002$, Bonferroni-corrected for multiple comparison, n = 8]. In

contrast with the FEF and SMA, neither the parietal cortex nor precuneus revealed significant

differences in BOLD signal strengths between the left-target and right-target conditions even

when a liberal threshold was used ($P < 0.05$, uncorrected). These results might reflect the

distinction in the functions of the frontal lobe (FEF and SMA) and parietal lobe (parietal

cortex and precuneus) with regard to processing egocentric information. The former might be

involved in an action planning (Tanji and Shima, 1994; Schall et al., 1995; Hanes et al., 1995;

Amador and Fried, 2004), while the latter might be associated with a spatial representation

(Colby et al., 1995; Merriam et al., 2003).

MRI functional-connectivity analysis

We subsequently investigated the brain regions that interacted with the parietal cortex and

precuneus for representing an egocentric location of a target object, when it was inside and

outside the visual field. To examine the interactions, we conducted whole-brain functional-

connectivity analysis under the left/right-target and back-target conditions by using the

anatomical ROIs of the parietal cortex and precuneus as the seeds based on the automated

anatomical labeling (AAL) (see method for details). The whole-brain functional-connectivity

analysis showed that, in addition to a mutual connectivity between these brain areas, both the

parietal cortex and precuneus exhibited a significantly higher connectivity to the FEF and

SMA under the left/right-target condition than under the back-target condition (Fig. 3a, $P <$

0.01, initial threshold, $P < 0.05$, cluster-corrected for multiple comparison). This result

indicates the preferential involvements of the FEF, SMA, parietal cortex, and precuneus for

processing a target object in the visual field even though the egocentric object location was

not perceived. In contrast, the parietal cortex but not the precuneus increased the functional

connectivity to the right ERC when a retrieved target location was beyond the visual field

(i.e., back condition) (Fig. 3b, $P < 0.01$, initial threshold, $P < 0.05$, cluster-corrected for

multiple comparison) although the ERC cluster survived only for a small volume correction

based on the mask for the MTL. We also examined the functional connectivity using the FEF

and SMA seeds and found that these two brain regions exhibited a strong connectivity to the

parietal cortex and precuneus in the left/right-target condition but did not increase functional

connectivity to any other brain regions, including the ERC, in the back-target condition.

These results suggest that a brain network consisting of the parietal cortex and ERC

represents a target object beyond the visual field, which is distinct from the other brain

network consisting of the FEF, SMA, precuneus, and parietal cortex that represents a target

object in the visual field.

MEG temporal-spatial analysis

Although the functional connectivity analysis suggests the occurrence of coupling between

the ERC and parietal cortex for the target representation outside the visual field, an increase

in BOLD signal strength was not detected in the ERC for the back-target condition. One

possible reason behind this result is that the ERC is involved in representing a target outside

the visual field more than inside for only a short duration, which might make it difficult for

the present fMRI study to detect the effect. To address this problem, we conducted a MEG

study using the same spatial-memory task that was used for the fMRI study except for the

time parameters (Fig. 1a).

Figure 4a shows the results of contrast analysis that compared the activity strength at

each sensor between the left/right-target and back-target conditions after the onset of

targeting period. We found a cluster of sensors in the left-posterior area, which showed

significantly stronger activity under the left/right-target condition than under the back-target

condition during 0.67–0.85 s after the onset of targeting period ($P < 0.05$, initial threshold,

two tailed; $P = 0.04$, spatial-temporal cluster-corrected for multiple comparison) (Fig. 4b).

Conversely, no cluster of sensors showed stronger activity under the back-target condition

than under the left/right-target condition. These MEG results were consistent with those of

the fMRI contrast analysis and suggest the predominant spatial-representation processing

within the visual field relative to that outside the visual field.

We subsequently examined the effect of egocentric target location by comparing the

activity strength under back-target, left-target, and right-target conditions with the control

trials, in which a targeting character was not presented and the participants were not required

to remember a target location (Fig. 4c). Figure 4d shows the time course of the topographic

activity map and the activity strengths under each condition relative to those under the

control conditions. An increased activity was observed on the left-posterior area of the head.

This left-posterior cluster showed a significant increase in activity during 0.5–0.6 s from the

onset of targeting period under the left-target and right-target conditions ($P < 0.05$, initial

threshold; $P < 0.05$, cluster-corrected for multiple comparison) although the same trend of

increase in activity was observed under the back-target condition [$t(11) = 2.42$, $P = 0.03$,

uncorrected]. To localize the brain regions contributing to the significant activity increase on

the left-posterior cluster, we conducted the source analysis of MEG signal (Dale et al., 2000).

The source powers were distributed largely in the parietal cortex and precuneus for the left-

target and right-target conditions (Fig. 4e). We also found the source-power for the back-

target condition to be distributed in the parietal cortex and precuneus although the level of

source-power was smaller than those under the left-target and right-target conditions.

To explore the brain regions exhibiting larger neural activity under the back-target

condition than under the left/right-target condition, we conducted a whole-brain analysis to

compare the source-power between the two conditions, every 0.2 s after the onset of targeting

period (Fig. 5a). We found a strong source-power for the back-target condition in the right

MTL including the ERC during 0.2–0.4 s after the onset of targeting period. Using

anatomical ROIs of each hemisphere of the whole MTL, we examined a precise time course

of the source-power for the back-target and left/right-target conditions separately, relative to

the control trials. The result indicated elevations in the source-power after the onset of

targeting period under both back-target and left/right-target conditions relative to the control

condition in both hemispheres of the MTL. However, only the right MTL exhibited a

significantly larger source-power for the back-target condition than for the left/right-target

condition in the early phase (0.25-0.37 s) after the onset of targeting period (Fig. 5b, $P < 0.05$,

initial threshold; $P < 0.05$, spatial-temporal cluster correction for multiple comparison, two

tailed). We further examined the source-power in the right MTL using the anatomical masks

of its subregions and found that the source-power was larger under the back-target condition

than under the left/right-target condition in all the subregions [HPC: $t(11) = 3.00$, $P = 0.048$;

PHC: $t(11) = 2.98$, $P = 0.049$; PRC: $t(11) = 3.22$, $P = 0.032$; ERC: $t(11) = 3.39$, $P = 0.024$,

Bonferroni-corrected for multiple comparisons (n = 4)] (Fig. 5c). Collectively, the MEG

source-power contrast analyses between the back-target and left/right-target conditions

indicate that the right MTL including the ERC was involved more under the back-target

condition than under the left/right-target condition in the early phase (0.25-0.37 s) after the

onset of targeting period, while the parietal cortex and precuneus were involved more under

the left/right-target condition than under the back-target condition in the late phase (0.67-0.85

s).

We examined the connectivity of the parietal cortex with the FEF, SMA, and MTL

areas by calculating PLI using the MEG data (Stam et al., 2007). We chose the two time-

windows of interests (0.08-0.48 s and 0.56-0.96 s after the onset of targeting period, see

method for details) that covered the early and late phases and were revealed from the MEG

source-power contrast analysis (Figs, 4&5), and subsequently, we included at least three

cycles of alpha-band waves. Figure 6 shows the differences in the connectivity between the

back-target and left/right-target conditions. The connectivity patterns differed significantly

between the two time-windows in alpha band (8-13 Hz) but not in other bands [(F(1, 132) =

8.24, P = 0.02, repeated-measures two-way ANOVA, Bonferroni-corrected for multiple

comparisons, n = 4 for frequency bands]. During the early time window, the parietal cortex

showed a larger connectivity with the right ERC and PRC of the MTL under the back-target

condition than under the left/right-target condition although it was statistically marginal

(ERC: $t(11) = 2.16$, $P = 0.06$; PRC: $t(11) = 1.91$, $P = 0.08$, uncorrected). Conversely, we

found a larger connectivity of the parietal cortex with the FEF ($t(11) = 2.61$, $P = 0.02$,

uncorrected) and SMA ($t(11) = 1.73$, $P = 0.11$, uncorrected) under the left/right-target

condition than under the back-target condition during the late time window. These results

were consistent with those of the functional connectivity analysis of fMRI (Fig. 3). In

contrast with that in the case of the parietal cortex, we did not find a change in the

connectivity of the precuneus with any of the ROIs in any frequency band across the two

time-windows of interests.

Discussion

The present combined fMRI and MEG study showed the larger involvements of the parietal cortex and its associated frontal areas (i.e., FEF and SMA) in the representations of target object within the visual field than outside the visual field although the egocentric spatial information was retrieved from short-term memory in both conditions and did not depend on the incoming visual sensory signal. Conversely, a larger interaction of the parietal cortex with the MTL including the ERC was revealed by both fMRI and MEG experiments, when a target object was behind a participant, although an increasing activity of the MTL was detected by only the MEG experiment with a high temporal resolution. These results suggest the two distinct networks that represent the egocentric space inside or outside the visual field (Fig. 7).

Results of both fMRI and MEG experiments exhibited disproportional spatial representations around the self-body, which showed a stronger bias toward a target object within the visual field than that behind the self-body. Preferential processing of a target location within the visual field by the fronto-parietal network may be consistent with the results of previous behavioral studies reporting "front facilitation", i.e., location of a target in front of a participant is more efficiently detected than that located behind them (Attneave, 1977; Sholl, 1987; Kelly, 2009).

One previous human fMRI study using multi-voxel pattern analysis (MVPA) reported that the parietal cortex codes the egocentric space both inside and outside of the visual field (Schindler and Barteles, 2013), which is consistent with the finding of our previous MVPA-based study that employed the same spatial-memory task as did the present one (Zhang and Naya, 2020). However, Schindler and Barteles (2013) did not find a relationship between mean BOLD signal strength and egocentric space that was demonstrated by the present study. The reason behind this discrepancy might be difference in the type of memory engaged in

locating a target object between the two studies. Schindler and Barteles (2013) intensively

trained participants for several days before the MRI scanning as they could remember the

location of the target object from the long-term memory, which stored the fixed spatial

relationship among eight objects. On the contrary, the present study prompted the participants

to encode the spatial relationship of the objects involuntarily in each trial (see Methods) and

remember the target location from the short-term memory. The distinct memory properties

between the two studies (i.e., 'long-term memory & voluntary encoding' and 'short-term

memory & involuntary encoding') may differentiate participants' retrieval strategies, which

may affect the representation of egocentric space in each study.

While the fronto-parietal network represents the target within the visual field, the

MTL-parietal network was more involved in the spatial representations outside the visual

field. However, it may also be important to note that the MEG source-power increased in the

MTL for targets both inside and outside of the visual field (Figs. 5b&c). These results suggest

the spreading of retrieved information from the MTL to parietal lobe (e.g., precuneus and

parietal cortex), when the participants remembered a target location either inside or outside of

the visual field. The spreading of memory retrieval signal from the MTL to the parietal lobe

may be supported by the early increase in the MEG source-power in the MTL (Fig. 5b) (Naya

et al., 2001). These findings are consistent with those of previous studies, which suggest

involvements of the parietal cortex and precuneus in the recollection of episodic memory as

members of the core brain system, which also includes the MTL (Schacter et al., 2007; Byrne

et al., 2007).

The preferred involvement of the MTL-parietal network in the mental representation

of the back space was supported by the fMRI connectivity analysis (Fig. 3b) in addition to the

MEG study. However, only the ERC coupled with the parietal cortex in the fMRI study while

the MEG study suggested the involvements of other MTL subregions in addition to the ERC.

This inconsistency might be due to the ill-posed nature of the MEG inverse problem (e.g. the

"source leakage"), considering that a limited number of magnetic-field sensors yielded

insufficient activity that discriminated among thousands of source points particularly for

neighboring regions (de Peralta-Menendez and Gonzalez-Andino, 1998; Sato et al., 2004;

Colclough et al., 2015). The MEG source-power in the other MTL subregions might thus be

caused by the signal leakage from the ERC. Another possibility might be that the MEG signal

reflected synchronized activity at each instantaneous time point and would be more sensitive

to the transient neuronal operation than is the fMRI connectivity analysis, which is based on

the averaged BOLD signal (Logothetis et al., 2001) in each TR (2 s). In either case, the ERC

might play a key role in the MTL-parietal network for the representation of a retrieved target

location, which is consistent with the results of previous studies that examined response

properties of the ERC neurons (e.g., grid cells, head direction cells) in both rodents (Hafting,

et al, 2005; Hargreaves et al., 2005; Howard et al. 2014; Chadwick et al. 2015; Wang et al,

2018) and primates (Doeller et al., 2010; Killian et al., 2012; Wilming et al., 2018).

Importantly, the primate ERC reportedly represents the external space according to the gaze

position and even in imagined navigation (Horner et al., 2016). In contrast to the ERC, other

MTL subregions might be involved in the retrieval of the target location (Yang and Naya,

2020). Considering the high performance of the present spatial-memory task and the post-

scan interview showing no strategic retrieval efforts by the participants, the retrieval process

might only be transient in the present experimental paradigm, which could be detected more

efficiently by MEG than by fMRI.

MEG connectivity analysis using PLI revealed the preferential increase in

synchronization at alpha-band frequency (8-13 Hz) for both fronto-parietal and MTL-parietal

networks inside and outside of the visual field, respectively. These findings were consistent

with previous MEG and EEG studies, which suggest functional roles of alpha-band phase

synchrony in long-range communications across distant brain regions including the parietal

cortex (Desburg et al., 2009; Sadaghiani et al., 2012; Takahashi and Kitazawa, 2017; Lobier

et al., 2018). One remaining question is that of the functional significance of increased

involvements of the MTL-parietal network in the representation of a retrieved target behind

the participants. In the present spatial-memory paradigm, the participants obtained the

egocentric location of a target located either inside or outside of the visual field from their

short-term memory (Zhang and Naya 2020). It should be noted here that the task required the

participants to identify the location of the target object that was present around them in the

virtual environment. In our daily life, we usually perceive and attend targets within the visual

field for the subsequent action (Ohbayashi et al., 2003 Science) while we, sometimes, retrieve

a target behind us from short-term memory, the contents of which have already passed by in

our spatiotemporal space (Naya and Suzuki, 2011; Eichenbaum, 2014; Tsao et al., 2018) (Fig.

7). We may thus combine the back scene, which we encoded just before retrieval, with the

mental representation of the present external world. This assumption suggests that the MTL-

parietal network may serve more for the representation of the back space than for that of the

front space although future studies should elucidate additional neuronal operations required

for representing the scene behind self-body. Collectively, our results indicate that the parietal

cortex may represent the whole space around self-body by coordinating the fronto-parietal

network for the target inside and the MTL-parietal network for the target outside the visual

field, corresponding to the future and past relative to the present time point.

Reference

Attneave F, Farrar P (1977) The visual world behind the head. Am J Psychol

90:549–563.

Amador N, Fried I (2004) Single-neuron activity in the human supplementary motor

area underlying preparation for action. J Neurosurg 100:250–259.

Burgess N (2006) Spatial memory: how egocentric and allocentric combine. Trends

Cogn Sci 10:551–557.

Bisiach E, Luzzatti C (1978) Unilateral neglect of representational

space. Cortex 14:129–133.

Byrne P, Becker S, Burgess N (2007) Remembering the past and imagining the

future: a neural model of spatial memory and imagery. Psychol Rev 114:340–375.

Chadwick MJ, Jolly AE, Amos DP, Hassabis D, Spiers HJ (2015) A goal direction

signal in the human entorhinal/subicular region. Curr Biol 25:87–92.

Colclough GL, Woolrich MW, Tewarie PK, Brookes MJ, Quinn AJ, Smith SM (2016)

How reliable are MEG resting-state connectivity metrics? Neuroimage 138:284–293.

Ciaramelli E, Rosenbaum RS, Solcz S, Levine B, Moscovitch M (2010) Mental space

travel: damage to posterior parietal cortex prevents egocentric navigation and

reexperiencing of remote spatial memories. J Exp Psychol Learn Mem Cogn 36:619.

Colby CL, Duhamel JR, Goldberg ME (1995) Oculocentric spatial representation in

parietal cortex. Cereb Cortex 5:470–481.

Colclough GL, Brookes MJ, Smith SM, Woolrich MW (2015) A symmetric

multivariate leakage correction for MEG connectomes. Neuroimage 117:439–448.

Dale AM, Liu AK, Fischl BR, Buckner RL, Belliveau JW, Lewine JD, Halgren E (2000)

Dynamic statistical parametric mapping: combining fMRI and MEG for high-

resolution imaging of cortical activity. Neuron 26:55–67.

Doesburg SM, Green JJ, McDonald JJ, Ward LM (2009) From local inhibition to

long-range integration: a functional dissociation of alpha-band synchronization

across cortical scales in visuospatial attention. Brain Res 1303:97–110.

Desmurget M, Epstein CM, Turner RS, Prablanc C, Alexander GE, Grafton ST (1999)

Role of the posterior parietal cortex in updating reaching movements to a visual

target. Nat Neurosci 2:563–567.

Donner T, Kettermann A, Diesch E, Ostendorf F, Villringer A, Brandt SA (2000)

Involvement of the human frontal eye field and multiple parietal areas in covert visual

selection during conjunction search. Eur J Neurosci 12:3407–3414.

Duvernoy HM (2005) The human hippocampus: functional anatomy, vascularization

and serial sections with MRI. New York, NY: Springer Science & Business Media.

de Peralta-Menendez RG, Gonzalez-Andino SL (1998) A critical analysis of linear

inverse solutions to the neuroelectromagnetic inverse problem. IEEE Trans Biomed

Eng 45:440–448.

Doeller CF, Barry C, Burgess N (2010) Evidence for grid cells in a human memory

network. Nature 463:657–661.

Eichenbaum H, Cohen NJ (2014) Can we reconcile the declarative memory and

spatial navigation views on hippocampal function? Neuron 83:764–770.

Fischl B, Sereno MI, Tootell RB, Dale AM (1999) High-resolution intersubject

averaging and a coordinate system for the cortical surface. Hum Brain Mapp 8:272–

284.

Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, Goj R,

Jas M, Brooks T, Parkkonen L, Hämäläinen M (2013) MEG and EEG data analysis

with MNE-Python. Front Neurosci 7:267.

Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C,

Parkkonen L, Hämäläinen MS (2014) MNE software for processing MEG and EEG

data. Neuroimage 86:446–460.

Goodale MA, Milner AD (1992) Separate visual pathways for perception and action.

Trends Neurosci 15:20–25.

Grenander U (1959) Probability and Statistics: The Harald Cramér Volume. New

York, NY: Almqvist & Wiksell.

Hu L, Zhang Z (2019) EEG Signal Processing and Feature Extraction. Singapore:
Springer Singapore.

Hanes DP, Thompson KG, Schall JD (1995) Relationship of presaccadic activity in

frontal eye field and supplementary eye field to saccade initiation in macaque:

Poisson spike train analysis. Exp Brain Res 103:85–96.

Hafting T, Fyhn M, Molden S, Moser MB, Moser EI (2005) Microstructure of a spatial

map in the entorhinal cortex. Nature 436:801–806.

Hargreaves EL, Rao G, Lee I, Knierim JJ (2005) Major dissociation between medial

and lateral entorhinal input to dorsal hippocampus. Science 308:1792–1794.

Horner AJ, Bisby JA, Zotow E, Bush D, Burgess N (2016) Grid-like processing of

imagined navigation. Curr Biol 26:842–847.

Insausti R, Juottonen K, Soininen H, Insausti AM, Partanen K, Vainio P, Laakso MP,

Pitkänen A (1998) MR volumetric analysis of the human entorhinal, perirhinal, and

temporopolar cortices. AJNR Am J Neuroradiol 19:659–671.

Jenkinson M, Smith S (2001) A global optimisation method for robust affine

registration of brain images. Med Image Anal 5:143–156.

Jenkinson M, Bannister P, Brady M, Smith S (2002) Improved optimization for the

robust and accurate linear registration and motion correction of brain

images. Neuroimage 17:825–841.

Knapp A (1938) An Introduction to Clinical Perimetry. Arch Ophthalmol 20:1116–

1117.

Kelly JW, McNamara TP (2009) Facilitated pointing to remembered objects in front:

Evidence for egocentric retrieval or for spatial priming? Psychon Bull Rev 16:295–

300.

Killian NJ, Jutras MJ, Buffalo EA (2012) A map of visual space in the primate

entorhinal cortex. Nature 491:761–764.

Khan AZ, Blangero A, Rossetti Y, Salemme R, Luauté J, Deubel H, Schneider WX,

Laverdure N, Rode G, Boisson D, Pisella L (2008) Parietal damage dissociates

saccade planning from presaccadic perceptual facilitation. Cereb Cortex 19:383–387.

Lobier M, Palva JM, Palva S (2018) High-alpha band synchronization across frontal,

parietal and visual cortex mediates behavioral and neuronal effects of visuospatial

attention. Neuroimage 165:222–237.

Logothetis NK, Pauls J, Augath M, Trinath T, Oeltermann A (2001)

Neurophysiological investigation of the basis of the fMRI signal. Nature 412:150–157.

Iriki A, Taoka M (2012) Triadic (ecological, neural, cognitive) niche construction: a

scenario of human brain evolution extrapolating tool use and language from the

control of reaching actions. Philos Trans R Soc Lond B Biol Sci 367:10–23.

Merriam EP, Genovese CR, Colby CL (2003) Spatial updating in human parietal

cortex. Neuron 39:361–373.

Nichols TE, Holmes AP (2002) Nonparametric permutation tests for functional

neuroimaging: a primer with examples. Hum Brain Mapp 15:1–25.

Presson CC, Hazelrigg MD (1984) Building spatial representations through primary

and secondary learning. J Exp Psychol Learn Mem Cogn 10:716–722.

Pruessner JC, Li LM, Serles W, Pruessner M, Collins DL, Kabani N, Lupien S, Evans

AC (2000) Volumetry of hippocampus and amygdala with high-resolution MRI and

three-dimensional analysis software: minimizing the discrepancies between

laboratories. Cereb Cortex 10:433–442.

Pruessner JC, Köhler S, Crane J, Pruessner M, Lord C, Byrne A, Kabani N, Collins

DL, Evans AC (2002) Volumetry of temporopolar, perirhinal, entorhinal and

parahippocampal cortex from high-resolution MR images: considering the variability

of the collateral sulcus. Cereb Cortex 12:1342–1353.

Rolls ET, Joliot M, Tzourio-Mazoyer N (2015) Implementation of a new parcellation

of the orbitofrontal cortex in the automated anatomical labeling

atlas. Neuroimage 122:1–5.

Stein JF (1989) Representation of egocentric space in the posterior parietal

cortex. Q J Exp Physiol 74:583–606.

Stein JF (1992) The representation of egocentric space in the posterior parietal

cortex. Behav Brain Sci 15:691–700.

Sadaghiani S, Scheeringa R, Lehongre K, Morillon B, Giraud AL, D'Esposito M,

Kleinschmidt A (2012) Alpha-band phase synchrony is related to activity in the

fronto-parietal adaptive control network. J Neurosci 32:14305–14310.

Sholl MJ (1987) Cognitive maps as orienting schemata. J Exp Psychol Learn Mem

Cogn 13:615–628.

Shelton AL, McNamara TP (1997) Multiple views of spatial memory. Psychon Bull

Rev 4:102–106.

Schindler A, Bartels A (2013) Parietal cortex codes for egocentric space beyond the

field of view. Curr Biol 23:177–182.

Suarez-Jimenez B, Bisby JA, Horner AJ, King JA, Pine DS, Burgess N (2018) Linked

networks for learning and expressing location-specific threat. Proc Natl Acad Sci U S

A 115:E1032–E1040.

Stam CJ, Nolte G, Daffertshofer A (2007) Phase lag index: assessment of functional

connectivity from multi channel EEG and MEG with diminished bias from common

sources. Hum Brain Mapp 28:1178–1193.

Sato MA, Yoshioka T, Kajihara S, Toyama K, Goda N, Doya K, Kawato M (2004)

Hierarchical Bayesian estimation for MEG inverse problem. Neuroimage 23:806–826.

Tanji J, Shima K (1994) Role for supplementary motor area cells in planning several

movements ahead. Nature 371:413–416.

Takahashi T, Kitazawa S (2017) Modulation of illusory reversal in tactile temporal

order by the phase of posterior α rhythm. J Neurosci 37:5298–5308.

Ungerleider LG, Haxby JV (1994) 'What' and 'where' in the human brain. Curr Opin

Neurobiol 4:157–165.

Vallar G (1998) Spatial hemineglect in humans. Trends Cogn Sci 2:87–97.

Wilming N, König P, König S, Buffalo EA (2018) Entorhinal cortex receptive fields are

modulated by spatial attention, even without movement. Elife 7:e31745.

Wang X, Fung CA, Guan S, Wu S, Goldberg ME, Zhang M (2016) Perisaccadic

receptive field expansion in the lateral intraparietal area. Neuron 90:400–409.

Yang C, Naya Y (2020) Hippocampal cells integrate past memory and present

perception for the future. BioRxiv doi: https://doi.org/10.1101/2020.03.08.982355

Zhang B, Naya Y (2020) Medial prefrontal cortex represents the object-based

cognitive map when remembering an egocentric target location. Cereb Cortex

bhaa117.

Zhou Y, Friston KJ, Zeidman P, Chen J, Li S, Razi A (2018) The hierarchical

organization of the default, dorsal attention and salience networks in adolescents

and young adults. Cereb Cortex 28:726–737.

Figure Legends

Figure 1. a) Spatial-memory task paradigm. Each trial consisted of four periods. Walking period: participants walked toward three human characters using the first-person perspective and stopped on a wood plate in the center. Facing period: one of the human characters was presented, indicating the participant's current self-orientation. Targeting period: a photo of target character was presented on the scrambled background. Choice period: the participants chose the direction of the target character relative to their body upon the presentation of a response cue. b) An example of the spatial relationship between a participant and the human characters. Left-target and right-target conditions (yellow) are inside the visual field. Back-target condition (blue) is outside the visual field. c) Performance of participants with regard to the three egocentric directions in MRI and MEG experiments. Error bar indicates SEM. n.s., no statistical significance detected by repeated-measures one-way ANOVA.

a

MRI contrast: Left/right - back

z = 42   z = 46   z = 50   z = 54   z = 58

SMA

FEF

Parietal cortex

Precuneus

b

FEF (L)

z = 52

FEF (R)

z = 56

SMA (L)

z = 48

SMA (R)

z = 54

Parietal (L)

z = 46

Parietal (R)

z = 54

Precuneus (L)

z = 56

Precuneus (R)

z = 54

Figure 2. a) MRI contrast left/right–back condition: significant clusters were revealed in the

parietal cortex, precuneus, FEF, and SMA, while no significant cluster was found for the

back-target condition relative to the left/right-target condition (cluster-corrected for multiple

comparison, initial threshold $P = 0.01$). b) BOLD signal in the functional ROIs derived from

Fig. 2a for each of the left-target, right-target, and back-target conditions. All ROIs showed

significant BOLD signal changes across the three conditions according to repeated-measures

one-way ANOVA ($P < 0.05$, Bonferroni-corrected for multiple comparisons for the ROIs, n

= 8). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, two-tailed t-test, Bonferroni-corrected for

multiple comparisons for the conditions (n = 3). Error bar indicates SEM.

a

MRI connectivity contrast: Left/right - back

b

MRI connectivity contrast: back - Left/right

Figure 3. MRI connectivity contrast of "left/right–back" and "back–left/right" conditions using the parietal cortex and precuneus as seeds. a) For the "left/right–back," condition significant connection was found in the parietal cortex, precuneus, FEF, and SMA for both seeds ($P < 0.01$, initial threshold, $P < 0.05$, cluster-corrected for multiple comparison); b) For the "back–left/right," an increased connectivity was revealed between the parietal cortex and right ERC, which survived for small volume correction based on the mask of bilateral MTL ($P < 0.01$, initial threshold, $P < 0.05$, cluster-corrected for multiple comparison); no connection was found between precuneus and MTL regions.

a

MEG contrast: Left/right - back

0-0.2 s   0.2-0.4 s   0.4-0.6 s   0.6-0.8 s   0.8-1.0 s

b

0.67-0.85 s

c

SMT (MEG)   Control (MEG)

d

MEG contrast: back, left, right - control

e

Source power in 0.5-0.6 s

Parietal cortex

Precuneus

Parietal cortex

Precuneus

Percentile

95th   99th

Figure 4. a) Mean topographic map for the MEG contrast of "left/right–back" condition for

every 0.2 s during the targeting period. b) Time courses of the signal strength on the left-

posterior cluster of sensors in the back-target and left/right-target conditions. A significantly

higher activity was found for the left/right-target condition relative to the back-target

condition 0.67–0.85 s after the onset of the targeting period ($P < 0.05$, initial threshold, $P <$

0.05, spatial-temporal cluster correction for multiple comparison, two tailed). c) A

comparison between the spatial-memory task (SMT) trials and the control trials in the MEG

experiment. d) Mean topographic map for each of the left-target, right-target, and back-target

conditions relative to the control condition for every 0.1 s during the targeting period.

Significant clusters were found in the left-posterior from 0.5 to 0.6 s for the left-target and

right-target conditions ($P < 0.05,$ initial threshold; $P < 0.05$, cluster-corrected for multiple

comparison, two tailed) but not for the back-target condition. e) Source-power distribution on

brain surface for each of the three conditions relative to the control within 0.5–0.6 s. Color

bar represents percentile rank of source-power strength.

Figure 5. a) MEG contrast of "back–left/right" condition in source-power for every 0.2 s

during the targeting period. Color bar represents percentile rank of source-power strength.. b)

ROI analysis for both hemispheres of the MTL. Shaded area in the top panel (right MTL)

show a significantly higher the source-power for the back-target condition than for the

left/right-target condition 0.25–0.37 s after the onset of targeting period ($P < 0.05$, spatial-

temporal cluster correction for multiple comparison, two tailed). c) ROI analysis of the

source-power in each of the right MTL subregions for each condition. * $P < 0.05$, t (11) =

3.00, 2.98, 3.22, and 3.39 for HPC, PHC, PRC, and ERC, respectively, two tailed,

Bonferroni-corrected for multiple comparisons (n = 4). Error bar represents SEM.

Figure 6. MEG connectivity contrast of the "back–left/right" conditions using the parietal

cortex as a seed. The connectivity with each of six anatomical ROIs was estimated for alpha,

beta, and gamma frequency bands using PLI during the early phase (0.08-0.48 s) and late

phase (0.56-0.96 s). * $P = 0.02$, $F (2, 132) = 8.24$, a main effect of time-windows (early vs.

late), repeated-measures two-way ANOVA with brain areas as another main effect,

Bonferroni-corrected for multiple comparisons of frequency bands (n = 4).

Figure 7. Two brain networks responsible for the egocentric space inside and outside of the

visual field. a) The fronto-parietal network represents target in the visual field (yellow shade).

The FEF and SMA are included in this network to represent the space in front for a

subsequent action. b) The MTL-parietal network represents a target outside the visual field

(blue shade). The right MTL, particularly ERC, is recruited in this network to represent a

target behind the self-body. In the present spatial memory task, the target location was

retrieved from the short-term memory, which a participant encoded just before retrieval

without a voluntary effort to memorize it.