Positive effect of combining high pressure and biopreservation treatments on lowering the

bacterial growth during storage of diced cooked ham with reduced nitrite salt

Stéphane Chaillou[1,§], Mihanta Ramaroson[2], Gwendoline Coeuret[1], Albert Rossero[2], Valérie Anthoine[2],

Marie Champomier-Vergès[1], Nicolas Moriceau[2], Sandrine Rezé[2], Jean-Luc Martin[3], Sandrine Guillou[2]

and Monique Zagorec[2].

[1] Université Paris-Saclay, INRAE, AgroParisTech, Micalis Institute, Jouy-en-Josas, France.

[2]INRAE, Oniris, SECALIM, Nantes, France

[3] IFIP, French Institute for the Pig and Pork Industry, Maisons-Alfort, France

§ Author for correspondence. **<u>Contact email:</u>** stephane.chaillou@inrae.fr

Keywords

Abstract

We investigated the combined effects of biopreservation and high pressure treatment on bacterial

communities of diced cooked ham prepared with diminished nitrite salt. First, bacterial communities

of four commercial brands of dice cooked ham from local supermarkets, sampled near the use-by-

date, were characterised. The four ham microbiota showed a relative low diversity but harboured

quite dissimilar communities. Two ham samples were dominated by different Proteobacteria

(*Pseudomonas*, *Serratia* for one, *Psychrobacter* and *Vibrio* for the other one) while the two others

were dominated by Firmicutes (*Latilactobacillus* and *Leuconostoc*). Second, sterile diced cooked ham,

prepared with reduced level of nitrite was inoculated with the two Proteobacteria-rich microbiota

collected from the aforementioned commercial samples together with a *Lactococcus lactis* protective

strain. Dices were then treated at 500 MPa for 5 minutes and bacterial dynamics was monitored

during storage at 8°C. Applied alone, none of the treatments stabilized durably the growth of hams

microbiota. Nevertheless, the combination of biopreservation and high pressure treatment was

efficient to reduce the growth of Proteobacteria spoilage species. However, this effect was

dependent on the nature of the initial microbiota, showing that use of biopreservation and high

pressure treatment as an alternative to nitrite reduction for ensuring cooked ham microbial safety

merits attention but still requires improvement.

1. Introduction

Nitrite salts are used since ancient times as curing agents for the production of cured meat products.

Nitrite (and eventually nitrate) salts are commonly added in the brine for the manufacturing of

cooked ham. Their role is important for the typical pinky/reddish color development of cured meats

(Alahakoon et al., 2015). Nitrite salts also participate to the hurdle technology for ensuring microbial

safety as bactericidal and bacteriostatic agents against several pathogenic bacteria occurring in meat

products, in particular *Clostridium botulinum* (Majou and Christieans, 2018). However, nitrite can

potentially lead to carcinogenic nitrosamine production in these products, which rises the concern of

a potential public health risk (Sindelar and Milkowski, 2012). In European Union (EU) all additives

authorized before the 20th of January 2009 have to be re-evaluated by 2020. In that context, the

European Food Safety Authority (EFSA) re-assessed the safety of nitrite and nitrate salts in 2017. The

current acceptable daily intakes for nitrite and nitrate was assessed and experts stated that

exposure to nitrites used as food additives was within safe levels for most of the population, except

for children (EFSA, 2017). Nevertheless, although experts evaluated the need for further scientific

information, a positive link could be established between dietary intake of nitrite (or of processed

meat containing both nitrite and nitrate) with some cancers (Mortensen et al., 2017a, 2017b).

Therefore, reducing nitrite levels in processed meats appears necessary in order to limit this sanitary

issue. Several hurdle technologies can be proposed in order to compensate a higher risk of microbial

safety in cooked ham with reduced nitrite salts. Among those, High Pressure Processing (HPP) has

been proposed in context of food additive reduction (Pottier et al., 2017) and was shown to

efficiently reduce bacterial contaminants in cooked ham (Han et al., 2011; Pietrasik et al., 2016;

Pingen et al., 2016). As well, biopreservation, a method using protective cultures, often lactic acid

bacteria (LAB) or their metabolites, was described more than 20 years ago for fighting undesired

bacterial contaminants (Stiles, 1996). Addition of such LAB protective cultures to cooked meat

products, and in particular cooked ham was indeed reported (Vermeiren et al., 2006, 2004). HPP

combined to biopreservation has also been studied (Oliveira et al., 2015). HPP treatment of cooked

ham treated with bacteriocins has been shown to reduce the population of several bacterial

pathogens or to prolong cold storage (Jofré et al., 2008a, 2008b; Liu et al., 2012; Marcos et al., 2008;

Teixeira et al., 2018). In the present study, we investigated the combined effect of HPP and

biopreservation on the dynamics of bacterial communities of cooked ham with reduced level of

nitrite salts. We had previously selected a *Lactococcus lactis* strain able to recover after HPP

treatment and producing the bacteriocin nisin (Ramaroson et al., 2018). Here, we first determined

and collected bacterial communities present on commercial diced cooked ham. Then the dynamics of

these bacterial communities was monitored, following their inoculation in sterile diced cooked ham

with reduced nitrite level, in the presence of the bioprotective strain, and after the application of the

HPP treatment.

2. Materials and methods

2.1. Bacterial strains, media and growth conditions

Lactococcus lactis CH-HP15 (Ramaroson et al., 2018) was first cultivated on M17 agar plates (Biokar

diagnostic, Beauvais, France) at 30°C for 72 h. One colony was inoculated for preculture in Medium

Modelling Ham (MMH) (Ramaroson et al., 2018, Modugno et al., 2019) at 30°C for 24 h, and then

cultured in MMH at 30°C under agitation at 65 rpm for 13h until early stationary phase. Bacterial

enumerations were performed by plating serial dilutions of bacterial cultures, microbiota or ham

stomached samples. The total viable counts were determined on Plate Count Agar (PCA) (Biokar

diagnostic, Beauvais, France). Psychrophilic and mesophilic counts were determined after incubation

for 4 days at 15°C or 24 h at 30°C, respectively. LAB were enumerated on Man Ragosa and Sharpe

(MRS) agar medium pH 5.2 (AES, France) containing bromocresol green (25 mg·l^{-1}) as previously

described after 4 days incubation at 25 °C (Najjari et al., 2008). In ham samples inoculated with *L.*

lactis CH-HP15, M17 agar plates were used for enumeration after 48 h incubation at 30°C. For

anaerobic conditions, plates were incubated in jars with anaerocult sachets (Anaerocult A, Merck,

Germany).

2.2. Ham sampling and microbiota recovery

Diced cooked ham (except for one sample consisting of sliced cooked ham) were purchased in 2015

and 2016 from local supermarkets in Nantes, France, transported at cold temperature to the

laboratory and reconditioned immediately. These samples, sold as ready-to-eat food were

conditioned in packs ranging from 120 to 200 g under modified atmosphere or without any indication

of any gas mixture used in the packs. At arrival in the laboratory (day 1), bags were opened and dices

were immediately reconditioned as 25 g aliquots under air or vacuum packaging and incubated at

4°C. Total counts and LAB counts were enumerated at day 1, 7, 14, 21, and 28. When PCA mesophilic

counts reached about 7 $\log_{10}$ CFU per gram of ham, bacterial communities were collected by mixing

25 g diced cooked ham in 75 ml peptone salt (AES, France) for 2 min in a stomacher (Masticator, IUL

Instruments, England). The homogenate was filtered through the bag filter and centrifuged through a

filter from Nucleospin Plant II Midi kit (Macherey Nagel, EURL, France) at 8000 × g during 10 min at

room temperature. The bacterial pellet was resuspended in 30 ml peptone salt and aliquoted as 1 ml

with glycerol 15% and stored at -80°C. In total, eight microbiota were recovered from ham dices

either conditioned under air or vacuum.

2.3. Challenge tests

Low-contaminated diced cooked ham containing 18 g/kg sodium chloride and reduced level of nitrite

(25 mg/kg; recommended max. level is 120 mg/kg) were used for challenge tests. Those were

produced as previously described (Ramaroson et al., 2018). Briefly, pork ham muscles were defatted,

trimmed and minced using a 20 mm grid. One kilogram of this grinding was mixed under vacuum

with 100 g of brine containing water 74.1 g, nitrite salt 4.6 g (0.6%), sodium chloride salt 15.2 g,

sodium ascorbate 0.6 g, and dextrose 5.5 g. The mix was melded and vacuum-packed. Hams (2.5 kg)

were cooked in a 100% humidity atmosphere (90 min at 55 °C, 60 min at 60 °C and 235 min at 67 °C),

cooled at ambient temperature (18 °C) for 20 min and then stored at 3 °C. Ham cubes of 1 cm × 1 cm

were prepared aseptically, aliquoted in 100 g portions, and stored vacuum-packed at −20 °C until

use. Prior to microbiota inoculation, cooked ham dices were defrosted at 4 °C for 24 h and first

inoculated with microbiota. Two different microbiota were inoculated in low-contaminated diced

cooked ham. Each challenge test was performed twice (two independent experiments). Microbiota

previously collected were defrosted rapidly at room temperature, diluted to a final concentration of

6 log CFU.mL^{-1}, inoculated at 4 log CFU g^{-1} in diced cooked ham (1% v/w inoculation rate), stored

overnight at 4°C for allowing bacterial recovery, and then inoculated with a fresh culture of *L. lactis*

CH-HP15. *L. lactis* culture in early stationary phase was centrifuged. The bacterial pellet was rinsed in

a sterile solution of NaCl 0.9% and resuspended at 9.3 log CFU g^{-1} and 0.5 ml were inoculated in 100

g dice cooked ham for an initial concentration of 7 log CFU g^{-1}. Diced cooked ham inoculated with

microbiota but without *L. lactis* were also used as controls.

2.4. HP treatments

Cooked ham dices were aliquoted as 20 g portions, vacuum-packed and High-Pressure treated at

500 MPa for 5 min. Samples were treated using high pressure in a 3 L vertical high pressure pilot unit

(ACB, Nantes, France) equipped with a temperature regulator device. The pressure transmitting fluid

used was distilled water. The samples were inserted into the vessel (whose internal temperature was

regulated to 20 °C) and processed at a compression rate of 3.4 MPa/s until the targeted pressure was

reached. The pressure level was held for 5 min, and decompression was nearly instantaneous (less

than 2 s). Water temperature reached 24 °C at 500 MPa because of the adiabatic heating. Once

treated, packs were then stored at 8°C and one portion was used at day 1, 5, 12, 30 for bacterial

enumeration and bacterial pellet collection for further DNA extraction. Unpressurized samples were

also included as controls.

2.5. DNA preparation and amplicon sequencing

Total genomic DNA was extracted from the bacterial pellet as described previously (Poirier et al.,

2018) using the PowerFood™ Microbial DNA Isolation kit (MoBio Laboratories Inc., Carlsbad, USA)

and the High Pure PCR Template Preparation kit (Roche Diagnostics Ltd, Burgess Hill, West Sussex,

UK). For each sample, both DNA extracts were pooled. Then, this DNA sample was used as template

for three independent amplifications using either the 16S V3-V4 region of the rRNA encoding gene or

an internal 280 bp fragment of the Gyrase B subunit encoding gene *gyrB*, as described previously

(Poirier et al., 2018). All PCRs were performed in triplicate. Replicates were pooled and the amplified

DNA was purified with a QIAquick kit (Qiagen, Hilden, Germany). Amplicon size, quality, and quantity

were checked on a DNA1000 chip (Agilent Technologies, Paris, France). The MiSeq Reagent Kit v2

(2x250 paired-end reads, 15 Gb output) was used according to the manufacturer's instructions for

library preparation and sequencing on a MiSeq 2 (Illumina, San Diego, USA). The quality of the

obtained sequences was checked with FastQ files generated at the end of the run (MiSeq Reporter

Software, Illumina, USA) and additional PhiX Control. The corresponding pairs of sequences were

then attributed to their respective samples using the individual multiplexing barcodes. Quality

controls indicated a PHRED quality score of at least 30 for 94% of the reads and a median number of

65,520 ± 14,180 reads per sample and 95,860 ± 11,300 reads per sample for 16S rDNA and *gyrB*

amplicons, respectively.

2.6. Operational Taxonomic Unit analysis and accession numbers.

For each sample, paired-end sequences were then loaded in the FROGS (Find Rapidly OTUs with Galaxy

Solutions) pipeline (Escudié et al., 2018), checked for quality and assembled. We retained merged

sequences with a size of 280 ± 50 bp for *gyrB* and 450 ± 50 bp for 16S rRNA gene. The merged

(assembled) sequences have been deposited in the Sequence Read Archive database under the

accession numbers SAMN13761947 to SAMN13762161, corresponding to BioProject PRJNA599607.

SWARM clustering (Mahé et al., 2014) was applied on the assembled sequences using a maximal

aggregation distance of three nucleotides for 16S rRNA gene sequences; for *gyrB* sequences, clustering

was more stringent, with a maximal aggregation distance of two nucleotides in order to potentially

assign OTUs to the subspecies level. After chimeras and spurious OTUs (low-abundance and low

frequency OTUs arising from sequencing artifacts) removal as described previously (Poirier et al.,

2018), the dataset comprised 69 OTUs for the 16S rDNA dataset and 252 OTUs for the *gyrB* dataset.

Taxonomic assignment of 16S rDNA OTUs was performed with the Blastn+ algorithm (Camacho et al.,

2009) on the SILVA 128 SSU database (Quast et al., 2012), using a threshold of 97% identity for species

assignment. For *gyrB* OTUs, it has been previously demonstrated that PCR amplification of *gyrB* can

also recover the paralogous *parE* gene, which encodes the β subunit of DNA topoisomerase IV. In order

to determine which of the two genes had been amplified for each species, the OTU sequences were

blasted against the *gyrB/parE* databases established by Poirier and colleagues (Poirier et al., 2018).

Both *gyrB* and *parE* OTUs were retained in our diversity analysis. We assigned taxonomy to the species

level when 95% of a sequence matched over 90% of length coverage found in the database. The last

step consisted in comparing the taxonomic assignments obtained from the 16S rDNA and the *gyrB*

amplicon sequencing. The taxonomic assignment of the 16S rDNA-based OTUs were then

homogenized and improved by those obtained from the *gyrB* analysis. This strategy was particularly

applied to 16S rRNA gene OTUs with no clear assignment to the species-level (several species included

at the threshold of 97% identity).

2.7. **Beta-diversity and statistical analysis**.

Bacterial diversity analyses were performed using the phyloseq package (v1.24.2) of R (McMurdie

and Holmes, 2013). For each of the samples, analysis of diversity was either performed at the OTU

level or at the species level. In studies at the species level, OTUs with similar taxonomic assignment

(both *gyrB* and 16S rDNA) were merged using the TAX_GLOM function of phyloseq package, and their

abundance (number of reads) was averaged. This resulted into a dataset of 56 species with different

taxonomic assignment. Similarly, comparative analysis between ham samples with different

experimental conditions was performed after technical (16S rDNA/*gyrB*) and biological (biological

repetition) data were merged with the phyloseq function MERGE_SAMPLES. In this case, species

abundance was averaged between the four replicates. Bacterial diversity was compared among

different groups of samples with permutational ANOVA, specifically using the adonis function within

the vegan package (Oksanen et al., 2019).

3.   Results

3.1. **Diced cooked ham selection for microbiota recovery.**

Preliminary tests were performed on diced cooked ham of different brands collected from local

supermarkets, as close as possible to the production date. These were then incubated under vacuum

or air packaging at 4°C or 8°C. We observed a large variation of initial total viable counts between

samples, ranging from 2 to 7 log CFU. g^{-1}. No strong influence of packaging was observed on

psychrophilic and mesophilic total counts, or MRS counts. Enumeration on MRS plates containing

bromocresol green enabled us to estimate the putative diversity of LAB species present in dices.

Only one to 3 types of colonies were detected indicating a poor diversity among the most dominant

LAB species.

Since our objective was to recover microbiota for further re-inoculation on cooked diced ham before

HPP treatment, we aimed at collecting standardized quantities of bacteria, as diverse as possible, and

enough DNA for further amplicon sequencing. For that purpose, we designed the sampling as

follows: i) diced cooked ham were sampled from supermarkets as close as possible to their

production date; ii) immediately after arrival in the laboratory dices were reconditioned under

vacuum packaging and under air to eventually increase the bacterial diversity to be collected; iii)

packs were then stored at 4°C and analysed every 7 days; iv) when total viable counts reached about

7 log CFU. g^{-1}, microbiota were collected and stored for further analyses and DNA was extracted for

amplicon sequencing. Through this strategy, diced cooked ham samples from four different brands

out of ten (further referred as HAM_A to HAM_D samples) were obtained with enough bacterial

diversity and DNA quality (Table 1). No significant difference in population level was observed

between air and vacuum packaging, except in HAM_B for which both PCA and MRS counts were

about 1 log higher (respectively 7.97 *vs* 6.53 log CFU g^{-1} and 7.74 *vs* 6.82 log CFU g^{-1}) after storage

under air.

3.2. The four types (brands) of diced cooked ham can be distinguished by the diversity of their

bacterial communities.

The bacterial diversity was estimated using a combination of two amplicon sequencing strategies,

one using V3-V4 region of the 16S rRNA gene and another one using an internal ~280 bp fragment of

the gyrase B subunit encoding gene (*gyrB*) in order to improve taxonomic assignment of the OTUs to

the species or sub-species level as described previously (Poirier et al., 2018). These two strategies

were also used as technical replicates for ensuring that no biases were obtained in the

characterization of the four different microbiota as observed in Figure 1A. Only slight differences

between both strategies were observed in the clustering of air samples of HAM_C and D. The

average number of species detected per sample was 38 ± 5. As shown in Figure 1B, HAM_C and

HAM_D samples showed a highly similar microbiota dominated by Firmicutes, in particular

Latilactobacillus sakei and *Leuconostoc carnosum*, two species commonly found on this type of cured

meat products (Raimondi et al., 2019; Vasilopoulos et al., 2010; Zagdoun et al., 2019). Furthermore,

storage type under vacuum or air packaging did not influence significantly the overall abundance of

the identified species. On the other hand, the bacterial communities characterized from samples of

HAM_A and HAM_B were significantly different from each other and from the aforementioned

products, in particular with a clear dominance of Proteobacteria over Firmicutes (Figure 1C). Ham_A

microbiota was mainly composed of *Pseudomonadaceae* and *Enterobacteriaceae* with *Pseudomonas*

lundensis and *Serratia grimesii* being the most abundant species for Proteobacteria and

Carnobacterium maltaromaticum being the most abundant species for Firmicutes. Samples from

Ham_B revealed to contain the most original microbiota among the four types of ham with the three

most abundant 16S rDNA or *gyrB* OTUs being assigned to unknown species as the identity

percentage of these OTUs to *gyrB* gene in databases was lower to the ANI (Average Nucleotide Index)

threshold of 95% classically used for bacterial species boundary (Qin et al., 2014). The most

abundant Firmicutes *gyrB* OTU was assigned to a *Carnobacterium sp.* showing only 89.0% identity (as

best match) to *gyrB* gene of *Carnobacterium funditum* strain DSM 5970. Among Proteobacteria, one

OTU was assigned to a *Psychrobacter sp.* with closest match was 89.9% to *gyrB* gene of

Psychrobacter cryohalolentis FDAARGOS_308, a strain isolated from Siberian permafrost (GenBank:

GCF_002208775.2). For the second Proteobacteria OTU, the best identity score found was 91.6 %

with *Vibrio sp. SM1977 gyrB* gene, a species isolated from coralline algae surface (GenBank:

CP045699.1). It should be noted that the abundance of these two uncharacterized species

(*Psychrobacter sp.* and *Vibrio sp.*) varied significantly according to the packaging type used for

storage, which may correlate with their 10 times higher counts on air stored samples (Table 1),

although we have no evidence for their cultivability in our growth conditions.

3.3. Combined effect of HPP and biopreservation on bacterial community dynamics.

For the further steps of our analysis dedicated to monitoring the effect of HPP and biopreservation

by *L. lactis* CH-HP15 on the ham microbiota dynamics during storage at 8°C, we decided to focus on

the two different microbiota from HAM_A and HAM_B. This was based on the fact that these two

microbiota showed a community structure enriched in bacterial species with highly spoilage

potential (Proteobacteria) in comparison to microbiotas of HAM_C and HAM_D enriched in

Latilactobacillus; those latter species being generally rather considered as positive microbial

component (Vermeiren et al., 2006, 2004). In addition, we chose HAM_B microbiota because of the

unexpected presence of the two uncharacterized *Psychrobacter* and *Vibrio* species. The experimental

strategy was based on re-inoculation at the level of 4 log CFU. g^{-1} of the original microbiota from both

HAM_A and HAM_B on minimally contaminated diced cooked ham with reduced nitrite content (25

mg/kg). For both types of microbiota, an equimolar amount mixture of microbiota from air and

vacuum storage previously stored at -80°C was used. The *L. lactis* CH-HP15 bioprotective strain was

inoculated at a higher level of 7 log CFU. g^{-1} to enable regrowth and ensure efficient biopreservation

activity taking into account a possible high inactivation following the HPP treatment (Ramaroson et

al., 2018). Table 2 is summarising the estimated bacterial population level at different storage times

after the several processing treatments. Similarly, Figure 2 is depicting the relative abundance of

each species via 16S rDNA and *gyrB* amplicon sequencing.

The control samples without any processing treatment revealed that at day 1 after inoculation, the

microbial communities were similar to the microbiota from HAM_A and HAM_B analyzed in Figure 1,

showing that the dices prepared for the challenge-tests were indeed poorly contaminated and that

the microbiota had recovered with no strong bias from freezing. HAM_A microbiota grew rapidly

under vacuum from ~4 log CFU. g^{-1} to reach ~7 log CFU. g^{-1} at day 5 and ~8 log CFU. g^{-1} at day 30. In

contrast, HAM_B microbiota showed a more reduced growth dynamics, reaching at most ~7 log CFU.

g^{-1} during the whole storage period. Interestingly, the structure of the communities also changed

during storage: in HAM_A samples, *P. lundensis*, a species abundant at the beginning of storage was

progressively overgrown by *S. grimesii* whereas, *C. maltaromaticum,* the third abundant species,

remained at a stable proportion within the community; in HAM_B, the dominant *Vibrio sp.* was also

overgrown over time by *Lactococcus piscium*, although this latter species was largely subdominant

(<0.1%) in the initial microbiota.

Processing with biopreservation by *L. lactis* CH-HP15 led to two different effects regarding samples

from HAM_A and HAM_B. In HAM_A samples, the addition of *L. lactis* had a strong reduction effect

on the abundance of Firmicutes species, in particular on *C. maltaromaticum*, which thus became

largely subdominant, whereas *B. thermosphacta* was temporally (up to 12 days of storage) and

presumably taking benefit from this change (figure 2A)*.* Meanwhile, the inoculation of *L. lactis* had no

obvious effect on Proteobacteria species whose pattern was unchanged overtime compared to

control. Indeed, after 30 days of storage, while the microbiota had reached a population level above

8.5 log CFU. g^{-1}, *P. lundensis* and *S. grimesii* species displayed a relative abundance of about 50% of

the microbiota; the remaining part of the microbiota being covered by *L. lactis,* although this strain

had been inoculated thousand-fold higher than the original microbiota. In HAM_B, the growth

dynamics of the dominant *Vibrio sp.* was also not affected in comparison to the untreated samples.

However, the addition of *L. lactis* induced a possible competition with *L. piscium*, resulting over time

into equilibrium of both species. More generally, *L. lactis* CH-HP15 was less competitive in microbiota

from HAM_B samples than in microbiota from HAM_A samples.

Whether or not biopreservation was applied, HPP treatment resulted in major changes in the growth

dynamics of both microbiota. HPP treatment alone on HAM_A samples reduced drastically the

viability of the original microbiota which remained below 4 log CFU. g^{-1} after 30 days of storage

(Table 2). It is likely that bacterial cells were severely injured, indeed no DNA could be recovered

from the bacterial pellet despite the use of several extraction protocol trials. Therefore, diversity

analysis by amplicon sequencing could not be performed. Nevertheless, samples treated with both

HPP and inoculation by *L. lactis* enabled better recovery of DNA and could provide an overview of the

HPP treatment on the HAM_A original microbiota.

Proteobacteria species were the most impacted by high pressure treatment, whereas *L. Lactis* CH-

HP15 could recover rapidly from this treatment reaching progressively a population level above 7.5

log CFU. g^{-1} after 30 days, and almost a complete domination of the microbiota. Interestingly, as

described above on samples only treated with *L. lactis*, the addition of the bioprotective

microorganism increased temporally (up to 12 days of storage) the relative proportion of *B.*

thermosphacta, thereby indicating that this latter species is quite resistant to HPP treatment. Unlike

microbiota from HAM_A samples, that from HAM_B samples recovered rapidly from HPP treatment

alone, almost reaching the population level of untreated samples at 30 days (6 log CFU.g in average).

The community structure was itself barely affected by the HPP treatment with only a higher

proportion of *Psychrobacter sp.* at longer storage time in comparison to untreated samples.

Processing with both HPP and *L. lactis* inoculation confirmed these observations as the population

level, combining both initial microbiota and *L. lactis* , recovered rapidly (12 days) to 7.0 log CFU. g^{-1}

and raised further to ~10 log CFU. g^{-1} at 30 days, which is almost 4 log higher than the level in

untreated ham. However, the HPP treatment was more favorable to Firmicutes species in the end,

leading to a domination of *L. lactis* CH-HP15 together with *B. thermosphacta.*

4.   Discussion

The use of an appropriate HPP treatment to preserve perishable food has been the focus of many

studies and reviews (Oliveira et al., 2015; Yamamoto, 2017). HPP is being used to enhance food

safety by reducing the microbial development in final products, meanwhile maintaining at best their

nutritional and sensorial values to a level acceptable to consumers (Pietrasik et al., 2016). However,

the effect of HPP on microbial growth dynamic and community structures has not been widely

studied and it is not known whether hurdle strategies should be applied in combination with HPP to

efficient microbiota inactivation or stabilisation. Indeed, a recent work carried out with a simplified

ham microbiota (five species including *Listeria monocytogenes, L. sakei, B. thermosphacta, C.*

maltaromaticum and *Leuconostoc gelidum*) showed that HPP treatment is not sufficient to inhibit

growth recovery of the microbiota over long storage time (Teixeira et al., 2018). This is pointing out

that there is a clear gap in our knowledge on the HPP efficiency towards various microbial

communities which may be present on cooked ham. Although the technology could be a very

promising strategy to improve the safety of nitrite-reduced cooked ham, it is necessary to investigate

whether it should be used as hurdle with the combination of other strategies such as biopreservation

(Ramaroson et al., 2018).

In this context, our study provides data in link with the following questions: does HPP display a

generic effect on microbial inactivation or is this inactivation a microbiota-dependent phenomenon?

Is the use of nisin-producing *L. lactis* mediated biopreservation a worth hurdle strategy when

combined with HPP, and, is this biopreservation effect also microbiota dependent?

Our first strategy was to recover microbiota from several types of cooked ham (different brands)

stored under air or vacuum. As previously demonstrated by Raimondi and colleagues (Raimondi et

al., 2019), the microbiota of the four types of cooked ham was poorly diverse but highly variable.

Some cooked hams with poor diversity were dominated by *L. sakei, Latilactobacillus curvatus* and *L.*

carnosum and their microbiota were not considered further as HPP treatments were previously

demonstrated to be efficient against the growth of these species (Han et al., 2011). On the other

hand, cooked hams from two other producers were characterized by a more diverse microbiota.

These microbiota were composed of a quite different mixture of species from Firmicutes and

Proteobacteria phyla, including not yet characterized dominant species. Therefore, these two types

of microbiota were chosen as model microbiota to test the efficiency of HPP and biopreservation.

Firstly, our results demonstrate that the reduction of viable cells population by HPP is slightly more

effective on Proteobacteria species than on Firmicutes species resulting in a small shift of the

dominant population from Proteobacteria to Firmicutes when HPP is applied. This finding

corroborates many previous observations made by comparing individual species of Gram-negative

and Gram-positive pathogenic bacteria resistance to HPP for instance (see Oliveira et al., 2015 for a

review). Although the cell surface morphology of both types of bacteria can explain such difference,

it is interesting to notice that the recovery of cells after HPP is species (and *per se* microbiota)

dependent. Although both microbiota from HAM_A and HAM_B were re-inoculated at the same

population level before HPP treatment, that from HAM_A, composed of *P. lundensis* and *S. grimesii*

revealed more sensitive (no or almost no recovery) to HPP than that from HAM_B composed of

uncharacterised *Psychrobacter* and *Vibrio* species. The species or even strain-dependent resistance

to HPP has already been pointed out for Firmicutes, in particular for *L. monocytogenes* and

Staphylococcus aureus (Jofré et al., 2008b). The underlying mechanisms are not fully understood as

HPP resistance and recovery of bacteria also depend on the cell physiology of bacteria present on the

food matrix before the treatment. As well, the matrix composition may influence bacterial recovery

(Tassou et al., 2007), as was shown for fat content influencing *L. monocytogenes* recovery from

thermal inactivation (Verheyen et al., 2020). Furthermore, the capacity of the bacteria to resist HPP

treatment should be dissociated from the ability of the bacterial cells to recover and thrive during

the storage conditions. Our data are a good illustration of this phenomenon. For instance, *S. grimesii*

revealed more competitive than *P. lundensis* to grow during 30 days at low temperature and perhaps

under vacuum packaging, leading to a switch of the two species during storage. Similarly, *L. piscium*,

a sub-dominant species in original HAM_B microbiota, outcompeted the initial dominant *Vibrio*

species.

Another finding from our work is that the *L. lactis* strain used for biopreservation is not competitive

towards ham original microbiota. Although, we noticed that it has perceivable effect towards the

reduction of other Firmicutes species (*e.g*. *C. maltaromaticum*), perhaps due to the production of

nisin. Indeed, this bacterium has been shown in previous studies to harbour sensitivity to this

bacteriocin in vacuum packed meat (La Storia et al., 2012; Teixeira et al., 2018). Albeit the *L. lactis*

CH-HP15 strain was inoculated with a level three orders of magnitude higher than the original ham

microbiota, the strain was found between 25% to 50% of overall relative abundance in HAM_A and

HAM_B, respectively. We previously observed that the inactivation level of this strain after HPP was

more important than that of other LAB species, which was compensated by its better ability to

recover and rapidly regrow after the treatment (Ramaroson et al., 2018). The lack of *L. lactis*

competitiveness is probably due to the specific ecology of ham. Indeed, *L. lactis* CH-HP15 was shown

able to grow in sterile diced cooked ham at 8°C, reaching 9 log CFU g^{-1} within 5 days with an initial

inoculum of 6 log CFU g^{-1} (Ramaroson et al., 2018). In the present study, the presence of species

belonging to the natural microbiota and thus potentially better adapted may explain the lack of

fitness of *L. lactis* CH-HP15. Nevertheless, the combination of high level of inoculation of *L. lactis* with

HPP treatment leads to an efficient stabilisation of the original cooked HAM_A microbiota, by limiting

the growth of *P. lundensis* and *S. grimesii*. Unlike this situation, results were different for HAM_B

samples for which the hurdle strategy could not trigger, over the storage time, the outcompetition of

L. lactis versus the original HAM_B microbiota. Therefore, it can be concluded that for microbiota of

HAM_B, the hurdle strategy failed in stabilizing and inactivating the microbial growth.

Furthermore, our results show that *B. thermosphacta* is a species with a very strong recovery

dynamics following the HPP treatment. Such observation has already been made by Teixiera and

colleagues using the simplified ham microbiota described above (Teixeira et al., 2018). This finding

raises the question of HPP treatment benefit on *B. thermosphacta* containing microbiota as this

species is a well-known meat spoilage micro-organism (Illikoud et al., 2019).

5. Conclusion

From our work, we thus conclude that both HPP treatment and *L. lactis*-based biopreservation is

strongly microbiota dependent and thus, the value of this strategy requires a specific assessment for

each type of cooked ham production. We recommend that HPP treatment should be evaluated not

only for pathogenic bacteria but also on putative spoilage bacteria in order to estimate the specific

selection of these undesirable micro-organisms, in particular *B. thermosphacta.*

Legends to the figures.

Figure 1. Comparative bacterial community composition between the four brand of cooked ham analysed.

Panel A: Cooked ham samples unsupervised clustering tree based on Bray-Curtis distance and Ward algorithm. Samples are colored according to the cooked ham brand. Both air (AIR) and vacuum packed (VPA) samples as well as 16 S rDNA-based or *gyrB* –based analysis are shown. **Panel B and C** show a bar plot composition of the top 10 species identified among Firmicutes and Proteobacteria phyla, respectively. Species are plotted according to their relative abundance in percentage of the whole microbiota. Novel genus nomenclature was applied for *Latilactobacillus* species (*L. sakei* and *L. curvatus*) as recently proposed by Zheng and colleagues (Zheng et al., 2020).

Figure 2. Time course dynamic barplot composition of bacterial communities in nitrite-reduced cooked ham samples upon various treatments.

Data for HAM_A and HAM_B samples are shown in panel A and B, respectively. Species are plotted according to their relative abundance in percentage of the whole microbiota. The *L. lactis* strain CH-HP15 used for biopreservation is highlighted in green in the left legend. Each bar plot is the mean of two biological replicates and each replicate value being the average of both 16S rDNA and *gyrB* amplicon sequencing data. As described in Figure 1 for *Latilactobacillus* species, novel genus nomenclature was applied for *Dellaglioa algida* (former *Lactobacillus algidus*) as recently proposed by Zheng and colleagues (Zheng et al., 2020).

Conflict of interest

None. The funders had no role in the study's design, data collection, and interpretation, or the decision to submit the work for publication.

Funding

This work was supported by the French National Research Agency (ANR) for their financial support in the framework of the BLAC-HP project (ANR-14-CE20-0004).

Acknowledgements

The authors express their gratitude to the INRAE MIGALE bioinformatics facility (MIGALE, INRAE, 2020.

Migale bioinformatics Facility, doi: 10.15454/1.5572390655343293E12) for providing computational

resources and data storage. We also thank the INRA @BRIDGe platform for carrying out the MiSeq

sequencing runs

References

Alahakoon, A.U., Jayasena, D.D., Ramachandra, S., Jo, C., 2015. Alternatives to nitrite in processed meat: Up to date. Trends Food Sci. Technol. 45, 37–49. https://doi.org/10.1016/j.tifs.2015.05.008

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., Madden, T.L., 2009. BLAST+: architecture and applications. BMC Bioinformatics 10, 421. https://doi.org/10.1186/1471-2105-10-421

EFSA, 2017. EFSA explains risk assessment. Nitrites and nitrates added to food.

Escudié, F., Auer, L., Bernard, M., Mariadassou, M., Cauquil, L., Vidal, K., Maman, S., Hernandez-Raquet, G., Combes, S., Pascal, G., 2018. FROGS: Find, Rapidly, OTUs with Galaxy Solution. Bioinforma. Oxf. Engl. 34, 1287–1294. https://doi.org/10.1093/bioinformatics/btx791

Han, Y., Jiang, Y., Xu, X., Sun, X., Xu, B., Zhou, G., 2011. Effect of high pressure treatment on microbial populations of sliced vacuum-packed cooked ham. Meat Sci. 88, 682–688. https://doi.org/10.1016/j.meatsci.2011.02.029

Illikoud, N., Gohier, R., Werner, D., Barrachina, C., Roche, D., Jaffrès, E., Zagorec, M., 2019. Transcriptome and Volatilome Analysis During Growth of *Brochothrix thermosphacta* in Food: Role of Food Substrate and Strain Specificity for the Expression of Spoilage Functions. Front. Microbiol. 10, 2527. https://doi.org/10.3389/fmicb.2019.02527

Jofré, A., Aymerich, T., Garriga, M., 2008a. Assessment of the effectiveness of antimicrobial packaging combined with high pressure to control Salmonella sp. in cooked ham. Food Control 19, 634–638. https://doi.org/10.1016/j.foodcont.2007.06.007

Jofré, A., Garriga, M., Aymerich, T., 2008b. Inhibition of *Salmonella sp. Listeria monocytogenes* and *Staphylococcus aureus* in cooked ham by combining antimicrobials, high hydrostatic pressure and refrigeration. Meat Sci. 78, 53–59. https://doi.org/10.1016/j.meatsci.2007.06.015

La Storia, A., Ferrocino, I., Torrieri, E., Di Monaco, R., Mauriello, G., Villani, F., Ercolini, D., 2012. A combination of modified atmosphere and antimicrobial packaging to extend the shelf-life of beefsteaks stored at chill temperature. Int. J. Food Microbiol. 158, 186–194. https://doi.org/10.1016/j.ijfoodmicro.2012.07.011

Liu, G., Wang, Y., Gui, M., Zheng, H., Dai, R., Li, P., 2012. Combined effect of high hydrostatic pressure and enterocin LM-2 on the refrigerated shelf life of ready-to-eat sliced vacuum-packed cooked ham. Food Control 24, 64–71. https://doi.org/10.1016/j.foodcont.2011.09.004

Mahé, F., Rognes, T., Quince, C., de Vargas, C., Dunthorn, M., 2014. Swarm: robust and fast clustering method for amplicon-based studies. PeerJ 2, e593. https://doi.org/10.7717/peerj.593

Majou, D., Christieans, S., 2018. Mechanisms of the bactericidal effects of nitrate and nitrite in cured meats. Meat Sci. 145, 273–284. https://doi.org/10.1016/j.meatsci.2018.06.013

Marcos, B., Jofré, A., Aymerich, T., Monfort, J.M., Garriga, M., 2008. Combined effect of natural antimicrobials and high pressure processing to prevent *Listeria monocytogenes* growth after a cold chain break during storage of cooked ham. Food Control 19, 76–81. https://doi.org/10.1016/j.foodcont.2007.02.005

McMurdie, P.J., Holmes, S., 2013. phyloseq: An R Package for Reproducible Interactive Analysis and
Graphics of Microbiome Census Data. PLOS ONE 8, e61217.
https://doi.org/10.1371/journal.pone.0061217
Modugno, C., Kmiha, S., Simonin, H., Aouadhi, C., Diosdado Cañizares, E., Lang, E., André, S., Mejri, S.,
Maaroufi, A., Perrier-Cornet, J. M., 2019. High pressure sensitization of heat-resistant and
pathogenic foodborne spores to nisin. Food Microbiol. 2019;84:103244.
https://doi.org/10.1016/j.fm.2019.103244
Mortensen, A., Aguilar, F., Crebelli, R., Domenico, A.D., Dusemund, B., Frutos, M.J., Galtier, P., Gott,
D., Gundert-Remy, U., Lambré, C., Leblanc, J.-C., Lindtner, O., Moldeus, P., Mosesso, P.,
Oskarsson, A., Parent-Massin, D., Stankovic, I., Waalkens-Berendsen, I., Woutersen, R.A.,
Wright, M., Brandt, P. van den, Fortes, C., Merino, L., Toldrà, F., Arcella, D., Christodoulidou,
A., Abrahantes, J.C., Barrucci, F., Garcia, A., Pizzo, F., Battacchi, D., Younes, M., 2017a. Re-
evaluation of potassium nitrite (E 249) and sodium nitrite (E 250) as food additives. EFSA J.
15, e04786. https://doi.org/10.2903/j.efsa.2017.4786
Mortensen, A., Aguilar, F., Crebelli, R., Domenico, A.D., Dusemund, B., Frutos, M.J., Galtier, P., Gott,
D., Gundert-Remy, U., Lambré, C., Leblanc, J.-C., Lindtner, O., Moldeus, P., Mosesso, P.,
Oskarsson, A., Parent-Massin, D., Stankovic, I., Waalkens-Berendsen, I., Woutersen, R.A.,
Wright, M., Brandt, P. van den, Fortes, C., Merino, L., Toldrà, F., Arcella, D., Christodoulidou,
A., Barrucci, F., Garcia, A., Pizzo, F., Battacchi, D., Younes, M., 2017b. Re-evaluation of sodium
nitrate (E 251) and potassium nitrate (E 252) as food additives. EFSA J. 15, e04787.
https://doi.org/10.2903/j.efsa.2017.4787
Najjari, A., Ouzari, H., Boudabous, A., Zagorec, M., 2008. Method for reliable isolation of *Lactobacillus*
sakei strains originating from Tunisian seafood and meat products. Int J Food Microbiol 121,
342–51. https://doi.org/S0168-1605(07)00617-4 [pii] 10.1016/j.ijfoodmicro.2007.11.045
Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R., O'Hara,
R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2019. vegan:
Community Ecology Package.
Oliveira, T.L.C. de, Ramos, A.L.S., Ramos, E.M., Piccoli, R.H., Cristianini, M., 2015. Natural
antimicrobials as additional hurdles to preservation of foods by high pressure processing.
Trends Food Sci. Technol. 45, 60–85. https://doi.org/10.1016/j.tifs.2015.05.007
Pietrasik, Z., Gaudette, N.J., Johnston, S.P., 2016. The use of high pressure processing to enhance the
quality and shelf life of reduced sodium naturally cured restructured cooked hams. Meat Sci.
116, 102–109. https://doi.org/10.1016/j.meatsci.2016.02.009
Pingen, S., Sudhaus, N., Becker, A., Krischek, C., Klein, G., 2016. High pressure as an alternative
processing step for ham production. Meat Sci. 118, 22–27.
https://doi.org/10.1016/j.meatsci.2016.03.014
Poirier, S., Rué, O., Peguilhan, R., Coeuret, G., Zagorec, M., Champomier-Vergès, M.-C., Loux, V.,
Chaillou, S., 2018. Deciphering intra-species bacterial diversity of meat and seafood spoilage
microbiota using *gyrB* amplicon sequencing: A comparative analysis with 16S rDNA V3-V4
amplicon sequencing. PLOS ONE 13, e0204629.
https://doi.org/10.1371/journal.pone.0204629
Qin, Q.-L., Xie, B.-B., Zhang, X.-Y., Chen, X.-L., Zhou, B.-C., Zhou, J., Oren, A., Zhang, Y.-Z., 2014. A
proposed genus boundary for the prokaryotes based on genomic insights. J. Bacteriol. 196,
2210–2215. https://doi.org/10.1128/JB.01688-14
Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glockner, F.O., 2012.
The SILVA ribosomal RNA gene database project: improved data processing and web-based
tools. Nucleic Acids Res 41, D590-6. https://doi.org/gks1219 [pii] 10.1093/nar/gks1219
Raimondi, S., Luciani, R., Sirangelo, T.M., Amaretti, A., Leonardi, A., Ulrici, A., Foca, G., D'Auria, G.,
Moya, A., Zuliani, V., Seibert, T.M., Søltoft-Jensen, J., Rossi, M., 2019. Microbiota of sliced
cooked ham packaged in modified atmosphere throughout the shelf life: Microbiota of sliced
cooked ham in MAP. Int. J. Food Microbiol. 289, 200–208.
https://doi.org/10.1016/j.ijfoodmicro.2018.09.017

Ramaroson, M., Guillou, S., Rossero, A., Rezé, S., Anthoine, V., Moriceau, N., Martin, J.-L., Duranton, F., Zagorec, M., 2018. Selection procedure of bioprotective cultures for their combined use with High Pressure Processing to control spore-forming bacteria in cooked ham. Int. J. Food Microbiol. 276, 28–38. https://doi.org/10.1016/j.ijfoodmicro.2018.04.010

Sindelar, J.J., Milkowski, A.L., 2012. Human safety controversies surrounding nitrate and nitrite in the diet. Nitric Oxide Biol. Chem. 26, 259–266. https://doi.org/10.1016/j.niox.2012.03.011

Stiles, M.E., 1996. Biopreservation by lactic acid bacteria. Antonie Van Leeuwenhoek 70, 331–345.Tassou, C. C., Galiatsatou, P., Samaras, F. J., and Mallidis, C. G., 2007. Inactivation kinetics of a piezotolerant *Staphylococcus aureus* isolated from high-pressure-treated sliced ham by high pressure in buffer and in a ham model system: Evaluation in selective and non-selective medium. *Innov Food Sci Emerg Technol* 8, 478-484. https://doi.org/10.1016/j.ifset.2007.04.002

Teixeira, J.S., Repková, L., Gänzle, M.G., McMullen, L.M., 2018. Effect of Pressure, Reconstituted RTE Meat Microbiota, and Antimicrobials on Survival and Post-pressure Growth of *Listeria monocytogenes* on Ham. Front. Microbiol. 9, 1979. https://doi.org/10.3389/fmicb.2018.01979

Vasilopoulos, C., De Maere, H., De Mey, E., Paelinck, H., De Vuyst, L., Leroy, F., 2010. Technology-induced selection towards the spoilage microbiota of artisan-type cooked ham packed under modified atmosphere. Food Microbiol 27, 77–84. https://doi.org/S0740-0020(09)00196-8 [pii] 10.1016/j.fm.2009.08.008

Verheyen, D., Govaert, M., Seow, T.K., Ruvina, J., Mukherjee, V., Baka, M., Skåra, T., Van Impe, J.F.M. 2020. The complex Effect ef food matrix fat content on thermal inactivation of *Listeria monocytogenes*: Case study in emulsion and gelled emulsion model systems. Front Microbiol 10:3149. https://doi: 10.3389/fmicb.2019.03149

Vermeiren, L., Devlieghere, F., Debevere, J., 2006. Co-culture experiments demonstrate the usefulness of *Lactobacillus sakei* 10A to prolong the shelf-life of a model cooked ham. Int. J. Food Microbiol. 108, 68–77. https://doi.org/10.1016/j.ijfoodmicro.2005.11.001

Vermeiren, L., Devlieghere, F., Debevere, J., 2004. Evaluation of meat born lactic acid bacteria as protective cultures for the biopreservation of cooked meat products. Int J Food Microbiol 96, 149–64. https://doi.org/10.1016/j.ijfoodmicro.2004.03.016 S0168160504001667

Yamamoto, K., 2017. Food processing by high hydrostatic pressure. Biosci. Biotechnol. Biochem. 81, 672–679. https://doi.org/10.1080/09168451.2017.1281723

Zagdoun, M., Coeuret, G., N'Dione, M., Champomier-Vergès, M.-C., Chaillou, S., 2020. Large microbiota survey reveals how the microbial ecology of cooked ham is shaped by different processing steps. Food Microbiol. 91. https://doi.org/10.1016/j.fm.2020.103547Zheng, J., Wittouck, S., Salvetti, E., Franz, CMAP., Harris, HMB., Mattarelli, P., O'Toole, PW., Pot, B., Vandamme, P., Walter, J., Watanabe, K., Wuyts, S., Felis, GE., Gänzle, MG., Lebeer, S., 2020. A taxonomic note on the genus *Lactobacillus*: Description of 23 novel genera, emended description of the genus *Lactobacillus* Beijerinck 1901, and union of *Lactobacillaceae* and *Leuconostocaceae*. Int J Syst Evol Microbiol. 70,2782-2858. https://doi.org/10.1099/ijsem.0.004107

Table 1 Description of ham specificities

Sample name according to brands	Sampling date	Additives[a]	Total counts[b] log CFU g^{-1}	LAB counts[c] log CFU g^{-1}
HAM_A	UBD + 6 days Dec 2 2015	Dextrose Sodium nitrite E250 Potassium chloride E508 Sodium erythorbate E316 Potassium lactate E326 Sodium ascorbate E301 Sodium acetate E262 Diphosphates E450 Triphosphates E451 Polyphosphates E452	8.6 ± 0.4	8.2 ± 0.0
HAM_B[d]	UBD + 5 days Mai 18 2016	Sugar Lactose Nitrite salt Common salt Spices	7.2 ± 0.7	7.3 ± 0.5
HAM_C	UBD – 1 day Nov 6 2015	Dextrose Sodium nitrite E250 Potassium chloride E508 Potassium lactate E326 Sodium acetate E262 Sodium ascorbate E316 Polyphosphates E452	8.6 ± 0.1	9.0 ± 0.1
HAM_D	UBD – 4 days Aug 21 2015	Sodium nitrite E250 Potassium chloride E508 Potassium lactate E326 Sodium diacetate E262 Sodium ascorbate E316	10.3 ± 0.1	8.1 ± 0.1

[a]: from information provided on the packaging. [b]: PCA count mean values from portions stored under vacuum and air packaging . [c]: MRS count mean values from portions stored under vacuum and air packaging. [d]: brand sold as organic.

Table 2.

	Population level in log CFU.g^{-1} of nitrite-reduced diced cooked ham			
	Storage time in days^A			
HAM_A samples	D01	D05	D12	D30
Without treatment	4.63 ± 0.09	7.32 ± 0.14	7.84 ± 0.10	8.15 ± 0.15
With HPP	1.20 ± 0.14	3.74 ± 0.01	3.00 ± 1.00	4.02 ± 0.59
With L. lactis	7.21 ± 0.62	9.18 ± 0.02	8.39 ± 0.12	8.87 ± 0.87
With L. lactis and HPP	2.54 ± 0.45	3.97 ± 0.26	5.42 ± 1.27	7.60 ± 0.03
HAM_B samples	D01	D05	D12	D30
Without treatment	5.24 ± 0.18	6.10 ± 0.13	6.90 ± 0.33	6.50 ± 0.20
With HPP	1.39 ± 0.09	ND^B	5.33 ± 0.34	6.23 ± 0.42
With L. lactis	7.25 ± 0.51	8.29 ± 0.01	7.82 ± 0.21	8.50 ± 0.50
With L. lactis and HPP	2.75 ± 0.44	3.64 ± 0.07	7.11 ± 0.13	10.2 ± 0.20

^{A} Population data were calculated and averaged from the two biological replicates and from two plating culture condition enumerating both total mesophilic bacteria and lactic acid bacteria (see M&M).

^{B} Not Determined.

Composition of top10 Firmicutes Species

Composition of top10 Proteobacteria Species

A Composition of top10 species in HAM_A samples

B Composition of top10 species in HAM_B samples