***In Vitro* Efficacy of Five Biofungicides against *Rhizoctonia solani* Kuhn in Rice**

Irish Mae Bauzon-Cantila[1*], Jaime C. Silvestre[2], Raquel B. Evangelista[2], Edralyn

Catubay[2,3]

[1]Graduate School, University of Southern Mindanao

[2]College of Agriculture, University of Southern Mindanao

[3]Institute of Agriculture and Related Sciences, Davao del Sur State College

Corresponding author: bauzon.irishmae@gmail.com

ABSTRACT

Rhizoctonia solani Kuhn*,* the causal pathogen of sheath blight is second most damaging

fungal disease in rice. While using chemical fungicides present high detriment to environment,

the study investigate the efficacy of treatments composing five biofungicides in three different

rates along with a biological agent, chemical check and untreated against the pathogen in *in vitro*

level. *In vitro* efficacy showed that *Melaleuca alternifolia* + terpenes at 3.00 ml/L of H_2O (T6),

Aloe vera powder (Manopol) + *Melaleuca* oil at 3.00 ml/L of H_2O (T15) and at 2.00 ml/L of H_2O

(T14) and *Melaleuca alternifolia* + terpenes at 2.00 ml/L of H_2O (T5) as very effective (0-10 mm

diameter zone of growth) treatments comparable to the chemical check (T17). Therefore,

attaining high yield rice while having low risk to environment can always be done.

KEYWORDS: Aloe vera, *Melaleuca,* Biofungicides, rice sheath blight

INTRODUCTION

Rhizoctonia solani Kuhn, causing sheath blight is one of the most economically

important fungal diseases that limits rice production [1], which its sclerotia or hyphae can easily

attach to the rice sheath and infecting blight in rice [2]. Further infections are caused by hyphae

growing upward towards uninfected plant parts and produce additional lesions and sclerotia on

leaf sheaths affecting productivity in plants [3, 4]. Depending on the plant stage upon infection

and type of environment, a range of 4 to 50% can be reduced to the yield, making sheath blight

as the second most damaging fungal disease in rice [1, 5-8].

Rice released varieties such as hybrid rice and foreign germplasm with low resistance to sheath blight [9] had been tested to high rainfall areas in the Philippines [10, 11], which is a disease prone environment. It could be advantageous then to apply biofungicides to the crop. Biofungicides as one of the methods that can manage the disease is gaining popularity because it is more environment-friendly [12, 13]. Moreover, regulations towards utilizing chemical fungicides could be stricter in the future as several organizations are putting pressure on taking out hazardous chemicals in the market [13]. Hence, this study was conducted to investigate the *in vitro* efficacy of biofungicides against *R. solani* Kuhn in rice.

MATERIALS AND METHODS

INOCULUM AND TREATMENT PREPARATION

Diseased sheaths of rice due to blight were collected in the field (Figure 1) and placed into plastic bag for diagnosis and isolation. Potato dextrose agar (PDA) was then used as medium for isolating sheath blight pathogen with the following components: 200 g of clean potato, 1 L of water, 20 g dextrose, and 20 g agar. The medium was dispensed in flat bottles, plugged with cotton and autoclaved at 15 *psi* or 121°C for 15-30 minutes. Tissue planting technique was used to transfer pathogen in PDA medium, which was incubated under room condition. Purification was done by transferring mycelial bits onto fresh PDA medium.

A total of 18 treatments composed of five biofungicides in three different rates, a biological control, chemical check, and untreated was used. The main component of the five biofungicides is *Bacillus subtilis*, *Melaleuca alternifolia* + terpenes, seaweed extract, *Lactobacillus plantarum*, aloe vera powder (Manopol) + *Melaleuca* oil. *Trichoderma* spp. was used as biological agent while Propiconazole + difenoconazole as chemical check. Other details of each treatment can be found in Table 1.

BIOASSAY TEST

In a 14 days old *R. solani* Kuhn culture (Figure 2), culture disc method was used for fungicidal assay with the following steps in Figure 3. Petri plates were rotated thoroughly for mixing and congealing. Culture disc of the pathogen was placed at the center of the plates, which were labelled and incubated in inverted position under room temperature. Measurement of the diameter zone of growth (mm) of the pathogen was done after 72 hrs of incubation.

DATA ANALYSIS

The *in vitro* test was laid-out in completely randomized design with 18 treatments in triplicates. The data gathered was diameter zone of growth (DZG), which is a measurement to determine the efficacy of treatment. The data (mm) was taken after 72 hrs of incubation, which efficacy follows an arbitrary scale range: 0-10 mm is very effective, 11-20 mm is effective, 21-30 mm is moderately effective, and 31 mm and above is not effective. The analysis of variance and mean separation were done using Statistix 9.0 [14].

RESULTS AND DISCUSSION

In vitro result showed the mean DZG (mm) of *R. solani* Kuhn was significantly affected by the application of different rates of five biofungicides after 72 hrs of incubation (Table 2). The most effective biofungicide was *Melaleuca alternifolia* + terpenes at 3.00 ml/L of H_2O (T6) with no observed growth of the pathogen (Figure 4). *Aloe vera* powder (Manopol) + *Melaleuca* oil at 3.00 ml/L of H_2O (T15) with 0.17 mm and at 2.00 ml/L of H_2O (T14) with 2.17 mm and *Melaleuca alternifolia* + terpenes at 2.00 ml/L of H_2O (T5) with 3.83 mm followed as among the most effective treatments (Figure 4). The effect of *Melaleuca alternifolia* + terpenes (T5 and T6) and *Aloe vera* powder (Manopol) + *Melaleuca* oil (T14 and T15) were comparable to the chemical check (T17) and rated as very effective (Figure 4). This implies that these two biofungicides were as effective as the chemical check in inhibiting the growth of *R. solani* Kuhn *in vitro*.

Previous reports showed that *Aloe vera* powder (Manopol) + *Melaleuca* oil effectively controlled *Corynespora* leaf fall disease of rubber caused by *Corynespora cassiicola* in *in vitro* study [15] while *Melaleuca alternifolia* + terpenes was found effective against *Pythium aphanidermatum* on pepper [16] and effectively control diseases *Citrus madurensis* [17]. The lowest rates at 1.00 ml/L of H_2O of *Melaleuca alternifolia* + terpenes (T4) and *Aloe vera* powder (Manopol) + *Melaleuca* oil (T13) were still rated effective, similar to the effect of lowest rate of *Bacillus subtilis* at 1.25 ml/L of H_2O (T1) (Figure 5). *Bacillus subtilis* at 2.5 ml/L of H_2O (T2) and 3.75 ml/L of H_2O (T3) were rated as moderately effective (Figure 6). Lastly, the three rates of seaweed extract and *Lactobacillus plantarum* were rated not effective with DZG means ranging

from 40.83 to 66.83 mm and their results were comparable to the untreated control and biological check with DZG means of 68.33 and 51.67 mm, respectively (Figure 7).

CONCLUSION

Biofungicides especially those containing *Melaleuca alternifolia* and *Aloe vera* are very effective in controlling *R. solani* Kuhn in rice in *in vitro*, which could be verified in *in vivo* study. Attaining high yield and less pesticides residue in rice is always possible, which should be promoted to uphold a safer environment.

ACKNOWLEDGEMENTS

The authors would like to thank Department of Plant Pathology, College of Agriculture for allowing the utility of laboratory equipment.

STATEMENT ON CONFLICT OF INTEREST

The authors declare no conflict of interest.

121   Table 1. Biofungicides and their corresponding details that has been used in the study.

Component	Commercial name	Reports	Source	Treatment (rate*)
Bacillus subtilis	Seranade 250	suppresses club root to canola on *in vitro* conditions and effective against crop damaging pathogens like powdery mildew, walnut blight and *Botrytis* bunch rot and fire blight	[18, 19]	T1 (1.25 ml)
				T2 (2.5 ml)
				T3 (3.75 ml)
Melaleuca alternifolia + terpenes	Timorex Gold	having prophylactic and curative activity against broad spectrum of diseases in crops	[20]	T4 (1 ml)
				T5 (2 ml)
				T6 (3 ml)
Seaweed extract	MPQ	produces metabolites with antiviral, antiprotozoal, antifungal, and antibacterial properties aiding in the protection	[21, 22]	T7 (0.25 g)
				T8 (0.5 g)
				T9 (0.75 g)
Lactobacillus plantarum	EM-1	produce antifungal substances such as benzoic acid, methylhydantoin, mevalonolactone against plant pathogens	[23-25]	T10 (1 ml)
				T11 (2 ml)
				T12 (3 ml)
Aloe vera powder (Manopol) + *Melaleuca* oil	AZ 41	effectively control diseases and insect pests in citrus	[17]	T13 (1 ml)
				T14 (2 ml)
				T15 (3 ml)
Biological control	*Trichoderma* spp.	induce systemic resistance against plant pathogens	[26]	T16 (0.5 ml)
Propiconazole + difenoconazole	Armure	controls *Rhizoctonia solani* Kuhn in rice very effectively	[27, 28]	T17 (1 ml)
Untreated				T18 (Untreated)

122   *diluted in L of H_2O, MPQ= Marvelous Premium Quality, EM= effective microorganism

128 Table 2. Diameter zone of growth (mm) of *R. solani* Kuhn as influenced by different rates of
129 biofungicides

Treatments	Mean (mm) [1]	Degree of Effectiveness
T1	18.17 [cde]	Effective
T2	20.50 [bcde]	Moderately effective
T3	22.17 [bcde]	Moderately effective
T4	14.83 [de]	Effective
T5	3.83 [e]	Very effective
T6	0.00 [e]	Very effective
T7	53.17 [ab]	Not effective
T8	66.83 [a]	Not effective
T9	48.83 [abcd]	Not effective
T10	44.00 [abcd]	Not effective
T11	40.83 [abcd]	Not effective
T12	41.33 [abcd]	Not effective
T13	20.33 [bcde]	Effective
T14	2.17 [e]	Very effective
T15	0.17 [e]	Very effective
T16	51.67 [abc]	Not effective
T17	0.00 [e]	Very effective
T18	68.33 [a]	Not effective

[1]Means in column followed by common letter superscripts are not significantly different at 1% level, Tukey's Test.

Figure 1. Rice plants showing symptoms of sheath blight on the (a) leaf (gray irregular lesions
banded with green brown coloration and grayish center) and (b) sheath (oval gray spots with
black brown margins that later enlarge).

Figure 2. Pure culture of *Rhizoctonia solani* Kuhn in 14 days old showing mycelia and matured
sclerotia on potato dextrose agar.
139

Figure 3. Culture disc method for biofungicidal assay of *Rhizoctonia solani* Kuhn. *a) Potato dextrose agar (PDA) was used as medium for isolating the pathogen, b) 1 ml of the treatment was added to each petri plate with 20 ml melted PDA, c) Ensure aseptic condition by flaming the corkborer, d) The isolated pathogen was removed aseptically, and e) The culture disc of the pathogen was placed at the center of each plate.

Figure 4. Diameter zone of growth of *R. solani* Kuhn showing very effective treatments in petri dish.

Figure 5. Diameter zone of growth of *R. solani* Kuhn showing effective treatments in petri dish.

Figure 6. Diameter zone of growth of *R. solani* Kuhn showing moderately effective treatments.

159 Figure 7. Diameter zone of growth of *R. solani* Kuhn showing not effective treatments.

References

1.   Molla, K.A., et al., *Understanding sheath blight resistance in rice: the road behind and the road*
ahead. Plant biotechnology journal, 2020. **18**(4): p. 895-915.
2.   Singh, R., S. Sunder, and P. Kumar, *Sheath blight of rice: Current status and perspectives.* 2016.
69: p. 340-351.
3.   Brooks, S.A., *Sensitivity to a Phytotoxin from Rhizoctonia solani Correlates with Sheath Blight*
Susceptibility in Rice. Phytopathology, 2007. **97**(10): p. 1207-1212.
4.   Webster, R.K., *Compendium of rice diseases.* 1992: The American Phytopathological Society
(APS).
5.   Bhukal, N., R. Singh, and N. Mehta, *Progression and development of sheath blight of rice in*
relation to weather variables J Mycol Pl Pathol 2015. **45**(2): p. 166-172.
6.   Cu, R., et al., *Effect of sheath blight on yield in tropical, intensive rice production system.* Plant
Disease, 1996.
7.   Singh, S.K., et al., *Current status and impact of sheath blight in rice (Oryza sativa L.) - a review*
Agricultural Reviews, 2004. **25**: p. 289-297.
8.   Zheng, A., et al., *The evolution and pathogenic mechanisms of the rice sheath blight pathogen.*
Nature communications, 2013. **4**: p. 1424-1424.
9.   https://www.pinoyrice.com/rice-varieties/. 12/01/2017].

10.   Cantila, A. and A. Quitel, *Statistical Analysis of Foreign Rice Phenotypes under Different Seasons*
of the Philippines. Journal of Advanced Applied Scientific Research, 2017. **1**(12).

11.   Cantila, A.Y., A.M.L. Fordan, and I.V. Boholano, *Phenotypic variation and correlation of traits in*
Philippine hybrid rice varieties. J Bangladesh Agril Univ, 2019. **17**(2): p. 179-186.

12.   Alabouvette, C., C. Olivain, and C. Steinberg, *Biological control of plant diseases: the European*
situation. European journal of plant pathology, 2006. **114**(3): p. 329-341.

13.   De Cal, A., et al., *Use of Biofungicides for Controlling Plant Diseases to Improve Food Availability.*
Agriculture, 2012. **2**(2).

14.   Analytical-software, *Statistix 9.0.* 2009, Analytical software: Florida State University, PO Box
12185, Tallahassee, Florida 32317, USA.

15.   Villanueva, J.D., et al., *Management of Corynespora Leaf Fall Disease of Rubber in the Nursery.*
USM R&D Journal, 2016. **24**(1).

16.   Mihajlović, M., et al., *In vitro and in vivo toxicity of several fungicides and Timorex gold*
biofungicide to Pythuim aphanidermatum. Pesticidi i fitomedicina, 2013. **28**(2): p. 117-123.

17.   Tangonan, N., E. Butardo, and C.P. Kabacan, *Effects of AZ41 organic foliar fertilizer on pests,*
diseases, and yield of calamansi (Citrus madurensis). Philippine Journal of Crop Science
(Philippines), 2004.

18.   Lahlali, R., et al., *Evidence that the biofungicide Serenade (Bacillus subtilis) suppresses clubroot*
on canola via antibiosis and induced host resistance. Phytopathology, 2013. **103**(3): p. 245-254.

19.   Peng, G., et al., *Potential biological control of clubroot on canola and crucifer vegetable crops.*
Plant Pathology, 2011. **60**(3): p. 566-574.

20.   STK, *Timorex Gold, a broad spectrum fungicide*, S.A.P. Incorporated, Editor. 2017, Stockton
group: Unit 12 1st Floor, MSA Prime Center, Madrigal Business Park, Ayala Alabang, Muntinlupa
City.

21.   Kumari, S.S., V.D. Kumar, and B. Priyanka, *Antifungal Efficacy of Seaweed Extracts Against*
Fungal Pathogen of Silkworm, Bombyx mori L. International Journal of Agricultural Research,
2017. **12**: p. 123-129.

22.   Pérez, M.J., E. Falqué, and H. Domínguez, *Antimicrobial Action of Compounds from Marine*
Seaweed. Marine drugs, 2016. **14**(3): p. 52.

23.   Prema, P., et al., *Production and characterization of an antifungal compound (3-phenyllactic*
acid) produced by Lactobacillus plantarum strain. Food and Bioprocess Technology, 2010. **3**(3):
p. 379-386.

24.   Renuka, R. and B. Parameswari, *Effective Microbes (EM) - An Organic Agricultural Technology.*
Research News for U, 2012. **9**: p. 102-105.

25.   Wang, H., et al., *Activity against plant pathogenic fungi of Lactobacillus plantarum IMAU10014*
isolated from Xinjiang koumiss in China. Annals of Microbiology, 2011. **61**(4): p. 879-885.

26.   Harman, G.E., *Myths and dogmas of biocontrol changes in perceptions derived from research on*
Trichoderma harzinum T-22. Plant disease, 2000. **84**(4): p. 377-393.

27.   Chahal, K., S. Sokhi, and G. Rattan, *Investigations on sheath blight of rice in Punjab.* Indian
Phytopathology, 2003. **56**(1): p. 22-26.

28.   Kandhari, J., *Management of sheath blight of rice through fungicides and botanicals.* Indian
Phytopathology, 2007. **60**(2): p. 214-217.

221