Isoprenoid biosynthesis regulation in poplars by methylerythritol

phosphate and mevalonic acid pathways

Ali Movahedi[1], Hui Wei[1,†], Boas Pucker[2,†], Tingbo Jiang[3], Weibo Sun[1], Dawei Li[1], Liming Yang[1,*],

and Qiang Zhuge[1,*]

[1] Co-Innovation Center for Sustainable Forestry in Southern China, Key Laboratory of Forest

Genetics & Biotechnology, Ministry of Education, College of Biology and the Environment,

Nanjing Forestry University, Nanjing, China; ali_movahedi@njfu.edu.cn (A.M.);

15850682752@163.com (H.W.); czswb@njfu.edu.cn (W.S.); dwli@njfu.edu.cn (D.L.);

yangliming@njfu.edu.cn (L.Y.); qzhuge@njfu.edu.cn (Q.Z.)

[2] Institute of Plant Biology, TU Braunschweig, Braunschweig, Germany; b.pucker@tu-

braunschweig.de (B.P.)

[3] State Key Laboratory of Tree Genetics and Breeding, Northeast Forestry University, Harbin,

China; tbjiang@yahoo.com (T.J.)

* Correspondence: yangliming@njfu.edu.cn; qzhuge@njfu.edu.cn

† These authors contributed equally as the first author.

Abstract

The isoprenoids found in plants are extremely important to survive with various human applications, such as flavoring, fragrance, dye, pharmaceuticals, and biomass used for biofuels. Methylerythritol phosphate (MEP) and mevalonic acid (MVA) pathways are critical in plants, responsible for isoprenoid biosynthesis. 1-deoxy-D-xylulose5-phosphate synthase (DXS) and 1-deoxy-D-xylulose5-phosphate reductoisomerase (DXR) catalyze the rate-limiting steps in the MEP pathway, while 3-hydroxy-3-methylglutaryl-CoA reductase (HMGR) catalyzes the rate-limiting step in the MVA pathway. Here, we showed while *PtHMGR* overexpressors (OEs) exhibited different MEP- and MVA-related gene expressions compared with non-transgenic poplars (NT), the *PtDXR-OEs* revealed upregulated MEP-related and downregulated MVA-related gene expressions. *PtDXR* and *PtHMGR* overexpressions caused changes in MVA-derived trans-zeatin-riboside, isopentenyl adenosine, castasterone, and 6-deoxocastasterone well as MEP-derived carotenoids and gibberellins. In *PtHMGR*-OEs, the accumulated geranyl diphosphate synthase (*GPS*) and geranyl pyrophosphate synthase (*GPPS)* transcript levels in the MEP pathway led to an accumulation of MEP-derived isoprenoids. In contrast, upregulation of farnesyl diphosphate synthase (*FPS*) expression in the MVA pathway contributed to increased levels of MVA-derived isoprenoids. In addition, *PtHMGR*-OEs increased MEP-related *GPS* and *GPPS* transcript levels, expanded MEP-derived isoprenoid levels, changed *FPS* transcript levels, and affected MVA-derived isoprenoid yields. These results demonstrate the contribution of MVA and MEP pathways regulating isoprenoid biosynthesis in poplars.

Keywords: MEP; MVA; HMGR; DXR; Isoprenoid biosynthesis

1. Introduction

Biosynthesis of isoprenoids (terpenoids) is essential for all living organisms. There are over 50,000 distinct molecules in living organisms that are isoprenoids, presenting many functional and structural properties (Thulasiram et al., 2007). Isoprenoids play vital roles in plant growth and development, as well as in membrane fluidity, photosynthesis, and respiration. As specific metabolites, they join in plant-pathogen and allelopathic interactions to preserve plants upon pathogens and herbivores, and they are also created to draw pollinators and seed-dispersing animals. Numerous isoprenoids are of commercial importance for rubber products and drugs, flavors, fragrances, agrochemicals, nutraceuticals, disinfectants, and pigments (Bohlmann and Keeling, 2008). A wide range of isoprenoid biochemical processes are involved in photosynthesis in plants, including electron transfer, quenching of excited chlorophyll triplets, light-harvesting, and energy conversion (Malkin R, 2000). Chlorophylls, consisting of the heme pathway-derived tetrapyrrole ring with an appended isoprenoid-derived phytol chain, exist in all reaction center and antenna complexes to absorb light energy and transfer electrons to the reaction centers. The linear or partially cyclized carotenes and their oxygenated derivative xanthophyll are isoprenoids that extinguish excess excitation energy through light-harvesting to preserve the light-harvesting system. However, these isoprenoids operate as attractants in flowers and fruits as well (Rodriguez-Concepcion, 2010). A large proportion of the isoprenoid flux in plants is conducted toward the synthesis of membrane sterol lipids. In contrast to vertebrates synthesizing cholesterol, higher plants synthesize a complex mix of sterol lipids called phytosterols (Boutte and Grebe, 2009).

Plants isoprenoids include gibberellins (GAs), carotene, Lycopene, cytokinins (CKs), strigolactones (GRs), and brassinosteroids (BRs) are produced through methylerythritol phosphate (MEP) and mevalonic acid (MVA) pathways (Henry et al., 2015; van Schie et al., 2006; Xie et al., 2008). The mentioned pathways are involved in plant growth, development, and response to environmental changes (Bouvier et al., 2005; Kirby and Keasling, 2009). The isopentenyl diphosphate isomerase (IDI) catalyzes the conversion of the isopentenyl diphosphate (IPP) into dimethylallyl diphosphate (DMAPP), leading to provide the basic materials for all isoprenoid productions (Hemmerlin, 2012; Lu et al., 2012; Zhang et al., 2019). The produced IPP and DMAPP play essential roles in MEP and MVA pathways crosstalk

(Huchelmann et al., 2014; Liao et al., 2016). The MVA pathway reactions appear in the

cytoplasm, endoplasmic reticulum (ER), and peroxisomes (Cowan et al., 1997; Roberts, 2007),

producing sesquiterpenoids and sterols. The 3-hydroxy-3-methylglutaryl-CoA reductase

(HMGR), a rate-limiting enzyme in the MVA pathway, catalyzes 3-hydroxy-3-methylglutary-

CoA (HMG-CoA) to form MVA (Cowan et al., 1997; Roberts, 2007).

Reactions of the MEP pathway occur in the chloroplast and produce carotenoids, GAs,

and diterpenoids. 1-deoxy-D-xylulose5-phosphate synthase (DXS) and 1-deoxy-D-xylulose5-

phosphate reductoisomerase (DXR) are rate-limiting enzymes in the MEP pathway that

catalyze the conversion of D-glyceraldehyde3-phosphate (D-3-P) and pyruvate into 2-C-

methyl-D-erythritol4-phosphate (MEP) (Cordoba et al., 2009; Perreca et al., 2020; Wang et al.,

2012; Yamaguchi, 2018). Terpenoids like phytoalexin and volatile oils play essential roles in

plant growth, development, and disease resistance (Hain et al., 1993; Ren et al., 2008).

Photosynthetic pigments convert organic carbon into plant biomass (Esteban et al., 2015). In

addition to an extensive range of natural functions in plants, terpenoids also consider the

potential for biomedical applications. Paclitaxel is one of the most effective chemotherapy

agents for cancer treatment, and artemisinin is an anti-malarial drug (Kim et al., 2016a; Kong

and Tan, 2015).

Previous metabolic engineering studies have proposed strategies to improve the

production of specific metabolites in plants (Ghirardo et al., 2014; Opitz et al., 2014). For

example, PMT and H6H encoding the putrescine N-methyltransferase and hyoscyamine 6 β-

hydroxylase respectively produced significantly higher scopolamine in transgenic henbane

hairy root. Also, HCHL encoding p-hydroxycinnamoyl-CoA hydratase/lyase accumulated the

glucose ester of p-hydroxybenzoic acid (pHBA) in Beta vulgaris hairy root (Rahman et al., 2009;

Zhang et al., 2004). The 3-hydroxy-3-methylglutaryl-coenzyme A synthase (HMGS) is the

second enzyme in the MVA pathway. Liao et al. (2018) confirmed that *HMGS* overexpression

of *Brassica juncea* upregulates carotenoid and phytosterol in tomatoes. HMGR has been

considered a critical factor in metabolically engineering terpenoids (Aharoni et al., 2005;

Dueber et al., 2009). In addition, *PgHMGR1* overexpression of ginseng increases ginsenosides

content, a necessary pharmaceutically active component (Kim, 2014).

Transgenic tobacco overexpressing the *Hevea brasiliensis HMGR* enhanced the

phytosterol levels (Schaller et al., 1995). It has been shown (Dai et al., 2011)

that *SmHMGR2* in *Salvia miltiorrhiza*, resulting in the improvement of squalene and tanshinone contents. Moreover, *Arabidopsis thaliana HMGR1 (AtHMGR1)* enhanced the phytosterol levels in the first generation of transgenic tomatoes (Enfissi et al., 2005). While the deaccumulation of *DXR* transcripts resulted in lower pigmentation and chloroplast appearance defects, the upregulated *DXR* expression caused the MEP-derived plastid isoprenoids to accumulate. Therefore, *DXR* can be genetically engineered to regulate the content of terpenoids and expressed *DXR* in *Arabidopsis* and observed enhanced flux through the MEP pathway (Carretero-Paulet et al., 2006). While the *A*. *thaliana DXR* overexpression caused the diterpene anthiolimine to accumulate in *Salvia sclarea* hairy roots (Vaccaro et al., 2014), the peppermint *DXR* overexpression resulted in essential oil inflation (about 50%) with no significant variations in monoterpene composition (Mahmoud and Croteau, 2001). Furthermore, previous studies have shown the exchange of metabolic intermediates included in the MVA- and MEP pathways through plastid membranes (Laule, 2003; Liao, 2006). In summary, the overexpression of genes involved in the MVA- and MEP pathways can change the abundances or activities of related enzymes and metabolic products, causing a new opportunity for plant breeding to enhance the accumulation of related metabolic products.

Poplars as an economic and energy species are widely used in industrial and agricultural production. Its fast growth characteristics and advanced resources in artificial afforestation play a vital role in the global ecosystem (Devappa, 2015).

This study investigates the poplar isoprenoid biosynthesis. We showed that the overexpression of the MVA-specific *PtHMGR* gene upregulated not only MVA- but MEP-related genes in the transcript levels. We also proved that the overexpression of the MEP-specific *PtDXR* gene caused downregulating MVA-related genes compared with upregulating MEP-related genes, enhancing terpenoid accumulation. Taken together, these results indicate that the MEP is a dominant pathway in contribution with the MVA pathway to produce isoprenoids secondary metabolites, and *HMGR* and *DXR* genes play key regulation points in these pathways.

2. Materials and Methods

2.1. *Plant materials and growth conditions*

Non-transgenic *P*. *trichocarpa* and *Populus × euramericana* cv. 'Nanlin 895' plants were

cultured in half-strength Murashige and Skoog (1/2 MS) medium (pH 5.8) under conditions of 24°C and 74% humidity (Movahedi et al., 2015). Subsequently, NT and transgenic poplars were cultured in 1/2 MS under long-day conditions (16 h light/8 h dark) at 24°C for 1 month (Movahedi et al., 2018).

2.2. *PtHMGR and PtDXR genes isolation and vector construction*

To produce cDNA, total RNA was extracted from *P. trichocarpa* leaves and processed with PrimeScript™ RT Master Mix, a kind of reverse transcriptase (TaKaRa, Japan). Forward and reverse primers (Supplementary Table 1: *PtHMGR*-F and *PtHMGR*-R) were designed, and the open reading frame (ORF) of *PtHMGR* was amplified via PCR. We then used the total volume of 50µl including 2 µl primers, 2.0 µl cDNA, 5.0 µl 10 × PCR buffer (Mg2+), 4µl dNTPs (2.5 mM), 0.5 µl rTaq polymerase (TaKaRa, Japan) for the following PCR reactions: 95°C for 7 min, 35 cycles of 95°C for 1 min, 58°C for 1 min, 72°C for 1.5 min, and 72°C for 10 min. Subsequently, the product of the *PtHMGR* gene was ligated into the pEASY-T3 vector (TransGen Biotech, China) based on blue-white spot screening, and the *PtHMGR* gene was inserted into the vector pGWB9 (Song et al., 2016) using Gateway technology (Invitrogen, USA). On the other hand, all steps to generate cDNA, RNA extraction, PCR, pEASY-T3 ligation, and vector construction (pGWB9-PtDXR) of *PtDXR* have been carried out according to Xu et al. (2019).

2.3. *phylogenetic analyses*

We applied the ClustalX for multiple sequence alignment of HMGR proteins, and MEGA5.0 software was used to construct a phylogenetic tree using 1000 bootstrap replicates. The amino acid sequences of HMGR from *Populus trichocarpa*, *Arabidopsis thaliana*, *Gossypium raimondii*, *Malus domestica*, *Manihot esculenta*, *Oryza sativa*, *Prunus persica*, *Theobroma cacao*, and *Zea mays* were obtained from the National Center for Biotechnology Information database (https://www.ncbi.nlm.nih.gov/) and Phytozome (https://phytozome-next.jgi.doe.gov/).

2.4. *Transgenic poplars: generation and confirmation*

Agrobacterium tumefaciens var. EHA105 was used for the infection of poplar leaves and petioles (Movahedi et al., 2014). Poplar buds were screened on differentiation MS medium

supplemented with 30 μg/mL Kanamycin (Kan). Resistant buds were planted in bud elongation

MS medium containing 20 μg/mL Kan and transplanted into 1/2 MS medium including 10

μg/mL Kan to generate resistant poplar trees. Genomic DNA has been extracted from putative

transformants one-month-old leaves grown on a kanamycin-containing medium using

TianGen kits (TianGen BioTech, China). The quality of the extracted genomic DNA (250–350

ng/μl) was determined by a BioDrop spectrophotometer (UK). PCR was carried out using

designed primers (Supplementary Table 1: CaMV35S as the forward and PtHMGR as the

reverse), Easy Taq polymerase (TransGene Biotech), and 50 ng of extracted genomic DNA as a

template to amplify about 2000 bp. In addition, total RNA was extracted from these one-

month-old leaves to produce cDNA, as mentioned above. These cDNA then were applied to

reverse transcription-quantitative PCR (RT-qPCR) (Supplementary Table 1: PtHMGR forward

and reverse) for comparing the transformants *PtHMGR-OEs* expressions with NT poplars and

transforming confirmation.

2.5.   *Phenotypic properties evaluation*

To evaluate phenotypic changes, we selected 45-day-old poplars from PtHMGR-and

PtDXR-OEs and NT poplars. We then simultaneously calculated the stem lengths (mm) and

stem diameters (mm) every day and recorded them. All recorded were analyzed by GraphPad

Prism 9, applying ANOVA one way (Supplementary Table 2).

2.6.   *Analyses via qRT-PCR*

12-month-old *PtDXR-OEs* (Xu et al., 2019) and *PtHMGR-OE* poplars (Soil-grown poplars)

have been used to extract total RNA. The qRT-PCR was performed to identify MVA- and MEP-

related gene expression levels in NT, *PtDXR-OE*, and *PtHMGR-OE* poplars. The qRT-PCR was

served with a StepOne Plus Real-time PCR System (Applied Biosystems, USA) and SYBR Green

Master Mix (Roche, Germany). Poplar *Actin* (*PtActin*) (XM-006370951.1) was previously tested

as a reference gene for this experiment (Zhang et al., 2013). The following conditions were

used for qRT-PCR reactions: pre-denaturation at 95°C for 10 min, 40 cycles of denaturation at

95°C for 15 s, and a chain extension at 60°C for 1 min. Three independent experiments were

conducted using gene-specific primers (Supplementary Table 1: PtHMGR forward and reverse).

2.7. *Metabolite analyses via high-performance liquid chromatography-tandem*
mass spectrometry

The isopropanol/acetic acid extraction method extracted poplar endogenous hormones
from NT, *PtDXR-OE*, and *PtHMGR-OE* leaves. GAs and CKs were extracted from, and then HPLC-
MS/MS (Qtrap6500, Agilent, USA) was used to quantify levels of GAs, zeatin, tZR, and IPA. Also,
methanol considered as solvent was used to extract 5-Deoxystrigol (5-DS), CS, and DCS, and
HPLC-MS/MS (Aglient1290, AB; SCIEX-6500Qtrap, Agilent; USA) was also used to determine
the contents of 5-DS, CS, and DCS. In addition, acetone, as a solvent, was used to isolate the
carotenoid component of poplar leaves. To identify the carotenoid contents, the peak areas
of carotenoids analyzed by HPLC (Symmetry Shield RP18, Waters, USA) were used to draw
standard carotenoid curves, including β-carotene, Lycopene, and Lutein. Also, the HPLC was
used to determine the contents of carotenoids, including β-carotene, Lycopene, and Lutein in
NT and OE lines.

3. Results

3.1. *Identification, analyses, and Isolation of PtHMGR and PtDXR genes*

Populus trichocarpa v3.1 (Phytozome genome ID: 444, NCBI taxonomy ID: 3694) has been
applied to download 595 amino acids (aa) PtHMGR (Potri.004G208500.1) and the other
species' HMGR to align and analyze. High similarity, including lots of conserved aa
accompanied by specific similar domains, HMG-CoA-binding motifs (EMPVGYVQIP' and
'TTEGCLVA), and NADPH-binding motifs (DAMGMNMV' and 'VGTVGGGT) (Ma et al., 2012)
(Supplementary Figure 1), confirmed the PtHMGR protein analytically. Consequently, a
phylogenetic tree based on the various species HMGR supported the PtHMGR candidate
identification (Supplementary Figure 2). The tblastn was then applied to reveal 2614
bp *PtHMGR* located on Chr04:21681480..21684242 with a 1785 bp CDS. After that, the
amplified 1857 bp of the *PtHMGR* from *Populus trichocarpa* cDNA confirmed the putative
transgenic lines (Supplementary Figure 3a), exhibiting amplicons in PCR identification
compared to NT poplar (Supplementary Figure 3b). The transgenic poplars (*PtHMGR-OEs*) also
showed enhanced *PtHMGR* expressions than NT (Supplementary Figure 3c), indicating
successful overexpression of *PtHMGR* in poplar. The *PtDXR* gene, which has been isolated,
sequenced, and analyzed previously by the authors (Xu et al., 2019), was then transferred into

poplar genome to generate PtDXR-OEs used in this study.

3.2.　*PtHMGR- and PtDXR overexpressions regulate MVA-related gene expressions*

MVA-related genes *AACT*, *MVK*, *MVD, and FPS,* except *HMGS,* were significantly
upregulated in *PtHMGR-OE* transgenics than NT poplars (Supplementary Figure 4a). In
contrast, while only *FPS* revealed significant upregulation by *PtDXR-OEs* in transgenics
compared with NT, the other MVA-related genes *AACT*, *HMGS*, *HMGR*, and *MVK* were
considerably downregulated (Supplementary Figure 4a). However, the mean comparison of
MVA-related gene expressions regulated by *HMGR* and *DXR* overexpressing exhibited
significant upregulated *MVK* through *PtHMGR-OEs* (Figure 1a). The *HMGS* was significantly
downregulated within *PtHMGR*-and *PtDXR-OEs*, and *MVK* was downregulated
by *PtDXR* overexpression (Figure 1a). The mean comparison of *AACT*, *MVD*, and *FPS* revealed
upregulation through *PtHMGR-OEs*, while *AACT* and *MVD* showed downregulation
through *PtDXR-OEs* (Figure 1a).

3.3.　*PtHMGR- and PtDXR overexpressions regulate MEP-related gene expressions*

While, the expression of MEP-related genes *DXS*, *DXR*, 1-hydroxy-2-methyl-2-(E)-butenyl-
4-diphosphate synthase (*HDS*), 1-hydroxy-2-methyl-2-(E)-butenyl-4-diphosphate reductase
(*HDR*), *IDI*, and *GPPS* were significantly upregulated in all *PtHMGR-OEs* transgenic poplars in
comparison with NT, the *GPS* overexpression was enhanced only by *PtHMGR-*
OE3 (Supplementary Figure 4c). In addition, 2-C-methyl-d-erythritol4-phosphate
cytidylyltransferase (*MCT*) and 4-diphosphocytidyl-2-C-methyl-D-erythritol kinase (*CMK*) have
been downregulated by *PtHMGR-OEs* (Supplementary Figure 4c). In contrast, all MEP-related
genes were upregulated significantly by *PtDXR* overexpression (Supplementary Figure 4d). In
total, the comparison of the effect of *PtHMGR* on MEP-related genes exhibited significant
upregulations of *HDR*, *IDI*, *GPS*, and *GPPS* (Figure 1b). These comparisons also revealed the
upregulation of *DXS* and *HDS* except *MCT* and *CMK*, downregulated by *PtHMGR*
overexpression (Figure 1b). In contrast, the mean comparison of the effect of *PtDXR*
overexpression on MEP-related genes exhibited upregulation of the all mentioned above
genes (Figure 1b).

3.4.   *MVA- and MEP-derived carotenoids are affected by PtHMGR-and PtDXR-OEs*

β-carotene is a carotenoid synthesis that has been broadly used in the industrial

composition of pharmaceuticals and as food colorants, animal supplies additives, and

nutraceuticals. MVA-and MEP pathways have been proved effective in the biosynthesis of β-

carotene (Yang, 2014). In addition, Lycopene is a carotenoid referring to C40 terpenoids and

is broadly found in various plants, particularly vegetables and fruits. It has been shown that

MVA and MEP-pathways directly influence the biosynthesis production of Lycopene (Kim et

al., 2019; Wei et al., 2018). While Wille et al. (2004) showed that β-carotene and Lutein are

synthesized using intermediates from the MEP pathway, Opitz et al. (2014) revealed that both

MVA and MPE pathways contribute to producing isoprenoids such as β-carotene and Lutein.

HPLC-MS/MS has analyzed the quantity of MVA and MEP derivatives. Our analyses revealed

that *HMGR*-OEs caused a significant enhancement in Lycopene (an average of ~ 0.08 ug/g), β-

carotene (an average of ~ 0.33 ug/g), and Lutein (an average of ~ 272 ug/g) production

compared with NT poplars (~0.02, ~0.08, and ~100 ug/g respectively) (Figure 2a, b, and c;

Supplementary Figure 5). The ABA-related gene expressions also have been calculated. Results

revealed a significantly increased *ZEP1*, *2*, and *3* relative gene expressions with averages of

~2.85, ~4.67, and ~2.92 compared to NT with an average of ~1 (Figure 2d). These results also

showed meaningful enhancements of *NCED1, 2,* and *3* relative gene expressions with the

averages of ~4.16, ~3.79, and ~3.4 compared to NT with an average of ~1 (Figure 2e).

On the other hand, the levels of the MEP-derived substances lycopene, β-carotene, and

Lutein were significantly increased in *PtDXR*-OEs with the averages of ~0.08, 0.22, 209.32 ug/g,

respectively compared to NT poplars (Figure 3a, b, and c; Supplementary Figure 6). The

analyses of ABA-related gene expressions revealed significantly increased *ZEP1*, *2*, and *3*

relative gene expressions with the averages of ~2.63, ~2.38, and ~3.86 compared to NT with

an average of ~1 (Figure 3d). These results also showed meaningful enhancements of *NCED2*

and *3* relative gene expressions with averages of ~2.25 and ~2.21 compared to NT with an

average of ~1 (Figure 2e). These results revealed a decreased average in *NCED1* relative gene

expression with an average of ~0.66 compared to NT poplars.

3.5.   *MVA and MEP-related derivatives are influenced by PtHMGR- and PtDXR-OEs*

The other MVA and MEP derivatives such as GAs, trans-zeatin-riboside (tZR), isopentenyl

adenosine (IPA), 6-deoxyocastasterone (DCS), and castasterone (CS) productions affected

by *PtHMGR-* and *PtDXR-OEs* have been analyzed. While Gibberellic acid (GA3) (a downstream

product of MEP) (an average of ~0.22 ng/g), tZR (an average of ~0.06 ng/g), IPA (an average

of ~0.59 ng/g), DCS (an average of 4.95 ng/g) revealed significantly more productions induced

by *HMGR*-OEs, the CS production (~0.095 ng/g) was decreased considerably compared to NT

poplars (~0.10, ~0.03, ~0.37, ~1.50, and ~0.20 ng/g respectively) (Figure 4a-j). These results

demonstrate that the *HMGR* gene interacts with MVA and MEP derivatives productions in

plants. On the other hand, the *PtDXR* overexpression significantly affected the contents of

MEP- and MVA-derived products except for CS. *PtDXR*-OEs showed a significant increase

~0.276 ng/g in the GA3 content (Figure 4a and f). The tZR content represented a 10-fold

increase (~0.304 ng/g) affected by *PtDXR-OEs* compared to NT poplars (0.032 ng/g) (Figure 4b

and g). The content of IPA in *PtDXR-OEs* meaningfully increased ~ 0.928 ng/g, compared to

0.363 ng/g in NT poplars (Figure 4c and h) with a 3-fold increase. In addition, the DCS content

considerably increased to ~3.36 ng/g, compared with ~1.50 ng/g in NT, representing a 3-fold

increase in *PtDXR-OEs* (Figure 4d and i). By contrast, the content of CS in *PtDXR-OEs*

significantly decreased (~0.137 ng/g) compared to NT poplar (0.203 ng/g), indicating

significant down-regulation in *PtDXR-OEs* (Figure 4e and j). The HPLC-MS/MS chromatograms

of GA, tZR, IPA, DCS, and CS standards are provided in Supplementary Figures 7–11.

3.6.   *Phenotypic properties*

To investigate the growth and development resulting from different produced

isoprenoids contents amongst the affected MVA-and MEP pathways contributions

by *PtHMGR*-and *PtDXR-OEs,* we decided to evaluate phenotypic stem lengths and diameters

changes. Results exhibited a significant increase in GA3 contents in *PtDXR-OEs* (Figure 4a)

associated with a considerable rise in cytokinin tZR (Figure 4b), resulting in significantly more

development in stem length compared to *PtHMGR-OEs* and NT poplars (Figure 5a and b).

Regarding increasing ABA-related gene expressions (*ZEP* and *NCED*) in *PtHMGR-OEs* than

PtDXR-OEs and NT poplars (Figure 5c and d) and also concerning insufficient increase cytokinin

tZR in *PtHMGR-OEs* compared with NT poplars (Figure 4b), *PtHMGR* transgenics showed a

shorter stem length that *PtDXR* transgenics compared with NT poplars (Figure 5a and b). We

also observed that only *PtDXR-OEs* revealed a few significant increases in stem diameters than

PtHMGR-OEs and NT poplars (Figure 5e).

4. Discussion

4.1. *The HMGR and DXR crucial roles in isoprenoid biosynthesis*

Several studies report that HMGR activity is regulated by isoprenoid outcomes when
stigmasterol and cholesterol reduce the HMGR activity by 35% (Russell and Davidson, 1982).
Utilization of the isoprenoid growth control abscisic acid also prevented HMGR activity in pea
by about 40%, while zeatin and gibberellin, other isoprenoid growth regulators, improved the
activity of HMGR (Russell and Davidson, 1982). Maurey et al. (1986) reported that the
alga *Ochromonas malhamensis* developed in mevinolin exhibited to 15-fold increase in
microsomal HMGR activity, slightly influencing cell growth. Moreover, the MEP pathway is the
primary precursor for required plastid isoprenoids (Wright et al., 2014). It has been shown
that volatile compounds made by the MEP pathway are involved in plant protection against
biotic and abiotic stresses (Gershenzon and Dudareva, 2007). By modifying the expression of
DXR, promising metabolite compounds have developed in the mint plant (Mahmoud and
Croteau, 2002). In addition, the DXR overexpression in Arabidopsis resulted in accumulating
isoprenoids such as tocopherols, carotenoids, and chlorophylls (Carretero-Paulet et al., 2006).
DXR overexpression has also been proven to improve diterpene contents in transgenic
bacteria (Morrone et al., 2010). Biotic stresses are vital in providing pharmaceutical
terpenoids by expanding the number of enzymes included in biosynthetic pathways by
controlling biosynthetic genes expression (Kang et al., 2009; Lu et al., 2016). Biotic stresses
caused to improve DXR expression followed by triptophenolide content in *Tripterygium*
wilfordii cell culture suspension (Tong et al., 2015).

4.2. *Overexpression of PtDXR results in upregulation of isoprenoid biosynthesis gene expression*

Liao et al. (2018) showed that overexpression of *BjHMGS1* affects the expression levels
of MEP- and MVA-related genes and slightly increases the transcript levels of *DXS* and *DXR* in
transgenic plants. However, *DXS*, *DXR*, *HDS*, and *HDR* expression levels have been
upregulated significantly in *PtHMGR-OE* poplars, while *MCT* and *CMK* are downregulated.
Similar to Liao et al. (2018) which the *BjHMGS1* overexpression in tomatoes significantly

increased the *GPS* and *GPPS* expressions, we exhibited that the *PtHMGR* overexpression

enhanced the farnesyl diphosphate synthase (*FPS*), *GPS*, and *GPPS* expressions may

stimulate the crosstalk between IPP and DMAPP, increasing the biosynthesis of plastidial C15

and C20 isoprenoid precursors. Xu et al. (2012) showed that *HMGR* overexpression

in *Ganoderma lucidum* caused upregulated *FPS*, squalene synthase (*SQS*), or lanosterol

synthase (LS) mRNA expressions and developed the contents of ganoderic acid and

intermediates, including squalene and lanosterol. In addition, the *BjHMGS1* overexpression in

tomatoes significantly increased transcript levels of *FPS*, *SQS*, squalene epoxidase (*SQE*), and

cycloartenol synthase (*CAS*) (Liao et al., 2018). This study exhibited that except

for *HMGS* downregulating, the *AACT*, *MVK*, and *MVD* transcript levels were significantly

upregulated in *PtHMGR-OE* poplars. We revealed that these enhanced gene expressions

mainly were associated with the MVA-related genes contributing to the biosynthesis of

sesquiterpenes and other C15 and universal C20 isoprenoid precursors.

4.3.   *Overexpression of PtDXR affects MEP- and MVA-related genes*

Zhang et al. (2018) showed that the *TwDXR* overexpression in *Tripterygium wilfordii*

increases the *TwHMGS*, *TwHMGR*, *TwFPS*, and *TwGPPS* expressions but decreases the *TwDXS*

expression. Moreover, Zhang et al. (2015) exhibited that the *NtDXR1* overexpression in

tobacco increases the transcript levels of eight MEP-related genes, indicating that the *NtDXR1*

overexpression led to upregulated MEP-related gene expressions. In *A. thaliana*, the *DXR*

transcript level changes do not affect *DXS* gene expression or enzyme accumulation, although

the *DXR* overexpression promotes MEP-derived isoprenoids such as carotenoids, chlorophylls,

and taxadiene (Carretero-Paulet et al., 2006).

On the other hand, the potato *DXS* overexpression in *A. thaliana* led to upregulation of

downstream *GGPPS* and phytoene synthase (*PSY*) genes (Henriquez et al., 2016). Furthermore,

(Simpson et al., 2016) exhibited that the *A. thaliana DXS* overexpression in Daucus carota

caused to enhance the *PSY* expression significantly.

In this study, while the *PtDXR-OEs* exposed higher MEP-related gene expressions than NT

poplars, the *PtDXR-OEs* revealed significant downregulated MVA-related gene expressions

than NT poplars. These findings illustrate that the MEP pathway regulates monoterpenes,

diterpenes, and tetraterpenoids biosynthesis and could affect the MVA pathway.

The diversity of biosynthetic pathways, the complexity of metabolic networks, and the

insufficient knowledge of gene regulation led to species-specific regulation patterns of MEP-

and MVA-related gene expression. One possible conclusion is that MEP- and MVA-related

genes often do not work alone but are co-expressed with upstream and downstream genes in

the MEP- and MVA- pathways to carry out a specific function.

4.4.   *Overexpression of HMGR promotes the formation of GAs, and carotenoids in*

393   *plastids and accumulation of tZR, IPA, and DCS in the cytoplasm*

HMGR, as the rate-limiting enzyme in the MVA-pathway of plants, plays a critical role in

controlling the flow of carbon within this metabolic pathway. The upregulation

of *HMGR* significantly increases isoprenoid levels in plants. Overexpression of *HMGRs* of

different plant species has been reported to raise isoprenoids levels significantly. The

heterologous expression of *Hevea brasiliensis HMGR1* in tobacco increased the sterol content

and accumulated intermediate metabolites (Schaller et al., 1995). The *A. thaliana HMGR*

*o*verexpression in *Lavandula latifolia* increased the levels of sterols in the MVA-and MEP-

derived monoterpenes and sesquiterpenes (Munoz-Bertomeu et al., 2007). In addition,

the *Salvia miltiorrhiza SmHMGR* overexpression in hairy roots developed MEP-derived

diterpene tanshinone (Kai et al., 2011). In our study, ABA synthesis-related genes

(*NCED1*, *NCED3*, *NCED6*, *ZEP1*, *ZEP2*, and *ZEP3*) and the contents of GA3 and carotenoids

were   upregulated   in *PtHMGR-OE*   poplar   seedlings.   This   finding   suggests   that

the *HMGR* overexpression may indirectly affect the biosynthesis of MEP-related isoprenoids,

including GA3 and carotenoids. The accumulation of MVA-derived isoprenoids including tZR,

IPA,   and   DCS   was   significantly   elevated   in *PtHMGR-OEs*,   indicating

that *PtHMGR* overexpression directly influences the biosynthesis of MVA-related isoprenoids.

Therefore, the *HMGR* gene directly affects MVA-derived isoprenoids and indirectly affects the

content of MEP-derived isoprenoids by changing the expression levels of MEP-related genes.

4.5.   *Higher levels of MEP- and MVA-derived products in PtDXR-OE seedlings*

DXR is the rate-limiting enzyme in the MEP pathway and an essential regulatory step in

the cytoplasmic metabolism of isoprenoid compounds (Takahashi et al., 1998). Mahmoud and

Croteau (2001) revealed that overexpression of *DXR* in *Mentha piperita* promoted the

synthesis of monoterpenes in the oil glands and increased the production of essential oil yield

by 50%. However, the up-regulation of *DXR* expression did not lead to change in the complex oil composition significantly. Hasunuma et al. (2008) exhibited that overexpression of *Synechocystis sp*. strain PCC6803 *DXR* in tobacco resulted in increased levels of β-carotene, chlorophyll, antheraxanthin, and Lutein. Xing et al. (2010) showed that the *A. thaliana dxr* mutants caused to lack of GAs, ABA, and photosynthetic pigments (REF57). These mutants showed pale sepals and yellow inflorescences (Xing et al., 2010). In our study, the relatively higher abundance of GA3 and carotenoids in *PtDXR-OE* poplar seedlings indicated an effect of *DXR* overexpression. Combined with the result described above of increased *DXS*, *HDS*, *HDR*, *MCT*, *CMK*, *FPS*, *GPS*, and *GPPS* expression levels, we postulate that overexpression of *DXR* not only affects the expression levels of MEP-related genes but also changes the field of GA3, and carotenoids.

4.6. *Communications exist between MVA- and MEP-pathways excess of IPP and DMAPP*

Although the substrates of MVA- and MEP pathways differ, there are common intermediates like IPP and DMAPP (Figure 6). Blocking only the MVA or the MEP pathway, respectively, does not entirely prevent the biosynthesis of terpenes in the cytoplasm or plastids, indicating that some MVA and MEP pathways products can be transported and/or move between cell compartments (Aharoni et al., 2003; Aharoni et al., 2004; Gutensohn et al., 2013). For example, it has been shown that the transferring IPP from the chloroplast to cytoplasm observed through 13C labeling, indicating that plentiful IPP is available for use in the MVA-pathway to produce terpenoids (Ma et al., 2017). In addition, segregation between the MVA- and MEP pathways is limited and might exchange some metabolites over the plastid membrane (Laule, 2003). Kim et al. (2016b) used clustered, regularly interspaced short palindromic repeats (CRISPR) technology to reconstruct the lycopene synthesis pathway and control the flow of carbon in the MEP-and MVA-pathways. The results showed that the expression of MVA-related genes was reduced by 81.6%, but the lycopene yield was significantly increased. By analyzing gene expression levels and metabolic outcome in *PtHMGR*-and *PtDXR-OEs*, we discovered that the correlation might exist between MVA- and MEP-related genes with MVA- and MEP-derived products, which are not restricted to crosstalk between IPP and DMAPP (Figure 6).

On the one hand, overexpression of *PtDXR* affected the transcript levels of MEP-related

genes and the contents of MEP-derived isoprenoids, including GAs and carotenoids. The

diminished accumulation of MVA-related gene products causes a reduction in the yields of

MVA-derived isoprenoids (including CS) but leads to increasing tZR, IPA, and DCS contents. We

hypothesize that IPP and DMAPP produced by the MEP pathway could enter the cytoplasm to

compensate for the lack of IPP and DMAPP, and the IPP and DMAPP as the precursors of the

MVA pathway are used to guide the synthesis of MVA-derived products. On the other

hand, *PtHMGR-OEs* exhibited higher transcript levels of *AACT*, *MVK*, and *MVD* and

higher *DXS*, *DXR*, *HDS*, and *HDR* than NT poplars, resulting in effect both MEP- and MVA-

related gene expressions. We successfully demonstrated that manipulation of *HMGR* in the

poplar MVA pathway results in dramatically enhanced yields of GAs and carotenoids. This

result illustrates that cytosolic *HMGR* overexpression expanded plastidial GPP- and GGPP-

derived products, such as carotenoids. Therefore, this study provides hints that

communications between the MVA-and MEP pathways increased the expression levels

of *GPS* and *GPPS* in *PtHMGR-OEs*, and elevated the contents of GA3 and carotenoids.

Moreover, changes in MEP- and MVA-related gene expressions affect MVA- and MEP-derived

isoprenoids.

In conclusion, overexpression of *PtHMGR* in poplars caused the accumulation of MVA-

derived isoprenoids and MEP-derived substances. The advanced insights into the regulation

of MVA- and MEP pathways in poplar add to the knowledge about these pathways in

Arabidopsis, tomato, and rice. In *PtHMGR-OE* poplars, most MEP- and MVA-related genes

associated with the biosynthesis of isoprenoid precursors were upregulated. In *PtDXR-*

OE poplars, elevated contents of GAs, carotenoids, and GRs were attributed to increased

expression of MEP-related genes as well as plastidial *GPP* and *GGPP*. Together, these results

show that manipulating *PtDXR and PtHMGR* is a novel strategy to influence poplar isoprenoids.

4.7.   *Communications between MVA- and MEP pathways affected by PtHMGR- and*
473   *PtDXR-OEs influence the plant growth and developments*

It has been shown that Abscisic acid (ABA) and GA3 perform essential functions in cell

division, shoot growth, and flower induction (Xing et al., 2016). It has also been demonstrated

that the cytokinin tZR, a variety of phytohormones, perform a crucial function as root to shoot

signals, directing numerous developmental and growth processes in shoots (Abul et al., 2010;

Sakakibara, 2006). Regarding these findings, we showed how the communications between

MVA- and MEP pathways and their changes affected by some stimulators (*HMGR*-and *DXR*-

OEs) influenced plant growth, especially in stem length. Finally, We figured out that the

gibberellic acid and cytokinin may be more effective in plant growth than inhibiting by ABA,

causing higher *PtDXR-OEs* than *PtHMGR-OEs* compared with NT poplars.

Author contributions

A.M. and H.W. conceived, planned, and coordinated the project, performed data analysis,

wrote the draft, and finalized the manuscript. B.P. validated and contributed to data analysis

and curation, revised and finalized the manuscript. T.J., W.S., and D.L. reviewed and edited the

manuscript. L.Y. and Q.Z. coordinated, contributed to data curation, finalized and funded this

research. A.M., H.W., and B.P. contributed equally as the first author.

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgments

This work was supported by the National Key Program on Transgenic Research

(2018ZX08020002), the National Natural Science Foundation of China (No. 31971682).

5. Reference

Abul, Y., Menendez, V., Gomez-Campo, C., Revilla, M.A., Lafont, F., Fernandez, H., 2010. Occurrence of plant
growth regulators in Psilotum nudum. J Plant Physiol 167, 1211-1213.

Aharoni, A., Giri, A.P., Deuerlein, S., Griepink, F., de Kogel, W.J., Verstappen, F.W., Verhoeven, H.A., Jongsma,
M.A., Schwab, W., Bouwmeester, H.J., 2003. Terpenoid metabolism in wild-type and transgenic Arabidopsis
plants. Plant Cell 15, 2866-2884.

Aharoni, A., Giri, A.P., Verstappen, F.W., Bertea, C.M., Sevenier, R., Sun, Z., Jongsma, M.A., Schwab, W.,
Bouwmeester, H.J., 2004. Gain and loss of fruit flavor compounds produced by wild and cultivated strawberry
species. Plant Cell 16, 3110-3131.

Aharoni, A., Jongsma, M.A., Bouwmeester, H.J., 2005. Volatile science? Metabolic engineering of terpenoids in
plants. Trends Plant Sci 10, 594-602.

Bohlmann, J., Keeling, C.I., 2008. Terpenoid biomaterials. Plant J 54, 656-669.

Boutte, Y., Grebe, M., 2009. Cellular processes relying on sterol function in plants. Curr Opin Plant Biol 12, 705-
713.

Bouvier, F., Rahier, A., Camara, B., 2005. Biogenesis, molecular regulation and function of plant isoprenoids.
Prog Lipid Res 44, 357-429.

Carretero-Paulet, L., Cairo, A., Botella-Pavia, P., Besumbes, O., Campos, N., Boronat, A., Rodriguez-Concepcion,
M., 2006. Enhanced flux through the methylerythritol 4-phosphate pathway in Arabidopsis plants overexpressing
deoxyxylulose 5-phosphate reductoisomerase. Plant Mol Biol 62, 683-695.

Cordoba, E., Salmi, M., Leon, P., 2009. Unravelling the regulatory mechanisms that modulate the MEP pathway in higher plants. J Exp Bot 60, 2933-2943.

Cowan, A.K., Moore-Gordon, C.S., Bertling, I., Wolstenholme, B.N., 1997. Metabolic Control of Avocado Fruit Growth (Isoprenoid Growth Regulators and the Reaction Catalyzed by 3-Hydroxy-3-Methylglutaryl Coenzyme A Reductase). Plant Physiol 114, 511-518.

Dai, Z., Cui, G., Zhou, S.F., Zhang, X., Huang, L., 2011. Cloning and characterization of a novel 3-hydroxy-3-methylglutaryl coenzyme A reductase gene from Salvia miltiorrhiza involved in diterpenoid tanshinone accumulation. J Plant Physiol 168, 148-157.

Devappa, R.K., Rakshit, S.K., Dekker, R.F.,, 2015. Forest biorefinery: Potential of poplar phytochemicals as value-added co-products. Biotechnol Adv 33, 681-716.

Dueber, J.E., Wu, G.C., Malmirchegini, G.R., Moon, T.S., Petzold, C.J., Ullal, A.V., Prather, K.L., Keasling, J.D., 2009. Synthetic protein scaffolds provide modular control over metabolic flux. Nat Biotechnol 27, 753-759.

Enfissi, E.M., Fraser, P.D., Lois, L.M., Boronat, A., Schuch, W., Bramley, P.M., 2005. Metabolic engineering of the mevalonate and non-mevalonate isopentenyl diphosphate-forming pathways for the production of health-promoting isoprenoids in tomato. Plant Biotechnol J 3, 17-27.

Esteban, R., Barrutia, O., Artetxe, U., Fernandez-Marin, B., Hernandez, A., Garcia-Plazaola, J.I., 2015. Internal and external factors affecting photosynthetic pigment composition in plants: a meta-analytical approach. New Phytol 206, 268-280.

Gershenzon, J., Dudareva, N., 2007. The function of terpene natural products in the natural world. Nat Chem Biol 3, 408-414.

Ghirardo, A., Wright, L.P., Bi, Z., Rosenkranz, M., Pulido, P., Rodriguez-Concepcion, M., Niinemets, U., Bruggemann, N., Gershenzon, J., Schnitzler, J.P., 2014. Metabolic flux analysis of plastidic isoprenoid biosynthesis in poplar leaves emitting and nonemitting isoprene. Plant Physiol 165, 37-51.

Gutensohn, M., Orlova, I., Nguyen, T.T., Davidovich-Rikanati, R., Ferruzzi, M.G., Sitrit, Y., Lewinsohn, E., Pichersky, E., Dudareva, N., 2013. Cytosolic monoterpene biosynthesis is supported by plastid-generated geranyl diphosphate substrate in transgenic tomato fruits. Plant J 75, 351-363.

Hain, R., Reif, H.J., Krause, E., Langebartels, R., Kindl, H., Vornam, B., Wiese, W., Schmelzer, E., Schreier, P.H., Stocker, R.H., et al., 1993. Disease resistance results from foreign phytoalexin expression in a novel plant. Nature 361, 153-156.

Hasunuma, T., Takeno, S., Hayashi, S., Sendai, M., Bamba, T., Yoshimura, S., Tomizawa, K., Fukusaki, E., Miyake, C., 2008. Overexpression of 1-Deoxy-D-xylulose-5-phosphate reductoisomerase gene in chloroplast contributes to increment of isoprenoid production. J Biosci Bioeng 105, 518-526.

Hemmerlin, A., Harwood, J. L., & Bach, T. J., 2012. A raison d'être for two distinct pathways in the early steps of plant isoprenoid biosynthesis? Prog. Lipid Res. 51, 95–148.

Henriquez, M.A., Soliman, A., Li, G., Hannoufa, A., Ayele, B.T., Daayf, F., 2016. Molecular cloning, functional characterization and expression of potato (Solanum tuberosum) 1-deoxy-d-xylulose 5-phosphate synthase 1 (StDXS1) in response to Phytophthora infestans. Plant Sci 243, 71-83.

Henry, L.K., Gutensohn, M., Thomas, S.T., Noel, J.P., Dudareva, N., 2015. Orthologs of the archaeal isopentenyl phosphate kinase regulate terpenoid production in plants. Proc Natl Acad Sci U S A 112, 10050-10055.

Huchelmann, A., Gastaldo, C., Veinante, M., Zeng, Y., Heintz, D., Tritsch, D., Schaller, H., Rohmer, M., Bach, T.J., Hemmerlin, A., 2014. S-carvone suppresses cellulase-induced capsidiol production in Nicotiana tabacum by interfering with protein isoprenylation. Plant Physiol 164, 935-950.

Kai, G., Xu, H., Zhou, C., Liao, P., Xiao, J., Luo, X., You, L., Zhang, L., 2011. Metabolic engineering tanshinone biosynthetic pathway in Salvia miltiorrhiza hairy root cultures. Metab Eng 13, 319-327.

Kang, S.M., Min, J.Y., Kim, Y.D., Karigar, C.S., Kim, S.W., Goo, G.H., Choi, M.S., 2009. Effect of biotic elicitors on the accumulation of bilobalide and ginkgolides in Ginkgo biloba cell cultures. J Biotechnol 139, 84-88.

Kim, M.J., Noh, M.H., Woo, S., Lim, H.G., Jung, G.Y., 2019. Enhanced Lycopene Production in Escherichia coli

by Expression of Two MEP Pathway Enzymes from Vibrio sp. Dhg. Catalysts 9.

Kim, M.S., Haney, M.J., Zhao, Y., Mahajan, V., Deygen, I., Klyachko, N.L., Inskoe, E., Piroyan, A., Sokolsky, M.,

Okolie, O., Hingtgen, S.D., Kabanov, A.V., Batrakova, E.V., 2016a. Development of exosome-encapsulated

paclitaxel to overcome MDR in cancer cells. Nanomedicine 12, 655-664.

Kim, S.K., Han, G.H., Seong, W., Kim, H., Kim, S.W., Lee, D.H., Lee, S.G., 2016b. CRISPR interference-guided

balancing of a biosynthetic mevalonate pathway increases terpenoid production. Metab Eng 38, 228-240.

Kim, Y.J., Lee, O. R., Ji, Y. O., Jang, M. G., & Yang, D. C., 2014. Functional analysis of HMGR encoding genes

in triterpene saponin-producing Panax ginseng Meyer. Plant Physiol 165, 373–387.

Kirby, J., Keasling, J.D., 2009. Biosynthesis of plant isoprenoids: perspectives for microbial engineering. Annu

Rev Plant Biol 60, 335-355.

Kong, L.Y., Tan, R.X., 2015. Artemisinin, a miracle of traditional Chinese medicine. Nat Prod Rep 32, 1617-1621.

Laule, O., Fürholz, A., Chang, H. S., Zhu, T., Wang, X., Heifetz, P. B., ... & Lange, M., 2003. Cross-talk between

cytosolic and plastidial pathways of isoprenoid biosynthesis in Arabidopsis thaliana. Proc. Natl. Acad. Sci. U. S.

A., 6866–6871.

Liao, P., Chen, X., Wang, M., Bach, T.J., Chye, M.L., 2018. Improved fruit alpha-tocopherol, carotenoid, squalene

and phytosterol contents through manipulation of Brassica juncea 3-HYDROXY-3-METHYLGLUTARYL-COA

SYNTHASE1 in transgenic tomato. Plant Biotechnol J 16, 784-796.

Liao, P., Hemmerlin, A., Bach, T.J., Chye, M.L., 2016. The potential of the mevalonate pathway for enhanced

isoprenoid production. Biotechnol Adv 34, 697-713.

Liao, Z.H., Chen, M., Gong, Y. F., Miao, Z. Q., Sun, X. F., & Tang, K. X., 2006. Isoprenoid biosynthesis in plants:

pathways, genes, regulation and metabolic engineering. J Biol Sci 6, 209–219.

Lu, X., Tang, K., Li, P., 2016. Plant Metabolic Engineering Strategies for the Production of Pharmaceutical

Terpenoids. Front Plant Sci 7, 1647.

Lu, X.M., Hu, X.J., Zhao, Y.Z., Song, W.B., Zhang, M., Chen, Z.L., Chen, W., Dong, Y.B., Wang, Z.H., Lai, J.S.,

2012. Map-based cloning of zb7 encoding an IPP and DMAPP synthase in the MEP pathway of maize. Mol Plant

5, 1100-1112.

Ma, D., Li, G., Zhu, Y., Xie, D.Y., 2017. Overexpression and Suppression of Artemisia annua 4-Hydroxy-3-

Methylbut-2-enyl Diphosphate Reductase 1 Gene (AaHDR1) Differentially Regulate Artemisinin and Terpenoid

Biosynthesis. Front Plant Sci 8, 77.

Ma, Y., Yuan, L., Wu, B., Li, X., Chen, S., Lu, S., 2012. Genome-wide identification and characterization of novel

genes involved in terpenoid biosynthesis in Salvia miltiorrhiza. J Exp Bot 63, 2809-2823.

Mahmoud, S.S., Croteau, R.B., 2001. Metabolic engineering of essential oil yield and composition in mint by

altering expression of deoxyxylulose phosphate reductoisomerase and menthofuran synthase. Proc Natl Acad Sci

U S A 98, 8915-8920.

Mahmoud, S.S., Croteau, R.B., 2002. Strategies for transgenic manipulation of monoterpene biosynthesis in plants.

Trends Plant Sci 7, 366-373.

Malkin R, a.N.K., 2000. Photosynthesis. American Society of Plant Physiologists, Rockville.

Maurey, K., Wolf, F., Golbeck, J., 1986. 3-Hydroxy-3-Methylglutaryl Coenzyme A Reductase Activity in

Ochromonas malhamensis: A System to Study the Relationship between Enzyme Activity and Rate of Steroid

Biosynthesis. Plant Physiol 82, 523-527.

Morrone, D., Lowry, L., Determan, M.K., Hershey, D.M., Xu, M., Peters, R.J., 2010. Increasing diterpene yield

with a modular metabolic engineering system in E. coli: comparison of MEV and MEP isoprenoid precursor

pathway engineering. Appl Microbiol Biotechnol 85, 1893-1906.

Movahedi, A., Zhang, J., Amirian, R., Zhuge, Q., 2014. An efficient Agrobacterium-mediated transformation

system for poplar. Int J Mol Sci 15, 10780-10793.

Movahedi, A., Zhang, J., Sun, W., Mohammadi, K., Almasi Zadeh Yaghuti, A., Wei, H., Wu, X., Yin, T., Zhuge,

Q., 2018. Functional analyses of PtRDM1 gene overexpression in poplars and evaluation of its effect on DNA

methylation and response to salt stress. Plant Physiol Biochem 127, 64-73.

Movahedi, A., Zhang, J.X., Yin, T.M., Qiang, Z.G., 2015. Functional Analysis of Two Orthologous NAC Genes,
CarNAC3, and CarNAC6 from Cicer arietinum, Involved in Abiotic Stresses in Poplar. Plant Molecular Biology
Reporter 33, 1539-1551.

Munoz-Bertomeu, J., Sales, E., Ros, R., Arrillaga, I., Segura, J., 2007. Up-regulation of an N-terminal truncated
3-hydroxy-3-methylglutaryl CoA reductase enhances production of essential oils and sterols in transgenic
Lavandula latifolia. Plant Biotechnol J 5, 746-758.

Opitz, S., Nes, W.D., Gershenzon, J., 2014. Both methylerythritol phosphate and mevalonate pathways contribute
to biosynthesis of each of the major isoprenoid classes in young cotton seedlings. Phytochemistry 98, 110-119.

Perreca, E., Rohwer, J., Gonzalez-Cabanelas, D., Loreto, F., Schmidt, A., Gershenzon, J., Wright, L.P., 2020.
Effect of Drought on the Methylerythritol 4-Phosphate (MEP) Pathway in the Isoprene Emitting Conifer Picea
glauca. Front Plant Sci 11, 546295.

Rahman, L., Kouno, H., Hashiguchi, Y., Yamamoto, H., Narbad, A., Parr, A., Walton, N., Ikenaga, T., Kitamura,
Y., 2009. HCHL expression in hairy roots of Beta vulgaris yields a high accumulation of p-hydroxybenzoic acid
(pHBA) glucose ester, and linkage of pHBA into cell walls. Bioresour Technol 100, 4836-4842.

Ren, D., Liu, Y., Yang, K.Y., Han, L., Mao, G., Glazebrook, J., Zhang, S., 2008. A fungal-responsive MAPK
cascade regulates phytoalexin biosynthesis in Arabidopsis. Proc Natl Acad Sci U S A 105, 5638-5643.

Roberts, S.C., 2007. Production and engineering of terpenoids in plant cell culture. Nat Chem Biol 3, 387-395.

Rodriguez-Concepcion, M., 2010. Supply of precursors for carotenoid biosynthesis in plants. Arch Biochem
Biophys 504, 118-122.

Russell, D.W., Davidson, H., 1982. Regulation of cytosolic HMG-CoA reductase activity in pea seedlings:
contrasting responses to different hormones, and hormone-product interaction, suggest hormonal modulation of
activity. Biochem Biophys Res Commun 104, 1537-1543.

Sakakibara, H., 2006. Cytokinins: activity, biosynthesis, and translocation. Annu Rev Plant Biol 57, 431-449.

Schaller, H., Grausem, B., Benveniste, P., Chye, M.L., Tan, C.T., Song, Y.H., Chua, N.H., 1995. Expression of the
Hevea brasiliensis (H.B.K.) Mull. Arg. 3-Hydroxy-3-Methylglutaryl-Coenzyme A Reductase 1 in Tobacco Results
in Sterol Overproduction. Plant Physiol 109, 761-770.

Simpson, K., Quiroz, L.F., Rodriguez-Concepcion, M., Stange, C.R., 2016. Differential Contribution of the First
Two Enzymes of the MEP Pathway to the Supply of Metabolic Precursors for Carotenoid and Chlorophyll
Biosynthesis in Carrot (Daucus carota). Front Plant Sci 7, 1344.

Song, X., Yu, X., Hori, C., Demura, T., Ohtani, M., Zhuge, Q., 2016. Heterologous Overexpression of Poplar
SnRK2 Genes Enhanced Salt Stress Tolerance in Arabidopsis thaliana. Front Plant Sci 7, 612.

Takahashi, S., Kuzuyama, T., Watanabe, H., Seto, H., 1998. A 1-deoxy-D-xylulose 5-phosphate reductoisomerase
catalyzing the formation of 2-C-methyl-D-erythritol 4-phosphate in an alternative nonmevalonate pathway for
terpenoid biosynthesis. Proc Natl Acad Sci U S A 95, 9879-9884.

Thulasiram, H.V., Erickson, H.K., Poulter, C.D., 2007. Chimeras of two isoprenoid synthases catalyze all four
coupling reactions in isoprenoid biosynthesis. Science 316, 73-76.

Tong, Y., Su, P., Zhao, Y., Zhang, M., Wang, X., Liu, Y., Zhang, X., Gao, W., Huang, L., 2015. Molecular Cloning
and Characterization of DXS and DXR Genes in the Terpenoid Biosynthetic Pathway of Tripterygium wilfordii.
Int J Mol Sci 16, 25516-25535.

Vaccaro, M., Malafronte, N., Alfieri, M., De Tommasi, N., Leone, A., 2014. Enhanced biosynthesis of bioactive
abietane diterpenes by overexpressing AtDXS or AtDXR genes in Salvia sclarea hairy roots. Plant Cell Tissue and
Organ Culture 119, 65-77.

van Schie, C.C., Haring, M.A., Schuurink, R.C., 2006. Regulation of terpenoid and benzenoid production in
flowers. Curr Opin Plant Biol 9, 203-208.

Wang, H., Nagegowda, D.A., Rawat, R., Bouvier-Nave, P., Guo, D., Bach, T.J., Chye, M.L., 2012. Overexpression
of Brassica juncea wild-type and mutant HMG-CoA synthase 1 in Arabidopsis up-regulates genes in sterol

biosynthesis and enhances sterol production and stress tolerance. Plant Biotechnol J 10, 31-42.

Wei, Y., Mohsin, A., Hong, Q., Guo, M., Fang, H., 2018. Enhanced production of biosynthesized lycopene via
heterogenous MVA pathway based on chromosomal multiple position integration strategy plus plasmid systems
in Escherichia coli. Bioresour Technol 250, 382-389.

Wille, A., Zimmermann, P., Vranova, E., Furholz, A., Laule, O., Bleuler, S., Hennig, L., Prelic, A., von Rohr, P.,
Thiele, L., Zitzler, E., Gruissem, W., Buhlmann, P., 2004. Sparse graphical Gaussian modeling of the isoprenoid
gene network in Arabidopsis thaliana. Genome Biol 5, R92.

Wright, L.P., Rohwer, J.M., Ghirardo, A., Hammerbacher, A., Ortiz-Alcaide, M., Raguschke, B., Schnitzler, J.P.,
Gershenzon, J., Phillips, M.A., 2014. Deoxyxylulose 5-Phosphate Synthase Controls Flux through the
Methylerythritol 4-Phosphate Pathway in Arabidopsis. Plant Physiol 165, 1488-1504.

Xie, Z., Kapteyn, J., Gang, D.R., 2008. A systems biology investigation of the MEP/terpenoid and
shikimate/phenylpropanoid pathways points to multiple levels of metabolic control in sweet basil glandular
trichomes. Plant J 54, 349-361.

Xing, L., Zhang, D., Zhao, C., Li, Y., Ma, J., An, N., Han, M., 2016. Shoot bending promotes flower bud formation
by miRNA-mediated regulation in apple (Malus domestica Borkh.). Plant Biotechnol J 14, 749-770.

Xing, S., Miao, J., Li, S., Qin, G., Tang, S., Li, H., Gu, H., Qu, L.J., 2010. Disruption of the 1-deoxy-D-xylulose-
5-phosphate reductoisomerase (DXR) gene results in albino, dwarf and defects in trichome initiation and stomata
closure in Arabidopsis. Cell Res 20, 688-700.

Xu, C., Wei, H., Movahedi, A., Sun, W., Ma, X., Li, D., Yin, T., Zhuge, Q., 2019. Evaluation, characterization,
expression profiling, and functional analysis of DXS and DXR genes of Populus trichocarpa. Plant Physiol
Biochem 142, 94-105.

Xu, J.W., Xu, Y.N., Zhong, J.J., 2012. Enhancement of ganoderic acid accumulation by overexpression of an N-
terminally truncated 3-hydroxy-3-methylglutaryl coenzyme A reductase gene in the basidiomycete Ganoderma
lucidum. Appl Environ Microbiol 78, 7968-7976.

Yamaguchi, S., Kamiya, Y., & Nambara, E., 2018. Regulation of ABA and GA levels during seed development
and germination in Arabidopsis. Annu Plant Rev 27, 224–247.

Yang, J., Guo, L., 2014. Biosynthesis of β-carotene in engineered E. coli using the MEP and MVA pathways.
Microb Cell Fact, 160.

Zhang, H., Niu, D., Wang, J., Zhang, S., Yang, Y., Jia, H., Cui, H., 2015. Engineering a Platform for Photosynthetic
Pigment, Hormone and Cembrane-Related Diterpenoid Production in Nicotiana tabacum. Plant Cell Physiol 56,
2125-2138.

Zhang, J., Li, J., Liu, B., Zhang, L., Chen, J., Lu, M., 2013. Genome-wide analysis of the Populus Hsp90 gene
family reveals differential expression patterns, localization, and heat stress responses. BMC Genomics 14, 532.

Zhang, K.K., Fan, W., Huang, Z.W., Chen, D.F., Yao, Z.W., Li, Y.F., Yang, Y.F., Qiu, D.Y., 2019. Transcriptome
analysis identifies novel responses and potential regulatory genes involved in 12-deoxyphorbol-13-phenylacetate
biosynthesis of Euphorbia resinifera. Industrial Crops and Products 135, 138-145.

Zhang, L., Ding, R., Chai, Y., Bonfill, M., Moyano, E., Oksman-Caldentey, K.M., Xu, T., Pi, Y., Wang, Z., Zhang,
H., Kai, G., Liao, Z., Sun, X., Tang, K., 2004. Engineering tropane biosynthetic pathway in Hyoscyamus niger
hairy root cultures. Proc Natl Acad Sci U S A 101, 6786-6791.

Zhang, Y., Zhao, Y., Wang, J., Hu, T., Tong, Y., Zhou, J., Song, Y., Gao, W., Huang, L., 2018. Overexpression and
RNA interference of TwDXR regulate the accumulation of terpenoid active ingredients in Tripterygium wilfordii.
Biotechnol Lett 40, 419-425.

6. Figure legends

Figure 1 | MVA- and MEP-related genes analyses in overexpressed *PtHMGR*- and *PtDXR-OE s* poplars. a, Mean comparison of relative expression of MVA-relate genes *AACT*, *HMGS*, *MVK*, *MVD*, and *FPS* (Indicated in red) affected by *PtHMGR* overexpressing. **b**, Mean comparison of relative expression of MEP-related genes *DXS*, *MCT*, *CMK*, *HDS*, *HDR*, *IDI*, *GPS*, and *GPPS* (Indicated in red) affected by *PtHMGR* overexpressing. *HMGR* and *DXR*, which were overexpressed respectively among *DXR*- and *HMGR-OEs*, were presented in Supplementary Figure 4. *PtActin* was used as an internal reference in all repeats; "ns" means not significant, * P < 0.05, ** P < 0.01, ***P < 0.001, ****P < 0.0001; Three independent replications were p erformed in this experiment.

Figure 2 | HPLC-MS/MS content analyses of lycopene, β-carotene, Lutein, and real-time PCR of *ZEP* and *NCED* genes family. HPLC-MS/MS content analyses have been performed to show the effect of *PtHMGR-OEs* on **a***,* lycopene **b**, β-carotene, and **c**, lutein. Relative expressions have been analyzed affected by *PtHMGR-OEs* compared with NT poplars of **d**, *ZEP*, and **e**, *NCED* genes family. Bars represent mean ± SD (n = 3); Stars reveal significant differences, * P < 0.05, ** P < 0.01, *** P < 0.001, ****P < 0.0001; Three independent experiments were performed in these analyses.

Figure 3 | HPLC-MS/MS content analyses of lycopene, β-carotene, Lutein, and real-time PCR of *ZEP* and *NCED* genes family. HPLC-MS/MS content analyses have been performed to show the effect of *PtDXR-OEs* on **a***,* lycopene **b**, β-carotene, and **c**, lutein. Relative expressions have been analyzed affected by *PtDXR-OEs* compared with NT poplars of **d**, *ZEP*, and **e**, *NCED* genes family. Bars represent mean ± SD (n = 3); Stars reveal significant differences, * P < 0.05, ** P < 0.01, *** P < 0.001, ****P < 0.0001; Three independent experiments were performed in these analyses.

Figure 4 | HPLC-MS/MS content analyses of MEP- and MVA-derived isoprenoids. a,b,c,d, and e, Violin plots reveal the contents of isoprenoids GA3, tZR, IPA, DCS, and CS obtained from MEP- and MVA pathways influenced by *PtHMGR*- and *PtDXR-OEs*. **f,g,h,i, and j**, the column plots reveal the effect of *PtHMGR-OE3* and *-7* and *PtDXR-OE1* and *-3* on the mentioned above isoprenoids separately; NT poplars have been used as the control. Bars represent mean ± SD (n = 3); Stars reveal significant differences, *P < 0.05, **P < 0.01, ***P < 0.001, ****P <

0.0001. k,l,m,n, and o, represent the HPLC-MS/MS chromatogram content analyses of GA3,

tZR, IPA, DCS, and CS, respectively affected by *PtHMGR-* and *PtDXR-OEs* compared with NT

poplars.

Figure 5 | Phenotypic changes resulted by affected communications of MVA- and MEP

pathways amongst *PtHMGR-* and *PtDXR-OEs* in 45-day-old poplars. a(I), The PtDXR

transgenic revealed a higher stem length than *PtHMGR-OEs* and NT poplars. **a**(II), The *PtHMGR*

transgenic presents an insignificantly more stem development than NT poplar. **a**(III), NT poplar

was used as a control; Scale bar represents 1 cm. **b**, The Box and Whisker mean comparison

plot of stem lengths revealed significantly higher lengths *PtDXR-OEs* than NT poplars

compared with *PtHMGR-OEs*. *PtHMGR* transgenics also revealed significantly higher lengths

than NT poplars. **c** and **d**, The Violin mean comparison plots of *ZEP* and *NCED* relative

expressions between *PtHMGR-*and *PtDXR-OEs* compared to NT poplars. **e**, The Box and

Whisker mean comparison plot of stem diameters revealed less significant differences

between *PtDXR-OEs* and NT poplars. Stars reveal significant differences, *P < 0.05, **P < 0.01,

***P < 0.001, ****P < 0.0001.

Figure 6 | Communications exist between MVA- and MEP-pathways excess of IPP and

DMAPP. The IPP and DMAPP are considered the common precursors of the MEP- and MVA

pathways between cytoplasm and plastid. In addition, the putative communication generated

between MVA- and MEP-related genes and MVA- and MEP-derived products. MVA: mevalonic

acid, MEP: methylerythritol phosphate, IPP: isopentenyl diphosphate, DMAPP: dimethylallyl

diphosphate, AACT: acetoacetyl CoA thiolase, HMGS: 3-hydroxy-3-methylglutaryl-CoA

synthase, HMG-CoA: 3-hydroxy-3-methylglutary-CoA, HMGR: 3-hydroxy-3-methylglutaryl-

CoA reductase, MVK: mevalonate kinase, MVD: mevalonate5-diphosphate decarboxylase, IPP:

isopentenyl diphosphate, IDI: IPP isomerase, GPP: geranyldiphosphate, FPP:

famesyldiphosphate, GPS: geranyl phosphate synthase, FPS: farnesyl-diphosphate synthase,

GPPS: geranyl diphosphate synthase, GGPPS: geranyl geranyl diphosphate synthase, DXS: 1-

deoxy-D-xylulose5-phosphate synthase, DXP: 1-deoxy-D-xylulose5-phosphate, DXR: 1-deoxy-

D-xylulose5-phosphate reductoisomerase, HDS: 1-hydroxy-2-methyl-2-(E)-butenyl4-

diphosphate synthase, HDR: 1-hydroxy-2-methyl-2-(E)-butenyl4-diphosphate reductase, MCT:

MEP cytidylyltransferase, CMK: 4-diphosphocytidyl-2-C-methyl-D-erythritol kinase.

Supplementary figures and table

Supplementary Figure 1 | Amino acid sequences alignment of PtHMGR protein and other known HMGR proteins. *A. thaliana* (NP_177775.2), *G. hirsutum* (XP_016691783.1), *M. domestica* (XP_008348952.1), *M. esculenta* (XP_021608133.1), *P. persica* (XM_020569919.1), *O. sativa* (XM_015768351.2), *T. cacao* (XM_007043046.2), *Z. mays* (PWZ28886.1). The HMG-CoA and NADPH binding domains are indicated in red rectangular.

Supplementary Figure 2 | Construction of a phylogenetic tree based on the HMGR sequences of various species. Accession numbers of the HMGR obtained from Phytozome are as follows: *A. thaliana* (AT1G76490 and AT2G17370) , *P. trichocarpa* (Potri.011G145000, Potri.005G257000, Potri.004G208500, Potri.001G457000, Potri.009G169900 and Potri.002G004000), *Gossypium raimondii* (Gorai.008G013000, Gorai.002G146000, Gorai.002G014700, Gorai.005G215800, Gorai.012G138100, Gorai.005G215500, Gorai.005G215600 and Gorai.005G215700), *Malus domestica* (MDP0000157996, MDP0000268909, MDP0000372490, MDP0000251253 and MDP0000312032), *Manihot esculenta* (Manes.15G114100, Manes.01G157500, Manes.03G096600, Manes.02G116900 and Manes.05G128600), *Oryza sativa* (LOC_Os09g31970, LOC_Os08g40180 and LOC_Os02g48330), *Prunus persica* (Prupe.7G187000, Prupe.7G187500 and Prupe.8G182300), *Theobroma cacao (*Thecc1EG000025, Thecc1EG007601 and Thecc1EG034814), and *Zea mays* (GRMZM2G393337, GRMZM2G058095, GRMZM2G136465, GRMZM2G001645 and GRMZM2G043503).

Supplementary Figure 3 | Molecular identification of *PtHMGR-OEs*. (**A**) PCR identification of *PtHMGR* in *PtHMGR-OEs* and NT poplars. Lane M: 15K molecular mass marker (TransGen, China); lane 1: genome DNA from WT as negative control; lanes 2–9: genome DNA from *PtHMGR-OE* lines (**B**) qRT-PCR identification of the transcript levels of *PtHMGR* in *PtHMGR-OEs* and NT poplars. Three independent experiments were performed; Stars reveal significant differences, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Supplementary Figure 4 | MEP- and MVA-related genes analyses in overexpressed *PtHMGR*- and *PtDXR-OEs* poplars. **a**, MVA-relate genes *AACT*, *HMGS*, *MVK*, *MVD,* and *FPS* affected by *PtHMGR* overexpressing. **b**, MVA-related genes *AACT*, *HMGS*, *HMGR*, *MVK*, *MVD,* and *FPS* affected by *PtDXR* overexpressing. **c**, MEP-related genes *DXS*, *MCT*, *CMK*, *HDS*, *HDR*, *IDI*, *GPS, GPPS, and DXR affected by PtHMGR overexpressing.* **d**, MEP-related genes *DXS*, *DXR*, *MCT*, *CMK*, *HDS*, *HDR*, *IDI*, *GPS,* and *GPPS* affected by *PtDXR* overexpressing. *PtActin* was used as an in

ternal reference in all repeats; * P < 0.05, ** P < 0.01, ***P < 0.001, ****P < 0.0001; Three in

dependent replications were performed in this experiment.

Supplementary Figure 5 | HPLC chromatograms of analyzing the contents of **(A)** β-carotene,

(B) lycopene, and **(C)** lutein in NT poplars and *PtHMGR-OEs*.

Supplementary Figure 6 | HPLC chromatograms of analyzing the contents of **(A)** β-carotene,

(B) lycopene, and **(C)** lutein in NT poplars and *PtDXR-OEs*.

Supplementary Figure 7 | Chromatogram analyses of GA3 standards via HPLC-MS/MS. The

chromatogram of standard GA3 at (**A**) 0.1, (**B**) 0.2, (**C**) 0.5, (**D**) 2, (**E**) 5, (**F**) 20, (**G**) 50, and (**H**)

200 ng/mL concentrations. (**I**) Equations for the GA3 standard curves.

Supplementary Figure 8 | Chromatogram analyses of tZR standards via HPLC-MS/MS. The

chromatogram of standard tZR at (**A**) 0.1, (**B**) 0.2, (**C**) 0.5, (**D**) 2, (**E**) 5, (**F**) 20, (**G**) 50, and (**H**)

200 ng/mL concentrations. (**I**) Equations for the tZR standard curves.

Supplementary Figure 9 | Chromatogram analyses of IPA standards via HPLC-MS/MS. The

chromatogram of standard IPA at (**A**) 0.2, (**B**) 0.5, (**C**) 2, (**D**) 5, (**E**) 20, (**F**) 50, and (**G**) 200 ng/mL

concentrations. (**H**) Equations for the IPA standard curves.

Supplementary Figure 10 | Chromatogram analyses of DCS standards via HPLC-MS/MS. The

chromatogram of standard DCS at (**A**) 0.5, (**B**) 2, (**C**) 10, (**D**) 20, and (**E**) 50 ng/mL concentrations.

(**F**) Equations for the DCS standard curves.

Supplementary Figure 11 | Chromatogram analyses of CS standards via HPLC-MS/MS. The

chromatogram of standard CS at (**A**) 0.5, (**B**) 5, (**C**) 10, (**D**) 20, and (**E**) 50 ng/mL concentrations.

(**F**) Equations for the CS standard curves.

Supplementary Table 1 | Primers were used in this study.

Supplementary Table 2 | Table of data analyses used in phenotypic changes evaluation. **a**,

Stem diameter data analyses. **b**, Stem length data analyses**.**

Figure 1 | MVA- and MEP-related genes analyses in overexpressed *PtHMGR-* and *PtDXR-OEs* poplars. a, Mean comparison of relative expression of MVA-relate genes *AACT*, *HMGS*, *MVK*, *MVD,* and *FPS* (Indicated in red) affected by *PtHMGR* overexpressing. **b**, Mean comparison of relative expression of MEP-related genes *DXS*, *MCT*, *CMK*, *HDS*, *HDR*, *IDI*, *GPS,* and *GPPS* (Indicated in red) affected by *PtHMGR* overexpressing; *HMGR* and *DXR*, which were overexpressed respectively by *DXR-* and *HMGR-OEs,* were presented in Supplementary Figure 4. *PtActin* was used as an internal reference in all repeats; "ns" means not significant, * P < 0.05, ** P < 0.01, ***P < 0.001, ****P < 0.0001; Three independent replications were performed in this experiment.

Figure 2 | HPLC-MS/MS content analyses of lycopene, β-carotene, lutein, and real-time PCR of *ZEP* and *NCED* genes family. HPLC-MS/MS content analyses have been performed to show the effect of *PtHMGR-OEs* on **a**, lycopene **b**, β-carotene, and **c**, lutein. Relative expressions have been analyzed affected by *PtHMGR-OEs* compared with NT poplars of **d**, *ZEP*, and **e**, *NCED* genes family. Bars represent mean ± SD (n = 3); Stars reveal significant differences, * P < 0.05, ** P < 0.01, *** P < 0.001, ****P < 0.0001; Three independent experiments were performed in these analyses.

Figure 3 | HPLC-MS/MS content analyses of lycopene, β-carotene, lutein, and real-time PCR of *ZEP* and *NCED* genes family. HPLC-MS/MS content analyses have been performed to show the effect of *PtDXR-OEs* on **a**, lycopene **b**, β-carotene, and **c**, lutein. Relative expressions have been analyzed affected by *PtDXR-OEs* compared with NT poplars of **d**, *ZEP*, and **e**, *NCED* genes family. Bars represent mean ± SD (n = 3); Stars reveal significant differences, * P < 0.05, ** P < 0.01, *** P < 0.001, ****P < 0.0001; Three independent experiments were performed in these analyses.

Figure 4 | HPLC-MS/MS content analyses of MEP- and MVA-derived isoprenoids. a,b,c,d, and e, Violin plots reveal the contents of isoprenoids GA3, tZR, IPA, DCS, and CS obtained from MEP- and MVA pathways influenced by *PtHMGR-* and *PtDXR-OEs*. **f,g,h,i, and j,** the column plots reveal the effect of *PtHMGR-OE3* and *-7* and *PtDXR-OE1* and *-3* on the mentioned above isoprenoids separately; NT poplars have been used as the control. Bars represent mean ± SD (n = 3); Stars reveal significant differences, *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. **k,l,m,n, and o,** represent the HPLC-MS/MS chromatogram content analyses of GA3, tZR, IPA, DCS, and CS, respectively affected by *PtHMGR- and PtDXR-OEs* compared with NT poplars.

Figure 5 | Phenotypic changes resulted by affected communications of MVA- and MEP pathways amongst *PtHMGR-* and *PtDXR-OEs* in 45-day-old poplars. a(I), The PtDXR transgenic revealed a higher stem length than *PtHMGR-OEs* and NT poplars. **a**(II), The *PtHMGR* transgenic presents an insignificantly more stem development than NT poplar. **a**(III), NT poplar was used as a control; Scale bar represents 1 cm. **b**, The Box and Whisker mean comparison plot of stem lengths revealed significantly higher lengths *PtDXR-OEs* than NT poplars compared with *PtHMGR-OEs*. *PtHMGR* transgenics also revealed significantly higher lengths than NT poplars. **c** and **d**, The Violin mean comparison plots of *ZEP* and *NCED* relative expressions between *PtHMGR-*and *PtDXR-OEs* compared to NT poplars. **e**, The Box and Whisker mean comparison plot of stem diameters revealed less significant differences between *PtDXR-OEs* and NT poplars. Stars reveal significant differences, *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Figure 6 | Communications exist between MVA- and MEP-pathways excess of IPP and DMAPP. The IPP and DMAPP are considered the common precursors of the MEP- and MVA pathways between cytoplasm and plastid. In addition, the putative communication generated between MVA- and MEP-related genes and MVA- and MEP-derived products. MVA: mevalonic acid, MEP: methylerythritol phosphate, IPP: isopentenyl diphosphate, DMAPP: dimethylallyl diphosphate, AACT: acetoacetyl CoA thiolase, HMGS: 3-hydroxy-3-methylglutaryl-CoA synthase, HMG-CoA: 3-hydroxy-3-methylglutary-CoA, HMGR: 3-hydroxy-3-methylglutaryl-CoA reductase, MVK: mevalonate kinase, MVD: mevalonate5-diphosphate decarboxylase, IPP: isopentenyl diphosphate, IDI: IPP isomerase, GPP: geranyldiphosphate, FPP: famesyldiphosphate, GPS: geranyl phosphate synthase, FPS: farnesyl-diphosphate synthase, GPPS: geranyl diphosphate synthase, GGPPS: geranyl geranyl diphosphate synthase, DXS: 1-deoxy-D-xylulose5-phosphate synthase, DXP: 1-deoxy-D-xylulose5-phosphate, DXR: 1-deoxy-D-xylulose5-phosphate reductoisomerase, HDS: 1-hydroxy-2-methyl-2-(E)-butenyl4-diphosphate synthase, HDR: 1-hydroxy-2-methyl-2-(E)-butenyl4-diphosphate reductase, MCT: MEP cytidylyltransferase, CMK: 4-diphosphocytidyl-2-C-methyl-D-erythritol kinase.